The latitudinal gradient in the rate of evolution of a biotic interaction trait

Benjamin G Freeman[1,2], Dolph Schluter[1,2*], Joseph A Tobias[3*]

[1] Biodiversity Research Centre, University of British Columbia, Vancouver, Canada.

[2] Department of Zoology, University of British Columbia, Vancouver, Canada.

[3] Department of Life Sciences, Imperial College London, London, UK.

* shared senior authorship

Abstract

Where is evolution fastest? The biotic interactions hypothesis proposes that greater

species richness creates more ecological opportunity, driving faster evolution at low

latitudes, whereas the "empty niches" hypothesis proposes that ecological opportunity is

greater where diversity is low, spurring faster evolution at high latitudes. Here we test

these contrasting predictions by analyzing rates of bird beak evolution for a global dataset

of 1141 sister pairs of birds. We find that beak size evolves at similar rates across

latitudes, while beak shape evolves faster in the temperate zone, consistent with the

empty niches hypothesis. By comparing rates of evolution in sympatric versus allopatric

sister lineages, we also find evidence that character displacement is similar between

latitudes. Our analyses focusing on recent speciation events finds that drivers of

evolutionary diversification are more potent at higher latitudes, thus calling into question

multiple hypotheses invoking faster tropical evolution to explain the latitudinal diversity

gradient.

Introduction

The biotic interactions hypothesis proposes strong tropical biotic interactions are both a cause and a consequence of the latitudinal diversity gradient (Schemske 2009). Imagine that diversity were equal in the tropics and temperate zone. If so, then stronger biotic interactions in the tropics are proposed to lead to faster adaptation and speciation at low latitudes, generating an excess of tropical species. This is because biotic interactions are thought to be the dominant drivers of selection for tropical organisms, and adaptation to biotic interactions is expected to lead to faster evolutionary change than adaptation to abiotic challenges (Dobzhansky 1950; MacArthur 1972). Now imagine that the latitudinal diversity gradient already exists. A greater number of species in the tropics could provide increased ecological opportunity via an increase in the diversity of possible interactions between species and greater spatial heterogeneity in these interactions, also leading to faster evolution (Schemske 2009). In this conception, diversity begets diversity in a positive feedback loop. Whether biotic interactions are stronger in the tropics has been tested by many studies, with mixed results (Schemske *et al.* 2009; Anstett *et al.* 2016; Moles & Ollerton 2016; Roslin *et al.* 2017; Baskett & Schemske 2018; Hargreaves *et al.* 2019; Freeman *et al.* 2020; Roesti *et al.* 2020). However, the prediction that stronger biotic interactions or higher diversity can drive faster evolution in the tropics remains largely unexplored (but see Benkman 2013).

An alternative is that evolutionary rates are instead greatest where diversity is low. According to standard niche theory, ecological opportunity should be negatively related to diversity (Hutchinson 1959), such that evolutionary rates are fastest where diversity is low (Schluter 2016; Schluter & Pennell 2017; Machac 2020). A geographic

prediction arising from this "empty niches" hypothesis is that evolutionary rates should

be faster in the species-poor temperate zone than the species-rich tropics.

Here, we investigate these contrasting predictions by analyzing latitudinal

differences in rates of bird beak evolution. Since birds use their beaks to feed on living

things, bird beaks can be considered a key trait mediating biotic interactions. Interactions

with the animals and plants that form bird's diets are widely thought to drive beak

morphological evolution (Smith 1987; Benkman 2003; Lerner *et al.* 2011; Grant & Grant

2014). Diet can also drive beak morphological evolution via interactions between

individuals competing for food, leading to character displacement between species

(Brown & Wilson 1956; Schluter 2000; Grant & Grant 2006). Bird beak morphology

may also be shaped by adaptation to abiotic factors such as thermoregulation (Tattersall

et al. 2009, 2017). However, beak morphological form is tightly coupled to dietary

ecological function across all ~10,000 species of birds, suggesting a pervasive role for

diet-related biotic interactions in driving bird beak evolution (Pigot *et al.* 2020).

We tested whether empirical patterns of bird beak evolution better matched

predictions of the biotic interactions or empty niche hypotheses, using new measurements

of beak size and shape for a previously published global dataset of bird sister pairs

(Cooney *et al.* 2017b). First, we tested whether beak size and shape evolution are fastest

for tropical sister pairs, as expected under the biotic interactions hypothesis, or fastest for

temperate sister pairs, as expected under the empty niches hypothesis. Second, we

examined one specific biotic interaction—competition between closely related species in

sympatry that results in greater trait divergence. This pattern includes both character

displacement, where competition directly drives trait divergence (Stuart & Losos 2013;

Weber & Strauss 2016), and ecological sorting, where competition prevents sympatry

between species with similar traits but permits sympatry between species with divergent

traits (Pigot & Tobias 2013; Tobias *et al.* 2014; Freeman 2015). If competition between

sympatric relatives leads to faster trait divergence in the tropics, as predicted by the biotic

interactions hypothesis, then evolutionary rates for sympatric sister pairs (relative to

baseline rates for allopatric sister pairs) should be faster in the tropics than the temperate

zone. The empty niches hypothesis makes no clear prediction about latitudinal gradients

in the strength of competition between close relatives.

Our analysis of global patterns of bird beak evolution sheds light on the more

general problem of why species richness is highest in the tropics. Dozens of hypotheses

have been proposed to explain the latitudinal diversity gradient. The biotic interactions

hypothesis is one of at least seven distinct hypotheses that invoke faster tropical evolution

to explain why species richness peaks near the equator (Mittelbach *et al.* 2007). These

"fast tropical evolution" hypothesis propose different mechanisms for why evolution is

most rapid in the tropics, including stronger biotic interactions (a focus of this

manuscript), faster rates of molecular evolution (the evolutionary speed hypothesis

(Rohde 1992; Gillman & Wright 2014)), narrower physiological tolerances (Janzen

1967), larger area (Rosenzweig 1995), greater climatic variation (Haffer 1969), greater

genetic drift (Fedorov 1966), and greater divergence in the face of gene flow (Moritz *et*

al. 2000). Finding faster rates of beak evolution in the tropics would therefore support the

biotic interactions hypothesis, but also be consistent with the entire set of fast tropical

evolution hypotheses. A result that beak evolution is faster in the temperate zone,

however, would support the empty niches hypothesis while calling into question all

hypotheses based on the concept of faster tropical evolution.

Materials and methods

Sister pair selection and geographic variables

We analyzed a previously published dataset of 1306 sister pairs of birds with information

on median latitude and range overlap for each pair (Cooney *et al.* 2017b). This dataset

defined sister pairs from a dated global bird phylogeny of species, subsetted to include

only species placed on the tree on the basis of genetic data (6670 species out of 9993)

(Jetz *et al.* 2012). Each sister pair in this dataset comprises two species that are each

other's closest relatives. Because of incomplete sampling, not all sister pairs are true

sister species. Nevertheless, each sister pair is a phylogenetically independent comparison

between two closely related species, and is therefore appropriate for our purposes.

Because our primary goal is to compare rates of beak evolution between tropical and

temperate sister pairs, we excluded 141 sister pairs where one species was classified as

tropical (median latitude < 23.4°) and the other as temperate (median latitude > 23.4°).

We further excluded the small number (1%) of sister pairs with estimated divergences

greater than 20 million years, and 11 sister pairs where morphological data was lacking

for one species within the pair. A total of 1141 sister pairs met these criteria.

Three key variables in this dataset are: (1) divergence time; (2) latitude; and (3)

range overlap, coded as allopatric vs. sympatric. We used divergence time—the time to

most recent common ancestor estimated using the dated phylogeny—as an estimate of

evolutionary age. We used the absolute latitude of each sister pair to classify it as tropical

(< 23.4°; N = 800) or temperate (> 23.4°; N = 341). Range overlap is a continuous

variable ranging from 0, where the two species in the pair do not overlap in range, to 1,

where the range of the smaller-ranged species is completely subsumed within the range

of the larger-ranged species. Following previous studies (Tobias *et al.* 2014; Cooney *et*

al. 2017b), we used a threshold of 20% range overlap to define whether sister pairs were

sympatric or allopatric (< 20% range overlap; N = 746) or sympatric (> 20% range

overlap; N = 395). We chose this threshold so that sister pairs were coded as sympatric

only when they had substantial range overlap (such that the two species likely interact).

To examine sensitivity of results, we conducted further tests using thresholds of > 0%

and > 50% range overlap to define sympatry/allopatry.

Morphological divergence

We extracted morphological data for each species in each species pair from a global

dataset of bird morphology (Pigot *et al.* 2020). We analyzed four traits, all measured in

mm: (1) bill length of the culmen; (2) bill length measured from the nares; (3) bill depth;

and (4) bill width. We excluded individuals that had missing values for any beak traits.

Our final morphological dataset contained 9,966 individuals (mean 4.4 individuals per

species; range = 1 – 89; 94% of species had four or more individuals measured).

132   We used a principal components analysis to place each of the 9,966 measured

individuals within a single multidimensional beak morphospace. To generate independent

axes of variation in beak morphology, we ran a principal component analysis (PCA) on

logged values of each of the four beak traits. The resulting morphospace is essentially

two dimensional: PC1, related to overall beak size, explained 82.1% of variation, while

PC2, a metric of beak shape, explained a further 14.3% of variation (Table S1, Figure

S1). We quantified the volume of beak morphospace for tropical and temperate zone

species by constructing a hypervolume using Gaussian kernel density estimation and

calculating its volume using the "hypervolume" package (Blonder *et al.* 2014; Blonder &

Harris 2019). Tropical species outnumber temperate species by ~ 3 to 1 in this dataset, as

in reality. However, the volumes of beak morphospace occupied by tropical and

temperate species in this dataset are nearly identical (four-dimensional volumes of 4.94

versus 4.86 and two-dimensional volumes of 1.00 versus 1.01, see Figure S2 for a two-

dimensional representation). This implies that tropical species in this dataset are more

densely packed within beak morphospace than temperate species, a feature found in other

avian diversity gradients (Pigot *et al.* 2016). We calculated divergence in beak size (PC1)

and shape (PC2) for each taxon pair as the absolute value of the difference in species

means along these axes. Sampling error will cause the raw difference in species' means

to be larger than the true difference in species means. We therefore estimated bias-

corrected divergence in beak size and shape for each sister pair using pooled sample

variance, calculated for all species (see demonstration of bias, as well as code to calculate

bias-corrected divergence, in a R script provided in the Supporting Information).

Following bias correction, some values of beak size and shape divergence between sister

pairs are negative.

Statistical analysis

We tested the hypothesis that evolutionary rates of beak divergence differ between

latitudes by fitting evolutionary and empirical models to our trait data in R version 3.9 (R

Development Core Team 2020). We fit distinct models to beak size (PC1) and beak

shape (PC2). For all beak size models, we removed two outlier sister pairs (the temperate

pair *Numenius minutus* and *N. arquata* (curlews) and the tropical pair *Geospiza*

magnirostris and *G. fuliginosa* (Darwin's finches from the Galápagos) with very low

estimated divergence dates and extremely high beak size divergence. These cases are

likely to represent phylogenetic error or instances where repeated introgression at target

loci generates spuriously short branch lengths among interacting lineages (Tobias et al.

2020). We first fit Brownian motion (BM) and Ornstein Uhlenbeck (OU) models. These

common models of trait evolution pass through the origin. In our dataset, however, both

beak size and beak shape divergence are relatively high even at very young evolutionary

ages (Figure 1a and 1b).

There are two likely explanations for this seemingly impossible initial rate of

evolution. The first is statistical: evolutionary age is estimated with error, and any species

pairs whose ages are underestimated will lead to models having an intercept fitting better

than models fit through the origin. The second explanation is biological: rapid evolution

(or phenotypic plasticity) can lead to trait differences that arise before any divergence at

the putatively neutral loci that are used to estimate evolutionary age. We therefore

explored fitting simple models with intercepts that better captured the patterns in our data

than formal evolutionary models. We interpret a high intercept as equivalent to a high

rate of evolution in the early stages of divergence, and hence we compare intercepts as

well as slopes of fitted models. We included simple power functions in the set of models

because they are more flexible than the evolutionary models, and can fit relationships

suggested by the loess regressions in Figure 1. In sum we fit BM and OU models, a BM

model with an intercept, a power function forced through the origin, and a power function

with an intercept, and compared model fit using AIC.

The response variable in our models (either beak size divergence or beak shape

divergence) was the signed square root of squared differences between bias-corrected

species means divided by two. BM and OU models estimate the variance between species

times two on the squared differences scale; we correct for this by dividing by two. We

used the signed square root of squared differences—the visual scale for analyzing trait

differences—because residuals are more normally distributed and had more homogenous

variance on this scale. All results from models fit to squared differences data, weighting

by the variance of residuals, were similar, but the highly skewed distributions of residuals

on the squared difference scale led us to be cautious in interpretation of AIC scores and

p-values (Figures S3 and S4, Tables S2 and S3).

We tested whether rates of trait evolution differed between the tropics and

temperate zone by fitting models with and without a latitudinal zone term (temperate or

tropical), and conducting F tests using the "anova" command in R. For the BM intercept

and power function with intercept, we fit three models in which (1) neither the intercept

nor the slope varied between tropical and temperate zone sister pairs; (2) the intercept

varied between tropical and temperate zone sister pairs; and (3) both slope and intercept

varied between tropical and temperate zone sister pairs.

Last, we analyzed the relationship between trait evolution and range overlap of

sister pairs in our dataset (Figure 1c). For both beak size and beak shape, we tested

whether rates of trait evolution differed between sympatric and allopatric sister pairs by

fitting three models using the power function with intercept (the best-fit model for both

beak size and beak shape) in which (1) neither the intercept nor the slope varied between

allopatric and sympatric sister pairs; (2) the intercept varied between allopatric and

sympatric sister pairs; and (3) both slope and intercept varied between allopatric and

sympatric sister pairs. We compared these models by conducting F tests using the

"anova" command in R. The key range overlap prediction arising from the biotic

interactions hypothesis is that evolutionary rates in sympatry are particularly fast in the

tropics. We tested this hypothesis by fitting three models of the best-fit power function

with intercept that included a sympatric/allopatric term in which (1) neither the intercept

nor the slope varied between tropical and temperate zone sister pairs; (2) the intercept

varied between tropical and temperate zone sister pairs; and (3) both slope and intercept

varied between tropical and temperate zone sister pairs with and without a

tropical/temperate term. We compared these models by conducting F tests using the

"anova" command in R.

Literature review

Last, we placed the results of our study in the broader context of latitudinal gradients in

rates of evolution. To do so, we reviewed the peer-reviewed published literature to find

previous studies of latitudinal gradients in evolutionary rates. BGF conducted Google

Scholar searches on 30 June 2020 for "molecular evolution AND latitudinal gradient",

"trait evolution AND latitudinal gradient", "speciation AND latitudinal gradient", and

"evolution AND latitudinal gradient", and examined the first 100 results for each search,

as well as additional studies cited within the papers uncovered via these searches.

Results

We found evolutionary rates were faster in the temperate zone for beak shape and similar

between tropical and temperate zone birds for beak size (Figures 2 and S5). For both beak

size and beak shape, the best-fit model was a power function with an intercept, and

standard evolutionary models (Brownian motion and Ornstein Uhlenbeck) were poor fits

(Δ AIC > 40; Table 1). For beak shape but not beak size, models that included a

latitudinal zone term (tropical/temperate) improved model fit (e.g., for the best-fit power

function with intercept, $p = 0.0034$; Table 2), with higher intercepts for beak shape

divergence in the temperate zone. We further found evidence that the slope of beak shape

divergence is steeper in the temperate zone ($p = 0.039$, model where both intercept and

slope differ between tropics and temperate zone plotted in Figure 2b): young lineages in

the temperate zone have greater beak shape divergence than equivalently aged lineages in

the tropics (higher intercept), but successively older lineages in the temperate zone have

even greater beak shape divergence than equivalently aged lineages in the tropics (steeper

slope).

Sympatric sister pairs have greater trait divergence than allopatric sister pairs for

beak size (Figure 3a, t-test $p = 0.020$, means = 0.17 and 0.14, respectively) and beak

shape (Figure 3b, t-test $p < 0.001$, means = 0.077 and 0.061, respectively). This pattern of

greater divergence in sympatry is consistent with a role for character displacement.

However, beak size and beak shape divergence also increase with evolutionary age

(Figure 1 a,b), and sympatric sister pairs are older than allopatric sister pairs. In this

dataset, median sister pair ages are 2.90 and 4.54 million years for tropical allopatric and

sympatric sister pairs, respectively (t-test on mean age; $p < 0.001$), and 2.30 and 3.60

million years for temperate zone allopatric and sympatric sister pairs (t-test on mean age;

$p = 0.0083$; Figure 3). For beak shape, including a range overlap term

(sympatry/allopatry) improved model fit ($p = 0.0022$, Figure 4b), consistent with a role

for character displacement in driving beak shape evolution. In contrast, models that

included a range overlap term did not improve model fit for beak size ($p = 0.20$, Figure

4a), indicating that greater divergence in beak size for sympatric sister pairs is primarily

due to their older age rather than character displacement. Models that included a

temperate/tropical term in addition to a sympatry/allopatry term further improved fit for

beak shape ($p = 0.0032$, Figure 4d) but not for beak size ($p = 0.69$, Figure 4c). However,

excess rates of divergence in sympatry compared to allopatry for beak shape were similar

between the tropics and temperate zone (Figure 4d). Results were similar when using

alternative thresholds to define sympatric versus allopatric sister pairs (Figures S6 and

S7, Tables S4 and S5).

Most studies measuring latitudinal variation in recent evolutionary rates,

particularly studies with large sample sizes, have found that rates are either fastest in the

temperate zone or similar between the temperate zone and tropics (Table 4).

Discussion

Our analyses of evolutionary rates across 1141 pairs of sister species reveal that bird beak

shape evolution is faster in the temperate zone whereas bird beak size evolution is similar

in rate across latitudes. In addition, sympatric tropical sister pairs do not have particularly

fast evolutionary rates, as predicted if one specific biotic interaction—competition

between closely related species that leads to character displacement—were stronger in

the species-rich tropics. These results contradict both aspects of the biotic interaction

hypothesis: that stronger biotic interactions in the tropics lead to faster evolution in biotic

interaction traits, and that greater species richness leads to faster evolution in the tropics.

Diversity does not beget diversity, at least for beak morphology. Instead, they are more

consistent with the empty niches hypothesis that proposes ecological opportunity is

greatest in the temperate zone where species richness is low.

It is perhaps unsurprising that ecological opportunity for bird beaks appears to be

greater in species-poor temperate regions. Bird beaks are well described by a two

dimensional morphospace, and there is little a priori reason to suspect the size of this

morphospace expands in response to increasing bird species richness. It is more likely

that a greater variety of resources in the tropics permits a larger beak morphospace

volume in tropical birds. However, beak morphospace is only slightly larger in the tropics

in this dataset. This implies that greater diversity is primarily associated with greater

niche packing rather than with niche expansion (Pigot *et al.* 2016), with this increased

saturation of available niche space offering fewer opportunities for rapid trait evolution

(Tobias *et al.* 2020). Diversity may beget diversity in other situations. For example,

greater diversity could spur rapid evolution (and coevolution) in biotic interaction traits

that are more multidimensional, such as behavioral traits or chemical traits associated

with defense or attraction. Studies that measured rates of evolution along the latitudinal

gradient for traits that differ in their dimensionality would test this idea.

It could also be argued that investigating rates of evolution among avian sister

species, in which both lineages universally share the same trophic level, runs counter to

some of the theoretical underpinnings of the biotic interactions hypothesis. For example,

earlier conceptualizations of this idea point towards diversity in lower trophic levels, i.e.

producers, having a knock-on effect on lineage diversification in consumers (Schemske et

al. 2009). Our results suggest that further studies of diversity-begets-diversity

mechanisms should not focus on within trophic levels, but across levels, to assess

whether the higher overall volume of tropical diversification (Schluter 2016) sends

ecological opportunities cascading through networks of interactions across the structure

of biodiversity as a whole.

A further caveat relates to the precision of our beak shape quantification. Our data

are based on a set of linear beak measurements and thus provide only a partial estimate of

aspects such as curvature. The beaks of two interacting hummingbird species may

potentially diverge in the extent of curvature linked to different floral niches while

appearing identical from the perspective of our measurements. It is clear that the linear

measurements we use nonetheless explain most of the ecological variation among

species, including relatively subtle behavioral differences (Pigot et al. 2020), but further

studies could revisit latitudinal effects on beak evolution using methods that account for

curvature, bill tip differences and other details of beak structure (Cooney *et al.* 2017a).

Sympatry is weakly associated with bird beak evolution

Based on linear measurements, we find that beak shape evolution is faster in sympatry

than allopatry across latitudes, while rates of beak size evolution are similar between

sympatric and allopatric sister pairs. We interpret faster evolution of beak shape in

sympatry to indicate a role for competition in driving divergence, either via character

displacement or ecological sorting (Tobias *et al.* 2020). Discriminating between these

alternative pathways is complicated by the fact they leave very similar signatures on

spatial patterns of trait divergence, but further sampling of sister pairs in allopatry vs.

sympatry across a range of latitudes may help to tease them apart. Beak size divergence is

greater for sympatric sister pairs than for allopatric sister pairs, but this appears to be

explained by the older age of sympatric sister pairs (Tobias *et al.* 2014). Given the lack of

differences in beak size evolution between sympatric and allopatric sister pairs, and the

relatively minor differences in beak shape evolution, our results provide further evidence

that character displacement (and ecological sorting) play a relatively minor role in bird

beak evolution (Tobias *et al.* 2020)

Comparison with previous studies of evolutionary rates across latitudes

Our study adds to a growing body of evidence documenting that recent evolutionary rates

are often faster in the temperate zone than in the tropics (see Table 4). While early

empirical tests reported faster molecular evolution in the tropics (e.g., Gillooly *et al.*

2005; Wright *et al.* 2006), a recent comprehensive analysis of ~8,000 taxon pairs from

six phyla of animals found that latitudinal differences in the rate of molecular evolution

are minimal (faster in the tropics in 51.6% of comparisons vs. faster in the temperate

zone in 48.4% of comparisons) (Orton *et al.* 2019). Consequently, the evolutionary speed

hypothesis, which proposes that higher temperatures lead to faster molecular evolution

via faster chemical reactions and faster generation times, appears unlikely to be generally

true. For trait evolution and speciation, a majority of studies report faster evolution in the

temperate zone. Song, plumage and body mass evolution in New World birds are fastest

in the temperate zone (Martin *et al.* 2010; Weir & Wheatcroft 2011; Lawson & Weir

2014), and speciation rates are fastest at high latitudes in marine fishes (Rabosky *et al.*

2018) and in New World birds and mammals (Weir & Schluter 2007) (but see (Rolland *et*

al. 2014). While evidence remains mixed—indeed, we find no latitudinal patterns for

beak size evolution and a fairly small difference for beak shape—the growing body of

available evidence suggests that rates of recent evolution are often faster in the temperate

zone, and seldom faster in the tropics (Table 4).

Conclusion

Explaining the latitudinal diversity gradient is a longstanding goal of evolutionary

ecology. The generality of the latitudinal diversity gradient across taxa, space and time

suggests the tantalizing possibility of a common explanation. Since Darwin and Wallace,

generations of biologists have added new hypotheses to a growing list of contenders

(reviewed by Pianka 1966; Rohde 1992; Mittelbach *et al.* 2007). We argue that, on the

strength of current evidence, hypotheses that invoke faster evolution in the tropics to

explain the latitudinal diversity gradient should be rejected. Not only do bird beaks

evolve either faster in the temperate zone or at similar rates across latitudes (this study),

but rates of molecular evolution are similar in animals across latitudes (Orton *et al.*

2019), and trait evolution and speciation are faster in the temperate zone for several

animal taxa (see Table 4). Evolutionary hypotheses that invoke slower extinction in the

tropics (rather than faster evolution) remain viable, though difficult to test. For example,

strong biotic interactions in the tropics may help explain the maintenance of high tropical

diversity (lower extinction in the tropics) rather than the origin of high tropical diversity

(faster evolution in the tropics). Last, we call attention to the repercussions of faster

evolution at high latitudes for the shape of the latitudinal diversity gradient. We might

expect that, all else equal, faster evolution and speciation at high latitudes would

gradually flatten of the latitudinal diversity gradient, all else equal. However, the large

majority of species currently inhabit the tropics. If per-capita speciation rates are only

somewhat faster in the temperate zone than in the tropics, then the latitudinal diversity

gradient may actually become steeper over time despite faster evolution and speciation in

the temperate zone.

Acknowledgements

BGF was supported by postdoctoral fellowships from the Biodiversity Research Centre

and Banting Canada (#379958). Morphological data collection was conducted by

Catherine Sheard, Monte Neate-Clegg, and Nico Alioravainen, among many others, and

funded by the UK Natural Environment Research Council (NE/I028068/1 and

NE/P004512/1 to JAT). None of our funders had any influence on the content of the

submitted manuscript, and none of our funders required approval of the final manuscript

to be published. Jason Weir provided helpful comments in response to an earlier

presentation of this work. Comments from the Schluter lab group and Ralf Yorque

greatly improved this manuscript.

References

Anstett, D.N., Nunes, K.A., Baskett, C. & Kotanen, P.M. (2016). Sources of controversy

surrounding latitudinal patterns in herbivory and defense. *Trends Ecol. Evol.*, 31,

789–802.

Baskett, C.A. & Schemske, D.W. (2018). Latitudinal patterns of herbivore pressure in a

temperate herb support the biotic interactions hypothesis. *Ecol. Lett.*, 21, 578–587.

Benkman, C.W. (2003). Divergent selection drives the adaptive radiation of crossbills.

Evolution, 57, 1176–1181.

Benkman, C.W. (2013). Biotic interaction strength and the intensity of selection. *Ecol.*

Lett., 16, 1054–1060.

Blonder, B. & Harris, D.J. (2019). hypervolume: high dimensional geometry and set

operations using kernel density estimation, support vector machines, and convex

hulls. https://CRAN.R-project.org/package=hypervolume

Blonder, B., Lamanna, C., Violle, C. & Enquist, B.J. (2014). The n-dimensional

hypervolume. *Glob. Ecol. Biogeogr.*, 23, 595–609.

Bromham, L. & Cardillo, M. (2003). Testing the link between the latitudinal gradient in

species richness and rates of molecular evolution. *J. Evol. Biol.*, 16, 200–207.

Brown, W.L. & Wilson, E.O. (1956). Character displacement. *Syst. Zool.*, 5, 49–64.

Cooney, C.R., Bright, J.A., Capp, E.J.R., Chira, A.M., Hughes, E.C., Moody, C.J.A., *et*

al. (2017a). Mega-evolutionary dynamics of the adaptive radiation of birds. *Nature*,

542, 344–347.

Cooney, C.R., Tobias, J.A., Weir, J.T., Botero, C.A. & Seddon, N. (2017b). Sexual

selection, speciation and constraints on geographical range overlap in birds. *Ecol.*

Lett., 20, 863–871.

Dobzhansky, T. (1950). Evolution in the tropics. *Am. Sci.*, 38, 209–221.

Fedorov, A.A. (1966). The structure of the tropical rain forest and speciation in the humid

tropics. *J. Ecol.*, 54, 1-11.

Freeman, B., Scholer, M., Boehm, M., Heavyside, J. & Schluter, D. (2020). Adaptation

and latitudinal gradients in species interactions: nest predation in birds. *Am. Nat.*, in

press.

Freeman, B.G. (2015). Competitive interactions upon secondary contact drive elevational

divergence in tropical birds. *Am. Nat.*, 186, 470–479.

Gillman, L.N., Keeling, D.J., Ross, H.A. & Wright, S.D. (2009). Latitude, elevation and

the tempo of molecular evolution in mammals. *Proc. R. Soc. B Biol. Sci.*, 276, 3353–

3359.

Gillman, L.N., Mccowan, L.S.C. & Wright, S.D. (2012). The tempo of genetic evolution

in birds: Body mass and climate effects. *J. Biogeogr.*, 39, 1567–1572.

Gillman, L.N. & Wright, S.D. (2014). Species richness and evolutionary speed: The

influence of temperature, water and area. *J. Biogeogr.*, 41, 39–51.

Gillooly, J.F., Allen, A.P., West, G.B. & Brown, J.H. (2005). The rate of DNA evolution:

Effects of body size and temperature on the molecular clock. *Proc. Natl. Acad. Sci.

U. S. A.*, 102, 140–145.

Grant, P.R. & Grant, B.R. (2006). Evolution of character displacement in Darwin's

finches. *Science*, 313, 224–226.

Grant, P.R. & Grant, B.R. (2014). *40 Years of Evolution: Darwin's Finches on Daphne

Major Island*. Princeton University Press, Princeton, N.J.

Haffer, J. (1969). Speciation in amazonian forest birds. *Science*, 165, 131–137.

Hargreaves, A.L., Suárez, E., Mehltreter, K., Myers-smith, I., Vanderplank, S.E., Slinn,

H.L., *et al.* (2019). Seed predation increases from the Arctic to the Equator and from

high to low elevations. *Sci. Adv.*, 5, eaau4403.

Hutchinson, G.E. (1959). Homage to Santa Rosalia or why are there so many kinds of

animals? *Am. Nat.*, 93, 145–159.

Janzen, D.H. (1967). Why mountain passes are higher in the tropics. *Am. Nat.*, 101, 233–

249.

Jetz, W., Thomas, G.H., Joy, J.B., Hartmann, K. & Mooers, A.O. (2012). The global

diversity of birds in space and time. *Nature*, 491, 444–448.

Lawson, A.M. & Weir, J.T. (2014). Latitudinal gradients in climatic-niche evolution

accelerate trait evolution at high latitudes. *Ecol. Lett.*, 17, 1427–1436.

Lerner, H.R.L., Meyer, M., James, H.F., Hofreiter, M. & Fleischer, R.C. (2011).

Multilocus resolution of phylogeny and timescale in the extant adaptive radiation of

Hawaiian honeycreepers. *Curr. Biol.*, 21, 1838–1844.

Lourenço, J.M., Glémin, S., Chiari, Y. & Galtier, N. (2013). The determinants of the

molecular substitution process in turtles. *J. Evol. Biol.*, 26, 38–50.

MacArthur, R.H. (1972). *Geographical ecology: patterns in the distribution of species*.

Princeton University Press, Princeton.

Machac, A. (2020). The dynamics of bird diversity in the New World. *Syst. Biol.*, 0, 1–

20.

Martin, P.R., Montgomerie, R. & Lougheed, S.C. (2010). Rapid sympatry explains

greater color pattern divergence in high latitude birds. *Evolution*, 64, 336–347.

Mittelbach, G.G., Schemske, D.W., Cornell, H. V, Allen, A.P., Brown, J.M., Bush, M.B.,

et al. (2007). Evolution and the latitudinal diversity gradient: speciation, extinction

and biogeography. *Ecol. Lett.*, 10, 315–331.

Moles, A.T. & Ollerton, J. (2016). Is the notion that species interactions are stronger and

more specialized in the tropics a zombie idea? *Biotropica*, 48, 141–145.

Moritz, C., Patton, J.L., Schneider, C.J. & Smith, T.B. (2000). Diversification of

rainforest faunas: an integrated molecular approach. *Annu. Rev. Ecol. Syst.*, 31, 533–

563.

Orton, M.G., May, J.A., Ly, W., Lee, D.J. & Adamowicz, S.J. (2019). Is molecular

evolution faster in the tropics? *Heredity*, 122, 513–524.

Pianka, E.R. (1966). Latitudinal gradients in species diversity : a review of concepts. *Am.*

Nat., 100, 33–46.

Pigot, A.L., Sheard, C., Miller, E.T., Bregman, T., Freeman, B., Seddon, N., *et al.* (2020).

Macroevolutionary convergence connects morphological form to ecological function

in birds. *Nat. Ecol. Evol.*, 4, 230-239.

Pigot, A.L. & Tobias, J.A. (2013). Species interactions constrain geographic range

expansion over evolutionary time. *Ecol. Lett.*, 16, 330–338.

Pigot, A.L., Trisos, C.H. & Tobias, J.A. (2016). Functional traits reveal the expansion

and packing of ecological niche space underlying an elevational diversity gradient in

passerine birds. *Proc. R. Soc. B Biol. Sci.*, 283, 1–9.

R Development Core Team. (2020). R: A language and environment for statistical

computing.

Rabosky, D.L., Chang, J., Title, P.O., Cowman, P.F., Sallan, L., Friedman, M., *et al.*

482 (2018). An inverse latitudinal gradient in speciation rate for marine fishes. *Nature*,

483 559, 392–395.

484 Rabosky, D.L., Title, P.O. & Huang, H. (2015). Minimal effects of latitude on present-

485 day speciation rates in New World birds. *Proc. R. Soc. B Biol. Sci.*, 282.

486 Roesti, M., Anstett, D.N., Freeman, B.G., Lee-Yaw, J.A., Schluter, D., Chavarie, L., *et*

487 *al.* (2020). Pelagic fish predation is stronger at temperate latitudes than near the

488 equator. *Nat. Commun.*, 11, 1527.

489 Rohde, K. (1992). Latitudinal gradients in species diversity : the search for the primary

490 cause. *Oikos*, 65, 514–527.

491 Rolland, J., Condamine, F.L., Jiguet, F. & Morlon, H. (2014). Faster speciation and

492 reduced extinction in the tropics contribute to the mammalian latitudinal diversity

493 gradient. *PLoS Biol.*, 12.

494 Rolland, J., Loiseau, O., Romiguier, J. & Salamin, N. (2016). Molecular evolutionary

495 rates are not correlated with temperature and latitude in Squamata: An exception to

496 the metabolic theory of ecology? *BMC Evol. Biol.*, 16, 1–6.

497 Rosenzweig, M.L. (1995). *Species Diversity in Space and Time*. Cambridge University

498 Press, Cambridge, UK.

499 Roslin, T., Hardwick, B., Novotny, V., Petry, W.K., Andrew, N.R., Asmus, A., *et al.*

500 (2017). Latitudinal gradients: Higher predation risk for insect prey at low latitudes

501 and elevations. *Science*, 356, 742–744.

502 Schemske, D. (2009). *Biotic interactions and speciation in the tropics*. In *Speciation and*

503 *patterns of Diversity*. Cambridge University Press.

504 Schemske, D.W., Mittelbach, G.G., Cornell, H. V, Sobel, J.M. & Roy, K. (2009). Is there

a latitudinal gradient in the importance of biotic interactions? *Annu. Rev. Ecol. Evol.*

Syst., 40, 245–269.

Schluter, D. (2000). Ecological character displacement in adaptive radiation. *Am. Nat.*,

156, 4–16.

Schluter, D. (2016). Speciation, ecological opportunity, and latitude. *Am. Nat.*, 187, 1–18.

Schluter, D. & Pennell, M.W. (2017). Speciation gradients and the distribution of

biodiversity. *Nature*, 546, 48–55.

Smith, T.B. (1987). Bill size polymorphism and intraspecific niche utilization in an

African finch. *Nature*, 329, 717–719.

Stuart, Y.E. & Losos, J.B. (2013). Ecological character displacement: glass half full or

half empty? *Trends Ecol. Evol.*, 28, 402–408.

Tattersall, G.J., Andrade, D. V. & Abe, A.S. (2009). Heat exchange from the toucan bill

reveals a controllable vascular thermal radiator. *Science*, 325, 468–470.

Tattersall, G.J., Arnaout, B. & Symonds, M.R.E. (2017). The evolution of the avian bill

as a thermoregulatory organ. *Biol. Rev.*, 92, 1630–1656.

Tobias, J.A., Cornwallis, C.K., Derryberry, E.P., Claramunt, S., Brumfield, R.T. &

Seddon, N. (2014). Species coexistence and the dynamics of phenotypic evolution in

adaptive radiation. *Nature*, 506, 359–363.

Tobias, J.A., Ottenburghs, J. & Pigot, A.L. (2020). Avian diversity: speciation,

macroevolution and ecological function. *Annu. Rev. Ecol. Evol. Syst.*, in press.

Weber, M.G. & Strauss, S.Y. (2016). Coexistence in close relatives: beyond competition

and reproductive isolation in sister taxa. *Annu. Rev. Ecol. Evol. Syst.*, 47, 359–381.

Weir, J.T. & Price, T.D. (2019). Song playbacks demonstrate slower evolution of song

discrimination in birds from Amazonia than from temperate North America. *PLoS*

Biol., 17, e3000478.

Weir, J.T. & Schluter, D. (2007). The latitudinal gradient in recent speciation and

extinction rates of birds and mammals. *Science*, 315, 1574.

Weir, J.T. & Wheatcroft, D. (2011). A latitudinal gradient in rates of evolution of avian

syllable diversity and song length. *Proc. R. Soc. B-Biological Sci.*, 278, 1713–1720.

Wright, S., Keeling, J. & Gillman, L. (2006). The road from Santa Rosalia: A faster

tempo of evolution in tropical climates. *Proc. Natl. Acad. Sci. U. S. A.*, 103, 7718–

7722.

Wright, S.D., Gillman, L.N., Ross, H.A. & Keeling, D.J. (2010). Energy and the tempo of

evolution in amphibians. *Glob. Ecol. Biogeogr.*, 19, 733–740.

Figure 1. Patterns of divergence in beak size (a), beak shape (b) and range overlap (c) in

1141 sister pairs of birds. Raw data are plotted. Loess regressions are shown in blue;

predictions from the best-fit models (power functions with intercepts) are shown as

dashed black lines. Beak size and shape divergence values are corrected for bias arising

from sampling error. Range overlap is the proportion of the smaller-ranged species that

falls within the range of the larger-ranged species. Sister pairs were coded as sympatric if

they had range overlaps > 0.20 (the dashed line).

547
Figure 2. Rates of beak size evolution are similar between tropics and temperate zone

(left panels), but beak shape evolution is faster in the temperate zone (right panels).

Model predictions are plotted for the two best models: power functions with an intercept

(a, b) and Brownian motion models with an intercept (c, d). *P*-values are from *F* tests

testing whether the inclusion of a tropical/temperate term improved model fit. Δ AIC

values compare different model fits separately for beak size and beak shape.

Figure 3. Sympatric sister pairs have greater beak size divergence (a), beak shape

divergence (b), and are older (c) than allopatric sister pairs. *P*-values are from t-tests of

trait divergence or ages between allopatric and sympatric sister pairs; separate t-tests for

age for temperate and tropical zones in panel (c). Median values of ages for temperate

and tropical sister pairs are plotted as vertical dashed lines.

Figure 4. Rates of beak size evolution are similar between allopatric (n = 746) and

sympatric (n = 395) sister pairs (left panels), but beak shape evolution is faster in

sympatry (right panels). *P*-values are from *F* tests testing whether the inclusion of an

allopatric/sympatric term to a power function with an intercept improved model fit (a, b),

or whether the inclusion of a tropical/temperate term to a power function with an

intercept and an allopatric/sympatric term improved model fit (c, d). The *p*-value for beak

shape (d) is from a *F* test comparing a reduced model with an allopatric/sympatric term to

a full model with terms allowing both the intercept and slope to differ between tropics

and temperate zone. Beak shape evolution is faster in the temperate zone in both allopatry

and sympatry compared to the tropics.

573 Table 1. Models that included intercepts (power function with an intercept and Brownian

574 motion model with an intercept) were better fits than models forced through the origin

575 (power function, Ornstein Uhlenbeck and Brownian motion models).

response variable	model	Δ AIC
beak size	power function intercept	0
	Brownian motion intercept	-15.11
	power function	-25.04
	Ornstein Uhlenbeck	-42.36
	Brownian motion	-42.38
beak shape	power function intercept	0
	Brownian motion intercept	-3.98
	power function	-12.99
	Ornstein Uhlenbeck	-40.15
	Brownian motion	-62.01

587   Table 2. Latitudinal zone is a predictor of beak shape evolution but not beak size

588   evolution. *P*-values are from *F* tests comparing model fit between full and reduced

589   models.

response variable	reduced model	full model	*p*-value
beak size	power intercept	power intercept + latzone (intercept)	0.52
	power intercept + latzone (intercept)	power intercept + latzone (intercept and slope)	0.70
	Brownian motion intercept	Brownian motion intercept + latzone (intercept)	0.41
	Brownian motion intercept + latzone (intercept)	Brownian motion intercept + latzone (intercept and slope)	0.25
	power	power + latzone	0.74
	Ornstein Uhlenbeck	Ornstein Uhlenbeck + latzone	0.58
	Brownian motion	Brownian motion + latzone	0.54
beak shape	power intercept	power intercept + latzone (intercept)	0.0034
	power intercept + latzone (intercept)	power intercept + latzone (intercept and slope)	0.039
	Brownian motion intercept	Brownian motion intercept + latzone (intercept)	0.0026
	Brownian motion intercept + latzone	Brownian motion intercept + latzone (intercept and slope)	0.91
	power	power + latzone	0.052
	Ornstein Uhlenbeck	Ornstein Uhlenbeck + latzone	0.0072
	Brownian motion	Brownian motion + latzone	0.0013

592 Table 3. Range overlap ("patry") is a predictor of beak shape evolution but not beak size

593 evolution. *P*-values are from *F* tests comparing model fit between full and reduced

594 models.

response variable	reduced model	full model	*p*-value
beak size	power intercept	power intercept + patry (intercept)	0.20
	power intercept + patry (intercept)	power intercept + patry (intercept and slope)	0.11
	power intercept + patry (intercept)	power intercept + patry (intercept) + lat zone (intercept)	0.64
	power intercept + patry (intercept) + lat zone (intercept)	power intercept + patry (intercept) + lat zone (intercept and slope)	0.59
beak shape	power intercept	power intercept + patry (intercept)	0.0022
	power intercept + patry (intercept)	power intercept + patry (intercept and slope)	0.10
	power intercept + patry (intercept)	power intercept + patry (intercept) + lat zone (intercept)	0.011
	power intercept + patry (intercept) + lat zone (slope & intercept)	power intercept + patry (intercept) + lat zone (intercept intercept and slope)	0.026

Table 4. Studies of latitudinal variation in rates of molecular evolution, trait evolution

and recent speciation compiled from a survey of the peer-reviewed literature. The key

information is whether the fastest evolutionary rate was found to occur in the tropics, the

temperate zone, or if was no latitudinal gradient in evolutionary rates. Most studies,

particularly those with large sample sizes, find either faster evolutionary rates in the

temperate zone or no latitudinal gradient in evolutionary rates.

study	taxa	fastest rate	citation
molecular evolution (mtDNA)	45 lineage pairs of birds	no gradient	(Bromham & Cardillo
molecular evolution (ITS)	45 lineage pairs of plants	tropics	(Wright *et al.* 2006)
molecular evolution (mtDNA)	130 lineage pairs of mammals	tropics	(Gillman *et al.* 2009)
molecular evolution (mtDNA)	94 lineage pairs of amphibians	tropics	(Wright *et al.* 2010)
molecular evolution (mtDNA)	30 lineage pairs of birds	tropics	(Gillman *et al.* 2012)
molecular evolution (mtDNA and nuclear DNA)	224 species of turtles	tropics	(Lourenço *et al.* 2013)
molecular evolution (branch lengths)	1651 species of squamates	no gradient	(Rolland *et al.* 2016)
molecular evolution (mtDNA)	8037 lineage pairs from six animal phyla (Arthropoda, Chordata, Mollusca, Annelida, Echinodermata, and Cnidaria)	tropics, but very small effect size (51.6% vs. 48.4%)	(Orton *et al.* 2019)
trait evolution (color)	78 sister pairs of New World songbirds	temperate zone	(Martin *et al.* 2010)
trait evolution (song)	116 sister pairs of New World	temperate	(Weir & Wheatcroft 2(

	songbirds	zone	
trait evolution (song perception)	109 pairs of songbirds in temperate North America vs. Amazon	temperate zone	(Weir & Price 2019)
trait evolution (climatic niche, body mass)	111 sister pairs of New World songbirds	temperate zone	(Lawson & Weir 2014)
trait evolution (beak size)	1141 sister pairs of birds (global)	no gradient	this study
trait evolution (beak shape)	1141 sister pairs of birds (global)	temperate zone	this study
speciation rate	118 sister pairs of New World mammals	temperate zone	(Weir & Schluter 2007)
speciation rate	191 sister pairs of New World birds	temperate zone	(Weir & Schluter 2007)
speciation rate	all mammals (N = 5,020)	tropics	(Rolland *et al.* 2014)
speciation rate	all New World songbirds (N = 2,571)	no gradient	(Rabosky *et al.* 2015)
speciation rate	all ray-finned fishes (global; N = 31,526)	temperate zone	(Rabosky *et al.* 2018)