Zooplankton community beta diversity in an Amazonian floodplain lake

Leonardo Fernandes Gomes[1]*, Ana Caroline Alcântara Missias Gomes[1], Carla Albuquerque de

Souza[1], Hasley Rodrigo Pereira[1], Marie-Paule Bonnet[2,3], Ludgero Cardoso Galli Vieira[1]

[1]Núcleo de Estudos e Pesquisas Ambientais e Limnológicas - NEPAL, Faculdade UnB de Planaltina,

Área Universitária 1, Vila Nossa Senhora de Fátima, 73.345-010 – Planaltina – DF, Brasil

[2]Geosciences Environnement Toulouse (UMR 5563 GET), IRD/ CNRS/ Université Toulouse III,

Toulouse, France.

[3]International Joint Laboratory Observatoire des Changements Environnementaux LMI OCE, IRD –

Universidade de Brasília, Campus Universitário Darcy Ribeiro, CEP 70910-900 Brasília, Brazil.

*Corresponding author: leof.ciamb@gmail.com (e-mail)

Keywords: Lago Grande do Curuai, hydrological cycle, Podani, flood pulse, beta diversity

partitioning

Abstract

Understanding the mechanisms that generate organism distribution patterns from the beta diversity perspective can assist in environmental monitoring strategies. In this study, we emphasized the limnic zooplankton due to the ability of these organisms to respond quickly to environmental variations. Therefore, we evaluated the following questions: (i) Do different regions of the same lake have the same importance in contributing to beta diversity? (ii) Do beta diversity and its components vary over the hydrological cycle? (iii) What is the importance of local and spatial predictors in beta diversity and its components? (iv) Do beta diversity and its components show a consistent pattern throughout the hydrological cycle? We found that the contribution of different sites to diversity was more associated with regions with low abundance and richness of organisms values, such as the littoral and *igarapes*, which shows the relevance of these areas for biological monitoring and for the delimitation of priority areas for the zooplankton diversity conservation. Despite the peculiarities of each hydrological period and regarding beta diversity components, we verified a species substitution and differences in abundance pattern in the lake. We also found low concordance patterns between the periods and low environmental and spatial variables prediction on beta diversity patterns.

Introduction

Species can present different distribution patterns in response to natural factors such as competition, predation, dispersive processes limitations, and/or local and regional environmental variables influences (Guisan & Thuiller, 2005). These factors may be intensified by human activities, which makes even more relevant to understand the mechanisms that generate such structuring patterns in biological communities. Thus, the understanding of these mechanisms can assist in the formulation of efficient environmental monitoring strategies and, even, in the delimitation of priority areas for conservation in several ecosystems (Socolar *et al.*, 2016). The comparative diversity across multiple sites, known as beta diversity (Whittaker, 1960), has undergone advances over the years both for understanding patterns of presence-absence of organisms and for density values per site (Baselga, 2010; Podani & Schmera, 2011; Podani, Ricotta & Schmera, 2013).

Both for organism occurrence and abundance, Podani family of beta diversity (Podani & Schmera, 2011; Podani *et al.*, 2013) can be partitioned into the following main components: (i) *species similarities*: commonly measured by the Jaccard index for presence-absence data and Ruzicka, for abundance data. High values of this partition mean that the pairs of sites put in comparison share many species or species with similar abundances; (ii) *difference in relative richness/abundance*: is the difference in species richness, or species abundance, between pairs of

sites. Therefore, high values of that partition show that the number of species or specimens between

compared sites is discrepant; (iii) *species replacement/abundance:* it can be maximized when there is

a high replacement of species, or species with equivalent abundances, along an environmental

gradient or between pairs of sites. Therefore, high replacement values for the abundance data mean

that, although the sites in comparison have similar abundances, the species composition is different.

Also, although the approach with abundance data may represent more subtle differences concerning

environmental variations, the values between the assessments for abundance and presence-absence

data can be quite different, even if evaluated with the same data set (Podani *et al.*, 2013).

The evaluation of the factors that influence beta diversity and its components can be even

more complex in floodplain lakes since they are predominantly dominated by the flood pulse that

controls the dynamics of entry and output of sediments, water and organisms that naturally contribute

for changes in biological diversity in the ecosystem (Junk *et al.*, 2012). These plains are continuously

or periodically flooded by direct precipitation or by the overflow of the main river and, depending on

the level of connectivity with the river, there may be a loss of connection between habitats during

periods of low water (Thomaz, Bini & Bozelli, 2007). However, as the cycle of extensive floodplains

is usually slow and monomodal, the biological dynamics of organisms can adapt in order to maximize

their performance according to hydrological cycles (Junk *et al.*, 2011).

In Amazonian rivers, the flow tends to be more intense and requires a high resilience capacity

of the organisms. Therefore, smaller aquatic organisms tend to be present with greater richness and

density in the lakes of these plains, where they can find shelter against predation and food (Junk,

Bayley & Sparks, 1989). Furthermore, according to the hydrological period, these organisms may

present beta diversity patterns that change over time (Bozelli *et al.*, 2015).

Assessing beta diversity and its components over space, but also highlighting whether the

pattern generated is consistent throughout the hydrological cycle is important in different aspects. For

example, due to the scarcity of financial resources and time allocated in environmental monitoring

programs and scientific research, if different hydrological periods show a concordant pattern of

diversity, there is a real possibility of adjustment in the sampling effort, reducing the number of

sampling campaigns, which would save financial resources and time. In the same way, it is possible

to use other alternatives as is the case of using lower taxonomic resolutions and or presence-absence

data instead of abundance data (Carneiro *et al.*, 2013; Vieira *et al.*, 2017; de Morais *et al.*, 2018).

In this study, we emphasized the limnic zooplankton due to the ability of these organisms to

respond quickly to environmental variations. Therefore, we evaluated the following questions: (i) Do

different regions of the same lake have the same importance in contributing to beta diversity? (ii) Do

beta diversity and its components vary over the hydrological cycle? (iii) What is the importance of

local (environmental characterization) and spatial (dispersive processes) predictors in beta diversity

and its components? (iv) Do beta diversity and its components show a consistent pattern throughout the hydrological cycle? Taking into account that the ecological dynamics of floodplains is temporally complex, we expected that the sites contribution to beta diversity would be different between hydrological periods. Besides, due to the spatial extent of the study area, we expected that species replacement patterns would be predominant, considering values of presence-absence of organisms, and patterns of differences in abundance, considering values of species abundance per site. Also, due to the complex interactions that dominate the occurrence of organisms, we expected that there would be a variation between environmental and spatial predictors in biological diversity patterns and, finally, as each period comprises a different hydrological dynamics, we did not expect to find many concordant values, being important to evelute in all hydrological periods to understand the distribution patterns of the zooplankton community.

Material and methods

Study area

The study area comprises an Amazonian floodplain lake called *Lago Grande do Curuai*, located in the State of Pará, Brazil. The majority of the water supply comes from the Amazon River (77%), while the others are subdivided between rainfall, runoff, and outcropping of groundwater (Bonnet *et al.*, 2008). The hydrological dynamics generate a monomodal cycle in this lake, comprising the periods of flooding (from January to the end of February), high water (from April to the end of June), flushing (from August to October) and low water (mid-October to November) (de Moraes Novo *et al.*, 2006).

The environmental characteristics of Lago Grande do Curuai are quite variable throughout the year, mainly concerning chlorophyll-*a* and dissolved oxygen. During the flooding period, chlorophyll-*a* levels are low enough for human consumption. However, the values in the flushing period rise to such an extent that water is not recommended for any type of activity (Affonso, Barbosa & Novo, 2011).

Sampling were carried out in 17 sample units (Fig. 1) in four campaigns: March / 2013 (flooding period), September / 2013 (flushing period), May / 2014 (high water period) and November / 2014 (low water period).

Fig. 1 Map of the study area and sampling units in Lago Grande do Curuai. Blue area: aquatic environments; beige area: terrestrial environments

Environmental variables

In each sampling unit, we used a multi-parameter YSY probe, model EXO2 to measure the variables dissolved oxygen (mg/L), blue-green algae (µg/L), fluorescent organic dissolved matter (raw), pH, water temperature (ºC), conductivity (µS/cm), total dissolved solids (mg/L), and turbidity (NTU). According to the protocol (APHA, 2005), water samples were obtained and frozen for further quantification in the laboratory of: alkalinity (mg/L), total chlorophyll (µg/L), total phosphorus (µg/L), total nitrogen (µg/L), total dissolved nitrogen (mg/L), ammonia (mg/L), nitrate (mg/L), and silica (mg/L).

Zooplankton

In each sampling unit, we sampled the zooplankton community on the subsurface (*ca* 50 cm). Therefore, we filtered 300 liters of water in a net with a 68 µm opening mesh. Samples were stored in polyethylene bottles, preserved with formaldehyde (5%), and buffered with sodium tetraborate. In the laboratory, the samples were concentrated in 75 mL. To quantify the densities of zooplanktonic organisms per sample unit, a 7.5 mL subsampling was performed with a *Hensen-Stempel* pipette. We read the subsampled organisms in a Sedgewick Rafter chamber for identification and counting using an optical microscope. Additionally, we carried out qualitative sampling to verify and record the existence of new *taxa* that were not identified during quantitative sampling (Bottrell *et al.*, 1976).

Data analysis

We performed a Local Contribution to Beta Diversity (LCBD) (Borcard, Gillet & Legendre,

2018) to obtain the degree of exclusivity of the sites in the species composition in each hydrological

period using the function *beta.div*, package *adespatial* (Dray *et al.*, 2018). To evaluate and partition

Podani family beta diversity by sample period, we used the function *beta.div.comp* of *adespatial*

package (Dray *et al.*, 2018). In both cases, we used the *Jaccard* index for presence and absence values

and *Ruzicka* for organism density data.

To verify if there were significant differences in the values resulting from the beta diversity

partitioning by period, we performed a Permutational Multivariate Analysis of Variance Using

Distance Matrices (PERMANOVA). We obtained these matrices using the *beta.div.comp* function for

both create a matrix encompassing all periods and generate matrices by pairs of periods. For

PERMANOVA, we use the *adonis2* function of the *vegan* package (Oksanen *et al.*, 2016) and the

matrices resulted from the partition were inserted in response to hydrological periods. Additionally,

we constructed triangular plots (simplex) to check the distributions of the pairs of sites concerning

the partitive components of beta diversity for both *Ruzicka* distance matrices and *Jaccard* in each

hydrological periods.

To assess the influence of environmental and spatial variables in the beta diversity partitions

of zooplankton community by hydrological period, we performed Distance-Based Redundancy

Analysis (dbRDA's) (Legendre & Andersson, 1999) with different matrices resulted from the beta

diversity partitioning (as response variables) and different environmental and spatial variables (as

predictor variables). To determine which variables would be inserted in the dbRDA, we performed

the analysis of variation inflation factor (VIF) (Borcard *et al.*, 2018), removing the environmental

variables that showed high collinearity in each sample period (VIF values greater than 20). To

determine the spatial predictors (geographic coordinates), we first converted the coordinates to

Cartesian distances using the *geoXY* function of the *SoDA* package (Chambers, 2013). Then, we

ordered the variables in a Distance-Based Moran's Eigenvector Maps (dbMEM) (Dray, Legendre &

Peres-Neto, 2006; Legendre & Legendre, 2012) using the dbmem function of the adespatial package

(Dray *et al.*, 2018).

To evaluate the temporal concordance in the distribution patterns of the different zooplankton

community beta diversity partitions between hydrological periods, we performed *Procrustes* tests

(Gower, 1975). For that, we ordered the matrices resulting from the beta diversity partitioning in

different Non-metric multidimensional scaling (NMDS), then we extracted the values from the

ordering scores and inserted them into the *protest* function, from *vegan* package (Oksanen *et al.*,

2013). To check the significance, 9999 permutations were performed.

For all the mentioned analyzes, we used the statistical software R (R Core Team, 2016).

Results

Regarding the contribution of sites to beta diversity (LCBD) using presence-absence data of

the zooplankton community, only the hydrological periods of flooding and low water presented sites

with significant contributions, with site 9 being important for the beta diversity in both periods (Fig.

2). All significant sites (8, 9, and 13) are located in the southern region of the lake. When we evaluated

the LCBD using abundance data (Fig. 3), the four periods presented significant sample units. In the

flooding and flushing periods, the significant sampling units were located in the north region of the

lake (sites 14 and 10, respectively); in the high waters, they were located in the south, and in the low

water period they were located in the west region of the lake.

Fig. 2 Map of the local contribution to beta diversity (LCBD) for zooplankton presence/absence data

with *Jaccard* matrix of the sample units by hydrological period. Filled circles represent sites with

significant contributions

Fig. 3 Map of the local contribution to beta diversity (LCBD) for zooplankton abundance data with Ruzicka matrix of the sample units by hydrological period. Filled circles represent sites with significant contributions

Evaluating the beta diversity partitions using presence and absence species data (Table 1), we verified a replacement dominant pattern (values comprised 73% to 81% of the beta diversity between hydrological periods), while we verified an abundance difference dominance pattern when using abundance data (values comprised 58% to 74% of the beta diversity between hydrological periods).

Table 1. Beta diversity partitioning for all hydrological periods with presence and absence and abundance values. BD = total beta diversity; Rep = replacement; RD = richness difference; AD = abundance difference; Rep/BD = ratio of replacement to total beta diversity; RD/BD = ratio of richness difference to total beta diversity; AD/BD = ratio of abundance difference to total beta diversity

	Period	BD	Rep	RD	Rep/BD	RD/BD
Presence-Absence (*Jaccard*)	Flooding	0.34	0.25	0.09	0.73	0.27
	Flushing	0.28	0.23	0.05	0.81	0.19
	High water	0.35	0.27	0.09	0.75	0.25
	Low water	0.28	0.22	0.06	0.78	0.22

	Period	BD	Rep	AD	Rep/BD	AD/BD
Abundance (*Ruzicka*)	Flooding	0.43	0.14	0.29	0.32	0.68
	Flushing	0.38	0.15	0.23	0.40	0.60
	High water	0.41	0.11	0.31	0.26	0.74
	Low water	0.36	0.15	0.21	0.42	0.58

When we compared the beta diversity partitions obtained by the hydrological periods (Table
2) using presence/absence data, the richness difference component was similar among all
hydrological periods, while the beta diversity and replacement component were different among them
all. When considering abundance data, the beta diversity component was different across all
hydrological periods, while the abundance difference component only was not different in flushing
and low water periods. There were no differences in the abundance replacement component.

Table 2. Permutational multivariate analysis of variance using distance matrices (PERMANOVA)
between the matrices resulting from the partition of the beta diversity for the different hydrological
periods. Significant values are in bold

	Period	Beta diversity (Jaccard)			Richness difference			Replacement		
		R^2	F	p	R^2	F	p	R^2	F	p
Presence-absence	Global	**0.31**	**9.78**	**0.001**	-0.04	-0.79	1.000	**0.38**	**12.84**	**0.001**
	Flooding x Flushing	**0.21**	**8.58**	**0.001**	-	-	-	**0.27**	**11.90**	**0.001**
	Flooding x High water	**0.18**	**7.15**	**0.001**	-	-	-	**0.22**	**9.27**	**0.001**
	Flooding x Low water	**0.25**	**10.66**	**0.001**	-	-	-	**0.32**	**15.26**	**0.001**
	Flushing x High water	**0.24**	**9.87**	**0.001**	-	-	-	**0.27**	**12.01**	**0.001**
	Flushing x Low water	**0.25**	**10.43**	**0.001**	-	-	-	**0.30**	**13.66**	**0.001**
	High water x Low water	**0.28**	**12.49**	**0.001**	-	-	-	**0.33**	**15.50**	**0.001**
	Period	Beta diversity (Ruzicka)			Abundance difference			Replacement		
		R^2	F	p	R^2	F	p	R^2	F	p
Abundance	Global	**0.20**	**5.32**	**0.001**	**0.27**	**7.92**	**0.001**	0.10	2.48	0.077
	Flooding x Flushing	**0.12**	**4.52**	**0.001**	**0.12**	**4.27**	**0.012**	-	-	-
	Flooding x High water	**0.10**	**3.63**	**0.001**	**0.14**	**5.35**	**0.004**	-	-	-
	Flooding x Low water	**0.13**	**4.66**	**0.001**	**0.09**	**3.19**	**0.035**	-	-	-
	Flushing x High water	**0.17**	**6.70**	**0.001**	**0.33**	**16.04**	**0.001**	-	-	-
	Flushing x Low water	**0.13**	**4.81**	**0.001**	-0.01	-0.27	0.997	-	-	-
	High water x Low water	**0.20**	**7.88**	**0.001**	**0.34**	**16.66**	**0.001**	-	-	-

In proportion, when we partitioned the beta diversity using presence-absence data, the pairs

of sample units were more associated with greater similarities and replacement values considering all

periods (Fig. 4). On the other hand, when we evaluated the partition using abundance data, the pairs

of sample units were more associated with abundance difference component and, secondly, with

higher replacement levels (Fig. 5).

Fig. 4 Triangular graph (simplex) of the proportion of elements of the beta diversity partition per pair

of sample units for values of presence-absence of organisms. RichDiff = richness difference and

Repl= species replacement.

Fig. 5 Triangular graph (simplex) of the proportion of elements in the beta diversity partition per pair of sample units for organism abundance values. AbDiff = abundance difference and Repl = species abundance replacement.

Because presented high collinearity or multicollinearity values, we removed the following environmental variables of each hydrological period: total chlorophyll, pH, conductivity and total dissolved solids (flooding); dissolved oxygen, pH, conductivity and total dissolved nitrogen (flushing); temperature, conductivity and total dissolved solids (high water) and dissolved oxygen, blue-green algae, pH, conductivity and total dissolved nitrogen (low water).

The environmental and spatial variables showed little influence on the distribution patterns of beta diversity and its components, regardless the hydrological period (Table 3). Considering the presence-absence species data, the environmental variables explained the beta diversity patterns in

flushing and low waters periods and the richness difference component in in the low water period.
Regarding the abundance data, the environmental variables explained the beta diversity and the
abundance difference component in the high water period (Table 3).

Concerning the presence-absence values, spatial variables explained the beta diversity
patterns in flooding, flushing, and low water periods, and replacement component in flushing and low
water periods. However, concerning the abundance data, spatial variables did not explain beta
diversity nor its components in any of the hydrological periods analyzed (Table 3).

 Table 3. Distance-based redundancy analysis (dbRDA) of the influence of environmental and spatial predictors on the matrices resulting from the

 beta diversity partition. Rep = replacement; RD = richness difference; AD = abundance difference. Significant values are in bold

Data	Season	Environmental variables									Spatial variables								
		Beta diversity (Jaccard)			Rep			RD			Beta diversity (Jaccard)			Rep			RD		
		R^2_{adj}	F	p	R^2_{adj}	F	p	R^2_{adj}	F	p	R^2_{adj}	F	p	R^2_{adj}	F	p	R^2_{adj}	F	p
Presence-Absence	Flooding	0.07	1.09	0.290	0.01	1.01	0.453	0.09	1.13	0.425	**0.05**	**1.275**	**0.034**	0.01	1.06	0.251	0.03	1.19	0.300
	Flushing	**0.31**	**1.61**	**0.008**	0.09	1.13	0.125	0.37	1.79	0.124	**0.14**	**1.848**	**0.001**	**0.06**	**1.33**	**0.003**	0.02	1.12	0.356
	High water	0.08	1.11	0.238	0.00	0.97	0.665	0.32	1.57	0.200	0.02	1.114	0.216	0.00	0.93	0.841	0.14	1.87	0.084
	Low water	**0.31**	**1.66**	**0.005**	0.07	1.11	0.181	**0.45**	**2.19**	**0.019**	**0.15**	**1.926**	**0.002**	**0.05**	**1.26**	**0.020**	0.13	1.83	0.072
		Beta diversity (Ruzicka)			Rep			AD			Beta diversity (Ruzicka)			Rep			AD		
		R^2_{adj}	F	p	R^2_{adj}	F	p	R^2_{adj}	F	p	R^2_{adj}	F	p	R^2_{adj}	F	p	R^2_{adj}	F	p
Abundance	Flooding	0.02	1.02	0.494	0.02	1.02	0.375	-0.04	0.95	0.574	0.03	1.156	0.257	0.01	1.04	0.316	-0.01	0.95	0.462
	Flushing	0.06	1.09	0.407	-0.01	0.98	0.615	0.07	1.10	0.430	0.08	1.474	0.080	0.01	1.06	0.252	0.02	1.12	0.325
	High water	**0.54**	**2.42**	**0.007**	-0.10	0.89	0.974	**0.63**	**3.06**	**0.008**	0.07	1.372	0.159	-0.01	0.93	0.831	0.09	1.51	0.138
	Low water	0.25	1.49	0.089	-0.03	0.96	0.746	0.23	1.44	0.163	0.03	1.168	0.256	0.02	1.09	0.090	-0.02	0.88	0.577

Regarding the concordance analyzes, zooplankton beta diversity and its components

showed low values between hydrological periods (Table 4). Taking into account the presence

and absence species data, there was concordance of beta diversity only in the comparisons

between low water and flushing periods, and low water and high water periods (Table 4).

Concerning the beta diversity components, there was concordance only in the comparisons

between high water and low water (richness difference component) and between flooding and

flushing (richness replacement component) and flooding and low water (richness replacement

component). On the other hand, the abundance data did not show concordant patterns between

the hydrological periods.

Table 4. Procrustes test evaluating the concordance of beta diversity and its components values

between hydrological periods. Significant values are in bold

	Season	Beta diversity (Jaccard)		Richness difference		Replacement	
		r	p	r	p	r	p
	Flooding x Flushing	0.39	0.153	0.38	0.173	**0.62**	**0.003**
Presence-absence	Flooding x High water	0.42	0.096	0.17	0.832	0.41	0.124
	Flooding x Low water	0.45	0.088	0.26	0.468	**0.49**	**0.031**
	Flushing x High water	0.42	0.110	0.13	0.917	0.46	0.058
	Flushing x Low water	**0.47**	**0.045**	0.27	0.404	0.36	0.242
	High water x Low water	**0.69**	**0.001**	**0.66**	**0.001**	0.41	0.125

	Season	Beta diversity (Ruzicka)		Abundance difference		Replacement	
		r	p	r	p	r	p
	Flooding x Flushing	0.26	0.522	0.06	0.909	0.27	0.574
Abundance	Flooding x High water	0.35	0.205	0.25	0.277	0.27	0.543
	Flooding x Low water	0.35	0.241	0.22	0.398	0.19	0.823
	Flushing x High water	0.35	0.208	0.34	0.196	0.19	0.847
	Flushing x Low water	0.44	0.075	0.18	0.602	0.36	0.253
	High water x Low water	0.23	0.555	0.09	0.974	0.17	0.890

Discussion

Local contributions to beta diversity

When evaluating the beta diversity local contributions patterns using the presence-

absence data, we found that the main contribution sites were located in the south during flooding

and low water periods. This lake region has a higher proportion of areas with pastoral use

(Peres, Gurgel & Laques, 2018) and also the highest proportion of *igarapés* area. On the other

hand, the northern region connects more predominantly with the Amazon River (Bonnet *et al.*,

2008). Given that the variation in species composition and abundance influence the LCBD

contributions, the land use may have influenced the difference in species composition between

sites, what, consequently, influenced the increase the beta diversity contribution.

The significant LCBD site located in the southern region presented the lowest richness

of individuals per sampling unit during the flooding period, while in the flushing period, it

showed a different occurrence of species comparing to the other sites. High LCBD values may

not be directly associated with high richness or abundance values, since areas with low richness

and occurrences of differentiated species may also present higher contribution values, which

may denote these areas as priorities for species conservation (Legendre & De Cáceres, 2013).

In the flooding and flushing periods, some sites in the northern region were

differentiated concerning organism abundance. There was low abundance of zooplanktonic

organisms at the significant site during the flooding period. Moreover, in flushing, the sampling

unit ten, which most contributed to the beta diversity, stood out for the occurrence of different

species compared to others sampling units in the same period (e.g., *Lecane elsa*, *Lecane luna*

and *Nebela collaris*). In the low water period, the sampling unit also showed distinct species

(e.g., *Difflugia elegans*). This distinction in the diversity patterns of the sampling units by

hydrological period showed that the flood pulse promoted different dynamics in the floodplain

lake. In the low water period, the sampling units were isolated from the main river, which means

that the considerable environmental heterogeneity may have been influenced differences in

species with different characteristics in each sampling period (Thomaz *et al.*, 2007). In this

case, as the sampling unit two, which has a higher LCBD value, is on the opposite side of the

most important contribution area of the river's water flow, located to the east, the isolation of

the site may have justified such differentiation.

Whereas zooplanktonic organisms respond effectively to environmental variations

(Vieira *et al.*, 2011; Wang *et al.*, 2016) and even greater impacts such as hydrological changes

in cases of dams (Souza *et al.*, 2019), we consider that the sampling units highlighted

accordingly to the criteria of uniqueness by the LCBD analysis, being always associated with

the marginal regions of the lake. These regions have higher interactions with *igarapés* and are
in contact with the aquatic-terrestrial transition zones. Therefore, despite the hydrological
importance of the flood pulse over the lake and the control over ecological dynamics, it is also
important to take into account the importance that these *igarapés* and vegetation areas have for
the existence of unique sites in relation to biodiversity for the Lago Grande do Curuai.
309
Beta diversity partition
Related to the presence-absence data, there was a predominance of replacement
concerning total beta diversity. It means that, despite a greater constancy in species richness per
sampling unit, the species composition between pairs of units was different. Species are
expected to show a substitution pattern over large environmental gradients, depending on other
factors such as ecological tolerance of species (Legendre, 2014). Some studies report the
sensitivity of organisms in the zooplankton community to environmental variations (Vieira *et*
al., 2011), in some cases responding through changes in the trophic structure of the community
(Ejsmont-Karabin *et al.*, 2018) and changes in reproductive rates and species composition in
the presence of other organisms (e.g., fish) (Feniova *et al.*, 2019).
The high water period showed the highest beta diversity values and species replacement
rate. It differed from our expectations, since we expected a greater environmental homogeneity
and consequent biological homogeneity, reflecting a higher biological similarity between the
sites due to the flood pulse in the high water period and due to the greater interconnectivity
between habitats (Thomaz *et al.*, 2007; Bozelli *et al.*, 2015). Despite this, the increase in beta
diversity values may have been attributable to a greater interaction area with the floodplain that
began during the flooding period (Junk *et al.*, 1989) and continued to settle during the high
water period. This same pattern may have justified the lower beta diversity and replacement
values in low water and flushing periods where, despite the isolation of habitats promoted by
the reduction in the water volume, consequently minimized the interaction with the floodplain
region and the main river.
On the other hand, although the beta diversity patterns using abundance data were the
same for the presence-absence data with the highest values in the high water and flooding
periods, the abundance difference component predominated over the replacement component.
These values denote that, despite a greater tendency to replace species along the environmental
gradient, these species had wide variations in abundance values. It highlighted the importance
of understanding the zooplankton community abundance variations that, despite the ability to
respond to environmental variations (e.g., variation in trophic status and phosphorus

concentration in water), is often overlooked in some ecological studies (García-Chicote,

Armengol & Rojo, 2018).

Environmental and spatial predictors

Despite the distinctions observed in the patterns of similarity and substitution of species

between hydrological periods, we observed that the environmental variables showed little

prediction about the diversity patterns of the zooplankton community for presence and absence

data. These variables explained only the patterns of similarity in the flushing and low water

periods and the richness difference in the low water period.

On the other hand, there was a higher pattern of prediction of spatial variables over

patterns of similarity in the composition of species, not explaining only in the high water period.

These values denote that spatial variation may have a greater control over the organisms

composition dynamics than environmental variation. Despite this, this control was only related

to presence-absence values. The patterns of organism abundance and presence-absence refer to

different factors. For example, for presence and absence data, beta diversity corresponding to

the inverse of similarity in the composition is prioritized (Podani & Schmera, 2011), while for

abundance data, besides the composition, variations in the number of individuals of each

species are also considered. Therefore, when abundance is taken into account, sites with high

species dissimilarity values are those that present a high distinction in species composition and

the corresponding organisms abundance (Podani *et al.*, 2013).

Therefore, the explanation obtained in the low water period using the presence-absence

data may be related to the heterogeneity of ecological niches (Legendre, 2014). The low water

period may have promoted the existence of different niches, some with more species and others

with fewer species, due to the isolation. The substitution of species explained spatially may also

be based on the isolation that makes the species of an environment unable to reach other places

(Thomaz *et al.*, 2007). For this reason, spatial isolation can drive the pattern of differentiation

of species within the habitat and this same pattern may explain the spatial prediction in the

period of flushing.

For the beta diversity components using abundance data, there was a low standard of

explanation for both environmental and spatial variables, which showed that there was a greater

complexity of factors (e.g., competition and predation) that may have been the most responsible

for these variations and that were not evaluated in this study. This low pattern of response shows

that the zooplankton community is not responding only to environmental variations at that time,

but to changes that occurred in other periods before the sampling carried out. Besides, as the

abundance and presence-absence data responded differently to different factors, we emphasize

that both approaches can be complementary when used for biological monitoring purposes.

Temporal concordance between beta diversity components

Despite the occurrence of significant values when evaluating the temporal concordance

between the beta diversity patterns using presence-absence values, no pair of periods showed

concordance between all the diversity patterns over the hydrological cycle. There was also no

concordance between the periods using the abundance values. These results are in agreement

with our expectations since even in other environments, there is a low standard of predictability

and synchrony of zooplankton with other variables that allow us to predict a constant and

predictable pattern for this community (Vieira *et al.*, 2019).

These results also show that the environmental and biological dynamics of the

floodplains are complex to be predictable and, depending on the hydrological period, which

changes the entrance of river sediments and the inflow or outflow of water in the floodplain,

and the evaluated group, the structuring of the communities can be different (Amoros &

Bornette, 2002). There are proposals that the dynamics are so distinct and susceptible to

hydrological variations that the high water period acts as a resumption of the successional

regime of the structure and composition of the zooplankton community (Baranyi *et al.*, 2002;

Bozelli *et al.*, 2015). Therefore, despite the economic advantages of sampling in less

hydrological periods, we found that, in order to understand the beta diversity patterns of the

zooplankton community, sampling are necessary to occur in all the hydrological periods of high

and low waters, as well as in the flood and ebb intermediate periods.

Conclusions

Hydrological variations govern the zooplankton community dynamics, thus the

contribution of different locations depending on the hydrological period evaluated. With some

exceptions, the sites that most contributed to the beta diversity presented less organism richness

or abundance and also showed proximity to the coastal regions of the lake, especially those

associated with *Igarapés* when using the organisms presence-absence data. This result denotes

the relevance of these areas for biological monitoring and for the delimitation of priority areas

for the conservation of zooplankton diversity.

405   Beta diversity was greatest in flooding and high water periods. Despite the differences
406 in the partition values by hydrological period, the species replacement was dominant in all
407 hydrological periods using the organisms presence-absence data, while the abundance
408 difference was dominant using the quantitative values of organisms per sample unit. Therefore,
409 the studies must evaluate both abundance and presence-absence data as a complementary way,
410 considering that they can portray different processes in the face of environmental and spatial
411 variations. Due to the complexity of factors that govern the distribution of zooplankton
412 organisms in floodplains, there was a little prediction of environmental and spatial variables on
413 the beta diversity distribution patterns for the community. Also, there was a low concordance
414 between the patterns for the different hydrological periods, which highlights the need to study
415 the hydrological periods of high and low waters, as well as the transient periods of flooding and
416 flushing to obtain an adequate assessment of the dynamics distribution patterns of the
417 zooplankton community from the perspective of beta diversity.

419 **Acknowledgments**

421 The authors thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES)
422 for providing financial assistance to LFG, ACAMG, CAS and HRP. LCGV was supported by
423 productivity fellowships of Conselho Nacional de Desenvolvimento Científico e Tecnológico
424 (CNPq). We also thank the Fondation pour la Recherche sur la Biodiversité (FRB) and the
425 Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq); these groups, in
426 partnership with Institut de Recherche pour le Développement (IRD), financed the project
427 number: 490634/2013-3.

429 **Data availability statement**

431 Data are available on request from the corresponding author.

433 **References**

435 Affonso A., Barbosa C. & Novo E. (2011). Water quality changes in floodplain lakes due to
436   the Amazon River flood pulse: Lago Grande de Curuaí (Pará). *Brazilian Journal of*
437   *Biology* **71**, 601–610. https://doi.org/10.1590/S1519-69842011000400004
438 Amoros C. & Bornette G. (2002). Connectivity and biocomplexity in waterbodies of riverine

floodplains. *Freshwater Biology* **47**, 761–776. https://doi.org/10.1046/j.1365-
2427.2002.00905.x

APHA (2005). *Standard Methods for the Examination of Water and Wastewater*, 21st edn.
American Public Health Association/American Water Works Association/Water
Environment Federation, Washington DC.

Baranyi C., Hein T., Holarek C., Keckeis S. & Schiemer F. (2002). Zooplankton biomass and
community structure in a Danube River floodplain system: effects of hydrology.
Freshwater Biology **47**, 473–482. https://doi.org/10.1046/j.1365-2427.2002.00822.x

Baselga A. (2010). Partitioning the turnover and nestedness components of beta diversity.
Global Ecology and Biogeography **19**, 134–143. https://doi.org/10.1111/j.1466-
8238.2009.00490.x

Bonnet M.P., Barroux G., Martinez J.M., Seyler F., Moreira-Turcq P., Cochonneau G., *et al.*
(2008). Floodplain hydrology in an Amazon floodplain lake (Lago Grande de Curuaí).
Journal of Hydrology **349**, 18–30. https://doi.org/10.1016/j.jhydrol.2007.10.055

Borcard D., Gillet F. & Legendre P. (2018). *Numerical Ecology with R*. Springer International
Publishing, Cham.

Bottrell H.H., Duncan A., Gliwicz Z.M., Grygierek E., Herzig A., Hillbrichtilkowska A., *et al.*
(1976). Review of some problems in zooplankton production studies. *Norwegian Journal*
of Zoology **24**, 419–456

Bozelli R.L., Thomaz S.M., Padial A.A., Lopes P.M. & Bini L.M. (2015). Floods decrease
zooplankton beta diversity and environmental heterogeneity in an Amazonian floodplain
system. *Hydrobiologia* **753**, 233–241. https://doi.org/10.1007/s10750-015-2209-1

Carneiro F.M., Nabout J.C., Vieira L.C.G., Lodi S. & Bini L.M. (2013). Higher Taxa Predict
Plankton Beta-diversity Patterns Across an Eutrophication Gradient. *Natureza &*
Conservação **11**, 43–47. https://doi.org/10.4322/natcon.2013.006

Chambers J.M. (2013). SoDA: Functions and Examples for "Software for Data Analysis"

Dray S., Blanchet G., Borcard D., Clappe S., Guenard G., Jombart T., *et al.* (2018). adespatial:
Multivariate Multiscale Spatial Analysis. *R package version 0.1-1*

Dray S., Legendre P. & Peres-Neto P.R. (2006). Spatial modelling: a comprehensive
framework for principal coordinate analysis of neighbour matrices (PCNM). *Ecological*
Modelling **196**, 483–493. https://doi.org/10.1016/j.ecolmodel.2006.02.015

Ejsmont-Karabin J., Feniova I., Kostrzewska-Szlakowska I., Rzepecki M., Petrosyan V.G. &
Dzialowski A.R. (2018). Factors influencing phosphorus regeneration by lake
zooplankton—An experimental approach. *Limnologica* **70**, 58–64.

https://doi.org/10.1016/j.limno.2018.01.003

Feniova I., Sakharova E., Karpowicz M., Gladyshev M.I., Sushchik N.N., Dawidowicz P., *et*

al. (2019). Direct and Indirect Impacts of Fish on Crustacean Zooplankton in

Experimental Mesocosms. *Water* **11**, 2090. https://doi.org/10.3390/w11102090

García-Chicote J., Armengol X. & Rojo C. (2018). Zooplankton abundance: A neglected key

element in the evaluation of reservoir water quality. *Limnologica* **69**, 46–54.

https://doi.org/10.1016/j.limno.2017.11.004

Gower J.C. (1975). Generalized procrustes analysis. *Psychometrika* **40**, 33–51.

https://doi.org/10.1007/BF02291478

Guisan A. & Thuiller W. (2005). Predicting species distribution: offering more than simple

habitat models. *Ecology Letters* **8**, 993–1009. https://doi.org/10.1111/j.1461-

0248.2005.00792.x

Junk W.J., Bayley P.B. & Sparks R.E. (1989). The flood pulse concept in river-floodplain

systems. *Canadian special publication of fisheries and aquatic sciences* **106**, 110–127

Junk W.J., Piedade M.T.F., Schöngart J., Cohn-Haft M., Adeney J.M. & Wittmann F. (2011).

A Classification of Major Naturally-Occurring Amazonian Lowland Wetlands. *Wetlands*

31, 623–640. https://doi.org/10.1007/s13157-011-0190-7

Junk W.J., Piedade M.T.F., Schöngart J. & Wittmann F. (2012). A classification of major

natural habitats of Amazonian white-water river floodplains (várzeas). *Wetlands Ecology*

and Management **20**, 461–475. https://doi.org/10.1007/s11273-012-9268-0

Legendre P. (2014). Interpreting the replacement and richness difference components of beta

diversity. *Global Ecology and Biogeography* **23**, 1324–1334.

https://doi.org/10.1111/geb.12207

Legendre P. & Andersson M.J. (1999). Distance-based redundancy analysis: Testing

multispecies responses in multifactorial ecological experiments. *Ecological Monographs*

69, 1–24. https://doi.org/10.1890/0012-9615(1999)069[0001:DBRATM]2.0.CO;2

Legendre P. & De Cáceres M. (2013). Beta diversity as the variance of community data:

dissimilarity coefficients and partitioning. *Ecology Letters* **16**, 951–963.

https://doi.org/10.1111/ele.12141

Legendre P. & Legendre L. (2012). *Numerical ecology*, Third. Elsevier, Amsterdam.

de Moraes Novo E.M.L., de Farias Barbosa C.C., de Freitas R.M., Shimabukuro Y.E., Melack

J.M. & Filho W.P. (2006). Seasonal changes in chlorophyll distributions in Amazon

floodplain lakes derived from MODIS images. *Limnology* **7**, 153–161.

https://doi.org/10.1007/s10201-006-0179-8

de Morais G.F., dos Santos Ribas L.G., Ortega J.C.G., Heino J. & Bini L.M. (2018).

Biological surrogates: A word of caution. *Ecological Indicators* **88**, 214–218.

https://doi.org/10.1016/j.ecolind.2018.01.027

Oksanen J., Blanchet, Guillaume F., Friendly M., Kindt R., Legendre P., Mcglinn D., *et al.*

(2016). Vegan: Community Ecology Package. *R package version 2.4-0*

Oksanen J., Blanchet F.G., Kindt R., Legendre P., Minchin P.R., O'hara R.B., *et al.* (2013).

Package 'vegan.' *Community ecology package, version* **2**

Peres L.G.M., Gurgel H. & Laques A.-E. (2018). Dinâmica da paisagem em planícies de

inundação amazônicas: o caso do Lago Grande do Curuai, Pará, Brasil. *Confins. Revue*

franco-brésilienne de géographie/Revista franco-brasilera de geografia

Podani J., Ricotta C. & Schmera D. (2013). A general framework for analyzing beta diversity,

nestedness and related community-level phenomena based on abundance data.

Ecological Complexity **15**, 52–61. https://doi.org/10.1016/j.ecocom.2013.03.002

Podani J. & Schmera D. (2011). A new conceptual and methodological framework for

exploring and explaining pattern in presence - absence data. *Oikos* **120**, 1625–1638.

https://doi.org/10.1111/j.1600-0706.2011.19451.x

R Core Team (2016). R: A Language and Environment for Statistical Computing

Socolar J.B., Gilroy J.J., Kunin W.E. & Edwards D.P. (2016). How Should Beta-Diversity

Inform Biodiversity Conservation? *Trends in Ecology & Evolution* **31**, 67–80.

https://doi.org/10.1016/j.tree.2015.11.005

Souza C.A. de, Vieira L.C.G., Legendre P., Carvalho P. de, Velho L.F.M. & Beisner B.E.

(2019). Damming interacts with the flood pulse to alter zooplankton communities in an

Amazonian river. *Freshwater Biology*, fwb.13284. https://doi.org/10.1111/fwb.13284

Thomaz S.M., Bini L.M. & Bozelli R.L. (2007). Floods increase similarity among aquatic

habitats in river-floodplain systems. *Hydrobiologia* **579**, 1–13.

https://doi.org/10.1007/s10750-006-0285-y

Vieira A.C.B., Medeiros A.M.A., Ribeiro L.L. & Crispim M.C. (2011). Population dynamics

of Moina minuta Hansen (1899), Ceriodaphnia cornuta Sars (1886), and Diaphanosoma

spinulosum Herbst (1967) (Crustacea: Branchiopoda) in different nutrients (N and P)

concentration ranges. *Acta Limnologica Brasiliensia* **23**, 48–56.

https://doi.org/10.4322/actalb.2011.018

Vieira M.C., Bini L.M., Velho L.F.M., Gomes L.F., Nabout J.C. & Vieira L.C.G. (2017).

Biodiversity shortcuts in biomonitoring of novel ecosystems. *Ecological Indicators* **82**,

505–512. https://doi.org/10.1016/j.ecolind.2017.07.025

Vieira M.C., Roitman I., de Oliveira Barbosa H., Machado Velho L.F. & Cardoso Galli Vieira

L. (2019). Spatial synchrony of zooplankton during the impoundment of amazonic

reservoir. *Ecological Indicators* **98**, 649–656.

https://doi.org/10.1016/j.ecolind.2018.11.040

Wang C., Wang L., Deng D. & Zhou Z. (2016). Temporal and spatial variations in rotifer

correlations with environmental factors in Shengjin Lake, China. *Environmental Science*

and Pollution Research **23**, 8076–8084. https://doi.org/10.1007/s11356-015-6009-y

Whittaker R.H. (1960). Vegetation of the Siskiyou Mountains, Oregon and California.

Ecological Monographs **30**, 279–338. https://doi.org/10.2307/1943563

550