1 **Morphological and genomic shifts in mole-rat 'queens' increase fecundity but reduce**
2 **skeletal integrity**
3

4 Rachel A. Johnston[1*], Philippe Vullioud[2], Jack Thorley[2], Henry Kirveslahti[4], Leyao Shen[3],
5 Sayan Mukherjee[4,5,6,7], Courtney Karner[3,8], Tim Clutton-Brock[2,9], Jenny Tung[1,10,11,12*]

6
[1]Department of Evolutionary Anthropology, Duke University, Durham, North Carolina 27708,
USA
[2]Department of Zoology, University of Cambridge, Cambridge CB23EJ, UK
[3]Department of Orthopaedic Surgery, Duke Orthopaedic Cellular, Developmental, and Genome
Laboratories, Duke University School of Medicine, Durham, NC 27710, USA
[4]Department of Statistical Science, Duke University, Durham, NC 27708, USA
[5]Department of Computer Science, Duke University, Durham, NC 27708, USA
[6]Department of Mathematics, Duke University, Durham, NC 27708, USA
[7]Department of Bioinformatics & Biostatistics, Duke University, Durham, NC 27708, USA
[8]Department of Cell Biology, Duke University, Durham, NC 27710, USA
[9]Department of Zoology and Entomology, Mammal Research Institute, University of Pretoria,
0028, Pretoria, South Africa
[10]Department of Biology, Duke University, Durham, North Carolina 27708, USA
[11]Institute of Primate Research, National Museums of Kenya, Nairobi 00502, Kenya
[12]Duke Population Research Institute, Duke University, Durham, NC 27708, USA
*Correspondence: racheljohnston7@gmail.com, jt5@duke.edu

Abstract
In some mammals and many social insects, highly cooperative societies are characterized by
reproductive division of labor, in which breeders and nonbreeders become behaviorally and
morphologically distinct. While differences in behavior and growth between breeders and
nonbreeders have been extensively described, little is known of their molecular underpinnings.
Here, we investigate the consequences of breeding for skeletal morphology and gene regulation
in highly cooperative Damaraland mole-rats. By experimentally assigning dominant breeding
'queen' status versus nonbreeder status to paired littermates, we confirm that queens experience
vertebral growth that likely confers advantages to fecundity. However, they also up-regulate
bone resorption pathways and show reductions in femoral mass, which predicts increased
vulnerability to fracture. Together, our results show that, as in eusocial insects, reproductive
division of labor in mole-rats leads to gene regulatory rewiring and extensive morphological
plasticity. However, in mole-rats, concentrated reproduction is also accompanied by costs to
bone strength.

Introduction
A hallmark of highly cooperative societies is reproductive division of labor. This
phenomenon is best understood in eusocial insects, where environmental cues lead to
reproductively and morphologically specialized castes, including one or few highly fecund
"queens"[1]. These changes help support the reproductive role of queens by differentiating them
from nonbreeding colony members, who forage, care for young, and engage in colony defense[1,2].
Queens are frequently much larger than their sterile colony mates (e.g., twice as large in honey
bees and Pharaoh ants[3,4]), reflecting dramatically altered growth and development programs that

are explained by changes in gene regulation[5]. Social insects thus exemplify the link between
reproductive division of labor, cooperative behavior, and extreme morphological plasticity.
Systems in which breeding is restricted to a single female supported by multiple
nonbreeding helpers are also observed in vertebrates, including birds and mammals[6]. Here,
breeding status is not determined during early development, but instead occurs in adulthood, for
those individuals who have the opportunity to transition into a reproductive role. In some
species, new breeders undergo a period of accelerated growth, which may be important either for
maintaining dominance or for supporting high fecundity[7-12]. While substantial gene regulatory
divergence with breeding status has been described for the brain and some peripheral organs[13-15],
we know little about the gene regulatory shifts responsible for breeder-associated growth
patterns. These processes are key to understanding both the basis for, and limits of, status-driven
differences in growth and development.
Here, we investigate the morphological and molecular consequences of experimental
transitions to breeding status in female Damaraland mole-rats (*Fukomys damarensis*). Like naked
mole-rats, Damaraland mole-rats are frequently classified as 'eusocial'[16-18], and female helpers
who transition to queens experience accelerated vertebral growth associated with increases in
fecundity[9,11]. However, no study has yet profiled gene regulation in mole-rat breeders and
helpers in skeletal tissue, investigated their consequences for skeletal growth potential and
integrity, or determined whether they extend beyond the vertebrae. To do so here, we randomly
assigned littermate sisters in captive colonies to become queens or remain as nonbreeders and
evaluated gene regulatory and morphological changes induced by the transition to queen status.
Our results indicate that queens not only experience lengthening of their lumbar vertebrae, but
also show reductions in the growth potential and structural integrity of their long bones. These
changes result from increased rates of bone resorption that may increase the risk of fracture,
indicating that the presence of helpers does not annul the costs of reproduction to queens. As in
eusocial insects, females that acquire breeding status experience substantial morphological
remodeling, associated with pathway-specific changes in gene regulation.
74
Results
Adaptive plasticity in the skeleton of Damaraland mole-rat queens
We randomly assigned female Damaraland mole-rats (16 sets of littermates; mean age =
19.0 ± 3.8 s.d. months at assignment; Supplementary Table 1) to either transition to queen status
in a newly founded colony (n = 12) or remain as nonbreeders for the duration of the experiment
(n = 18) (Figure 1A). Nonbreeders were either kept in their natal colonies as helpers or placed
into solitary housing, which recapitulates extended periods of dispersal in this species[18] (n = 10
helpers and n = 8 solitaires). Six non-experimental animals (1 queen and 5 nonbreeders) were
also included in the sample, resulting in a total sample size of 13 breeders and 23 nonbreeders
(Supplementary Table 1).
Females assigned to the queen treatment were each transferred to a new tunnel system
containing only an unrelated adult male, simulating the natural process of dispersal and new
colony formation in the wild[18]. Pairing with an unrelated male leads to rapid activation of the
reproductive axis, including initiation of ovulation, sexual activity, and the potential for
conception[19,20]. At the start of experimental treatment, animals assigned to the queen treatment
and to the nonbreeding treatment did not significantly differ in body mass (unpaired t-test, t =
0.894, df = 16.828, p = 0.384; means = 116.3 and 110.9 grams), age (unpaired t-test, t = 0.746, df
= 22.137, p = 0.466; means = 628 and 580 days), or vertebral length (as measured by lumbar

vertebra 5 [LV5] following [11]; unpaired t-test, t = -1.053, df = 20.131, p = 0.305; means = 5.0
and 5.13 mm). Queens gave birth to a mean of 6.92 ± 5.57 s.d. live offspring during the 12 – 22
month follow up period, produced in a mean of 2.85 ± 1.75 s.d. litters (Supplementary Table 1).

Figure 1. Transition to queen status leads to lumbar vertebral lengthening. (a) Experimental design:
nonbreeding adult female (♀) littermates were randomly assigned to transition to queen status (purple ⚨) by being
paired with an unrelated male (♂), or to remain in a nonbreeding treatment (cyan). Duration of treatment ranged
from 12 – 22 months. **(b)** Queens show more rapid growth in lumbar vertebra 5 (LV5) in the first four months of the
experiment, relative to nonbreeders (treatment by time point interaction: β = 0.078, n = 49, p = 3.47 x 10^{-3}). Dots
show means +/- standard errors (bars). **(c)** At the start of the experiment (0 months, left panel), the lumbar vertebrae
of breeders do not differ from those of nonbreeders (unpaired t-tests, all p > 0.05). However, at 12 months (right
panel), queens have longer lumbar vertebrae relative to nonbreeders (unpaired t-tests, * indicates p < 0.05). Dots
show means +/- standard errors (bars). Lengths of lumbar vertebrae above the plots are scaled to indicate the mean
lengths of queens (top) and nonbreeders (bottom) at each time point; vertebrae highlighted in purple are significantly
longer in queens relative to nonbreeders. **(d)** Litter size is positively correlated with maternal body length in the
Damaraland mole-rat colony (β = 0.353, n = 328 litters, p = 1.35 x 10^{-3}).

As previously described[11], queens showed rapid growth in the lumbar vertebrae in the
first 12 months post-pairing, relative to nonbreeders (Figure 1B). In contrast to naked mole-rats,
where the lumbar vertebrae at the cranial end exhibit the most growth in queens[21], the lumbar
vertebrae toward the caudal end (LV5 and LV6) were most responsive to breeding status in our
experiment, suggesting potential convergent evolution of the vertebral lengthening phenotype.
Based on longitudinal measurements, most of this differential growth was concentrated soon
after the breeding status transition, concordant with findings in naked mole-rats[22]. Specifically,
we observed a significant interaction between breeding status (queen versus nonbreeder) and
post-pairing time point in the first four months of the experiment (Figure 1B; β = 0.0784, p =
3.47 x 10^{-3}; n = 49 x-rays from 28 animals), but not for x-rays taken in later time point intervals
(4 months versus 8 months; 8 versus 12 months, all p > 0.05). However, due to the early
increased growth in queens, size differences persisted, such that the absolute length of LV5 in
queens was, on average, 4.8% longer than nonbreeders after 12 months (Figure 1C; LV5:
unpaired t-test, t = 2.509, df = 21.095, p = 0.020), and the absolute length of the lumbar vertebral
column in queens relative to nonbreeders was 3.5% longer, although the latter difference was not
significant (unpaired t-test, t = 1.945, df = 22.49, p = 0.064). Differences between queens and
nonbreeders were even more apparent if lumbar vertebrae measures were scaled by zygomatic

arch (head) width, as done in previous studies[9,11,22] (LV5: 9.3% longer, unpaired t-test, t = 4.12,
df = 15.135, p = 8.87 x 10^{-4}; lumbar vertebral column length: 7.9% longer, unpaired t-test, t =
4.34, df = 15.37, p = 5.58 x 10^{-4}). This difference highlights changes in allometry across the
queen skeleton due to the combination of lumbar vertebral growth and narrower zygomatic arch
width in queens (4% narrower at 12 months, unpaired t-test, t = -2.448, df = 24.126, p = 0.022).
Consistent with previous results[11], queens in the colony with longer bodies (which
correlates with longer lumbar vertebrae, Pearson's r = 0.856, p = 5.99 x 10^{-59}; Supplementary
Figure 1) had more pups per litter (Figure 1D; Supplementary Table 2; β = 0.353, p = 1.35 x 10^{-3},
n = 328 litters from all breeding groups maintained in the same breeding facility). Controlling for
litter size, longer queens also had larger pups: for every additional centimeter of maternal body
length, pup body mass increased by 2.9% (β = 0.28, p = 0.032, n = 971 pups). Thus, the
elongated queen phenotype is a strong candidate for adaptive plasticity that supports increased
fertility in queen mole-rats.
139
Breeding status induces gene regulatory changes in the queen mole-rat skeleton
To identify the gene regulatory changes associated with skeletal plasticity, we cultured
cells enriched for bone marrow-derived mesenchymal stromal cells (bMSCs) isolated from the
lumbar vertebrae (pooled LV1-LV5) of both queens and nonbreeders (n = 5 queens, 11
nonbreeders). bMSC cultures include multipotent skeletal stem cells, the precursor of the
osteoblast and chondrocyte lineages responsible for bone growth. In parallel, we cultured cells
enriched for bMSCs from the pooled long bones (humerus, ulna, radius, left femur, and left tibia)
of the same animals, which do not show increased elongation in queens (for both tibia length and
femur length, all p > 0.05 for interactions between breeding status and time intervals [0 months
versus 4 months, 4 months versus 8 months, 8 months versus 12 months]; femur at 12 months:
unpaired t-test, t = -0.202, df = 19.326, p = 0.842; tibia at 12 months: unpaired t-test, t = -0.860,
df = 16.759, p = 0.402). To evaluate the potential role of sex steroid hormone signaling on bone
growth, we treated cells from each bone sample for 24 hours with either 10 nM estradiol or
vehicle control, resulting in 47 total samples. We then performed RNA-seq on each sample to
screen for genes that were systematically differentially expressed in the bone cells of queens
versus nonbreeders.
Of 10,817 detectably expressed genes, 171 genes showed a significant effect of breeding
status at a false discovery rate (FDR) threshold of 10% in the long bones (329 at an FDR of 20%;
Supplementary Table 3). Surprisingly, no genes showed a significant effect of breeding status in
the lumbar vertebrae at either FDR threshold. However, effect sizes were highly correlated
between bone types overall (R^2 = 0.75, p = 4.60 x 10^{-53}), with more pronounced effects of
breeding status in the long bone samples than in the lumbar vertebrae (paired t-test on breeding
status effects in long bone versus vertebrae: t = 3.97, df = 317.67, p = 8.73 x 10^{-5}). Importantly,
breeding status-related differences were not readily attributable to differences in bone cell
composition. Based on both canonical markers of bMSC lineage cells and deconvolution of the
RNA-seq data using data from 27 mesenchymal or hematopoietic lineage mouse cell types, the
majority cell type in both queen and nonbreeder samples was most similar to cells from the
bMSC lineage[23-25] (Supplementary Figures 2 and 3). Additionally, estimated cell type
proportions (as summarized by the top three principal components) did not differ between
queens and nonbreeders (all FDR > 0.1, Supplementary Table 4), and we identified no cases in
which the effects of breeding status on gene expression were significantly mediated by the first
principal component of cell composition (p > 0.05 for all 171 queen-associated genes at 10%

FDR; Supplementary Table 5). Because we observed no significant effect of estradiol treatment
on gene expression in either bone type (all FDR > 10%), we did not evaluate the effects of
estradiol further.
The majority of breeding status-associated genes were up-regulated in queens (151 of 171
genes, 88%). In support of their role in skeletal plasticity, up-regulated genes were enriched for
"bone remodeling" ($\log_2[OR] = 4.07$, $p = 5.07$ x 10^{-6}), a process that involves the balanced cycle
between bone formation by osteoblasts and bone resorption by osteoclasts[26] (Figure 2).
Surprisingly, however, enriched terms were specifically concentrated in pathways and processes
related to bone resorption, not formation (Supplementary Table 6), including "positive regulation
of bone resorption" (Figure 2A, C; $\log_2[OR] = 6.51$, $p = 1.55$ x 10^{-6}) and "superoxide anion
generation" (Figure 2A, C; $\log_2[OR] = 5.29$, $p = 1.4$ x 10^{-5}), which is involved in osteoclast
activity and degradation of bone matrix[27-30]. Differentially expressed genes were also enriched
for immune-related processes (e.g., "cytokine secretion", "chemotaxis", "leukocyte activation
involved in immune response"; Supplementary Table 6). These observations suggest that
transitions to queen status also involve changes in immunoregulatory signaling (osteoclast cells,
like macrophages, are derived from monocytes). Omni-ATAC-seq profiling of open chromatin
regions further supports a central role for bone resorption and osteoclast activity in the queen
skeleton (n = 8; Supplementary Table 7). Specifically, transcription factor binding motifs
(TFBMs) located in accessible chromatin near queen up-regulated genes were enriched for PU.1
and MITF, two transcription factors that are essential for osteoclast differentiation[31] (Figure 2B,
C; PU.1 $\log_2[OR] = 1.041$, $p = 2.84$ x 10^{-4}; MITF $\log_2[OR] = 0.707$, $p = 7.36$ x 10^{-3}; see
Supplementary Table 7 for complete list of enriched TFBMs). *MITF* was also among the 151
genes that were differentially expressed between queens and nonbreeders and up-regulated in
both queen long bones and lumbar vertebrae.

Figure 2. Queen status drives increased regulatory activity of bone resorption pathways. (a) Gene Ontology (GO) terms enriched in queen up-regulated genes, relative to the background set of all genes tested. Bars represent 95% confidence intervals. Processes highlighted in purple are also depicted in (c). Highest-level (most general) terms are shown; for full GO enrichment results, see Supplementary Table 6. **(b)** Accessible transcription factor binding site motifs enriched near queen up-regulated genes, relative to all genes tested. Bars represent 95% confidence intervals. Transcription factors highlighted in purple are also depicted in (c). **(c)** Schematic of the balance between bone formation and bone resorption, showing key regulators and markers for mesenchymal stem cell differentiation into osteoblasts and hematopoietic stem cell differentiation into osteoclasts[26,31]. Queen up-regulated genes or proteins are highlighted as purple ovals (FDR < 10%) and transcription factors with binding motifs enriched near queen up-regulated genes are highlighted as purple hexagons. Inset for osteoclasts shows the NADPH oxidase system, which generates superoxide radicals (O_2^-) necessary for bone resorption and is highly enriched for queen up-regulated genes (purple ovals).

Extensive skeletal remodeling in queen mole-rats

The gene expression data suggest that queen status-driven changes to the skeleton extend beyond the lumbar vertebrae to the long bones. Further, they suggest that bone resorption—an important counterpoint to bone formation that is required for normal skeletal maintenance—also distinguishes breeding and nonbreeding females. To investigate this possibility, we performed high-resolution micro-computed tomography (μCT) scanning to generate 3D reconstructions of LV6, LV7, right femur, and right tibia of queens and female nonbreeders (n = 140 bones from 36 animals; Figure 3A; Supplementary Figure 4). This approach substantially increases the level of resolution for investigating breeding status-linked differences in skeletal morphology, as previous studies relied on x-ray data alone[9,11].

We first asked whether breeding status could be predicted from morphological differences in the 3D reconstructions. We found that it could for the lumbar vertebra, but not for the femur: by applying the smooth Euler characteristic transform[32], we were able to predict queen versus nonbreeder status in LV6 (77.8% accuracy, p = 0.01, n = 36), but not the femur (52.8% accuracy, p = 0.53, n = 36). Including only highly fecund queens (≥6 total offspring)

improved predictive accuracy in the femur (70% accuracy, p = 0.12, n = 30). Although these
predictions did not reach statistical significance, they raised the possibility that morphological
changes in femurs become enhanced with increasing reproductive effort.
We next tested whether the transition to queen status affects the ability to continue bone
lengthening. Lengthening requires the presence of a growth plate, a region of cartilage in the
bone where longitudinal growth occurs through proliferation of cartilage cells (chondrocytes)
(Figure 3A-B; Supplementary Figure 4). Closure of the growth plate, which indicates that bone
lengthening potential has terminated, typically occurs in mammals after reaching sexual
maturation, when energy begins to be invested in reproduction instead of growth[33]. To test
whether the transition to queen status alters bone lengthening potential, we performed Safranin-
O staining on sections of the right tibia and LV7 to visualize growth plates (Figure 3B). In the
proximal tibia but not LV7, queens were less likely to have open growth plates (Figure 3C;
Supplementary Figure 4; Supplementary table 8; tibia: two-sided binomial test, p = 0.019; LV7:
two-sided binomial test, p = 0.422). The increased probability of growth plate closure in the tibia
of queens is linked to the number of offspring a female has produced: females with more
offspring showed a higher expanse of closure across the growth plate ($\beta = 0.050$, p = 4.51×10^{-3},
n = 12, controlling for age). This pattern may be due in part to reduced chondrocyte proliferation,
as females that produced more offspring had fewer chondrocyte columns in the remaining
growth plate (Figure 3D; $\beta = -0.132$, p = 0.020, n = 12, controlling for age). Thus, offspring
production in queens is associated with loss of the ability to lengthen the long bones, but not the
lumbar vertebrae, consistent with the importance of abdominal lengthening for supporting larger
litters.

Figure 3. Queen status reduces growth potential in the tibia. (a) μCT scans of female Damaraland mole-rat tibias. Boxes
indicate the location of the proximal growth plate, which is either unfused (left) or fully fused (right). **(b)** Example Safranin-O
stained histological sections of the proximal tibia, in which the growth plate is unfused (top), partially fused (middle), or fully
fused (bottom). Values in the top left of each box indicate percent growth plate fusion across the width of the bone. The
cartilaginous growth plate is stained deep pink and calcified bone is stained green. **(c)** Queens, and specifically queens that gave
birth to more offspring, show increased growth plate fusion ($\beta = 0.050$, p = 4.51×10^{-3}, n = 12, controlling for age) and **(d)**
decreased number of chondrocyte columns within the remaining growth plate ($\beta = -0.132$, p = 0.020, n = 12, controlling for age).
Each box represents the interquartile range, with the median value depicted as a horizontal bar. Whiskers extend to the most
extreme values within 1.5x of the interquartile range. In (c) and (d), dots represent individual animals, and shading indicates each
animal's total offspring number. Ages of queens and nonbreeders do not significantly differ (unpaired t-test, t = 0.489, n = 12, p =
0.644).

A major demand on reproductively active female mammals is a high requirement for
calcium, particularly during lactation when mothers support rapid offspring bone growth.
Maternal skeletons are remodeled to meet this demand, although in most mammals, these
changes are not permanent (reviewed in [34]). Because of the particularly intense reproductive
investment made by cooperatively breeding mole-rat queens, we therefore also evaluated the
effect of queen status on trabecular and cortical bone volumes, which are thought to be important
in satisfying short-term and long-term calcium demands, respectively. We found no effect of
queen status on the amount of trabecular bone in the femur, tibia, LV6, or LV7 (all p > 0.05 for
bone volume/total volume). However, we found that cortical bone was significantly thinner at
the femoral midshaft, but not in the lumbar vertebrae, in queens compared to their nonbreeding
sisters (Figure 4A; Supplementary Figure 5; femur: paired t-test of cortical area/total area, t = -
4.067, df = 8, p = 3.60 x 10^{-3}; LV6: paired t-test of cortical area/total area, t = -0.741, df = 6, p =
0.487). Cortical thinning in queens appears to be specifically due to increased bone resorption,
which typically occurs on the endosteal (internal) surface of long bones in the marrow cavity.
Indeed, queens had a larger marrow cavity (paired t-test, t = 5.355, df = 8, p = 6.82 x 10^{-4}; Figure
4B) but showed no difference in periosteal area compared to their nonbreeding sisters (paired t-
test, t = 1.539, df = 8, p = 0.162; Figure 4C).

Figure 4. Queens exhibit reduced cortical area and increased marrow area at the femoral midshaft. Femoral cross-sections
with area highlighted in gray show the measures represented in the corresponding plots below. Each line represents an age-
matched, nonbreeder and queen littermate pair. **(a)** Queens have less cortical bone (relative to the total area of the femoral
midshaft cross-section), compared to their paired nonbreeding littermates (paired t-test of cortical area/total area, t = -4.07, df = 8,
p = 3.60 x 10^{-3}). **(b)** Queens also have enlarged marrow cavities (paired t-test, t = 5.36, df = 8, p = 6.82 x 10^{-4}), where bone
resorption typically occurs, but **(c)** show no difference in overall periosteal area (paired t-test, t = 1.54, df = 8, p = 0.162).
280

Because changes in cortical bone are thought to reflect accumulated demands over long
time frames, we hypothesized that cortical thinning in queens is a consequence of repeated
cycles of pregnancy and lactation over time, which can occur simultaneously in Damaraland
mole-rat queens. In support of this idea, we found that, within queens, the relative amount of
cortical bone is not predicted by the number of pups in a queen's most recent litter (β = -0.024, n
= 13, p = 0.287), but instead by the total number of pups she produced in her lifetime.
Specifically, queens who had more live births had reduced cortical bone thickness along the
entire shaft of the femur (Figure 5 and Supplementary Table 9; across decile sections of the
femur: all p < 0.05, controlling for mother's litter as a random effect; at midsection: β = -0.011, n
= 13, p = 0.031). Thus, cortical thinning does not commence with the transition to queen status

per se (i.e., it is not a correlate of *achieving* breeder status), but instead appears to be a
consequence of repeated investment in pregnancy and lactation. Notably, thinning is particularly
marked in queens who had at least six offspring, which usually occurs by 14 months after a
breeding status transition (i.e., by the third litter; Supplementary Table 9). Given that wild
Damaraland mole-rat queens can maintain their status for many years[35], our results suggest that
long-lived queens may experience substantial morphological change.

Figure 5. Offspring production in queens leads to cortical thinning across the femoral shaft. (a) Queens (top row) relative to
their same-aged nonbreeding littermates (bottom row) present thinner cortical bone across the femur, particularly in females that
have many offspring (top right). Number of offspring is indicated above each femur, and vertical arrows indicate littermate pairs.
(b) Within each decile section across the femoral shaft, number of offspring is negatively correlated with average cortical
thickness (linear mixed model with littermate pair as random effect). Full results are presented in Supplementary Table 9.
Asterisk indicates $p < 0.05$.

Skeletal remodeling predicts increased risk of femur breakage in queens

In humans, accelerated bone resorption is a central cause of osteoporosis-related bone
fragility[36]. We therefore hypothesized that cortical thinning in queen mole-rat femurs would be
linked to decreased bone strength. To test this hypothesis, we calculated two key indicators of
femoral structural integrity: cortical area (CA) and the minimum moment of inertia (I_{min}, a
predictor of resistance to bending). In nonbreeders, both are positively correlated with body mass
(I_{min}: $R^2 = 0.368$, n = 24, p = 9.92 x 10^{-4}; CA: $R^2 = 0.407$, n = 24, p = 4.80 x 10^{-4}). However, in
queens, I_{min} is not significantly predicted by body mass ($R^2 = 0.088$, n = 13, p = 0.17), but is
instead a function of number of offspring produced ($R^2 = 0.283$, p = 0.035). Queen CA is
predicted by both offspring number and body mass, but offspring number explains almost twice
the variance (offspring number $R^2 = 0.634$, p = 6.83 x 10^{-4}; body mass $R^2 = 0.385$, p = 0.014).

To evaluate the effects of reproductive activity on the risk of bone failure, we drew on
data on the relationship between CA and bone mechanical failure in a large data set of mouse
femurs[37]. In this data set, CA is the best predictor of maximum load (the maximum force a bone
can withstand prior to failure), and, crucially, the CA-max load relationship is highly linear
(Supplementary Figure 6; $R^2 = 0.88$, $p = 6.64 \times 10^{-38}$). Scaling the mole-rat CA data to mouse
suggests that transitions to queen significantly increase the risk of bone failure (Figure 6A;
hazard ratio (95% confidence interval) = 2.67 (1.20, 5.93), n = 36, p = 0.016). Similar to growth
potential and cortical thinning, this effect is driven by highly fertile queens, such that those who
had at least six offspring showed the highest predicted risk of bone failure (Figure 6B; queens
with ≥ 6 offspring relative to nonbreeders: HR = 3.81 (1.47, 9.83), n = 30, p = 0.006). The risk of
bone failure is thus predicted to increase by 21% for each additional pup (HR = 1.21 (1.10, 1.33),
n = 36, $p = 8.85 \times 10^{-5}$).

Figure 6. Effect of reproductive status on the probability of bone failure. (a) Survival curves for femurs from queens versus
nonbreeders, based on predictions from the midshaft cortical area and data from [37] (Wald test, p = 0.02, n = 36; see Methods). **(b)**
Survival curves for femurs from queens with ≥ 6 offspring versus nonbreeders (Wald test, p = 0.006, n = 30). Vertical dashed
lines indicate group medians, with the median failure time for nonbreeders fixed at a value of 1.
331

Discussion

Our results demonstrate that transitions to breeding status in Damaraland mole-rat queens
drive widespread skeletal remodeling linked to shifts in gene regulation. In addition to previously
described vertebral growth[11], queen Damaraland mole-rats up-regulate bone resorption-related
pathways, lose bone lengthening potential in the long bones, and develop thinner femurs that are
predicted to be more prone to mechanical failure. While vertebral growth is triggered
immediately after transition to queen status, these other changes are a function of repeated
offspring production. Thus, queens quickly progress from traits typically associated with pre-

reproductive and pubertal growth in mammals (e.g., body elongation), to traits typically linked to
aging (e.g., marrow cavity expansion and cortical thinning). This progression is supported by the
signature of bone resorption rather than bone growth at the time of sampling, which occurred 1 –
2 years post-transition. Combined, these results suggest some key differences between
Damaraland mole-rats and eusocial insects. Specifically, while abdominal lengthening allows
queen mole-rats to increase fecundity per reproductive effort, loss of cortical bone in the femur is
unlikely to directly benefit either fertility or survival. Instead, it reflects the cumulative burden of
continuous cycles of pregnancy and lactation[34]. Thus, unlike eusocial insect queens[38,39],
Damaraland mole-rat queens incur morphological costs to concentrated reproduction in addition
to morphological changes that facilitate increased fitness.
The typical mammalian pattern of bone remodeling, in which bone mineral density
decreases during pregnancy and lactation but recovers after offspring are weaned[34], may explain
bone thinning in Damaraland mole-rats queens: queens often lactate for one litter while gestating
another, leaving little to no time for recovery. One possible reason that this fast rate of breeding
is achievable is that queens in colonies with more helpers work less and rest more[40], consistent
with studies in other cooperative mammals that show that helpers alleviate breeding-associated
loss of condition in queens[41]. However, the extent to which helpers reduce the costs of breeding
to queens may differ between species depending on the relative numbers of helpers to breeders.
For example, in eusocial insects, large colonies and the high ratio of helpers to queens reduce the
costs of reproduction to queens to low levels[1,2]. Similarly, in naked mole-rats (where colonies
can include hundreds of animals, compared to up to dozens in Damaraland mole-rat
colonies[17,18,43]), a small sample of queens (n = 6) suggests increased rather than decreased
femoral cortical thickness relative to age-matched nonbreeders[44]. Testing how the costs and
benefits of reproduction are resolved across different levels of cooperativity, including the
molecular mechanisms that mediate these differences, is an important next step towards
understanding the evolution of cooperative breeding in mammals.
366

Acknowledgements
We thank Tim Vink, Dave Gaynor, and the mole-rat house staff and volunteers for their
tremendous contributions to the Kalahari Mole-rat Project. We also thank Irene Garcia, Mari
Cobb, Brianna Bowman, Anna Luiza Wolf, Alice Zhou, Yilin Yu, B.J. Nielsen, and Tawni
Voyles for their contributions to sample collection and data generation, Graham Treece for
guidance on quantifying bone thickness with Stradview, Karl Jepsen for sharing data on mouse
femurs, Lou DeFrate for guidance on estimating bone strength, and members of the Tung lab for
feedback on earlier versions of this manuscript. Support for this work was provided by the
European Research Council (Grants 294494 and 742808 to T.C.B), the Human Frontier Science
Program (RGP0051-2017 to J.T., S.M., and T.C.B.), the National Science Foundation (IOS-
7801004 to J.T.), the National Institutes of Health (F32HD095616 to R.A.J.), a Sloan Foundation
Early Career Research Fellowship to J.T., a Foerster-Bernstein Postdoctoral Fellowship to
R.A.J., and a Natural Environmental Research Council Doctoral Training Program to Ja.T. High-
performance computing resources were supported by the North Carolina Biotechnology Center
(Grant Number 2016-IDG-1013).
382

383 **Author Contributions**
384 Conceptualization, R.A.J., T.C.B., and J.T.; Investigation, R.A.J., Ja.T., P.V., H.K., L.S., S.M.,
385 C.K., T.C.B., and J.T. Formal Analysis, R.A.J., H.K., S.M.; Writing—Original Draft, R.A.J. and

386 J.T.; Writing—Reviewing & Editing, R.A.J., P.V., Ja.T., H.K., L.S., S.M., C.K., T.C.B., and J.T.
387 Funding Acquisition, J.T. and T.C.B. Supervision, T.C.B. and J.T.
388
389 **Competing Interests**
390 The authors declare no competing interests.
391

Methods
Study system and experimental design

Damaraland mole-rats (*Fukomys damarensis*) were maintained in a captive colony at the Kuruman River Reserve in the Northern Cape Province of South Africa, within the species' natural range. Only animals born in captivity, with known birthdates and litter composition, were used in this study. Animals were maintained in artificial tunnel systems built from PVC pipes with compartments for a nest-box and waste-box and transparent windows to allow behavioral observation. Animals were fed *ad libitum* with sweet potatoes and cucumbers.

Adult females were randomly assigned to be either nonbreeders or queens, such that females assigned to queen status had matched siblings who were assigned to the nonbreeding condition (age-matched littermates where possible). Nonbreeders were either maintained in their natal colony as helpers or maintained alone, which models the social condition experienced by dispersing females. Females assigned to the breeder condition were transferred to a new tunnel system with an unrelated male from a separate social group. Ten new breeding colonies, with corresponding age-matched nonbreeding littermates where possible (6 helper and 8 solitaire age-matched littermates, and 4 non-littermate helpers), were set up in December 2015-June 2017[11]. With one exception (G10F026), animals maintained their breeding status for 14-22 months before sample collection. Breeding females were at different stages of pregnancy at time of sacrifice. An additional 4 breeding colonies (with 4 matched littermate helpers) were formed in October 2017 and followed for 11-12 months (Supplementary Table 1). One queen died before sample collection, and one non-experimental helper was euthanized during the course of the study and included in sample collection. The final sample size included 13 breeders, 15 helpers, and 8 solitaire females.

X-ray data

For a subset of study subjects, full body X-rays were taken every two months during the first 12 months of the experiment and at the time of sacrifice. From these X-rays, an experimenter blind to animal breeding status measured the length of each lumbar vertebra from vertebra 1 to 7, the right femur, the right tibia, body length, and the width of the zygomatic arch using ImageJ[45]. The caudal-most lumbar vertebra was labeled as LV7. We tested for an effect of breeding status on LV5 using a linear mixed model in which post-pairing time point, breeding status, and the interaction of time point by breeding status were modeled as fixed effects and animal ID as a random effect.

Effects of queen body length on fertility

To test the effect of maternal body length on litter size and pup size, we used body length measurements obtained during routine colony monitoring of all queens maintained in the captive colony (i.e., not restricted to experimental animals). Following [11], we used body length measurements obtained nearest to, and no more than 90 days from, the date of parturition. The resulting dataset included 328 litters (971 pups) from 76 mothers (a 76% increase from the data

set in Thorley *et al*, 2018). We fit two linear mixed effects models. In the first model, we
modeled litter size as a function of maternal body length, controlling for whether the litter was
the female's first litter, and included maternal ID as a random effect. In the second model, we
modeled pup mass as a function of maternal body length, controlling for litter size and whether
the litter was the female's first litter as fixed effects, and maternal ID and litter ID as random
effects.
438
Sample collection and cell culture from lumbar vertebrae and long bones
Animals were deeply anesthetized with isoflurane and sacrificed with decapitation
following USGS National Wildlife Health Center guidelines and under approval from the
Animal Ethics Committee of the University of Pretoria. Immediately upon sacrifice, the lumbar
vertebrae and long bones were dissected, and attached muscle tissue was removed with forceps.
Lumbar vertebrae 6 and 7 and the right femur and tibia were collected into 50% ethanol for 24
hours, then transferred to 70% ethanol and stored at 4° C for µCT scans and histochemistry.
To isolate bone cells for culture, lumbar vertebrae 1-5 were incubated in 2% Collagenase
P (Roche, Switzerland) for 30 minutes at 30° C. Each bone was then cut in half and transferred
to a 1.5 ml microcentrifuge tube containing a G-Tube microcentrifuge tube (VWR, Radnor, PA,
USA) that had been punctured at the bottom with a 15 gauge needle. Tubes were spun at 3,000
RCF for 5 seconds, allowing the marrow to collect into the 1.5 mL microcentrifuge tube. Cell
pellets were resuspended in red blood cell lysis buffer, pooled, and incubated for 3 minutes at
room temperature. 10 ml bMSC media (MEM-alpha [ThermoFisher, Waltham, MA, USA] +
15% fetal bovine serum [Hyclone, Logan, UT, USA] + 1% penicillin/streptomycin + 2 ng/ml
recombinant human fibroblast growth factor-basic [Biocam, Centurion, Gauteng, South Africa] +
10 nM ROCK inhibitor Y-27632 [RI; Cayman Chemical, Ann Arbor, MI, USA]) was added to
stop lysis, and the tubes were spun for 5 minutes at 300 RCF. The cell pellet was resuspended in
1 ml bMSC media and strained through a 70 µm cell strainer. Cells were plated at 1.6×10^5 cells
per cm^2. The long bones (excluding right femur and tibia) were processed to enrich for bMSCs
following the same procedure, but without incubation in Collagenase P. Cells were cultured at
37° C and 5% CO_2. Twenty-four hours post plating, plates were carefully washed three times
with 1x PBS and supplied with fresh media to remove non-adherent cells. Once bMSC clusters
were visible (2-9 days post plating), plates were fed bMSC media without RI or fed bMSC media
without RI + 10 nM estradiol (E2). Twenty-four hours later, cells were collected into buffer RLT
and stored at -80° C. Samples were shipped on dry ice to Duke University for RNA extraction
using the Qiagen RNeasy Micro Kit. RNA-Seq libraries were generated using the NEBNext
Single Cell/Low Input RNA Library Prep Kit for Illumina.
467
Gene expression analysis
RNA-Seq libraries were sequenced on an Illumina HiSeq 4000 (100 base pair single end
reads) to a mean coverage of 16.1 ± 3.9 s.d. million reads. Reads were trimmed with *cutadapt*
version 2.3[46] (parameters: -q 20 -e 0.2 --times 5 --overlap 2 -a AGATCGGAAGAGC -a "T" --
minimum-length=20). Trimmed reads were then mapped to the Damaraland mole-rat v1.0
genome[47] (DMR_v1.0) using two pass mapping with STAR[48]. Only uniquely mapped reads were
retained. HTseq[49] was used to quantify read counts mapping to genes (using the v1.0.92 gtf file
from Ensembl; we extended the genomic coordinates of the *SERPINE1* gene by 2000 basepairs
in both directions due to very high expression directly adjacent to the annotated coordinates). We
transformed read counts to transcripts per million (TPM)[50], and retained only genes with TPM $\geq$

2 in at least 25% of samples. We performed voom normalization[51] on the raw counts, using
normalization factors produced by the trimmed mean of M-values (TMM) method[52] in DESeq[53].
We used the *limma*[54] function *lmFit* to regress out the proportion of uniquely mapped reads in
genes (which controls for efficiency of mRNA selection during RNA-Seq library preparation)
and animal natal colony (which controls for littermate sets and date of sacrifice) to obtain
normalized, batch-corrected gene expression values for downstream analysis. We used the mixed
effects model approach in *emmreml*[55] to estimate, for each gene, the effect of breeding status on
gene expression within lumbar vertebrae and within long bones using the following model:
486

$$y_i = \mu + d_i\beta_1 + b_i\beta_2 + q_i\beta_3 * I(b = 0) + q_i\beta_4 * I(b = 1) + s_i\beta_5 * I(b = 0) + s\beta_6 * I(b = 1) + Zu + \varepsilon_i,$$

487                                  $$u \sim MVN\ 0, \sigma_u^2 K,$$
                                     $$\varepsilon \sim MVN\ 0, \sigma_e^2$$

488
where *y* is the vector of gene expression levels for $n = 47$ samples (indexed by *i*); μ is the
intercept; *d* is the number of days in culture and β_1 its effect size; *b* is bone type (i.e., long bones
or lumbar vertebrae) and β_2 its effect size; and *q* is a 0/1 variable representing breeder status and
β_3 and β_4 its effect size in long bones and lumbar vertebrae, respectively. *I* is an indicator
variable for bone type (0 = long bone; 1 = lumbar vertebrae). *s* is a 0/1 variable representing
whether the cells were cultured with estradiol and β_5 and β_6 are its effect sizes in long bones and
lumbar vertebrae, respectively. Z is an incidence matrix that maps samples to animal ID to take
into account repeated sampling from the same individual, and *u* is a random effect term that
controls for relatedness. *K* is an *m* by *m* matrix of pairwise relatedness estimates (derived from
the genotype data described below) between all *m* animals. ε is the residual error, σ_u^2 is the
genetic variance component, and σ_e^2 is the environmental variance component. To control for
multiple testing, we calculated the false discovery rate following Storey and Tibshirani[56] using
an empirical null distribution derived from 100 permutations of each variable of interest.
We used g:profiler[57] to perform gene ontology enrichment analysis of the genes up-
regulated with queen status in lumbar vertebrae and long bones (151 of 171 genes significantly
associated with queen status at a 10% FDR). All genes in the original analysis set were used as
the background gene set. We set both the minimum size of the functional category and the
minimum size of the query/term intersection to 3. Finally, we retained categories that passed a
Bonferroni corrected p-value of 0.05.
508
Genotyping to estimate relatedness
To control for relatedness when modeling the gene expression data, we performed single
nucleotide polymorphism (SNP) genotyping of the RNA-Seq data using the Genome Analysis
Toolkit[58] (GATK). We used the SplitNCigarReads function on the trimmed, uniquely mapped
reads and performed GATK indel realignment. Base recalibration was performed by using all
SNPs with GQ ≥ 4 in an initial UnifiedGenotyper run on the full data set as a reference.
Genotypes were called on the recalibrated bam files using HaplotypeCaller. Variants were
filtered with the following GATK VariantFiltration parameters: QUAL < 100.0, QD < 2.0, MQ <
35, FS > 30, HaplotypeScore > 13, MQRankSum < -12.5, ReadPosRankSum < -8. Variants were
further filtered with *vcftools*[59] to only retain biallelic SNPs in Hardy-Weinberg equilibrium (p >
0.05) with minor allele frequency ≥ 0.1, minimum mean depth of 5, max missing count of 2, and
minimum GQ of 99. Finally, SNPs were thinned to a distance of 10 kb basepairs, resulting in a

final dataset of 1,965 stringently filtered biallelic SNPs. Missing values were imputed using
beagle[60], and the resulting vcf file was used to create a kinship matrix using *vcftools*[59]. Values of
the kinship matrix were confirmed to be higher in known siblings compared to non-siblings
(unpaired t-test, t = 27.939, p = 2.23 x 10^{-12}; means = 0.513 and -0.097). Two pairs of siblings
were found to have different fathers (G1F022 and G1F025; G4F020 and G4F019).
526
Cell type heterogeneity
Although selection for adherent cells from bone marrow enriches for bMSCs, other cell
types are also present[61]. To assess whether cell type heterogeneity accounts for queen-associated
differential expression, we used CIBERSORT to deconvolve the proportion of component cell
types from the RNA-Seq data[25]. We trained CIBERSORT on a data set of quantile normalized
gene expression values from mouse purified primary cell populations[24]. Specifically, we subset
the training data to 27 purified cell populations of mesenchymal or hematopoietic origin
(Supplementary Figure 3) and to genes that were included in our mole-rat gene expression
dataset. We then predicted the composition of the cells that contributed to the mole-rat quantile
normalized gene expression data set, for each sample.
To test whether cell type heterogeneity was significantly explained by queen status, we
also modeled cell type proportion (as summarized by the first principal component of
CIBERSORT-estimated proportions for all 27 potential cell types; PC1 explains 50.9% of the
overall variation) following the same method used for gene-by-gene expression analysis but with
PC1 included as an explanatory variable. We then performed mediation analysis on each of the
171 genes that showed a significant effect of breeding status at FDR < 10%. To do so, we first
estimated the indirect effect of breeding status on gene expression through the mediating variable
(CIBERSORT PC1). The indirect effect of breeding status through CIBERSORT PC1 was
estimated by calculating the difference in the effect of breeding status between two models: one
model that did not include the mediator (i.e., β_3 and β_4 from equation 1 above) and the same
model with the addition of the mediating variable. We performed 1000 iterations of bootstrap
resampling to obtain the 95% confidence interval for the indirect effect, and considered an
indirect effect for a gene significant if the 95% interval did not overlap 0.
550
ATAC-seq data and transcription factor binding site analysis
To investigate whether differentially expressed genes were associated with accessible
binding motifs for specific transcription factors, we generated Omni-ATAC-seq data to profile
regions of open chromatin[62,63]. We performed Omni-ATAC-seq on both lumbar vertebrae
bMSCs and long bone bMSCs from two female nonbreeding and two queen mole-rats (n = 8
libraries total), following the published protocol[63] with the following modifications: 5,000 cells
were centrifuged at 500 RCF for 5 minutes at 4°C. The pellet was resuspended in 50 ul
transposition mix (25 ul 2xTD buffer, 16.5 ul PBS, 6.75 ul water, 1 ul 10% NP40, 1 ul 10%
Tween-20, 1 ul 1% digitonin, and 0.25 ul Tn5 transposase). The reaction was incubated at 37°C
for 30 minutes without mixing, followed by a 1.5x Ampure bead cleanup. Omni-ATAC libraries
were sequenced on a NovaSeq 6000 as 100 basepair paired-end reads to a mean coverage (± SD)
of 26.9 (±4.4) million reads (range: 16.8-38.3). Reads were trimmed with Trim Galore![64] to
remove adapter sequence and low quality basepairs (Phred score < 20; reads ≥ 25 bp). Read pairs
were mapped to the DMR_v1.0 genome using *bwa-mem*[65] with default settings. Only uniquely
mapped reads were retained. The alignment bam files for each treatment (breeding or
nonbreeding) were merged, and open chromatin regions were identified using MACS2 v2.1.2[66]

with the following parameters: "-nomodel -keep-dup all -q 0.05". We combined open chromatin
peaks with regions in the DMRv1.0 genome that match sequences of vertebrate transcription
factor binding site motifs, using motifs defined in the HOMER database[67]. We used Fisher's
exact tests (using a p-value threshold of 0.01) to test if transcription factor binding motifs
belonging to the same transcription factor were enriched in open chromatin regions within 2,000
bp of the 5' most transcription start site of queen up-regulated genes.
573
Micro-CT scans and analysis
We performed μCT scans of LV6, LV7, right femur, and right tibia using a VivaCT 80
scanner (Scanco Medical AG, Brüttisellen, Switzerland) set at 55 kVp and 145 μA, with voxel
size 10.4 μm. Trabecular bone was quantified from the 100 μCT slices below the proximal tibia
growth plate, the 100 μCT slices above the distal femur growth plate, and the 100 μCT slices
medial to the caudal growth plate of LV6. To obtain midshaft cross-sections of the femur, tibia,
and LV6, we first reduced each bone mesh to 100,000 faces using Avizo Lite version 9.7.0.
Mesh files from the same bone type were auto-aligned using Auto3dgm[68] in Matlab. Aligned
mesh files were then back scaled to their original sizes in Matlab, and the midshaft cross-section
was generated using Rhinoceros version 6. The MomentMacro plugin in ImageJ was used to
calculate bone area and moment of inertia.
585
Classification of breeding status from bone shape
To predict breeding status from bone shape, we first applied the smooth Euler
characteristic transform[32] to the aligned right femur and LV6. We then performed leave-one-out
predictions, running each bone type separately, using the linear kernel and c-classification with
the support vector machine (SVM) implemented by the R package *e1071*[69]. The SVM classifier
was equipped with 1:100,000 weighting to achieve class balanced predictions. The empirical p-
values were estimated for each bone type by running 100 permutations of the queen/nonbreeder
labels[70].
594
Histochemistry
For a subset of individuals (Supplementary Table 8), the tibia and LV7 were plasticized,
sectioned, and stained with Safranin O by the Washington University Musculoskeletal Research
Center. The proportion of the tibia proximal growth plate that was fused, and the mean
proportion of the LV7 cranial and caudal growth plates that were fused, were measured in
ImageJ from Safranin O stained sections. To quantify growth plate activity, we calculated the
number of chondrocyte columns (defined as linear stacks of at least three chondrocytes)
controlling for length of open growth plate. For each bone type (tibia and LV7), we ran two
models: proportion growth plate fusion or chondrocyte columns per mm growth plate as the
dependent variable, and number offspring born and age as the independent variables.
605
Cortical thickness across the femur
We used Stradview[71,72] on dicom images from the μCT scans to measure and visualize, in
an automated manner, cortical thickness across the surface of the femur. Bone surface was
defined in Stradview by thresholding pixel intensity and contouring the bone at every 14
sections, with the following parameters: resolution = medium, smoothing = standard, strength =
very low, contour accuracy = 6. To measure cortical thickness, we used the auto threshold
method in Stradview, with line width set to 5, smooth set to 1, and line length set to 3 mm. The

smoothed thickness values of each femur were then registered (i.e., mapped) to a single
"canonical" femur surface (mole-rat GRF002) using wxRegSurf v18
(http://mi.eng.cam.ac.uk/~ahg/wxRegSurf/). We sectioned the cortical thickness values into
deciles according to location along the length of the femur. The top and bottom deciles were
removed, because cortical and trabecular bone towards the ends of the femur could not be easily
differentiated by the automated method. Deciles were then recreated for the remaining length of
the bone (i.e., the central 80%). From each bone decile, we estimated cortical thickness as the
mean of all cortical thickness measures within that interval. For each decile across animals, we
used a linear mixed model to model cortical thickness as a function of breeding status and
number offspring, with litter pair as a random effect.
623
Modeling the probability of bone failure
Previous research on mechanical properties of rodent femurs found that, among several
morphological and compositional traits measured in eight morphologically varying mouse
strains, cortical area (CA) at the midshaft was the best predictor of maximum load (defined as
the greatest force attained prior to bone failure, measured via four-point bending; published
Pearson's $r = 0.95$)[37]. We therefore used cortical area at the femoral midshaft to predict max load
of Damaraland mole-rat femurs. To do so, we first fit a linear model of max load as a function of
cortical area (unadjusted for body weight) using published mouse data ($R^2 = 0.877$, n = 81, p =
6.64×10^{-38})[37]. We extrapolated from this linear fit to predict max load from cortical area at the
midshaft of Damaraland mole-rat femurs. Predicted max loads were then used as input for Cox
proportional hazards models, comparing either all queens to nonbreeders or queens with ≥ 6
offspring to nonbreeders. Models were fit using the R function *coxph*, and were confirmed to
meet the proportional hazards assumption using the *cox.zph* function in the R package *survival*[73].
Because max load was not directly measured in the Damaraland mole-rats, we used the Cox
proportional hazards models to specifically evaluate the relative hazard of bone failure
depending on queen status/number of offspring. We therefore report the results in Figure 6 based
on relative force (with the median predicted failure value for nonbreeders set to 1) instead of
absolute force in Newtons.
642
643   **Data Availability**
644   All RNA sequencing data generated during this study are available in the NCBI Gene Expression
645   Omnibus (series accession GSE152659). ATAC-Seq data are available in the NCBI Sequence
646   Read Archive (BioProject accession number PRJNA649596). µCT data from this study are
647   available on MorphoSource (http://www.morphosource.org, project 1056).
648

References

1   Wilson, E. O. The insect societies. *The insect societies.* (1971).

2   Keller, L. & Genoud, M. Extraordinary lifespans in ants: a test of evolutionary
theories of ageing. *Nature* **389**, 958-960 (1997).

3   Berndt, K. P. & Eichler, W. Die Pharaoameise, Monomorium pharaonis (L.)(Hym.,
Myrmicidae). *Mitteilungen aus dem Museum für Naturkunde in Berlin. Zoologisches
Museum und Institut für Spezielle Zoologie (Berlin)* **63**, 3-186 (1987).

4   Page Jr, R. E. & Peng, C. Y.-S. Aging and development in social insects with emphasis
on the honey bee, Apis mellifera L. *Experimental gerontology* **36**, 695-711 (2001).

5   Smith, C. R., Toth, A. L., Suarez, A. V. & Robinson, G. E. Genetic and genomic analyses
of the division of labour in insect societies. *Nature Reviews Genetics* **9**, 735-748
(2008).

6   Koenig, W. D. & Dickinson, J. L. *Cooperative breeding in vertebrates: studies of
ecology, evolution, and behavior.* (Cambridge University Press, 2016).

7   Clutton-Brock, T. *et al.* Intrasexual competition and sexual selection in cooperative
mammals. *Nature* **444**, 1065-1068 (2006).

8   Huchard, E., English, S., Bell, M. B., Thavarajah, N. & Clutton-Brock, T. Competitive
growth in a cooperative mammal. *Nature* **533**, 532-534 (2016).

9   O'Riain, M., Jarvis, J., Alexander, R., Buffenstein, R. & Peeters, C. Morphological castes
in a vertebrate. *Proceedings of the National Academy of Sciences* **97**, 13194-13197
(2000).

10   Russell, A. F., Carlson, A. A., McIlrath, G. M., Jordan, N. R. & Clutton‐Brock, T.
Adaptive size modification by dominant female meerkats. *Evolution* **58**, 1600-1607
(2004).

11   Thorley, J., Katlein, N., Goddard, K., Zöttl, M. & Clutton-Brock, T. Reproduction
triggers adaptive increases in body size in female mole-rats. *Proceedings of the Royal
Society B: Biological Sciences* **285**, 20180897 (2018).

12   Young, A. J. & Bennett, N. C. Morphological divergence of breeders and helpers in
wild Damaraland mole‐rat societies. *Evolution: International Journal of Organic
Evolution* **64**, 3190-3197 (2010).

13   Bens, M. *et al.* Naked mole-rat transcriptome signatures of socially suppressed
sexual maturation and links of reproduction to aging. *BMC biology* **16**, 77 (2018).

14   Mulugeta, E. *et al.* Molecular insights into the pathways underlying naked mole-rat
eusociality. *bioRxiv*, 209932 (2017).

15   Sahm, A. *et al.* Status-dependent aging rates in long-lived, social mole-rats are
shaped by HPA stress axis. *bioRxiv* (2020).

16   Bennett, N. C. & Faulkes, C. G. *African mole-rats: ecology and eusociality.* (Cambridge
University Press, 2000).

17   Jarvis, J. Eusociality in a mammal: cooperative breeding in naked mole-rat colonies.
Science **212**, 571-573 (1981).

18   Jarvis, J. & Bennett, N. Eusociality has evolved independently in two genera of
bathyergid mole-rats—but occurs in no other subterranean mammal. *Behavioral
Ecology and Sociobiology* **33**, 253-260 (1993).

19   Bennett, N., Faulkes, C. & Molteno, A. Reproductive suppression in subordinate, non-
breeding female Damaraland mole-rats: two components to a lifetime of socially

694   induced infertility. *Proceedings of the Royal Society of London. Series B: Biological*
695   *Sciences* **263**, 1599-1603 (1996).
696 20   Snyman, P., Jackson, C. R. & Bennett, N. C. Do dispersing non-reproductive female
697   Damaraland mole-rats, Cryptomys damarensis (Rodentia: Bathyergidae) exhibit
698   spontaneous or induced ovulation? *Physiology & behavior* **87**, 88-94 (2006).
699 21   Henry, E. C., Dengler-Crish, C. M. & Catania, K. C. Growing out of a caste-reproduction
700   and the making of the queen mole-rat. *Journal of Experimental Biology* **210**, 261-268
701   (2007).
702 22   Dengler‐Crish, C. M. & Catania, K. C. Cessation of reproduction‐related spine
703   elongation after multiple breeding cycles in female naked mole‐rats. *The*
704   *Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology:*
705   *Advances in Integrative Anatomy and Evolutionary Biology* **292**, 131-137 (2009).
706 23   Dominici, M. *et al.* Minimal criteria for defining multipotent mesenchymal stromal
707   cells. The International Society for Cellular Therapy position statement. *Cytotherapy*
708   **8**, 315-317 (2006).
709 24   Hume, D. A., Summers, K. M., Raza, S., Baillie, J. K. & Freeman, T. C. Functional
710   clustering and lineage markers: insights into cellular differentiation and gene
711   function from large-scale microarray studies of purified primary cell populations.
712   *Genomics* **95**, 328-338 (2010).
713 25   Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression
714   profiles. *Nature methods* **12**, 453 (2015).
715 26   Redlich, K. & Smolen, J. S. Inflammatory bone loss: pathogenesis and therapeutic
716   intervention. *Nature reviews Drug discovery* **11**, 234-250 (2012).
717 27   Darden, A. G., Ries, W. L., Wolf, W. C., Rodriguiz, R. M. & Key Jr, L. L. Osteoclastic
718   superoxide production and bone resorption: stimulation and inhibition by
719   modulators of NADPH oxidase. *Journal of Bone and Mineral Research* **11**, 671-675
720   (1996).
721 28   Datta, H., Rathod, H., Manning, P., Turnbull, Y. & McNeil, C. Parathyroid hormone
722   induces superoxide anion burst in the osteoclast: evidence for the direct
723   instantaneous activation of the osteoclast by the hormone. *Journal of endocrinology*
724   **149**, 269-275 (1996).
725 29   Key, L., Ries, W., Taylor, R., Hays, B. & Pitzer, B. Oxygen derived free radicals in
726   osteoclasts: the specificity and location of the nitroblue tetrazolium reaction. *Bone*
727   **11**, 115-119 (1990).
728 30   Key, L., Wolf, W., Gundberg, C. & Ries, W. Superoxide and bone resorption. *Bone* **15**,
729   431-436 (1994).
730 31   Segeletz, S. & Hoflack, B. Proteomic approaches to study osteoclast biology.
731   *Proteomics* **16**, 2545-2556 (2016).
732 32   Crawford, L., Monod, A., Chen, A. X., Mukherjee, S. & Rabadán, R. Functional data
733   analysis using a topological summary statistic: the smooth Euler characteristic
734   transform. *arXiv preprint arXiv:1611.06818* (2016).
735 33   Kilborn, S. H., Trudel, G. & Uhthoff, H. Review of growth plate closure compared with
736   age at sexual maturity and lifespan in laboratory animals. *Journal of the American*
737   *Association for Laboratory Animal Science* **41**, 21-26 (2002).

34   Kovacs, C. S. Maternal mineral and bone metabolism during pregnancy, lactation,
and post-weaning recovery. *Physiological reviews* **96**, 449-547 (2016).
35   Schmidt, C. M., Jarvis, J. U. & Bennett, N. C. The long-lived queen: reproduction and
longevity in female eusocial Damaraland mole-rats (Fukomys damarensis). *African*
Zoology **48**, 193-196 (2013).
36   Szulc, P., Seeman, E., Duboeuf, F., Sornay‐Rendu, E. & Delmas, P. D. Bone fragility:
failure of periosteal apposition to compensate for increased endocortical resorption
in postmenopausal women. *Journal of bone and mineral research* **21**, 1856-1863
(2006).
37   Jepsen, K. J., Akkus, O. J., Majeska, R. J. & Nadeau, J. H. Hierarchical relationship
between bone traits and mechanical properties in inbred mice. *Mammalian Genome*
14, 97-104 (2003).
38   Rodrigues, M. A. & Flatt, T. Endocrine uncoupling of the trade-off between
reproduction and somatic maintenance in eusocial insects. *Current opinion in insect*
science **16**, 1-8 (2016).
39   Rueppell, O., Aumer, D. & Moritz, R. F. Ties between ageing plasticity and
reproductive physiology in honey bees (Apis mellifera) reveal a positive relation
between fecundity and longevity as consequence of advanced social evolution.
Current opinion in insect science **16**, 64-68 (2016).
40   Houslay, T. M., Vullioud, P., Zöttl, M. & Clutton-Brock, T. H. Benefits of cooperation in
captive Damaraland mole-rats. *Behavioral Ecology* (2020).
41   Clutton-Brock, T. H. & Manser, M. Meerkats: cooperative breeding in the Kalahari.
Cooperative breeding in vertebrates **294**, 317 (2016).
42   Rosa, N., Simoes, R., Magalhães, F. D. & Marques, A. T. From mechanical stimulus to
bone formation: a review. *Medical engineering & physics* **37**, 719-728 (2015).
43   Jarvis, J. U., O'Riain, M. J., Bennett, N. C. & Sherman, P. W. Mammalian eusociality: a
family affair. *Trends in Ecology & Evolution* **9**, 47-51 (1994).
44   Pinto, M., Jepsen, K., Terranova, C. & Buffenstein, R. Lack of sexual dimorphism in
femora of the eusocial and hypogonadic naked mole-rat: a novel animal model for
the study of delayed puberty on the skeletal system. *Bone* **46**, 112-120 (2010).
45   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of
image analysis. *Nature methods* **9**, 671-675 (2012).
46   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet. journal* **17**, 10-12 (2011).
47   Fang, X. *et al.* Adaptations to a subterranean environment and longevity revealed by
the analysis of mole rat genomes. *Cell reports* **8**, 1354-1364 (2014).
48   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).
49   Anders, S., Pyl, P. T. & Huber, W. HTSeq–A Python framework to work with high-
throughput sequencing data. *Bioinformatics*, btu638 (2014).
50   Wagner, G. P., Kin, K. & Lynch, V. J. Measurement of mRNA abundance using RNA-seq
data: RPKM measure is inconsistent among samples. *Theory in biosciences* **131**, 281-
285 (2012).
51   Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. Voom: precision weights unlock linear
model analysis tools for RNA-seq read counts. *Genome biology* **15**, R29 (2014).

52   Robinson, M. D. & Oshlack, A. A scaling normalization method for differential
expression analysis of RNA-seq data. *Genome biology* **11**, R25 (2010).
53   Anders, S. & Huber, W. Differential expression analysis for sequence count data.
Genome Biol **11**, R106, doi:10.1186/gb-2010-11-10-r106 (2010).
54   Smyth, G. K. in *Bioinformatics and computational biology solutions using R and*
Bioconductor   397-420 (Springer, 2005).
55   Akdemir, D. & Okeke, U. EMMREML: Fitting mixed models with known covariance
structures. *R package version* **3** (2015).
56   Storey, J. D. & Tibshirani, R. Statistical significance for genomewide studies. *Proc*
Natl Acad Sci U S A **100**, 9440-9445, doi:10.1073/pnas.1530509100 (2003).
57   Raudvere, U. *et al.* g: Profiler: a web server for functional enrichment analysis and
conversions of gene lists (2019 update). *Nucleic acids research* **47**, W191-W198
(2019).
58   McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome research* **20**, 1297-1303
(2010).
59   Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-
2158 (2011).
60   Browning, S. R. & Browning, B. L. Rapid and accurate haplotype phasing and
missing-data inference for whole-genome association studies by use of localized
haplotype clustering. *The American Journal of Human Genetics* **81**, 1084-1097
(2007).
61   Phinney, D. G., Kopen, G., Isaacson, R. L. & Prockop, D. J. Plastic adherent stromal
cells from the bone marrow of commonly used strains of inbred mice: variations in
yield, growth, and differentiation. *Journal of cellular biochemistry* **72**, 570-585
(1999).
62   Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J. Transposition
of native chromatin for fast and sensitive epigenomic profiling of open chromatin,
DNA-binding proteins and nucleosome position. *Nature methods* **10**, 1213 (2013).
63   Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background and enables
interrogation of frozen tissues. *Nature methods* **14**, 959 (2017).
64   Krueger, F. Trim Galore.
http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/ (2015).
65   Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows–Wheeler
transform. *Bioinformatics* **26**, 589-595 (2010).
66   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome biology* **9**, R137
(2008).
67   Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors
prime cis-regulatory elements required for macrophage and B cell identities.
Molecular cell **38**, 576-589 (2010).
68   Boyer, D. M. *et al.* A new fully automated approach for aligning and comparing
shapes. *The Anatomical Record* **298**, 249-276 (2015).
69   Dimitriadou, E., Hornik, K., Leisch, F., Meyer, D. & Weingessel, A. Misc functions of
the Department of Statistics (e1071), TU Wien. *R package* **1**, 5-24 (2008).

70   Golland, P., Liang, F., Mukherjee, S. & Panchenko, D. Permutation tests for
classification. *International Conference on Computational Learning Theory*, 501-515
(2005).
71   Treece, G. M. http://mi.eng.cam.ac.uk/~gmt11/stradview. (2019).
72   Treece, G. M., Gee, A. H., Mayhew, P. & Poole, K. E. High resolution cortical bone
thickness measurement from clinical CT data. *Medical image analysis* **14**, 276-290
(2010).
73   Therneau, T. M. A Package for Survival Analysis in R. R package version 3.1-12,
available at https://CRAN.R-project.org/package=survival. (2020).
836
837

838 **Supplementary Figure 1. Body length is positively correlated with the lengths of lumbar**
839 **vertebrae (LV) 1 - 7.** Pearson correlations between the length of each lumbar vertebra and body
840 length from mole-rat x-ray data. Narrower ovals with darker shades of red indicate larger
841 Pearson correlations; correlation values are also given in the lower left triangle.
842

Supplementary Figure 2. Mole-rat RNA-Seq samples cluster closest with purified mouse osteoblasts, based on canonical bMSC markers. Clustering was performed using Ward's hierarchical clustering method on Euclidean distances of the quantile normalized expression of the seven bMSC markers (out of 11 described[23]) that were quantified in both the mole-rat and reference mouse[24] data sets. The black box indicates mole-rat samples, and the blue box indicates mouse osteoblasts, a bMSC lineage cell type.

Supplementary Figure 3. Estimated cell proportions for the 47 mole-rat RNA-Seq samples.
Each box represents the interquartile range, with the median value depicted as a horizontal bar.
Whiskers extend to the most extreme estimates within 1.5x the interquartile range. Cell
proportions were estimated with CIBERSORT[25], based on reference gene expression levels for
412 marker genes in 27 purified mouse cell types[24]. The predicted predominant cell type in the
mole-rat samples is most similar to early stage osteoblasts, which are cells from the bMSC
lineage.

Supplementary Figure 4. Growth potential in lumbar vertebra 7 (LV7). (a) µCT scan of LV7 of a female Damaraland mole-rat. The boxes indicate the locations of the caudal and cranial growth plates. (b) The number of offspring produced by queens does not significantly predict growth plate fusion (quantified as the average of the caudal and cranial growth plates; $\beta = 2.745$ x 10^{-4}, p = 0.970, n = 12, controlling for age) or chondrocyte proliferation within the remaining growth plate ($\beta = -0.033$, p = 0.293, n = 12, controlling for age). Each box represents the interquartile range, with the median value depicted as a horizontal bar. Whiskers extend to the most extreme values within 1.5x of the interquartile range. Dots represent individual animals, and shading indicates each animal's total offspring number.

Supplementary Figure 5. Queens do not exhibit reduced cortical area at the midsection of
LV6. At top, cross-section with area highlighted in gray shows the measure represented in the
plot. Each line represents an age-matched nonbreeder and queen littermate pair. Queens and
nonbreeders show no difference in cortical area/total area at the LV6 midsection (paired t-test of
cortical area/total area, t = -0.741, df = 6, p = 0.487).

Supplementary Figure 6. Relationship between max load and cortical area in mouse
femurs. Max load shows a highly linear relationship with cortical area across mouse femurs (R^2
$= 0.88$, p $= 6.64$ x 10^{-38}). Each dot represents a single mouse femur. Solid line indicates the best
fit line. Data are from [37].