1 **Morphological and genomic shifts in mole-rat 'queens' increase fecundity but reduce**
2 **skeletal integrity**
3

4 Rachel A. Johnston[1*], Philippe Vullioud[2], Jack Thorley[2], Henry Kirveslahti[4], Leyao Shen[3],
5 Sayan Mukherjee[4,5,6,7], Courtney Karner[3,8], Tim Clutton-Brock[2,9], Jenny Tung[1,10,11*]
6

[1]Department of Evolutionary Anthropology, Duke University, Durham, North Carolina 27708,
USA
[2]Department of Zoology, University of Cambridge, Cambridge CB23EJ, UK
[3]Department of Orthopaedic Surgery, Duke Orthopaedic Cellular, Developmental, and Genome
Laboratories, Duke University School of Medicine, Durham, NC 27710, USA
[4]Department of Statistical Science, Duke University, Durham, NC 27708, USA
[5]Department of Computer Science, Duke University, Durham, NC 27708, USA
[6]Department of Mathematics, Duke University, Durham, NC 27708, USA
[7]Department of Bioinformatics & Biostatistics, Duke University, Durham, NC 27708, USA
[8]Department of Cell Biology, Duke University, Durham, NC 27710, USA
[9]Department of Zoology and Entomology, Mammal Research Institute, University of Pretoria,
0028, Pretoria, South Africa
[10]Department of Biology, Duke University, Durham, North Carolina 27708, USA
[11]Duke Population Research Institute, Duke University, Durham, NC 27708, USA
*Correspondence: racheljohnston7@gmail.com, jt5@duke.edu
22

Abstract

In some mammals and many social insects, highly cooperative societies are characterized by
reproductive division of labor, in which breeders and nonbreeders become behaviorally and
morphologically distinct. While differences in behavior and growth between breeders and
nonbreeders have been extensively described, little is known of their molecular underpinnings.
Here, we investigate the consequences of breeding for skeletal morphology and gene regulation
in highly cooperative Damaraland mole-rats. By experimentally assigning breeding 'queen'
status versus nonbreeder status to age-matched littermates, we confirm that queens experience
vertebral growth that likely confers advantages to fecundity. However, they also up-regulate
bone resorption pathways and show reductions in femoral mass, which predicts increased
vulnerability to fracture. Together, our results show that, as in eusocial insects, reproductive
division of labor in mole-rats leads to gene regulatory rewiring and extensive morphological
plasticity. However, in mole-rats, concentrated reproduction is also accompanied by costs to
bone strength.

Introduction
A hallmark of highly cooperative societies is reproductive division of labor. This
phenomenon is best understood in eusocial insects, where environmental cues lead to
reproductively and morphologically specialized castes, including one or few highly fecund
"queens" [1]. These changes help support the reproductive role of queens by differentiating them
from nonbreeding colony members, who forage, care for young, and engage in colony defense
[1, 2]. Queens are frequently much larger than their sterile colony mates (e.g., twice as large in
honey bees and Pharaoh ants [3, 4]), reflecting dramatically altered growth and development
programs that are explained by changes in gene regulation [5]. Social insects thus exemplify the

tight evolutionary links between reproductive division of labor, cooperative behavior, and
extreme morphological plasticity.
Systems in which breeding is restricted to a single female supported by multiple
nonbreeding helpers are also observed in vertebrates, including birds and mammals [6]. Here,
breeding status is not determined during early development, but instead occurs in adulthood, and
breeding is only achieved by those individuals who have the opportunity to transition into a
reproductive role. In some species, new breeders undergo a period of accelerated growth, which
may be important either for maintaining dominance or for supporting high fecundity [7-12].
While substantial gene regulatory divergence with breeding status has been described for the
brain and some peripheral organs [13-15], we know little about the gene regulatory shifts
responsible for breeder-associated growth patterns. Because morphological change is often
crucial for ramping up offspring production, these processes are key to understanding both the
basis for, and limits of, status-driven differences in growth and development.
Here, we investigate the morphological and molecular consequences of experimental
transitions to breeding status in female Damaraland mole-rats (*Fukomys damarensis*). Like naked
mole-rats, Damaraland mole-rats are frequently classified as 'eusocial' [16-18], and female
helpers who transition to queens experience accelerated vertebral growth associated with
increases in fecundity [9, 11]. However, it is not clear what triggers skeletal remodeling, where it
is localized within the vertebral column, or whether it extends to other parts of the skeleton.
Further, the gene regulatory changes that support skeletal remodeling in mole-rat queens are not
known, nor are their consequences for skeletal growth potential and integrity. To address these
questions, we experimentally assigned age-matched, female littermates to become queens or
remain as nonbreeders and evaluated gene regulatory and morphological changes induced by the
transition to queen status. Our results indicate that, as in eusocial insects, females that acquire
breeding status experience substantial morphological remodeling, associated with pathway-
specific changes in gene regulation. Notably, we found that queens not only experience
lengthening of their lumbar vertebrae, but also show reductions in the growth potential and
structural integrity of their long bones. These changes result from increased rates of bone
resorption that may increase the risk of fracture, indicating that the presence of helpers does not
annul the costs of reproduction to queens.
77
Results
Adaptive plasticity in the skeleton of Damaraland mole-rat queens
Adult female Damaraland mole-rats were randomly assigned to either transition to queen
status (n = 12) or remain as nonbreeders (n = 18) for the duration of the experiment (Figure 1A;
Supplementary Table S1). Age at assignment (mean age = 19.4 ± 4.4 s.d. months) was consistent
with the age at dispersal observed in wild Damaraland mole-rats (1 – 3 years, Thorley and
Clutton-Brock, unpublished data). To resolve whether skeletal changes are a function of the
queen transition *per se* versus release from reproductive suppression in the natal colony,
nonbreeders were either kept in their natal colonies as helpers or placed into solitary housing in
the absence of a breeding queen, recapitulating extended periods of dispersal in this species [18]
(n = 10 helpers and n = 8 solitaires). At the time of assignment, females assigned to the queen,
helper, and solitaire treatments were statistically indistinguishable in body mass, age, or vertebral
length (as measured by lumbar vertebra 5 (LV5); unpaired t-tests between all pairwise
combinations of treatments: p > 0.05; Supplementary Figure S1). When possible, we assigned
age-matched littermates to queen versus nonbreeding treatments (26 of 30 experimental animals

were in sets of littermate sisters; Supplementary Table S1). Six non-experimental animals (1
queen and 5 nonbreeders) were also included in the sample, resulting in a total sample size of 13
breeders and 23 nonbreeders (Supplementary Table S1).
Females assigned to the queen treatment were each transferred to a new tunnel system
containing only an unrelated adult male, simulating the natural process of dispersal and new
colony formation in the wild [18]. This pairing procedure, which defines the queen treatment,
typically leads to immediate sexual activity and rapid activation of the reproductive axis,
including initiation of ovulation and the potential for conception [19, 20]. Queens gave birth to a
mean of 6.92 ± 5.57 s.d. live offspring during the 12 – 22 month follow up period, produced in a
mean of 2.85 ± 1.75 s.d. litters (range: 0 – 6; Supplementary Table S1). As expected, helpers and
solitaires produced no offspring, and did not differ from each other in body mass or vertebral
length after the 12 – 22 month follow-up period (unpaired t-tests, all p > 0.05; Supplementary
Figure S2). Because helpers and solitaires were morphologically indistinguishable, and also
exhibited no differences in gene expression in our subsequent genomic assays (Supplementary
Table S2), we grouped them together into a single "nonbreeder" treatment for the remainder of
our analyses.

Figure 1. Transition to queen status leads to lumbar vertebral lengthening. (a) Experimental design:
nonbreeding adult female (☿) littermates were randomly assigned to transition to queen status (purple ♀) by being
paired with an unrelated male (♂), or to remain in a nonbreeding treatment (cyan). Duration of treatment ranged
from 12 – 22 months. **(b)** Queens show more rapid growth in lumbar vertebra 5 (LV5) in the first four months of the
experiment, relative to nonbreeders (treatment by time point interaction: β = 0.078, n = 49, p = 3.47 x 10^{-3}). Dots
show means +/- standard errors (bars). **(c)** At the start of the experiment (0 months, left panel), the lumbar vertebrae
of breeders do not differ from those of nonbreeders (unpaired t-tests, all p > 0.05). However, at 12 months (right
panel), queens have longer lumbar vertebrae relative to nonbreeders (unpaired t-tests, * indicates p < 0.05). Dots
show means +/- standard errors (bars). Lengths of lumbar vertebrae above the plots are scaled to indicate the mean
lengths of queens (top) and nonbreeders (bottom) at each time point; vertebrae highlighted in purple are significantly
longer in queens relative to nonbreeders. **(d)** Litter size is positively correlated with maternal body length in the
Damaraland mole-rat colony (β = 0.353, n = 328 litters, p = 1.35 x 10^{-3}).
121
Compared to nonbreeders, queens showed rapid growth in the lumbar vertebrae in the
first 12 months post-pairing (Figure 1B), especially in the vertebrae toward the caudal end of the

vertebral column (LV5 and LV6). Based on longitudinal measurements, most of this differential
growth was concentrated soon after the breeding status transition. Specifically, we observed a
significant interaction between breeding status (queen versus nonbreeder) and post-pairing time
point in the first four months of the experiment (Figure 1B; $\beta = 0.0784$, $p = 3.47$ x 10^{-3}; n = 49 x-
rays from 28 animals), but not for measurements taken in later time point intervals (4 months
versus 8 months; 8 versus 12 months, all p > 0.05). Moreover, in the first four months, only
queens that had already experienced pregnancy showed accelerated vertebral lengthening relative
to nonbreeders (unpaired t-test; LV5 of pregnant queens vs. nonbreeders: t = -5.735, df = 16.871,
$p = 2.50$ x 10^{-5}; LV5 of queens not yet pregnant vs. nonbreeders: t = -0.789, df = 13.007, p =
0.444; n = 14 nonbreeders, 5 pregnant queens, and 2 queens not yet pregnant).
As a result of accelerated vertebral growth in queens post-transition, size differences
persisted throughout the study. After 12 months, the absolute length of LV5 in queens was, on
average, 4.8% longer than nonbreeders (Figure 1C; LV5: unpaired t-test, t = 2.509, df = 21.095,
p = 0.020), and the absolute length of the lumbar vertebral column in queens relative to
nonbreeders was 3.5% longer, although the latter difference was not significant (unpaired t-test, t
= 1.945, df = 22.49, p = 0.064). Differences between queens and nonbreeders were even more
apparent if lumbar vertebrae measures were scaled by zygomatic arch (head) width, as in
previous studies [9, 11, 21] (LV5: 9.3% longer, unpaired t-test, t = 4.12, df = 15.135, p = 8.87 x
10^{-4}; lumbar vertebral column length: 7.9% longer, unpaired t-test, t = 4.34, df = 15.37, p = 5.58
x 10^{-4}). Thus, transitions to queen status induce reproductive investment, which in turn leads to
organism-wide allometric changes that generate an elongated phenotype.
The elongated phenotype appears to subsequently facilitate future fecundity. Queens with
longer bodies (which correlates with longer lumbar vertebrae, Pearson's $r = 0.856$, $p = 5.99$ x 10^{-}
^{59}; Supplementary Figure S3) had more pups per litter (Figure 1D; $\beta = 0.353$, $p = 1.35$ x 10^{-3}, n =
328 litters from all breeding groups maintained in the same breeding facility; Supplementary
Table S3). Controlling for litter size, longer queens also had larger pups: for every additional
centimeter of maternal body length, pup body mass increased by 2.9% ($\beta = 0.28$, p = 0.032, n =
971 pups). Thus, the elongated queen phenotype is a strong candidate for adaptive plasticity that
supports increased fertility in queen mole-rats.
153
Breeding status induces gene regulatory changes in the queen mole-rat skeleton
To identify the gene regulatory changes associated with skeletal plasticity, we cultured
cells enriched for bone marrow-derived mesenchymal stromal cells (bMSCs) isolated from the
lumbar vertebrae (pooled LV1 – LV5) of queens and nonbreeders (n = 5 queens, 11
nonbreeders). bMSC cultures include multipotent skeletal stem cells, the precursor of the
osteoblast and chondrocyte lineages responsible for bone growth. In parallel, we cultured cells
enriched for bMSCs from the pooled long bones (humerus, ulna, radius, left femur, and left tibia)
of the same animals, which do not show increased elongation in queens (femur at 12 months:
unpaired t-test, t = -0.202, df = 19.326, p = 0.842; tibia at 12 months: unpaired t-test, t = -0.860,
df = 16.759, p = 0.402). To evaluate the potential role of sex steroid hormone signaling on bone
growth, we treated cells from each bone sample for 24 hours with either 10 nM estradiol or
vehicle control, resulting in 47 total samples. We then performed RNA-seq on each sample to
screen for genes that were systematically differentially expressed in the bone cells of queens
versus nonbreeders.
Of 10,817 detectably expressed genes, 171 genes showed a significant effect of breeding
status at a false discovery rate (FDR) threshold of 10% in the long bones (329 at an FDR of 20%;

Supplementary Table S4). Surprisingly, no genes showed a significant effect of breeding status
in the lumbar vertebrae at either FDR threshold. However, effect sizes were highly correlated
between bone types overall ($R^2 = 0.75$, p = 4.60 x 10^{-53}), with more pronounced effects of
breeding status in the long bone samples than in the lumbar vertebrae (paired t-test on breeding
status effects in long bone versus vertebrae: t = 3.97, df = 317.67, p = 8.73 x 10^{-5}). Importantly,
breeding status-related differences were not readily attributable to differences in bone cell
composition. Based on both canonical markers of bMSC lineage cells and deconvolution of the
RNA-seq data using data from 27 mesenchymal or hematopoietic lineage mouse cell types, the
majority cell type in both queen and nonbreeder samples was most similar to cells from the
bMSC lineage [22-24] (Supplementary Figures S4 and S5). Additionally, the top three principal
components summarizing estimated cell type proportions did not differ between queens and
nonbreeders (all FDR > 10%, Supplementary Table S5), and we identified no cases in which the
effects of breeding status on gene expression were significantly mediated by the first principal
component of cell composition (p > 0.05 for all 171 queen-associated genes at 10% FDR;
Supplementary Table S6).
The majority of breeding status-associated genes were up-regulated in queens (151 of 171
genes, 88%). In support of their role in skeletal plasticity, up-regulated genes were enriched for
bone remodeling ($\log_2$[OR] = 4.07, p = 5.07 x 10^{-6}), a process that involves the balanced cycle
between bone formation by osteoblasts and bone resorption by osteoclasts [25] (Figure 2).
Surprisingly, however, enriched pathways were specifically related to bone resorption, not
formation (Supplementary Table S7), including "positive regulation of bone resorption" (Figure
2A, C; $\log_2$[OR] = 6.51, p = 1.55 x 10^{-6}) and "superoxide anion generation," which is involved in
osteoclast activity and degradation of bone matrix (Figure 2A, C; $\log_2$[OR] = 5.29, p = 1.4 x 10^{-5}
) [26-29]. Differentially expressed genes were also enriched for immune-related processes (e.g.,
"cytokine secretion", "chemotaxis", "leukocyte activation involved in immune response";
Supplementary Table S7). These observations suggest that transitions to queen status also
involve changes in immunoregulatory signaling (osteoclast cells are derived from monocytes).
Omni-ATAC-seq profiling of open chromatin regions further supports a central role for
bone resorption and osteoclast activity in the queen skeleton (n = 8; Supplementary Table S8).
Specifically, transcription factor binding motifs (TFBMs) located in accessible chromatin near
queen up-regulated genes were enriched for PU.1 and MITF, two transcription factors that are
essential for osteoclast differentiation [30] (Figure 2B, C; PU.1 $\log_2$[OR] = 1.041, p = 2.84 x 10^{-4}
; MITF $\log_2$[OR] = 0.707, p = 7.36 x 10^{-3}; see Supplementary Table S8 for complete list of
enriched TFBMs). *MITF* was also among the 151 genes that were differentially expressed
between queens and nonbreeders and up-regulated in both queen long bones and lumbar
vertebrae. Surprisingly, given the role of sex steroid hormones in bone growth and elevated
estradiol levels in queen versus helper Damaraland mole rats [31], we observed no significant
effects of estradiol treatment on gene expression in either bone type (all FDR > 10%). Queen up-
regulated genes were also not in closer proximity to androgen response elements (ARE) or
estrogen response elements (ERE) than expected by chance (ARE $\log_2$[OR] = 0.207, p = 0.627;
ERE $\log_2$[OR] = 0.196, p = 0.652). Consistent with this observation, transcription factor
footprinting analysis showed no evidence of queen-associated differences in transcription factor
activity of the androgen receptor, estrogen receptor 1 (ESR1), or estrogen receptor 2 (ESR2), in
either the long bones or lumbar vertebrae (all paired t-tests: p > 0.05; Supplementary Figure S6).
Thus, our data point to the involvement of non-sex steroid-mediated signaling pathways in
remodeling queen mole-rat bones, at least after one year post-transition.

Figure 2. Queen status drives increased regulatory activity of bone resorption pathways. (a) Gene Ontology (GO) terms enriched in queen up-regulated genes, relative to the background set of all genes tested. Bars represent 95% confidence intervals. Processes highlighted in purple are also depicted in (c). Highest-level (most general) terms are shown; for full GO enrichment results, see Supplementary Table S7. **(b)** Accessible transcription factor binding site motifs enriched near queen up-regulated genes, relative to all genes tested. Bars represent 95% confidence intervals. Transcription factors highlighted in purple are also depicted in (c). **(c)** Schematic of the balance between bone formation and bone resorption, showing key regulators and markers for mesenchymal stem cell differentiation into osteoblasts and hematopoietic stem cell differentiation into osteoclasts [25, 30]. Note that not all genes or proteins in gray were tested for differential expression (e.g., because they were not annotated in the Damaraland mole-rat genome or were too lowly expressed in our sample); see Supplementary Table S4 for full set of tested genes. Queen up-regulated genes or corresponding proteins (FDR < 10%) are highlighted as purple ovals, and transcription factors with binding motifs enriched near queen up-regulated genes are highlighted as purple hexagons. Inset for osteoclasts shows the NADPH oxidase system, which generates superoxide radicals (O_2^-) necessary for bone resorption and is highly enriched for queen up-regulated genes (purple ovals).

Extensive skeletal remodeling in queen mole-rats

The gene expression data suggest that queen status-driven changes to the skeleton extend beyond the lumbar vertebrae to the long bones. Further, they suggest that bone resorption—an important counterpoint to bone formation that is required for normal skeletal maintenance—also distinguishes breeding and nonbreeding females. To investigate this possibility, we performed high-resolution micro-computed tomography (μCT) scanning to generate 3D reconstructions of LV6, LV7, right femur, and right tibia of queens and female nonbreeders (n = 140 bones from 36 animals; Figure 3A; Supplementary Figure S7). This approach substantially increases the level of resolution for investigating breeding status-linked differences in skeletal morphology, as previous studies relied on x-ray data alone [9, 11].

We first asked whether breeding status could be predicted from morphological differences in the 3D reconstructions. We found that it could for the lumbar vertebra, but not for the femur: by applying the smooth Euler characteristic transform [32], we were able to predict

queen versus nonbreeder status in LV6 (77.8% accuracy, p = 0.01, n = 36), but not the femur
(52.8% accuracy, p = 0.53, n = 36). Including only highly fecund queens (≥6 total offspring)
improved predictive accuracy in the femur (70% accuracy, p = 0.12, n = 30). Although these
predictions did not reach statistical significance, they raised the possibility that morphological
changes in femurs become enhanced with increasing reproductive effort.
We next tested whether the transition to queen status affects the ability to continue bone
lengthening. Lengthening requires the presence of a growth plate, a region of cartilage in the
bone where longitudinal growth occurs through proliferation of cartilage cells (chondrocytes)
(Figure 3A, B; Supplementary Figure S7). Closure of the growth plate, which indicates that bone
lengthening potential has terminated, typically occurs in mammals after reaching sexual
maturation, when energy begins to be invested in reproduction instead of growth [33]. To test
whether the transition to queen status alters bone lengthening potential, we performed Safranin-
O staining on sections of the right tibia and LV7 to visualize growth plates (Figure 3B). In the
proximal tibia but not LV7, queens were less likely to have open growth plates (Figure 3C;
Supplementary Figure S7 and Table S9; tibia: two-sided binomial test, p = 0.019; LV7: two-
sided binomial test, p = 0.422). The increased probability of growth plate closure in the tibia of
queens is linked to the number of offspring a female has produced: females with more offspring
showed a higher expanse of closure across the growth plate (β = 0.050, p = 4.51 x 10^{-3}, n = 12,
controlling for age). This pattern may be due in part to reduced chondrocyte proliferation, as
females that produced more offspring had fewer chondrocyte columns in the remaining growth
plate (Figure 3D; β = -0.132, p = 0.020, n = 12, controlling for age). Thus, offspring production
in queens is associated with loss of the ability to lengthen the long bones, but not the lumbar
vertebrae, consistent with the importance of abdominal lengthening for supporting larger litters.
A major demand on reproductively active female mammals is a high requirement for
calcium, particularly during lactation when mothers support rapid offspring bone growth.
Maternal skeletons are remodeled to meet this demand, although in most mammals, these
changes are not permanent (reviewed in [34]). Because of the particularly intense reproductive
investment made by cooperatively breeding mole-rat queens, we therefore also evaluated the
effect of queen status on trabecular and cortical bone volumes, which are thought to be important
in satisfying short-term and long-term calcium demands, respectively. We found no effect of
queen status on the amount of trabecular bone in the femur, tibia, LV6, or LV7 (all p > 0.05 for
bone volume/total volume). However, we found that cortical bone was significantly thinner at
the femoral midshaft, but not in the lumbar vertebrae, in queens compared to their nonbreeding
sisters (Figure 3E; Supplementary Figure S8; femur: paired t-test of cortical area/total area, t = -
4.067, df = 8, p = 3.60 x 10^{-3}; LV6: paired t-test of cortical area/total area, t = -0.741, df = 6, p =
0.487). Cortical thinning in queens appears to be specifically due to increased bone resorption,
which typically occurs on the endosteal (internal) surface of long bones in the marrow cavity.
Indeed, queens had a larger marrow cavity (paired t-test, t = 5.355, df = 8, p = 6.82 x 10^{-4}; Figure
3F) but showed no difference in periosteal area compared to their nonbreeding sisters (paired t-
test, t = 1.539, df = 8, p = 0.162; Figure 3G).
284

Figure 3. Queen status leads to reduced growth potential in the tibia and reduced cortical area at the femoral midshaft. (a) μCT scans of Damaraland mole-rat tibias. Boxes indicate the location of the proximal growth plate, which varies between unfused (left) to fully fused (right). **(b)** Example Safranin-O stained histological sections of the proximal tibia, in which the growth plate is unfused (top), partially fused (middle), or fully fused (bottom). Values indicate percent growth plate fusion across the width of the bone. The cartilaginous growth plate is stained deep pink, and calcified bone is stained green. **(c)** Queens, and specifically queens that gave birth to more offspring, show increased growth plate fusion ($\beta = 0.050$, $p = 4.51 \times 10^{-3}$, n = 12, controlling for age) and **(d)** decreased number of chondrocyte columns within the remaining growth plate ($\beta = -0.132$, $p = 0.020$, n = 12, controlling for age). Each box represents the interquartile range, with the median value depicted as a horizontal bar. Whiskers extend to the most extreme values within 1.5x of the interquartile range. In (c) and (d), dots represent individual animals, and shading indicates each animal's total offspring number. Ages of queens and nonbreeders do not significantly differ (unpaired t-test, $t = 0.489$, n = 12, $p = 0.644$). **(e-g)** Femoral cross-sections with area highlighted in gray show the measures represented in the corresponding plots below. Each line represents an age-matched, nonbreeder and queen littermate pair. **(e)** Queens have less cortical bone (relative to the total area of the femoral midshaft cross-section), compared to their paired nonbreeding littermates (paired t-test, $t = -4.07$, df = 8, $p = 3.60 \times 10^{-3}$). **(f)** Queens also have enlarged marrow cavities (paired t-test, $t = 5.36$, df = 8, $p = 6.82 \times 10^{-4}$) but **(g)** show no difference in overall periosteal area (paired t-test, $t = 1.54$, df = 8, $p = 0.162$).

Because changes in cortical bone are thought to reflect accumulated demands over long time frames, we hypothesized that cortical thinning in queens is a consequence of repeated cycles of pregnancy and lactation over time, which can occur simultaneously in Damaraland mole-rat queens. In support of this idea, we found that, within queens, the relative amount of

cortical bone is not predicted by the number of pups in a queen's most recent litter (β = -0.024, n
= 13, p = 0.287), but instead by the total number of pups she produced in her lifetime.
Specifically, queens who had more live births had reduced cortical bone thickness along the
entire shaft of the femur (Figure 4 and Supplementary Table S10; across decile sections of the
femur: all p < 0.05, controlling for mother's litter as a random effect). Thus, cortical thinning
does not commence with the transition to queen status *per se* (i.e., it is not a correlate of
achieving breeder status), but instead appears to be a consequence of repeated investment in
pregnancy and lactation. Notably, thinning is particularly marked in queens who had at least six
offspring, which usually occurs by 14 months after a breeding status transition (i.e., by the third
litter; Supplementary Table S10). Given that wild Damaraland mole-rat queens can maintain
their status for many years [35], our results suggest that long-lived queens may experience
substantial morphological change.

Figure 4. Offspring production in queens leads to cortical thinning across the femoral shaft. (a) Queens (top
row) relative to their same-aged female nonbreeding littermates (bottom row) present thinner cortical bone across
the femur, particularly in females that have many offspring (top right). Number of offspring is indicated above each
femur, and vertical arrows indicate littermate pairs. **(b)** Within each decile section across the femoral shaft, number
of offspring is negatively correlated with average cortical thickness (linear mixed model with littermate pair as
random effect). Full results are presented in Supplementary Table S10. Asterisk indicates p < 0.05.
325

Skeletal remodeling predicts increased risk of femur breakage in queens
In humans, accelerated bone resorption is a central cause of osteoporosis-related bone
fragility [36]. We therefore hypothesized that cortical thinning in queen mole-rat femurs would
be linked to decreased bone strength. To test this hypothesis, we calculated two key indicators of
femoral structural integrity: cortical area (CA) and the minimum moment of inertia (I_{min}, a
predictor of resistance to bending). In nonbreeders, both are positively correlated with body mass
(I_{min}: $R^2 = 0.368$, n = 24, p = 9.92 x 10^{-4}; CA: $R^2 = 0.407$, n = 24, p = 4.80 x 10^{-4}). However, in
queens, I_{min} is not significantly predicted by body mass ($R^2 = 0.088$, n = 13, p = 0.17), but is

instead a function of number of offspring produced (R^2 = 0.283, p = 0.035). Queen CA is
predicted by both offspring number and body mass, but offspring number explains almost twice
the variance (offspring number R^2 = 0.634, p = 6.83 x 10^{-4}; body mass R^2 = 0.385, p = 0.014).
To evaluate the effects of reproductive activity on the risk of bone failure, we drew on
data on the relationship between CA and bone mechanical failure in a large data set of mouse
femurs [37]. In this data set, CA is the best predictor of maximum load (the maximum force a
bone can withstand prior to failure), and, crucially, the CA-max load relationship is highly linear
(Supplementary Figure S9; R^2 = 0.88, p = 6.64 x 10^{-38}). Scaling the mole-rat CA data to mouse
suggests that transitions to queen significantly increase the risk of bone failure (Figure 5; hazard
ratio (95% confidence interval) = 2.67 (1.20, 5.93), n = 36, p = 0.016). Similar to growth
potential and cortical thinning, this effect is driven by highly fertile queens, such that those who
had at least six offspring showed the highest predicted risk of bone failure (Figure 5; queens with
≥ 6 offspring relative to nonbreeders: HR = 3.81 (1.47, 9.83), n = 30, p = 0.006). The risk of
bone failure is thus predicted to increase by 21% for each additional pup (HR = 1.21 (1.10, 1.33),
n = 36, p = 8.85 x 10^{-5}).

Figure 5. Effect of reproductive status on the probability of bone failure. Survival curves for femurs from nonbreeders versus
queens (Wald test, p = 0.02, n = 36) and versus queens with ≥ 6 offspring (Wald test, p = 0.006, n = 30), based on predictions
from the midshaft cortical area and data from [37]. Vertical dashed lines indicate group medians, with the median failure time for
nonbreeders fixed at a value of 1.
353

Discussion

Our results demonstrate that transitions to breeding status in Damaraland mole-rat queens
lead to a cascade of skeletal changes linked to shifts in gene regulation. The vertebral
lengthening observed in Damaraland mole-rat queens is concordant with previous reports of
vertebral lengthening in both Damaraland mole-rats [11] and naked mole-rats [9]. Like naked
mole-rats, our analyses show that most growth is concentrated soon after the breeding status
transition, especially in connection with the first post-transition pregnancies [21, 38]. However,
our findings also suggest subtle differences: for instance, while the growth phenotype in naked
mole-rats occurs at the cranial end of the lumbar vertebrae [38], it is concentrated at the caudal
end of the vertebral column in Damaraland mole-rats. Given that Damaraland mole-rats and
naked mole-rats independently evolved a similar, highly cooperative social structure [18, 39],
this difference suggests potential convergent evolution of the vertebral lengthening phenotype in
queens, presumably in response to the selection pressure for increased fertility.

In addition to previously described vertebral growth, we found that queen Damaraland
mole-rats lose bone lengthening potential in the long bones and develop thinner femurs that are
predicted to be more prone to mechanical failure. Moreover, gene expression levels in queens
reflect a signature of bone resorption, rather than bone growth, at the time of sampling, which
occurred 1 – 2 years post-transition. The molecular signature of bone resorption temporally
aligns with changes in morphology, in which accelerated vertebral growth primarily occurs
during a female's first pregnancy, whereas cortical thinning in the long bones are a function of
repeated cycles of offspring production. Thus, queens quickly progress from traits typically
associated with pre-reproductive and pubertal growth in mammals (e.g., body elongation), to
traits typically linked to aging (e.g., marrow cavity expansion and cortical thinning).
The complex pattern of bone growth and bone resorption in queens likely involves
multiple regulatory mechanisms. Because estrogen is known to impact bone growth and
maintenance [40, 41], and estrogen levels are higher in mole-rat queens relative to nonbreeding
females [31], we hypothesized that queens and nonbreeders would differ in their response to
estradiol in bone marrow-derived cells. Surprisingly, we observed no gene expression response
to estradiol treatment. By itself, this result could be a function of the specific concentration or
duration of estradiol treatment we applied. However, we also observed no enrichment for
estrogen receptor binding motifs near queen up-regulated genes, and no evidence that estrogen or
androgen receptor binding sites are differentially bound in cells from queens versus non-
breeders. Thus, our results suggest a role for other, as-yet unknown signaling pathways in the
queen-specific signature of long bone cortical resorption (although it does not exclude the
importance of estrogen signaling in other phenotypes, such as bone elongation or growth plate
closure, [42]).
Bone loss in Damaraland mole-rat queens may be an extreme of the typical mammalian
pattern of bone remodeling, in which bone mineral density decreases during pregnancy and
lactation, but recovers once offspring are weaned [34]. Thinning in mole-rats may be sustained,
however, because queens begin gestating soon after lactating for the previous litter, leaving little
to no time for recovery. One possible reason that this fast rate of breeding is achievable is that
queens in colonies with more helpers work less and rest more [43], consistent with studies in
other cooperative mammals that show that helpers alleviate breeding-associated loss of condition
in queens [44]. Paradoxically, helpers might not only help offset costs of, but also contribute to,
decreased bone mass in queens, given that large numbers of helpers are themselves produced via
high queen fecundity, and reduced physical activity can also lead to decreases in bone mass [45].
The extent to which helpers reduce the costs of breeding to queens may also differ
between species depending on the relative numbers of helpers to breeders. For example, in
eusocial insects, large colonies and the high ratio of helpers to queens reduce the costs of
reproduction to queens to very low levels [1, 2]. Similarly, in naked mole-rats (where colonies
can include hundreds of animals, compared to dozens in Damaraland mole-rat colonies [17, 18,
46]), a small sample of queens (n = 6) suggests increased rather than decreased femoral cortical
thickness relative to age-matched nonbreeders [47]. Testing how the costs and benefits of
reproduction are resolved across different levels of cooperativity, including the molecular
mechanisms that mediate these differences, is an important next step towards understanding the
evolution of cooperative breeding in mammals.
Finally, despite frequent analogies between Damaraland mole-rats and eusocial insects
[18, 46, 48], our results suggest some key points of differences. Specifically, while abdominal
lengthening allows queen mole-rats to increase fecundity per reproductive effort, loss of cortical

bone in the femur is unlikely to directly benefit either fertility or survival. Instead, it reflects the
cumulative burden of continuous cycles of pregnancy and lactation [34]. Thus, unlike eusocial
insect queens [49, 50], Damaraland mole-rat queens incur morphological costs to concentrated
reproduction in addition to morphological changes that facilitate increased fitness. How these
costs translate into fertility or survival outcomes in natural populations remains a fascinating,
unanswered question.
419
Materials and Methods
Study system and experimental design
Damaraland mole-rats (*Fukomys damarensis*) were maintained in a captive colony at the
Kuruman River Reserve in the Northern Cape Province of South Africa, within the species'
natural range. Only animals born in captivity, with known birthdates and litter composition, were
used in this study, so that exact ages were known. Animals were maintained in artificial tunnel
systems built from PVC pipes with compartments for a nest-box and waste-box and transparent
windows to allow behavioral observation [51]. Animals were fed *ad libitum* with sweet potatoes
and cucumbers.
Adult females (> 1 year) from 16 natal colonies were randomly assigned to be either
nonbreeders or queens, such that females assigned to queen status had age-matched littermates,
where possible, who were assigned to the nonbreeding condition. To distinguish the effects of
queen status from release from reproductive suppression, nonbreeders were either maintained in
their natal colony as helpers or maintained alone, which models the social condition experienced
by dispersing females. Females assigned to the breeder condition were transferred to a new
tunnel system with an unrelated male from a separate social group. Nine new breeding females, 6
helpers, and 8 solitaire females (age-matched littermates where possible; Supplementary Table
S1) were set up in December 2015 – July 2016 [11]. With one exception (animal G10F026),
animals maintained their breeding status for 14 – 22 months before sample collection. One queen
and five helpers that were siblings, but not age-matched littermates, of experimental animals
were also included in sample collection. To increase the final sample size, an additional 4
breeding colonies, matched against 4 age-matched littermate helpers, were formed in October
2017 and followed for 11-12 months (Supplementary Table S1). One queen died before sample
collection, and one non-experimental helper was euthanized during the course of the study and
included in sample collection. The final sample size included 13 queens, 15 helpers, and 8
solitaire females.
446
X-ray data
For a subset of study subjects, full body X-rays were taken using the Gierth TR 90/20
battery-operated generator unit with portable Leonardo DR Mini plate (OR Technology,
Rostock, Germany) every two months during the first 12 months of the experiment and at the
time of sacrifice. From these X-rays, an experimenter blind to animal breeding status measured
the length of each lumbar vertebra (from vertebra 1 to 7), the right femur, the right tibia, body
length, and the width of the zygomatic arch using ImageJ [52]. The caudal-most lumbar vertebra
was labeled as LV7. We tested for an effect of breeding status on LV5 using a linear mixed
model in which post-pairing time point, breeding status, and the interaction of time point by
breeding status were modeled as fixed effects and animal ID as a random effect.
457
Effects of queen body length on fertility

To test the effect of maternal body length on litter size and pup size, we used body length
measurements obtained during routine colony monitoring of all queens maintained in the captive
colony (i.e., not restricted to experimental animals). Following [11], we used body length
measurements obtained nearest to, and no more than 90 days from, the date of parturition. The
resulting dataset included 328 litters (971 pups) from 76 mothers, which represents a 76%
increase over an earlier analysis of this relationship in [11]. We fit two linear mixed effects
models. In the first model, we modeled litter size as a function of maternal body length,
controlling for whether the litter was the female's first litter, and included maternal ID as a
random effect. In the second model, we modeled pup mass as a function of maternal body length,
controlling for litter size and whether the litter was the female's first litter as fixed effects, and
maternal ID and litter ID as random effects.
470
Sample collection and cell culture from lumbar vertebrae and long bones
Animals were deeply anesthetized with isoflurane and sacrificed with decapitation
following USGS National Wildlife Health Center guidelines and under approval from the
Animal Ethics Committee of the University of Pretoria. Immediately upon sacrifice, the lumbar
vertebrae and long bones were dissected, and attached muscle tissue was removed with forceps.
Lumbar vertebrae 6 and 7 and the right femur and tibia were collected into 50% ethanol for 24
hours, then transferred to 70% ethanol and stored at 4° C for μCT scans and histochemistry.
To isolate bone cells for culture, lumbar vertebrae 1 – 5 were incubated in 2%
Collagenase P (Roche, Switzerland) for 30 minutes at 30° C. Each bone was then cut in half and
transferred to a 1.5 ml microcentrifuge tube containing a G-Tube microcentrifuge tube (VWR,
Radnor, PA, USA) that had been punctured at the bottom with a 15 gauge needle. Tubes were
spun at 3,000 RCF for 5 seconds, allowing the marrow to collect into the 1.5 mL microcentrifuge
tube. Cell pellets were resuspended in red blood cell lysis buffer, pooled, and incubated for 3
minutes at room temperature. 10 ml bMSC media (MEM-alpha [ThermoFisher, Waltham, MA,
USA] + 15% fetal bovine serum [Hyclone, Logan, UT, USA] + 1% penicillin/streptomycin + 2
ng/ml recombinant human fibroblast growth factor-basic [Biocam, Centurion, Gauteng, South
Africa] + 10 nM ROCK inhibitor Y-27632 [RI; Cayman Chemical, Ann Arbor, MI, USA]) was
added to stop lysis, and the tubes were spun for 5 minutes at 300 RCF. The cell pellet was
resuspended in 1 ml bMSC media and strained through a 70 μm cell strainer. Cells were plated at
1.6×10^5 cells per cm^2. The long bones (excluding right femur and tibia) were processed to
enrich for bMSCs following the same procedure, but without incubation in Collagenase P. Cells
were cultured at 37° C and 5% CO_2. Twenty-four hours post plating, plates were carefully
washed three times with 1x PBS and supplied with fresh media to remove non-adherent cells.
Once bMSC clusters were visible (2 – 9 days post plating), plates were fed bMSC media without
RI or fed bMSC media without RI + 10 nM estradiol (E2). Twenty-four hours later, cells were
collected into buffer RLT and stored at -80° C. Samples were shipped on dry ice to Duke
University for RNA extraction using the Qiagen RNeasy Micro Kit. RNA-Seq libraries were
generated using the NEBNext Single Cell/Low Input RNA Library Prep Kit for Illumina.
499
Gene expression analysis
RNA-Seq libraries were sequenced on an Illumina HiSeq 4000 (100 base pair single end
reads) to a mean coverage of 16.1 ± 3.9 s.d. million reads. Reads were trimmed with *cutadapt*
version 2.3 [53] (parameters: -q 20 -e 0.2 --times 5 --overlap 2 -a AGATCGGAAGAGC -a "T" -
-minimum-length=20). Trimmed reads were then mapped to the Damaraland mole-rat v1.0

genome [54] (DMR_v1.0) using two pass mapping with STAR [55]. Only uniquely mapped
reads were retained. HTseq [56] was used to quantify read counts mapping to genes (using the
v1.0.92 gtf file from Ensembl; we extended the genomic coordinates of the *SERPINE1* gene by
2000 basepairs in both directions due to very high expression directly adjacent to the annotated
coordinates). We transformed read counts to transcripts per million (TPM) [57], and retained
only genes with TPM ≥ 2 in at least 25% of samples. We performed voom normalization [58] on
the raw counts, using normalization factors produced by the trimmed mean of M-values (TMM)
method [59] in DESeq [60]. We used the *limma* [61] function *lmFit* to regress out the proportion
of uniquely mapped reads in genes (which controls for efficiency of mRNA selection during
RNA-Seq library preparation) and animal natal colony (which controls for littermate sets and
date of sacrifice) to obtain normalized, batch-corrected gene expression values for downstream
analysis. We used the mixed effects model approach in *emmreml* [62] to estimate, for each gene,
the effect of breeding status on gene expression within lumbar vertebrae and within long bones
using the following model:
519

$$y_i = \mu + d_i\beta_1 + b_i\beta_2 + q_i\beta_3 * I(b=0) + q_i\beta_4 * I(b=1) + s_i\beta_5 * I(b=0) + s\beta_6 * I(b=1)$$
$$+ Zu + \varepsilon_i,$$

520
$$u \sim MVN\ 0, \sigma_u^2 K,$$
$$\varepsilon \sim MVN\ 0, \sigma_e^2$$

521
where *y* is the vector of gene expression levels for *n* = 47 samples (indexed by *i*); μ is the
intercept; *d* is the number of days in culture and β_1 its effect size; *b* is bone type (i.e., long bones
or lumbar vertebrae) and β_2 its effect size; and *q* is a 0/1 variable representing breeder status and
β_3 and β_4 its effect size in long bones and lumbar vertebrae, respectively. *I* is an indicator
variable for bone type (0 = long bone; 1 = lumbar vertebrae). *s* is a 0/1 variable representing
whether the cells were cultured with estradiol and β_5 and β_6 are its effect sizes in long bones and
lumbar vertebrae, respectively. Z is an incidence matrix that maps samples to animal ID to take
into account repeated sampling from the same individual, and *u* is a random effect term that
controls for relatedness. *K* is an *m* by *m* matrix of pairwise relatedness estimates (derived from
the genotype data, described below) between all *m* animals. ε is the residual error, σ_u^2 is the
genetic variance component, and σ_e^2 is the environmental variance component. We also ran an
identical model but with an additional fixed effect of solitaire status in long bones and in lumbar
vertebrae, to test for a difference in gene expression between helpers and solitaires. To control
for multiple testing, we calculated the false discovery rate following Storey and Tibshirani [63]
using an empirical null distribution derived from 100 permutations of each variable of interest.
We used g:profiler [64] to perform gene ontology enrichment analysis of the genes up-
regulated with queen status in lumbar vertebrae and long bones (151 of 171 genes significantly
associated with queen status at a 10% FDR). All genes in the original analysis set were used as
the background gene set. We set both the minimum size of the functional category and the
minimum size of the query/term intersection to 3. Finally, we retained categories that passed a
Bonferroni corrected p-value of 0.05.
543
Genotyping to estimate relatedness
To control for relatedness when modeling the gene expression data, we performed single
nucleotide polymorphism (SNP) genotyping of the RNA-Seq data using the Genome Analysis
Toolkit [65] (GATK). We used the SplitNCigarReads function on the trimmed, uniquely mapped

reads and performed GATK indel realignment. Base recalibration was performed by using all
SNPs with GQ ≥ 4 in an initial UnifiedGenotyper run on the full data set as a reference.
Genotypes were called on the recalibrated bam files using HaplotypeCaller. Variants were
filtered with the following GATK VariantFiltration parameters: QUAL < 100.0, QD < 2.0, MQ <
35, FS > 30, HaplotypeScore > 13, MQRankSum < -12.5, ReadPosRankSum < -8. Variants were
further filtered with *vcftools* [66] to only retain biallelic SNPs in Hardy-Weinberg equilibrium (p
> 0.05) with minor allele frequency ≥ 0.1, minimum mean depth of 5, max missing count of 2,
and minimum GQ of 99. Finally, SNPs were thinned to a distance of 10 kb basepairs, resulting in
a final dataset of 1,965 stringently filtered biallelic SNPs. Missing values were imputed using
beagle [67], and the resulting vcf file was used to create a kinship matrix using *vcftools* [66].
Values of the kinship matrix were confirmed to be higher in known siblings compared to non-
siblings (unpaired t-test, t = 27.939, p = 2.23 x 10^{-12}; means = 0.513 and -0.097). Two pairs of
siblings were found to have different fathers (G1F022 and G1F025; G4F020 and G4F019).
561
Cell type heterogeneity
Although selection for adherent cells from bone marrow enriches for bMSCs, other cell
types are also present [68]. To assess whether cell type heterogeneity accounts for queen-
associated differential expression, we used CIBERSORT to deconvolve the proportion of
component cell types from the RNA-Seq data [24]. We trained CIBERSORT on a data set of
quantile normalized gene expression values from mouse purified primary cell populations [23].
Specifically, we subset the training data to 27 purified cell populations of mesenchymal or
hematopoietic origin (Figure S3) and to genes that were included in our mole-rat gene expression
dataset. We then predicted the composition of the cells that contributed to the mole-rat quantile
normalized gene expression data set, for each sample.
To test whether cell type heterogeneity was significantly explained by queen status, we
also modeled cell type proportion (as summarized by the first principal component of
CIBERSORT-estimated proportions for all 27 potential cell types; PC1 explains 50.9% of the
overall variation) following the same method used for gene-by-gene expression analysis but with
PC1 included as an explanatory variable. We then performed mediation analysis on each of the
171 genes that showed a significant effect of breeding status at FDR < 10%. To do so, we first
estimated the indirect effect of breeding status on gene expression through the mediating variable
(CIBERSORT PC1). The indirect effect of breeding status through CIBERSORT PC1 was
estimated by calculating the difference in the effect of breeding status between two models: one
model that did not include the mediator (i.e., β_3 and β_4 from equation 1 above) and the same
model with the addition of the mediating variable. We performed 1000 iterations of bootstrap
resampling to obtain the 95% confidence interval for the indirect effect, and considered an
indirect effect for a gene significant if the 95% interval did not overlap 0.
585
ATAC-seq data and transcription factor binding site analysis
To investigate whether differentially expressed genes were associated with accessible
binding motifs for specific transcription factors, we generated Omni-ATAC-seq data to profile
regions of open chromatin [69, 70]. We performed Omni-ATAC-seq on both lumbar vertebrae
bMSCs and long bone bMSCs from two female nonbreeding and two queen mole-rats (n = 8
libraries total), following the published protocol [70] with the following modifications: 5,000
cells were centrifuged at 500 RCF for 5 minutes at 4°C. The pellet was resuspended in 50 ul
transposition mix (25 ul 2xTD buffer, 16.5 ul PBS, 6.75 ul water, 1 ul 10% NP40, 1 ul 10%

Tween-20, 1 ul 1% digitonin, and 0.25 ul Tn5 transposase). The reaction was incubated at 37°C
for 30 minutes without mixing, followed by a 1.5x Ampure bead cleanup. Omni-ATAC libraries
were sequenced on a NovaSeq 6000 as 100 basepair paired-end reads to a mean coverage (± SD)
of 26.9 (±4.4) million reads (range: 16.8 – 38.3). Reads were trimmed with Trim Galore! [71] to
remove adapter sequence and low quality basepairs (Phred score < 20; reads ≥ 25 bp). Read pairs
were mapped to the DMR_v1.0 genome using *bwa-mem* [72] with default settings. Only
uniquely mapped reads were retained. The alignment bam files for each treatment (breeding or
nonbreeding) were merged, and open chromatin regions were identified using MACS2 v2.1.2
[73] with the following parameters: "-nomodel -keep-dup all -q 0.05". We combined open
chromatin peaks with regions in the DMRv1.0 genome that match sequences of vertebrate
transcription factor binding site motifs, using motifs defined in the HOMER database [74]. We
used Fisher's exact tests (using a p-value threshold of 0.01) to test if transcription factor binding
motifs belonging to the same transcription factor were enriched in open chromatin regions within
2,000 bp of the 5' most transcription start site of queen up-regulated genes.
To compare genome-wide signatures of DNA-transcription factor binding for androgen
receptor (AR), estrogen receptor 1 (ESR1), and estrogen receptor 2 (ESR2), we characterized
transcription factor footprints in queens and nonbreeders, in both the lumbar vertebra and long
bones, using HINT-ATAC from the Regulatory Genomics Toolbox (RGT) with default
parameters [75]. We focused on the subset of peak regions called using MACS2 [73]. We
identified TF footprints by merging reads within each bone type-breeding status combination and
calling footprints on the combined data. For each bone type, we then created a meta-footprint set
by merging the respective footprint calls across queens and nonbreeders using the *bedtools*
function *merge* [76]. We identified transcription factor motifs in the *DMR_v1.0* genome that fell
within meta-footprints, based on the JASPAR CORE Vertebrates set of curated position
frequency matrices [77]. Finally, we tested for differential footprints of AR, ESR1, and ESR2
binding using the RGT *differential* function, using the activity score metric described in [75] and
default parameters.
621
Micro-CT scans and analysis
We performed μCT scans of LV6, LV7, right femur, and right tibia using a VivaCT 80
scanner (Scanco Medical AG, Brüttisellen, Switzerland) set at 55 kVp and 145 μA, with voxel
size 10.4 μm. Trabecular bone was quantified from the 100 μCT slices below the proximal tibia
growth plate, the 100 μCT slices above the distal femur growth plate, and the 100 μCT slices
medial to the caudal growth plate of LV6. To obtain midshaft cross-sections of the femur, tibia,
and LV6, we first reduced each bone mesh to 100,000 faces using Avizo Lite version 9.7.0.
Mesh files from the same bone type were auto-aligned using Auto3dgm [78] in Matlab. Aligned
mesh files were then back scaled to their original sizes in Matlab, and the midshaft cross-section
was generated using Rhinoceros version 6. The MomentMacro plugin in ImageJ was used to
calculate bone area and moment of inertia.
633
Classification of breeding status from bone shape
To predict breeding status from bone shape, we first applied the smooth Euler
characteristic transform [32] to the aligned right femur and LV6. We then performed leave-one-
out predictions, running each bone type separately, using the linear kernel and c-classification
with the support vector machine (SVM) implemented by the R package *e1071* [79]. The SVM
classifier was equipped with 1:100,000 weighting to achieve class balanced predictions. The

empirical p-values were estimated for each bone type by running 100 permutations of the
queen/nonbreeder labels [80].
642

Histochemistry
For a subset of individuals (Supplementary Table S9), the tibia and LV7 were plasticized,
sectioned, and stained with Safranin O by the Washington University Musculoskeletal Research
Center. The proportion of the tibia proximal growth plate that was fused, and the mean
proportion of the LV7 cranial and caudal growth plates that were fused, were measured in
ImageJ from Safranin O stained sections. To quantify growth plate activity, we calculated the
number of chondrocyte columns (defined as linear stacks of at least three chondrocytes)
controlling for length of open growth plate. For each bone type (tibia and LV7), we ran two
models: proportion growth plate fusion or chondrocyte columns per mm growth plate as the
dependent variable, and number offspring born and age as the independent variables.
653

Cortical thickness across the femur
We used Stradview [81, 82] on dicom images from the μCT scans to measure and
visualize, in an automated manner, cortical thickness across the surface of the femur. Bone
surface was defined in Stradview by thresholding pixel intensity and contouring the bone at
every 14 sections, with the following parameters: resolution = medium, smoothing = standard,
strength = very low, contour accuracy = 6. To measure cortical thickness, we used the auto
threshold method in Stradview, with line width set to 5, smooth set to 1, and line length set to 3
mm. The smoothed thickness values of each femur were then registered (i.e., mapped) to a single
"canonical" femur surface (mole-rat GRF002) using wxRegSurf v18
(http://mi.eng.cam.ac.uk/~ahg/wxRegSurf/). We sectioned the cortical thickness values into
deciles according to location along the length of the femur. The top and bottom deciles were
removed, because cortical and trabecular bone towards the ends of the femur could not be easily
differentiated by the automated method. Deciles were then recreated for the remaining length of
the bone (i.e., the central 80%). From each bone decile, we estimated cortical thickness as the
mean of all cortical thickness measures within that interval. For each decile across animals, we
used a linear mixed model to model cortical thickness as a function of breeding status and
number offspring, with litter pair as a random effect.
671

Modeling the probability of bone failure
Previous research on mechanical properties of rodent femurs found that, among several
morphological and compositional traits measured in eight morphologically varying mouse
strains, cortical area (CA) at the midshaft was the best predictor of maximum load (defined as
the greatest force attained prior to bone failure, measured via four-point bending; published
Pearson's $r = 0.95$) [37]. We therefore used cortical area at the femoral midshaft to predict max
load of Damaraland mole-rat femurs. To do so, we first fit a linear model of max load as a
function of cortical area (unadjusted for body weight) using published mouse data ($R^2 = 0.877$, n
$= 81$, p = 6.64×10^{-38}) [37]. We extrapolated from this linear fit to predict max load from cortical
area at the midshaft of Damaraland mole-rat femurs. Predicted max loads were then used as input
for Cox proportional hazards models, comparing either all queens to nonbreeders or queens with
≥ 6 offspring to nonbreeders. Models were fit using the R function *coxph*, and were confirmed to
meet the proportional hazards assumption using the *cox.zph* function in the R package *survival*
[83]. Because max load was not directly measured in the Damaraland mole-rats, we used the Cox

proportional hazards models to specifically evaluate the relative hazard of bone failure
depending on queen status/number of offspring. We therefore report the results in Figure 5 based
on relative force (with the median predicted failure value for nonbreeders set to 1) instead of
absolute force in Newtons.
690
691 **Data Availability**
692 All RNA sequencing data generated during this study are available in the NCBI Gene Expression
693 Omnibus (series accession GSE152659). ATAC-Seq data are available in the NCBI Sequence
694 Read Archive (BioProject accession number PRJNA649596). µCT data from this study are
695 available on MorphoSource (http://www.morphosource.org, project 1056).
696

Acknowledgements
We thank Tim Vink, Dave Gaynor, and the mole-rat house staff and volunteers for their
tremendous contributions to the Kalahari Mole-rat Project. We also thank Irene Garcia, Mari
Cobb, Brianna Bowman, Anna Luiza Wolf, Alice Zhou, Yilin Yu, B.J. Nielsen, and Tawni
Voyles for their contributions to sample collection and data generation, Graham Treece for
guidance on quantifying bone thickness with Stradview, Karl Jepsen for sharing data on mouse
femurs, Lou DeFrate for guidance on estimating bone strength, Saideep Gona and Luis Barreiro
for their contributions on the footprint analysis, and members of the Tung lab for feedback on
earlier versions of this manuscript. Support for this work was provided by the European
Research Council (Grants 294494 and 742808 to T.C.B), the Human Frontier Science Program
(RGP0051-2017 to J.T., S.M., and T.C.B.), the National Science Foundation (IOS-7801004 to
J.T.), the National Institutes of Health (F32HD095616 to R.A.J.), a Sloan Foundation Early
Career Research Fellowship to J.T., a Foerster-Bernstein Postdoctoral Fellowship to R.A.J., and
a Natural Environmental Research Council Doctoral Training Program to Ja.T. High-
performance computing resources were supported by the North Carolina Biotechnology Center
(Grant Number 2016-IDG-1013).

713
714 **Author Contributions**
715 Conceptualization, R.A.J., T.C.B., and J.T.; Investigation, R.A.J., Ja.T., P.V., H.K., L.S., S.M.,
716 C.K., T.C.B., and J.T. Formal Analysis, R.A.J., H.K., S.M.; Writing—Original Draft, R.A.J. and
717 J.T.; Writing—Reviewing & Editing, R.A.J., P.V., Ja.T., H.K., L.S., S.M., C.K., T.C.B., and J.T.
718 Funding Acquisition, J.T. and T.C.B. Supervision, T.C.B. and J.T.
719
Competing Interests
The authors declare no competing interests.

Supplementary Figure S1. Queen, helper, and solitaire female mole-rats do not differ in
mass, age, or lumbar vertebra 5 length at the start of the experiment. Each box represents
the interquartile range, with the median value depicted as a horizontal bar. Whiskers extend to
the most extreme values within 1.5x of the interquartile range. Dots represent individual animals.
P-values are from unpaired t-tests between each pairwise comparison between helpers (n = 10),
queens (n = 12), and solitaires (n = 8). For LV5 length, data were only available for 10 queens.
Raw data values are provided in Supplementary Table S1.

Supplementary Figure S2. Mass and LV5 length of helper and solitaire female mole-rats
after 12 – 22 months of experimental treatment of social status. Helper and solitaire female
mole-rats do not differ in (a) mass (unpaired t-test; t = 0.496, df = 12.733, p = 0.629) or (b)
lumbar vertebra 5 length (unpaired t-test; t = -0.358, df = 8.391, p = 0.729) after 12 – 22 months
of experimental treatment of social status. Each box represents the interquartile range, with the
median value depicted as a horizontal bar. Whiskers extend to the most extreme values within
1.5x of the interquartile range. Dots represent individual animals. Raw data values are provided
in Supplementary Table S1.

Supplementary Figure S3. Body length is positively correlated with the lengths of lumbar
vertebrae (LV) 1 – 7. Pearson correlations between the length of each lumbar vertebra and body
length from mole-rat x-ray data. Narrower ovals with darker shades of red indicate larger
Pearson correlations; correlation values are also given in the lower left triangle.
741

Supplementary Figure S4. Mole-rat RNA-Seq samples cluster closest with purified mouse osteoblasts, based on canonical bMSC markers. Clustering was performed using Ward's hierarchical clustering method on Euclidean distances of the quantile normalized expression of the seven bMSC markers (out of 11 described [22]) that were quantified in both the mole-rat and reference mouse [23] data sets. The black box indicates mole-rat samples, and the blue box indicates mouse osteoblasts, a bMSC lineage cell type.

Supplementary Figure S5. Estimated cell proportions for the 47 mole-rat RNA-Seq samples. Each box represents the interquartile range, with the median value depicted as a horizontal bar. Whiskers extend to the most extreme estimates within 1.5x the interquartile range. Cell proportions were estimated with CIBERSORT [24], based on reference gene expression levels for 412 marker genes in 27 purified mouse cell types [23]. The predicted predominant cell type in the mole-rat samples is most similar to early stage osteoblasts, which are cells from the bMSC lineage.

Supplementary Figure S6. Footprint profiles of transcription factors androgen receptor
(AR), estrogen receptor 1 (ESR1), and estrogen receptor 2 (ESR2). Transcription factor
footprints were profiled separately for lumbar vertebrae (n = 4; 2 queens and 2 nonbreeders) and
for long bones (n = 4; 2 queens and 2 nonbreeders). Transcription factor activity was not
significantly different between queens and nonbreeders in any of the three transcription factors,
in either bone type (paired t-tests; all p > 0.05).

Supplementary Figure S7. Growth potential in lumbar vertebra 7 (LV7). (a) μCT scan of LV7 of a female Damaraland mole-rat. The boxes indicate the locations of the caudal and cranial growth plates. (b) The number of offspring produced by queens does not significantly predict growth plate fusion (quantified as the average of the caudal and cranial growth plates; β = 2.745 x 10^{-4}, p = 0.970, n = 12, controlling for age) or chondrocyte proliferation within the remaining growth plate (β = -0.033, p = 0.293, n = 12, controlling for age). Each box represents the interquartile range, with the median value depicted as a horizontal bar. Whiskers extend to the most extreme values within 1.5x of the interquartile range. Dots represent individual animals, and shading indicates each animal's total offspring number.

Supplementary Figure S8. Queens do not exhibit reduced cortical area at the midsection of LV6. At top, cross-section with area highlighted in gray shows the measure represented in the plot. Each line represents an age-matched nonbreeder and queen littermate pair. Queens and nonbreeders show no difference in cortical area/total area at the LV6 midsection (paired t-test of cortical area/total area, t = -0.741, df = 6, p = 0.487).

Supplementary Figure S9. Relationship between max load and cortical area in mouse
femurs. Max load shows a highly linear relationship with cortical area across mouse femurs (R^2
$= 0.88$, p $= 6.64 \times 10^{-38}$). Each dot represents a single mouse femur. Solid line indicates the best
fit line. Data are from [37].

References
1. Wilson, E.O. (1971). The insect societies. The insect societies.
2. Keller, L., and Genoud, M. (1997). Extraordinary lifespans in ants: a test of
evolutionary theories of ageing. Nature *389*, 958-960.
3. Berndt, K.P., and Eichler, W. (1987). Die Pharaoameise, Monomorium pharaonis
(L.)(Hym., Myrmicidae). Mitteilungen aus dem Museum für Naturkunde in Berlin.
Zoologisches Museum und Institut für Spezielle Zoologie (Berlin) *63*, 3-186.
4. Page Jr, R.E., and Peng, C.Y.-S. (2001). Aging and development in social insects with
emphasis on the honey bee, Apis mellifera L. Experimental gerontology *36*, 695-711.
5. Smith, C.R., Toth, A.L., Suarez, A.V., and Robinson, G.E. (2008). Genetic and genomic
analyses of the division of labour in insect societies. Nature Reviews Genetics *9*, 735-
748.
6. Koenig, W.D., and Dickinson, J.L. (2016). Cooperative breeding in vertebrates:
studies of ecology, evolution, and behavior, (Cambridge University Press).
7. Clutton-Brock, T., Hodge, S., Spong, G., Russell, A., Jordan, N., Bennett, N., Sharpe, L.,
and Manser, M. (2006). Intrasexual competition and sexual selection in cooperative
mammals. Nature *444*, 1065-1068.
8. Huchard, E., English, S., Bell, M.B., Thavarajah, N., and Clutton-Brock, T. (2016).
Competitive growth in a cooperative mammal. Nature *533*, 532-534.
9. O'Riain, M., Jarvis, J., Alexander, R., Buffenstein, R., and Peeters, C. (2000).
Morphological castes in a vertebrate. Proceedings of the National Academy of
Sciences *97*, 13194-13197.
10. Russell, A.F., Carlson, A.A., McIlrath, G.M., Jordan, N.R., and Clutton‐Brock, T.
(2004). Adaptive size modification by dominant female meerkats. Evolution *58*,
1600-1607.
11. Thorley, J., Katlein, N., Goddard, K., Zöttl, M., and Clutton-Brock, T. (2018).
Reproduction triggers adaptive increases in body size in female mole-rats.
Proceedings of the Royal Society B: Biological Sciences *285*, 20180897.
12. Young, A.J., and Bennett, N.C. (2010). Morphological divergence of breeders and
helpers in wild Damaraland mole‐rat societies. Evolution: International Journal of
Organic Evolution *64*, 3190-3197.
13. Bens, M., Szafranski, K., Holtze, S., Sahm, A., Groth, M., Kestler, H.A., Hildebrandt, T.B.,
and Platzer, M. (2018). Naked mole-rat transcriptome signatures of socially
suppressed sexual maturation and links of reproduction to aging. BMC biology *16*,
77.
14. Mulugeta, E., Marion-Poll, L., Gentien, D., Ganswindt, S.B., Ganswindt, A., Bennett,
N.C., Blackburn, E.H., Faulkes, C.G., and Heard, E. (2017). Molecular insights into the
pathways underlying naked mole-rat eusociality. bioRxiv, 209932.
15. Sahm, A., Hoffmann, S., Koch, P., Henning, Y., Bens, M., Groth, M., Burda, H., Begall, S.,
Ting, S., and Goetz, M. (2020). Status-dependent aging rates in long-lived, social
mole-rats are shaped by HPA stress axis. bioRxiv.
16. Bennett, N.C., and Faulkes, C.G. (2000). African mole-rats: ecology and eusociality,
(Cambridge University Press).
17. Jarvis, J. (1981). Eusociality in a mammal: cooperative breeding in naked mole-rat
colonies. Science *212*, 571-573.

18.   Jarvis, J., and Bennett, N. (1993). Eusociality has evolved independently in two
826   genera of bathyergid mole-rats—but occurs in no other subterranean mammal.
827   Behavioral Ecology and Sociobiology *33*, 253-260.
828   19.   Bennett, N., Faulkes, C., and Molteno, A. (1996). Reproductive suppression in
829   subordinate, non-breeding female Damaraland mole-rats: two components to a
830   lifetime of socially induced infertility. Proceedings of the Royal Society of London.
831   Series B: Biological Sciences *263*, 1599-1603.
832   20.   Snyman, P., Jackson, C.R., and Bennett, N.C. (2006). Do dispersing non-reproductive
833   female Damaraland mole-rats, Cryptomys damarensis (Rodentia: Bathyergidae)
834   exhibit spontaneous or induced ovulation? Physiology & behavior *87*, 88-94.
835   21.   Dengler‐Crish, C.M., and Catania, K.C. (2009). Cessation of reproduction‐related
836   spine elongation after multiple breeding cycles in female naked mole‐rats. The
837   Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology:
838   Advances in Integrative Anatomy and Evolutionary Biology *292*, 131-137.
839   22.   Dominici, M., Le Blanc, K., Mueller, I., Slaper-Cortenbach, I., Marini, F., Krause, D.,
840   Deans, R., Keating, A., Prockop, D., and Horwitz, E. (2006). Minimal criteria for
841   defining multipotent mesenchymal stromal cells. The International Society for
842   Cellular Therapy position statement. Cytotherapy *8*, 315-317.
843   23.   Hume, D.A., Summers, K.M., Raza, S., Baillie, J.K., and Freeman, T.C. (2010).
844   Functional clustering and lineage markers: insights into cellular differentiation and
845   gene function from large-scale microarray studies of purified primary cell
846   populations. Genomics *95*, 328-338.
847   24.   Newman, A.M., Liu, C.L., Green, M.R., Gentles, A.J., Feng, W., Xu, Y., Hoang, C.D., Diehn,
848   M., and Alizadeh, A.A. (2015). Robust enumeration of cell subsets from tissue
849   expression profiles. Nature methods *12*, 453.
850   25.   Redlich, K., and Smolen, J.S. (2012). Inflammatory bone loss: pathogenesis and
851   therapeutic intervention. Nature reviews Drug discovery *11*, 234-250.
852   26.   Darden, A.G., Ries, W.L., Wolf, W.C., Rodriguiz, R.M., and Key Jr, L.L. (1996).
853   Osteoclastic superoxide production and bone resorption: stimulation and inhibition
854   by modulators of NADPH oxidase. Journal of Bone and Mineral Research *11*, 671-
855   675.
856   27.   Datta, H., Rathod, H., Manning, P., Turnbull, Y., and McNeil, C. (1996). Parathyroid
857   hormone induces superoxide anion burst in the osteoclast: evidence for the direct
858   instantaneous activation of the osteoclast by the hormone. Journal of endocrinology
859   *149*, 269-275.
860   28.   Key, L., Ries, W., Taylor, R., Hays, B., and Pitzer, B. (1990). Oxygen derived free
861   radicals in osteoclasts: the specificity and location of the nitroblue tetrazolium
862   reaction. Bone *11*, 115-119.
863   29.   Key, L., Wolf, W., Gundberg, C., and Ries, W. (1994). Superoxide and bone resorption.
864   Bone *15*, 431-436.
865   30.   Segeletz, S., and Hoflack, B. (2016). Proteomic approaches to study osteoclast
866   biology. Proteomics *16*, 2545-2556.
867   31.   Bennett, N. (1994). Reproductive suppression in social Cryptomys damarensis
868   colonies—a lifetime of socially‐induced sterility in males and females (Rodentia:
869   Bathyergidae). Journal of Zoology *234*, 25-39.

32.   Crawford, L., Monod, A., Chen, A.X., Mukherjee, S., and Rabadán, R. (2016).
Functional data analysis using a topological summary statistic: the smooth Euler
characteristic transform. arXiv preprint arXiv:1611.06818.
33.   Kilborn, S.H., Trudel, G., and Uhthoff, H. (2002). Review of growth plate closure
compared with age at sexual maturity and lifespan in laboratory animals. Journal of
the American Association for Laboratory Animal Science *41*, 21-26.
34.   Kovacs, C.S. (2016). Maternal mineral and bone metabolism during pregnancy,
lactation, and post-weaning recovery. Physiological reviews *96*, 449-547.
35.   Schmidt, C.M., Jarvis, J.U., and Bennett, N.C. (2013). The long-lived queen:
reproduction and longevity in female eusocial Damaraland mole-rats (Fukomys
damarensis). African Zoology *48*, 193-196.
36.   Szulc, P., Seeman, E., Duboeuf, F., Sornay‐Rendu, E., and Delmas, P.D. (2006). Bone
fragility: failure of periosteal apposition to compensate for increased endocortical
resorption in postmenopausal women. Journal of bone and mineral research *21*,
1856-1863.
37.   Jepsen, K.J., Akkus, O.J., Majeska, R.J., and Nadeau, J.H. (2003). Hierarchical
relationship between bone traits and mechanical properties in inbred mice.
Mammalian Genome *14*, 97-104.
38.   Henry, E.C., Dengler-Crish, C.M., and Catania, K.C. (2007). Growing out of a caste-
reproduction and the making of the queen mole-rat. Journal of Experimental Biology
210, 261-268.
39.   Faulkes, C.G., and Bennett, N.C. (2016). Damaraland and naked mole-rats:
convergence of social evolution. Cooperative breeding in vertebrates, 338-352.
40.   Cutler Jr, G. (1997). The role of estrogen in bone growth and maturation during
childhood and adolescence. The Journal of steroid biochemistry and molecular
biology *61*, 141-144.
41.   Khalid, A.B., and Krum, S.A. (2016). Estrogen receptors alpha and beta in bone. Bone
87, 130-135.
42.   Juul, A. (2001). The effects of oestrogens on linear bone growth. Apmis *109*, S124-
S134.
43.   Houslay, T.M., Vullioud, P., Zöttl, M., and Clutton-Brock, T.H. (2020). Benefits of
cooperation in captive Damaraland mole-rats. Behavioral Ecology.
44.   Clutton-Brock, T.H., and Manser, M. (2016). Meerkats: cooperative breeding in the
Kalahari. Cooperative breeding in vertebrates *294*, 317.
45.   Morseth, B., Emaus, N., and Jørgensen, L. (2011). Physical activity and bone: The
importance of the various mechanical stimuli for bone mineral density. A review.
Norsk epidemiologi *20*.
46.   Jarvis, J.U., O'Riain, M.J., Bennett, N.C., and Sherman, P.W. (1994). Mammalian
eusociality: a family affair. Trends in Ecology & Evolution *9*, 47-51.
47.   Pinto, M., Jepsen, K., Terranova, C., and Buffenstein, R. (2010). Lack of sexual
dimorphism in femora of the eusocial and hypogonadic naked mole-rat: a novel
animal model for the study of delayed puberty on the skeletal system. Bone *46*, 112-
120.

48.   Burda, H., Honeycutt, R.L., Begall, S., Locker-Grütjen, O., and Scharff, A. (2000). Are
naked and common mole-rats eusocial and if so, why? Behavioral Ecology and
Sociobiology *47*, 293-303.
49.   Rodrigues, M.A., and Flatt, T. (2016). Endocrine uncoupling of the trade-off between
reproduction and somatic maintenance in eusocial insects. Current opinion in insect
science *16*, 1-8.
50.   Rueppell, O., Aumer, D., and Moritz, R.F. (2016). Ties between ageing plasticity and
reproductive physiology in honey bees (Apis mellifera) reveal a positive relation
between fecundity and longevity as consequence of advanced social evolution.
Current opinion in insect science *16*, 64-68.
51.   Zöttl, M., Vullioud, P., Mendonça, R., Ticó, M.T., Gaynor, D., Mitchell, A., and Clutton-
Brock, T. (2016). Differences in cooperative behavior among Damaraland mole rats
are consequences of an age-related polyethism. Proceedings of the National
Academy of Sciences *113*, 10382-10387.
52.   Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25
years of image analysis. Nature methods *9*, 671-675.
53.   Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet. journal *17*, 10-12.
54.   Fang, X., Seim, I., Huang, Z., Gerashchenko, M.V., Xiong, Z., Turanov, A.A., Zhu, Y.,
Lobanov, A.V., Fan, D., and Yim, S.H. (2014). Adaptations to a subterranean
environment and longevity revealed by the analysis of mole rat genomes. Cell
reports *8*, 1354-1364.
55.   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15-21.
56.   Anders, S., Pyl, P.T., and Huber, W. (2014). HTSeq–A Python framework to work with
high-throughput sequencing data. Bioinformatics, btu638.
57.   Wagner, G.P., Kin, K., and Lynch, V.J. (2012). Measurement of mRNA abundance
using RNA-seq data: RPKM measure is inconsistent among samples. Theory in
biosciences *131*, 281-285.
58.   Law, C.W., Chen, Y., Shi, W., and Smyth, G.K. (2014). Voom: precision weights unlock
linear model analysis tools for RNA-seq read counts. Genome biology *15*, R29.
59.   Robinson, M.D., and Oshlack, A. (2010). A scaling normalization method for
differential expression analysis of RNA-seq data. Genome biology *11*, R25.
60.   Anders, S., and Huber, W. (2010). Differential expression analysis for sequence
count data. Genome Biol *11*, R106.
61.   Smyth, G.K. (2005). Limma: linear models for microarray data. In Bioinformatics and
computational biology solutions using R and Bioconductor. (Springer), pp. 397-420.
62.   Akdemir, D., and Okeke, U. (2015). EMMREML: Fitting mixed models with known
covariance structures. R package version *3*.
63.   Storey, J.D., and Tibshirani, R. (2003). Statistical significance for genomewide
studies. Proc Natl Acad Sci U S A *100*, 9440-9445.
64.   Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., and Vilo, J.
(2019). g: Profiler: a web server for functional enrichment analysis and conversions
of gene lists (2019 update). Nucleic acids research *47*, W191-W198.

65. McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A.,
Garimella, K., Altshuler, D., Gabriel, S., and Daly, M. (2010). The Genome Analysis
Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing
data. Genome research *20*, 1297-1303.
66. Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker,
R.E., Lunter, G., Marth, G.T., and Sherry, S.T. (2011). The variant call format and
VCFtools. Bioinformatics *27*, 2156-2158.
67. Browning, S.R., and Browning, B.L. (2007). Rapid and accurate haplotype phasing
and missing-data inference for whole-genome association studies by use of localized
haplotype clustering. The American Journal of Human Genetics *81*, 1084-1097.
68. Phinney, D.G., Kopen, G., Isaacson, R.L., and Prockop, D.J. (1999). Plastic adherent
stromal cells from the bone marrow of commonly used strains of inbred mice:
variations in yield, growth, and differentiation. Journal of cellular biochemistry *72*,
570-585.
69. Buenrostro, J.D., Giresi, P.G., Zaba, L.C., Chang, H.Y., and Greenleaf, W.J. (2013).
Transposition of native chromatin for fast and sensitive epigenomic profiling of
open chromatin, DNA-binding proteins and nucleosome position. Nature methods
10, 1213.
70. Corces, M.R., Trevino, A.E., Hamilton, E.G., Greenside, P.G., Sinnott-Armstrong, N.A.,
Vesuna, S., Satpathy, A.T., Rubin, A.J., Montine, K.S., and Wu, B. (2017). An improved
ATAC-seq protocol reduces background and enables interrogation of frozen tissues.
Nature methods *14*, 959.
71. Krueger, F. (2015). Trim Galore.
http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/.
72. Li, H., and Durbin, R. (2010). Fast and accurate long-read alignment with Burrows–
Wheeler transform. Bioinformatics *26*, 589-595.
73. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum,
C., Myers, R.M., Brown, M., and Li, W. (2008). Model-based analysis of ChIP-Seq
(MACS). Genome biology *9*, R137.
74. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,
Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B
cell identities. Molecular cell *38*, 576-589.
75. Li, Z., Schulz, M.H., Look, T., Begemann, M., Zenke, M., and Costa, I.G. (2019).
Identification of transcription factor binding sites using ATAC-seq. Genome biology
20, 45.
76. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for
comparing genomic features. Bioinformatics *26*, 841-842.
77. Sandelin, A., Alkema, W., Engström, P., Wasserman, W.W., and Lenhard, B. (2004).
JASPAR: an open‐access database for eukaryotic transcription factor binding
profiles. Nucleic acids research *32*, D91-D94.
78. Boyer, D.M., Puente, J., Gladman, J.T., Glynn, C., Mukherjee, S., Yapuncich, G.S., and
Daubechies, I. (2015). A new fully automated approach for aligning and comparing
shapes. The Anatomical Record *298*, 249-276.

79.   Dimitriadou, E., Hornik, K., Leisch, F., Meyer, D., and Weingessel, A. (2008). Misc
functions of the Department of Statistics (e1071), TU Wien. R package *1*, 5-24.
80.   Golland, P., Liang, F., Mukherjee, S., and Panchenko, D. (2005). Permutation tests for
classification. International Conference on Computational Learning Theory, 501-
515.
81.   Treece, G.M. (2019). http://mi.eng.cam.ac.uk/~gmt11/stradview.
82.   Treece, G.M., Gee, A.H., Mayhew, P., and Poole, K.E. (2010). High resolution cortical
bone thickness measurement from clinical CT data. Medical image analysis *14*, 276-
290.
83.   Therneau, T.M. (2020). A Package for Survival Analysis in R. R package version 3.1-
12, available at https://CRAN.R-project.org/package=survival.
1013