Evidence that non-pathogenic microorganisms drive sea star wasting disease through boundary layer oxygen diffusion limitation

Citlalli A. Aquino[1*], Ryan M. Besemer[2*], Christopher M. DeRito[3], Jan Kocian[4], Ian R. Porter[5], Peter Raimondi[6], Jordan E. Rede[3], Lauren M. Schiebelhut[7], Jed P. Sparks[8], John P. Wares[9], Ian Hewson[3†]

[1]Department of Biology, Estuary and Ocean Science Center, San Francisco State University, Romberg Tiburon Campus, 3150 Paradise Dr, Tiburon CA 94132 USA

[2]Center for Marine Science, University of North Carolina Wilmington, 5600 Marvin K Moss Ln, Wilmington, NC 28409 USA

[3]Department of Microbiology, Cornell University, Wing Hall 403, Ithaca NY 14853 USA

[4]Unaffiliated Researcher, Leland WA 98376 USA

[5]Department of Clinical Sciences, College of Veterinary Medicine, Cornell University, 602 Tower Rd, Ithaca, NY 14853 USA

[6]Institute of Marine Sciences, Department of Ecology & Evolutionary Biology, University of California Santa Cruz, 1156 High Street, Santa Cruz, CA 95064 USA

[7]Life and Environmental Sciences, University of California Merced, 5200 North Lake Rd. Merced, CA 95343 USA

[8]Department of Ecology & Evolutionary Biology, Cornell University, Corson Hall, Ithaca NY 14853 USA

[9]Department of Genetics, University of Georgia, 30602 120 W Green St, Athens, GA 30602 USA

*Denotes equal author contribution

†Corresponding Author: **hewson@cornell.edu**

ABSTRACT

Sea star wasting disease describes a condition affecting asteroids that resulted in significant Northeastern Pacific population decline following a mass mortality event in 2013. The etiology of sea star wasting is unresolved. We hypothesized that asteroid wasting is a sequela of microbial organic matter remineralization near respiratory surfaces which leads to boundary layer oxygen diffusion limitation (BLODL). Wasting lesions were induced in *Pisaster ochraceus* by enrichment with a variety of organic matter (OM) sources and by experimentally reduced oxygen conditions. Microbial assemblages inhabiting tissues and at the asteroid-water interface bore signatures of copiotroph proliferation before wasting onset, followed by the proliferation of putatively facultative and strictly anaerobic taxa. These results together illustrate that suboxic conditions at the animal-water interface may be established by heterotrophic bacterial activity in response to organic matter loading. Wasting susceptibility was significantly and positively correlated with rugosity, a key determinant of boundary layer thickness. At a semi-continuously monitored field site (Langley Harbor), wasting predictably occurred at annual peak or decline in phytoplankton biomass over 5 years, suggesting that primary production-derived OM may contribute to BLODL. Finally, wasting individuals from 2013 – 2014 bore stable isotopic signatures reflecting anaerobic processes which suggests that this phenomenon may have affected asteroids during mass mortality. The impacts of BLODL may be more pronounced under higher temperatures due to lower O_2 solubility, in more rugose asteroid species due to restricted hydrodynamic flow, and in larger specimens due to their lower surface area to volume ratios which affects diffusive respiratory potential. Moreover, our results demonstrate that marine invertebrate disease may result from heterotrophic microbial activity that occurs adjacent to respiratory tissues, which raises important questions about the etiology of marine diseases in other benthic taxa.

Keywords: **Sea Star Wasting, Oxygen, Heterotroph, Remineralization, Phytoplankton**

INTRODUCTION

Sea star wasting (SSW) disease describes a suite of clinical signs in asteroids including loss of turgor, arm twisting, epidermal ulceration, limb autotomy, and death. The condition gained prominence in 2013 when it caused mass mortality of >20 asteroid species in the Northeastern Pacific (Hewson et al., 2014) with continuous observations since (Miner et al., 2018;Jaffe et al., 2019). However, lesions compatible with SSW in various asteroid species have been reported since at least 1896 in the Eastern US (Mead, 1898), and at many locations globally (reviewed in Hewson et al., 2019). The cause of SSW is unresolved. Early reports that SSW was associated with a densovirus (Hewson et al., 2014) were refuted by subsequent investigation that failed to show a consistent association between the virus and presence of disease (Hewson et al., 2018), and recent description of persistent and phylogenetically widespread infection by related densoviruses (Jackson et al., 2020a;Jackson et al., 2020b) suggest this virus to be a component of normal microbiome. Furthermore, wasting is not consistently associated with any bacterial or microbial eukaryotic organism (Hewson et al., 2018). Environmental conditions, including elevated water temperatures (Eisenlord et al., 2016;Kohl et al., 2016), lower water temperatures and higher pCO_2 (Menge et al., 2016), and meteorological conditions (Hewson et al., 2018) correspond with wasting at distinct locations. Recent modelling studies suggest repeated sea surface temperature anomalies may correlate with wasting (Aalto et al., 2020). Reports of SSW spread between adjacent geographic locations, through public aquarium intakes, and challenge experiments with tissue homogenates suggested a transmissible etiology (Hewson et al., 2014;Bucci et al., 2017). However, there is a lack of mechanistic understanding how SSW is generated in affected individuals.

Here we provide convergent evidence that asteroid wasting is a sequela of boundary layer oxygen diffusion limitation (BLODL; Fig. 1). In this model, elevated organic matter (OM) concentrations stimulate the growth of copiotrophic microorganisms adjacent to animal surfaces, driving down dissolved O_2 concentrations and causing suboxic conditions at the animal-seawater interface within the diffusive boundary layer. Over time, these conditions do not meet respiratory O_2 demand of tissues, resulting in their damage and decomposition, which further enriches near-asteroid pools of organic material. This in turn results in the proliferation of anaerobic taxa on and within tissues. Here we provide evidence for this effect through study of microbiome

composition during organic matter amendment and as animals waste in the absence of external

stimuli. Further, we demonstrate that wasting can result from suboxic water column conditions.

Next, we illustrate that wasting susceptibility is related to inherent asteroid properties, including

rugosity (a key physical characteristic influencing diffusive boundary layer height), animal size

and respiratory demand. We explored the relationship between primary production and wasting

during a 5 year time series at a field site and find significant correlation between wasting

occurrence and trends in chlorophyll a, which is one potential source of OM for bacterial

nutrition. Finally, we demonstrate that wasting asteroids from the 2013-2014 mass mortality

event bore stable isotopic signatures that reflect anaerobic microbial processes compared to their

asymptomatic sympatric counterparts.

METHODS

Gross wasting definition and general specimen characteristics: Sea star wasting (SSW) in field

populations is reported to comprise a wide suite of signs, including loss of turgor (deflation),

discoloration, puffiness, arm twisting/curling, limb autotomy, body wall lesions and erosions and

protrusion of pyloric caeca and gonads (Hewson et al., 2014). There is currently no case

definition for any species of asteroid, nor is there progression of disease signs characteristic of

wasting (i.e. wasting has no pathognomic signs; Hewson et al., 2019). Because many of these

disease signs are subject to observer bias, we sought to standardize occurrence of wasting

between experiments and surveys. We considered wasting as the appearance of non-focal body

wall lesions across all experiments. We used the time between experiment initiation and body

wall lesion appearance (i.e. lesion genesis) as a standardized parameter to assess the speed of

wasting. Experiments were performed where individuals were physically isolated from each

other while in the same aquarium water body through plastic containers which had been punched

with holes (4mm diameter) at a density of ~ 2 holes cm^{-2}. However, in the May – June 2018

observation of wasting in the absence of external stimuli, individuals were identified by the

unique "fingerprint" of apical circlet paxillae and spines and were not physically separated from

each other. In all experiments, lesion position was standardized by clockwise naming of rays

from the madreporite. Individual asteroids were considered dead when tube feet did not move

during observation for 30s, or when all limbs autotomized away from the central disc. The mass

of individuals was determined using a top pan balance, and each ray measured from tip to the

center of central disc using a ruler or calipers.

Impact of organic matter enrichment on wasting in* Pisaster ochraceus*: We examined the

impact of OM enrichment on asteroid wasting to test whether enrichment caused bacterial

abundance and composition shifts consistent with the boundary layer oxygen diffusion limitation

(BLODL) hypothesis. Twenty *P. ochraceus* (mass 303 ± 26 g) were obtained from the jetty at

Bodega Bay in July 2019 and transported to the Bodega Bay Marine Lab (UC Davis), where they

were placed in flow-through large volume sea tables for 7 d prior to commencement of the

experiment. Asteroids were placed individually into baskets and suspended in each of 4 sea

tables with flow rates of 60 ± 15 mL s^{-1}. A dense ($\sim 10^4$ cells mL^{-1}) culture of *Dunaliella*

tertiolecta was prepared in 20 L artificial seawater, which was filtered onto 0.2 µm Durapore

filters and resuspended in 1 L artificial seawater. The resuspended matter was divided into 21 x

45 mL aliquots and frozen prior to use. Coastal particulate OM (POM) was prepared by filtering

40 L seawater from the unfiltered intake pipe at Bodega Marine Laboratory through 0.2 µm

Durapore filters, which was then resuspended in 1 L seawater. This was also divided into 21 x 35

ml aliquots and frozen before use. Individuals (n = 5 per treatment in separate sea tables) were

amended with 3 OM sources: 1) peptone (approx 300 mg individual^{-1}); 2) 45mL of *Dunaliella*

tertiolecta filtered culture; and 3) Coastal POM (6mL) daily. Individual asteroids were observed

daily (aboral and ventral surface) for the presence of lesions and they were weighed. There was

no significant difference in the rate of animal mass change over the course of the experiment

between treatments (Fig. S1), where individuals lost 11.0 ± 0.2 % of their initial mass over the

course of the experiment. Daily samples (2 mL of water) for surface bacterial abundance were

withdrawn using 3mL syringes which were pressed onto the aboral surface of individual stars

while immersed, and preserved in 5% formalin and kept at 4°C in darkness prior to processing.

Surface microbial communities were sampled every 48 h by collecting a swab (sterile cotton-

tipped, dry transport system; Puritan) from each aboral surface which were then frozen at -20°C

prior to processing. Microbiome compositional analyses was performed as described later in this

section.

Wasting progression in the absence of external stimuli: We sought to investigate longitudinal

microbiome composition over wasting progression in lesion margins and in artificial scar tissues,

in an attempt to discern opportunistic taxa from potential pathogens. This work was performed to

target epidermis-associated microorganisms which were minimally sampled (i.e. by surface

scrape), and in a separate study to target body wall tissues (i.e. by transverse body wall biopsy

punch). We anecdotally observed in previous work (Hewson et al., 2018) that sampling stress,

especially by collection of tissues by biopsy punch, may influence wasting in aquarium studies

so sought to minimize sampling before the appearance of lesions.

To examine wasting in epidermal tissues (which presumably include microorganisms inhabiting

tissue surfaces as well as those beneath the epidermis and cuticle but excluding body wall and

mutable collagenous tissue-associated microorganisms), we examined temporal progression in *P.

ochraceus* specimens from central California collected on 19 July 2018. Six specimens (mean

mass 290 ± 54 g and ray length 11.2 ± 0.9 cm) were collected from the intertidal zone at

Davenport, CA ($37^o1'19''$N, $122^o12'56''$W) at low tide and transported in insulated coolers to the

Long Marine Laboratory at UC Santa Cruz, where they were housed in flow-through aquaria

(3.81 ± 0.05 mL s^{-1}; mean water residence time 37 min) indoors in individual containers.

Individuals were maintained under these conditions for the duration of the experiment.

Temperature, salinity, and dissolved O_2 were measured daily using a YSI-3000 handheld meter.

After a 48 h acclimation period, a small scar (~ 5 mm long) was made on a single ray using a

sterile 4 mm biopsy punch. Individual asteroids were monitored daily for the presence of lesions

(as defined above). A small tissue sample (~ 3 mm x 2 mm) was taken from the margin of

artificial scars using sterile 4 mm biopsy punches after 24 h. New artificial scars were made on

adjacent rays each day, and sampled after 24h. When lesions away from these artificial scars

were observed (i.e. lesions not caused by physical scarring), their margin tissues were sampled

using the same approach. All tissue samples were placed into sterile 1.2 mL cryovials and

immediately frozen in liquid N_2 or in a -80^{o}C freezer.

To examine wasting in all body wall + epidermal tissues (i.e. biopsy punch), we examined

temporal progression in *P. ochraceus* specimens collected on 24 May 2018 from Davenport, CA

($37°1'19''$N, $122°12'56''$W), which were transported in an insulated container to the Long

Marine Laboratory at UC Santa Cruz where they were placed into a single, flow-through sea

table. The individuals were measured, weighed and photographed to fingerprint apical circlet

pattern. A single biopsy punch (4 mm) was retrieved from each individual and preserved in

liquid N_2. After 312 h, 10 of the 12 had lesions on their surfaces, and after 360 h, all individuals

had lesions. After 360 h, lesions were sampled by biopsy punch and preserved in liquid N_2, and

again at time of death, which occurred between 360 h and 480 h. Because we did not capture the

exact time of lesion occurrence, we restricted analysis of these individuals to samples collected

at initial (T0), 360 h (TI – i.e. shortly after lesions had formed) and at time of death (TF).

Impact of suboxic conditions on wasting in* Asterias forbesi*: To test whether wasting could

result from suboxic conditions, we incubated asteroids in incubations where O_2 was depleted and

examined surface microbiome abundance and composition. Twenty-four *Asterias forbesi* (mass

63 ± 7 g) were obtained from Bar Harbor, ME ($44^o25.7$'N, $68^o12.0$'W) on 29 June 2019 and

transported in insulated coolers to the laboratory at Cornell University. There, the individuals

were initially placed into a single, 320 L aquarium for 24 h, before being placed into individual

flow-through baskets and divided into two treatments. One large volume (230 L) sump

containing artificial seawater (Instant Ocean) was set up for each treatment. One sump served as

a control, while another sump was continuously sparged with medical-grade N_2 ($\sim$ 5-6 L min^{-1})

to lower pO_2. Continuous flow between sump and aquarium systems was maintained by non-

self-priming pumps and removed by gravity through a standpipe. O_2 and temperature was

continuously monitored in the experimental tanks using HOBO O_2 Loggers (U26; Onset).

O_2 was depleted in N_2-sparged sump waters (5.87 ± 0.29 mgL^{-1}) compared to control incubations

(9.62 ± 0.06 mg L^{-1}), representing a mean concentration decrease of ~39%. Individual asteroids

were monitored daily for the gross appearance of lesions on aboral and ventral surfaces. There

was no significant difference in the rate of mass change between treatments, where all

individuals lost on average 30.0 ± 6.5% of the initial mass over the course of the experiment as

they were not fed (Fig. S1). At experiment initiation, surface bacterial abundance samples were

taken and preserved as described above. The mass of individuals was recorded daily; stars

remained without prey during the experiment. Surface microbial swabs were sampled every 48h

following the approach outlined above. To reduce sampling stress on individuals, half of

individuals in each treatment were biopsied (2 x 3 mm biopsy samples collected using sterile

biopsy punches) on their aboral surface at experiment initiation, and then every 5 d until

experiment termination (biopsied specimen lesion genesis time was not significantly different to

non-biopsied specimen lesion genesis time). Biopsy punches (1 each) were preserved in

RNALater or in 10% neutral buffered formalin. Upon appearance of lesions, their margins were

sampled using a 5 mm biopsy punch to scrape a ~ 3 mm x 2 mm tissue sample. Additionally, a 3

mm biopsy punch was used to obtain a sample through body wall tissues on the lesion margins.

Lesion margin tissues were stored at -20^{o}C until analysis.

Bacterial abundance: Bacterial abundance in suboxic and organic matter amendment

experiments was determined by SYBR Gold staining and epifluorescence microscopy (Porter

and Feig, 1980;Noble and Fuhrman, 1998;Shibata et al., 2006). An aliquot (1 mL) of each

sample was first stained with SYBR Gold (2 µl mL^{-1} of the 10,000X stock) for 2 min, then

samples were filtered through 25mm diameter 0.2 µm black cyclopore filters mounted on 25 mm

Type AA Millipore filters to even flow. The filters were removed from the backing filter,

adhered to clean glass slides and mounted in 30 µL of PBS:Glycerol (50:50) containing 0.1% p-

phenylenediamine. The slides were visualized on an Olympus BX-51 epifluorescence

microscope under blue light excitation. Over 200 cells were counted in >10 fields. Bacterial

abundance was calculated by multiplying mean abundance per reticle grid by total grids per filter

area, and divided by volume passed through the filter. For some samples, high background

fluorescence precluded accurate counts and so are not included in downstream analyses.

Microbial assemblage analyses: Microbial assemblages inhabiting body wall samples (i.e.

biopsy punch; wasting in the absence of external stimuli), lesion margins (i.e epidermal scrapes;

wasting in the absence of external stimuli), and at the animal/water interface (i.e. swabs; suboxic

and organic matter enrichment experiments) were examined by 16S rRNA amplicon sequencing.

Nucleic acids were extracted from frozen 3mm biopsies, lesion margin scrapes, and frozen

surface swabs using a Quick-DNA Fungal/Bacterial MiniPrep Kit (Zymo Research, cat# D6005)

according to the manufacturer's protocol. Bacterial DNA was quantified using a Quant-IT

dsDNA Assay (Invitrogen, cat# Q33120) in conjunction with a StepOnePlus™ Real-Time PCR

system (Applied Biosystems). Bacterial community composition was examined via PCR

amplification sequencing of the V4 region of the 16S rRNA gene using a modified version from

Caporaso et al., 2011. Fifty µl PCR reactions contained 1x 5PRIME HotMasterMix (QuantaBio,

cat# 2200400) and 0.1µM each primer (515f/barcoded-806r). Template DNA quantity varied

between experiments. For examining microbiome composition during wasting progression, 1 µl

of extract (containing from below detection limit of 0.1 ng to 80 ng [mean = 17 ng]) was used as

PCR template. For *Asterias forbesi* hypoxia and *Pisaster ochraceus* OM enrichment, 5 pg of

genomic DNA (determined by Femto Bacterial DNA Quantification Kit; Zymo Research, cat#

E2006) was used to standardize prokaryotic template amounts (Hewson, 2019). PCR products

from duplicate reactions were pooled for each sample and cleaned using a Mag-Bind RxnPure

Plus Kit (Omega Bio-tek, cat# M1386-01). Ten ng of bacterial DNA from each sample were

pooled, libraries prepared using the NextFLEX prep, and sequenced on 2 lanes of Illumina

MiSeq (2 x 250 paired end) at the Cornell Biotechnology Resource Center. Sequence libraries

are available at QIITA under studies 12131 and 13061 and at NCBI under BioProject

PRJNA637333.

Raw sequences were uploaded to QIITA and processed using the native 16S rRNA pipeline

(Gonzalez et al., 2018). After demultiplexing, reads were trimmed to 150 bp and sub-OTUs

(sOTUs) were configured using Deblur (Amir et al., 2017). The SEPP phylogenetic tree and

BIOM/FA files were downloaded from the deblur reference hit table and converted into qiime2

(v2019.10) artifacts, after which taxonomy was assigned using the Silva 132 release (Quast et al.,

2012) and q2-feature-classifier plugin. All files were then imported into R (v3.6.1) using qiime2R

v(0.99) (Caporaso et al., 2010) and compiled into a phyloseq (v1.28) (McMurdie and Holmes,

2013) object for downstream analyses.

Amplicon data was transformed using the PhILR (Phylogenetic Isometric Log-Ratio Transform,

v1.1) package for ordination (Silverman et al., 2017). PhILR transforms compositional data (i.e.,

proportional, or relative data) into a new matrix of 'balances' that incorporates phylogenetic

information. Here, a balance is defined as the isometric log-ratio (ILR) between two clades that

share a common node. The ILR is a common tool used in compositional data analysis that

transforms constrained data into an unconstrained space (i.e., Euclidean space), thereby allowing

standard statistical tools to be applied. For each experiment, low abundance sOTUs were filtered

based on sequencing depth/evenness and a pseudocount of 0.65 was applied to all 0 counts. The

ILR analyses in this study do not require an even sampling depth, and therefore, count

normalization techniques like rarefying, which lead to a loss of information (McMurdie and

Holmes, 2014), were not utilized. Phyloseq (1.32.0) (McMurdie and Holmes, 2013) was used to

perform principal coordinate analyses (PCoA) on the Euclidean distances between PhILR

transformed samples, and the adonis function in the vegan package (v2.5-6) (Oksanen et al.,

2019) was used to perform a PERMANOVA on relevant PCoAs. For June 2018 *Pisaster*

ochraceus samples, a principal coordinate analysis based on Weighted Unifrac distances

(Lozupone et al., 2011) was used in lieu of the PhILR transformation due to higher explained

variance from a PERMANOVA. For these samples, low abundant sOTUs were removed and the

data was transformed to an even sampling depth. Multiple filtering strategies were applied to all

analyses and did not affect results.

Balances were used to quantify differential abundance in order to address the issue of

compositionality in amplicon data. Two approaches were used that rely on the ILR

transformation. For comparisons between two categorical variables (diseased/non-diseased tissue

and surface swabs from specimens immediately before lesions appear with earlier specimens), a

sparse logistic regression with an l_1 penalty of $\lambda=0.15$ was applied to PhILR balances using the

glmnet package v(3.0-2) (Friedman et al., 2010). For time-course experiments, the PhyloFactor

package (v0.0.1) was used. PhyloFactor calculates balances in a similar fashion to PhILR, but

instead of using nodes to contrast clades, Phylofactor bisects a phylogenetic tree along its edges

in an iterative manner. Each iteration, or 'factor,' is regressed using a generalized linear model.

Edges were maximized using the F statistic and a Kolmogorov-Smirnov test was used to break

the iterations.

Additional experimental challenges of* Pisaster ochraceus:*** We also sought further evidence of

BLODL by examining wasting in context of water flow rates, desiccation stress, and challenge

with tissue homogenates from a wasting specimen. *P. ochraceus* (n = 24) were collected from

Mitchell Cove, Santa Cruz (36º57.1N, 122º2.51'W), on 25 June 2018 at low tide (mean mass

357 ± 37 g), and transported in a cooler to the Long Marine Laboratory. Metadata on their size

and weight, along with mean flow rates in incubations and change in mass over time is provided

in Table S1. The temperature in aquarium settings for all experiments was measured by Onset

Hobo Spot loggers (n = 2) which were deployed into aquarium outflows for the first 4 days of

the experiment, and after 6 days was measured using a YSI Handheld Instrument.

The impact of water flow on asteroid wasting was examined in 12 specimens. Asteroids (n = 6

each treatment) were placed into individual plastic boxes which were subject to high (7.06 ± 0.39

ml s^{-1}; water residence time in container ~ 20 min) and low (2.84 ± 0.26 ml s^{-1}; water residence

time 50 min) flow-through rates. Temperature and salinity were monitored daily using a

handheld YSI Probe (YSI-3000). Individuals were visually inspected daily for the presence of

body wall lesions and were weighed to determine changes in their overall mass over the course

of the experiment.

The experiments were performed over a 21 day period during which mean water flow-through

temperatures ranged from 14.4 – 17.6°C (Fig. S2), with day-night variation of 1.8 – 1.9°C. Water

temperatures increased mostly between 10 and 15 days of incubation. Salinity did not vary by

more than 0.2 over the course of the experiment. Additionally, we deployed two loggers into the

intertidal zone at Davenport, CA (37°1'19"N, 122°12'56"W) to examine variation between

experimental temperature conditions and conditions experienced by asteroids at the collection

site. In contrast to experimental flow-through systems, field-deployed HOBO loggers revealed

strong changes in temperature accompanying tidal cycles and cycles of immersion/emersion

(Fig. S3). The maximum temperature variation recorded *in situ* occurred during a low tide in the

morning of 29 June, when the temperature swung from from 11.0°C at 6:00am to a maximum of

33.7°C at 10:08am, then back to 13.3°C by 11:00am.

The impacts of desiccation under both high and low flow was examined by first placing 6

individuals onto plastic trays in sunlight for 1 h. During this period, air temperature was 33.8°C

(mean flow-through incubation temperature was 14.5°C). After desiccation for 1 h, individuals

were placed into individual flow-through plastic boxes and monitored per the variable flow rate

experiments described above. Comparison between desiccation and tissue homogenate challenge

(described below) were performed against controls high/low flow as described above.

The effects of challenge with wasting tissue homogenates was examined in 4 individuals. A

single wasting star was collected at Davenport, CA (37°1'19"N, 122°12'56"W) at low tide on 25

June 2018 and transported to the Long Marine Laboratory. Tissue surrounding lesions (~ 2 g

total) was excised using a sterile razor blade, and placed into 40 mL of seawater from the lab's

inflow system. The tissue was then homogenized in a sterilized mortar and pestle for 10 min.

Half of the tissue homogenate was treated with 10,000U of proteinase k (Sigma-Aldrich) and

incubated for 1 h at 37°C. Two *Pisaster ochraceus* were inoculated with 10 mL of crude tissue

homogenate and two *P. ochraceus* inoculated with the proteinase-k treated homogenate by direct

injection into their coelomic cavity. The inoculated asteroids were then placed in individual

plastic flow-through aquaria (under low-flow conditions). Individuals were monitored per the

flow-rate experiments described above.

Comparison of wasting time (lesion genesis) with experimental parameters: We modeled the

time to wasting (i.e. lesion genesis time) across all experiments against available parameters,

which varied between experiments (Table S2). All statistical analyses were performed using

XLStat version 2019.4.1 in Microsoft Excel. Response of lesion time to treatment in OM

addition experiments were examined by least squares mean ANOVA. The relationship between

lesion time and comparison variables was performed by multiple linear regression, forward or

backward selection procedure and change in Akaike's AIC as entry criterion.

Association of wasting susceptibility with rugosity and surface area to volume ratio: We

investigated whether inherent asteroid properties related to wasting susceptibility in context of

BLODL by examining their rugosity (i.e. corrugated-ness), which is a key determinant of

boundary layer extent. Individual, intact whole-animal specimens of asteroids were collected

from several locations (Table S3) and immediately preserved in 20% neutral buffered formalin.

All individuals were transported to the lab at Cornell University. Computed tomography was

performed on whole specimens at the Cornell University Equine Hospital without contrast to

estimate surface area: volume using a Toshiba Aquillon computed tomographic multi-slice

scanner.

The relative rugosity between wasting-affected and less wasting-affected asteroid species was

examined by calculating the ratio of 3D (determined by computed tomography) to 2D (as

calculated below) for each asteroid specimen and comparing between species. Asteroid species

were categorized based on prevalence of wasting (less or not affected = *Dermasterias imbricata*,

Henricia leviuscula, *Patiria miniata*; wasting affected = *Pisaster ochraceus*, *Solaster stimpsoni*,

Pycnopodia helianthoides, *Leptasterias* sp., *Asterias forbesi*, *Orthasterias kohleri*, *Pisaster*

giganteus and *Pisaster brevispinus*) as reported elsewhere (Montecino-Latorre et al., 2016;Bucci

et al., 2017;Miner et al., 2018;Jaffe et al., 2019;Konar et al., 2019). These were likewise

compared between animal volume, surface area:volume and 2D area.

Because the resolution of computed tomography is only 400 µm, which is larger than some

surface features (e.g. papulae), we performed micro-CT (µCT) analyses on one large and one

small individual of several key species after staining for at least 24 h in IKI solution (Fig. S4). X-

Ray µCT data were analyzed using the Avizo version 2019.4 software (ThermoFisher

Scientific). Briefly, 2-D image slices were uploaded and stacked to reconstruct a 3-D volume for

each specimen. A median filter was applied to each 3-D reconstruction to reduce noise and

smooth edges. The volume of interest was isolated from surrounding background using a

thresholding approach. Next, the total volume for each specimen was segmented into 1 cm

sections (each composed of 500 stacked 20 µm slices). After eliminating holes from each 1 cm

sub-volume, the total volume, surface area, and rugosity were determined using the "Label

Analysis" module of the Avizo software. The surface areas reported were calculated by

subtracting the 2-D surface areas of the two, flat end slices from the total 3-D surface area for

each 1-cm segment. The relationship between ray length and surface area was investigated by

linear regression. We first calculated total two-dimensional area for each specimen by taking into

consideration central disc radius and assuming triangular shape of rays, accounting for total

height of central disc and height of ray tip (see Fig. S5). We then used the ratio of total surface

area (determined by CT) to calculate surface area as a measure of rugosity. All image stacks for

micro-CT analysis are available from MorphoSource (Duke University) under project accession

P1047.

Asteroid Specimen Respiration: We hypothesized that specimens with greater respiratory O_2

demand relative to their calculated O_2 flux would be more susceptible to wasting. The respiration

rates of individual asteroids was measured upon experiment initiation for *Asterias forbesi* and

Pisaster ochraceus, and for additional species at the Bodega Marine Laboratory (Table S4).

HOBO O_2 Loggers (U26; Onset) were placed into sealable plastic containers to which

individuals were added. The incubations were circulated using a battery-operated submersible

DC motor and propeller. The containers were then filled by immersion in flow-through seawater,

and sealed, excluding all visible bubbles. Individuals were incubated for 1 – 2 hr in containers

before retrieval of O_2 probe. Respiration rate was calculated by the linear change in O_2

concentration over time in incubations. Respiration rates were compared to calculated maximum

diffusion rates based on overall surface area determined by computed tomography using Fick's

second law of diffusion ($J = -D * \partial C / \partial d$ where J = flux across the membrane, D = diffusivity

constant of O_2 in seawater, C = concentration difference between coelomic fluid and seawater –

in this case assuming completely anoxic coelomic fluid and saturated seawater, and d = thickness

of outer epithelium -assumed to be 20 µm.)

Time series analyses of wasting intensity and chlorophyll a at Whidbey Island: To understand
the relationship between primary producer biomass (chlorophyll a), physico-chemical parameters
(temperature, salinity, dissolved O_2), and occurrence of wasting, we examined data obtained
from the Penn Cove Shellfish data buoy and compared this to observations of wasting frequency
at Coupeville Wharf and Langley Harbor as reported previously (Hewson et al., 2018) from
August 2014 to June 2019 (i.e. 5 years). We also compared wasting frequency with precipitation
data obtained from the National Center for Environmental Information (NOAA), which may be
seen as a proxy for potential terrestrial runoff. We first calculated the mean time of wasting over
the 5-year period and compared this to 5-year mean values of all parameters. We then performed
at-time-of-wasting to prior to wasting comparison following a shifting window approach
comparing the 3-month window immediately before wasting with 3-month windows in earlier
months (Fig. S6).

Stable isotopic signatures in historical wasting asteroid specimens: The natural abundance of
^{15}N and ^{13}C was determined in 71 individual starfish specimens, including 50 individuals
representing paired asymptomatic/wasting affected species at distinct sites and sampling times
which were collected as part of prior work (Hewson et al., 2014) (Table S5). We included an
additional 21 individuals of different species to provide context of stable isotopic composition.
Samples were collected and frozen at -20°C prior to analysis. Thawed tissue samples were
subsectioned for analysis by scraping tube feet into sterile 1.2 ml cryovials. Samples were freeze-
dried at -45°C for one week then ground with mortar and pestle. A subsample of 1 mg of tissue
was encapsulated into tin and subsequently analyzed on a Carlo Erba NC2500 Elemental
Analyzer coupled to a Thermo Scientific Delta V Advantage IRMS (Bremen, Germany).

RESULTS AND DISCUSSION

The results of our work provide support for our hypothesis that sea star wasting is associated
with the formation of anaerobic conditions adjacent to asteroid surfaces. First, we show that
organic matter amendment leads to faster lesion genesis than untreated stars, which is preceded
by increased bacterial abundance in some treatments and the proliferation of copiotrophic taxa
on and above asteroid surfaces in all treatments. In asteroids which wasted in the absence of
external stimuli, epidermal and body wall tissues showed a similar progression of copiotrophic

bacterial orders, and at the time of lesion and until death the proliferation of strict and facultative

anaerobes. Next, we show that lesions form as a consequence of exposure to suboxic water

column conditions. We also demonstrate that wasting is correlated with primary production

trends over 5 years at a field site. Finally, we provide further evidence of predominantly

anaerobic conditions during mass mortality in 2013 – 2014 by way of enriched ^{15}N pools in

affected tissues relative to asymptomatic individuals.

Organic matter amendment stimulates boundary layer microorganisms and results in

rapid wasting

We sought to examine the impact of elevated heterotrophic bacterial respiration on animal

surfaces through amendment with various sources of OM which we hypothesized would fuel

microbial remineralization. We performed laboratory experiments in which *P. ochraceus* was

amended with peptone, *Dunaliella tertiolecta*-derived particulate OM (POM), and coastal

seawater POM and examined their impacts on SSW progression and boundary layer bacterial

abundance and composition. The addition of organic substrates (peptone and *Dunaliella*

tertiolecta-derived POM) induced significantly faster lesion genesis than control incubations

(p=0.012 for peptone and p=0.04 for *Dunaliella*-POM, Student's t-test, df=5), but lesion genesis

time was not significantly different for the addition of coastal-POM (Fig. 2). Collective treatment

temporal pattern of lesion genesis was only significantly different from controls with amendment

with peptone (p = 0.0154, log-rank test, df=5) and *Dunaliella*-POM (p=0.0339, log-rank test,

df=5). Variation in dissolved O_2 in incubations varied over the course of the experiment from 9.6

– 10.2 mg L^{-1} and were never under-saturated. Temperature varied from 12 – 14ºC, but variation

did not correspond with wasting in any treatment.

The boundary layer microbiota of *P. ochraceus* during the organic matter amendment

experiment changed over time in all treatments (Figs. 3E-G; Figs S7), but the most prominent

changes were distinguished by the copiotrophic orders Flavobacteriales and Rhodobacterales

(Flavobacteriales; control: p < 0.001, *Dunaliella*: p = 0.002, peptone: p = 0.001, coastal POM: p

< 0.001, ANOVA fit with a Generalized Linear Model) (Rhodobacterales; control: p < 0.001,

Dunaliella: p < 0.001, peptone: p = 0.010, coastal POM: p < 0.001, ANOVA with a Generalized

Linear Model), which increased uniformly in all incubations, indicating that captivity alone may

stimulate these groups (i.e. containment affect; Fig. 4). We also observed evidence for treatment-

specific proliferation of genera with OM enrichment. Unamended and peptone supplemented *P.*

ochraceus experienced the most consistent change in Flavobacteriales, with both conditions

exhibiting a linear increase in population mean relative to the mean of all other sub-OTUs.

Flavobacteriales in coastal POM-supplemented *P. ochraceus* were elevated from the first to final

timepoints, but were primarily distinguished by a large boom and bust after 96 h (c.f. bacterial

abundance below). This spike was due to an increase in the family *Crocinitomicaceae*, which, in

addition to the family *Flavobacteriaceae*, comprised the majority of Flavobacteriales across all

treatments. Rhodobacterales, which primarily consisted of the family *Rhodobacteraceae*,

increased in all experimental conditions.

Bacterial cell abundance on surfaces (relative to both initial values and controls) illustrated large

swings prior to wasting onset (Fig. 5). Individuals that did not waste over the course of the

experiment maintained abundances of $0.7 – 2.6 \times 10^6$ cells mL^{-1}, which was enriched 53 to

1743% above bacterioplankton abundances in incubation treatments (Fig. 5). On aggregate,

wasting stars had higher bacterial abundances than non-wasting stars (Fig. S8), however the

relationship was not significant because of high variation between treatments with OM.

Treatment bacterial abundances remained no different to controls over the first 48 h of

incubation, but increased relative to controls in peptone and coastal-POM treated asteroids after

72 and 96 h, respectively. However, by 96 h for peptone and 120 h for coastal POM both

amendments had again declined, and remained no different to controls after this time (Fig. 5). In

contrast, bacterial abundance in *Dunaliella*-POM incubations were no different to controls over

the first 48 h of incubation, and were far less than controls after 72 h. Since bacterial abundance

increased prior to lesion genesis in at least two OM treatments, we posit that wasting is

influenced by copiotroph proliferation on animal surfaces. The decrease in bacterial abundance

after initial increase in both peptone, coastal POM, and consistently lower bacterial abundance in

Dunaliella-POM incubations may be evidence of heterotrophic remineralization-fueled O_2

deficit over time on wasting asteroids, similar to the effect observed in our experiments with

Asterias forbesi incubated in hypoxic water (see below). Facultative and strict anaerobes

generally experience slow growth rates compared to aerobic taxa because it is less energetically

efficient to grow on reduced electron acceptors. While standing stock of aquatic bacteria may be

higher in anaerobic conditions than in aerobic conditions, population growth rates are typically

lower (Cole and Pace, 1995).

Shifts in heterotrophic bacterial and archaeal communities during wasting progression in the absence of external stimuli

Because sampling by biopsy punch imparts stress on animals that may elicit wasting, sampling in the absence of external stimuli focused on samples collected at the time of lesion genesis (August 2018) and samples collected after lesion genesis (May – June 2018), and are distinct from samples collected by surface swab (collected in organic matter enrichment experiment). Hence, compositional changes observed in these surveys likely reflect a combination of taxa that change prior to lesion formation and those that degrade tissues.

In the August 2018 study, *Pisaster ochraceus* developed lesions without stimuli beginning 5 d after isolation, and by 8 d more than half of incubated stars were symptomatic (Fig. 6). Lesions formed initially concomitant with an approx. $2^{o}C$ swing in temperature, however continued in other stars as temperatures progressively decreased over the course of the experiment. Initial genesis of lesions was not accompanied by variation in either DO or pH. Lesions were grossly characterized by nonfocal loss of epidermal tissues, which exposed underlying body wall tissues. Lesion margins were not remarkable in terms of coloration which would otherwise indicate melanization. Epidermal samples from these specimens revealed no significant difference in microbial composition between artificial and natural lesions in 3 specimens that wasted (PERMANOVA; $P = 0.139$; Fig. S9). Between initial samples and the time of lesion genesis, most taxa identified as differentially abundant were less abundant sub-OTUs. In the May-June 2018 study, we observed a progressive increase in copiotrophic orders between initial samples and those taken shortly after lesions had formed, including Campylobacterales ($p<0.001$), Flavobacteriales ($p<0.001$) and Vibrionales ($p<0.001$; ANOVA fit with a Generalized Linear Model) (Fig. 3A). This occurred concomitant with an increase in *Nitrosopumilus* and obligate anaerobes (Deltaproteobacteria) relative to a large clade of typically fast-growing phyla (Alpha- and Gammaproteobacteria) (Fig. 3B-D).

Following lesion formation, bacteria on surfaces likely experience a complex milieu of OM molecules, including those from decaying tissues. We found evidence for the proliferation of anaerobic taxa under these conditions by comparing microbial composition between lesion formation and animal death. Between lesion genesis and animal death, we observed a further increase in copiotrophs in body wall (May-June 2018) and epidermal (August 2018) samples,

and a proliferation in microaerophiles (*Arcobacter* spp.), facultative anaerobes (*Moritella* spp.),

and obligate anaerobic families (Clostridia, Fusobacteria and Bacteroidia) at time of death (Fig.

3A; Fig. S10). The two sub-OTUs with the largest F-statistic from regression over the entire

course of wasting from initial samples to animal death belonged to the families

Desulfobulbaceae (p<0.001) and *Desulfovibrionacea* (p=0.002; ANOVA fit with a Generalized

Linear Model). Both of these families are strictly anaerobic sulfate reducers. These results are

consistent with the pattern of microbial assemblage variation observed in the organic matter

amendment experiment. Previous study comparing wasting (specimens already had lesions) and

asymptomatic asteroid-associated community gene transcription also noted the increase in

transcripts from *Propionibacterium*, *Lachnospiraceae* and *Methanosarcina*, which are strict

anaerobes, as well as *Stigmatella* and *Staphylococcus*, which are facultative anaerobes, as a

proportion of total transcripts (Gudenkauf and Hewson, 2015).

Microbial assemblage composition and abundance variation illustrates an increasingly anaerobic environment near asteroid surfaces

Taken together, these studies suggest that lesion formation is associated with a general

proliferation of bacteria, including well-known copiotrophic orders (including *Flavobacteriaceae*

and *Vibrio* spp.) of marine bacteria. Several studies have observed the proliferation of

copiotrophic taxa longitudinally during wasting, including genera within the families

Flavobacteriaceae, Rhodobacteriacaea (Lloyd and Pespeni, 2018), Actinobacteria, and genera in

the orders Altermonadales (Nunez-Pons et al., 2018), Vibrionales and Oceanospiralles (Hoj et

al., 2018). These taxonomic groups are amongst the most active constituents of bacterioplankton

and major players in marine OM degradation, some of which have facultative anaerobic

metabolisms (Pinhassi et al., 2004;Choi et al., 2010;Buchan et al., 2014;Thiele et al.,

2017;Pohlner et al., 2019). While it is tempting to ascribe pathogenicity traits to groups that are

enriched on disease-affected tissues (based on members of the same family or genus causing

pathology), or infer their role in community dysbiosis (i.e. the microbial boundary effect), this is

not possible in the absence of demonstrated pathogenicity or strain-level assignment (Hewson,

2019). The observation of strict anaerobic bacteria in underlying tissues after lesions had formed

suggest the creation of a depleted oxygen environment on asteroid surfaces in response to

organic matter loading.

All aquatic surfaces are coated with a thin film of water (i.e. diffusive boundary layer) that

impedes gas and solute exchange, and, provided aerobic respiration is sufficiently high, suboxic

conditions can form on a surface despite oxygen saturated water circulating above (Jørgensen

and Revsbech, 1985). This may result in the proliferation of facultative and obligate anaerobes

until asteroid death.   Stimulation of bacteria and subsequent O_2 diffusion limitation is well

described in mammalian respiratory systems as well, and is especially pronounced in cystic

fibrosis patients. Heterotrophic bacteria inhabiting mammalian lungs thrive on mucins and

generate biofilms which further restrict O_2 diffusion into tissues. O_2 consumption by biofilms

and by neutrophils may result in hypoxia and reduced diffusion of O_2 across alveolar tissues (Wu

et al., 2018). This in turn leads to the proliferation of anaerobes, which are present in clinically

normal lungs (reviewed in Guilloux et al., 2018) and elevated in diseased lungs (Denner et al.,

2016;Wang et al., 2019;Spence et al., 2020). This phenomenon is also observed in fish gills

(Legrand et al., 2018;Meyer et al., 2019) which are inhabited by copiotrophic and potentially

facultatively anaerobic taxa (Reverter et al., 2017;Rosado et al., 2019).

Rapid mineralization of dissolved OM by bacteria near asteroids has been noted previously in

studies of epidermal amino acid uptake by *Asterias rubrens*, which ultimately led to decreased

animal weight (Siebers, 1979), and may be responsible for very low ambient DOM

concentrations adjacent to asteroids (Siebers, 2015). Remineralization of OM by heterotrophic

microorganisms fuels oxygen consumption, which may in turn lead to oxygen deficit when

consumption is not matched by gas diffusion. For example, excess phytoplankton-fueled

bacterial respiration, caused by eutrophication and enrichment from terrestrial sources and

upwelling zones may result in 'dead zones' (e.g. Mississippi River Plume, Peruvian upwelling

zone, Benguela current; reviewed in (Diaz and Rosenberg, 2008) and may be exacerbated by

seasonal temperature changes (Murphy et al., 2011) and restricted bathymetry (Diaz, 2001). We

posit that OM amendment stimulates bacterial abundance immediately adjacent to asteroid

respiratory surfaces (i.e. within boundary layers) leading to suboxic microzones and ultimately

limiting gas diffusion potential (Gregg et al., 2013).

Bacterial stimulation and enhanced wasting in asteroids is paralleled by the DDAM (dissolved

organic carbon, disease, algae, microorganism) positive feedback loop in tropical corals

(Dinsdale et al., 2008;Barott and Rohwer, 2012;Silveira et al., 2019). Coral disease is associated

with OM enrichment (David et al., 2006;Smith et al., 2006), some of which originates from

sympatric primary producers (Haas et al., 2010;Haas et al., 2011), which in turn are more labile

than OM released from the corals themselves (Haas et al., 2016;Nakajima et al., 2018) and

results in both elevated bacterial abundance on coral surfaces (Dinsdale and Rohwer, 2011;Haas

et al., 2016), and enhanced remineralization rates (Haas et al., 2016). Bacteria at the coral-water

interface have higher energetic demands than those in plankton (Roach et al., 2017), and are

highly adapted to organic carbon availability in their local environment (Kelly et al., 2014). The

spatial scale on which bacteria react to OM is primarily at water-surface interfaces (Brocke et al.,

2015). Hypotheses for the mechanism of coral mortality caused by heterotrophic bacteria include

disruption in the balance between corals and their associated microbiota (David et al., 2006),

introduction of pathogens that have reservoirs on macroalgae (Nugues et al., 2004), or dysbiosis

resulting in invasion by opportunistic pathogens (Barott and Rohwer, 2012). In black band

disease, DOC released from primary production causes micro-zones of hypoxia which result in

production of toxic sulfides, which in turn result in opening of niches for cyanobacteria (Sato et

al., 2017). In asteroid wasting, the proliferation of heterotrophic bacteria and wasting disease

may be due to any of these effects.

Asteroid wasting is induced by suboxic conditions

Our data demonstrate that SSW is induced by suboxic water column conditions. We incubated *A.*

forbesi in suboxic water and observed patterns of wasting progression, boundary layer bacterial

abundance and microbial assemblage β-diversity. Dissolved oxygen (DO) concentrations were

controlled in an aquarium setting by continuous sparging with N_2, which were on average 39%

lower than untreated control incubations (Fig. 7). All individuals remained asymptomatic in

control incubations over the 13 day experiment, while 75% of individuals in hypoxic conditions

developed lesions (mean time to lesion genesis = 9.58 ± 0.89 d; Fig. 7). Development of lesions

over time was strongly related to treatment (p = 0.006, log-rank test, df=12). Bacterial

abundance on animal surfaces (which we define as abundance in surface samples) corrected for

aquarium water values increased in both control and suboxic treatments over the first 6d of

incubation, but by day 13, abundance of bacteria in suboxic treatments was significantly lower

(p<0.001, Student's t-test, df=12) on suboxic treated individuals than in control individuals (Fig.

S11). *Asterias forbesi* treated with suboxic waters demonstrated consistent shifts in microbial

communities with treatment (Fig. S9). However, no single bacterial taxonomic organization

strongly differentiated normoxic from suboxic conditions.

Further evidence for the role of oxygen in SSW was observed in experiments comparing wasting

speed under variable incubation flow rates (i.e. water replenishment rates). While we did nt

measure DO concentrations in these experiments, higher flow rates likely had higher DO

concentrations than lower flow rates, in addition to reducing OM (e.g. mucus) and toxic exudates

in animal waste (notably NH_3 [Propp et al., 1983] and S^- [Vistisen and Vismann, 1997]).

Furthermore, higher flow rates may have experienced less extensive boundary layers than lower

flow rates (Fonseca and Kenworthy, 1987). The time to lesion genesis in *Pisaster ochraceus*

was faster for asteroids under low-flow conditions than those under high flow conditions (p =

0.006, Student's t-test, df=3), however survival was not significantly different between flows

(log-rank test, ns; Table S6). Desiccation, which was used to insult asteroids and simulate

emersion during low tide events in warmer temperatures, resulted in faster lesion formation (p =

0.05, Student's t-test, df=3) under low flow conditions, but not different under high flow

conditions (log-rank test, ns; Fig. 8). Addition of tissue homogenates, which was initially used to

assess transmissibility of tissue-derived agents, resulted in faster wasting than addition of

proteinase k treated tissue homogenates (p = 0.04, Student's t-test, df=3); however, survival was

no different between control treatment low flow and the addition of proteinase K-treated or

untreated tissue homogenates (log rank test, ns; Fig. 8).

The mechanism by which asteroids are particularly sensitive to ambient O_2 concentrations is not

well constrained by empirical studies, especially as it relates to SSW. Asteroids mostly rely on

passive respiration (c.f. ventilation) and gas diffusion across outer membranes to meet

respiratory demand, a point illustrated by mass mortality events of benthic invertebrates,

including asteroids, correlated to low O_2 conditions (reviewed in Diaz and Rosenberg,

1995;Levin, 2003;Levin et al., 2009). Together, these data point to significant influence of O_2

conditions on asteroid wasting. While water column hypoxia events were not observed in concert

with SSW in 2013 and beyond, spatially localized hypoxia may occur near surfaces experiencing

limited hydrodynamic flow (Gregg et al., 2013).

Analysis of covariance revealed that lesion time was explained best by different parameters,

depending on the experiment. For *Asterias forbesi*, lesion genesis time variation was best

explained by both overall animal mass change during the experiment and initial bacterial

abundance (p = 0.12), while in *Pisaster ochraceus* experiments, both OM addition and combined

with flow rate was best explained by initial animal mass (p=0.006). In OM addition, variation in

lesion genesis time was also explained by change in bacterial abundance over the first 3 d of

incubation (p= 0.018) (Table S7). While these observations are entirely correlative – in other

words, cannot reveal causality – they further indicate a key role of heterotrophic bacteria

inhabiting the asteroid-seawater interface in wasting.

Wasting is related to inherent asteroid properties that dictate boundary layer extent, gas

diffusion, and respiration

Inter- and intra-species susceptibility to asteroid wasting is extensively recorded in previous

study, including a significant and positive relationship between individual size and wasting

(Hewson et al., 2014), and shifts in size structure after wasting from larger to smaller individuals

of *Pisaster ochraceus*, which was believed to be due to recruitment of juveniles (Bates et al.,

2009;Eisenlord et al., 2016;Menge et al., 2016;Kay et al., 2019). Wasting in 2013-2014 affected

> 20 species of asteroid (Hewson et al., 2014), however the magnitude of SSW impact varied

between species. Comparison of community structure before and after wasting suggests inter-

species variability in wasting mortality. Asteriid taxa (*Pycnopodia helianthoides, Pisaster* spp.,

and *Evasterias troschelii*) experienced considerable declines in the Salish Sea (Montecino-

Latorre et al., 2016;Schultz et al., 2016) and Southeast Alaska (Konar et al., 2019), while

Dermasterias imbricata maintained or increased in abundance after mass mortality (Eckert et al.,

1999;Montecino-Latorre et al., 2016;Schultz et al., 2016;Konar et al., 2019). In the Channel

Islands, SSW disproportionately affected Asteriid taxa relative to *D. imbricata* and *Patiria*

miniata (Eckert et al., 1999). Inter-species differences in wasting intensity have been noted in

citizen science data accumulated by MARINe (Miner et al., 2018). The potential causes of inter-

and within-species wasting susceptibility remain poorly constrained.

We hypothesized that wasting susceptibility may relate to both inter-species variation in rugosity

(i.e. degree of corrugation), which dictates diffusive boundary layer thickness, and intra-species

surface area-to-volume ratio, which determines total gas flux potential, which are ultimately

reflected in patterns of population change since 2013 (Eckert et al., 1999;Montecino-Latorre et

al., 2016). Mean and turbulent flow structure around aquatic animals and plants relates to the

mean height, density and shape of structures as they compare to flat surfaces (Koch, 1994;Nepf,

2011;Brodersen et al., 2015). Asteroid surfaces bear numerous spines and processes, including

papulae, spines, paxillae and pedicellaria. These structures impart rugosity and thus generate

diffusive boundary layers proportional to their relative height under both mean and turbulent

flow. For example, the boundary layer height above the urchin *Evechinus chloroticus.* can be 4-5

mm under low (1.5 cm s^{-1}) flow conditions, which was approximately 2 – 6 X greater than

sympatric macroalgae (Hurd et al., 2011). We speculate that more extensive boundary layers

may result in a greater deficit in O_2 due to entrapment of OM adjacent to animal tissues, and also

increase the potential for hypoxia at the animal surface since the effective distance over which

O_2 must diffuse is higher in specimens with greater boundary layer extent. Direct measurement

of oxygen concentration in boundary layers as they relate to bacterial remineralization are

precluded by the sensitivity of instruments (e.g. microelectrodes) to physical damage in non-

immobilized specimens.

To explore the relationship between species rugosity and wasting susceptibility, we examined

specimens of similar size (n = 26 individual specimens) representing wasting-affected (n = 3)

and less/not affected species (n = 5) using whole-animal computed tomography. CT-derived

volume was significantly and positively correlated to overall mass across all specimens (p =

0.00001, R^2=0.9999). To estimate overall surface area of specimens for which respirometry was

measured (which were not imaged by CT), we modeled the surface area to ray length, and found

it followed an exponential function (mean ray length = $0.0825e^{1.0536*LOG(Surface area)}$, $R^2 = 0.93$.

Log (Surface area) was significantly and linearly correlated to Log (Volume; LOG(Volume) =

0.7319*LOG(Surface Area) + 0.7662; $R^2 = 0.9703$); Fig. S20). The surface area: volume was

significantly and negatively correlated with a logarithmic function defined as SA:Vol = $4.6242e^{-0.549[LOG(Volume)]}$) (Fig. S12). The mean rugosity (defined as 3D:2D surface area) was significantly

(p = 0.015, Student's t-test, df = 14) lower in less affected species than more affected species

(Fig. 9). Surface area:volume, individual specimen mass, and overall surface area were not

significantly different between categories among similarly-sized animals. Because analysis of

large animal specimens is limited to a resolution of 400 µm (which is potentially larger than fine-

scale features, e.g. papulae on echinoderm surfaces), we performed further analysis on rays of a

subset (n = 16) of individuals using micro-computed tomography, which has a resolution of 20

µm. The rugosity of wasting-affected taxa was significantly (p = 0.0002, Student's t-test, df = 4)

greater than less-wasting affected species (Fig. 9). Our observation that more rugose species

were more affected by wasting supports the idea that these individuals may be more susceptible

because of their greater extent (physical distance) of diffusive boundary layers on respiratory

surfaces.

Much of the intra-species wasting susceptibility may also be explained by inherent variation in

diffusive flux potential. We observed a significant and positive relationship between wasting

lesion genesis rate and initial animal mass (p = 0.006; analysis of covariance). Larger individuals

have a much lower surface area:volume ratio, where surface area is related to gas flux potential.

Under near-surface hypoxic conditions, or when diffusion is impeded by extensive boundary

layers, larger individuals are more strongly affected than smaller individuals. We also posit that

these observations are the result of more extensive boundary layer height above larger

specimens. It is also important to note that ossicle density varies between species (Blowes et al.,

2017), and those taxa with lower densities (e.g. *Pycnopodia helianthoides*) were more affected

than those with higher densities (e.g. *Pisaster ochraceus*). Species with lower ossicle densities

may be differentially susceptible to wasting since their structure may be broken down faster by

microbial decomposition or apoptotic processes.

Wasting risk susceptibility may furthermore result from differential diffusive flux potential

compared to respiratory demand. We measured the respiration rate (i.e. oxygen demand) of

individuals at the start of each experiment, as well as in individuals of several species that were

both affected by wasting and those that were less or not affected by wasting that were not a part

of experiments to explore whether susceptibility was related to oxygen demand. Mass-

normalized measured respiration rates of asteroids were greatest for *Asterias forbesi*, and least

for *Dermasterias imbricata* and *Patiria miniata* (Fig. S4). Both *Pisaster ochraceus* and *Asterias*

forbesi respiration rates were considerably more than for other specimens. Measured respiration

rates for entire animals was compared to theoretical maximum diffusion rates into coelomic

fluids (hereafter abbreviated RR:TD). RR:TD was greatest in *Asterias forbesi* and *Pisaster*

ochraceus (which were both >1 in most specimens) and least in *Patiria miniata* and

Dermasterias imbricata (which were always < 0.1). The observed RR:TD corresponds with

wasting susceptibility. Perturbation of O_2 availability in animal surface boundary layers may

skew diffusive flux by elongating diffusive path length or reducing differences in O_2 between

tissues and surrounding seawater. Hence, specimens with a higher RR:TD may be more affected

by the condition than those with lower RR:TD. We cannot account for variable permeability of

outer epidermis between individuals (not measured), and assume that all surface area of asteroids

is involved in respiration (which may be over-estimated, since presumably some component of

this area comprises mineral structures). Some asteroid species inhabiting typically suboxic

environments employ morphological and behavioral strategies to meet O_2 demand, including

nidamental cavities (Johansen and Petersen, 1971;Nance, 1981), cribiform organs (Shick et al.,

1981), epiproctoral cones (Shick, 1976), active ventilation of burrows and decreased size of

internal organs (Mironov et al., 2016). However, it is unlikely asteroids typically occurring in

normoxic intertidal or subtidal conditions have the ability to morphologically adapt to hypoxic

conditions.

Potential sources of OM fueling BLODL

Heterotrophic bacteria in marine environments remineralize OM that originates from

autochthonous and allochthonous sources (Ducklow, 1983;Benner et al., 1992;Amon and

Benner, 1996). We hypothesize there are two primary sources fueling BLODL: OM from

primary production (phytoplankton and macroalgae), and OM from decaying asteroids. Most

asteroid wasting is reported in late fall or summer, with fewer reports during other times of the

year (Eckert et al., 1999;Bates et al., 2009;Eisenlord et al., 2016;Menge et al., 2016;Montecino-

Latorre et al., 2016;Hewson et al., 2018;Miner et al., 2018;Harvell et al., 2019;Hewson et al.,

2019). Among the myriad of OM sources in seawater, phytoplankton-derived OM are highly

labile (Ochiai et al., 1980;Ogawa and Tanoue, 2003;Thornton, 2014). We propose that wasting is

associated with peak or post-peak declines in phytoplankton production in overlying waters,

which subsequently results in peak dissolved OM availability. The mean time of wasting mass

mortality observed at Langley and Coupeville, Whidbey Island between 2014 and 2019 fell at or

within 1 month after the mean annual maximum of chlorophyll a, minimum DO concentration,

maximum temperature, and minimum rainfall (Fig. 10). Multiple linear regression (stepwise,

backwards selection criteria) revealed a significant model (R^2=0.866; p=0.001) where

temperature (p = 0.006), chlorophyll a (p = 0.027) and salinity (p = 0.044) explained most

variation in wasting mass mortality, while forward selection (R^2=0.774; p=0.0002) revealed that

monthly variation in wasting was significantly explained by DO alone. Mass mortality was

significantly related (one-way ANOVA, p<0.0001) to elevated chlorophyll in the previous 3

months relative to non-mass mortality months, to elevated salinity, and reduced rainfall (Fig.

S6). Because wasting occurs seasonally in late summer and early autumn, correlation between

these parameters alone does not necessarily indicate a direct link between primary production

and wasting. However, since we also observed that asteroids challenged with a phytoplankton

(*Dunalella tertiolecta*) formed lesions faster than unamended controls, it is entirely possible that

wasting is related to phytoplankton-derived DOM.

The coherence of wasting with primary production in the Salish Sea raises the question of why

wasting mass mortality in the northeast Pacific occurred in the 12 month period following June

2013, especially when asteroids normally persist at sites experiencing very high phytoplankton

biomass and only experienced wasting in 2014 (e.g. Cape Perpetua, OR; Leslie et al., 2005).

Suchy et al. (2019) observed a prolonged (10 month) period of decreased water column

stratification, concomitant with strong predominately southerly winds in fall 2013 and spring

2014 in the northern Strait of Georgia. Chlorophyll a concentrations in the region was also higher

in the region in late 2013 compared to the previous 8 years (Suchy et al., 2019) and the 1981-

2010 average (Moore et al., 2014) and was marked by a significant peak in late August (later

than previous years (Moore et al., 2014), concomitant with the lowest NH_3 concentration

measured over several years at the Seattle Aquarium, presumably a consequence of

phytoplankton uptake (Olsen et al. 2016). Mean monthly precipitation was anomalously lower

in mid-summer compared to 2005-2012 means, but then increased dramatically in late

September 2013, prior to wasting onset in October (Hewson et al., 2018).

Elsewhere, there is evidence that wasting in 2013 – 2014 was tied to elevated primary

production. The high pCO_2 but low temperature-wasting positive relationship noted in Oregon

indicates that upwelling may have stimulated primary production at this location (Menge et al.,

2016). The CALCoFi program observed highest coastal upwelling on record in 2013 in central

California during wasting onset (Leising et al., 2014). These observations suggest that primary

production intensity and timing in 2013-2014 departed from inter-annual variation in prior years,

and has followed seasonal patterns since 2014. The discontinuous latitudinal emergence of

wasting in 2013-2014 and regional apparent longshore sequence of SSW occurrence is consistent

with regional and basin-scale patterns of organic matter availability. The spatial scale of

phytoplankton blooms sustained solely by terrestrial runoff and groundwater discharge ranges

from 880-3600 km^2 in the Southern California Bight (Santoro et al., 2010). Assuming these

blooms are constrained within 10 km of shore, the areal extent of phytoplankton-derived organic

matter inputs is well within the reported longshore spread of SSW (Hewson et al., 2014).

Upwelling, on the other hand, may affect wider coastal productivity patterns. In 2013, strong

upwelling was recorded between 36ºN and 48ºN (i.e. 1,332 km). It is interesting to note that

mass mortality in *Heliaster kubiniji* in the Gulf of California occurred during a prolonged period

of heavy rainfall and elevated temperatures prior to El Niño (Dungan et al., 1982). Such rainfall

may have caused elevated terrestrial discharge, which in turn may have fueled primary

production.

Another potential source of organic matter fueling BLODL is macroalgal detritus (Krumhansl

and Scheibling, 2012) and exudates (Abdullah and Fredriksen, 2004). Macroalgae experience

seasonal increases in biomass during spring and fall due to elevated temperature and elevated

nutrient conditions, but may also experience nutrient limitation in summer (Brown et al., 1997).

For example, a study by (Van Alstyne, 2016) found that *Ulva lactuca* abundance was greatest in

July when compared to both May and September in Penn Cove. Interestingly the author found

that the primary source of nutrients for algal growth during the summer and fall was water from

a nearby river, as well as wastewater effluent from a facility at Coupeville (Van Alstyne, 2016).

Laminaria hyperborean in the North Atlantic Ocean has a pronounced seasonal productivity

cycle including an active growing season from February through May during which previous

year lamina are shed, followed by a non-growing season until November (Kain, 1979).

Exudation of dissolved OM is greatest during *L. hyperborean*'s non-growing season (Abdullah

and Fredriksen, 2004). Both macroalgal detritus (Robinson et al., 1982) and exuded OM (Zhang

and Wang, 2017) are highly labile and rapidly assimilated by bacteria. While there have been no

studies of seasonal exudation or detrital release in the regions affected by SSW, the seasonal

variation of SSW and coherence with OM production and detrital breakdown (Krumhansl and

Scheibling, 2012) warrants further investigation.

OM originating from decaying asteroids may also generate BLODL. Experimental challenge

with asteroid tissue homogenates in this study (Fig. 8), and reported previously (Hewson et al.,

2014;Bucci et al., 2017), suggest that wasting may also be associated with decomposition of

nearby asteroid individuals via assimilation of tissue-derived compounds and subsequent

BLODL. We previously isolated heterotrophic bacteria using sea star tissue homogenates as

nutritional source (Hewson et al., 2018). These bacteria include well-known copiotrophic genera.

Enrichment of near-benthic OM pools by wasting-affected individuals may have resulted in the

apparent density dependence of wasting observed in 2014 in some populations (Hewson et al.,

2014). Indeed, challenge with tissue homogenates by direct injection into coelomic cavities

likely enriches within-and near animal organic matter pools, which in turn may stimulate

heterotrophic remineralization. Hence, challenge experiments, such as those performed

previously (Hewson et al., 2014;Bucci et al., 2017) and in this study, may be a consequence of

BLODL induced by organic matter availability (and possibly protein-bearing material). The

apparent transmissibility of SSW in field sites is based on observations of density dependence at

some sites, along with geographic spread between adjacent sites and through public aquaria

intake pipes (Hewson et al. 2014). These observations may be inaccurately ascribed to

transmissible pathogenic microorganisms, since they may also be explained by enrichment of

surrounding habitats and through intake pipes of organic matter pools from decaying individuals.

Wasted asteroids in 2013-2017 bore stable isotopic signatures of anaerobic processes

Because wasting has no pathognomic signs and has been reported for over a century (reviewed in

Hewson et al., 2019), an obvious question is whether BLODL was related to asteroid mass

mortality observed from 2013. While retrospective analyses of O_2 status of asteroids during this

event is not possible, hypoxic conditions impart elemental signatures in tissues of preserved

specimens. We examined the natural abundance of stable isotopes comparing wasting-affected

and grossly normal individuals at the same location and time, in 2013 and 2014. The natural

abundance of 15N ($\delta^{15}N$) and $\delta^{13}C$ varied between species, with highest values for *Hippasteria*

spinosa and lowest for *Pteraster tesselatus* (Fig. S13). There was no correspondence between

known diet of asteroids (and, hence, food web position) and relative stable isotope composition

between species. The elemental composition of asteroids, like all animals, largely reflects

nutritional source, who obtain anabolic material from consumed prey. Furthermore, asteroids

may take up DOM directly from the water column and use these materials for soft body parts,

like tube feet (Ferguson, 1967b;a). The half-life of isotopic signatures in tissues relates to tissue

turnover and is most stable in ectotherms (Vander Zanden et al., 2015). Dissimilatory anaerobic

nitrogen cycling processes, such as denitrification, shift the balance between ^{15}N and ^{14}N (i.e.

selecting against ^{15}N), resulting in higher $\delta^{15}N$ (ratio of tissue ^{15}N to atmospheric ^{15}N) in

environments. Thus, we restricted our analysis of tissue $\delta^{15}N$ to fast-growing, regenerative tube

feet which will therefore reflect the most recent environmental conditions prior to collection.

Wasting asteroids (including *Pisaster ochraceus*, *Pycnopodia helianthoides*, and *Evasterias*

troschelii), had generally higher $\delta^{15}N$ in their tissues than asymptomatic tissues at the same site

within-species (ns) except for *Leptasterias* sp., which had significantly lower $\delta^{15}N$ in wasting

tissues than in asymptomatic individuals (Fig. 11). On average, $\delta^{15}N$ was enriched by 3.9 ± 3.3

% for each species (7.0 ± 2.8 % excluding *Leptasterias* sp.) in wasted compared to asymptomatic

stars. Ellipse analysis, which can be used to infer isotopic niches or metabolic differences

between populations (Jackson et al., 2011) suggested that in all paired site-species comparisons

wasted stars have altered C and N metabolisms compared to asymptomatic individuals (Fig.

S14).

Translocation of consumed elements to growing tissues is accomplished through continual flux

from digestive glands to these tissues through coelomic fluid (Ferguson, 1964). Internal tissues

of asteroids are inhabited by a suite of bacteria and archaea (Jackson et al., 2018) including

abundant spirochaetes (Holland and Nealson, 1978;Kelly et al., 1995;Kelly and McKenzie,

1995;Nakagawa et al., 2017). Hence, asteroid tube feet tissues, which are distal from digestive

glands, may be influenced by heterotrophic microbial activities which enrich for ^{15}N over ^{14}N.

Our finding of higher $\delta^{15}N$ in most wasted asteroids supports the hypothesis that wasting is

associated with enhanced anaerobic dissimilatory respiration of nitrogen species, perhaps during

translocation of materials between organs or tissues within asteroids, (Ferguson, 1964) or during

uptake of enriched ^{15}N in DOM pools surrounding affected asteroids (Ferguson, 1967b;a).

To the best of our knowledge, there has only been one previous report on the effects of hypoxia

on stable isotopic composition in animal tissues. Oysters affected by hypoxic water conditions

demonstrated $\delta^{15}N$ enrichment, which they propose was due to hypoxia-induced starvation

responses resulting in recycling of internal tissues (Patterson and Carmichael, 2018). Asteroids

likewise have similar autophagous responses to starvation, prioritizing somatic maintenance over

reproduction (AQUINAS and P., 1976;Harrold and Pearse, 1980). Under typical food

availability, reproductive and digestive tissues demonstrate inverse relationships in overall size

relating to spawning and feeding time in most asteroid species. However, the ratio between

reproductive and digestive tissues in *Leptasterias* spp. is synchronous over time in females (but

not so in males), which is different from other starfish species (Menge, 1975). We speculate that

the lower $\delta^{15}N$ observed in wasting *Leptasterias* spp., an opposite trend to other species, may

relate to timing of autophagous transfer of materials within individuals and timing of predicted

hypoxia (which peaks in late summer) relative to autophagy within animals. It is also possible

that asymptomatic and wasting affected specimens were different species of *Leptasterias* sp.

since they form a cryptic species complex (Melroy et al., 2017), which may affect comparison

between disease states.

Further evidence for BLODL association with wasting

Wasting imparts transcriptional and population genetic changes in asteroids and surviving

populations, respectively. In a metatranscriptomic studies comparing gene expression between

wasting and asymptomatic individuals, the relative transcription of high affinity cytochrome c

oxidase (ccb3; Preisig et al., 1996) was higher in symptomatic individuals (Gudenkauf and

Hewson, 2015). Furthermore, cytochrome P450 2J6, which plays a dual role in both oxidation

and detoxification of H_2S (Tobler et al., 2014), was expressed in at least two studies of wasting

asteroids (Fuess et al., 2015;Gudenkauf and Hewson, 2015). Surviving juvenile recruits are

genetically distinct to asteroids before 2013 (Schiebelhut et al., 2018). Loci selected for in

surviving populations correspond to those heightened in experiments with elevated temperature

(Ruiz-Ramos et al., 2020). In particular, Ruiz-Ramos et al (2020) found a synchronous decrease

in expression of ND5 (NADH dehydrogenase 5) among field-wasting specimens and those

subject to temperature challenge in aquaria, and corresponding mutation in ND5 in surviving

populations. Extracellular hypoxia causes downregulation of NADH dehydrogenase in vertebrate

cells (Piruat and López-Barneo, 2005), and variation in mt ND5 genes is related to hypoxia

sensitivity in humans (Sharma et al., 2019). Elevated temperatures may reduce overall O_2

concentrations in seawater and cause faster microbial growth rates. Hence, previous observations

of enhanced temperature corresponding to wasting (Eisenlord et al., 2016;Kohl et al.,

2016;Montecino-Latorre et al., 2016;Miner et al., 2018;Harvell et al., 2019) and with periodic

temperature excursion frequency (Aalto et al., 2020) are consistent with the BLODL model

proposed in our work.

Conclusion

Here we present evidence to support our hypothesis that wasting is a sequela of BLODL. We

provide evidence that this condition may relate to bacterial abundance/compositional shifts on

asteroid respiratory surfaces, and that this results from enrichment with OM. While we cannot

definitiely identify a specific source of OM that may lead to BLODL, inter-annual wasting in the

field corresponds with phytoplankton biomass, and in controlled experiments we demonstrate

that algal-derived OM stimulates wasting. We also provide evidence for this effect as occurring

in specimens from the 2013 – 2014 mass mortality event. BLODL may be exacerbated under

warmer ocean conditions, or conditions in which labile OM from terrestrial sources (which may

include anthropogenic nutrient pollution) may be present in coastal environments. Holothurian

wasting, bearing similarity to asteroid wasting in gross disease signs, was reported in the Puget

Sound and southeast Alaska beginning in 2014 concomitant with asteroid mass mortality

(Hewson et al., 2020) suggesting that this phenomena may affect sympatric benthic

invertebrates. Most urchin diseases are associated with diverse bacteria capable of anaerobic

metabolism (reviewed in Hewson, 2019). Hence, BLODL may help explain the variation in

etiologies observed between echinoderms and between other invertebrate groups, especially

those that rely on diffusion for respiratory activities.

ACKNOWLEDGEMENTS

The authors are grateful to Jim Nagel (Penn Cove Shellfish Company), Karl Menard (Bodega

Bay Marine Laboratory), Taylor White (UC Santa Cruz), Joe Gaydos, Lizzy Ashley (Seadoc

Society), Martin Haulena (Vancouver Aquarium), Lesanna Lahner (Minnesota Zoo) and Kipp

Quinby for provision of specimens and data; Betsy Steele (UC Santa Cruz), Joel Markis and

Marnie Chapman (University of Alaska Southeast) for use of laboratory space; Kim Sparks and

Elliot Jackson (Cornell University) for laboratory assistance; Mary Sewell (U Auckland),

Christopher Mah (Smithsonian); Tom Mumford (University of Washington) for helpful

conversations on macroalgae; and Thierry Work (USGS) for assistance with experiments and

comments on an early manuscript draft. Asteroid collections were performed under permits CF-

19-107 from the Alaska Department of Fish and Game, 19-149a from the Washington State

Department of Fish and Wildlife and SC-13144 from the California Department of Fish and

Wildlife. This manuscript has been released as a pre-print at biorxiv (Aquino et al., 2020). This

work was supported by US National Science Foundation Grants OCE-1537111 and OCE-

1737127 awarded to IH and USGS Contract G19AC00434 awarded to IH and T. Work. The

authors declare no conflict of interest in publication of this work.

Author Contributions: LMS and IH designed research; CA, RMB, CMD, JK, IRP, PR, JER,

LMS, JPS and IH performed research; CA, RMB, CMD, IRP, PR, JER, LMS, JPS, JPW and IH

conducted review and wrote the manuscript; PR and IH provided funding for the research.

References

Aalto, E.A., Lafferty, K.D., Sokolow, S.H., Grewelle, R.E., Ben-Horin, T., Boch, C.A.,

Raimondi, P.T., Bograd, S.J., Hazen, E.L., Jacox, M.G., Micheli, F., and De Leo, G.A.

(2020). Models with environmental drivers offer a plausible mechanism for the rapid

spread of infectious disease outbreaks in marine organisms. *Sci Rep* 10**,** 5975.

Abdullah, M.I., and Fredriksen, S. (2004). Production, respiration and exudation of dissolved

organic matter by the kelp *Laminaria hyperborea* along the west coast of Norway. *J Mar

Biol Assoc UK* 84**,** 887-894.

Amir, A., Mcdonald, D., Navas-Molina, J.A., Kopylova, E., Morton, J.T., Zech Xu, Z., Kightley,

E.P., Thompson, L.R., Hyde, E.R., Gonzalez, A., and Knight, R. (2017). Deblur Rapidly

Resolves Single-Nucleotide Community Sequence Patterns. *mSystems* 2**,** e00191-00116.

Amon, R.M.W., and Benner, R. (1996). Bacterial utilization of different size classes of dissolved

organic matter. *Limnol Oceanogr* 41**,** 41-51.

Aquinas, S.M., and P., N.O. (1976). Histochemical changes in gonadal nutrient reserves

correlated with nutrition in the sea stars, *Pisaster ochraceus* and *Patiria miniata*. *Biol

Bull* 151**,** 357-369.

Aquino, C.A., Besemer, R.M., Derito, C.M., Kocian, J., Porter, I.R., Raimondi, P.T., Rede, J.E.,

Schiebelhut, L.M., Sparks, J.P., Wares, J.P., and Hewson, I. (2020). Evidence for

boundary layer oxygen diffusion limitation as a key driver of asteroid wasting. *bioRxiv*,
doi: https://doi.org/10.1101/2020.1107.1131.231365

Barott, K.L., and Rohwer, F.L. (2012). Unseen players shape benthic competition on coral reefs.
Trend Microbiol 20**,** 621-628.

Bates, A.E., Hilton, B.J., and Harley, C.D.G. (2009). Effects of temperature, season and locality
on wasting disease in the keystone predatory sea star *Pisaster ochraceus*. *Dis Aquat Org*
86**,** 245-251.

Benner, R., Pakulski, J.D., Mccarthy, M., Hedges, J.I., and Hatcher, P.G. (1992). Bulk chemical
characteristics of dissolved organic matter in the ocean. *Science* 255**,** 1561-1564.

Blowes, L.M., Egertová, M., Liu, Y., Davis, G.R., Terrill, N.J., Gupta, H.S., and Elphick, M.R.
(2017). Body wall structure in the starfish *Asterias rubens*. *J Anatom* 231**,** 325-341.

Brocke, H.J., Polerecky, L., De Beer, D., Weber, M., Claudet, J., and Nugues, M.M. (2015).
Organic matter degradation drives benthic cyanobacterial mat abundance on Caribbean
coral reefs. *PLoS One* 10**,** e0125445.

Brodersen, K., Lichtenberg, M., Paz, L.-C., and Kühl, M. (2015). Epiphyte-cover on seagrass
(*Zostera marina* L.) leaves impedes plant performance and radial O_2 loss from the below-
ground tissue. *Front Mar Sci* 2.

Brown, M.T., Nyman, M.A., Keogh, J.A., and Chin, N.K.M. (1997). Seasonal growth of the
giant kelp *Macrocystis pyrifera* in New Zealand. *Mar Biol* 129**,** 417-424.

Bucci, C., Francoeur, M., Mcgreal, J., Smolowitz, R., Zazueta-Novoa, V., Wessel, G.M., and
Gomez-Chiarri, M. (2017). Sea Star Wasting Disease in *Asterias forbesi* along the
Atlantic Coast of North America. *PLoS One* 12**,** 20.

Buchan, A., Lecleir, G.R., Gulvik, C.A., and González, J.M. (2014). Master recyclers: features
and functions of bacteria associated with phytoplankton blooms. *Nat Rev Microbiol* 12**,**
686-698.

Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K.,
Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights,
D., Koenig, J.E., Ley, R.E., Lozupone, C.A., Mcdonald, D., Muegge, B.D., Pirrung, M.,
Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T.,
Zaneveld, J., and Knight, R. (2010). QIIME allows analysis of high-throughput
community sequencing data. *Nat Method* 7**,** 335-336.

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J.,
Fierer, N., and Knight, R. (2011). Global patterns of 16S rRNA diversity at a depth of
millions of sequences per sample. *Proc Nat Acad Sci USA* 108**,** 4516-4522.

Choi, E.J., Kwon, H.C., Sohn, Y.C., and Yang, H.O. (2010). *Kistimonas asteriae* gen. nov., sp.
nov., a gammaproteobacterium isolated from *Asterias amurensis*. *Internat J System Evol*
Microbiol 60**,** 938-943.

Cole, J.J., and Pace, M.L. (1995). Bacterial secondary production in oxic and anoxic freshwaters.
Limnol Oceanogr 40**,** 1019-1027.

David, I.K., Neilan, M.K., Mya, B., Nancy, K., and Forest, R. (2006). Role of elevated organic
carbon levels and microbial activity in coral mortality. *Mar Ecol Progr Ser* 314**,** 119-125.

Denner, D.R., Sangwan, N., Becker, J.B., Hogarth, D.K., Oldham, J., Castillo, J., Sperling, A.I.,
Solway, J., Naureckas, E.T., Gilbert, J.A., and White, S.R. (2016). Corticosteroid therapy
and airflow obstruction influence the bronchial microbiome, which is distinct from that of
bronchoalveolar lavage in asthmatic airways. *J Allerg Clin Immunol* 137**,** 1398-+.

Diaz, R.J. (2001). Overview of hypoxia around the world. *J Environ Qual* 30**,** 275-281.

Diaz, R.J., and Rosenberg, R. (1995). Marine benthic hypoxia: A review of its ecological effects
and the behavioural responses of benthic macrofauna. *Oceanogr Mar Biol Ann Rev* 33**,**
245-303.

Diaz, R.J., and Rosenberg, R. (2008). Spreading dead zones and consequences for marine
ecosystems. *Science* 321**,** 926-929.

Dinsdale, E.A., Pantos, O., Smriga, S., Edwards, R.A., Angly, F., Wegley, L., Hatay, M., Hall,
D., Brown, E., Haynes, M., Krause, L., Sala, E., Sandin, S.A., Thurber, R.V., Willis,
B.L., Azam, F., Knowlton, N., and Rohwer, F. (2008). Microbial ecology of four coral
atolls in the northern Line Islands. *PLoS One* 3**,** e1584.

Dinsdale, E.A., and Rohwer, F. (2011). "Fish or germs? Microbial dynamics associated with
changing trophic structures on Coral Reefs," in *Coral Reefs: An Ecosystem in Transition,*
eds. Z. Dubinsky & N. Stambler. (Dordrecht: Springer Netherlands), 231-240.

Ducklow, H.W. (1983). Production and fate of bacteria in the oceans. *Bioscience* 33**,** 494-501.

Dungan, M.L., Miller, T.E., and Thomson, D.A. (1982). Catastrophic decline of a top carnivore
in the Gulf of California rocky intertidal zone. *Science* 216**,** 989-991.

Eckert, G., Engle, J.M., and Kushner, D. (Year). "Sea star disease and population declines at the
Channel Islands", in: *Proceedings of the 6th California Islands Symposium*, 435-441.
Eisenlord, M.E., Groner, M.L., Yoshioka, R.M., Elliott, J., Maynard, J., Fradkin, S., Turner, M.,
Pyne, K., Rivlin, N., Van Hooidonk, R., and Harvell, C.D. (2016). Ochre star mortality
during the 2014 wasting disease epizootic: role of population size structure and
temperature. *Phil Transac Roy Soc B* 371**,** 20150212.
Ferguson, J.C. (1964). Nutrient transport in starfish. I. Properties of the coelomic fluid. *Biol Bull*
126**,** 33-53.
Ferguson, J.C. (1967a). An autoradiographic study of the utilization of free exogenous amino
acids by starfishes. *Biol Bull* 133**,** 317-329.
Ferguson, J.C. (1967b). Utilization of dissolved exogenous nutrients by the starfishes, *Asterias
forbesi* and *Henricia sanguinolenta*. *Biol Bull* 132**,** 161-173.
Fonseca, M.S., and Kenworthy, W.J. (1987). Effects of current on photosynthesis and
distribution of seagrasses. *Aquat Bot* 27**,** 59-78.
Friedman, J., Hastie, T., and Tibshirani, R. (2010). Regularization paths for generalized linear
models via coordinate descent. *J Statistic Software* 33**,** 1-22.
Fuess, L.E., Eisenlord, M.E., Closek, C.J., Tracy, A.M., Mauntz, R., Gignoux-Wolfsohn, S.,
Moritsch, M.M., Yoshioka, R., Burge, C.A., Harvell, C.D., Friedman, C.S., Hewson, I.,
Hershberger, P.K., and Roberts, S.B. (2015). Up in arms: Immune and nervous system
response to sea star wasting disease. *PLoS One* 10**,** e0133053.
Gonzalez, A., Navas-Molina, J.A., Kosciolek, T., Mcdonald, D., Vázquez-Baeza, Y.,
Ackermann, G., Dereus, J., Janssen, S., Swafford, A.D., Orchanian, S.B., Sanders, J.G.,
Shorenstein, J., Holste, H., Petrus, S., Robbins-Pianka, A., Brislawn, C.J., Wang, M.,
Rideout, J.R., Bolyen, E., Dillon, M., Caporaso, J.G., Dorrestein, P.C., and Knight, R.
(2018). Qiita: rapid, web-enabled microbiome meta-analysis. *Nat Method* 15**,** 796-798.
Gregg, A.K., Hatay, M., Haas, A.F., Robinett, N.L., Barott, K., Vermeij, M.J.A., Marhaver, K.L.,
Meirelles, P., Thompson, F., and Rohwer, F. (2013). Biological oxygen demand optode
analysis of coral reef-associated microbial communities exposed to algal exudates. *PeerJ*
1**,** e107.
Gudenkauf, B.M., and Hewson, I. (2015). Metatranscriptomic analysis of *Pycnopodia
helianthoides* (Asteroidea) affected by sea star wasting disease. *PLoS One* 10**,** e0128150.

Guilloux, C.A., Lamoureux, C., and Hery-Arnaud, G. (2018). Anaerobic bacteria, the unknown
members of the lung microbiota. *M S-Medecine Sciences* 34**,** 253-260.

Haas, A.F., Fairoz, M.F.M., Kelly, L.W., Nelson, C.E., Dinsdale, E.A., Edwards, R.A., Giles, S.,
Hatay, M., Hisakawa, N., Knowles, B., Lim, Y.W., Maughan, H., Pantos, O., Roach,
T.N.F., Sanchez, S.E., Silveira, C.B., Sandin, S., Smith, J.E., and Rohwer, F. (2016).
Global microbialization of coral reefs. *Nat Microbiol* 1**,** 16042.

Haas, A.F., Naumann, M.S., Struck, U., Mayr, C., El-Zibdah, M., and Wild, C. (2010). Organic
matter release by coral reef associated benthic algae in the Northern Red Sea. *J Exper*
Mar Biol Ecol 389**,** 53-60.

Haas, A.F., Nelson, C.E., Wegley Kelly, L., Carlson, C.A., Rohwer, F., Leichter, J.J., Wyatt, A.,
and Smith, J.E. (2011). Effects of coral reef benthic primary producers on dissolved
organic carbon and microbial activity. *PloS One* 6**,** e27973-e27973.

Harrold, C., and Pearse, J.S. (1980). Allocation of pyloric caecum reserves in FED and starved
sea stars, *Pisaster giganteus* (Stimpson): somatic maintenance comes before
reproduction. *J Exper Mar Biol Ecol* 48**,** 169-183.

Harvell, C.D., Montecino-Latorre, D., Caldwell, J.M., Burt, J.M., Bosley, K., Keller, A., Heron,
S.F., Salomon, A.K., Lee, L., Pontier, O., Pattengill-Semmens, C., and Gaydos, J.K.
(2019). Disease epidemic and a marine heat wave are associated with the continental-
scale collapse of a pivotal predator (*Pycnopodia helianthoides*). *Sci Advan* 5**,** eaau7042.

Hewson, I. (2019). Technical pitfalls that bias comparative microbial community analyses of
aquatic disease. *Dis Aquat Organ* 137**,** 109-124.

Hewson, I., Bistolas, K.S.I., Carde, E.M.Q., Button, J.B., Foster, P.J., Flanzenbaum, J.M.,
Kocian, J., and Lewis, C.K. (2018). Investigating the complex association between viral
ecology, environment, and Northeast Pacific sea star wasting. *Front Mar Sci* 5.
https://doi.org/10.3389/fmars.2018.00077.

Hewson, I., Button, J.B., Gudenkauf, B.M., Miner, B., Newton, A.L., Gaydos, J.K., Wynne, J.,
Groves, C.J., Hendler, G., Murray, M., Fradkin, S., Breitbart, M., Fahsbender, E.,
Lafferty, K.D., Kilpatrick, A.M., Miner, C.M., Raimondi, P., Lahner, L., Friedman, C.S.,
Daniels, S., Haulena, M., Marliave, J., Burge, C.A., Eisenlord, M.E., and Harvell, C.D.
(2014). Densovirus associated with sea-star wasting disease and mass mortality. *Proc Nat*
Acad Sci USA 111**,** 17276-17283.

Hewson, I., Johnson, M.R., and Tibbetts, I.R. (2020). An unconventional flavivirus and other
RNA viruses in the sea cucumber (Holothuroidea; Echinodermata) virome. *Viruses* 12**,**
1057.

Hewson, I., Sullivan, B., Jackson, E.W., Xu, Q., Long, H., Lin, C., Quijano Cardé, E.M.,
Seymour, J., Siboni, N., Jones, M.R.L., and Sewell, M.A. (2019a). Perspective:
Something old, something new? Review of wasting and other mortality in Asteroidea
(Echinodermata). *Front Mar Sci* 6.

Hoj, L., Levy, N., Baillie, B.K., Clode, P.L., Strohmaier, R.C., Siboni, N., Webster, N.S.,
Uthicke, S., and Bourne, D.G. (2018). Crown-of-Thorns sea star *Acanthaster* cf. *solaris*
has tissue characteristic microbiomes with potential roles in health and reproduction.
Appl Environ Microbiol 84.

Holland, N.D., and Nealson, K.H. (1978). Fine-structure of the echinoderm cuticle and the sub-
cuticular bacteria of echinoderms. *Acta Zool* 59**,** 169-185.

Hurd, C.L., Cornwall, C.E., Currie, K., Hepburn, C.D., Mcgraw, C.M., Hunter, K.A., and Boyd,
P.W. (2011). Metabolically induced pH fluctuations by some coastal calcifiers exceed
projected 22nd century ocean acidification: a mechanism for differential susceptibility?
Global Change Biology 17**,** 3254-3262.

Jackson, A.L., Inger, R., Parnell, A.C., and Bearhop, S. (2011). Comparing isotopic niche widths
among and within communities: SIBER - Stable Isotope Bayesian Ellipses in R. *J Anim*
Ecol 80**,** 595-602.

Jackson, E.W., Pepe-Ranney, C., Debenport, S.J., Buckley, D.H., and Hewson, I. (2018). The
microbial landscape of sea stars and the anatomical and interspecies variability of their
microbiome. *Front Microbiol* 9**,** 12.

Jackson, E.W., Pepe-Ranney, C., Johnson, M.R., Distel, D.L., and Hewson, I. (2020a). A highly
prevalent and pervasive densovirus discovered among sea stars from the North American
Atlantic coast. *Appl Environ Microbiol* 86.

Jackson, E.W., Wilhelm, R.C., Johnson, M.R., Lutz, H.L., Danforth, I., Gaydos, J.K., Hart,
M.W., and Hewson, I. (2020b). Diversity of sea star-associated densoviruses and
transcribed endogenized viral elements of densovirus origin. *bioRxiv***,**
2020.2008.2005.239004.

Jaffe, N., Eberl, R., Bucholz, J., and Cohen, C.S. (2019). Sea star wasting disease demography
and etiology in the brooding sea star *Leptasterias* spp. *PloS One* 14**,** e0225248-e0225248.

Johansen, K., and Petersen, J.A. (1971). Gas exchange and active ventilation in a starfish,
Pteraster tesselatus. Zeitschrift für vergleichende Physiologie 71**,** 365-381.

Jørgensen, B.B., and Revsbech, N.P. (1985). Diffusive boundary layers and the oxygen uptake of
sediments and detritus. *Limnol Oceanogr* 30**,** 111-122.

Kain, J. (1979). A view of the genus *Laminaria. Oceanogr Mar Biol Ann Rev* 17**,** 101-161.

Kay, S.W.C., Gehman, A.-L.M., and Harley, C.D.G. (2019). Reciprocal abundance shifts of the
intertidal sea stars, *Evasterias troschelii* and *Pisaster ochraceus*, following sea star
wasting disease. *Proc Roy Soc B* 286**,** 20182766.

Kelly, L.W., Williams, G.J., Barott, K.L., Carlson, C.A., Dinsdale, E.A., Edwards, R.A., Haas,
A.F., Haynes, M., Lim, Y.W., Mcdole, T., Nelson, C.E., Sala, E., Sandin, S.A., Smith,
J.E., Vermeij, M.J.A., Youle, M., and Rohwer, F. (2014). Local genomic adaptation of
coral reef-associated microbiomes to gradients of natural variability and anthropogenic
stressors. *Proc Nat Acad Sci USA* 111**,** 10227-10232.

Kelly, M.S., Barker, M.F., Mckenzie, J.D., and Powell, J. (1995). The incidence and morphology
of subcuticular bacteria in the echinoderm fauna of New Zealand. *Biol Bull* 189**,** 91-105.

Kelly, M.S., and Mckenzie, J.D. (1995). Survey of the occurrence and morphology of sub-
cuticular bacteria in shelf echinoderms from the north-east Atlantic Ocean. *Mar Biol* 123**,**
741-756.

Koch, E.W. (1994). Hydrodynamics, diffusion-boundary layers and photosynthesis of the
seagrasses *Thalassia testudinum* and *Cymodocea nodosa. Mar Biol* 118**,** 767-776.

Kohl, W.T., Mcclure, T.I., and Miner, B.G. (2016). Decreased temperature facilitates short-term
sea star wasting disease survival in the keystone intertidal sea star *Pisaster ochraceus*.
PLoS One 11**,** e0153670.

Konar, B., Mitchell, T.J., Iken, K., Coletti, H., Dean, T., Esler, D., Lindeberg, M., Pister, B., and
Weitzman, B. (2019). Wasting disease and static environmental variables drive sea star
assemblages in the Northern Gulf of Alaska. *J Exper Mar Biol Ecol* 520**,** 151209.

Krumhansl, K.A., and Scheibling, R.E. (2012). Production and fate of kelp detritus. *Mar Ecol*
Progr Ser 467**,** 281-302.

Legrand, T., Catalano, S.R., Wos-Oxley, M.L., Stephens, F., Landos, M., Bansemer, M.S.,
Stone, D.a.J., Qin, J.G., and Oxley, A.P.A. (2018). The inner workings of the outer
surface: Skin and gill microbiota as indicators of changing gut health in Yellowtail
Kingfish. *Front Microbiol* 8**,** 17.

Leising, A.W., Schroeder, I.D., Bograd, S.J., Bjorkstedt, E.O., Field, J., Sakuma, K., Abell, J.,
Robertson, R.R., Tuberczy, J., and Al., E. (2014). State of the caliofrnia current 2013-
2014: El Nino Looming. *CalCOFI Report* 55**,** 51-87.

Leslie, H.M., Breck, E.N., Chan, F., Lubchenco, J., and Menge, B.A. (2005). Barnacle
reproductive hotspots linked to nearshore ocean conditions. *Proc Nat Acad Sci USA* 102**,**
10534-10539.

Levin, L.A. (2003). Oxygen minimum zone benthos: Adaptation and community response to
hypoxia. *Oceanogr Mar Biol Ann Rev* 41**,** 1-45.

Levin, L.A., Ekau, W., Gooday, A.J., Jorissen, F., Middelburg, J.J., Naqvi, S.W.A., Neira, C.,
Rabalais, N.N., and Zhang, J. (2009). Effects of natural and human-induced hypoxia on
coastal benthos. *Biogeosciences* 6**,** 2063-2098.

Lloyd, M.M., and Pespeni, M.H. (2018). Microbiome shifts with onset and progression of sea
star wasting disease revealed through time course sampling. *Sci Rep* 8**,** 12.

Lozupone, C., Lladser, M.E., Knights, D., Stombaugh, J., and Knight, R. (2011). UniFrac: an
effective distance metric for microbial community comparison. *ISME J* 5**,** 169-172.

Mcmurdie, P.J., and Holmes, S. (2013). phyloseq: An R package for reproducible interactive
analysis and graphics of microbiome census data. *PLoS One* 8**,** e61217.

Mcmurdie, P.J., and Holmes, S. (2014). Waste not, want not: why rarefying microbiome data is
inadmissible. *PLoS Comput Biol* 10**,** e1003531.

Mead, A.D. (1898). "Twenty-eighth Annual Report of the Commissioners of Inland Fisheries,
Made to the General Assembly at Its January Session, 1898", (eds.) J.M.K. Southwick,
H.T. Root, C.W. Willard, W.M.P. Morton, A.D. Roberts & H.C. Bumpus.).

Melroy, L.M., Smith, R.J., and Cohen, C.S. (2017). Phylogeography of direct-developing sea
stars in the genus *Leptasterias* in relation to San Francisco Bay outflow in central
California. *Mar Biol* 164.

Menge, B.A. (1975). Brood or broadcast? The adaptive significance of different reproductive
strategies in the two intertidal sea stars *Leptasterias hexactis* and *Pisaster ochraceus*.
Mar Biol 31**,** 87-100.

Menge, B.A., Cerny-Chipman, E.B., Johnson, A., Sullivan, J., Gravem, S., and Chan, F. (2016).
Sea star wasting disease in the keystone predator *Pisaster ochraceus* in Oregon: Insights
into differential population impacts, recovery, predation rate, and temperature effects
from long-term research. *PLoS One* 11**,** e0153994.

Meyer, J.L., Castellanos-Gell, J., Aeby, G.S., Häse, C.C., Ushijima, B., and Paul, V.J. (2019).
Microbial community shifts associated with the ongoing stony coral tissue loss disease
outbreak on the Florida Reef Tract. *Front Microbiol* 10.

Miner, C.M., Burnaford, J.L., Ambrose, R.F., Antrim, L., Bohlmann, H., Blanchette, C.A.,
Engle, J.M., Fradkin, S., Gaddam, R., Hardley, C.D.G., Miner, B.G., Murray, S.N.,
Smith, J.R., Whitaker, S.G., and Raimondi, P.T. (2018). Large-scale impacts of sea star
wasting disease (SSWD) on intertidal sea stars and implications for recovery. *PLoS One*
13**,** e0192870.

Mironov, A.N., Dilman, A.B., Vladychenskaya, I.P., and Petrov, N.B. (2016). Adaptive strategy
of the Porcellanasterid sea stars. *Biol Bull* 43**,** 503-516.

Montecino-Latorre, D., Eisenlord, M.E., Turner, M., Yoshioka, R., Harvell, C.D., Pattengill-
Semmens, C.V., Nichols, J.D., and Gaydos, J.K. (2016). Devastating transboundary
impacts of sea star wasting disease on subtidal asteroids. *PLoS One* 11**,** e0163190.

Moore, S.K., Stark, K., Bos, J., Williams, P., Newton, J., and Dzinbal, K. (2014). "Puget sound
marine waters: 2013 overview.", P.S.E.M.P.M.W. Workgroup. (Seattle, Washington).

Murphy, R.R., Kemp, W.M., and Ball, W.P. (2011). Long-term trends in Chesapeake Bay
seasonal hypoxia, stratification, and nutrient loading. *Estuaries Coasts* 34**,** 1293-1309.

Nakagawa, S., Saito, H., Tame, A., Hirai, M., Yamaguchi, H., Sunata, T., Aida, M., Muto, H.,
Sawayama, S., and Takaki, Y. (2017). Microbiota in the coelomic fluid of two common
coastal starfish species and characterization of an abundant *Helicobacter*-related taxon.
Sci Rep 7**,** 8764.

Nakajima, R., Haas, A.F., Silveira, C.B., Kelly, E.L.A., Smith, J.E., Sandin, S., Kelly, L.W.,
Rohwer, F., Nakatomi, N., and Kurihara, H. (2018). Release of dissolved and particulate

organic matter by the soft coral *Lobophytum* and subsequent microbial degradation. *J Exper Mar Biol Ecol* 504**,** 53-60.

Nance, J.M. (1981). Respiratory water flow and production of mucus in the cushion star, *Pteraster tesselatus* Ives (Echinodermata: Asteroidea). *J Exper Mar Biol Ecol* 50**,** 21-31.

Nepf, H.M. (2011). "2.13 - Flow Over and Through Biota," in *Treatise on Estuarine and Coastal Science,* eds. E. Wolanski & D. Mclusky. (Waltham: Academic Press), 267-288.

Noble, R.T., and Fuhrman, J.A. (1998). Use of SYBR Green I rapid epifluoresence counts of marine viruses and bacteria. *Aquat Microb Ecol* 14**,** 113-118.

Nugues, M.M., Smith, G.W., Van Hooidonk, R.J., Seabra, M.I., and Bak, R.P.M. (2004). Algal contact as a trigger for coral disease. *Ecol Lett* 7**,** 919-923.

Nunez-Pons, L., Work, T.M., Angulo-Preckler, C., Moles, J., and Avila, C. (2018). Exploring the pathology of an epidermal disease affecting a circum-Antarctic sea star. *Sci Rep* 8**,** 12.

Ochiai, M., Nakajima, T., and Hanya, T. (1980). Chemical composition of labile fractions in DOM. *Hydrobiol* 71**,** 95-97.

Ogawa, H., and Tanoue, E. (2003). Dissolved organic matter in oceanic waters. *J Oceanogr* 59**,** 129-147.

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., Mcglinn, D., Minchin, P.R., O'hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., and Wagner, H. (2019). "vegan: Community Ecology Package". R package version 2.5-6 ed.).

Olsen, A.Y., Smith, A., and Larson, S. (2016) "A temporal analysis of water quality variability at the Seattle Aquarium in Elliott Bay, Puget Sound, WA," in *Water Challenges of an Urbanizing World,* ed. M. Glavan. (DOI: 10.5772/intechopen.68339: IntechOpen), 55-69.

Patterson, H.K., and Carmichael, R.H. (2018). Dissolved oxygen concentration affects $\delta15N$ values in oyster tissues: implications for stable isotope ecology. *Ecosphere* 9**,** e02154.

Pinhassi, J., Sala, M.M., Havskum, H., Peters, F., Guadayol, Ò., Malits, A., and Marrasé, C. (2004). Changes in bacterioplankton composition under different phytoplankton regimens. *Appl Environ Microbiol* 70**,** 6753-6766.

Piruat, J.I., and López-Barneo, J. (2005). Oxygen tension regulates mitochondrial DNA-encoded complex I gene expression. *J Biol Chem* 280**,** 42676-42684.

Pohlner, M., Dlugosch, L., Wemheuer, B., Mills, H., Engelen, B., and Reese, B.K. (2019). The
majority of active *Rhodobacteraceae* in marine sediments belong to uncultured genera: A
molecular approach to link their distribution to environmental conditions. *Front*
Microbiol 10.

Porter, K.G., and Feig, Y.S. (1980). The use of DAPI for identifying and counting aquatic
microflora. *Limnol Oceanogr* 25**,** 943-948.

Preisig, O., Zufferey, R., Thöny-Meyer, L., Appleby, C.A., and Hennecke, H. (1996). A high-
affinity cbb3-type cytochrome oxidase terminates the symbiosis-specific respiratory
chain of *Bradyrhizobium japonicum*. *J Bacteriol* 178**,** 1532-1538.

Propp, M.V., Ryabushko, V.I., Zhuchikhina, A.A., and Propp, L.N. (1983). Seasonal changes in
respiration, ammonia and phosphate excretion, and activity of carbohydrate-metabolism
enzymes in four echinoderm species from the sea of Japan. *Comp Biochem Physiol B* 75**,**
707-711.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner,
F.O. (2012). The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucl Acid Res* 41**,** D590-D596.

Reverter, M., Sasal, P., Tapissier-Bontemps, N., Lecchini, D., and Suzuki, M. (2017).
Characterisation of the gill mucosal bacterial communities of four butterflyfish species: a
reservoir of bacterial diversity in coral reef ecosystems. *FEMS Microb Ecol* 93**,** 10.

Roach, T.N.F., Abieri, M.L., George, E.E., Knowles, B., Naliboff, D.S., Smurthwaite, C.A.,
Kelly, L.W., Haas, A.F., and Rohwer, F.L. (2017). Microbial bioenergetics of coral-algal
interactions. *PeerJ* 5**,** e3423.

Robinson, J.D., Mann, K.H., and Novitsky, J.A. (1982). Conversion of the particulate fraction of
seaweed detritus to bacterial biomass. *Limnol Oceanogr* 27**,** 1072-1079.

Rosado, D., Pérez-Losada, M., Severino, R., Cable, J., and Xavier, R. (2019). Characterization of
the skin and gill microbiomes of the farmed seabass (*Dicentrarchus labrax*) and
seabream (*Sparus aurata*). *Aquaculture* 500**,** 57-64.

Ruiz-Ramos, D.V., Schiebelhut, L.M., Hoff, K.J., Wares, J.P., and Dawson, M.N. (2020). An
initial comparative genomic autopsy of wasting disease in sea stars. *Molec Ecol* 29**,** 1087-
1102.

Santoro, A.E., Nidzieko, N.J., Dijken, G.L.V., Arrigo, K.R., and Boehma, A.B. (2010). Contrasting spring and summer phytoplankton dynamics in the nearshore Southern California Bight. *Limnol Oceanogr* 55**,** 264-278.

Sato, Y., Ling, E.Y.S., Turaev, D., Laffy, P., Weynberg, K.D., Rattei, T., Willis, B.L., and Bourne, D.G. (2017). Unraveling the microbial processes of black band disease in corals through integrated genomics. *Sci Rep* 7**,** 40455.

Schiebelhut, L.M., Puritz, J.B., and Dawson, M.N. (2018). Decimation by sea star wasting disease and rapid genetic change in a keystone species, *Pisaster ochraceus*. *Proc Nat Acad Sci USA* 115**,** 7069-7074.

Schultz, J.A., Cloutier, R.N., and Cote, I.M. (2016). Evidence for a trophic cascade on rocky reefs following sea star mass mortality in British Columbia. *PeerJ* 4**,** 19.

Sharma, S., Singh, S., Gupta, R., Ganju, L., Singh, S., Kumar, B., and Singh, Y. (2019). Mitochondrial DNA sequencing reveals association of variants and haplogroup M33a2′3 with High altitude pulmonary edema susceptibility in Indian male lowlanders. *Sci Rep* 9.

Shibata, A., Goto, Y., Saito, H., Kikuchi, T., Toda, T., and Taguchi, S. (2006). Comparison of SYBR Green I and SYBR Gold stains for enumerating bacteria and viruses by epifluorescence microscopy. *Aquat Microb Ecol* 43**,** 223-231.

Shick, J.M. (1976). Physiological and behavioral responses to hypoxia and hydrogen sulfide in the infaunal asteroid *Ctenodiscus crispatus*. *Mar Biol* 37**,** 279-289.

Shick, J.M., Edwards, K.C., and Dearborn, J.H. (1981). Physiological ecology of the deposit feeding sea star *Ctenodiscus cripsatus* - ciliated surfaces and animal-sediment interactions. *Mar Ecol Progr Ser* 5**,** 165-184.

Siebers, D. (1979). Transintegumentary uptake of dissolved amino acids in the sea star *Asterias rubens* - Reassessment of its nutritional role with special reference to the significance of heterotrophic bacteria. *Mar Ecol Progr Ser* 1**,** 169-177.

Siebers, D. (2015). Bacterial—invertebrate interactions in uptake of dissolved organic matter. *Amer Zool* 22**,** 723-733.

Silveira, C.B., Luque, A., Roach, T.N.F., Villela, H., Barno, A., Green, K., Reyes, B., Esther, R.-P., Le, T., Mead, S., Hatay, M., Bermeij, M.J., Takeshita, Y., Haas, A.F., Bailey, B., and Rohwer, F. (2019). Biophysical and physiological processes causing oxygen loss from coral reefs. *eLife* 8**,** e49114.

Silverman, J.D., Washburne, A.D., Mukherjee, S., and David, L.A. (2017). A phylogenetic

transform enhances analysis of compositional microbiota data. *eLife* 6**,** e21887.

Smith, J.E., Shaw, M., Edwards, R.A., Obura, D., Pantos, O., Sala, E., Sandin, S.A., Smriga, S.,

Hatay, M., and Rohwer, F.L. (2006). Indirect effects of algae on coral: algae-mediated,

microbe-induced coral mortality. *Ecol Lett* 9**,** 835-845.

Spence, C.D., Vanaudenaerde, B., Einarsson, G.G., Mcdonough, J., Lee, A.J., Johnston, E.,

Verleden, G.M., Elborn, J.S., Dupont, L.J., Van Herck, A., Gilpin, D.F., Vos, R., Tunney,

M.M., and Verleden, S.E. (2020). Influence of azithromycin and allograft rejection on the

post-lung transplant microbiota. *J Heart Lung Transplant* 39**,** 176-183.

Suchy, K.D., Le Baron, N., Hilborn, A., Perry, R.I., and Costa, M. (2019). Influence of

environmental drivers on spatio-temporal dynamics of satellite-derived chlorophyll a in

the Strait of Georgia. *Progr Oceanogr* 176**,** 102134.

Thiele, S., Richter, M., Balestra, C., Glöckner, F.O., and Casotti, R. (2017). Taxonomic and

functional diversity of a coastal planktonic bacterial community in a river-influenced

marine area. *Mar Genom* 32**,** 61-69.

Thornton, D.C.O. (2014). Dissolved organic matter (DOM) release by phytoplankton in the

contemporary and future ocean. *Eur J Phycol* 49**,** 20-46.

Tobler, M., Henpita, C., Bassett, B., Kelley, J.L., and Shaw, J.H. (2014). H_2S exposure elicits

differential expression of candidate genes in fish adapted to sulfidic and non-sulfidic

environments. *Comp Bioechem and Physiol A* 175**,** 7-14.

Van Alstyne, K.L. (2016). Seasonal changes in nutrient limitation and nitrate sources in the

green macroalga *Ulva lactuca* at sites with and without green tides in a northeastern

Pacific embayment. *Mar Poll Bull* 103**,** 186-194.

Vander Zanden, M.J., Clayton, M.K., Moody, E.K., Solomon, C.T., and Weidel, B.C. (2015).

Stable isotope turnover and half-life in animal tissues: A literature synthesis. *PLoS One*

10**,** e0116182.

Vistisen, B., and Vismann, B. (1997). Tolerance to low oxygen and sulfide in *Amphiura*

filiformis and *Ophiura albida* (Echinodermata: Ophiuroidea). *Mar Biol* 128**,** 241-246.

Wang, K., Huang, Y., Zhang, Z., Liao, J., Ding, Y., Fang, X., Liu, L., Luo, J., and Kong, J.

(2019). A preliminary study of microbiota diversity in saliva and bronchoalveolar lavage

fluid from patients with primary bronchogenic carcinoma. *Med Sci Monit* 25**,** 2819-2834.

Washburne, A.D., Silverman, J.D., Leff, J.W., Bennett, D.J., Darcy, J.L., Mukherjee, S., Fierer,

N., and David, L.A. (2017). Phylogenetic factorization of compositional data yields

lineage-level associations in microbiome datasets. *PeerJ* 5**,** e2969.

Wu, Y., Klapper, I., and Stewart, P.S. (2018). Hypoxia arising from concerted oxygen

consumption by neutrophils and microorganisms in biofilms. *Pathogen Dis* 76**,** fty043.

Zhang, T., and Wang, X. (2017). Release and microbial degradation of dissolved organic matter

(DOM) from the macroalgae *Ulva prolifera*. *Mar Pollut Bull* 125**,** 192-198.

1300

1301

1302

1303

Fig. 1: Conceptualization of BLODL. Under typical conditions (A), boundary layer conditions are normoxic because of limited dissolved organic matter inputs, and consequently lower heterotrophic bacterial respiration. When organic matter (possibly from primary production) increases in surrounding waters (B; e.g. during heightened primary productivity, from terrestrial runoff, or from decaying asteroid carcasses), this stimulates bacterial heterotrophic respiration and abundance and results in the formation of suboxic waters within the limited water motion boundary layer, which in turn results in longer distances over which diffusion must occur to maintain animal respiratory demand. Over time (C) suboxic conditions in the boundary layer results in tissue damage, and prevalence of strict and facultative anaerobes. Because their growth is less efficient than aerobic metabolisms their abundance is less than at hypoxia onset. Release of labile organic matter from decaying tissues (D) and persistent organic-matter rich conditions within the asteroid boundary layer result in animal mortality.

1316

1317

1318

Fig 2: Proportion of asymptomatic *P. ochraceus* (n = 5 each treatment) incubated in flow-

through conditions at the Bodega Marine Lab in response to organic matter enrichments

(Peptone, *Dunaliella tertiolecta* culture POM, and coastal POM collected from the inflow at the

Bodega Marine Laboratory in August 2019. Dissolved O_2 and temperature were measured in

flow-through sea tables bearing each OM treatment. Error Bars = SD. The red arrows above the

top panel indicate sampling for microbiome analyses. * indicates wasting speed (i.e. time to

1325   appearance of first lesion) was significantly (p<0.05, Student's t-test) faster than control. ‡

1326   indicates that the the overall trend in lesion formation was significantly different to controls (p<

1327   0.05, log-rank test).

1328

Fig. 3: Differential abundance of bacterial taxa from body wall samples (A; *P. ochraceus* June 2018) and surface swabs (B-F; *P. ochraceus* August 2019). (A) Boxplots were derived using PhyloFactor (Washburne et al. 2017), which uses a generalized linear model to regress the isometric log-ratio (ILR balance) between opposing clades (contrasted by an edge) on a phylogenetic tree. This was done iteratively, with each iteration, or factor, maximizing the F statistic from regression. Shown taxa represent either a single factor or combination of factors (when, for example, multiple factors identified different sOTUs with the same taxonomic classification). Labels represent either the highest taxonomic resolution or the highest classification shared by all sOTUs of a given clade. T0 = experiment commencement, TI = lesion genesis, TF = time of death. (B-D) Balance contrast of early (before lesion genesis) samples compared to late (immediately prior to lesion genesis) samples. Samples were transformed using the Phylogenetic Isometric Log-Ratio (PhILR; Silverman et al. 2017) transform, which uses a phylogenetic tree (B) to convert an sOTU table into a new matrix of coordinates derived from the ILR of clades that descend from a common node. We used a sparse logistic regression with an l_1 penalty of $\lambda=0.15$ (Silverman et al. 2017) to analyze the ILR at each node, and included a select number of 'balances' with positive coefficients (C-D). (C) is the balance of *Nitrosopumilus* (colored blue in (B), comprises the thin sliver on the right side of the tree) relative to the rest of the dataset (also shown in blue in (B)). A positive shift indicates an increase in *Nitrosopumilus* relative to its denominator. (D) is the balance between a clade of Alpha/Gammaproteobacteria (large, red clade in (B)) and Deltaproteobacteria (Bdellovibrionales and Desulfobacterales; small, red clade in (B)). A negative shift indicates that the denominator, Deltaproteobacteria, is increasing relative to Alpha/Gammaproteobacteria. (E) and (F) were derived from a PhyloFactor object and show the ILR balance of Flavobacteriales (E) and Rhodobacterales (F) relative to all other sOTUs. Organic amendment is given above boxplots. Time 0 = 0 h, Time 2 = 48 h, Time 4 = 96 h; Time 8 = 192 h; Time 10 = 240 h. Total sample numbers for each treatment (which varied due to the loss of asteroids over the course of the experiment to wasting) is given in Fig. 4. The vertical red line in panels E and F indicate the average time at which asteroids formed lesions.

1358

1359

Fig 4: Relative abundance of bacterial orders derived from *P. ochraceus* epidermal swabs.

Specimens were enriched with the indicated organic material and sampled until lesion genesis.

Time 0 represents initial sampling and each subsequent time indicates the respective day. n

values reflect the number of healthy specimens at each given timepoint. The solid blue line on

each panel indicates when lesions first formed per treatment, and the solid red line on each panel

indicates the mean lesion time within the treatment. Time 0 = 0 h, Time 2 = 48 h, Time 4 = 96 h;

Time 8 = 192 h; Time 10 = 240 h.

1367

1368

1369

Fig 5: Abundance of bacteria in proximity to *P. ochraceus* surfaces (top), percentage change from initial (middle), and relative to controls (bottom). during first 10 days of experiment in response to organic matter enrichment (n = 5 for each treatment) as assessed by SYBR Gold staining and epifluorescence microscopy. Control specimens are indicated in red, while the mean of stars that wasted in Peptone, *Dunaliella tertiolecta* POM, and Coastal POM are indicated

separately. The solid vertical line on the top panel represents the mean time that asteroids

developed lesions in the coastal POM treatment, the solid line on the middle panel represents the

mean time for lesion development in *Dunaliella tertiolecta* POM treatments, and solid line on

the bottom panel represents the mean time for lesion development in peptone treatments

(separated between panels for clarity). The shaded regions represent lesion development standard

error for respective treatments.

1381

1382

Fig. 6: Proportion of asymptomatic *P. ochraceus* (n = 5) remaining over time during longitudinal

study of microbiome composition in the absence of external stimuli. The mean flow rate into

aquariums was 3.81 ± 0.05 mL s^{-1} (average residence time in aquariums 37 min).

1386

1387

1388 **Fig. 7:** Proportion of asymptomatic *Asterias forbesi* incubated in normoxic and suboxic water

1389 (top) and variation in temperature and O₂ in incubation aquaria (bottom) over time. The red

1390 arrows at top indicate samples which were included in analysis of microbiome composition (see

1391 Fig. S9).

1392

Fig. 8: Proportion asymptomatic *P. ochraceus* over time in response to treatment (top)
desiccation (n = 3), and (bottom) treatment with crude and proteinase k-treated tissue
homogenates (n = 2). * indicates that low flow lesion genesis time was significantly (p<0.05,
Student's t-test) compared to high flow rate; ‡ indicates that the overall trend of desiccation
under low vs high flow rate and with the addition of proteinase-k treated homogenate vs low
flow controls was significant (p< 0.05, log-rank test).

1402

Fig. 9: Rugosity of similarly-sized animals between wasting-affected and less wasting affected

species as determined by whole animal computed tomography (top) and of an asteroid ray by

micro-computed tomography (bottom). a, b denote significant difference at $p < 0.001$. More and

less wasting affected were based upon previous work and defined in the text.

1407

1408

1409 **Fig 10:** Correspondence between mean time of wasting mass mortality (indicated by solid

1410 orange line (SE range indicated by lighter orange bar) compared with physico-chemical

1411 parameters (top) and chlorophyll a concentration (bottom) at Penn Cove, Whidbey Island from

1412 2014 - 2019. Temp = temperature; DO = dissolved oxygen; Sal = salinity.

1413

1414

Fig. 11: Comparison of asymptomatic and wasting δ^{15}N values between species. ONP = Starfish
point, Olympic national park and SC = Davenport, Santa Cruz, CA. * indicates p< 0.05.
Numbers above box plots indicate *n* of specimens used in comparison.

1415
1416
1417

1418

1419