The distribution of plasmid fitness effects explains plasmid persistence in bacterial communities

Aida Alonso-del Valle[1], Ricardo León-Sampedro[1,2], Jerónimo Rodríguez-Beltrán[1,2], Javier DelaFuente[1], Marta Hernández-García[1,3], Patricia Ruiz-Garbajosa[1,3], Rafael Cantón[1,3], Rafael Peña-Miller[4,*], Álvaro San Millán[1,2,5]*.

[1] *Servicio de Microbiología. Hospital Universitario Ramón y Cajal* and *Instituto Ramón y Cajal de Investigación Sanitaria*. Madrid, Spain.

[2] *Centro de Investigación Biológica en Red. Epidemiología y Salud Pública*, *Instituto de Salud Carlos III*. Madrid. Spain.

[3] *Red Española de Investigación en Patología Infecciosa*. *Instituto de Salud Carlos III*. Madrid. Spain.

[4] Center for Genomic Sciences, *Universidad Nacional Autónoma de México*, Cuernavaca, Mexico.

[5] Current address: *Centro Nacional de Biotecnología–CSIC*, Madrid, Spain.

* Correspondence: Álvaro San Millán, alvsanmillan@gmail.com ORCID: 0000-0001-8544-0387 and Rafael Peña-Miller, rafael.penamiller@gmail.com ORCID: 0000-0002-2767-0640

Introductory paragraph

Plasmid persistence in bacterial populations is strongly influenced by the fitness effects associated with plasmid carriage. However, plasmid fitness effects in wild-type bacterial hosts remain largely unexplored. In this study, we determined the distribution of fitness effects (DFE) for the major antibiotic resistance plasmid pOXA-48 in wild-type, ecologically compatible enterobacterial isolates from the human gut microbiota. Our results show that although pOXA-48 produced an overall reduction in bacterial fitness, the DFE was dominated by quasi-neutral effects, and beneficial effects were observed in several isolates. Incorporating these data into a simple population dynamics model revealed a new set of conditions for plasmid stability in bacterial communities, with plasmid persistence increasing with bacterial diversity and becoming less dependent on conjugation. Moreover, genomic results showed a link between plasmid fitness effects and bacterial phylogeny, helping to explain pOXA-48 epidemiology. Our results provide a simple and general explanation for plasmid persistence in natural bacterial communities.

Introduction

Plasmids are extra-chromosomal mobile genetic elements able to transfer between bacteria through conjugation[1]. Plasmids carry accessory genes that help their hosts to adapt to a myriad of environments and thus play a key role in bacterial ecology and evolution[2]. A key example of the importance of plasmids in bacterial evolution is their central role in the spread of antibiotic resistance mechanisms among clinical pathogens over recent decades[3,4]. Some of the most clinically relevant resistance genes, such those encoding carbapenemases (ß-lactamase enzymes able to degrade carbapenem antibiotics), are carried on conjugative plasmids that spread across high-risk bacterial clones[5,6].

Despite the abundance of plasmids in bacterial populations and the potential advantages associated with their acquisition, these genetic elements generally produce physiological alterations in their bacterial hosts that lead to a reduction in fitness[7–9]. These fitness costs make it difficult to explain how plasmids are maintained in bacterial populations over the long-term in the absence of selection for plasmid-encoded traits, a puzzle known as "the plasmid-paradox"[10]. Different solutions to this paradox have been proposed. For example, compensatory evolution contributes to plasmid persistence by alleviating the costs associated with plasmid-carriage, and a high conjugation rate can promote the survival of plasmids as genetic parasites[11–18].

Over the past decades, many studies have investigated the existence conditions for plasmids in bacterial populations[14,18–23]. However, understanding of plasmid population biology is held in check by limitations of the model systems used for its study. First, most experimental reports of fitness costs have studied arbitrary associations between plasmids and laboratory bacterial strains[7,24]. These examples do not necessarily replicate plasmid fitness effects in natural bacterial hosts, which remain largely unexplored. Second, studies tend to analyse the fitness effects of a single plasmid in a single bacterium. However, plasmid fitness effects can differ between bacteria[25–28], and this variability may impact plasmid persistence in bacterial communities (for a relevant example see[29]). Third, most mathematical models of plasmid population biology study clonal or near-clonal populations. However, bacteria usually live in complex communities in which conjugative plasmids can spread between different bacterial hosts[30–32]. To fully understand plasmid persistence in natural bacterial populations, it will be necessary to address these limitations.

In this study, we provide the first description of the distribution of fitness effects (DFE) of a plasmid in

wild-type bacterial hosts. We used the clinically relevant carbapenem-resistance conjugative plasmid

pOXA-48 and 50 enterobacteria strains isolated from the gut microbiota of patients admitted to a large

tertiary hospital in Madrid. Incorporation of the experimentally determined DFE into a population biology

model provides new key insights into the existence conditions of plasmids in bacterial communities.

Results

Construction of a pOXA-48 transconjugant collection

We studied the DFE of the plasmid pOXA-48 in a collection of ecologically compatible bacterial hosts.

pOXA-48 is an enterobacterial, broad-host-range, conjugative plasmid that is mainly associated with *K.*

pneumoniae and *Escherichia coli*[33–35]. pOXA-48 encodes the carbapenemase OXA-48 and is

distributed worldwide, making it one of the most clinically important carbapenemase-producing

plasmids[6,34]. The gut microbiota of hospitalised patients is a frequent source of enterobacteria clones

carrying pOXA-48[6]. In recent studies, we described the in-hospital epidemiology of pOXA-48 in a large

collection of extended-spectrum ß-lactamase (ESBL)- and carbapenemase-producing enterobacteria

isolated from more than 9,000 patients in our hospital over a period of two years (R-GNOSIS collection,

see methods)[31,36–38]. pOXA-48-carrying enterobacteria were the most frequent carbapenemase-

producing enterobacteria in the hospital, with 171 positive isolates, and they colonised 1.13% of the

patients during the study period (105/9,275 patients). In this study we focused on plasmid pOXA-

48_K8, which is a recently described pOXA-48-like plasmid isolated from a *K. pneumoniae* in our

hospital[31] (Figure 1a, for simplicity we will refer to pOXA-48_K8 and pOXA-48-like plasmids as pOXA-

48 throughout the text).

To study the DFE of pOXA-48, we selected 50 isolates from the R-GNOSIS collection as bacterial

hosts. Our criteria were to select (i) pOXA-48-free isolates, to avoid selecting clones in which

compensatory evolution had already reduced plasmid-associated costs; (ii) isolates from the most

frequent pOXA-48-carrying species, *K. pneumoniae* and *Escherichia coli*; and (iii) strains isolated from

patients located in wards in which pOXA-48-carrying enterobacteria were commonly reported[31]. The

underlying rationale was to select clones which were naïve to pOXA-48 but ecologically compatible with

it (*i.e.* isolated from patients coinciding on wards with others who were colonised with pOXA-48-carrying

clones). We selected 25 *K. pneumoniae* and 25 *E. coli* isolates that are representative of the R-

GNOSIS study and cover the *K. pneumoniae* and *E. coli* phylogenetic diversity in the collection (see

methods, Figure 1b and Supplementary Table 1). It is important to note that, because of the nature of

the R-GNOSIS collection, the isolates used in this study produce ESBLs. However, ESBL-producing

enterobacteria are widespread not only in hospitals but also in the community[39], and most pOXA-48-

carrying enterobacteria isolated in our hospital also produce ESBLs[36].

pOXA-48 was introduced into the collection of recipient strains by conjugation (see Methods), and the

presence of the plasmid was confirmed by PCR and antibiotic susceptibility testing (Supplementary

Table 2). The presence of the entire pOXA-48 plasmid was confirmed by sequencing the complete

genomes of the 50 transconjugant clones, which also revealed the genetic relatedness of the isolates

(Figure 1b). In line with previous studies[31,40], the sequencing results revealed that a subset of isolates

initially identified as *K. pneumoniae* in fact belonged to the species *Klebsiella quasipneumoniae* (n= 4)

and *Klebsiella variicola* (n= 1). These species are also pOXA-48 hosts in our hospital[31] and so were

maintained in the study (Figure 1b).

Figure 1. Experimental model system. Representation of pOXA-48 plasmid and the enterobacteria

strains used in this study. (a) pOXA-48_K8 (accession number MT441554). Reading frames are shown

as arrows, indicating the direction of transcription. Colours indicate gene function classification (see

legend). The *bla*OXA-48 gene is shown in pink. (b) Unrooted phylogeny of whole-genome assemblies

from *E. coli* clones (left) and *Klebsiella* spp. clones (right). Branch length gives the inter-assembly mash

distance (a measure of k-mer similarity). The grouping of multi-locus sequence types (ST) is also

indicated (*E. coli* ST6217 belongs to the ST10 group). Note that the sequencing results revealed that a

subset of isolates initially identified as *K. pneumoniae* were in fact *Klebsiella quasipneumoniae* (n= 4)

and *Klebsiella variicola* (n= 1).

Measuring pOXA-48 fitness effects

To measure pOXA-48 fitness effects, we performed growth curves and competition assays for all the

plasmid-carrying and plasmid-free clones in the collection. We first performed growth curves in pure

cultures to calculate maximum growth rate (μ_{max}) and maximum optical density (OD_{max}), which can be

used to estimate the intrinsic population growth rate (r) and carrying capacity (K), respectively

(Supplementary Figure 1). We also measured the area under the growth curve (AUC), which integrates

information about r and K. To estimate plasmid-associated fitness effects, we compared these

parameters between each plasmid-carrying and plasmid-free pair of isogenic isolates (Figure 2a). The

results showed that, as expected, pOXA-48 produced an overall decrease in the parameters extracted

from the growth curves. However, in a substantial subset of clones, plasmid acquisition was not

associated with a reduction in these parameters (Figure 2a).

Competition assays allow measurement of the relative fitness (w) of two bacteria competing for

resources in the same culture[41]. Competition between otherwise isogenic plasmid-carrying and

plasmid-free clones thus provides a quantitative assessment of the fitness costs associated with

plasmid carriage. For the competition assays, we used flow cytometry; strains were labelled using an

in-house developed small, non-conjugative plasmid vector, called pBGC, that encodes an inducible

green fluorescent protein (GFP) (Supplementary Figure 2). pBGC was introduced into the wild-type

isolate collection by electroporation, and all pOXA-48-carrying and pOXA-48-free clones were

competed against their pBGC-carrying parental strain. We were unable to introduce pBGC into eight of

the isolates; in those cases, for the competitor, we used *E. coli* strain J53 carrying the pBGC vector

(see Methods for details). Data from the competition assays were used to calculate the competitive

fitness of pOXA-48-carrying clones relative to their plasmid-free counterparts (Figure 2b). There were

no significant differences between the fitness effects of pOXA-48 in *Klebsiella* spp*.* and *E. coli* isolates

(ANOVA effect of Species x Plasmid interaction; F=0.088, df=1, P=0.767).

To validate our results, we compared the values obtained from growth curves and competition assays.

This analysis revealed a significant correlation between relative fitness values and the parameters

extracted from the growth curves (Supplementary Figure 3).

13:

Figure 2. pOXA-48 fitness effects in a set of ecologically compatible wild-type enterobacteria. (a) Relative values of growth-curve parameters (plasmid-carrying/plasmid-free isogenic clones): maximum optical density (OD_{max}, pink), maximum growth rate (μ_{max}, yellow), and area under the curve (AUC, green). Dots represent each relative value (red, *E. coli*; blue, *Klebsiella* spp.). Values below 1 indicate a reduction in these parameters associated with plasmid-acquisition. Five biological replicates were performed for each growth curve. (b) Relative fitness (*w*) of plasmid-carrying clones compared with plasmid-free clones obtained by competition assays (red, *E. coli*; blue, *Klebsiella* spp.). Values below 1 indicate a reduction in *w* due to plasmid acquisition; values above 1 indicate an increase in *w*. Bars represent the mean of five independent experiments, and error bars represent the standard error of the mean. Two horizontal lines indicate those clones showing significant costs or benefits associated with carrying pOXA-48 plasmid.

The distribution of pOXA-48 fitness effects

Results from the competition assays showed that the overall effect of pOXA-48 was a small but significant reduction in relative fitness (mean *w*= 0.971, ANOVA effect of plasmid; F=70.04, df=1, P=1.02x10^{-15}). However, plasmid fitness effects varied greatly between the isolates in the collection, producing a normal distribution ranging from a >20% reduction to almost a 20% increase in relative fitness (Figure 2b and 3a; Shapiro-Wilk normality test, P= 0.14). Indeed, plasmid acquisition was associated with a significant fitness decrease in only 14 strains, and 7 isolates showed a significant increase in fitness (Bonferroni corrected two sample t-test, *P*< 0.05). These results revealed a highly

dynamic scenario in which a plasmid produces a wide distribution of fitness effects in different bacterial

hosts, ranging from costs to benefits.

To place our results in context with previous reports, we compared the DFE for pOXA-48 with the

results from a recent meta-analysis of plasmid fitness effects by Vogwill and MacLean[24] (Figure 3).

These authors recovered data for 50 plasmid-bacterium pairs from 16 studies. The DFE constructed

from those reports showed a higher mean plasmid cost (mean w= 0.91) and differed significantly from

the DFE we report here for pOXA-48 in wild-type enterobacteria (Wilcoxon signed rank test, V= 922, P=

0.006). The discrepancy between these distributions may, at least in part, reflect the different nature of

plasmid-bacterium associations considered in the different studies. Although the plasmids studied in

earlier reports were isolated from natural sources, they were introduced into laboratory bacterial strains,

and the detected fitness effects may not be fully representative of wild-type plasmid-bacterium

associations. Our study, on the other hand, analysed the fitness effects of pOXA-48 in ecologically

compatible bacterial hosts. Taken together, the data suggest that the distribution of plasmid fitness

effects is likely influenced by the ecological compatibility between plasmids and their bacterial hosts.

Figure 3. Distribution of plasmid fitness effects. Comparison between the DFE obtained in this study

and the DFE from previous studies. (a) DFE for pOXA-48 in the ecologically compatible collection of

enterobacteria isolates. Bars indicate the number of *E. coli* (red) and *Klebsiella* spp. (blue) strains in

each relative fitness category. The grey dotted line indicates the mean relative fitness of the population.

Note that relative fitness values are normally distributed (w= 0.971, var= 0.0072). (b) DFE for plasmids

in bacterial hosts obtained in a previous meta-analysis[24]. Most of the included studies were based on

arbitrary associations between plasmids and laboratory strains. Bars indicate the number of plasmid-

bacterium associations in each relative fitness category. The grey dotted line indicates the mean

relative fitness across studies. Relative fitness values are not normally-distributed (w= 0.91, var= 0.029;

Shapiro-Wilk normality test, P= 0.0006).

---

pOXA-48 fitness effects across bacterial phylogeny

A key limit to the prediction of plasmid-mediated evolution is the inability to anticipate plasmid fitness

effects in new bacterial hosts. This is particularly relevant to the evolution of antibiotic resistance

because some of the most concerning multi-resistant clinical pathogens arise from very specific

associations between resistance plasmids and high-risk bacterial clones[4,6,42]. Interestingly, a recent

study in an important pathogenic *E. coli* lineage (ST131) showed that the acquisition and maintenance

of resistance plasmids is associated with specific genetic signatures[43]. Pursuing this idea, we analysed

the DFE for pOXA-48 across the whole-genome phylogeny of our isolates, with the aim of determining if

genetic content could help to predict plasmid fitness effects (Figure 4). We calculated the genetic

relatedness of *Klebsiella* spp. and *E. coli* isolates by reconstructing their core genome phylogeny

(Figure 4a). Plasmid fitness effects can also be strongly influenced by the accessory genome. For

example, the presence of further mobile genetic elements can deeply impact the costs of plasmids[44,45].

Therefore, we also constructed trees from the distance matrix of the accessory gene network[46], which

includes plasmid content (Figure 4b).

For each group of isolates, we scanned the fitness effects of pOXA-48 across the core and accessory

genome using the local indicator of phylogenetic association index[47,48] (LIPA, see Supplementary

Figure 4, Supplementary Table 3, and methods for the complete analysis). For the *E. coli* isolates, the

results showed no association of pOXA-48 fitness effects with the core or accessory phylogenies (LIPA,

P> 0.1). In contrast, for *Klebsiella* spp., LIPA indices revealed a significant phylogenetic signal in four

clones in which pOXA-48 produced a high fitness cost, all of them belonging to ST1427 (Kpn01, Kpn04,

Kpn08, and Kpn13, accounting for 4 of the 5 ST1427 clones analysed in this study; LIPA, P< 0.05).

Three of these ST1427 clones also produced a significant signal in the analysis of fitness effects across

the accessory genome (Kpn01, Kpn08, and Kpn13; LIPA, P< 0.05). The results thus reveal that pOXA-

48 tended to produce a high cost in *K. pneumoniae* clones belonging to ST1427. Interestingly, although

K. pneumoniae ST1427 is relatively common in our hospital (4.8% of ESBL-producing *K.*

pneumoniae[38]), none of the 103 pOXA-48-carrying *K. pneumoniae* isolates recovered in the R-GNOSIS

collection belong to this ST[31] (Fisher's exact test for count data, 8/166 vs 0/103, P= 0.025). These

results suggest that the high cost associated with plasmid acquisition in this clade may limit in-hospital

spread of pOXA-48-carrying *K. pneumoniae* ST1427. Conversely, pOXA-48 is commonly associated

with *K. pneumoniae* ST11 in our hospital[31,36], and in the four ST11 clones tested in this study, pOXA-48

produced neutral (Kpn07, Kpn20, Kpn23) or even beneficial fitness effects (Kpn22, Figure 4A) (pOXA-

48 fitness effects in ST1427 [n=5] vs. in ST11 [n=4], Welch's unequal variances two-tailed t-test, t= -

2.39, df= 7, P= 0.048).

Figure 4. Fitness effects of pOXA-48 across bacterial genome content. An association was found

between pOXA-48 fitness effects and bacterial host genomic content for four *K. pneumoniae* ST1427

isolates. (a) Core genome relationships among *E. coli* (left) and *Klebsiella* spp. (right). Tree construction

is based on polymorphisms in the core genome. The outer circle indicates the relative fitness of pOXA-

48-carrying bacterial hosts (see legend for colour code; red indicates fitness costs and green indicates

fitness benefits associated with pOXA-48 carriage). Asterisks denote clones with a phylogenetic signal

associated with plasmid fitness effects (LIPA, P< 0.05). (b) Accessory genome relationships among *E.*

coli (left) and *Klebsiella* spp. (right). Tree construction is based on the distance matrix of the accessory

gene network of each group. The outermost circle indicates relative fitness as in (a). The intermediate

circles indicate presence/absence of plasmids belonging to the different plasmid families named in the

figure. Asterisks denote clones with a significant phylogenetic signal associating accessory genome

composition with pOXA-48 fitness effects (LIPA, P< 0.05).

Modelling the role of DFE in plasmid stability

In general, mathematical models of plasmid population biology consider a clonal population in which the plasmid produces a constant reduction in growth rate[14,18–23]. These models usually include the rate of plasmid loss through segregation[49,50] and the rate of horizontal plasmid transfer by conjugation[19,20,51], and some of them also incorporate a rate of compensatory mutations that alleviate plasmid fitness costs over time[14,23]. Our results show that plasmids produce a wide DFE in naturally compatible bacterial hosts, and this distribution could strongly influence plasmid stability in polyclonal bacterial communities. To assess the effect of the DFE on plasmid stability in bacterial communities, we developed a simple mathematical model based on Stewart and Levin's pioneering work on plasmid existence conditions[19].

The model describes the population dynamics of multiple subpopulations competing for a single exhaustible resource in well-mixed environmental conditions, assuming that transition between plasmid-bearing and plasmid-free cells is driven by segregation events. The growth rate of each subpopulation is determined by a substrate-dependent Monod term that depends on the extracellular resource concentration, and therefore each strain can be described by two structurally identifiable parameters[49]: the resource conversion rate (ρ) and the specific affinity for the resource (V_{max}/K_m). These parameters were estimated from the optical densities of each strain growing in monoculture (with and without plasmids) using a Markov chain Monte Carlo (MCMC) method with a Metropolis-Hastings sampler (See Methods, Figure 5a and Supplementary Figure 5).

By solving the system of differential equations (described in Methods), we were able to evaluate the final frequency of plasmid-bearing cells in an experiment of duration *T* time units and quantify the fitness effect of the plasmid on the strain. Figure 5b shows the DFE obtained after performing *in silico* pair-wise competition experiments between plasmid-bearing and plasmid-free subpopulations (with parameter values shown in Supplementary Table 5, Supplementary Figure 6), resulting in a theoretical DFE (*w*= 0.985, var= 0.0070) that is consistent with the experimentally measured DFE presented in Figure 3 (*w*= 0.971, var= 0.0072). Moreover, comparison of model predictions with relative fitness values obtained by flow cytometry are consistent (R^2= 0.603; Figure 5c), showing that the population dynamics model can accurately predict the outcome of a competition experiment from the individual growth dynamics.

Previous studies showed that the probability of plasmid fixation is correlated with the rate of horizontal

transmission[19,20,49]. As previous models, we consider horizontal transmission of plasmids as a function

of the densities of donor and recipient cells, with conjugation events occurring at a constant rate.

Competition experiments for a range conjugation rates are illustrated in Figure 5d; while at low

horizontal transmission rates plasmid-free cells outcompete plasmid-bearing cells, at higher conjugative

rates, plasmid-bearing cells increase in frequency.

Figure 5. Modelling the DFE for pOXA-48. (a) Distribution of parameter values obtained using Bayesian

inference to estimate growth kinetic parameters from OD measurements obtained for each strain in

isolation. Diamonds represent *Klebsiella* spp. strains and circles *E. coli* clones; filled symbols denote

plasmid-bearing strains and empty symbols plasmid-free cells. The ellipses represent standard

deviations of best-fit Normal distributions (green for plasmid-bearing strains and orange for plasmid-free

cells). (b) Bars represent a DFE obtained from *in silico* competition experiments with parameter values

determined from experimental growth curves. The solid curve represents the computationally estimated

DFE obtained by randomly sampling wild-type and transconjugant parameter distributions obtained using the MCMC algorithm and numerically solving the model to evaluate the relative fitness associated with plasmid carriage. (c) Comparison of relative fitness values obtained experimentally and using the population dynamics model (R^2= 0.603). (d) Fraction of plasmid-bearing cells as a function of the rate of horizontal transfer for random plasmid-host associations sampled from the MCMC parameter distribution. The dotted line illustrates the mean of 10^4 pair-wise competition experiments under the assumption plasmid-bearing is associated with a constant reduction in fitness in different clones (*w*= 0.985, var= 0), while the solid line is obtained by considering a wide DFE (*w*= 0.985, var= 0.0070). The arrow denotes the difference in the conjugation threshold that positively selects for plasmids in the population, supporting the tenet that the DFE maintains plasmids in the population at lower conjugation rates.

Community complexity promotes plasmid persistence

To explore how plasmid stability is affected by increasing community complexity and rates of horizontal transmission, we randomly sampled *N*= 10^4 plasmid-free cells from the distribution of growth parameters estimated using the MCMC algorithm. These random communities were used to study the population dynamics of plasmids transmitting vertically and horizontally in multi-strain communities. The fitness cost (or benefit) of bearing plasmids was modelled as a random variable that modifies the wild-type (plasmid-free) growth rate by a factor σ, such that if σ= 0, the DFE has zero variance (Figure 6a), but if σ> 0, the resulting DFE is a symmetrical heavy-tailed distribution with a right-hand tail expanding towards positive fitness effects (Figure 6b), indicating the existence of plasmid-host associations in which plasmid carriage produces a fitness benefit.

To assess how DFE influences plasmid persistence in polymicrobial communities, we extended the model to consider populations composed of subsets of 1, 2, 3, 4, ..., *M* ≤ *N* cell types sampled randomly from the wild-type parameter distribution (see Methods and Supplementary Figure 7 and 8). This enabled us to estimate the relative frequency of plasmid-bearing cells at the end of a long-term experiment and evaluate the stability of the plasmid in multi-strain communities with different population structures. Initial bacterial densities were determined by first running the system forward (with all strains initially present at equal densities and carrying pOXA-48) for *T*= 24 time units, and then clearing all plasmid-free cells from the population. This assumption is akin to patients receiving an antibiotic

therapy that clears all plasmid-free cells from the bacterial community. The results obtained after 5,000

computer simulations over a range of conjugation rates and numbers of cell types in the community are

shown in Figure 6. The simulations either assumed an identical fitness cost for all strains (w= 0.985,

var= 0; Figure 6a,c) or allowed plasmid fitness effects to vary according to the experimentally

determined DFE (w= 0.985, var= 0.0070; Figure 6b,d).

Although the mean fitness cost was the same in both conditions, the results of the computational

experiments suggest that allowing fitness effects to vary between members of the population markedly

increases the chances of plasmid persistence, especially at low conjugation rates. More importantly, in

the numerical simulations, plasmid frequency decreased as a function of the number strains in the

community when plasmid acquisition was associated with a constant fitness cost, but increased with

community complexity for DFEs with larger variance (Figure 6c,d,e). The explanation for this effect is

that if plasmid fitness cost is identical for all community members, diversity simply means extra

competition for plasmid-carrying cells, and plasmid persistence becomes more dependent on a high

conjugation rate. In contrast, if the fitness effects vary, a larger number of available bacterial hosts in

the population increases the probability of the plasmid arriving to a host in which it produces a neutral

or beneficial fitness effect. This is an important result because it implies that increasing bacterial

community complexity could increase the probability of plasmid persistence in natural environments.

Given that most natural microbiota are complex and plasmids can usually conjugate and replicate in

different clones, this may explain the high prevalence of plasmids in nature. Our results also indicate

that the threshold conjugation rate for plasmid persistence may be lower than previously thought. In

fact, once plasmids are present in multiple members of a community, they may be able to persist even

in the absence of conjugation (Figure 6d).

Figure 6. Modelling plasmid persistence in polymicrobial communities, assuming fixed (a,c) or variable (b,d) plasmid fitness effects. (a-b) Relative fitness histogram obtained by randomly sampling 10^4 parameter values from the parameter distribution shown in the inset plot (points illustrate the expected values of each distribution and ellipses their standard deviation; green, plasmid-bearing bacteria; orange, plasmid-free bacteria). The green ellipse in b is larger as a consequence of considering that the cost of plasmid-bearing is normally-distributed with variance 0.007. As a result, the DFE also has higher variance, with a considerable fraction of plasmid-host associations producing a benefit to the host.

Dotted red lines indicate mean relative fitness of plasmid carrying cells. (c-d) Colour gradient represents

the percentage of cells carrying plasmids at the end of 5,000 stochastic simulations; orange indicates a

population without plasmids and green a community composed of plasmid-carrying cells. If plasmid-

bearing is associated with a fixed fitness cost for all members of the community, plasmid maintenance

requires a high conjugation rate. The increased proportion of plasmid-bearing cells in d indicates that a

DFE with high variance reduces the critical conjugation rate needed to maintain plasmids in the

population, enabling plasmids to persist at low conjugation rates. (e) Mean fraction of plasmid-bearing

cells as a function of the number of strains in the community with a conjugation rate $\gamma = 1.5 \times 10^{-11}$. If

the plasmid always produces a reduction in host fitness (mean w <1 and low variance), plasmid

frequency decreases as the number of strains in the community increases (green line). In contrast, for

higher variance at the same mean w, the fraction of plasmid-bearing cells increases with community

complexity (orange line).

Discussion

The DFE for new mutations is a central concept in genetics and evolutionary biology, with implications

ranging from population adaptation rates to complex human diseases[52]. The fitness effects of new

spontaneous mutations in bacteria follow a heavy-tailed distribution dominated by quasi-neutral

mutations with infrequent strongly deleterious mutations[53,54]. Horizontally acquired genes also produce

a distribution of fitness effects in new bacterial hosts[55,56]. However, horizontal gene transfer in bacteria

is frequently mediated by entire mobile genetic elements, such as plasmids, that carry multiple genes.

Numerous studies have measured the fitness effects of individual plasmids in a bacterial host[24], but the

DFE of a plasmid in multiple, ecologically compatible bacterial hosts had not been reported before.

Here, we determined the DFE of a carbapenem resistance plasmid in wild-type enterobacteria

recovered from the human gut microbiota. Unsurprisingly, the DFE of pOXA-48 differed from the DFE of

spontaneous mutations. As with spontaneous mutations, the pOXA-48 DFE was also dominated by

quasi-neutral effects and was slightly shifted towards fitness costs; however, instead of a single heavy

tail of deleterious effects, it had a symmetrical shape, with tails expanding both towards negative and

positive fitness effects (Figure 3a).

Two key implications of the experimentally determined DFE in this study are that, according to our

simple mathematical model, the probability of plasmid persistence becomes less dependent on a high

conjugation rate and increases with the number of bacterial strains in the population. The complex and

multi-clonal nature of most natural bacterial communities attests the likely relevance of our findings to

the extremely high prevalence of plasmids in bacterial populations[57]. The human gut microbiota, for

example, includes a great variety of bacteria from hundreds of species[58], including several strains from

the *Enterobacterales* order alone[59]. Our experimental system is in fact inspired by the dynamics of

pOXA-48 in the gut microbiota of hospitalised patients. In a recent study, we observed that once

patients are colonised by a pOXA-48-carrying clone, the plasmid spreads through conjugation to other

resident enterobacteria present in the gut microbiota[31]. Crucially, pOXA-48 usually persists in the gut of

patients throughout the hospital stay and can be detected in subsequent hospital admissions months or

years later, and not necessarily in the original colonizing strain[31]. Our results indicate that the pOXA-48

DFE could explain the long-term persistence of this and other plasmids in the human gut microbiota.

Another interesting result of this study is that pOXA-48 produced a particularly elevated cost in *K.*

pneumoniae isolates belonging to ST1427 (Figure 4A). ST1427 is under-represented among the pOXA-

48-carrying *K. pneumoniae* isolates in our hospital, which are dominated by ST11[31,36]. Remarkably, in

the four *K. pneumoniae* ST11 clones tested in this study, pOXA-48 produced neutral (Kpn07, Kpn20,

Kpn23) or even beneficial fitness effects (Kpn22, Figure 4A). Therefore, despite the small number of *K.*

pneumoniae clones analysed, our results suggest that phylogeny might influence fitness compatibility

between plasmids and bacteria at the clonal level, dictating the epidemiology of plasmid-bacterium

associations in clinical settings. Further analysis of a larger sample of *K. pneumoniae* isolates from the

different STs will be needed to elucidate the genetic basis underlying these specific interactions

between bacterial phylogeny and pOXA-48 fitness effects.

The main experimental limitation of our study is that plasmid fitness effects were determined *in vitro*,

using planktonic cultures in LB medium. This is the standard practise in the field, and previous studies

have shown that plasmid fitness effects measured in laboratory conditions correlate with those

measured in animal models[24]; however, our results may not be fully representative of pOXA-48 fitness

effects in the human gut. Future studies will need to explore more complex *in vitro* systems[60], as well as

in vivo animal models[61]. Another important limitation of our study is that we modelled bacterial

communities with a simple resource competition model that does not consider spatial structure[62],

complex ecological interactions between community members[63], plasmid-host co-evolution[64], or

differential rates of horizontal transmission[28]. Although more complex models[65] will be needed to

integrate bacterial community complexity and plasmid fitness effects, consideration of diverse

polymicrobial populations with complex spatiotemporal interactions would likely only increase DFE

variance, therefore promoting plasmid stability.

Methods

Strains, pOXA-48 plasmid, and culture conditions

We selected 50 representative ESBL-producing clones form the R-GNOSIS collection (Supplementary

Table 1). This collection was constructed in our hospital as part of an active surveillance-screening

program for detecting patients colonised by ESBL/carbapenemase-producing enterobacteria, from

March 4th, 2014, to March 31st, 2016 (R-GNOSIS-FP7-HEALTH-F3-2011-282512, www.r-gnosis.eu/,

approved by the Ramón y Cajal University Hospital Ethics Committee, Reference 251/13)[36,38]. The

screening included a total of 28,089 samples from 9,275 patients admitted at 4 different wards

(gastroenterology, neurosurgery, pneumology and urology) in the Ramon y Cajal University Hospital

(Madrid, Spain). The characterisation of samples was performed during the R-GNOSIS study

period[36,66]; rectal swabs were plated on Chromo ID-ESBL and Chrom-CARB/OXA-48 selective agar

media (BioMérieux, France) and bacterial colonies able to grow on these media were identified by

MALDI-TOF MS (Bruker Daltonics, Germany) and further characterized by pulsed-field gel

electrophoresis (PFGE). For the present study, we selected 25 *E. coli* and 25 *K. pneumoniae* ESBL-

producing isolates from the R-GNOSIS collection. The strains were representative of *E. coli* and *K.*

pneumoniae diversity in the R-GNOSIS collection (randomly chosen form the most common pulsed-

field gel electrophoresis profiles[38]), they did not carry any carbapenemase gene and they were

recovered from patients not colonised by other pOXA-48-carrying clones. To construct the

transconjugants, we used the most common pOXA-48 plasmid variant from the R-GNOSIS collection in

our hospital, according to plasmid genetic sequence (pOXA-48_K8, accession number MT441554)[30].

Bacterial strains were cultured in lysogeny broth (LB) at 37°C in 96-well plates with continuous shaking

(250 rpm) and on LB agar plates at 37°C.

Construction of transconjugants collection

We performed an initial conjugation round to introduce pOXA-48_K8 plasmid from wild type *E. coli*

C609 strain[31], into *E. coli* β3914[67], a diaminopimelic acid (DAP) auxotrophic laboratory mutant of *E. coli*

K-12 (kanamycin, erythromycin and tetracycline resistant, Supplementary Table 1), which was used as

the common counter-selectable donor. The pOXA-48-carrying wild type *E. coli* C609 and *E. coli* β3914

were streaked from freezer stocks onto solid LB agar medium with ertapenem 0.5 µg/ml and DAP 0.3

mM, respectively, and incubated overnight at 37°C. Donor and recipient colonies were independently

inoculated in 2 ml of LB in 15-ml culture tubes and incubated overnight. After growth, donor and

recipient cultures were collected by centrifugation (15 min, 1,500 g) and cells were re-suspended in

each tube with 300 µl of sterile NaCl 0.9%. Then, the suspensions were mixed in a 1:1 proportion,

spotted onto solid LB medium with DAP 0.3 mM and incubated at 37ºC overnight. Transconjugants

were selected by streaking the conjugation mix on LB with ertapenem (0.5 µg/ml), DAP 0.3 mM,

tetracycline (15 µg/ml), and kanamycin (30 µg/ml). The presence of pOXA-48 was checked by PCR,

using primers for bla_{OXA-48} gene and for the replication initiation protein gene *repC* (Supplementary

Table 4).

We used the counter-selectable *E. coli* β3914/pOXA-48_K8 donor to conjugate plasmid pOXA-48 in the

50 wild type strains. We used the same protocol described above, but the final conjugation mix was

plated on LB with no DAP (to counter-select the donor) and with amoxicillin-clavulanic acid (to select for

transconjugants). The optimal concentration of amoxicillin-clavulanic acid was experimentally

determined for each isolate in the collection and ranged from 64 µg/ml to 384 µg/ml. The presence of

pOXA-48 in the transconjugants was checked by PCR, as described above, and by antibiotic

susceptibility testing and whole genome sequencing (see below). To test the stability of plasmid pOXA-

48 in the transconjugants we propagated cultures in LB with no antibiotic selection (two consecutive

days, 1:10,000 dilution) and plated cultures on LB agar. After ON incubation at 37ºC, 100 independent

colonies of each transconjugant were replicated both on LB agar and LB agar with amoxicillin-

clavulanic acid to identify pOXA-48-carrying colonies (including negative controls of plasmid-free wild

type clones). Results showed that the plasmid was overall stable in the transconjugants; 100% stable in

43 isolates, and ≥ 90% stable in the 7 remaining isolates.

Antibiotic susceptibility testing

Antibiotic susceptibility profiles were determined for every wild-type and transconjugant strain by the

disc diffusion method following the EUCAST guidelines (www.eucast.org) (Supplementary Table 2). We

used the following antimicrobials agents: imipenem (10 µg), ertapenem (10 µg), amoxicillin-clavulanic

acid (20/10 µg), rifampicin (30 µg), streptomycin (300 µg), chloramphenicol (30 µg) and amikacin (30

µg) (Bio-Rad, CA, USA). pOXA-48-carrying and pOXA-48-free strains were pre-cultured in Müller-

Hinton (MH) broth at 37 ºC in 15 ml test tubes with continuous shaking (250 rpm), and disc diffusion

antibiograms were performed on MH agar plates (BBL, Becton Dickinson, MD, USA).

Growth curves

Pre-cultures of plasmid-free and plasmid-carrying strains (5 replicates of each) were prepared by

inoculating single independent colonies into LB broth and overnight incubation at 37 ºC with continuous

shaking (250 rpm). Overnight cultures were diluted 1:1,000 into fresh LB in 96-well plates, which were

incubated during 22 h at 37 ºC with shaking (250 rpm) in a plate reader (Synergy HTX Multi-Mode

Reader, BioTek Instruments, Inc, VT, USA). Optical densities (OD) were measured every 15 minutes

during the incubation. The maximum growth rate (μ_{max}), maximum optical density (OD_{max}), and area

under the growth curve (AUC) were determined using Gen5™ Microplate Reader and Imager Software

and the *growthrates* package in R. We calculated the relative OD_{max}, μ_{max}, and AUC, by dividing the

average value of each parameter for the pOXA-48-carrying isolate between that of the pOXA-48-free

isolate using the follow formula:

$$Relative_{ODmax,Vmax,AUC} = \frac{Plasmid - carrying\ _{ODmax,Vmax,AUC}}{Plasmid - free\ _{ODmax,Vmax,AUC}}$$

Construction of pBGC, a GFP-expressing non-mobilizable plasmid.

To fluorescently label the wild type isolates for competition assays using flow cytometry, we constructed

the pBGC plasmid, a non-mobilizable version of the *gfp*-carrying small plasmid pBGT[68] (Supplementary

Figure 2, accession number MT702881). The pBGT backbone was amplified, except for the region

including the oriT and *bla*$_{TEM1}$ gene, using the pBGC Fw/Rv primers. The *gfp* terminator region was

independently amplified using the GFP-Term Fw/Rv primers (Supplementary Table 4). PCR

amplifications were made with Phusion Hot Start II DNA Polymerase at 2 U/µL (ThermoFisher

Scientific, MA, USA), and PCR products were digested with DpnI to eliminate plasmid template before

setting up the assembly reaction (New England BioLabs, MA, USA). Finally, pBGC was constructed by

joining the amplified pBGT backbone and the *gfp* terminator region using the Gibson Assembly Cloning

Kit (New England BioLabs, MA, USA). Resulting reaction was transformed by heat shock into NEB 5-

alpha Competent *E. coli* (New England BioLabs, MA, USA), following manufacturer's instructions.

Transformation product was plated on LB agar with arabinose 0.1% and chloramphenicol 30 μg/ml, and

incubated overnight at 37 ºC. Plasmid-bearing colonies were selected by green fluorescence. The *gfp*

gene in pBGC is under the control of the P_{BAD} promoter, so GFP production is generally repressed and

induced by the presence of arabinose. pBGC was completely sequenced using primers described in

Supplementary Table 4. We confirmed that neither pOXA-48, nor helper plasmid pTA-Mob[69], could

mobilized pBGC by conjugation using the conjugation protocol described above, confirming that pBGC

plasmid is not mobilizable. Finally, pBGC plasmid was introduced into our isolate collection by

electroporation (Gene Pulser Xcell Electroporator, BioRad, CA, USA). Of note, we were not able to

obtain pBGC-carrying transformants in eight of the isolates due to a pre-existing high chloramphenicol

resistance phenotype.

Competition assays using flow cytometer

We performed competition assays[41], using flow cytometry, to obtain the relative fitness of pOXA-48-

carrying isolates compared to their pOXA-48-free parental counterparts. We used the collection of

pBGC transformed wild type isolates as competitors against their isogenic pOXA-48-carrying and

pOXA-48-free isolates. Specifically, two sets of competitions were performed for each isolate: pOXA-

48-free vs. pBGC-carrying, and pOXA-48-carrying vs. pBGC-carrying. Five biological replicates of each

competition were performed. Pre-cultures were incubated overnight in LB in 96-well plates at 225 rpm

an 37ºC, then mixed 1:1 and diluted 10,000-fold in 200 μl of fresh LB in in 96-well plates. Mixtures were

competed for 24 h in LB at 37ºC and 250 rpm (the low initial cell density and the strong shaking hinders

pOXA-48 conjugation, see control experiment below). To determine the initial proportions, initial 1:1

mixes were diluted 2,000-fold in 200 μl of NaCl 0.9 % with L-arabinose 0.1 %, and incubated at 37 ºC at

250 rpm during 1.5 h to induce *gfp* expression. The measurements were performed via flow cytometry

using a CytoFLEX Platform (Beckman Coulter Life Sciences, IN, US) with the following parameters: 50

μl min⁻¹ flow rate, 22 μm core size, and 10,000 events recorded per sample (Supplementary Figure 9).

After 24 hours of incubation, final proportions were determined as described above, after 2,000-fold

dilution of the cultures. The fitness of each strain relative to its pBGC-carrying parental isolate was

determined using the formula:

$$w = \frac{\ln(N_f / N_i)}{\ln(N_{f,pBGC+} / N_{i,pBGC+})}$$

where w is the relative fitness of the pOXA-48-carrying ($w_{pOXA-48+}$) or pOXA-48-free ($w_{pOXA-48-}$) isolates

compared to the pBGC-bearing parental clone, N_i and N_f are the number of cells of the pBGC-free

clone at the beginning and end of the competition, and $N_{i, pBGC}$ and $N_{f, pBGC}$ are the number of cells of the

pBGC-carrying clone at the beginning and end of the competition, respectively. The fitness of the

pOXA-48-carrying isolates relative to the pOXA-48-free parental isolates were calculated with the

formula, $w_{pOXA-48+}$ / $w_{pOXA-48-}$ to correct for the fitness effects of pBCG (see Supplementary Figure 10 for

pBGC fitness effects), and the error propagation method was used to calculate the standard error of the

resulting value. Note that the fitness effects of pBGC did not correlate with those form pOXA-48

(Pearson's correlation, R= 0.11, t= 0.66, df= 39, P= 0.51). For the 8 strains where pBGC plasmid could

not be introduced (Ec13, Kpn10, Kpn11, Kpn19-Kpn23), pOXA-48-carrying and pOXA-48-free isolates

were competed against a pBGC-carrying $E. coli$ J53[70] (a sodium azide resistant laboratory mutant of $E.$

$coli$ K-12), following the same protocol described above. In general, we prefer to perform competitions

assays between isogenic bacteria to avoid interactions between clones that may affect the outcome of

the competition for reasons beyond the presence of the plasmid under study (such as bacteriocin

production). However, we did not observe any evidence of growth inhibition between the 8 wild type

isolates and $E. coli$ J53 in the flow cytometry data, and the relative fitness results obtained with these

competitions were comparable to those obtained in the isogenic competitions (two-tailed t-test, t= 1.64,

df= 11.2, P= 0.13). To confirm that the isogenic competitions and those against $E. coli$ J53/pBGC

produced similar results, we selected 10 random isolates from the 42 isolates with fitness data

calculated from isogenic competitions, and repeated their competitions against $E. coli$ J53/pBGC

(Supplementary Figure 11). Results showed that relative fitness values calculated with isogenic

competitions and those using $E. coli$ J53/pBGC presented a good correlation (Pearson's correlation, R=

0.81, t= 3.96, df= 8, P= 0.004, Supplementary Figure 11). Finally, we performed controls to test for the

potential conjugative transfer of pOXA-48 during head-to-head competitions by plating the final time

points of the competition assays on amoxicillin-clavulanic acid (with the adequate concentration for

each isolate), and chloramphenicol (30 µg/ml). No transconjugants were detected in these controls,

showing that the low initial inoculum size we used in the competitions (10,000-fold dilution), and the

vigorous shaking of the liquid cultures prevented pOXA-48 conjugation.

DNA extraction and genome sequencing

Genomic DNA of all the pOXA-48 bearing strains was isolated using the Wizard genomic DNA

purification kit (Promega, WI, USA), and quantified using the QuantiFluor dsDNA system (Promega, WI,

USA), following manufacturers' instructions. Whole genome sequencing was conducted at the

Wellcome Trust Centre for Human Genetics (Oxford, UK), using the Illumina HiSeq4000 platform with

125 base pair (bp) paired-end reads and at MicrobesNG (Birmingham, UK), using Illumina platforms

(MiSeq or HiSeq2500) with 250 bp paired-end reads.

Bioinformatic analyses

The Illumina sequence reads were trimmed using the Trimmomatic v0.33 tool[71]. SPAdes v3.9.0[72] was

used to generate *de novo* assemblies from the trimmed sequence reads with the –cov-cutoff flag set to

'auto'. QUAST v4.6.0[73] was used to generate assembly statistics. Three genomes were dropped from

the analysis because of the poor quality of the sequences (2 *E. coli* [Ec09, Ec17] and 1 *K. pneumoniae*

[Kpn05]). All the *de novo* assemblies used reached enough quality including total size of 5–7 Mb, and

the total number of contigs over 1 kb was lower than 200. Prokka v1.5[74] was used to annotate the *de*

novo assemblies with predicted genes. The seven-gene ST of all the isolates was determined using the

multilocus sequence typing (MLST) tool (https://github.com/tseemann/mlst). The plasmid content of

each genome was characterised using PlasmidFinder 2.1[75], and the antibiotic resistance gene content

was characterised with ResFinder 3.2[76] (Supplementary Table 1).

In order to confirm the presence of the entire pOXA-48_K8 plasmid, the sequences belonging to pOXA-

48 plasmid in the transconjugants were mapped using as reference the complete sequence of plasmid

from the donor strain, which had been previously sequenced by PacBio[31] (from *K. pneumoniae* k8 –

GenBank Accession Number MT441554). Snippy v3.1 (https://github.com/tseemann/snippy) was used

to check that no SNPs or indels accumulated in pOXA-48_K8 during strain construction . Coding

sequences in pOXA-48 were predicted and annotated using Prokka 1.14.6 software[74]. Plasmid

annotation was complemented with the National Center for Biotechnology Information (NCBI)

Prokaryotic Genome Annotation Pipeline[77].

To determine distances between genomes we used Mash v2.0[78] with the raw sequence reads, and a

phylogeny was constructed with mashtree v0.33[79]. For the analysis of the core genome we calculated

the genetic relatedness of isolates belonging to *Klebsiella* spp. and to *E. coli* by reconstructing their

core genome phylogeny with an alignment of the SNPs obtained with Snippy v3.1

(https://github.com/tseemann/snippy). A maximum-likelihood tree was generated using IQ-TREE with

automated detection of the best evolutionary model[80]. The tree was represented with midpoint root

using the phylotools package in R (https://github.com/helixcn/phylotools) and visualised using the iTOL

tool[81]. We also constructed a distance matrix of the accessory gene network to analyse the accessory

genome. To this end, we used AccNET, a tool that allows to infer the accessory genome from the

proteomes and cluster them based on protein similarity[46]. The set of representative proteins was used

to build a binary matrix (presence/absence of proteins in the accessory genome) in the R-environment

and a cladogram to classify the strains according to the accessory genomes. The Euclidean distance

was calculated by the 'dist' function and a hierarchical clustering was performed with UPGMA using the

'hclust' function in the R environment. This cladogram was represented with midpoint root using the

phylotools package in R (https://github.com/helixcn/phylotools) and visualised using the iTOL tool[81].

Analysis of plasmid fitness effects across bacterial phylogeny

We tested for the presence of phylogenetic signal in core and accessory genomes of *E. coli* and *K.*

pneumoniae using several statistical tests available in the *phylosignal* R package[47]. In essence, these

analyses are designed to identify statistical dependence between a given continuous trait (relative

fitness) and the phylogenetic tree of the taxa from which the trait is measured. Therefore, a positive

phylogenetic signal indicates that there is a tendency for related taxa to resemble each other[82]. Several

indices have been proposed to identify phylogenetic signal, but the choice among them is not

straightforward[83]. We first assayed the methods implemented in the *phyloSignal* function, which

produce global measures of phylogenetic signal (*i.e.* across the whole phylogeny). The methods

employed were Abouheif's C_{mean}, Moran's I index, Bloomberg's K and K*, and Pagel's λ[47]. All methods

except Pagel's λ detected a marginally significant phylogentic signal in the *K. pneumoniae* core genome

(Supplementary table 3 [first tab]; 0.11>P>0.02). Abouheif's C_{mean} and Moran's I (but not Bloomberg's K

and K*, and Pagel's λ) also detected a marginally significant signal in the *K. pneumoniae* accessory

genome tree (Supplementary table 3 [first tab]; P<0.056 for both cases). Intrigued by these results, we

used the Local Indicator of Phylogenetic Association (LIPA) based on local Moran's I, which is meant to

detect local hotspots of phylogenetic signal[47,48]. LIPA, implemented in the *lipaMoran* function, computes

local Moran's I indexes for each tip of the phylogeny and a non-parametric test to ascertain statistical

significance (Supplementary Figure 4 and Supplementary table 3 [second tab]).

Plasmid population dynamics model

We used a simple mathematical model of microbial growth under resource limitation to study the role of

the DFE in the ecological dynamics of a plasmid spreading in a bacterial population[14]. Bacterial growth

rate was modelled as a saturating function of the environmental resource concentration, R,

$$G(R) = \rho \cdot \frac{V_{max}R}{K_m+R} = \rho \cdot u(R),$$

where ρ denotes the cell's efficiency to convert resource molecules into biomass and $u(R)$ a resource

uptake function that depends on the maximum uptake rate (V_{max}) and a half-saturation constant (K_m).

If we denote with B_p the density of plasmid-bearing cells and with B_0 the density of plasmid-free cells

(each with its own growth kinetic parameters and growth functions denoted $G_p(R)$ and $G_0(R)$,

respectively), then the density of each subpopulation can be described by a system of ordinary

differential equations:

$$\frac{dR}{dt} = -u_p(R) - u_0(R) - dR,$$

$$\frac{dB_p}{dt} = (1-\lambda)\,G_p(R)B_p + \gamma B_0 B_p - dB_p,$$

$$\frac{dB_0}{dt} = G_0(R)B_0 + \lambda\,G_p(R)B_p - \gamma B_0 B_p - dB_0.$$

where λ represents the rate of segregational loss rate and d a dilution parameter. Moreover, we

represent with γ the rate of conjugative transfer, and therefore we model plasmid conjugation as a

function of the densities of donor and recipient cells. By numerically solving the system of equations

(using standard differential equations solvers in Matlab), we obtain the final density of each bacterial

type in an experiment of $T = 24$ units of time with $d = 0$ (to replicate the batch culture conditions used

to estimate the DFE experimentally).

Growth kinetic parameters were determined with a Markov chain Monte Carlo method (MCMC; scripts

coded in R and available at http://www.github.com/esb-lab/pNUK73/) applied to growth curves of each

strain growing in isolation, with and without plasmids. This data fitting algorithm implements a

Metropolis-Hastings sampler with a burn-in parameter of 0.2 and executed for 5×10^6 iterations, or until

achieving convergence of the Markov chains (see Supplementary Figure 5 for an example and

Supplementary Table 5 for parameters values estimated for each strain).

Stochastic simulations of polymicrobial communities

Numerical experiments were performed by randomly sampling $N = 1 \times 10^4$ cells from the parameter

distribution obtained after applying the MCMC algorithm to all 50 strains and fitting a bivariate Normal

distribution. We then assembled $5{,}000$ synthetic communities composed of a random subset of $M < N$

different strains sampled from this distribution, and solved a multi-strain extension of the population

dynamics model. For each numerical experiment, the total density of strain i would be $B^i(t) = B_p^i(t) +$

$B_0^i(t)$, where B_p^i and B_0^i denote, respectively, the densities of plasmid-bearing and plasmid-free cells of

type $1 \le i \le M$. To model the fitness effects of bearing plasmids, we introduced a parameter, σ, such

that when $\sigma = 0$, the fitness difference between B_p^i and B_0^i corresponds to a fixed reduction in growth

rate (corresponding to a DFE with variance 0 and mean $w = 0.985$). Conversely, if $\sigma > 0$, then growth

kinetic parameters for each plasmid-bearing strain in the community were determined by sampling s_i

from a Normal distribution, $N(0, \sigma^2)$, and multiplying both V_{max}^i and ρ^i by a factor of $(1 + s_i)$.

As with the single-strain model, we consider segregational loss as a transition from B_p^i to B_0^i occurring

at a rate λ, but now we also consider that plasmid-free cells can acquire plasmids via conjugation from

any plasmid-bearing strain in the community, at a constant rate γ, and with equal probability of

transferring between different bacterial hosts. Therefore, we obtain a system of $2M + 1$ differential

equations that can be written, for each strain i, as follows:

633
$$\frac{dB_p^i}{dt} = (1 - \lambda)\, G_p^i(R)\, B_p^i \; + \gamma \sum_{j=1}^{M} B_p^j B_0^i \; - \; dB_p^i,$$

634
$$\frac{dB_0^i}{dt} = G_0^i(R)\, B_0^i \; + \lambda G_p^i(R)\, B_p^i \; - \gamma \sum_{j=1}^{M} B_p^j B_0^i \; - \; dB_0^i.$$

635 Furthermore, if $\hat{R}$ represents the input of resource into the system, then

636
$$\frac{dR}{dt} = - \sum_{i=1}^{M} \left(u_p^i(R) + u_0^i(R)\right) - d\left(R - \hat{R}\right).$$

Initial bacterial densities were determined by first running the system forward (with all strains initially

present at equal densities) for $T = 24$ time units, and then clearing all plasmid-free cells from the

population. This assumption is consistent with patients receiving antimicrobial therapy that clears all

susceptible (plasmid-free) cells from the microbiota or, in an experimental microcosm, to a round of

growth in selective media after an overnight culture. As we are interested in the long-term population

dynamics, we ran each simulation starting from the aforementioned initial condition until the plasmid

fraction was below a threshold $\epsilon > 0$ (i.e. plasmid extinction), the plasmid fraction was near 100% and

the total plasmid-free density was below ϵ (i.e. plasmid fixation), or wild-type and transconjugant sub-

populations appeared to co-exist indefinitely in the population (either in equilibrium or exhibiting

oscillatory behaviour, as illustrated in Supplementary Figures 7 and 8).

Statistical analyses

The statistical tests used are indicated in the text. Analyses were performed using R (v. 3.5.0).

References

1.   LEDERBERG, J. Cell genetics and hereditary symbiosis. *Physiol. Rev.* **32**, 403–430 (1952).

2.   Frost, L. S., Leplae, R., Summers, A. O. & Toussaint, A. Mobile genetic elements: The agents of
open source evolution. *Nature Reviews Microbiology* vol. 3 722–732 (2005).

3.   Partridge, S. R., Kwong, S. M., Firth, N. & Jensen, S. O. Mobile genetic elements associated with
antimicrobial resistance. *Clinical Microbiology Reviews* vol. 31 (2018).

4.   San Millan, A. Evolution of Plasmid-Mediated Antibiotic Resistance in the Clinical Context. *Trends in*
Microbiology (2018) doi:10.1016/j.tim.2018.06.007.

5.   Bonomo, R. A. *et al.* Carbapenemase-Producing Organisms: A Global Scourge. *Clinical Infectious*
Diseases vol. 66 1290–1297 (2018).

6.   David, S. *et al.* Epidemic of carbapenem-resistant Klebsiella pneumoniae in Europe is driven by
nosocomial spread. *Nat. Microbiol.* **4**, 1919–1929 (2019).

7.   San Millan, A. & MacLean, R. C. Fitness Costs of Plasmids: a Limit to Plasmid Transmission.
Microbiol. Spectr. (2017) doi:10.1128/microbiolspec.MTBP-0016-2017.

8.   Baltrus, D. A. Exploring the costs of horizontal gene transfer. *Trends in Ecology and Evolution* vol. 28
489–495 (2013).

9.   San Millan, A. *et al.* Integrative analysis of fitness and metabolic effects of plasmids in Pseudomonas
aeruginosa PAO1. *ISME J.* **12**, 3014–3024 (2018).

10.   Harrison, E. & Brockhurst, M. A. Plasmid-mediated horizontal gene transfer is a coevolutionary
process. *Trends in Microbiology* (2012) doi:10.1016/j.tim.2012.04.003.

11.   Bouma, J. E. & Lenski, R. E. Evolution of a bacteria/plasmid association. *Nature* **335**, 351–352
(1988).

12.   Dahlberg, C. & Chao, L. Amelioration of the Cost of Conjugative Plasmid Carriage in Eschericha coli

K12. *Genetics* **165**, 1641–1649 (2003).

13. Dionisio, F., Conceiçã, I. C., Marques, A. C. R., Fernandes, L. & Gordo, I. The evolution of a
conjugative plasmid and its ability to increase bacterial fitness. *Biol. Lett* **1**, 250–252 (2005).

14. San Millan, A. *et al.* Positive selection and compensatory adaptation interact to stabilize non-
transmissible plasmids. *Nat. Commun.* **5**, 5208 (2014).

15. Harrison, E., Guymer, D., Spiers, A. J., Paterson, S. & Brockhurst, M. A. Parallel Compensatory
Evolution Stabilizes Plasmids across the Parasitism-Mutualism Continuum. *Curr. Biol.* **25**, 2034–2039
(2015).

16. Loftie-Eaton, W. *et al.* Compensatory mutations improve general permissiveness to antibiotic
resistance plasmids. *Nat. Ecol. Evol.* (2017) doi:10.1038/s41559-017-0243-2.

17. Wein, T., Hülter, N. F., Mizrahi, I. & Dagan, T. Emergence of plasmid stability under non-selective
conditions maintains antibiotic resistance. *Nat. Commun.* **10**, 1–13 (2019).

18. Lopatkin, A. J. *et al.* Persistence and reversal of plasmid-mediated antibiotic resistance. *Nat.*
Commun. **8**, 1689 (2017).

19. Stewart, F. M. & Levin, B. R. The Population Biology of Bacterial Plasmids: A PRIORI Conditions for
the Existence of Conjugationally Transmitted Factors. *Genetics* **87**, 209–28 (1977).

20. Bergstrom, C. T., Lipsitch, M. & Levin, B. R. Natural selection, infectious transfer and the existence
conditions for bacterial plasmids. *Genetics* **155**, 1505–1519 (2000).

21. Zwanzig, M. *et al.* Mobile Compensatory Mutations Promote Plasmid Survival. *mSystems* **4**, (2019).

22. Lili, L. N., Britton, N. F. & Feil, E. J. The persistence of parasitic plasmids. *Genetics* **177**, 399–405
(2007).

23. Ponciano, J. M., De Gelder, L., Top, E. M. & Joyce, P. The population biology of bacterial plasmids: A
hidden Markov model approach. *Genetics* **176**, 957–968 (2007).

24. Vogwill, T. & Maclean, R. C. The genetic basis of the fitness costs of antimicrobial resistance: A
meta-analysis approach. *Evol. Appl.* **8**, 284–295 (2015).

25. De Gelder, L., Ponciano, J. M., Joyce, P. & Top, E. M. Stability of a promiscuous plasmid in different
hosts: no guarantee for a long-term relationship. *Microbiology* **153**, 452–463 (2007).

26. Humphrey, B. *et al.* Fitness of Escherichia coli strains carrying expressed and partially silent IncN and
IncP1 plasmids. *BMC Microbiol* **12**, 53 (2012).

27. Di Luca, M. C. *et al.* Low biological cost of carbapenemase-encoding plasmids following transfer from
Klebsiella pneumoniae to Escherichia coli. *J. Antimicrob. Chemother.* **72**, 85–89 (2017).

28. Gama, J. A., Kloos, J., Johnsen, P. J. & Samuelsen, Ø. Host dependent maintenance of a bla NDM-
1-encoding plasmid in clinical Escherichia coli isolates. *Sci. Rep.* **10**, 1–7 (2020).

29. Li, L. *et al.* Plasmids persist in a microbial community by providing fitness benefit to multiple
phylotypes. *ISME J.* **14**, 1170–1181 (2020).

30. Klümper, U. *et al.* Broad host range plasmids can invade an unexpectedly diverse fraction of a soil
bacterial community. *ISME J.* **9**, 934–945 (2015).

31. Leon-Sampedro, R. *et al.* Dissemination routes of the carbapenem resistance plasmid pOXA-48 in a
hospital setting. *bioRxiv* 2020.04.20.050476 (2020) doi:10.1101/2020.04.20.050476.

32. Dahlberg, C. *et al.* Interspecies bacterial conjugation by plasmids from marine environments
visualized by gfp expression. *Mol. Biol. Evol.* **15**, 385–390 (1998).

33. Poirel, L., Héritier, C., Tolün, V. & Nordmann, P. Emergence of Oxacillinase-Mediated Resistance to

Imipenem in Klebsiella pneumoniae. *Antimicrob. Agents Chemother.* **48**, 15–22 (2004).

34.   Pitout, J. D. D., Peirano, G., Kock, M. M., Strydom, K.-A. & Matsumura, Y. The Global Ascendency of

OXA-48-Type Carbapenemases. *Clin. Microbiol. Rev.* **33**, (2020).

35.   Ledda, A. *et al.* Hospital outbreak of carbapenem-resistant Enterobacteriales associated with an

OXA-48 plasmid carried mostly by Escherichia coli ST399. *bioRxiv* 2020.06.15.148189 (2020)

doi:10.1101/2020.06.15.148189.

36.   Hernández-García, M. *et al.* Characterization of carbapenemase-producing Enterobacteriaceae from

colonized patients in a university hospital in Madrid, Spain, during the R-GNOSIS project depicts

increased clonal diversity over time with maintenance of high-risk clones. *J. Antimicrob. Chemother.*

73, 3039–3043 (2018).

37.   Diaz-Agero, C. *et al.* Local prevalence of extended-spectrum beta-lactamase (ESBL) producing

Enterobacteriaceae intestinal carriers at admission and co-expression of ESBL and OXA-48

carbapenemase in Klebsiella pneumoniae: A prevalence survey in a Spanish University Hospital.

BMJ Open **9**, (2019).

38.   Maechler, F. *et al.* Contact isolation versus standard precautions to decrease acquisition of extended-

spectrum β-lactamase-producing Enterobacterales in non-critical care wards: a cluster-randomised

crossover trial. *Lancet Infect. Dis.* **20**, 575–584 (2020).

39.   Karanika, S., Karantanos, T., Arvanitis, M., Grigoras, C. & Mylonakis, E. Fecal Colonization with

Extended-spectrum Beta-lactamase-Producing Enterobacteriaceae and Risk Factors among Healthy

Individuals: A Systematic Review and Metaanalysis. *Clin. Infect. Dis.* **63**, 310–318 (2016).

40.   Holt, K. E. *et al.* Genomic analysis of diversity, population structure, virulence, and antimicrobial

resistance in Klebsiella pneumoniae, an urgent threat to public health. *Proc. Natl. Acad. Sci. U. S. A.*

112, E3574–E3581 (2015).

41.   Lenski, R. E., Rose, M. R., Simpson, S. C. & Tadler, S. C. *Long-Term Experimental Evolution in*

Escherichia coli. I. Adaptation and Divergence During 2,000 Generations. Source: The American

Naturalist vol. 138 (1991).

42.   Stoesser, N. *et al.* Evolutionary history of the global emergence of the Escherichia coli epidemic clone

ST131. *MBio* **7**, (2016).

43.   McNally, A. *et al.* Combined Analysis of Variation in Core, Accessory and Regulatory Genome

Regions Provides a Super-Resolution View into the Evolution of Bacterial Populations. *PLOS Genet.*

12, e1006280 (2016).

44.   San Millan, A., Heilbron, K. & MacLean, R. C. Positive epistasis between co-infecting plasmids

promotes plasmid survival in bacterial populations. *ISME J.* **8**, 601–612 (2014).

45.   San Millan, A., Toll-Riera, M., Qi, Q. & MacLean, R. C. Interactions between horizontally acquired

genes create a fitness cost in Pseudomonas aeruginosa. *Nat. Commun.* **6**, (2015).

46.   Lanza, V. F., Baquero, F., De La Cruz, F. & Coque, T. M. AcCNET (Ac cessory Genome C

onstellation Net work): Comparative genomics software for accessory genome analysis using

bipartite networks. *Bioinformatics* **33**, 283–285 (2017).

47.   Keck, F., Rimet, F., Bouchez, A. & Franc, A. Phylosignal: An R package to measure, test, and

explore the phylogenetic signal. *Ecol. Evol.* **6**, 2774–2780 (2016).

48.   Anselin, L. Local Indicators of Spatial Association-LISA. *Geogr. Anal.* **27**, 93–115 (2010).

49.   Cooper, N. S., Brown, M. E. & Caulcott, C. A. A mathematical method for analysing plasmid stability

in micro-organisms. *J. Gen. Microbiol.* **133**, 1871–1880 (1987).

50.   Paulsson, J. & Ehrenberg, M. Noise in a minimal regulatory network: Plasmid copy number control.

Quarterly Reviews of Biophysics vol. 34 1–59 (2001).

51.   Simonsen, L., Gordon, D. M., Stewart, F. M. & Levin, B. R. Estimating the rate of plasmid transfer: An

end-point method. *J. Gen. Microbiol.* **136**, 2319–2325 (1990).

52.   Eyre-Walker, A. & Keightley, P. D. The distribution of fitness effects of new mutations. *Nat. Rev.* **8**,

608–618 (2007).

53.   Bataillon, T. & Bailey, S. F. Effects of new mutations on fitness: insights from models and data. *Ann.*

N. Y. Acad. Sci. **1320**, 76–92 (2014).

54.   Robert, L. *et al.* Mutation dynamics and fitness effects followed in single cells. *Science (80-.).* **359**,

1283–1286 (2018).

55.   Vos, M., Hesselman, M. C., te Beek, T. A., van Passel, M. W. J. & Eyre-Walker, A. Rates of Lateral

Gene Transfer in Prokaryotes: High but Why? *Trends in Microbiology* vol. 23 598–605 (2015).

56.   Sorek, R. *et al.* Genome-wide experimental determination of barriers to horizontal gene transfer.

Science (80-.). **318**, 1449–1452 (2007).

57.   Acman, M., van Dorp, L., Santini, J. M. & Balloux, F. Large-scale network analysis captures biological

features of bacterial plasmids. *Nat. Commun.* **11**, 2452 (2020).

58.   Almeida, A. *et al.* A new genomic blueprint of the human gut microbiota. *Nature* **568**, 499–504 (2019).

59.   Martinson, J. N. V. *et al.* Rethinking gut microbiome residency and the Enterobacteriaceae in healthy

human adults. *ISME J.* **13**, 2306–2318 (2019).

60.   Stalder, T. *et al.* Evolving Populations in Biofilms Contain More Persistent Plasmids. *Mol. Biol. Evol.*

(2020) doi:10.1093/molbev/msaa024.

61.   Gumpert, H. *et al.* Transfer and persistence of a multi-drug resistance plasmid in situ of the infant gut

microbiota in the absence of antibiotic treatment. *Front. Microbiol.* **8**, (2017).

62.   Nadell, C. D., Drescher, K. & Foster, K. R. Spatial structure, cooperation and competition in biofilms.

Nature Reviews Microbiology vol. 14 589–600 (2016).

63.   Coyte, K. Z., Schluter, J. & Foster, K. R. The ecology of the microbiome: Networks, competition, and

stability. *Science (80-.).* **350**, 663–666 (2015).

64.   Jordt, H. *et al.* Coevolution of host–plasmid pairs facilitates the emergence of novel multidrug

resistance. *Nat. Ecol. Evol.* **4**, (2020).

65.   Lopatkin, A. J. & Collins, J. J. Predictive biology: modelling, understanding and harnessing microbial

complexity. *Nature Reviews Microbiology* (2020) doi:10.1038/s41579-020-0372-5.

66.   Hernández-García, M. *et al.* Intestinal co-colonization with different carbapenemase-producing

Enterobacterales isolates is not a rare event in an OXA-48 endemic area. *EClinicalMedicine* **15**, 72–

79 (2019).

67.   Le Roux, F., Binesse, J., Saulnier, D. & Mazel, D. Construction of a Vibrio splendidus mutant lacking

the metalloprotease gene vsm by use of a novel counterselectable suicide vector. *Appl. Environ.*

Microbiol. **73**, 777–784 (2007).

68.   San Millan, A., Escudero, J. A., Gifford, D. R., Mazel, D. & MacLean, R. C. Multicopy plasmids

potentiate the evolution of antibiotic resistance in bacteria. *Nat. Ecol. Evol.* (2016)

doi:10.1038/s41559-016-0010.

69.   Strand, T. A., Lale, R., Degnes, K. F., Lando, M. & Valla, S. A New and Improved Host-Independent

Plasmid System for RK2-Based Conjugal Transfer. *PLoS One* **9**, e90372 (2014).

70. Matsumura, Y., Peirano, G. & Pitout, J. D. D. Complete genome sequence of Escherichia coli J53, an

azide-resistant laboratory strain used for conjugation experiments. *Genome Announc.* **6**, (2018).

71. Bolger, M. A., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data |

Bioinformatics | Oxford Academic. *Bioinformatics (Oxford, England)* vol. 30 2114–2120

https://academic.oup.com/bioinformatics/article/30/15/2114/2390096 (2014).

72. Bankevich, A. *et al.* SPAdes: A new genome assembly algorithm and its applications to single-cell

sequencing. *J. Comput. Biol.* **19**, 455–477 (2012).

73. Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: Quality assessment tool for genome

assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

74. Seeman, T. Prokka: rapid prokaryotic genome annotation | Bioinformatics | Oxford Academic.

Bioinformatics **30**, 2068–2069 (2014).

75. Carattoli, A. *et al.* In Silico detection and typing of plasmids using plasmidfinder and plasmid

multilocus sequence typing. *Antimicrob. Agents Chemother.* **58**, 3895–3903 (2014).

76. Zankari, E. *et al.* Identification of acquired antimicrobial resistance genes. *J. Antimicrob. Chemother.*

67, 2640–2644 (2012).

77. Haft, D. H. *et al.* RefSeq: An update on prokaryotic genome annotation and curation. *Nucleic Acids*

Res. **46**, D851–D860 (2018).

78. Ondov, B. D. *et al.* Mash: fast genome and metagenome distance estimation using MinHash.

Genome Biol. **17**, 132 (2016).

79. Katz, L. S. *et al.* Mashtree: a rapid comparison of whole genome sequence files. (2019)

doi:10.21105/joss.01762.

80. Nguyen, L. T., Schmidt, H. A., Von Haeseler, A. & Minh, B. Q. IQ-TREE: A fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**, 268–274

(2015).

81. Letunic, I. & Bork, P. Interactive Tree of Life v2: Online annotation and display of phylogenetic trees

made easy. *Nucleic Acids Res.* **39**, (2011).

82. Blomberg, S. P., Garland, T. & Ives, A. R. Testing for phylogenetic signal in comparative data:

Behavioral traits are more labile. *Evolution (N. Y).* **57**, 717–745 (2003).

83. Münkemüller, T. *et al.* How to measure and test phylogenetic signal. *Methods Ecol. Evol.* **3**, 743–756

(2012).

Acknowledgements

This work was supported by the European Research Council under the European Union's Horizon 2020

research and innovation programme (ERC grant agreement no. 757440-PLASREVOLUTION) and by

the *Instituto de Salud Carlos III* (co-funded by European Development Regional Fund "a way to achieve

Europe") grant PI16-00860. RC acknowledges financial support from European Commission (grant R-

GNOSIS-FP7-HEALTH-F3-2011-282512) and *Plan Nacional de I+D+i2013–2016* and *Instituto de Salud*

Carlos III, Subdirección General de Redes y Centros de Investigación Cooperativa, Ministerio de

Economía, Industria y Competitividad, Spanish Network for Research in Infectious Diseases (REIPIR

D16/0016/0011) co-financed by European Development Regional Fund "A way to achieve Europe"

(ERDF), Operative program Intelligent Growth 2014–2020. ASM is supported by a Miguel Servet

Fellowship (MS15-00012). JRB is a recipient of a Juan de la Cierva-Incorporación Fellowship (IJC2018-

035146-I) co-funded by *Agencia Estatal de Investigación del Ministerio de Ciencia e Innovación*. MH-G

was supported with a contract from *Instituto de Salud Carlos III*, Spain (iP-FIS program, ref.

IFI14/00022). RPM was supported by PAPIIT-UNAM (IN209419) and CONACYT (Ciencia Básica grant

A1-S-32164). We thank the Oxford Genomics Centre at the Wellcome Centre for Human Genetics

(funded by Wellcome Trust grant reference 203141/Z/16/Z) for the generation and initial processing of

the sequencing data.

Author Contributions

ASM, AAdV and RPM conceived the study. RC designed and supervised sampling and collection

of R-GNOSIS bacterial isolates. MHG, PRG collected the bacterial isolates and performed

bacterial characterization. AAdV performed the experimental work with help from JRB and JdLF.

AAdV, JRB and JdLF analysed experimental results. RLS and JRB performed the

bioinformatic/phylogenetic analyses. RPM developed the mathematical model and computer

simulations. ASM supervised the study. ASM, AAdV and RPM wrote the initial draft of the

manuscript and all the authors contributed to the final version of the manuscript and approved it.

Competing Interests statement

Authors declare no competing interests.

Data availability

The sequences generated and analysed during the current study are available in the Sequence Read

Archive (SRA), BioProject ID: PRJNA641166, https://www.ncbi.nlm.nih.gov/sra/PRJNA641166.

Code availability

The code generated during the current study is available in GitHub, http://www.github.com/esb-lab/

pOXA48/

Supplementary Information

The distribution of plasmid fitness effects explains plasmid persistence in bacterial

communities

Aida Alonso-del Valle[1], Ricardo León-Sampedro[1,2], Jerónimo Rodríguez-Beltrán[1,2], Javier

DelaFuente[1], Marta Hernández-García[1,3], Patricia Ruiz-Garbajosa[1,3], Rafael Cantón[1,3], Rafael

Peña-Miller[4,*], Álvaro San Millán[1,2,5*].

[1] *Servicio de Microbiología. Hospital Universitario Ramón y Cajal* and *Instituto Ramón y Cajal de*

Investigación Sanitaria. Madrid, Spain.

[2] *Centro de Investigación Biológica en Red. Epidemiología y Salud Pública, Instituto de Salud Carlos*

III. Madrid. Spain.

[3] *Red Española de Investigación en Patología Infecciosa. Instituto de Salud Carlos III*. Madrid. Spain.

[4] Center for Genomic Sciences, *Universidad Nacional Autónoma de México*, Cuernavaca, Mexico.

[5] Current address: *Centro Nacional de Biotecnología–CSIC*, Madrid, Spain.

* Correspondence: Álvaro San Millán, alvsanmillan@gmail.com ORCID: 0000-0001-8544-0387 and

Rafael Peña-Miller, rafael.penamiller@gmail.com ORCID: 0000-0002-2767-0640

Supplementary Figure 1. Growth curves of wild-type isolates and pOXA-48-carrying transconjugants.

a

b

21
22

Growth curves of pOXA48-free (wild-type, blue) and pOXA-48-carrying (transconjugant, red) for

every (a) *E. coli* and (b) *Klebsiella* spp*.* analysed in this study. The lines represent the average of

five biological replicates and the shaded area indicates 95% confidence intervals.

Supplementary Figure 2. Construction of plasmid pBGC.

Schematic representation of the construction of plasmid pBGC (accession number MT702881) form

plasmid pBGT[68]. Two segments of pBGT were amplified using primers with added cohesive ends

(pBGC Fw/Rv and GFP Term Fw/Rv, Supplementary Table 4). pBGC plasmid resulted from the Gibson

assembly of the amplified fragments. The reading frames for genes are shown as arrows, with the

direction of transcription indicated by the arrowhead. The origin of replication (oriV), origin of transfer

(oriT), and P$_{BAD}$ promoter are also indicated.

Supplementary Figure 3. Correlation between relative growth curve parameters and relative fitness.
35

Correlation between relative growth curve parameters (a) maximum optical density (OD_{max}), (b)

maximum growth rate (μ_{max}), and (c) area under the growth curve (AUC), and relative fitness values

obtained from competition assays for each strain. The blue line represents the linear regression model

and the grey shading represents 95% confidence intervals. Points represent each relative value (red, *E.*

coli and blue, *Klebsiella* spp.). Pearson's correlation (R) and p-value are indicated. As expected,

maximum optical density, maximum growth rate and area under the growth curve are positively

correlated with relative fitness.

 Supplementary Figure 4. Local Indicator of Phylogenetic Association (LIPA) analyses.

46
47

Phylogenetic trees for core (upper panels) and accessory (lower panels) genomes obtained for *E. coli*

(left) and *Klebsiella* spp. (right). Bar plots show the LIPA score associated with each tip of the

phylogeny, with higher values representing a stronger phylogenetic signal. Red colour indicates

statistically significant LIPA scores (*i.e.* phylogenetic signal, Supplementary Table 3).

52   Supplementary Figure 5. Model parametrization.
53

(a) Bacterial density of strain Kpn18 as a function of time obtained from the optical density of wild-type

(red) and transconjugant (black) strains growing in isolation. (b) Traces of chains for parameters

V_{max}/K_m (above) and ρ (below) obtained by fitting a simple Monod model to growth curve data using a

Metropolis-Hastings Markov chain Monte Carlo method (MCMC). (c) 2-dimensional posterior

distributions obtained for each strain (top: B_0, bottom: B_p). (d) Numerical solutions of the model using

parameters selected randomly from the posterior distribution and with initial conditions determined from

the experimental growth curves.

Supplementary Figure 6. *In silico* competition experiments.

Each box represents a theoretical pair-wise competition experiment between plasmid-free (open

circles) and plasmid-bearing cells (filled circles). The diameter of each circle is proportional to the

relative fraction of the population, a value estimated by numerically solving the model for $T = 24$ with

parameter values obtained from the posterior distribution of each strain. Horizontal axis represents the

specific affinity (V_{max}/K_m) and the vertical axis the cell's resource conversion rate (ρ).

    Supplementary Figure 7. Effect of conjugation and community complexity in plasmid population

70   dynamics in the absence of a DFE.

Numerical simulations of the population dynamics model performed over a range of conjugation rates

and number of strains in the community. In this case, all strains exhibit a reduction of fitness when

carrying the plasmid (a DFE with mean w= 0.985 and variance 0). The colour of each box in the grid

corresponds to the percentage of $5,000$ random communities that exhibited: a) plasmid extinction (total

plasmid frequency was below a threshold), b) plasmid-bearing and plasmid-free cells co-exist in the

population, and c) every cell in the population carries the plasmid at the end of the experiment. (d-g)

Example of relative abundances over time for a range of conjugation rates in a community composed of

1 (d,e) and 15 (f,g) strains, with segregation rate $\lambda = 1 \times 10^{-8}$ and conjugation rate $\gamma = 10^{-10}$ (d,f) or

$\gamma = 0$ (e,g). The left-hand column illustrates the growth kinetic parameters for each strain (empty

circles denote plasmid-free cells and filled circles plasmid-bearing cells, with diameters proportional to

their final relative abundances). Middle column shows the density of each subpopulation as a function

of time (dotted lines denote plasmid-free strains and solid lines subpopulations carrying the

plasmid). Right-hand column shows semilog plots with the total fraction of cells with and without

plasmids (solid and dotted lines, respectively). As plasmid-bearing is associated with a fitness cost,

then the plasmid is only maintained in the population at high conjugation rates.

   Supplementary Figure 8. Effect of conjugation and community complexity in plasmid population

dynamics in the presence of a DFE.

Numerical simulations of the population dynamics model performed over a range of conjugation rates

and number of strains in the community. This case corresponds to a wide DFE (mean $w = 0.985$ and

variance 0.007). The colour of each box in the grid corresponds to the percentage of 5,000 random

communities that exhibited: a) plasmid extinction (total plasmid frequency was below a threshold), b)

plasmid-bearing and plasmid-free cells co-exist in the population, and c) every cell in the population

carries the plasmid at the end of the experiment. d-g) Example of relative abundances over time for a

range of conjugation rates in a community composed of 1 (d,e) and 15 (f,g) strains, with segregation

rate $\lambda = 1 \times 10^{-8}$ (d,f) or $\gamma = 0$ (e,g). The left-hand column illustrates

the growth kinetic parameters for each strain (empty circles denote plasmid-free cells and filled circles

plasmid-bearing cells, with diameters proportional to their final relative abundances). Middle column

shows the density of each subpopulation as a function of time (dotted lines denote plasmid-free strains

and solid lines subpopulations carrying the plasmid). Right-hand column shows semilog plots with the

total fraction of cells with and without plasmids (solid and dotted lines, respectively). Note how a wide

DFE allows plasmids to persist, even at very low conjugation rates.

Supplementary Figure 9. Determination of different cells types in competition assays using flow

cytometry.

We used flow cytometry to differentiate between GFP-producing and -non-producing cells. (a) we used

forward versus side scatter (FSC vs SSC) gating to identify bacterial cells in the sample. (b-c) GFP-

producing (bright green) and -non-producing (dark green) cells were differentiated using the FITC-A

(fluorescein isothiocyanate) channel, allowing us to measure the proportion of each competitor in the

mix.

Supplementary Figure 10. Distribution of pBGC fitness effects.

Relative fitness (*w*) of pBGC-carrying clones compared to plasmid-free clones, obtained from

competition assays (red, *E. coli* and blue, *Klebsiella* spp.). Values below 1 indicate a reduction in *w* and

values above 1 indicate an increase in *w* due to pBGC acquisition. Bars represent the average of five

independent experiments and error bars represent the standard error of the mean. Note that the fitness

effects of pBGC did not correlate with those form pOXA-48 (Pearson's correlation, R= 0.11, t= 0.66, df=

39, *P*= 0.51).

Supplementary Figure 11. Correlation between relative fitness values calculated in competitions vs. *E.*

coli J53/pBGC or isogenic clones with pBGC.

Correlation between relative fitness values obtained from competitions assays using pBGC-carrying

isogenic isolates and pBGC-carrying *E.* coli J53 for ten different isolates. The blue line represents the

linear regression model and the grey shading represents 95% confidence intervals. Blue points

correspond to *Klebsiella* spp. isolates and red points to *E. coli* isolates. Labels indicate isolates names.

Pearson's correlation (R) and p-value are indicated.

Supplementary Table 4. Primers used in this study.

Primers		
Name	**Sequence 5'->3'**	**Use**
Oxa-48 Fw	TTGGTGGCATCGATTATCGG	Amplification of *bla*$_{TEM1}$ *gene*
Oxa-48 Rv	GAGCACTTCTTTTGTGATGGC	
IncL Fw	CGGAACCGACATGTGCCTACT	Amplification of *repC* gene
IncL Rv	GAACTCCGGCGAAAGACCTTC	
pBGC Fw	CGTTGATCGGCACGTAAG	Amplification of pBGC backbone for Gibson cloning (see Supplementary Figure 2)
pBGC Rv	GCTGTCTAGACTATTTGTATAGTTCATCCATGC	
GFP-Term Fw	atacaaatagtctagacagcGGGAATCCTGCTCTGCGAG[1]	
GFP-Term Rv	ctcttacgtgccgatcaacgGGGTTATTGTCTCATGAGCGG[1]	
pBGC_Seq1_Fw	AGTTAAAAGGTATTGATTTTAA	pBGC sequencing (3503-3524)[2]
pBGC_Seq1_Rv	GCCACATCTTGCGAATA	pBGC sequencing (464-480)[2]
pBGC_Seq2_Fw	ATAAGATCACTACCGGGC	pBGC sequencing (44-61)[2]
pBGC_Seq2_Rv	ACCCGACAGGACTATAAAGATA	pBGC sequencing (1285-1306)[2]
pBGC_Seq3_Fw	GAGGTAACTGGCTTGGAGG	pBGC sequencing (931-949)[2]
pBGC_Seq3_Rv	GTCGCGTCTGTCACATCT	pBGC sequencing (2124-2141)[2]
pBGC_Seq4_Fw	GTTTCCCGACTGGAAAGC	pBGC sequencing (1703-1720)[2]
pBGC_Seq4_Rv	CTTTGGTCCCGCTTTGTTAC	pBGC sequencing (2894-2913)[2]
pBGC_Seq5_Fw	GTCGGTCGATAAAAAAATCGAG	pBGC sequencing (2650-2671)[2]
pBGC_Seq5_Rv	ATGTGGTCTCTCTTTTCGTTGG	pBGC sequencing (3764-2671)[2]

134 1. Lower case nucleotides correspond with the added cohesive ends.

135 2. Numbers correspond with primers binding sites in pBGC (according to sequence available with

136 GenBank Accession Number MT702881).

137   Supplementary Table 5. Growth kinetic parameters of each strain obtained using the MCMC algorithm.

Strain	Wild-type		Transconjugant	
	V_{max}/K	ρ	V_{max}/K	ρ
Ec01	4.895×10^{-10}	9.844×10^{8}	5.283×10^{-10}	8.776×10^{8}
Ec02	7.603×10^{-10}	5.504×10^{8}	6.648×10^{-10}	6.786×10^{8}
Ec03	7.597×10^{-10}	8.900×10^{8}	6.538×10^{-10}	8.151×10^{8}
Ec04	8.373×10^{-10}	7.764×10^{8}	6.576×10^{-10}	7.702×10^{8}
Ec05	6.299×10^{-10}	7.727×10^{8}	7.831×10^{-10}	6.416×10^{8}
Ec06	6.852×10^{-10}	9.518×10^{8}	6.839×10^{-10}	6.567×10^{8}
Ec07	3.786×10^{-10}	1.040×10^{9}	4.154×10^{-10}	1.013×10^{9}
Ec08	5.304×10^{-10}	1.050×10^{9}	5.053×10^{-10}	9.900×10^{8}
Ec09	6.610×10^{-10}	9.195×10^{8}	7.311×10^{-10}	8.694×10^{8}
Ec10	5.660×10^{-10}	1.071×10^{9}	7.368×10^{-10}	8.472×10^{8}
Ec11	7.248×10^{-10}	5.064×10^{8}	7.248×10^{-10}	6.483×10^{8}
Ec12	5.200×10^{-10}	8.042×10^{8}	4.752×10^{-10}	8.016×10^{8}
Ec13	6.831×10^{-10}	8.528×10^{8}	7.711×10^{-10}	7.932×10^{8}
Ec14	6.515×10^{-10}	6.589×10^{8}	8.356×10^{-10}	7.500×10^{8}
Ec15	6.322×10^{-10}	9.106×10^{8}	6.998×10^{-10}	8.528×10^{8}
Ec16	4.908×10^{-10}	1.122×10^{9}	5.550×10^{-10}	1.040×10^{9}
Ec17	6.089×10^{-10}	8.298×10^{8}	6.366×10^{-10}	8.647×10^{8}
Ec18	6.655×10^{-10}	8.963×10^{8}	6.939×10^{-10}	6.250×10^{8}
Ec19	6.080×10^{-10}	9.812×10^{8}	6.200×10^{-10}	9.424×10^{8}
Ec20	5.033×10^{-10}	1.024×10^{9}	5.296×10^{-10}	9.133×10^{8}
Ec21	5.546×10^{-10}	1.021×10^{9}	4.897×10^{-10}	8.386×10^{8}
Ec22	6.148×10^{-10}	9.736×10^{8}	6.619×10^{-10}	9.886×10^{8}
Ec23	5.443×10^{-10}	9.635×10^{8}	5.186×10^{-10}	9.094×10^{8}
Ec24	6.010×10^{-10}	1.008×10^{9}	4.808×10^{-10}	9.768×10^{8}
Ec25	6.784×10^{-10}	9.237×10^{8}	5.465×10^{-10}	1.004×10^{9}
Kpn01	4.841×10^{-10}	8.746×10^{8}	3.652×10^{-10}	8.673×10^{8}
Kpn02	6.638×10^{-10}	7.905×10^{8}	6.352×10^{-10}	7.952×10^{8}
Kpn03	4.802×10^{-10}	9.639×10^{8}	3.686×10^{-10}	9.975×10^{8}
Kpn04	4.844×10^{-10}	9.288×10^{8}	4.303×10^{-10}	8.936×10^{8}
Kpn05	5.151×10^{-10}	9.303×10^{8}	5.155×10^{-10}	8.513×10^{8}
Kpn06	5.240×10^{-10}	8.952×10^{8}	5.664×10^{-10}	7.742×10^{8}
Kpn07	4.498×10^{-10}	8.682×10^{8}	3.785×10^{-10}	9.458×10^{8}
Kpn08	7.163×10^{-10}	8.211×10^{8}	4.334×10^{-10}	7.188×10^{8}
Kpn09	6.207×10^{-10}	8.064×10^{8}	7.302×10^{-10}	7.772×10^{8}
Kpn10	4.833×10^{-10}	7.990×10^{8}	4.467×10^{-10}	7.967×10^{8}
Kpn11	5.380×10^{-10}	9.710×10^{8}	5.620×10^{-10}	8.490×10^{8}
Kpn12	5.620×10^{-10}	8.490×10^{8}	5.620×10^{-10}	8.490×10^{8}

Kpn13	7.086×10^{-10}	9.195×10^8	7.364×10^{-10}	7.206×10^8
Kpn14	4.162×10^{-10}	8.994×10^8	5.828×10^{-10}	7.306×10^8
Kpn15	6.243×10^{-10}	8.854×10^8	7.489×10^{-10}	8.917×10^8
Kpn16	5.386×10^{-10}	8.523×10^8	4.359×10^{-10}	8.465×10^8
Kpn17	4.144×10^{-10}	1.037×10^9	4.002×10^{-10}	9.786×10^8
Kpn18	6.906×10^{-10}	7.173×10^8	4.977×10^{-10}	9.954×10^8
Kpn19	5.578×10^{-10}	8.319×10^8	6.083×10^{-10}	7.739×10^8
Kpn20	4.857×10^{-10}	1.022×10^9	4.287×10^{-10}	1.046×10^9
Kpn21	4.674×10^{-10}	1.077×10^9	5.487×10^{-10}	9.948×10^8
Kpn22	4.708×10^{-10}	1.006×10^9	5.224×10^{-10}	1.006×10^9
Kpn23	7.897×10^{-10}	6.709×10^8	7.201×10^{-10}	6.328×10^8
Kpn24	5.736×10^{-10}	7.568×10^8	8.670×10^{-10}	6.601×10^8
Kpn25	4.873×10^{-10}	9.630×10^8	4.255×10^{-10}	9.896×10^8