Chromosome painting in cultivated banana and their wild relatives (*Musa* spp.) reveals differences in chromosome structure

Šimoníková D[1], Němečková A[1], Čížková J[1], Brown A[2], Swennen R[.2,3,4], Doležel J[1], Hřibová E[1]*

[1]Institute of Experimental Botany of the Czech Academy of Sciences, Centre of the Region Hana for Biotechnological and Agricultural Research, Olomouc, Czech Republic

[2]International Institute of Tropical Agriculture, Banana Breeding, Arusha, Tanzania

[3]Bioversity International, Banana Genetic Resources, Heverlee, Belgium

[4]Division of Crop Biotechnics, Laboratory of Tropical Crop Improvement, Katholieke Universiteit Leuven, Leuven, Belgium

*Corresponding author:

Eva Hřibová

Institute of Experimental Botany

Centre of the Region Hana for Biotechnological and Agricultural Research

Šlechtitelů 31

CZ-77900 Olomouc

Czech Republic

E-mail: hribova@ueb.cas.cz

Abstract

Edible banana cultivars are diploid, triploid or tetraploid hybrids which originated by natural cross hybridization between subspecies of diploid *Musa acuminata*, or between *M. acuminata* and diploid *M. balbisiana*. Participation of two wild diploid species *M. schizocarpa* and *M. textilis* was also indicated by molecular studies. Fusion of gametes with structurally different chromosome sets may give rise to progenies with structural chromosome heterozygosity and reduced fertility due to aberrant chromosome pairing and unbalanced chromosome segregation. Only a few translocations have been classified on the genomic level so far and a comprehensive molecular cytogenetic characterization of cultivars and species of the family *Musaceae* is still lacking. FISH with chromosome-arm specific oligo painting probes was used for comparative karyotype analysis in a set of wild *Musa* species and edible banana clones. The results revealed large differences in chromosome structure discriminating individual accessions and permitted identification of putative progenitors of cultivated clones.

Keywords

Introduction

Banana represents one of the major staple foods and is one of the most important cash crops with the estimated value of $25 billion for the banana industry. An annual global production of bananas reached 114 million tons in 2017 (FAOSTAT, 2017) with about 26 million tons exported in 2019 (International Trade Statistics). Two types of bananas are known - sweet bananas, serving as a food supplement, and cooking bananas, which are characteristic by starchier fruits (*Price, 1995*). Edible banana cultivars are vegetatively propagated diploid, triploid and tetraploid hybrids which originated after natural cross hybridization between wild diploids *Musa acuminata* (2n=2x=22, AA genome) and *M. balbisiana* (2n=2x=22, BB genome) and their hybrid progenies. To some extent, also other *Musa* species such as *M. schizocarpa* (2n=22=22, SS genome) and *M. textilis* (2n=2x=20, TT genome) contributed to the origin of some edible banana clones (*Carreel et al., 1994, Čížková et al., 2013, Němečková et al., 2018*).

Based on morphology and geographical distribution, *M. acuminata* has been divided into nine subspecies (*banksii, burmannica, burmannicoides, errans, malaccensis, microcarpa, siamea, truncata* and *zebrina*) and three varieties (chinensis, sumatrana, tomentosa) (*Perrier et al., 2011, Martin et al., 2017, WCSP, 2018*). It has been estimated that least four subspecies of *M. acuminata* contributed to the origin of cultivated bananas (*Perrier et al., 2011, Rouard et al., 2018*). Out of them, *M. acuminata* ssp. *banksii*, with the original center of diversity in New Guinea played a major role in this process (*Sharrock, 1990, Perrier et al., 2009, Němečková et al., 2018*). Other subspecies were *M. acuminata* ssp. *burmannica* with the center of diversity in Myanmar (*Cheesman, 1948*), ssp. *malaccensis* with the origin in Malay peninsula (*De Langhe et al., 2009, Perrier et al., 2011*) and ssp. *zebrina* which originated in Indonesia (*Rouard et al., 2018*).

It is believed that human migration together with a different geography of the present archipelago in South-East Asia during glacial period, when drop of sea level resulted in interconnection of current islands in South-East Asia into one land mass (*Sand, 1989, Denham, 2004, Denham, 2010, Kagy et al., 2016*), brought different *M. acuminata* subspecies to a close vicinity, enabling cross hybridization and giving rise to diploid intra-specific hybrids that were subjected to human selection and propagation (*Perrier et al., 2011, Martin et al., 2017*). Fusion of unreduced gametes produced by diploid edible and partially sterile cultivars with normal haploid gametes from fertile diploid (*Simmonds, 1962, Carreel et al., 1994, Raboin et al., 2005*) would give rise to triploids. The number one export dessert banana type Cavendish as well as

other important dessert bananas, such as Gros Michel and Pome types, originated according to this scenario by hybridization of a diploid representative of sub-group 'Mchare' (originally named 'Mlali') (AA genome; *zebrina / microcarpa* and *banksii* ascendance) which served as a donor of an unreduced diploid gamete with haploid gamete of 'Khai' (*malaccensis* ascendance) (*Perrier et al., 2009, Perrier et al., 2019, Martin et al., 2020*).

Another group of edible triploid bananas, clones with AAB (so called plantains) or ABB constitution, cover nearly 40 % of global banana production, whereas plantains stand for 18 % of total banana production (*Baurens et al., 2019*). These interspecific triploid cultivars originated after fusion of an unreduced gamete from interspecific AB hybrid with haploid gamete from diploid *M. acuminata* ssp. *banksii* or *M. balbisiana* (*Perrier et al., 2011, Baurens et al., 2019*). Their evolution most probably involved several backcrosses (*De Langhe et al., 2010*). Two important AAB subgroups of starchy bananas evolved in two centers of diversity, the African Plantains and the Pacific (Maia Maoli / Popoulu) Plantains (*De Langhe et al., 2009*).

East African Highland bananas (EAHB) represent an important starchy type of banana for over 80 million people living in the Great Lakes region of East Africa, which is considered as a secondary center of banana diversity (*Cooper et al., 2001, Tugume et al., 2003*). EAHBs are a sub-group of triploid bananas with AAA constitution, which arose from hybridization between diploid *M. acuminata* ssp. *banksii* and *M. acuminata* ssp. *zebrina*. However, also *M. schizocarpa* contributed to the formation of these hybrids (*Němečková et al., 2018*). Interestingly, different EAHB varieties have relatively low genetic diversity, on the contrary to morphological variation which occurred most probably due to the accumulation of somatic mutations and epigenetic changes (*Perrier et al., 2009, Perrier et al., 2011, Kitavi et al., 2016, Christelová et al., 2017, Němečková et al., 2018*).

Cultivated clones originating from inter-subspecific and inter-specific hybridization and with a contribution of unreduced gametes in case of triploid clones have reduced or zero production of fertile gametes. This is a consequence of aberrant chromosome pairing during meiosis due to structural chromosome heterozygosity and/or odd ploidy levels. Reduced fertility greatly hampers the efforts to breed improved cultivars (*Burke et Arnold, 2001, Martin et al., 2017, Baurens et al., 2019, Batte et al., 2019*), which are needed to satisfy the increasing demand for dessert and starchy bananas under the conditions of climate change and increasing pressure of pests and diseases (*Christelová et al., 2017*).

Traditional breeding strategies for triploid banana cultivars involve development of tetraploids (4x) from 3x × 2x crosses, followed by production of secondary triploid hybrids (3x) from 4x × 2x crosses (*Bakry et Horry, 1992, Tomepke et al., 2004, Ortiz ,2013, Nyine et al., 2017*). Similarly, diploids play important role also in breeding strategy of diploid cultivars, 4x × 2x crosses are used to create improved cultivars (*Ortiz, 2013*). It is thus necessary to identify cultivars that produce seeds under specific condition, followed by breeding for target traits and re-establishing seed-sterile end product. Low seed yield after pollination (e.g. 4 seeds per Matooke (genome AAA) bunch and only 1.6 seeds per Mchare (genome AA) bunch followed by embryo rescue, with very low germination rate (~ 2%) illustrates the serious bottleneck for the breeding processes (*Brown and Swennen, unpublished*).

Since banana breeding programs use diploids as the principle vehicle for introducing genetic variability (*e.g. Amorim et al., 2011, Amorim et al., 2013, Tenkouano et al., 2003*), the knowledge of their genome structure at chromosomal level is critical to reveal possible causes of reduced fertility, and presence of non-recombining haplotype blocks, provide data to identify parents of cultivated clones, which originated spontaneously without a direct human intervention, and to select parents for improvement programs. In order to provide insights into the genome structure at chromosome level, we employed FISH with chromosome-arm specific oligo painting probes in a set of wild *Musa* species and edible banana clones potentially useful in banana improvement. Chromosome painting in twenty representatives of the Eumusa section of genus *Musa*, which included subspecies of *M. acuminata*, *M. balbisiana* and their inter-subspecific and inter-specific hybrids, revealed chromosomal rearrangements discriminating subspecies of *M. acuminata* and structural chromosome heterozygosity of cultivated clones. Identification of chromosome translocations pointed to particular *Musa* subspecies as putative parents of cultivated clones and provided an independent support for hypotheses on their origin.

Results

Karyotype of *M. acuminata* and *M. balbisiana*

The first part of the study focused on comparative karyotype analysis in six subspecies of *M. acuminata* and in *M. balbisiana*. Oligo painting FISH in structurally homozygous *M. acuminata* ssp. *banksii* 'Banksii' and *M. acuminata* ssp. *microcarpa* 'Borneo' did not reveal detectable chromosome translocations (*Fig. 1A*) as compared to the reference banana genome of *M. acuminata* ssp. *malaccensis* 'DH Pahang' (*Šimoníková et al., 2019*). Thus, the three

subspecies share the same overall organization of their chromosome sets. On the other hand, different chromosome translocations, which were found to be subspecies-specific were identified in the remaining four subspecies of *M. acuminata* (ssp. *zebrina*, ssp. *burmannica*, ssp. *burmannicoides*, ssp. *siamea).*

In *M. acuminata* ssp. *zebrina*, a reciprocal translocation between the short arm of chromosome 3 and the long arm of chromosome 8 was identified (*Figs. 1B, 2A*). Three phylogenetically closely related subspecies *M. acuminata* ssp. *burmannica*, *M. acuminata* ssp. *burmannicoides* and *M. acuminata* ssp. *siamea* shared two translocations (*Figs. 1C, 3A, 3B, 3E, 3F; Suppl. Figs. S1A, S1B*). The first of these involved a transfer of a segment of the long arm of chromosome 8 to the long arm of chromosome 2; the second was a reciprocal translocation involving a large segment of the short arm of chromosome 9 and the long arm of chromosome 1. In addition to the translocations shared by representatives of the three subspecies, chromosome painting revealed additional subspecies-specific translocations.

M. acuminata ssp. *banksii* 'Banksii' ITC0806 and ssp. *microcarpa* 'Borneo' ITC0253

B

M. acuminata ssp. *zebrina* 'Maia Oa' ITC0728

C

M. acuminata ssp. *burmannicoides* 'Calcutta 4' ITC0249

Figure 1: Idiograms of four structurally homozygous diploid (2n=2x=22) subspecies of *M. acuminata*: (A) *M. acuminata* ssp. *banksii* 'Banksii' and ssp. *microcarpa* 'Borneo'; (B) *M. acuminata* ssp. *zebrina* 'Maia Oa'; (C) *M.*

acuminata ssp. *burmannicoides* 'Calcutta 4'. Chromosomes are oriented with their short arms of at the top. Chromosome paints were not used for short arms of chromosomes 1, 2 and 10.

In *M. acuminata* ssp. *siamea* 'Pa Rayong' translocation of a small segment of the long arm of chromosome 3 to the short arm of chromosome 4 was detected. Importantly, this translocation was visible only on one of the homologs, indicating structural chromosome heterozygosity and a hybrid origin (*Fig. 3C; Suppl. Fig. S1A*). Subspecies-specific translocations involving only one of the homologs were also found in *M. acuminata* ssp. *burmannica* 'Tavoy'. They included a reciprocal translocation between chromosomes 3 and 8, which was also detected in *M. acuminata* ssp. *zebrina,* and a Robertsonian translocation between chromosomes 7 and 8, which gave rise to a chromosome comprising long arms of chromosomes 7 and 8 and a chromosome made of short arms of chromosomes 7 and 8 (*Figs. 2B, 3D; Suppl. Fig. S1B*).

In *M. balbisiana* 'Pisang Klutuk Wulung' translocation of a small segment of the long arm of chromosome 3 to the long arm of chromosome 1 was observed (*Suppl. Fig. S1C*), similar to the translocation identified in our previous work (*Šimoníková et al., 2019*) in *M. balbisiana* 'Tani'. In agreement with a low level of genetic diversity of *M. balbisiana* (*De Langhe et al., 2015, Kagy et al., 2016*), no detectable differences in karyotypes were found between both accessions of the species.

Karyotype structure of edible clones which originated as intra-specific hybrids

Chromosome painting in diploid cooking banana cultivars belonging to the Mchare group with AA genome, confirmed their hybrid origin. All three accessions analyzed in this work comprised two reciprocal translocations, with each of them observed only in one chromosome set. The first translocation involved short segments of long arms of chromosomes 4 and 1 (*Suppl. Fig. S1E*), while the second one involved short arms of chromosomes 3 and long arm of chromosome 8. Both translocations were observed in heterozygous state in all three analyzed Mchare banana representatives (*Fig. 2D; Suppl. Fig. S1E*). Reciprocal translocation involving short segments of long arms of chromosomes 4 and 1 was also identified in diploid cultivar 'Pisang Lilin' with AA genome (*Fig. 4C; Suppl. Fig. S1D*). This clone is used in breeding programs as a donor of useful agronomical characteristics, such as female fertility or resistance to yellow Sigatoka (*do Amaral et al., 2015*). Also this accession was heterozygous for the translocation, indicating a hybrid origin (*Suppl. Fig. S1D*).

Two representatives of triploid dessert banana cultivars 'Cavendish' and 'Gros Michel' with AAA genome constitution displayed identical chromosome structure as assessed by chromosome painting. One of the three chromosome sets was characterized by reciprocal translocation between short segments of the long arms of chromosomes 4 and 1 (*Fig. 4D; Suppl. Fig. S1F*). It is worth mentioning that the same translocation was identified in Mchare cultivars and in 'Pisang Lilin' (*Fig. 4C; Suppl. Figs. S1D, S1E*). Another chromosome set of 'Cavendish' (*Fig. 2F*) and 'Gros Michel' (*Fig. 2E*) contained a reciprocal translocation between the short arm of chromosome 3 and the long arm of chromosome 8, which was also identified in *M. acuminata* ssp. *zebrina* (*Figs. 1B, 2A*). In addition to the two translocations, a translocation of the short arm of chromosome 7 to the long arm of chromosome 1 was observed in both accessions of desert banana. The translocation resulted in formation of a small telocentric chromosome consisting of only the long arm of chromosome 7 (*Fig. 4E; Suppl. Fig. S1F*).

East African Highland bananas (EAHB) represent an important group of triploid cultivars with genome constitution AAA. We have analyzed two accessions of these cooking bananas. Both cultivars ('Imbogo' and 'Kagera') contained a reciprocal translocation between the short arm of chromosome 3 and the long arm of chromosome 8, which was also observed in *M. acuminata* ssp. *zebrina*. This translocation was identified only in two out of the three chromosome sets (*Fig. 2C; Suppl. Figs. S1G, S1H*). Karyotype analysis revealed that cultivar 'Imbogo' lacked one chromosome and was aneuploid (2n = 32). Chromosome painting facilitated identification of the missing chromosome and suggested the origin of the aneuploid karyotype (*Suppl. Figs. S1G, S2*), which involved a Robertsonian translocation between chromosomes 7 and 1, giving rise to a recombined chromosome containing long arms of chromosomes 7 and 1. Our observation suggests that short arms of the two chromosomes were lost (*Suppl. Figs. S1G, S2*). The loss of a putative chromosome comprising two short arms is a common consequence of the Robertsonian translocation (*Robertson, 1916*).

Karyotype structure of inter-specific banana clones

Plantains are an important group of starchy type of bananas with AAB genome constitution and originated as hybrids between *M. acuminata* (A genome) and *M. balbisiana* (B genome). Chromosome painting in three cultivars representing two plantain morphotypes - Horn (cultivar '3 Hands Planty') and French type (cultivars 'Obino l'Ewai' and 'Amou') confirmed the presence of B-genome specific translocation in one chromosome set in all three accessions, i.e. the translocation of a small segment of the long arm of chromosome 3 to the

long arm of chromosome 1 (*Fig. 4B; Suppl. Figs. S1I, S1J*). No other translocation was found in these cultivars. However, the clone 'Amou' was found to be aneuploid (2n = 32) as it missed one copy of chromosome 2 (*Fig. 4F; Suppl. Fig. S1J*).

In agreement with their predicted ABB genome constitution, B genome-specific chromosome translocation was also observed in two chromosome sets in triploid cultivars 'Pelipita' and 'Saba sa Hapon' (*Fig. 4A, Suppl. Fig. S1K*). No other translocation was found in these cultivars.

Figure 2: Examples of oligo painting FISH on mitotic metaphase plates of six *Musa* accessions: (A) 'Maia Oa' (2n=2x=22, AA genome), probes for long arm of chromosome 3 and short arm of chromosome 8 were labeled in red and green, respectively; (B) 'Tavoy' (2n=2x=22, AA genome), probes for long arm of chromosome 3 and short arm of chromosome 8 were labeled in red and green, respectively; (C) 'Imbogo' (2n=3x - 1= 32, AAA genome): probes for long arm of chromosome 3 and short arm of chromosome 8 were labeled in greed and red, respectively; (D) 'Huti (Shumba nyeelu)' (2n=2x=22, AA genome), probes for short arm of chromosome 3 and long arm of chromosome 8 were labeled in red and green, respectively; (E) 'Gros Michel' (2n=3x=33, AAA genome), probes for short arm of chromosome 3 and long arm of chromosome 8 were labeled in red and green, respectively; (F) 'Poyo' (2n=2x=33, AAA genome), probes for long arm of chromosome 3 and short arm of chromosome 8 were labeled in red and green, respectively. Chromosomes were counterstained with DAPI (light grey pseudocolor). Arrows point translocation chromosomes. Bars = 5 μm.

Figure 3: Examples of oligo painting FISH on mitotic metaphase plates of six *Musa* accessions: (A) 'Calcutta 4' (2n=2x=22, AA genome), probes for long arm of chromosome 1, long arm of chromosome 9 and short arm of chromosome 9 were labeled in green, red and purple, respectively; (B) 'Calcutta 4' (2n=2x=22, AA genome), probes for long arm of chromosome 2, long arm of chromosome 8 and short arm of chromosome 8 were labeled in green, red and purple, respectively; (C) 'Pa Rayong' (2n=2x=22, AA genome), probes for long arm of chromosome 3 and short arm of chromosome 4 were labeled in red and green, respectively; (D) 'Tavoy' (2n=2x=22, AA genome), probes for long arm of chromosome 7 and long arm of chromosome 8 were labeled in green and red, respectively; (E) 'Tavoy' (2n=2x=22, AA genome), probes for long arm of chromosome 2 and chromosome 8 were labeled in red and green, respectively; (F) 'Tavoy' (2n=2x=22, AA genome), probes for long arm of chromosome 1 and long arm of chromosome 9 were labeled in red and green, respectively. Chromosomes were counterstained with DAPI (light grey pseudocolor). Arrows point translocation chromosomes. Bars = 5 μm.

Figure 4. Examples of oligo painting FISH on mitotic metaphase plates of six structurally heterozygous *Musa* accessions: (A) 'Saba sa Hapon' (2n=3x=33, ABB genome), probes for long arm of chromosome 1, long arm of chromosome 2 and chromosome 3 were labeled in red, purple and green, respectively; (B) '3 Hands Planty' (2n=3x=33, AAB genome), probes for long arm of chromosome 1 and chromosome 3 were labeled in green and red, respectively; (C) 'Pisang Lilin' (2n=2x=22, AA genome), probes for long arm of chromosome 1 and chromosome 4 were labeled in red and green, respectively; (D) 'Gros Michel' (2n=3x=33, AAA genome), probes for long arm of chromosome 1 and long arm of chromosome 4 were labeled in red and green, respectively; (E) 'Gros Michel' (2n=3x=33, AAA genome), probes for long arm of chromosome 1, long arm of chromosome 2 and chromosome 7 were labeled in green, purple and red, respectively; (F) 'Amou' (2n=3x – 1=32, AAB): probes for long arm of chromosome 2 and chromosome 4 were labeled in green and red, respectively. Chromosomes were counterstained with DAPI (light grey pseudocolor). Arrows point translocation chromosomes. Bars = 5 µm.

Discussion

Until recently, cytogenetic analysis in plants was hindered by the lack of available DNA probes suitable for fluorescent painting of individual chromosomes (*Schubert et al., 2001, Jiang, 2019*). The only option was to use pools of BAC clones which were found useful in the plants species with small genomes (*e.g. Lysák et al., 2001, Idziak et al., 2014*). The development of oligo painting FISH (*Han et al., 2015*) changed this situation dramatically and it is now possible to label individual plant chromosomes and chromosomal regions in many species (*Qu et al., 2017, Braz et al., 2018, He et al., 2018, Machado et al., 2018, Albert et al., 2019, Bi et al., 2020*). Chromosome arm-specific oligo painting probes were recently developed also for banana (*Musa* spp.) by Šimoníková *et al*. (*2019*) who demonstrated the utility of this approach for anchoring DNA pseudomolecules of a reference genome sequence to individual chromosomes *in situ*, and for identification of chromosome translocations.

In this work we used oligo painting FISH for comparative karyotype analysis in a set of *Musa* accessions comprising wild species used in banana breeding programs and economically significant edible cultivars (*Suppl. Tab. S1*). These experiments revealed chromosomal translocations in subspecies of *M. acuminata* (A genome), their intra-specific hybrids as well as in *M. balbisiana* (B genome) and in inter-specific hybrid clones originating from cross hybridization between *M. acuminata* and *M. balbisiana* (*Figure 5*). A difference in chromosome structure among *M. acuminata* subspecies was suggested earlier by Shepherd *et al*. (*1999*) who identified seven translocation groups in *M. acuminata* based on chromosome pairing during meiosis. An independent confirmation of this classification was the observation

of segregation distortion during genetic mapping in inter-subspecific hybrids of *M. acuminata* (*Fauré et al., 1993, Hippolyte et al., 2010, Mbanjo et al., 2012, Noumbissie et al., 2016*).

Figure 5. Overview of common translocation events revealed by oligo painting FISH in *Musa*: Diversity tree, constructed using SSR markers according to Christelová *et al.* (*2017*), shows the relationships among *Musa*

species, subspecies and hybrid clones. Lineages of closely related accessions and groups of edible banana clones are highlighted in different colors. Individual chromosome structures (displayed as chromosome schemes) are depicted in rows, and their number correspond to the number of chromosomes bearing the rearrangement in the nuclear genome in somatic cell lines (2n).

Structural genome variation in diploid *M. acuminata*

In this work we analyzed one representative of each of six subspecies of *M. acuminata*. We cannot exclude differences in chromosome structure within individual subspecies. However, given the large genetic homogeneity of the six subspecies as clearly demonstrated by molecular markers (*Carreel et al., 2002, Perrier et al., 2009, Hřibová et al., 2011, Christelová et al., 2017, Němečková et al., 2018, Dupouy et al., 2019*), this does not seem probable. We observed a conserved genome structure in *M. acuminata* ssp. *banksii* and *M. acuminata* ssp. *microcarpa* which did not contain any translocation chromosome when compared to the reference genome of *M. acuminata* ssp. *malaccensis* 'DH Pahang' (*Martin et al., 2016*). The genome structure shared by the three subspecies was also observed in *M. schizocarpa* (*Šimoníková et al., 2019*) and corresponds to the standard translocation (ST) group as defined by Shepherd (*1999*).

Other chromosome translocation group defined by Shepherd (*1999*), the Northern Malayan group (NM), is characteristic for *M. acuminata* ssp. *malaccensis* and some AA cultivars, including 'Pisang Lilin'. Our results revealed a reciprocal translocation between chromosomes 1 and 4 in one chromosome set of 'Pisang Lilin', thus confirming Shepherd's characterization of this clone as heterozygous having ST x NM genome structure. Before the *Musa* genome sequence became available, Hippolyte *et al.* (*2010*) assumed a presence of a duplication of distal region of chromosome 1 on chromosome 4 in this clone based on comparison of high dense genetic maps. Taking the advantage of the availability of reference genome sequence and after resequencing genomes of a set of *Musa* species, Martin *et al.* (*2017*) described heterozygous reciprocal translocation between chromosomes 1 and 4, involving 3 Mb of long arm of chromosome arm 1 and 10 Mb segment of long arm of chromosome arm 4 in *M. acuminata* ssp. *malaccensis*. Further experiments indicated preferential transmission of the translocation to the progeny, and its frequent presence in triploid banana cultivars (*Martin et al., 2017*). Yet, it is not clear if the reciprocal translocation between chromosomes 1 and 4, which we observed in the heterozygous state only, originated in ssp. *malaccensis*, or if it was transmitted to genomes of some *malaccensis* accessions by ancient hybridization events

(*Martin et al., 2017*). Three phylogenetically closely related subspecies *M. acuminata* ssp. *burmannica*, *M. acuminata* ssp. *burmannicoides* and *M. acuminata* ssp. *siamea*, which have similar phenotype and geographic distribution (*Simmonds, 1962, Perrier et al., 2009*) share a translocation of a part of the long arm of chromosome 8 to the long arm of chromosome 2, and a reciprocal translocation between chromosomes 1 and 9. These translocations were identified recently by Dupouy *et al.* (*2019*) after mapping mate-paired Illumina sequence reads to the reference genome of 'DH Pahang'. The authors estimated the size of translocated region of chromosome 8 to chromosome 2 to be 7.2 Mb, while the size of the distal region of chromosome 2, which was found translocated to chromosome 8 in wild diploid clone 'Calcutta 4' (ssp. *burmannicoides*) was estimated to be 240 kb (*Dupouy et al., 2019*). The size of the translocated regions of chromosomes 1 and 9 was estimated to be 20.8 Mb and 11.6 Mb respectively (*Dupouy et al., 2019*). Using oligo painting, we did not detect the 240 kb distal region of chromosome 2 translocated to chromosome 8, and this may reflect the limitation in the sensitivity of whole chromosome arm oligo painting.

The shared translocations in all three subspecies of *M. acuminata* (*burmannicoides*, *burmannica* and *siamea*) support their close phylogenetic relationship as proposed by Shepherd (*1999*) and later verified by molecular studies (*Carreel et al., 2002, Perrier et al., 2009, Hřibová et al., 2011, Christelová et al., 2017, Němečková et al., 2018*). Dupouy *et al.* (*2019*) coined the idea of a genetically uniform *burmannica* group. However, our data indicate a more complicated evolution of the three genotypes recognized as representatives of different *acuminata* subspecies. First, the characteristic translocations between chromosomes 2 and 8, and 1 and 9 were detected only on one chromosome set in *burmannica* 'Tavoy', as compared to *M. acuminata* 'Calcutta 4' (ssp. *burmannicoides*) and 'Pa Rayong' (ssp. *siamea*). Second, we observed two subspecies-specific translocations in ssp. *burmannica* only in one chromosome set, and we detected additional subspecies-specific translocation in *M. acuminata* ssp. *siamea* only in one chromosome set, indicating its hybrid origin. Based on these results we hypothesize that ssp. *burmannicoides* could be a progenitor of the clones characterized by structural chromosome heterozygosity. Divergence in genome structure between the three subspecies (*burmannica, burmannicoides* and *siamea*) was demonstrated also by Shepherd (*1999*), who classified some *burmannica* and *siamea* accessions as Northern 2 translocation group of *Musa*, differing from the Northern 1 group (*burmannicoides* and other *siamea* accessions) by one additional translocation.

We observed subspecies-specific translocations also in *M. acuminata* ssp. *zebrina* 'Maia Oa'. In this case, chromosome painting revealed a Robertsonian translocation between chromosomes 3 and 8. Interestingly, Dupouy *et al.* (*2019*) failed to detect this translocation after sequencing mate-pair libraries using Illumina technology. The discrepancy may point to the limitation of the sequencing approach to identify translocations arising by a breakage of (peri)centromeric regions. As these regions comprise mainly DNA repeats, they may not be assembled properly in a reference genome, thus preventing their identification by sequencing. In fact, this problem may also be encountered if subspecies-specific genome regions are absent in the reference genome sequence. We revealed the *zebrina*-type translocation (a reciprocal translocation between chromosomes 3 and 8) also in all three analyzed cultivars of diploid Mchare banana. The presence of a translocation between long arm of chromosome 1 and long arm of chromosome 4 on one chromosome set of Mchare indicates a hybrid origin of Mchare, with ssp. *zebrina* being one of the progenitors of this banana group. This agrees with the results obtained by genotyping using molecular markers (*Christelová et al., 2017*). Most recently, complex hybridization scheme of Mchare bananas was supported also by application of transcriptomic data for identification of specific SNPs in 23 *Musa* species and edible cultivars (*Martin et al., 2020*).

Genome structure and origin of cultivated triploid *Musa* clones

Plantains are an important group of triploid starchy bananas with AAB genome constitution, which originated after hybridization between *M. acuminata* and *M. balbisiana*. As expected, chromosome painting in '3 Hands Planty', 'Amou' and 'Obino l'Ewai' cultivars revealed B-genome specific translocation of 8 Mb segment from the long arm of chromosome 3 to the long arm of chromosome 1. Unlike the B-genome chromosome set, the two A-genome chromosome sets of plantains lacked any detectable translocation. Genotyping using molecular markers revealed that the A genomes of the plantain group are related to *M. acuminata* ssp. *banksii* (*Horry, 1989, Lebot et al., 1993, Carreel et al., 2002, Kagy et al., 2016*). This is in line with the absence of chromosome translocations we observed in *M. acuminata* ssp. *banksii*.

Triploid cultivar 'Pisang Awak', a representative of the ABB group, is believed to contain one A genome chromosome set closely related to *M. acuminata* ssp. *malaccensis* (*Perrier et al., 2009*). However, after SSR genotyping Christelová *et al.* (*2017*) found, that some ABB clones from Saba and Bluggoe-Monthan groups clustered together with the representatives of Pacific banana Maia Maoli / Popoulu (AAB). These results point to *M. acuminata* ssp. *banksii* as their most probable progenitor. Unfortunately, in this case chromosome painting did not

bring useful hints on the nature of the A subgenomes in these inter-specific hybrids, as *M. acuminata* ssp. *banksii* and ssp. *malaccensis* representatives do not differ in the presence of specific chromosome translocation.

The presence of B genome-specific translocation of a small region of long arm of chromosome 3 to the long arm of chromosome 1 (*Šimoníková et al., 2019*), observed in all *M. balbisiana* accessions and their inter-specific hybrids with *M. acuminata*, seems to be a useful cytogenetic landmark of the presence of B sub-genome. In our study, the number of chromosome sets containing B genome specific translocation agreed with the predicted genomic constitution (AAB or ABB) of the hybrids. Clearly, one B genome-specific landmark is not sufficient for analysis of complete genome structure of inter-specific hybrids at the cytogenetic level. Further work is needed to identify additional cytogenetic landmarks to uncover the complexities of genome evolution after inter-specific hybridization in *Musa*. Recently, Baurens *et al*. (*2019*) analyzed genome composition of banana inter-specific hybrid clones using whole genome sequencing strategies followed by bioinformatic analysis based on A- and B-genome specific SNPs calling and they also detected the B genome-specific translocation in inter-specific hybrids.

Chromosome painting confirmed a small genetic difference between triploid clones 'Gros Michel' and 'Cavendish' (AAA genomes) as previously determined by various molecular studies (*Raboin et al., 2005, Christelová et al., 2017*). Both clones share the same reciprocal translocation between long arms of chromosomes 1 and 4 in one chromosome set. Interestingly, also diploid Mchare cultivars contain the same translocation in one chromosome set. The presence of the translocation was identified also by sequencing genomic DNA both in 'Cavendish' and 'Gros Michel', as well as in Mchare banana 'Akondro Mainty' (*Martin et al., 2017*). These observations confirm close genetic relationship between both groups of edible bananas as noted previously (*Raboin et al., 2005, Perrier et al., 2009*). According to Martin *et al*. (*2017, 2020*), 2n gamete donor, which contributed to the origin of dessert banana clones with AAA genomes, including 'Cavendish' and 'Gros Michel', belongs to the Mchare (Mlali) sub-group. The genome of this ancient sub-group, which probably originated somewhere around Java, Borneo and New Guinea, but today is only found in East Africa, is based on *zebrina / microcarpa* and *banksii* subspecies (*Perrier et al., 2009*).

The third chromosome set in triploid 'Cavendish' and 'Gros Michel' contains a reciprocal translocation between chromosomes 3 and 8, which was detected by oligo painting FISH in the

diploid *M. acuminata* ssp. *zebrina*. Our observations indicate that heterozygous representatives of *M. acuminata* ssp. *malaccensis* and *M. acuminata* spp. *zebrina* contributed to the origin of 'Cavendish' and 'Gros Michel' as their ancestors as suggested earlier (*Perrier et al., 2009, Hippolyte et al., 2012, Christelová et al., 2017*). According to Perrier *et al.* (*2009*), *M. acuminata* ssp. *banksii* was one of the two progenitors of 'Cavendish' / 'Gros Michel' group of cultivars. However, using chromosome arm-specific oligo painting, we observed a translocation of the short arm of chromosome 7 to the long arm of chromosome 1, resulting in a small telocentric chromosome made only of the long arm of chromosome 7, which was, up to now, identified only in these cultivars. The translocation, which gave arise to the small telocentric chromosome, could be a result of processes accompanying the evolution of this group of triploid AAA cultivars. Alternatively, another wild diploid clone, possibly structurally heterozygous, was involved in the origin of 'Cavendish' / 'Gros Michel' bananas.

We observed reciprocal translocation between chromosomes 3 and 8, which is typical for *M. acuminata* ssp. *zebrina*, in the economically important group of triploid East African Highland bananas (EAHB) with AAA genome constitution. An important role of *M. acuminata* ssp. *zebrina* and *M. acuminata* ssp. *banksii,* as the most probable progenitors of EAHB, was suggested previously (*Carreel et al., 2002, Li et al., 2013, Kitavi et al., 2016, Christelová et al., 2017, Němečková et al., 2018, Martin et al., 2020*). Our results, which indicate that EAHB contained two chromosome sets from ssp. *zebrina* and one chromosome set from ssp. *banksii*, point to the most probable origin of EAHB. Hybridization between *M. acuminata* ssp. *zebrina* and ssp. *banksii* could give arise to intra-specific diploid hybrid with a reduced fertility. Triploid EAHB cultivars then could originate by backcross of the intra-specific hybrid (a donor of non-reduced gamete) with *M. acuminata* ssp. *zebrina*, or with another diploid, most probably a hybrid of *M. acuminata* ssp. *zebrina*. Moreover, phylogenetic analysis of Němečková *et al.* (*2018*) surprisingly revealed also possible contribution of *M. schizocarpa* to EAHB formation, thus indicating a more complicated origin. Further investigation is needed, and the availability of EAHB genome sequence in particular, to shed more light on the origin and evolution of these triploid clones.

The origin of aneuploidy

To date, aneuploidy in *Musa* has been identified by chromosome counting (*Sandoval et al., 1996, Shepherd and Da Silva, 1996, Bartoš et al., 2005, Čížková et al., 2013, Čížková et al., 2015, Němečková et al., 2018*). Although this approach is laborious and low throughput, it

cannot be replaced by flow cytometric estimation of nuclear DNA amounts because of the differences in genome size between *Musa* species, sub-species and their hybrids (*e.g. Čížková et al., 2013, Christelová et al., 2017, Němečková et al., 2018*). A more laborious approach to achieve high resolution flow cytometry as used by Roux *et al.* (*2003*) is too slow and laborious to be practical. Thus, traditional chromosome counting (*Sandoval et al., 1996, Shepherd and Da Silva, 1996, Bartoš et al., 2005, Čížková et al., 2013, Čížková et al., 2015, Němečková et al., 2018*) remains the most reliable approach. Obviously, it is not suitable to identify the chromosome(s) involved in aneuploidy and the origin of the aberrations.

Here we employed chromosome painting to shed light on the nature of aneuploids among the triploid *Musa* accessions. One of the aneuploid clones was identified in plantain 'Amou', in which one copy of chromosome 2 was lost. The origin of aneuploidy in clone 'Imbogo' (AAA genome), a representative of EAHB, involved structural chromosome changes involving breakage of chromosomes 1 and 7 in centromeric regions, followed by fusion of long arms of chromosomes 1 and 7 and subsequent loss of short arms of both chromosomes (*Suppl. Fig. S2*). It needs to be noted that these plants were obtained from the International Musa Germplasm Transit Centre (ITC, Leuven, Belgium), where the clones are stored *in vitro*. The loss of whole chromosome in plantain 'Amou' could occur during long-term culture.

To conclude, the application of oligo painting FISH improved the knowledge on genomes of cultivated banana and their wild relatives at chromosomal level. For the first time, a comparative molecular cytogenetic analysis of twenty representatives of the Eumusa section of genus *Musa*, including accessions commonly used in banana breeding, was performed using chromosome painting. However, as only a single accession from each of the six subspecies of *M. acuminata* was used in this study, our results will need to be confirmed by analyzing more representatives from each taxon. The identification of chromosome translocations pointed to particular *Musa* subspecies as putative parents of cultivated clones and provided an independent support for hypotheses on their origin. While we have unambiguously identified a range of translocations, a precise determination of breakpoint positions will have to be done using a long read DNA sequencing technology (*Sedlazeck et al., 2018, Hu et al., 2020, Soto et al., 2020*).

The discrepancies in genome structure of banana diploids observed in our study and published data then point to alternative scenarios on the origin of the important crop. The observation on structural chromosome heterozygosity confirmed the hybrid origin of cultivated banana and some of the wild diploid accessions which were described as individual subspecies,

and informs breeders on possible causes of reduced fertility. The knowledge on genome structure at chromosomal level and characterization structural chromosome heterozygosity will aid breeders in selecting parents for improvement programs.

Materials and Methods

Plant material and diversity tree construction

Representatives of twenty species and clones from the section Eumusa of genus *Musa* were obtained as *in vitro* rooted plants from the International *Musa* Transit Centre (ITC, Bioversity International, Leuven, Belgium). *In vitro* plants were transferred to soil and kept in a heated greenhouse. Table 1 lists the accessions used in this work. Genetic diversity analysis of banana accessions used in the study was performed using SSR data according to Christelová *et al.* (*2017*). Dendrogram was constructed based on the results of UPGMA analysis implemented in DARwin software v6.0.021 (*Perrier and Jacquemoud-Collet, 2006*) and visualized in FigTree v1.4.0 (http://tree.bio.ed.ac.uk/software/figtree/).

Table 1: List of *Musa* accessions analyzed in this work and their genomic constitution

Species	Subspecies / Subgroup	Accession name	ITC code[a]	Genomic constitution	Chromosome number (2n)
M. acuminata	*banksii*	Banksii	0806	AA	22
	microcarpa	Borneo	0253	AA	22
	zebrina	Maia Oa	0728	AA	22
	burmannica	Tavoy	0072	AA	22
	burmannicoides	Calcutta 4	0249	AA	22
	siamea	Pa Rayong	0672	AA	22
M. balbisiana	-	Pisang Klutuk Wulung	-	BB	22
Cultivars	unknown	Pisang Lilin	-	AA	22
	Mchare	Huti White		AA	22

	Mchare	Huti (Shumba nyeelu)	1452	AA	22
	Mchare	Ndyali	1552	AA	22
	Cavendish	Poyo	1482	AAA	33
	Gros Michel	Gros Michel	0484	AAA	33
	Mutika/Lujugira	Imbogo	0168	AAA	32*
	Mutika/Lujugira	Kagera	0141	AAA	33
	Plantain (Horn)	3 Hands Planty	1132	AAB	33
	Plantain (French)	Amou	0963	AAB	32*
	Plantain (French)	Obino l'Ewai	0109	AAB	33
	Pelipita	Pelipita	0472	ABB	33
	Saba	Saba sa Hapon	1777	ABB	33

[a]Code assigned by the International Transit Centre (ITC, Leuven, Belgium).

*Aneuploidy observed in our study.

Preparation of oligo painting probes and mitotic metaphase chromosome spreads

Chromosome-arm specific painting probes were prepared as described by Šimoníková *et al*. (*2019*). Briefly, sets of 20,000 oligomers (45-nt) covering individual chromosome arms were synthesized as immortal libraries by Arbor Biosciences (Ann Arbor, Michigan, USA) and then labeled directly by CY5 fluorochrome, or by digoxigenin or biotin according to Han *et al*. (*2015*). N.B.: In the reference genome assembly of *M. acuminata* DH Pahang (*D'Hont et al., 2012*), pseudomolecules 1, 6, 7 are oriented inversely to the traditional way karyotypes are presented, where the short arms are on the top (*Suppl. Fig. S3, see also Šimoníková et al., 2019*).

Actively growing root tips were collected and pre-treated in 0.05% (w/v) 8-hydroxyquinoline for three hours at room temperature, fixed in 3:1 ethanol:acetic acid fixative overnight at -20°C and stored in 70% ethanol at -20°C. After washing in 75-mM KCl and 7.5-mM EDTA (pH 4), root tip segments were digested in a mixture of 2% (w/v) cellulase and 2%

(w/v) pectinase in 75-mM KCl and 7.5-mM EDTA (pH 4) for 90 min at 30°C. The suspension of protoplasts thus obtained was filtered through a 150-µm nylon mesh, pelleted and washed in 70% ethanol. For further use, the protoplast suspension was stored in 70% ethanol at -20°C. Mitotic metaphase chromosome spreads were prepared by a dropping method from protoplast suspension according to Doležel *et al.* (*1998*), the slides were postfixed in 4% (v/v) formaldehyde solution in 2 × SSC solution, air dried and used for FISH.

Fluorescence *in situ* hybridization and image analysis

Fluorescence *in situ* hybridization and image analysis were performed according to Šimoníková *et al.* (*2019*). Hybridization mixture containing 50% (v/v) formamide, 10% (w/v) dextran sulfate in 2 × SSC and 10 ng/µl of labeled probes was added onto slide and denatured for 3 min at 80°C. Hybridization was carried out in a humid chamber overnight at 37°C. The sites of hybridization of digoxigenin- and biotin-labeled probes were detected using anti-digoxigenin-FITC (Roche Applied Science, Penzberg, Germany) and streptavidin-Cy3 (ThermoFisher Scientific/Invitrogen, Carlsbad, CA, USA), respectively. Chromosomes were counterstained with DAPI and mounted in Vectashield Antifade Mounting Medium (Vector Laboratories, Burlingame, CA, USA). The slides were examined with Axio Imager Z.2 Zeiss microscope (Zeiss, Oberkochen, Germany) equipped with Cool Cube 1 camera (Metasystems, Altlussheim, Germany) and appropriate optical filters. The capture of fluorescence signals, merging the layers and measurement of chromosome length were performed with ISIS software 5.4.7 (Metasystems). The final image adjustment and creation of idiograms were done in Adobe Photoshop CS5.

Acknowledgements

We thank Dr. Ines van den Houwe for providing the plant material. This work was supported by the Czech Science Foundation (award No. 19-20303S) and by the Ministry of Education, Youth and Sports of the Czech Republic (Program INTER-EXCELLENCE, INTER-TRANSFER, grant award LTT19, and ERDF project "Plants as a tool for sustainable global development" No. CZ.02.1.01/0.0/0.0/16_019/0000827). The computing was supported by the National Grid Infrastructure MetaCentrum (grant No. LM2010005 under the program Projects of Large Infrastructure for Research, Development, and Innovations). The authors thank all donors who supported this work also through their contributions to the CGIAR Fund

Author contributions

EH and JD conceived the project. DŠ, AN and JČ conducted the study and processed the data. AB and RS provided the banana materials. DŠ and EH wrote the manuscript. EH, DŠ, AB, RS and JD discussed the results and contributed to manuscript writing. All authors have read and approved the final manuscript.

Competing interests

The authors declare no conflicts of interest

Data statement

Data supporting the findings of this work are available within the paper and its Supporting Information files. All other data generated and analyzed during the current study are available from the corresponding author upon reasonable request.

Supporting information

Figure S1: Idiograms of *Musa* accessions. (A) *M. acuminata* ssp. *siamea* 'Pa Rayong' (genome AA); (B) *M. acuminata* ssp. *burmannica* 'Tavoy' (genome AA); (C) *M. balbisiana* 'Pisang Klutuk Wulung' (genome BB); (D) *M. acuminata* 'Pisang Lilin' (genome AA); (E) *M. acuminata* subgr. Mchare, clones 'Huti white', 'Huti (Shumba nyeelu)' and 'Ndyali' (genomes AA); (F) triploid clones 'Gros Michel' and 'Poyo' (genomes AAA); (G) East African Highland Banana (EAHB) clone 'Imbogo' (genome AAA); (H) EAHB clone 'Kagera' (genome AAA); (I) plantains '3 Hands Planty' and 'Obino l'Ewai' (genomes AAB); (J) aneuploid plantain clone 'Amou'; (K) triploid clones 'Pelipita' and 'Saba sa Hapon' (genomes ABB). Short arms of the chromosomes are on the top and the long arms on the bottom in all idiograms. For better orientation, translocated parts of the chromosomes contain extra labels, which include chromosome number and chromosome arm which was involved in the rearrangement.

Figure S2: Robertsonian translocation as a mechanism leading to structural chromosome change (formation of a translocation chromosome 7L.1L) and aneuploidy (loss of the short arms of chromosomes 1 and 7) in East African Highland Banana clone 'Imbogo'.

Figure S3: Anchoring of chromosomes to *M. acuminata* 'DH Pahang' pseudomolecules which were used to develop oligo painting probes. Note that in the reference genome assembly, pseudomolecules 1, 6, 7 are oriented inversely to the traditional way karyotypes are presented, where the short arms are on the top.

Table S1: Comparison of genome structure in selected *Musa* accessions compared to the *Musa* reference genome sequence (*M. acuminata* ssp. *malaccensis* 'DH Pahang').

References

Albert PS, Zhang T, Semrau K, Rouillard JM, Kao YH, Wang CJR, Danilova TV, Jiang JM, Birchler JA. 2019. Whole-chromosome paints in maize reveal rearrangements, nuclear domains, and chromosomal relationships. *Proceedings of the National Academy of Sciences of the United States of America* **116:** 1679–1685. DOI: https://doi.org/10.1073/pnas.1813957116

Amorim EP, Silva SO, Amorim VBO, Pillay M. 2011. Quality improvement of cultivated *Musa*. In: *Banana Breeding: Progress and Challenges* (Pillay M, Tenkouano A, eds), pp 252–280. New York: CRC Press. DOI: https://doi.org/10.1201/b10514-14

Amorim EP, dos Santos-Serejo JA, Amorim VBO, Ferreira CF, Silva SO. 2013. Banana breeding at embrapa cassava and fruits. *Acta Horticulturae* **968:** 171–176. DOI: https://doi.org/10.17660/ActaHortic.2013.986.18

Bakry F, Horry JP. 1992. Tetraploid hybrids from interploid 3x/2x crosses in cooking bananas. *Fruit*s **47**: 641–655.

Bartoš J, Alkhimova O, Doleželová M, De Langhe E, Doležel J. 2005. Nuclear genome size and genomic distribution of ribosomal DNA in *Musa* and *Ensete* (*Musaceae*): taxonomic implications. *Cytogenetic and Genome Research* **109:** 50–57. DOI: https://doi.org/10.1159/000082381

Batte M, Swennen R, Uwimana B, Akech V, Brown A, Tumuhimbise R, Hovmalm HP, Geleta M, Ortiz R. 2019. Crossbreeding East African Highland Bananas: Lessons learnt relevant to the botany of the crop after 21 years of genetic enhancement. *Frontiers in Plant Science* **10:** 81. DOI: https://doi.org/10.3389/fpls.2019.00081

Baurens FC, Martin G, Hervouet C, Salmon F, Yohomé D, Ricci S, Rouard M, Habas R, Lemainque A, Yahiaoui N, D'Hont A. 2019. Recombination and large structural variations shape interspecific edible bananas genomes. *Molecular Biology and Evolution* **36:** 97–111. DOI: https://doi.org/10.1093/molbev/msy199

Bi Y, Zhao Q, Yan W, Li M, Liu Y, Cheng C, Zhang L, Yu X, Li J, Qian C, Wu Y, Chen J, Lou Q. 2020. Flexible chromosome painting based on multiplex PCR of oligonucleotides and

its application for comparative chromosome analyses in *Cucumis*. *Plant Journal* **102:** 178–186. DOI: http://doi: 10.1111/tpj.14600

Braz GT, He L, Zhao H, Zhang T, Semrau K, Rouillard JM, Torres GA, Jiang J. 2018. Comparative oligo-FISH mapping: An efficient and powerful methodology to reveal karyotypic and chromosomal evolution. *Genetics* **208:** 513–523. DOI: http://doi: 10.1534/genetics.117.300344

Burke JM, Arnold ML. 2001. Genetics and the fitness of hybrids. *Annual Review of Genetics* **35:** 31–52. DOI: https://doi.org/10.1146/annurev.genet.35.102401.085719

Carreel F, Fauré S, Gonzalez de Leon D, Lagoda PJL, Perrier X, Bakry F, Tezenas du Montcel H, Lanaud C, Horry JP. 1994. Evaluation of the genetic diversity in diploid bananas (*Musa* sp.). *Genetics Selection Evolution* **26:** 125–136. DOI: https://doi.org/10.1051/gse:19940709

Carreel F, Gonzalez de Leon D, Lagoda P, Lanaud C, Jenny C, Horry J, Tezenas du Montcel H. 2002. Ascertaining maternal and paternal lineage within *Musa* by chloroplast and mitochondrial DNA RFLP analyses. *Genome* **45:** 679–692. DOI: https://doi.org/10.1139/G02-033

Cheesman EE. 1948. Classification of the bananas. *Kew Bulletin* **3:** 145–153. DOI: https://doi.org/10.2307/4119749

Christelová P, De Langhe E, Hřibová E, Čížková J, Sardos J, Hušáková M, Van den Houwe I, Sutanto A, Kepler AK, Swennen R, Roux N, Doležel J. 2017. Molecular and cytological characterization of the global *Musa* germplasm collection provides insights into the treasure of banana diversity. *Biodiversity and Conservation* **26:** 801–824. DOI: https://doi.org/10.1007/s10531-016-1273-9

Čížková J, Hřibová E, Christelová P, Van den Houwe I, Häkkinen M, Roux N, Swennen R, Doležel J. 2015. Molecular and cytogenetic characterization of wild *Musa* species. *PLoS One*, **10:** e0134096. DOI: https://doi.org/10.1371/journal.pone.0134096

Čížková J, Hřibová E, Humplíková L, Christelová P, Suchánková P, Doležel J. 2013. Molecular analysis and genomic organization of major DNA satellites in banana (*Musa* spp.). *PLoS One*, **8:** e54808. DOI: https://doi.org/10.1371/journal.pone.0054808

Cooper H, Spillane C, Hodgkin T. 2001. *Broadening the genetic bases of crop production*. IPGRI, FAO, CABI, Rome (Italy), Wallingford (UK). DOI: https://doi.org/10.1079/9780851994116.0000

De Langhe E, Hřibová E, Carpentier S, Doležel J, Swennen R. 2010. Did backcrossing contribute to the origin of hybrid edible bananas? *Annals of Botany* **106:** 849–857. DOI: https://doi.org/10.1093/aob/mcq187

De Langhe E, Perrier X, Donohue M, Denham T. 2015. The original Banana Split: Multi-disciplinary implications of the generation of African and Pacific Plantains in Island Southeast Asia. *Ethnobotany Research and Applications* **14:** 299–312. DOI: https://doi.org/10.17348/era.14.0.299-312

De Langhe E, Vrydaghs L, De Maret P, Perrier X, Denham TP. 2009. Why bananas matter: An introduction to the history of banana domestication. *Ethnobotany Research and Applications* **7:** 165–177. DOI: https://doi.org/10.17348/era.7.0.165-17.7

Denham T. 2010. From Domestication Histories to Regional Prehistory: Using Plants to Re-evaluate Early and Mid-Holocene Interaction between New Guinea and Southeast Asia. *Food & History* **8:** 3–22. DOI: https://doi.org/10.1484/J.FOOD.1.100971

Denham T, Haberle S, Lentfer C. 2004. New evidence and revised interpretations of early agriculture in Highland New Guinea. *Antiquity* **78:** 839–857. DOI: https://doi.org/10.1017/S0003598X00113481

D'Hont A, Denoeud F, Aury JM, Baurens FC, Carreel F, Garsmeur O, Noel B, Bocs S, Droc G, Rouard M, Da Silva C, Jabbari K, Cardi C, Poulain J, Souquet M, Labadie K, Jourda C, Lengellé J, Rodier-Goud M, Alberti A, Bernard M, Correa M, Ayyampalayam S, Mckain MR, Leebens-Mack J, Burgess D, Freeling M, Mbéguié-A-Mbéguié D, Chabannes M, Wicker T, Panaud O, Barbosa J, Hřibová E, Heslop-Harrison P, Habas R, Rivallan R, Francois P, Poiron C, Kilian A, Burthia D, Jenny C, Bakry F, Brown S, Guignon V, Kema G, Dita M, Waalwijk C, Joseph S, Dievart A, Jaillon O, Leclercq J, Argout X, Lyons E, Almeida A, Jeridi M, Doležel J, Roux N, Risterucci A.-M, Weissenbach J, Ruiz M, Glaszmann J-C, Quétier F, Yahiaoui N, Wincker P. 2012. The banana (*Musa acuminata*) genome and the evolution of monocotyledonous plants. *Nature* **488:** 213–217. DOI: https://doi.org/10.1038/nature11241

do Amaral CM, de Almeida dos Santos-Serejo J, de Oliveira e Silva S, da Silva Ledo CA, Amorim EP. 2015. Agronomic characterization of autotetraploid banana plants derived from 'Pisang Lilin' (AA) obtained through chromosome doubling. *Euphytica* **202:** 435–443. DOI: https://doi.org/10.1007/s10681-014-1320-0

Doležel J, Doleželová M, Roux N, Van den Houwe I. 1998. A novel method to prepare slides for high resolution chromosome studies in *Musa* spp. *Infomusa* **7:** 3–4.

Dupouy M, Baurens FC, Derouault P, Hervouet C, Cardi C, Cruaud C, Istace B, Labadie K, Guiougou C, Toubi L, Salmon F, Mournet P, Rouard M, Yahiaoui N, Lemainque A, Martin G, D'Hont A. 2019. Two large reciprocal translocations characterized in the disease resistance-rich *burmannica* genetic group of *Musa acuminata*. *Annals of Botany* **124:** 319–329. DOI: https://doi.org/10.1093/aob/mcz078

FAOSTAT, Agriculture Organization of the United Nations, FAO, 2017. Available from: www.fao.org.

Fauré S, Noyer JL, Horry JP, Bakry F, Lanaud C, de León DG. 1993. A molecular marker-based linkage map of diploid bananas (*Musa acuminata*). *Theoretical and Applied Genetics* **87:** 517–526. DOI: https://doi.org/10.1007/BF00215098

Han Y, Zhang T, Thammapichai P, Weng Y, Jiang J. 2015. Chromosome-specific painting in *Cucumis* species using bulked oligonucleotides. *Genetics* **200:** 771–779. DOI: https://doi.org/10.1534/genetics.115.177642

He L, Braz GT, Torres GA, Jiang JM. 2018. Chromosome painting in meiosis reveals pairing of specific chromosomes in polyploid *Solanum* species. *Chromosoma* **127:** 505–513. DOI: https://doi.org/10.1007/s00412-018-0682-9

Hippolyte I, Bakry F, Seguin M, Gardes L, Rivallan R, Risterucci AM, Jenny C, Perrier X, Carreel F, Argout X, Piffanelli P, Khan IA, Miller RNG, Pappas GJ, Mbéguié-A-Mbéguié D, Matsumoto T, De Bernardinis V, Huttner E, Kilian A, Baurens FC, D'Hont A, Cote F, Courtois B, Glaszmann JC. 2010. A saturated SSR/DArT linkage map of *Musa acuminata* addressing genome rearrangements among bananas. *BMC Plant Biology* **10:** 65. DOI: https://doi.org/10.1186/1471-2229-10-65.

Hippolyte I, Jenny C, Gardes L, Bakry F, Rivallan R, Pomies V, Cubry P, Tomekpe K, Risterucci AM, Roux N, Rouard M, Arnaud E, Kolesnikova-Allen M, Perrier X. 2012. Foundation characteristics of edible *Musa* triploids revealed from allelic distribution of SSR markers. *Annals of Botany* **109:** 937–951. DOI: https://doi.org/10.1093/aob/mcs010

Horry JP. 1989. Chimiotaxonomie et organisation génétique dans le genre *Musa* (III). *Fruits* **44:** 573–578.

Hřibová E, Čížková J, Christelová P, Taudien S, De Langhe E, Doležel J. 2011. The ITS-5.8S-ITS2 sequence region in the *Musaceae*: structure, diversity and use in molecular phylogeny. *PLoS One* **6:** e17863. DOI: https://doi.org/10.1371/journal.pone.0017863

Hu L, Liang F, Cheng D, Zhang Z, Yu G, Zha J, Wang Y, Xia Q, Yuan D, Tan Y, Wang D, Liang Y, Lin G. 2020. Location of balanced chromosome-translocation breakpoints by long-read sequencing on the Oxford Nanopore platform. *Frontiers in Genetics* **10:** 1330. DOI: https://doi.org/10.3389/fgene.2019.01313

Idziak D, Hazuka I, Poliwczak B, Wiszynska A, Wolny E, Hasterok R. 2014. Insight into the karyotype evolution of *Brachypodium* species using comparative chromosome barcoding. *PloS One* **9:** e93503. DOI: https://doi.org/10.1371/journal.pone.0093503

International trade statistics. Available from: www.wto.org.

Jiang J. 2019. Fluorescence *in situ* hybridization in plants: recent developments and future applications. *Chromosome Research* **27:** 153–165. DOI: https://doi.org/10.1007/ s10577-019-09607-z

Kagy V, Wong M, Vandenbroucke H, Jenny C, Dubois C, Ollivier A, Cardi C, Mournet P, Tuia V, Roux N, Doležel J, Perrier X. 2016. Traditional banana diversity in Oceania: An endangered heritage. *PLoS One* **11:** e0151208. DOI: https://doi.org/10.1371/journal.pone.0151208

Kitavi M, Downing T, Lorenzen J, Karamura D, Onyango M, Nyine M, Ferguson M, Spillane C. 2016. The triploid East African Highland Banana (EAHB) genepool is genetically uniform arising from a single ancestral clone that underwent population expansion by vegetative propagation. *Theoretical and Applied Genetics* **129:** 547–561. DOI: https://doi.org/10.1007/s00122-015-2647-1

Lebot V, Aradhya MK, Manshardt RM, Meilleur BA. 1993. Genetic relationships among cultivated bananas and plantains from Asia and the Pacific. *Euphytica* **67:** 163–175. DOI: https://doi.org/10.1007/BF00040618

Li LF, Wang HY, Zhang C, Wang XF, Shi FX, Chen WN, Ge XJ. 2013. Origins and domestication of cultivated banana inferred from chloroplast and nuclear genes. *PLoS One*, **8:** e80502. DOI: https://doi.org/10.1371/journal.pone.0080502

Lysák MA, Fransz PF, Ali HBM, Schubert I. 2001. Chromosome painting in *A. thaliana*. *Plant Journal* **28:** 689–697. DOI: https://doi.org/10.1046/j.1365-313x.2001.01194.x

Machado MA, Pieczarka JC, Silva FHR, O'Brien PCM, Ferguson-Smith MA, Nagamachi CY. 2018. Extensive karyotype reorganization in the fish *Gymnotus arapaima* (*Gymnotiformes, Gymnotidae*) highlighted by zoo-FISH analysis. *Frontiers in Genetics* **9:** 8. DOI: https://doi.org/10.3389/fgene.2018.00008

Martin G, Baurens FC, Droc G, Rouard M, Cenci A, Kilian A, Hastie A, Doležel J, Aury JM, Alberti A, Carreel F, D'Hont A. 2016. Improvement of the banana "*Musa acuminata*" reference sequence using NGS data and semi-automated bioinformatics methods. *BMC Genomics*, **17:** 1–12. DOI: https://doi.org/10.1186/s12864-016-2579-4

Martin G, Carreel F, Coriton O, Hervouet C, Cardi C, Derouault P, Roques D, Salmon F, Rouard M, Sardos J, Labadie K, Baurens FC, D'Hont A. 2017. Evolution of the banana genome (*Musa acuminata*) is impacted by large chromosomal translocations. *Molecular Biology and Evolution* **34:** 2140–2152. DOI: https://doi.org/10.1093/molbev/msx164

Martin G, Cardi C, Sarah G, Ricci S, Jenny C, Fondi E, Perrier X, Glaszmann J-C, D'Hont A, Yahiaoui N. 2020. Genome ancestry mosaics reveal multiple and cryptic contributors to cultivated banana. *Plant Journal* **102:** 1008–1025. DOI: https://doi.org/10.1111/tpj.14683

Mbanjo EGN, Tchoumbougnang F, Mouelle AS, Oben JE, Nyine M, Dochez C, Ferguson ME, Lorenzen J. 2012. Molecular marker-based genetic linkage map of a diploid banana population (*Musa acuminata* Colla). *Euphytica* **188:** 369–386. DOI: https://doi.org/10.1007/s10681-012-0693-1

Němečková A, Christelová P, Čížková J, Nyine M, Van den Houwe I, Svačina R, Uwimana B, Swennen R, Doležel J, Hřibová E. 2018. Molecular and cytogenetic study of East African Highland Banana. *Frontiers in Plant Science* **9:** 1371. DOI: https://doi.org/10.3389/fpls.2018.01371

Noumbissié GB, Chabannes M, Bakry F, Ricci S, Cardi C, Njembele JC, Yohoume D, Tomekpe K, Iskra-Caruana ML, D'Hont A, Baurens FC. 2016. Chromosome segregation in an allotetraploid banana hybrid (AAAB) suggests a translocation between the A and B genomes and results in eBSV-free offsprings. *Molecular Breeding* **36:** 38–52. DOI: https://doi.org/10.1007/s11032-016-0459-x

Nyine M, Uwimana B, Swennen R, Batte M, Brown A, Christelová P, Hřibová E, Lorenzen J, Doležel J. 2017. Trait variation and genetic diversity in a banana genomic selection training population. *PLoS One* **12:** e0178734. DOI: https://doi.org/10.1371/journal.pone.0178734

Ortiz R. 2013. Conventional banana and plantain breeding. *Acta Horticulturae* **986:** 177-194. DOI: https://doi.org/10.17660/ActaHortic.2013.986.19

Perrier X, Bakry F, Carreel F, Jenny F, Horry JP, Lebot V, Hippolyte I. 2009. Combining biological approaches to shed light on the evolution of edible bananas. *Ethnobotany Research and Applications* **7:** 199–216. DOI: https://doi.org/10.17348/era.7.0.199-216

Perrier X, De Langhe E, Donohue M, Lentfer C, Vrydaghs L, Bakry F, Carreel F, Hippolyte I, Horry JP, Jenny C, Lebot V, Risterucci AM, Tomekpe K, Doutrelepont H, Ball T, Manwaring J, de Maret P, Denham T. 2011. Multidisciplinary perspectives on banana (*Musa* spp.) domestication. *Proceedings of the National Academy of Sciences of the United States of America* **108:** 11311–11318. DOI: https://doi.org/10.1073/pnas.1102001108

Perrier X, Jacquemoud-Collet JP. 2006. DARwin software. http://darwin.cirad.fr/

Perrier X, Jenny C, Bakry F, Karamura D, Kitavi M, Dubois C, Hervouet C, Philippson G, De Langhe E. 2019. East African diploid and triploid bananas: a genetic complex transported from South-East Asia. *Annals of Botany* **123:** 19–36. DOI: https://doi.org/10.1093/aob/mcy156.

Price NS. 1995. The origin and development of banana and plantain cultivation. In: *Bananas and Plantains. World Crop Series* (Gowen S, eds)*,* Dordrecht: Springer. DOI: https://doi.org/10.1007/978-94-011-0737-2_1

Qu M, Li K, Han Y, Chen L, Li Z, Han Y. 2017. Integrated karyotyping of woodland strawberry (*Fragaria vesca*) with oligopaint FISH probes. *Cytogenetic and Genome Research* **153:** 158–164. DOI: https://doi.org/10.1159/000485283

Raboin LM, Carreel F, Noyer JL, Baurens FC, Horry JP, Bakry F, Tezenas Du Montcel H, Ganry J, Lanaud C, Lagoda PJL. 2005. Diploid ancestors of triploid export banana cultivars: Molecular identification of 2n restitution gamete donors and n gamete donors. *Molecular Breeding*, **16:** 333–341. DOI: https://doi.org/10.1007/s11032-005-2452-7

Robertson WRB. 1916. Chromosome studies. I. Taxonomic relationships shown in the chromosomes of *Tettigidae* and *Acrididae*: V-shaped chromosomes and their significance in *Acrididae*, *Locustidae*, and *Gryllidae*: Chromosomes and variation. *Journal of Morphology* **27:** 179–331. DOI: https://doi.org/10.1002/jmor.1050270202

Rouard M, Droc G, Martin G, Sardos J, Hueber Y, Guignon V, Cenci A, Geigle B, Hibbins MS, Yahiaoui N, Baurens F-C, Berry V, Hahn MW, D'Hont A, Roux N. 2018. Three new genome assemblies support a rapid radiation in *Musa acuminata* (wild banana). *Genome Biology and Evolution* **10:** 3129–3140. DOI: https://doi.org/10.1093/gbe/evy227

Roux N, Toloza A, Radecki Z, Zapata-Arias FJ, Doležel J. 2003. Rapid detection of aneuploidy in *Musa* using flow cytometry. *Plant Cell Reports* **21:** 483–490. DOI: https://doi.org/10.1007/s00299-002-0512-6

Sand C. 1989. Petite histoire du peuplement de l'Océanie. In: Ricard M, ed. Migrations et identités—Colloque CORAIL, Nouméa (PYF), 1988/11/21-22. Papeete: Université Française du Pacifique; pp. 39–40. Available: http://www.documentation.ird.fr/hor/fdi:27803.

Sandoval JA, Côte FX, Escoute J. 1996. Chromosome number variations in micropropagated true-to-type and off-type banana plants (*Musa* AAA Grande Naine cv.). *In Vitro Cellular & Developmental Biology – Plant* **32:** 14–17. DOI: https://doi.org/10.1007/BF02823007

Sedlazeck FJ, Rescheneder P, Smolka M, Fang H, Nattestad M, von Haeseler A, Schatz MC. 2018. Accurate detection of complex structural variations using single-molecule sequencing. *Nature Methods* **15:** 461–468. DOI: https://doi.org/10.1038/s41592-018-0001-7

Schubert I, Fransz PF, Fuchs J, De Jong JH. 2001. Chromosome painting in plants. *Methods in Cell Science* **23:** 57–69. DOI: https://doi.org/10.1023/A:1013137415093

Sharrock S. 1990. Collecting *Musa* in Papua New Guinea. Identification of genetic diversity in the genus *Musa*, ed Jarret RL (International Network for the Improvement of Banana and Plantain, Montpellier, France), pp. 140–157.

Shepherd K. 1999. Cytogenetics of the genus *Musa*. International Network for the Improvement of Banana and Plantain, Montpellier, France. ISBN: 2-910810-25-9

Shepherd K, Da Silva KM. 1996. Mitotic instability in banana varieties. Aberrations in conventional triploid plants. *Fruit*s **51:** 99–103.

Simmonds NW. 1962. The evolution of the bananas. Tropical Science Series. Longmans, London (GBR). 170p.

Šimoníková D, Němečková A, Karafiátová M, Uwimana B, Swennen R, Doležel J, Hřibová E. 2019. Chromosome painting facilitates anchoring reference genome sequence to chromosomes in situ and integrated karyotyping in banana (*Musa* spp.). *Frontiers in Plant Science* **10:** 1503. DOI: https://doi.org/10.3389/fpls.2019.01503

Soto DC, Shew C, Mastoras M, Schmidt JM, Sahasrabudhe R, Kaya G, Andrés AM, Dennis MY. 2020. Identification of structural variations in chimpanzees using optical mapping and nanopore sequencing. *Genes* **11:** 276. DOI: https://doi.org/10.3390/genes11030276

Tenkouano A, Vuylsteke D, Agogbua JU, Makumbi D, Swennen R, Ortiz R. 2003. Diploid banana hybrids TMB2x5105-1 and TMB2x9128-3 with good combining ability, resistance to black sigatoga and nematodes. *HortScience* **38:** 468–472. DOI: https://doi.org/10.21273/HORTSCI.38.3.468

Tomepke K, Jenny C, Escalant J-V. 2004. A review of conventional improvement strategies for *Musa*. *InfoMusa* **13:** 2-6.

Tugume AK, Lubega GW, Rubaihayo PR. 2003. Genetic diversity of East African Highland bananas using AFLP. *Infomusa* **11:** 28–32.

WCSP. 2018. World Checklist of Selected Plant Families. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet; http://wcsp.science.kew.org/ Retrieved.

A

M. acuminata ssp. *siamea* 'Pa Rayong' ITC0672

B

M. acuminata ssp. *burmannica* 'Tavoy' ITC0072

C

M. balbisiana 'Pisang Klutuk Wulung'

D

clone 'Pisang Lilin'

E

Mchare clones 'Huti white', 'Huti (Shumba nyeelu)' ITC1452, 'Ndyali' ITC1552

F

Gros Michel and Cavendish clones
'Gros Michel' ITC0484, 'Poyo' ITC1482

G

Mutika/Lujugira clone 'Imbogo' ITC0168

H

Mutika/Lujugira clone 'Kagera' ITC0141

Plantain clones '3 Hands Planty' ITC1132, 'Obino l'Ewai' ITC0109

J

Plantain clone 'Amou' ITC0963

K

Pelipita and Saba clones 'Pelipita' ITC0472, 'Saba sa Hapon' ITC1777

Supplementary Figure S2.

Supplementary Figure S3.

Supplementary Table S1.

Chromosome change	Diploid accessions														Triploid accessions								
	M. acuminata ssp. (AA)								Mchare cultivars (AA)			M. balbisiana (BB)		M. schizocarpa (SS)	AAA				AAB			ABB	
															Cavendish	Gros Michel	EAHB		Plantains			Pelipita	Saba
	malaccensis 'Pahang'**	*banksii* 'Banksii'	*microcarpa* 'Borneo'	*Zebrina* 'Maia Oa'	*burmannicoides* 'Calcutta 4'	*burmannica* 'Tavoy'	*siamea* 'Pa Rayong'	*P. lilin*	Huti White	Huti (Shumba nyeelu)	Ndyali	Tani**	Pisang Klutuk Wulung	Schizocarpa* *	Poyo	Gros Michel	Imbogo*	Kagera	3 Hands Planty	Amou*	Obino l'Ewai	Pelipita	Saba sa Hapon
3S/8L + 8S/3L				●●		●			●	●	●				●	●	●●	●●					
9S/1L + 1L/9S					●●	●	●●																
8L/2L					●●	●	●●																
7S/8S + 7L/8L						●																	
3L/4S							●																
4L/1L + 1L/4L								●	●	●	●				●	●							
7S/1L + telocentric 7L															●	●							
1L/3L (B genome specific)												●●	●●						●	●	●	●●	●●
7L/1L + missing 7S and 1S																	●						
missing chromosome 2																				●			

* aneuploid clones

** data from Šimoníková *et al*. (2019)