# Control of aperture closure during reach-to-grasp movements in immersive haptic-free virtual reality

**Madhur Mangalam[1,*], Mathew Yarossi[1,2,*], Mariusz P. Furmanek[1,3,*], and Eugene Tunik[1,2,4]**

[1]Department of Physical Therapy, Movement and Rehabilitation Science, Northeastern University, Boston, Massachusetts 02115, USA

[2]Department of Electrical and Computer Engineering, Northeastern University, Boston, Massachusetts 02115, USA

[3]Institute of Sport Sciences, The Jerzy Kukuczka Academy of Physical Education in Katowice, 40-065 Katowice, Poland

[4]Department of Bioengineering, Northeastern University, Boston, Massachusetts 02115, USA

*These authors contributed equally to this work.

Madhur Mangalam

m.mangalam@northeastern.edu

**Abstract**

Virtual reality (VR) has garnered much interest as a training environment for motor skill
acquisition, including for neurological rehabilitation of upper extremities. While the focus has
been on gross upper limb motion, VR applications that involve reaching for, and interacting
with, virtual objects are growing. The absence of true haptics in VR when it comes to hand-
object interactions raises a fundamentally important question: can haptic-free immersive
virtual environments (hf-VEs) support naturalistic coordination of reach-to-grasp movements?
This issue has been grossly understudied, and yet is of significant importance in the
development and application of VR across a number of sectors. In a previous study
(Furmanek et al. 2019), we reported that reach-to-grasp movements are similarly coordinated
in both the physical environment (PE) and hf-VE. The most noteworthy difference was that the
closure phase—which begins at maximum aperture and lasts through the end of the
movement—was longer in hf-VE than in PE, suggesting that different control laws might
govern the initiation of closure between the two environments. To do so, we reanalyzed data
from Furmanek et al. (2019), in which the participants reached to grasp three differently sized
physical objects, and matching 3D virtual object renderings, placed at three different
locations. Our analysis revealed two key findings pertaining to the initiation of closure in PE
and hf-VE. First, the respective control laws governing the initiation of aperture closure in PE
and hf-VE both included state estimates of transport velocity and acceleration, supporting a
general unified control scheme for implementing reach-to-grasp across physical and virtual
environments. Second, aperture was less informative to the control law in hf-VE. We suggest
that the latter was likely because transport velocity at closure onset and aperture at closure
onset were less independent in hf-VE than in PE, ultimately resulting in aperture at closure
onset having a weaker influence on the initiation of closure. In this way, the excess time and
muscular effort needed to actively bring the fingers to a stop at the interface of a virtual object
was factored into the control law governing the initiation of closure in hf-VE. Crucially, this
control law remained applicable, albeit with different weights in hf-VE, despite the absence of
terminal haptic feedback and potential perceptual differences.

***Keywords:*** collision detection; coordination, haptic feedback, prehension, rehabilitation,
virtual environment

## Introduction

Virtual reality (VR) has garnered much interest as a training environment for motor skill acquisition, including for neurological rehabilitation of upper extremities (Schultheis and Rizzo 2001; Sveistrup 2004; Holden 2005; Rizzo and Kim 2005; Rose et al. 2005; Adamovich et al. 2009; Cheung et al. 2014). The ability to reach-to and grasp virtual objects is paramount to any VR-based manual training. Devices that provide haptic feedback in VR tend to be expensive, bulky, and restrict natural motion in the virtual environment (Borst and Volz 2005; Pacchierotti et al. 2017; Culbertson et al. 2018). Haptic-free immersive virtual environments (hf-VEs) are a more viable option. However, the extent to which hf-VEs support naturalistic coordination of reach-to-grasp movements is mostly unknown. To fill this gap, in a previous study (Furmanek et al. 2019), we asked participants to reach to grasp physical objects of different sizes located at different distances, and to grasp virtual renderings of these objects, in an immersive hf-VE. We found that reach-to-grasp movements are similarly coordinated in both the physical environment (PE) and immersive hf-VE. The most noteworthy difference was that the closure—which begins at maximum aperture and lasts through the end of the movement—was longer in hf-VE than in PE. Understanding the factors underlying this difference is essential to modify a given hf-VE so that it provides for more naturalistic reach-to-grasp movements.

Much attention has been given to the processes that govern/trigger the initiation of closure during reach-to-grasp movements. Based on the observation that maximum aperture occurs approximately when hand transport deceleration is maximum, it was initially suggested that the initiation of closure is triggered by the reduction of hand transport velocity (Jeannerod 1984; Paulignan et al. 1991a, b). This suggestion was expanded by Rand and colleagues (2006a, b, 2008), who proposed that the initiation of closure is governed by a control law describing the dependence of the control action on the dynamics of the arm and hand in relation to the object location and size. According to this view, the distance of the hand from the object at closure is predicted by a linear function combining the state estimates of transport velocity, transport acceleration, and aperture. This control law has been shown to adequately describe the initiation of closure across different movement speeds (Rand et al. 2006b), object distances (Rand et al. 2010), and disease states, such as Parkinson's disease (Rand et al. 2006a).

To our knowledge, a comparison of the control laws governing the initiation of closure in PE and hf-VEs has not been made before despite reasons to believe that altered perceptual information and/or task constraints in hf-VEs might affect the planning, execution, and control of manual actions (Harris et al. 2019). For example, egocentric depth cues in VE may be processed differently by brain circuits than the natural depth cues in the real world, contributing to uncertainty in perceptual estimates of the object size and distance. Specific to hf-VEs, an absence of terminal haptic feedback may increase uncertainty in perception of the object size as haptics cannot be used to calibrate grasp (Bingham et al. 2007; Fukui and Inui 2013; Renner et al. 2013). Furthermore, the control law governing the initiation of closure in hf-VE might also differ because in PE, the closure is stopped instantaneously as soon as the fingers contact a physical object, but when reaching to grasp a virtual object, voluntary effort must be exerted to arrest finger movement to prevent interpenetration into that object (Prachyabrued and Borst 2012). Knowledge about whether lack of, or uncertainty about, key perceptual information in hf-VE compared to PE changes the control law used to govern closure is crucial as we aim to improve the fidelity of VR applications for manual skills training and rehabilitation.

In the present study, we reanalyzed data from Furmanek et al. (2019) to investigate whether the same control law governs the initiation of closure for reach-to-grasp movements performed in PE and our immersive hf-VE. We hypothesized that slower closure initiated farther from the object in the hf-VE than the PE condition that was noted by Furmanek et al. (2019) resulted from a difference in the control law governing the initiation of closure between the two environments.

To investigate the source of variation in the control low governing the initiation of closure in PE and hf-VE, we followed an information-theoretic approach to model selection (Burnham and Anderson 2002). This approach chooses the best model from a given set of a priori candidate models—by comparing the models against each other, instead of the null hypothesis, and balances the goodness of fit with simplicity. We considered the null model (containing only the intercept) and all the different combinations of the state estimates (aperture, transport velocity, and transport acceleration) to test whether the control law relies on each of these estimates in both PE and hf-VE and whether the strength of reliance on each of these estimates differs between the two environments.

## Methods

### *Participants*

Fifteen right-handed subjects (13 male, 12 female; mean±1*SD* age: 23.9±6.8 years old) with no reported muscular, orthopedic, or neurological health concerns, voluntarily participated in Furmanek et al. (2019) after providing verbal and written consent. Data from these participants were reanalyzed. The experimental task and procedure are described in brief below. Further details about the experimental task and procedure can be found in Furmanek et al. (2019).

### *Experimental task*

The participants performed the task of reaching toward and grasping three differently-sized physical objects and matching 3D virtual renderings. The objects were cuboids of identical height (8 cm) and depth (2.5 cm) but different widths: small: 3.6 cm; medium: 5.4 cm; large: 7.2 cm (Fig. 1). The physical objects were 3D printed using PLA thermoplastic (mass: small: 30 g; medium: 44 g; large: 59 g) and covered with glow-in-the-dark paint. The participants viewed blue-colored virtual renderings of these objects in a custom 3D immersive hf-VE designed in UNITY (ver. 5.6.1f1, Unity Technologies, San Francisco, CA) and presented via head-mounted display (Oculus Rift DK2, Oculus Inc., Menlo Park, CA). The objects, in PE or hf-VE, were placed at three different distances from the starting position—24 cm (near), 30 cm (middle), 36 cm (far)—and oriented at a 65° angle along the vertical axis to minimize excessive wrist extension (Fig. 1).

### *Experimental procedure*

The participants were seated in a chair with the right arm and hand placed on a table in front of them (Fig. 1). At the start position, the thumb and index finger straddled a 1.5 cm wide wooden peg located 12 cm in front and 24 cm to the right of the sternum, with the thumb depressing a switch. Lifting the thumb off the switch marked movement onset. A six-camera motion tracking system (PPT Studio N™, WorldViz, Inc., Santa Barbara, CA; sampling rate: 75 Hz) recorded the 3D motion of IRED markers attached to the participant's wrist (at the center of the segment running between the ulnar and radial styloid process), thumb nail, and index fingernail. A pair of IRED markers attached to the HMD co-registered the participant's

head motion to render the virtual environment. When reaching to grasp a virtual object in hf-
VE, the participants viewed the tips of their thumb and index finger as two 3D spheres (green
in color, 0.8 cm diameter), reflecting the real-time 3D position of the respective IRED marker.
When reaching to grasp a physical object, all lights were turned off so that only the illuminated
IRED markers attached to the thumb and index finger, as well as the glow-in-the-dark object,
were visible to the participants. The overhead light was turned on after every block of 12 trials
to prevent acclimatization of the vision to the dark environment.

Prior to the experiment, the participants were allowed to practice the reach-to-grasp
task in both PE and hf-VE until they felt comfortable in both environments. The experiment
consisted of 216 trials, each trial lasting 3 seconds: 108 trials in PE and 108 trials in hf-VE (3
objects × 3 distances/object × 12 trials/ or each size–distance combination in each
environment). The trials were blocked for each size-distance combination and each
environment. The order of the presentation of the three objects and their placement was
randomized for each block of size-distance combination.

In each trial, an auditory cue—a beep—signaled the participants to reach for, grasp,
and lift the object. The participants were considered to have grasped a physical object the
moment the tips of their thumb and index finger came into contact with that object's lateral
surface, and aperture did not change more than 1 mm for at least 50 ms. The participants
were considered to have grasped a virtual object the instance a custom collision detection
algorithm detected that both 3D spheres had come in contact with the virtual object. Once
grasp of the virtual object was detected, the object changed color from blue to red. The
participants lifted and raised each object briefly before returning their hand to the starting
position.

### Data processing and feature extraction

All data were analyzed offline using custom MATLAB routines (Mathworks Inc., Natick,
MA). For each trial, the time series data were cropped from movement onset (and the
moment the switch was released) to movement offset (the moment aperture velocity became
zero in PE, and the moment the collision detection criterion was met in hf-VE) and filtered (6
Hz, fourth-order low-pass Butterworth filter).

For each condition (environment, object size, and object distance), mean values were
calculated on a per-participant basis for the following outcome variables: (a) Movement time,
defined as the interval between movement onset and movement offset; (b) Maximum
aperture, which marked the initiation of closure or closure onset (CO) and which we refer to
as aperture at CO; (c) Opening time, defined as the interval from movement onset to CO; (d)
Closure time, defined as the interval from CO to movement offset; (e) Transport distance,
defined as the distance between the initial and final positions of the wrist; (f) Opening
distance, defined as the distance between the wrist's position at movement onset and the
object; (g) Closure distance, defined as the distance between the wrist's position at CO and
the object; (h) Maximum closure velocity; (i) Maximum closure deceleration; (j) Transport
velocity at CO; and (k) Transport acceleration at CO.

### Statistical analysis

A 2 × 3 × 3 repeated measures analysis of variance (rm-ANOVA) with within-subject
factors of the Environment (PE, hf-VE), Object size (Small, Medium, Large), and Object
distance (levels: Near, Middle, Far) was used to evaluate the effects of the Environment,
Object size, and Object distance on each kinematic variable. All data mostly fulfilled the
conditions of normality, homogeneity of variance, and sphericity; when the assumption of

sphericity was not met, Greenhouse-Geisser correction was applied, and in these cases, the
corrected *p*-values and the Greenhouse-Geisser correction factor ($\varepsilon$) are reported. Pearson's
correlation tests were used to examine the relationship between opening/closure time and
movement time in both PE and hf-VE, as well as between opening/closure distance and
transport distance in both PE and hf-VE. Each test was performed in SPSS 21 (IBM Inc.,
Chicago, IL), and each test statistic was considered significant at the two-tailed alpha level of
0.05. All effect sizes are reported as partial eta-squared ($\eta_p^2$).

To investigate which state estimates best explained variation in closure distance in PE
and hf-VE, we followed an information-theoretic approach to model selection using RStudio.
This approach uses the Akaike Information Criterion (AIC; or quasi-AIC (QAICc) for over-
dispersed data to choose a set of plausible models from a given set of *a priori* candidate
models (Burnham and Anderson 2002). AICc for a given model is computed as:
$AIC_c = -\ln[l(\hat{\theta}|data)] + 2K$ , where $l(\hat{\theta}|data)$ is the likelihood of the parameters ( $\hat{\theta}$ ) in
the model and $K$ = number of estimable parameters in the model. For example, a simple
linear regression model, y = *a* + *b*x, has three estimable parameters: in intercept *a*, the slope
b, and the variance ($s^2$). This definition of AIC is applicable for large samples (*n*). In the small-
sample case (i.e., *n*/*K* < 40), $AIC_c = -\ln[l(\hat{\theta}|data)] + 2K + \dfrac{2K(K+1)}{n-K+1}$ . To compute QAICc, the
number of model parameters is increased by 1 to account for overdispersion.

QAICc serves as an estimator of out-of-sample prediction error and thereby the relative
quality of statistical models for a given set of data. QAICc estimates the quality of each model
relative to each of the other models. Specifically, a smaller QAICc value reflects better
performance/complexity trade-off. Thus, QAICc provides a means for selecting the model with
the best performance/complexity trade-off. In the present study, we considered eight
candidate models, including the null model (containing only the intercept) and all the subset
models of the following full model: CD = *a* + *b*($A_{CO}$) + *c*($TV_{CO}$) + *d*($TA_{CO}$), by allowing the
intercept to differ by participant, where CD = closure distance, $A_{CO}$ = aperture at CO, $TV_{CO}$ =
transport velocity at CO, and $TA_{CO}$ = transport acceleration at CO.

**Results**

Figure 2 provides spatial trajectories of the reach (left) and grasp (right) components of
the reach-to-grasp movements performed in PE and hf-VE for a representative object size-
distance pair (medium object, middle distance). Table 1 summarizes the mean (±1*SD*) values
of each movement variable across all participants (*n* = 13). Table 2 summarizes the outcomes
of rm-ANOVAs. Test statistics only for significant effects are presented below (see Table 2 for
non-significant outcomes.

**Closure lasted longer and was initiated farther from the object, with lower peak**
**deceleration, in hf-VE than in PE**

Closure lasted longer in hf-VE than in PE (rm-ANOVA: $F_{1,12}$ = 19.24, *p* = 0.001, $\eta_p^2$ =
0.62), and this effect of environment was mediated by object size ($F_{2,12}$ = 7.69, *p* = 0.003, $\eta_p^2$ =
0.39; Fig. 3A). Closure was initiated farther from the object in hf-VE than in PE ($F_{1,12}$ = 15.63,
*p* = 0.002, $\eta_p^2$ = 0.57), an effect also mediated by object size ($F_{2,12}$ = 6.77, *p* = 0.005, $\eta_p^2$ =
0.36; Fig. 3B). The maximum closure velocity did not differ between the two environments
($F_{1,12}$ = 1.77, *p* = 0.208, $\eta_p^2$ = 0.13; Fig. 3C), but maximum closure deceleration was higher in

hf-VE than in PE ($F_{1,12}$ = 8.55, $p$ = 0.015, $\eta_p^2$ = 0.42; Fig. 3D). Thus, closure lasted longer and
was initiated farther from the object, with lower peak deceleration, in hf-VE than in PE.

**Closure scaled in time but not distance from the object in both PE and hf-VE**

To examine whether the initiation of closure occurred at a consistent time relative to
movement onset or grasping of the object in PE and hf-VE, we analyzed the relationship
between opening/closing time and total movement time. Opening time showed a strong
positive relationship with movement time in both PE (Pearson's correlation, $\rho$ = 0.85, $p$ <
0.001; Fig. 4A) and hf-VE ($\rho$ = 0.76, $p$ < 0.001; Fig. 4B), and closing time showed a weak
positive relationship with movement time in both PE ($\rho$ = 0.50, $p$ < 0.001; Fig. 4A) and hf-VE
($\rho$ = 0.43, $p$ < 0.001; Fig. 4B). The slopes of the regression lines for both opening time (0.72
in PE vs. 0.71 in hf-VE) and closing time (0.27 in PE vs. 0.29 in hf-VE) were similar between
the two environments.

To examine whether the initiation of closure occurred at a consistent distance relative
to the initial or final transport position in PE and hf-VE, we analyzed the relationship between
opening/closing distance and total transport distance. Opening distance showed a strong
positive relationship with transport distance in both PE ($\rho$ = 0.98, $p$ < 0.001; Fig. 4C) and hf-
VE ($\rho$ = 0.91, $p$ < 0.001; Fig. 4D). In contrast, closing distance did not show a significant
relationship with transport distance in PE ($\rho$ = 0.11, $p$ = 0.505; Fig. 4C) or hf-VE ($\rho$ = 0.14, $p$ =
0.306; Fig. 4D). The slopes of the regression lines for both opening distance (0.98 in PE vs.
0.94 in hf-VE) and closing distance (0.024 in PE vs. 0.060 in hf-VE) were similar between the
two environments.

**Similar control laws governed the initiation of closure in PE and hf-VE**

Of the eight models used to identify the control law governing the initiation of closure in
PE, only two models yielded nonzero weights, $w_i$, (the probability that a given model is the
best model among all models). Support for the first model was 999 times stronger than the
second model (evidence ratio = $w_1/w_2$ = 0.9900/0.0010 = 999; $\chi^2$ = 16.66, $p$ < 0.001, Table 3).
The first model yielded both the best fit (log-likelihood closest to zero) and the best
performance-to-complexity tradeoff (lowest QAICc). This model included a coefficient for $A_{CO}$,
$TV_{CO}$, and $TA_{CO}$, supporting the idea presented by Rand and coworkers (Rand et al. 2006a, b)
that the distance at closure is predicted by a linear function combining the state estimates of
transport velocity, transport acceleration, and aperture.

Of the eight models used to identify the control law governing the initiation of closure in
hf-VE, only two models yielded non-negligible $w_i$. Support for the first model was 3.2 times
stronger than the second model (evidence ratio = $w_1/w_2$ = 0.73/0.27 = 2.50; Table 3). While
the model with the best fit (model 2, log-likelihood closest to zero) included a coefficient for all
three variables, the model with the best performance-to-complexity tradeoff (model 1, lowest
QAICc) did not include a coefficient for aperture at CO. Nonetheless, the first and second
models yielded close goodness of fit (log-likelihood values: –85.05 and –84.86, respectively)
and the inclusion of $A_{CO}$ in the model did not significantly improve the fit ($\chi^2$ = 0.38, $p$ = 0.537).
In short, the initiation of closure in hf-VE was not as dependent on aperture at CO as it did in
PE.

To further explore the contribution of different state variables to the respective control
laws, we analyzed the relationship between the values of closure distance and each state
variable averaged across all participants. Closure distance showed a negative relationship
with aperture at CO in both environments (PE: Pearson's $r$ = –0.78, $p$ = 0.013; hf-VE: $r$ = –
0.81, $p$ = 0.008; Fig. 5A, left), a strong positive relationship with transport velocity at CO (PE:

$r = 0.94$, $p < 0.001$; hf-VE: $r = 0.98$, $p < 0.001$; Fig. 5B, left), and a weak negative relationship
with transport acceleration at CO (PE: $r = -0.60$, $p = 0.088$; hf-VE: $r = -0.64$, $p = 0.067$; Fig.
5C, left). The strong relationship between closure distance and transport velocity at CO in
both PE and hf-VE indicates that it is the most important state variable contributing to the
control laws. This result confirms that of the model selection in that the model with the best fit
and performance-to-complexity tradeoff for PE, and all four models with nonzero weight, $w_i$,
for hf-VE, included a coefficient for transport velocity at CO (Table 3).

To substantiate that the state estimate of aperture contributed to the variability in the
control law governing the initiation of closure between PE and hf-VE, we compared the
relationship between each state variable and closure distance between PE and hf-VE. In both
PE and hf-VE, closure distance showed a strong negative relationship with aperture at CO
(PE: Pearson's $r = -0.78$, $p = 0.013$; hf-VE: $r = -0.81$, $p = 0.008$; Fig. 5A, left) and a very
strong positive relationship with transport velocity at CO (PE: $r = 0.94$, $p < 0.001$; hf-VE: $r = $
$0.98$, $p < 0.029$; Fig. 5B, left) but did not show a significant relationship with transport
acceleration at CO (PE: $r = -0.060$, $p = 0.088$; hf-VE: $r = -0.064$, $p = 0.067$; Fig. 5C, left). At
the individual participant level, the slope of the regression line between closure distance and
aperture at CO was steeper in hf-VE than in PE (paired $t$-test: $t_{2,13} = 4.40$, $p = 0.029$; Fig. 5A,
right). Crucially, the slope of the regression line did not differ for transport velocity at CO ($t_{2,13}$
$= -1.90$, $p = 0.082$; Fig. 5B, right) and transport acceleration at CO ($t_{2,13} = 1.81$, $p = 0.095$; Fig.
5C, right).

Finally, to test whether a difference in the relationship between aperture at CO and
transport velocity at CO might be a possible reason for why the initiation of closure in hf-VE
did not depend on aperture at CO as it did in PE, we compared relationships among the three
state parameters between PE and hf-VE. In both PE and hf-VE, aperture at CO showed a
strong relationship with transport velocity at CO (PE: Pearson's $r = -0.72$, $p = 0.023$; hf-VE: $r = $
$-0.88$, $p = 0.002$; Fig. 6A, left) but no significant relationship with transport acceleration at CO
(PE: $r = 0.38$, $p = 0.309$; hf-VE: $r = 0.21$, $p = 0.594$; Fig. 6B, left). Transport velocity at CO
showed a strong negative relationship with transport acceleration at CO in PE ($r = -0.72$, $p = $
$0.029$) but no significant relationship in hf-VE ($r = -0.62$, $p = 0.072$; Fig. 6C, left). At the
individual participant level, the slope of the regression line between aperture at CO and
transport velocity at CO was steeper in hf-VE than in PE (paired $t$-test: $t_{2,13} = 4.47$, $p < 0.001$;
Fig. 6A, right). Crucially, the slope of the regression line between aperture at CO and
transport acceleration at CO did not differ between PE and hf-VE ($t_{2,13} = -0.11$, $p = 0.917$; Fig.
6B, right), and the slope of the regression line between transport velocity at CO and transport
acceleration at CO was only marginally steeper in hf-VE than in PE ($t_{2,13} = 2.21$, $p = 0.047$;
Fig. 6C, right).

## Discussion

In the present study, reanalysis of the Furmanek et al. (2019) data indicated that the
closure phase was also initiated farther from the object, with smaller peak deceleration, in
immersive hf-VE than in PE. These differences presumably reflect the fact that when reaching
to grasp in hf-VE, the voluntary effort needed to actively bring the fingers to a stop at the
interface of a virtual object. PE does not impose this physical constraint, as the fingers
automatically stop when they contact the physical object. However, the goal of this study was
not merely to identify differences in the initiation of closure between the two environments; the
goal of the present study was to investigate potential differences and similarities in the control
laws governing the initiation of closure in PE and an immersive hf-VE. To do so, a linear
regression model—previously suggested by Rand et al. (2008) to predict the distance from

the object at which the closure is initiated from the object from the limb state parameters of transport velocity, transport acceleration, and peak aperture—was applied to kinematic data from each environment. The model comparisons yielded two important insights: (1) Transport velocity was the dominant predictor of closure distance in both environments, and the relationship between transport velocity at CO and closure distance was nearly identical between environments, and (2) Aperture was less informative to the control law in hf-VE.

In hf-VE, two factors significantly influence the speed-accuracy tradeoff in reach-to-grasp movements. The first is the well-documented finding that VR increases the perceptual uncertainty about object distance and size (Armbrüster et al. 2008; Ogawa et al. 2018; Harris et al. 2019). The increase in perceptual uncertainty about the spatial properties of objects, which is hypothesized to cause slower movements in hf-VE (Viau et al. 2004; Magdalon et al. 2011), was likely a major factor in smaller peak transport velocity found in the present study. The second factor influencing the speed-accuracy tradeoff is the precision demand of grasping. The lack of physical properties of the object in hf-VE not only requires the voluntary effort needed to actively bring the fingers to a stop at the interface of a virtual object, but also increases the demand for precision of the finger aperture in the closure phase of the movement.

Investigations of the speed-accuracy tradeoff for reach to grasp have shown that the full reach-to-grasp movement is poorly described by the Fitts law (Bootsma et al. 1994). Instead, it may be better characterized by Welford's (1968) theoretical position that object distance and size place separate constraints upon movement time (McIntosh et al. 2018). However, Rand et al. (2008) have suggested that the precision of the closure phase specifically may be reasonably well described by principles of speed-accuracy tradeoff as described by the standard form of the Fitts' law (Fitts 1954). Based on the finding that controlled manipulations of movement speed did not alter the relationship between transport velocity at CO and closure distance, Rand et al. (2006b) reasoned that variability in the opening process might be more acceptable as long as the transport-aperture coordination at the time of the phase transition to closure is tightly controlled. Our data support this assertion by Rand for movements conducted in hf-VE with one notable exception. Rand et al. (2006b) found faster movements (i.e., movements with greater peak transport velocity) were associated with higher transport velocity at CO, and the closure initiated farther from the object. In contrast, we observed that the movements performed in hf-VE, despite having significantly lower peak transport velocity than those performed in PE (Furmanek et al. 2019), were characterized by higher transport velocity at CO and closure distance farther from the object. Indeed, this finding appears to be consistent with previous empirical results suggesting that precision demands of grasping are associated with an extended deceleration phase, suggesting an important role for feedback-based correction (MacKenzie et al. 1987; Bootsma et al. 1994; MacKenzie and Graham 1997). We suggest that a parsimonious explanation for this difference between our result and that of Rand et al. (2006b) is that movement plan at the time of movement initiation factored the specific task and precision demands of the closure in hf-VE to extend the relationship between transport velocity at CO and closure distance learned over a lifetime of practice in PE. The participants treated the virtual object as if it were a physical object with greater precision demands for grasping (Gentilucci et al. 1991; Iyengar et al. 2009; Magdalon et al. 2011). The striking similarity of the relationship between transport velocity at CO and closure distance offers valuable insights into the critical importance of preserving this aspect of the phase transition between the opening and closing of the grasp for successful reach to grasp. We suggest that the preservation of this relationship may be a critical benchmark for any VE in which reach-to-grasp tasks are performed.

Our second insight was that aperture was less informative to the control law in hf-VE.
This, too, appears to relate directly to uncertainty about spatial properties of the object and
precision demands of the task. It has been shown that calibration of grasp size with reach
distance allows executing naturalistic reach-to-grasp movements even without haptic terminal
feedback (e.g., Bingham et al. 2007). A lack of any opportunity to calibrate grasp size with
reach distance in the present study might have led to uncertainty about object size in hf-VE.
Indeed, the finding that closure distance scaled differently with object size in PE and hf-VE
speaks directly to the perceptual uncertainty about object size in hf-VE. Furthermore, aperture
at CO was less independent in hf-VE than in PE, ultimately resulting in the aperture at CO
having a weaker influence on the initiation of closure. Indeed, the strong negative relationship
between transport velocity and aperture at CO in hf-VE (Fig. 6A, left) speaks to the deliberate
slowing of hand transport in situations with higher precision demand, as expected of hf-VE.

The present findings, along with those of Furmanek et al. (2019), show that immersive
hf-VE can partially support naturalistic reach-to-grasp movements. These results are
especially relevant to the study of the role that terminal feedback may have in naturalistic
reach-to-grasp movements and to reproducing naturalistic reach-to-grasp movements in
immersive hf-VEs. When reaching to grasp a physical object, one of the important roles of
terminal haptic feedback is to convey, via somatosensory signals, that the object has been
grasped successfully. This feedback may benefit subsequent actions by updating and
recalibrating forward models controlling reach-to-grasp movements so that the grasp size and
reach distance can be scaled appropriately (Coats et al. 2008; Cavina-Pratesi and Hesse
2013). Notably, in the present study, when a collision detection algorithm detected that both
3D spheres (denoting the 3D position of the fingertips in hf-VE) had come into contact with
the virtual object, the object's color changed from blue to red to provide [terminal] visual
feedback that the object had been grasped. This visual feedback could presumably substitute
for the haptic feedback available upon grasping a physical object, and thus the reach-to-grasp
movements were coordinated similarly in both PE and hf-VE (Prachyabrued and Borst 2014;
Geiger et al. 2018). Our future work will investigate how bio-inspired collision detection
algorithms can help thinning the remaining gap between prehensile actions performed in PE
and immersive hf-VE.

To conclude, the present findings addresses the concern that the planning, execution,
and control of manual actions in hf-VEs than in  PE may involve distinct mechanisms (Harris
et al. 2019). Such a distinction implies that VR might have certain limitations in its utility for
training and rehabilitation. However, we show that albeit with different weights in hf-VE, a
control law that involves sensorimotor integration of the internal state estimates of the velocity
and acceleration of the arm and perceptual estimate object size dictates the initiation of
closure in both PE and hf-VE. This finding suggests that haptic-free immersive virtual
environments support naturalistic coordination of reach-to-grasp movements and can be a
viable option for low-cost home-based personalized training and rehabilitation, as well as for
behavioral and neurophysiological investigations of reach-to-grasp movements and it's
underlying control mechanisms.

## Acknowledgments

This work was supported by NIH grants #R01NS085122 and #2R01HD058301, and NSF grants #CBET-1804550 and #CMMI-M3X-1935337, to Eugene Tunik. We thank Alex Hunton and Samuel Berin for developing the VR platform.

## Author contributions

M.M., M.Y., M.P.F., and G.T. conceived and designed research; M.P.F. performed experiments; M.M., M.P.F., and M.P.F. analyzed data; M.M., M.Y., M.P.F., and G.T. interpreted results of experiments; M.M. prepared figures; M.M. drafted manuscript; M.M., M.Y., M.P.F., and G.T. edited and revised manuscript; M.M., M.Y., M.P.F., and G.T. approved final version of manuscript.

## Conflict of interest

The authors declare that no competing interests exist.

## References

Adamovich S V, Fluet GG, Eugene T, Merians AS (2009) Sensorimotor training in virtual reality: A review. NeuroRehabilitation 25:25–44. https://doi.org/10.3233/NRE-2009-0497

Armbrüster C, Wolter M, Kuhlen T, et al (2008) Depth perception in virtual reality: Distance estimations in peri- and extrapersonal space. CyberPsychology Behav 11:9–15. https://doi.org/10.1089/cpb.2007.9935

Bingham G, Coats R, Mon-Williams M (2007) Natural prehension in trials without haptic feedback but only when calibration is allowed. Neuropsychologia 45:288–294. https://doi.org/10.1016/j.neuropsychologia.2006.07.011

Bootsma RJ, Marteniuk RG, MacKenzie CL, Zaal FTJM (1994) The speed-accuracy trade-off in manual prehension: Effects of movement amplitude, object size and object width on kinematic characteristics. Exp Brain Res 98:535–541. https://doi.org/10.1007/BF00233990

Borst CW, Volz RA (2005) Evaluation of a haptic mixed reality system for interactions with a virtual control panel. Presence Teleoperators Virtual Environ 14:677–696. https://doi.org/10.1162/105474605775196562

Burnham KP, Anderson DR (2002) Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach. Springer-Verlag, New York, NY

Cavina-Pratesi C, Hesse C (2013) Why do the eyes prefer the index finger? Simultaneous recording of eye and hand movements during precision grasping. J Vis 13:15. https://doi.org/10.1167/13.5.15

Cheung KL, Tunik E, Adamovich S V, Boyd LA (2014) Neuroplasticity and virtual reality. In: Weiss PL (Tamar), Keshner EA, Levin MF (eds) Virtual Reality for Physical and Motor Rehabilitation. Springer New York, New York, NY, pp 5–24

Coats R, Bingham GP, Mon-Williams M (2008) Calibrating grasp size and reach distance: Interactions reveal integral organization of reaching-to-grasp movements. Exp Brain Res 189:211–220. https://doi.org/10.1007/s00221-008-1418-5

Culbertson H, Schorr SB, Okamura AM (2018) Haptics: The present and future of artificial touch sensation. Annu Rev Control Robot Auton Syst 1:385–409. https://doi.org/10.1146/annurev-control-060117-105043

Fitts PM (1954) The information capacity of the human motor system in controlling the amplitude of movement. J Exp Psychol 47:381–391. https://doi.org/10.1037/h0055392

Fukui T, Inui T (2013) How vision affects kinematic properties of pantomimed prehension movements. Front Psychol 4:44. https://doi.org/10.3389/fpsyg.2013.00044

Furmanek MP, Schettino LF, Yarossi M, et al (2019) Coordination of reach-to-grasp in physical and haptic-free virtual environments. J Neuroeng Rehabil 16:78. https://doi.org/10.1186/s12984-019-0525-9

Geiger A, Bewersdorf I, Brandenburg E, Stark R (2018) Visual feedback for grasping in virtual reality environments for an interface to instruct digital human models. In: Ahram T, Falcão C (eds) Advances in Usability and User Experience. Springer International Publishing, Cham, Switzerland, pp 228–239

Gentilucci M, Castiello U, Corradini ML, et al (1991) Influence of different types of grasping on the transport component of prehension movements. Neuropsychologia 29:361–378. https://doi.org/10.1016/0028-3932(91)90025-4

Harris DJ, Buckingham G, Wilson MR, Vine SJ (2019) Virtually the same? How impaired sensory information in virtual reality may disrupt vision for action. Exp Brain Res 237:2761–2766. https://doi.org/10.1007/s00221-019-05642-8

Holden MK (2005) Virtual environments for motor rehabilitation: Review. Cyberpsychology Behav 8:187–211. https://doi.org/10.1089/cpb.2005.8.187

Iyengar V, Santos MJ, Aruin AS (2009) Role of movement velocity on the magnitude of grip force while lifting an object with touch from the contralateral finger. Motor Control 13:130–141. https://doi.org/10.1123/mcj.13.2.130

Jeannerod M (1984) The timing of natural prehension movements. J Mot Behav 16:235–254. https://doi.org/10.1080/00222895.1984.10735319

MacKenzie CL, Graham ED (1997) Separating A and W effects: Pointing to targets on computer displays. Behav Brain Sci 20:316–318. https://doi.org/DOI: 10.1017/S0140525X97361445

MacKenzie CL, Marteniuk RG, Dugas C, et al (1987) Three-dimensional movement trajectories in Fitts' task: Implications for control. Q J Exp Psychol Sect A 39:629–647. https://doi.org/10.1080/14640748708401806

Magdalon EC, Michaelsen SM, Quevedo AA, Levin MF (2011) Comparison of grasping movements made by healthy subjects in a 3-dimensional immersive virtual versus physical environment. Acta Psychol (Amst) 138:126–134. https://doi.org/10.1016/j.actpsy.2011.05.015

McIntosh RD, Mon-Williams M, Tresilian JR (2018) Grasping at laws: Speed-accuracy trade-offs in manual prehension. J Exp Psychol Hum Percept Perform 44:1022–1037. https://doi.org/10.1037/xhp0000512

Ogawa N, Narumi T, Hirose M (2018) Object size perception in immersive virtual reality: Avatar realism affects the way we perceive. 2018 IEEE Conf Virtual Real 3D User Interfaces 647–648. https://doi.org/10.1109/VR.2018.8446318

Pacchierotti C, Sinclair S, Solazzi M, et al (2017) Wearable haptic systems for the fingertip and the hand: Taxonomy, review, and perspectives. IEEE Trans Haptics 10:580–600. https://doi.org/10.1109/TOH.2017.2689006

Paulignan Y, Jeannerod M, MacKenzie C, Marteniuk R (1991a) Selective perturbation of visual input during prehension movements. 2. The effects of changing object size. Exp Brain Res 87:407–420. https://doi.org/10.1007/BF00231858

Paulignan Y, MacKenzie C, Marteniuk R, Jeannerod M (1991b) Selective perturbation of visual input during prehension movements. 1. The effects of changing object position. Exp Brain Res 83:502–512. https://doi.org/10.1007/BF00229827

Prachyabrued M, Borst CW (2012) Visual interpenetration tradeoffs in whole-hand virtual grasping. 2012 IEEE Symp 3D User Interfaces 39–42. https://doi.org/10.1109/3DUI.2012.6184182

Prachyabrued M, Borst CW (2014) Visual feedback for virtual grasping. 2014 IEEE Symp 3D User Interfaces 19–26. https://doi.org/10.1109/3DUI.2014.6798835

Rand MK, Shimansky YP, Hossain ABMI, Stelmach GE (2008) Quantitative model of transport-aperture coordination during reach-to-grasp movements. Exp Brain Res 188:263–274. https://doi.org/10.1007/s00221-008-1361-5

Rand MK, Shimansky YP, Hossain ABMI, Stelmach GE (2010) Phase dependence of transport–aperture coordination variability reveals control strategy of reach-to-grasp movements. Exp Brain Res 207:49–63. https://doi.org/10.1007/s00221-010-2428-7

Rand MK, Smiley-Oyen AL, Shimansky YP, et al (2006a) Control of aperture closure during reach-to-grasp movements in Parkinson's disease. Exp Brain Res 168:131–142. https://doi.org/10.1007/s00221-005-0073-3

Rand MK, Squire LM, Stelmach GE (2006b) Effect of speed manipulation on the control of aperture closure during reach-to-grasp movements. Exp Brain Res 174:74–85. https://doi.org/10.1007/s00221-006-0423-9

Renner RS, Velichkovsky BM, Helmert JR (2013) The perception of egocentric distances in
virtual            environments-a         review.         ACM        Comput        Surv        46:1–40.
https://doi.org/10.1145/2543581.2543590

Rizzo A "Skip", Kim GJ (2005) A SWOT analysis of the field of virtual reality rehabilitation and
therapy.        Presence        Teleoperators        Virtual        Environ        14:119–146.
https://doi.org/10.1162/1054746053967094

Rose FD, Brooks BM, Rizzo AA (2005) Virtual reality in brain damage rehabilitation: Review.
CyberPsychology Behav 8:241–262. https://doi.org/10.1089/cpb.2005.8.241

Schultheis MT, Rizzo AA (2001) The application of virtual reality technology in rehabilitation.
Rehabil Psychol 46:296–311. https://doi.org/10.1037/0090-5550.46.3.296

Sveistrup H (2004) Motor rehabilitation using virtual reality. J Neuroeng Rehabil 1:10.
https://doi.org/10.1186/1743-0003-1-10

Viau A, Feldman AG, McFadyen BJ, Levin MF (2004) Reaching in reality and virtual reality: A
comparison of movement kinematics in healthy subjects and in adults with hemiparesis. J
Neuroeng Rehabil 1:11. https://doi.org/10.1186/1743-0003-1-11

Welford AT (1968) The Fundamentals of Skill. Methuen, London, UK

**Table 1** Mean (±1*SD*) values of each movement variable across all participants (*n* = 13) for reach-to-grasp movements
performed in PE and hf-VE

| Variable | Object size /distance | PE | | | hf-VE | | |
|---|---|---|---|---|---|---|---|
| | | Near | Middle | Far | Near | Middle | Far |
| Closure time [ms] | | | | | | | |
| | Small | 344±64 | 361±96 | 392±108 | 418±120 | 411±114 | 421±119 |
| | Medium | 314±63 | 324±56 | 346±78 | 413±91 | 415±111 | 441±98 |
| | Large | 300±59 | 319±58 | 314±48 | 416±128 | 423±119 | 430±97 |
| Closure distance [cm] | | | | | | | |
| | Small | 3.1±1.2 | 3.1±1.0 | 3.4±1.4 | 4.7±2.0 | 5.1±2.4 | 5.6±2.7 |
| | Medium | 2.4±0.9 | 2.6±1.0 | 3.0±1.2 | 4.0±1.7 | 4.3±2.3 | 4.8±2.9 |
| | Large | 2.5±0.8 | 2.6±0.9 | 2.6±0.8 | 3.3±1.7 | 3.6±1.9 | 4.2±2.2 |
| Maximum closure velocity [cm/s] | | | | | | | |
| | Small | −15.8±5.7 | −16.0±6.8 | −15.6±6.3 | −12.8±4.6 | −16.0±4.6 | −15.6±4.1 |
| | Medium | −15.8±5.5 | −15.2±6.6 | −15.3±5.6 | −13.0±4.4 | −13.9±5.4 | −13.5±3.8 |
| | Large | −14.1±4.5 | −13.7±5.0 | −13.2±4.6 | −13.2±5.9 | −12.4±5.1 | −13.0±4.4 |
| Maximum closure deceleration [cm/s$^2$] | | | | | | | |
| | Small | −143.3±63.5 | −134.7±77.9 | −134.2±62.8 | −101.7±43.7 | −105.7±36.8 | −102.0±39.6 |
| | Medium | −147.7±61.0 | −145.3±63.1 | −135.4±59.9 | −105.0±46.0 | −110.8±65.6 | −113.3±42.2 |
| | Large | −133.8±35.9 | −113.8±47.1 | −119.7±44.0 | −108.4±52.8 | −96.9±45.9 | −95.4±39.6 |

537 **Table 2** Outcomes of rm-ANOVAs examining the effects of the Environment (PE, hf-VE),
538 Object size (Small, Medium, Large), and Object distance (Near, Middle, Far) on each
539 movement variable of reach-to-grasp movements

| Variable | Effect | $F$ | df | $\varepsilon^*$ | $p^\dagger$ | $\eta_p^2$ |
|---|---|---|---|---|---|---|
| Closure time [ms] | | | | | | |
| | Environment | 19.24 | 1,12 | | **0.001** | 0.62 |
| | Object size | 3.77 | 2,12 | | **0.038** | 0.24 |
| | Object distance | 11.29 | 2,12 | | **< 0.001** | 0.49 |
| | Environment × Object size | 7.69 | 2,12 | | **0.003** | 0.39 |
| | Environment × Object distance | 1.11 | 2,12 | 0.69 | 0.329 | 0.09 |
| | Object size × Object distance | 0.78 | 4,12 | 0.53 | 0.475 | 0.06 |
| | Environment × Object size × Object distance | 0.76 | 4,12 | | 0.559 | 0.06 |
| Closure distance [cm] | | | | | | |
| | Environment | 15.63 | 1,12 | | **0.002** | 0.57 |
| | Object size | 20.25 | 2,12 | | **0.001** | 0.63 |
| | Object distance | 13.66 | 2,12 | 0.70 | **0.007** | 0.53 |
| | Environment × Object size | 6.77 | 2,12 | | **0.005** | 0.36 |
| | Environment × Object distance | 2.64 | 2,12 | | 0.092 | 0.18 |
| | Object size × Object distance | 0.31 | 4,12 | 0.65 | 0.788 | 0.03 |
| | Environment × Object size × Object distance | 0.66 | 4,12 | | 0.622 | 0.05 |
| Maximum closure velocity [cm/s] | | | | | | |
| | Environment | 1.77 | 1,12 | | 0.208 | 0.13 |
| | Object size | 5.84 | 2,12 | | **0.009** | 0.33 |
| | Object distance | 0.12 | 2,12 | | 0.886 | 0.01 |
| | Environment × Object size | 0.62 | 2,12 | 0.69 | 0.492 | 0.05 |
| | Environment × Object distance | 1.83 | 2,12 | | 0.182 | 0.13 |
| | Object size × Object distance | 1.16 | 2,12 | | 0.343 | 0.09 |
| | Environment × Object size × Object distance | 0.80 | 4,12 | | 0.533 | 0.06 |
| Maximum closure deceleration [cm/s$^2$] | | | | | | |
| | Environment | 8.55 | 1,12 | | **0.015** | 0.42 |
| | Object size | 3.52 | 2,12 | | **0.046** | 0.23 |
| | Object distance | 2.86 | 2,12 | | 0.077 | 0.19 |
| | Environment × Object size | 0.76 | 2,12 | 0.65 | 0.428 | 0.06 |
| | Environment × Object distance | 1.78 | 2,12 | | 0.179 | 0.13 |
| | Object size × Object distance | 1.67 | 2,12 | | 0.173 | 0.12 |
| | Environment × Object size × Object distance | 0.14 | 4,12 | | 0.965 | 0.01 |

*Greenhouse-Geisser correction factor for violation of the assumption of sphericity.

540 †Boldfaced values indicate statistical significance at the two-tailed alpha level of 0.05.

541 **Table 3** Summary of model selection

| Physical environment (PE) | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| Model (i) | $B$ | $A_{CO}$ | $TV_{CO}$ | $TA_{CO}$ | $K$ | Log-Likelihood | QAICc | $\Delta_i$ | $w_i$ |
| 1 | 1.46 | −0.084 | 0.13 | 0.0030 | 6 | −25.56 | 63.9 | 0.00 | 0.9990 |
| 2 | 1.37 | −0.092 | 0.12 | | 5 | −33.89 | 78.3 | 14.43 | 0.0010 |
| 3 | 0.55 | | 0.14 | 0.0034 | 5 | −34.95 | 80.4 | 16.56 | 0.00 |
| 4 | 0.34 | | 0.12 | | 4 | −43.91 | 96.2 | 32.30 | 0.00 |
| 5 | 3.81 | −0.18 | | −0.0042 | 5 | −104.88 | 220.3 | 156.43 | 0.00 |
| 6 | 4.60 | −0.19 | | | 4 | −110.97 | 230.3 | 166.42 | 0.00 |
| 7 | 2.03 | | | −0.0048 | 4 | −116.60 | 241.6 | 177.67 | 0.00 |
| 8 | 2.82 | | | | 3 | −123.00 | 252.2 | 188.33 | 0.00 |
| MAP | 1.45 | −0.084 | 0.13 | 0.0030 | | | | | |
| 2.5% CI | 0.94 | −0.12 | 0.12 | 0.0016 | | | | | |
| 97.5% CI | 1.96 | −0.047 | 0.15 | 0.0044 | | | | | |
| Haptic-free virtual environment (hf-VE) | | | | | | | | | |
| Model (i) | $B$ | $A_{CO}$ | $TV_{CO}$ | $TA_{CO}$ | $K$ | Log-Likelihood | QAICc | $\Delta_i$ | $w_i$ |
| 1 | −0.42 | | 0.19 | 0.0048 | 5 | −85.05 | 124.0 | 0.00 | 0.73 |
| 2 | −0.74 | 0.028 | 0.20 | 0.0050 | 6 | −84.86 | 126.0 | 2.01 | 0.27 |
| 3 | −0.74 | | 0.18 | | 4 | −94.61 | 134.3 | 10.29 | 0.0040 |
| 4 | −0.43 | −0.027 | 0.17 | | 5 | −94.45 | 136.1 | 12.30 | 0.0020 |
| 5 | 7.20 | −0.47 | | −0.0080 | 5 | −180.95 | 249.5 | 125.44 | 0.00 |
| 6 | 8.77 | −0.49 | | | 4 | −187.73 | 560.1 | 132.09 | 0.00 |
| 7 | 2.85 | | | −0.0091 | 4 | −195.24 | 265.9 | 141.92 | 0.00 |
| 8 | 4.40 | | | | 3 | −202.08 | 272.7 | 148.68 | 0.00 |
| MAP | −0.50 | 0.0073 | 0.19 | 0.0048 | | | | | |
| 2.5% CI | −1.31 | −0.056 | 0.18 | 0.0028 | | | | | |
| 97.5% CI | 0.30 | 0.11 | 0.21 | 0.0070 | | | | | |

Parameters include the intercept ($B$); the coefficients for each of the continuous predictors:
aperture at CO ($A_{CO}$), transport velocity at CO ($TV_{CO}$), and transport acceleration at CO ($TA_{CO}$);
the number of free parameters ($K$); Log-Likelihood; the Akaike information criterion corrected
for overdispersion (QAICc); the difference in QAICc between the $i$th model and the best
model ($\Delta_i$); and model weight or the probability that a given model is the best model among all
models ($w_i$). Models are arranged in order from best (lowest $\Delta_i$) to worst (highest $\Delta_i$). The
bottom rows describe the model-averaged parameter estimates (MAP) with lower (2.5%) and
upper (97.5%) bounds of 95% confidence intervals (CI).

**Fig. 1.** Schematic illustration of the experimental setup and procedure. After wearing an
Oculus™ head-mounted display (HMD), the participants sat on a chair in front of the
experimental rig, with their thumb pressing a clipper switch (indicated in yellow). IREDs
markers were attached to the participant's wrist, and the tips of the thumb and index finger. An
auditory cue—a beep—signaled the participant to reach to grasp the object: black (small,
3.6×2.5×8 cm), dark gray (medium, 5.4×2.5×8 cm), and light gray (large, 7.2×2.5×8 cm),
placed at three different distances relative to the clipper switch (near: 24 cm, middle: 30 cm,
and far: 36 cm).

**Fig. 2.** Mean spatial profiles of the reach (left) and grasp (right) components of the reach-to-grasp movements performed in PE and hf-VE for a representative object size-distance pair (medium object, middle distance). (**A**) Transport distance. (**B**) Aperture. (**C**) Transport velocity. (**D**) Aperture velocity. (**E**) Transport acceleration. (**F**) Aperture acceleration. Solid circles in each plot denote closure onset (CO). Shaded areas indicate ±1*SEM* (*n* = 13).

**Fig. 3.** Mean values of closure variables for the reach-to-grasp movements performed in PE and hf-VE. (**A**) Closure time. (**B**) Closure distance. (**C**) Maximum closure velocity. (**D**) Maximum closure deceleration. Error bars indicate $\pm 1 SEM$ ($n$ = 13), and individual points represent the participant means for the near, middle, and far distances (left) and the small, medium, and large sizes. *$p < 0.05$; **$p < 0.005$; ***$p < 0.001$.

**Fig 4.** The relationship between opening/closure time and movement time, and between opening/closure distance and transport distance, in PE and hf-VE. (**A**) Opening/closure time vs. movement time in PE. (**B**) Opening/closure time vs. movement time in hf-VE. (**C**) Opening/ closure distance vs. transport distance in PE. (**D**) Opening/closure distance vs. transport distance in hf-VE.

**Fig. 5.** Relationship between closure distance and each of the three state parameters in PE (red) and hf-VE (blue). (**A**) Closure distance and aperture at CO. (**B**) Closure distance and transport velocity at CO. (**C**) Closure distance and transport acceleration at CO. Left panels show pairwise relationship in PE and hf-VE. Right panels show comparisons of the slope of the regression line between PE and hf-VE using paired samples *t*-test. Horizontal and vertical bars indicate ±1*SEM* (*n* = 13).

Fig. 6. Pairwise relationships among the three state parameters in PE (red) and hf-VE (blue). (A) Transport velocity at CO and aperture at CO. (B) Transport acceleration at CO and aperture at CO. (C) Transport velocity at CO and transport acceleration at CO. Left panels pairwise relationship in PE and hf-VE. Right panels show comparisons of the slope of the regression line between PE and hf-VE using paired samples $t$-test. Horizontal and vertical bars indicate $\pm 1 SEM$ ($n$ = 13).