1   **Title:**   Predictive visual motion extrapolation emerges spontaneously and without
2   supervision at each layer of a hierarchical neural network with spike-timing-
3   dependent plasticity
4

5   **Abbreviated Title:**   Predictive visual motion through spike-based learning
6

7   **Authors:**   Anthony N. Burkitt*[1] & Hinze Hogendoorn[2]
8

Affiliations:

[1]Department of Biomedical Engineering, The University of Melbourne, Melbourne, VIC 3010, Australia

[2]Melbourne School of Psychological Sciences, The University of Melbourne, Melbourne, VIC 3010,
Australia

***Correspondence**: aburkitt@unimelb.edu.au

Number of pages: 35

Number of figures: 9

Number of words for abstract: 220

Number of words for introduction: 929

Number of words for discussion: 2188

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgments: HH acknowledges support from the Australian Research Council's Discovery
Projects funding scheme project DP180102268. ANB acknowledges funding from the Australian
Government, via grant AUSMURIB000001 associated with ONR MURI grant N00014-19-1-2571. We
thank Hamish Meffin and Stefan Bode for helpful comments on the manuscript.

Abstract

The fact that the transmission and processing of visual information in the brain takes time presents a

problem for the accurate real-time localisation of a moving object. One way this problem might be

solved is extrapolation: using an object's past trajectory to predict its location in the present moment.

Here, we investigate how a simulated *in silico* layered neural network might implement such

extrapolation mechanisms, and how the necessary neural circuits might develop. We allowed an

unsupervised hierarchical network of velocity-tuned neurons to learn its connectivity through spike-

timing dependent plasticity. We show that the temporal contingencies between the different neural

populations that are activated by an object as it moves causes the receptive fields of higher-level

neurons to shift in the direction opposite to their preferred direction of motion. The result is that

neural populations spontaneously start to represent moving objects as being further along their

trajectory than where they were physically detected. Due to the inherent delays of neural

transmission, this effectively compensates for (part of) those delays by bringing the represented

position of a moving object closer to its instantaneous position in the world. Finally, we show that this

model accurately predicts the pattern of perceptual mislocalisation that arises when human observers

are required to localise a moving object relative to a flashed static object (the flash-lag effect).

Significance Statement

Our ability to track and respond to rapidly changing visual stimuli, such as a fast moving tennis ball,

indicates that the brain is capable of extrapolating the trajectory of a moving object in order to predict

its current position, despite the delays that result from neural transmission. Here we show how the

neural circuits underlying this ability can be learned through spike-timing dependent synaptic

plasticity, and that these circuits emerge spontaneously and without supervision. This demonstrates

how the neural transmission delays can, in part, be compensated to implement the extrapolation

mechanisms required to predict where a moving object is at the present moment.

1. Introduction

The transmission and processing of information in the nervous system takes time. In the case of visual input to the eyes, for example, it takes up to ~50-70 milliseconds for information from the retina to reach the primary visual cortex (Maunsell & Gibson, 1992; Lamme & Roelfsema, 2000), and up to ~120 milliseconds before human observers are able to initiate the first actions based on that information (Thorpe et al., 1996; Kirchner & Thorpe, 2006). Because events in the world continue to unfold during this time, visual information becomes progressively outdated as it travels up the visual hierarchy.

Although this is inconsequential when visual stimuli are unchanging on this time scale, these delays pose a problem when input is time-varying, for instance in the case of visual motion. If neural delays were not somehow compensated, we would consistently mislocalise moving objects behind their true positions. However, humans and many other visual animals are strikingly accurate at interacting with even fast moving objects (Smeets et al., 1998), suggesting that the brain implements some kind of mechanism to compensate for neural delays.

One candidate mechanism by which the brain might compensate for delays is prediction (Nijhawan, 2008). In the case of motion, the brain might use an object's previous trajectory to extrapolate its current position, even though actual sensory input about the object's current position will not become available for some time. Consistent with this interpretation, motion extrapolation mechanisms have been demonstrated in multiple levels of the visual hierarchy, including the retina of salamanders, mice, and rabbits (Berry et al., 1999; Hosoya et al., 2005; Schwartz et al., 2007), cat lateral geniculate nucleus (Sillito et al., 1994), and both cat and macaque primary visual cortex (Jancke et al., 2004; Subramaniyan et al., 2018; Benvenuti et al., 2019). In humans, recent EEG and MEG studies using apparent motion similarly revealed predictive activation along motion trajectories (Hogendoorn & Burkitt, 2018; Aitken et al., 2020; Blom et al., 2020; Robinson et al., 2020), and motion extrapolation mechanisms have been argued to be the cause of the so-called flash-lag effect (Hogendoorn, 2020; Nijhawan, 1994; Khoei et al., 2017).

The existence of predictive mechanisms at multiple stages of the visual hierarchy is reminiscent of

hierarchical predictive coding, a highly influential model of cortical organisation (Rao & Ballard, 1999).

In this model, multiple layers of a sensory hierarchy send predictions down to lower levels, which in

turn send prediction errors up to higher levels. In this way, the hierarchy essentially infers the

underlying causes of incoming sensory input, using prediction errors to correct and update that

inference. It is important to note, however, that the "predictions" in predictive coding are hierarchical,

rather than temporal: predictive coding networks 'predict' (or reconstruct) activity patterns in other

layers, rather than predicting the future. Consequently, the conventional formulation of predictive

coding cannot compensate for neural delays. In fact, we previously argued that neural delays pose a

specific problem for hierarchical predictive coding, because descending hierarchical predictions will

be misaligned in time with ascending sensory input (Hogendoorn and Burkitt, 2019). For any time-

varying stimulus (such as a moving object) this would lead to significant (and undesirable) prediction

errors.

To address this, we previously proposed a Real-Time Temporal Alignment hypothesis, which extends

the predictive coding framework to account for neural transmission delays (Hogendoorn and Burkitt,

2019). In this hypothesis, both forward and backward connections between hierarchical layers

implement extrapolation mechanisms to compensate for the incremental delay incurred at that

particular step. Without these extrapolation mechanisms, delays progressively accumulate as

information flows through the visual hierarchy, such that information at higher hierarchical layers is

outdated relative to information at lower hierarchical layers. Conversely, the striking consequence of

the Real-Time Temporal Alignment hypothesis is that for a predictable stimulus trajectory, different

layers of the visual hierarchy become aligned in time. The hypothesis posits that extrapolation

mechanisms are implemented at multiple stages of the visual system, which is consistent with the

neurophysiological findings outlined above, as well as with human behavioural experiments (van

Heusden et al., 2019). However, a key question that remains is how such extrapolation mechanisms

are implemented at the circuit level, and how those neural circuits arise during development.

Here, we address those two questions by simulating *in silico* the first two layers of a feedforward

hierarchical neural network sensitive to visual motion. We present the network with simulated moving

objects, and allow neurons to learn their connections through spike-time dependent plasticity (STDP;

Markram et al., 1997; Bi & Poo, 1998), a synaptic learning rule that strengthens and weakens synapses

contingent on the relative timing of input and output action potentials. We focus on the first two

layers of the hierarchical network to explore the key mechanisms, which would be expected to occur

at each higher level of the hierarchy.

We show that when a motion-sensitive hierarchical network is allowed to learn its connectivity

through STDP (without supervision), the temporal contingencies between the different neural

populations that are activated by the object as it moves cause the receptive fields of higher-level

neurons to spontaneously shift in the direction opposite to their preferred direction of motion. As a

result, they start to encode the extrapolated position of a moving object along its trajectory, rather

than its physical position. However, due to the delays inherent in neural transmission, this mechanism

actually brings the represented position of the object closer to its instantaneous position in the world,

effectively compensating for (part of) those delays. Finally, we show that the behaviour of the resulting

network predicts the pattern of velocity-dependence in the perceptual localisation of moving objects.

2. Methods

2.1. Network Architecture

The network architecture considered here is one in which a moving object generates a visual input

stimulus that is encoded at each layer of the network by a population code that represents both the

position and velocity of the stimulus. This population code includes sub-populations of neurons tuned

to both position and velocity, as has previously been proposed by Khoei et al. (2017) and consistent

with the known velocity-tuning of a large proportion of visual neurons in the early visual system

(Orban et al., 1986). This sub-population coding of the velocity of the stimulus at each layer is inherited

from the lower layers, beginning in the retina (Berry et al., 1999; Ravello et al, 2019) and LGN (Sillito

et al., 1994), and then passed on to the primary visual cortex (Janke et al., 2004; Benvenuti et al.,

2020), and it is further enhanced in the motion-processing centers of areas MT and MST (Maunsell &

Van Essen, 1983; Koch et al., 1989; Perrone & Thiele, 2001; Inaba et al., 2011).

Three layers of the network are shown schematically in Figure 1A, in which there are N_n position sub-

populations at each layer. Although both classical predictive coding and the Real-Time Temporal

Alignment hypothesis posit both feed-forward and feedback connections, here we consider only feed-

forward connections as proof-of-principle. In addition, in a more general scheme lateral weights at

each layer could also be included, as previously proposed by Jancke and Erlhagen (Jancke & Erlhagen,

2010), but these are neglected here.

Figure 1: A. Schematic illustration of a portion of the first three layers of the hierarchical network

architecture. The circles denote the neural population at each of the N_n positions (fifteen shown, with

the central population in black). The straight lines indicate the possible non-zero weights connecting

the neural populations between layers, which have a limited spatial spread. The arrows indicate the

direction of the neural connectivity (pre-synaptic to post-synaptic) and only the weights for a subset of

neurons is illustrated. B. Each position population is further divided into M velocity-tuned sub-

populations. Each velocity-tuned sub-population projects to sub-populations in the subsequent layer

with the same velocity-tuning. Note that the analysis in this study focusses upon two adjacent layers.

The neural activation of each stage of processing feeds forward to the following stage. An important

aspect of the network architecture is that each neural population receives input from a limited

receptive field of neural populations at the preceding stage. In this way, the receptive field size of

neural populations increases as the activity propagates to higher stages of processing.

2.2 Neural Model

The Poisson neuron model is used, in which the spikes of each neuron i ($i =1,...,N$) are generated

stochastically with a spiking-rate function, $r_i(t)$, that is time-dependent and described by a Poisson

point process. The instantaneous probability of a spike is given by this Poisson spiking-rate function,

$r_i(t)$, so that in numerical simulations with discrete time-steps of Δt the probability of neuron i firing a

spike at time t is $r_i(t)\Delta t$. This is a stochastic neuron model that is widely used for both analytical and

computational studies (Gerstner et al., 2014 [Ch.7]). A more complete mathematical description and

analysis of this model is given in Appendix A of (Kempter et al., 1998).

The population place code of the stimulus at every stage of processing is described by a set of N_n units

representing overlapping place fields, equally distributed over the interval [0, 1] and each with an

identical Gaussian distribution width σ_p. This width represents the number of *independent* place fields,

N_p, in the input layer, given by $N_p = 1/ \sigma_p$. In this network, the Gaussian distribution represents the

activity evoked by a stimulus, as illustrated in Figure 2, in which the neural activity of each population

of active input neurons corresponding to a particular place field (i.e., an object at a particular location)

is represented by a different colour, and the Gaussian curve represents the amplitude of the firing-

rate of each neuron in that population. The firing rate, which represents the rate of action potentials,

is described by a Poisson process, with a base firing rate of 5Hz in the absence of stimulation. Note

that periodic boundary conditions are used, so that the position code can equivalently be represented

by place on a circle. For simplicity, we consider the situation where only one object activates the input

at any time, so that the relative activations of the input neurons give a neural representation of the

position of the object.

A layered network structure is considered, in which the units at the input layer feed their activity

forward to the following layer, which has the same number of units, N_n. For simplicity, the layers are

taken to have an equal spatial separation and the propagation delay time for activity between layers

has a constant value t_{delay}. This neural transmission delay is of the order of several milliseconds

between layers of the hierarchy (Maunsell & Gibson, 1992).

To incorporate velocity, each place field is further subdivided into M distinct sub-populations,

corresponding to M different velocities (or velocity intervals) for the input stimulus, as illustrated in

Figure 1B. The velocity of the object is encoded by the activity in the corresponding sub-population of

the place fields. Consequently, an object moving at a constant velocity will primarily activate one

velocity sub-population within each place field, i.e., for simplicity a discrete (quantised) representation

Figure 2: Place code in first (input) layer: Each colour represents the neural activity of the population

of neurons, illustrated here for N_p = 32 positions corresponding to the location of the peak of each

Gaussian curve, with width $\sigma_p = 1/N_p$. A stimulus at the position indicated on the plot by Δ generates

firing rates in the neural population, whose amplitude is indicated by the crosses on the corresponding

place curves. The place distribution of population activity for a stimulus centred at the position $x = 0.5$

is shown in bold-red.

of velocity is considered rather than a continuous representation. The velocity is assumed here to have

been encoded by an earlier stage of neural processing, such as in the retina, so that the details of how

this encoding occurs are not incorporated into the model. It suffices that velocity is encoded at each

stage of the processing, which is a reasonable assumption as it is known that velocity is present in

higher stages of the visual pathway such as area MT (Movshon & Newsome, 1996). In addition, the

velocity sub-populations are assumed for simplicity to be precise, with no diffusion in velocity space.

A more complete description of this process would include lateral connections between the neurons

in each layer (both across positions and velocities, as implemented in Khoei et al. (2017) and Khoei et

al. (2013)). However, the dynamics and interactions of these lateral connections are not the focus of

this paper, since we are concerned here with the first wave of feed-forward activity through the

network. Note that the place field at time t is defined as the average marginalized over the velocity

sub-populations: $P(x(t)) = \sum_v P(x(t)|v)$.

2.3 Neural Learning

We require that the computations that underlie learning in the network must be based upon known

principles of synaptic plasticity, namely that the change in a synaptic strength is activity dependent

and local. The locality constraint of synaptic plasticity requires that changes in the synaptic strength

(i.e., the weight of a connection) can only depend on the activity of the presynaptic neuron and the

postsynaptic neuron. Consequently, the spatial distribution of the synaptic changes in response to a

stimulus are confined to the spatial extent of the position representation of the stimulus, which has

important consequences for the structure of the network that emerges as a result of learning.

In the full network the weights are described by a matrix, W between every pair of successive layers

and these are taken to be excitatory, in keeping with the excitatory nature of the long-range pyramidal

neurons in cortex. Since our focus here will be upon the first two layers, W is taken to be a N_nxN_n

matrix in which the elements w_{ji} are the weights between the first two layers. The locality constraint

is implemented in the network by requiring that the weights from a neuron at location i in the first

layer has a probability of being connected to a neuron at location j in the second layer that is Gaussian,

namely that

$$\text{Prob}\{w_{ji} \neq 0\} = G\left(\frac{x_i - y_j}{\sigma_W}\right) \equiv \frac{1}{\sigma_W\sqrt{2\pi}}\exp\left(\frac{(x_i - y_j)^2}{2\sigma_W^2}\right),$$

(1)

where σ_W is the width of the Gaussian distribution, $G(\cdot)$, and x_i and y_j are the locations of the neurons

i and j, in the first- and second-layers resp. The width σ_W is chosen throughout to be sufficiently small

relative to the spatial extent of the network (i.e., $\sigma_W \ll 1$) that it is effectively equivalent to the van

Mises circular distribution. In simulations, the amplitudes of the non-zero weights are initialized

randomly and given small positive initial values, $w_{\text{init}} = 0.01$, while the zero-valued weights are fixed

throughout (i.e., corresponding to the absence of any synaptic connection between the two neurons).

In simulations this Gaussian distribution was truncated to zero at $5\sigma_W$. The non-zero weights then

evolve according to Spike-Timing Dependent Plasticity (STDP), namely a change of the weight, Δw_{ji} ,

will occur when an input spike generated by neuron i at time t_i arrives after a time-delay t_{delay} at a

synapse on neuron j , and an output spike occurs at this neuron at time t_j , so that the time difference

Δt lies within the STDP time-window $F_{STDP}(\Delta t)$

$$\Delta w_{ji} = \rho\, F_{STDP}\left(t_j - t_i - t_{\text{delay}}\right) ,$$

(2)

231   where ρ is the learning rate and

$$F_{\text{STDP}}(\Delta t) = \begin{cases} c_{\text{p}}e^{\Delta t/\tau_{\text{p}}} , & \Delta t > 0 \\ 0 , & \Delta t = 0 \\ -c_{\text{d}}e^{-\Delta t/\tau_{\text{d}}} , & \Delta t < 0 , \end{cases}$$

(3)

with coefficients of potentiation and depression given by c_{p} and c_{d}, and time constants of potentiation

and depression given by τ_{p} and τ_{d} resp., and $\Delta t = \left(t_j - t_i - t_{\text{delay}}\right)$ is the difference between the

spike output of the post-synaptic neuron at time t_j and the incoming spike (generated by neuron i at

time t_i) that arrives at the synapse at time $t_i + t_{\text{delay}}$. Throughout the simulations balanced STDP was

used, i.e., the STDP time-window has an equal amount of potentiation and depression, with $c_p = c_d =$

1 and $\tau_p = \tau_d$ = 20ms, and a value of t_{delay} = 20 ms was chosen. An upper bound on the individual

weights, w_{max}, ensures that the weights do not grow unbounded. Note that the STDP learning

described here is translationally invariant, since each neuron in a layer of the hierarchy receives a

time- and space-shifted version of the input received by other neurons in the layer. Consequently, the

learning is convolutional, because the weights connecting a neuron to neurons in the preceding layer

of the hierarchy evolve to have a translational invariance across the layer.

Consider how an arbitrary weight w_{ij} between unit i in Layer 1 at position x_i and unit j in Layer 2 at

position y_j (chosen here for simplicity to be $y_j = 0$) evolves according to STDP due the motion of an

input stimulus moving from left to right with velocity v. Initially each weight has a small value w_0 and

the distribution of possible weights is given by Eq.(1). Suppose that the stimulus of unit i is given by

$r_i(t)$, then the change in weight Δw_{ij} will depend upon both the timing of the pre-synaptic (input)

spikes, with probability distribution $P(t_i) \propto r_i(t)$, and the timing of the post-synaptic (output) spikes,

with probability distribution $P(t_j)$:

$$\Delta w_{ij} \propto \int_{-\infty}^{\infty} dt_i P(t_i) \int_{-\infty}^{\infty} dt_j P(t_j) F_{\text{STDP}}(t_j - t_i - t_{delay}) . \tag{4}$$

250 Consider now a stimulus that is a very localized point-like object, such as an insect or the spot or light

251 from a laser pointer or a dot in a circular motion on a screen, moving at velocity v. This input can be

252 approximated as a Dirac delta function, $r_i(t) = \delta(t - t_i^*)$, where $t_i^* = \frac{x_i}{v}$. Then the probability

253 distribution $P(t_j)$ becomes

$$P(t_j) \propto w_0 G\left(\frac{x_i - y_j}{\sigma_w}\right) = w_0 G\left(\frac{vt_i}{\sigma_w}\right) = w_0 G\left(\frac{v(t_j - t_{delay})}{\sigma_w}\right) \tag{5}$$

254 where $G(\cdot)$ is the Gaussian distribution given in Eq.(1). Consequently

$$\Delta w_{ij} \propto \int_{-\infty}^{\infty} dt_i r(t_i) \int_{-\infty}^{\infty} dt_j w_0 G\left(\frac{v(t_j - t_{delay})}{\sigma_w}\right) F_{STDP}(t_j - t_i - t_{delay}) \tag{6}$$

255 The time integral can be separated into the potentiation and depression contributions as

$$\Delta w_{ij} \propto \int_{-\infty}^{\infty} dt_i \delta(t_i - t_i^*) w_0 \left[c_P \int_{t_i + t_{delay}}^{\infty} dt_j G \left(\frac{v(t_j - t_{delay})}{\sigma_w} \right) exp \left\{ \frac{(t_j - t_i - t_{delay})}{\tau_P} \right\} \right.$$

$$\left. - c_D \int_{-\infty}^{t_i + t_{delay}} dt_j G \left(\frac{v(t_j - t_{delay})}{\sigma_w} \right) exp \left\{ \frac{(t_j - t_i - t_{delay})}{\tau_P} \right\} \right]$$ (7)

256 Integrating over t_i using the properties of the Dirac delta function gives

$$\Delta w_{ij} \propto w_0 \left[c_P \int_{t_i^* + t_{delay}}^{\infty} dt_j G \left(\frac{v(t_j - t_{delay})}{\sigma_w} \right) exp \left\{ \frac{(t_j - t_i^* - t_{delay})}{\tau_P} \right\} \right.$$

$$\left. - c_D \int_{-\infty}^{t_i^* + t_{delay}} dt_j G \left(\frac{v(t_j - t_{delay})}{\sigma_w} \right) exp \left\{ \frac{(t_j - t_i^* - t_{delay})}{\tau_P} \right\} \right]$$ (8)

257 Assuming STDP with equal potentiation and depression $(\tau = \tau_P = \tau_D$ and $c = c_P = c_D)$ gives

$$\Delta w_{ij} \propto cw_0 \int_0^{\infty} dt e^{-t/\tau} \left[G \left(\frac{vt + x_i}{\sigma_w} \right) - G \left(\frac{vt - x_i}{\sigma_w} \right) \right]$$ (9)

This change in the weight w_{ij} as a function of the position x_i of the input neuron is plotted in Figure 3

for velocities v = 0.05, 0.1, 0.15, 0.2 cycles/sec. The weights on the leading (incoming) side of the

weight distribution experience more potentiation than depression due to STDP, whereas the weights

on the tailing (outgoing) side of the weight distribution experience more depression than potentiation.

This effect becomes larger as the velocity increases and the relative change in weights becomes larger.

For more realistic two-dimensional motion, as occurs with neural activation of the retina, the above

analysis can be extended in a straightforward fashion to two dimensions. Note that this expression

does not depend upon the time delay t_{delay}, which is to be expected since the delay is the same for all

neurons between Layers 1 and 2 and it is only the relative times of the pre-synaptic and post-synaptic

events at the synapses that play a role in STDP. While this expression for Δw_{ij} describes the essence

of the mechanism by which the shift in weight distribution is in the direction opposite to the direction

of motion of the stimulus, there are a number of approximations that will influence the magnitude of

this effect. The above calculation uses the continuum expressions for the weight density, whereas in

Figure 3: Relative change in weights generated by STDP for motion of the point-stimulus to the right.

A) Each plotted line represents the change in weight, Δw, generated by a stimulus with velocity,

v = 0.05 (blue dash-dot), 0.1 (red dotted), 0.15 (cyan dashed), 0.2 (magenta solid) cycles/sec, plotted

relative to the x-coordinate of the input (pre-synaptic) neuron. The $\Delta w = 0$ line is plotted as a black

solid line, with potentiation (pot) indicated for $\Delta w > 0$ and depression (dep) indicated for $\Delta w < 0$.

B) The initial Gaussian probability distribution of weights for one post-synaptic neuron in the upper

layer neuron receiving pre-synaptic inputs from the lower layer (as shown in Fig.2).

reality each neuron has discrete number of synaptic inputs, which introduces a discretization error to

the above calculation. Also, for larger stimuli it is necessary to use the more complete expression,

Eq.(6), as this expression takes account of the variability in spiking associated with stimuli with greater

spatial extent. In addition, this expression provides a description of only the initial phase of learning,

as there are two features of the neural behaviour that make it more difficult to give an analytical

expression for the complete weight evolution. First, the weights have an upper bound (i.e., once a

weight reach this upper bound, it cannot be further potentiated), which introduces a non-linearity

into the weight evolution. Second, the probability of the Level 2 neuron spiking depends upon the

complete distribution of all its input weights, which are themselves constantly changing through STDP.

In the above expression only the initial Gaussian distribution is used, whereas when learning

progresses the distributions will become increasingly non-Gaussian, as described by the receptive

fields observed in the numerical simulations in the Results, Section 3.2.

2.4 Neural Simulations

Simulations of a network were carried out using MATLAB with N_n = 2000 neurons at both the input

layer and the first layer. The input layer had N_p = 32 place-fields, i.e., the width σ_p of the Gaussian

place-fields on the input layer was chosen to be $\sigma_p = 1/N_p$ in order to give place-fields that were both

localized and had a reasonable amount of overlap. The weight distribution width, σ_W, was also chosen

to be $\sigma_W = 1/N_p$, and the width of the resulting place field $\sigma_{RF,j}$ of each neuron in the second layer was

measured from the distribution of weights after learning by spatially binning the weight amplitudes

and finding the width of the resulting histogram, fit to a normal distribution. The width of the place

fields in the second layer are distributed around a value that will shift depending upon the velocity of

the stimulus, so these are labelled as $\sigma_{RF,v}$, to indicate this velocity-dependence. A time-step of 1ms

was used for the simulations, and velocities over the range $v \in [0, 5]$, where the units of velocity are

represented in terms of the inverse time (sec^{-1}) taken to traverse the full spatial range $x \in [0, 1]$.

3. Results

The analysis here focuses upon the first two layers of the network, since the structure and function of

the network follows the same principles at each successive level of the layered network.

3.1. Stationary input stimulus

In order to illustrate the effects of a moving stimulus and to have a baseline for comparison, we

consider first the case with stationary inputs, i.e., in which the stimulus velocity is zero. We use a single

point stimulus with an amplitude 20 times greater than the base firing rate. In this case, there is no

change in stimulus position over time, but rather stationary stimuli are presented for short periods of

time at random positions. When the stimulus activates the input, it will generate activity in the units

at the first layer, as described in Section 2.2 and illustrated in Figure 2. Because this generates constant

input to layer two, and a balanced STDP window is used, convolution by the STDP function would not

be expected to systematically change synaptic weights. Consequently, the network maintains a stable

position code in each layer of the network, namely a localized (Gaussian-like) place field at each layer

that arises through the variance of the STDP learning of the weights (Kempter et al., 1999).

The organisation of the receptive fields of the neural populations in the second layer therefore simply

reflects the input in the first layer, which has a spatial spread σ_p, and the activity transmitted through

the weights, which has a spatial spread of σ_W. Figure 4 shows the results of a simulation for this case,

in which the neural population in the second layer generates a place field representation of the input,

as expected. The weights are initialized with a small value and then evolve under STDP, as described

in the Methods. In this way, the width of the place field distribution at any layer depends upon the

width of the place field at the preceding layer and the spatial spread of the synaptic connections that

connect the two layers.

3.2. Moving input stimulus

We now consider the case where the same point-stimulus is moving. The velocity is chosen to have a

discrete representation (i.e., a discrete number of velocities are chosen for simplicity, rather than a

continuous representation), the place input distribution is as described above by a Gaussian

distribution of width σ_p , as outlined above, and a simulation time step Δt=1ms is used. A stimulus

Figure 4: Place code in the second layer generated by learning: Each colour represents the neural

activity of a population of neurons, illustrated here for a simulation with N_p = 32 position populations

in the second layer. The place distribution for the population centred at the position x = 0.5 is shown in

bold.

moving at a velocity v has a place representation that changes over time so that at a time Δt later it

has shifted a distance $\Delta x = v\,\Delta t$. A moving object will sequentially activate successive populations of

Level 1 neurons, which in turn project to Level 2. Importantly, a neuron in level 2 receiving input from

level 1 neurons driven by this moving object will tend to fire more as the stimulus moves towards the

centre of its place field. Due to STDP, inputs that arrive at the Level 2 neuron relatively early (before

its peak firing rate) will be potentiated, whereas inputs that arrive relatively late are likely to be

depressed. Consequently, the synapses connecting a given Level 2 neuron to Level 1 neurons centred

on the direction from which the stimulus is arriving will tend to be potentiated by STDP. Conversely,

the synapses on the other side (i.e., where the stimulus departs from the place field), will tend to be

depressed, since the inputs on average arrive after the peak in output spiking activity.

We therefore hypothesise that for neural populations tuned to visual motion, the pattern of arrival of

synaptic inputs, together with STDP, will tend to potentiate the synapses in the incoming direction of

the stimulus and depress synapses in the departing direction of the stimulus. This would then lead to

an overall shift of the place field in the direction towards the incoming stimulus. Moreover, due to the

limited temporal window of STDP, the shift in the place fields of the level 2 neurons would be expected

to be larger for larger velocities.

To investigate this hypothesis, we simulated the activity of the neural network when it was presented

with simulated objects moving at a range of velocities, and investigated the evolution of the receptive

fields of Level 2 populations over time. We investigated neural populations tuned to 26 velocities,

from 0 to 5 cycles per second in steps of 0.2 cycles per second (since we used periodic boundaries,

one cycle is equivalent to traversing the full range of positions once). Because of the symmetry in our

neural model, we only considered rightwards velocities, but the network behaves equivalently for

leftward velocities. Each simulation ran for 5 simulated seconds (5,000 timesteps of Δt =1ms). The

simulated object at a single location provided input to the Level 1 neurons according to their

respective place fields.

To evaluate whether receptive fields indeed shifted as a result of learning, we calculated the mean

receptive field of all Level 2 neurons at each timestep by aligning the 32 Level 2 place fields and

averaging their receptive fields at that timestep. This yielded a mean receptive field as a function of

simulation time for each velocity. Figure 5 shows the evolution of receptive field position over time

for 6 evenly-spaced velocities (0-5 cycles/s).

Figure 5: Simulated receptive field position of level 2 neural populations as a function of learning, for 6 different rightward velocities. Top row: In each panel, the colours in each row of pixels show the horizontal receptive field position at a single time-step, with hot colours indicating stronger connections to incoming signals from level 1 populations centred on that position. The center of the receptive field at each timepoint is marked by a black line. For clarity, only the first 2500 ms of the simulation are shown. Bottom row: mean receptive fields before (blue) and after learning (red). Altogether, these plots reveal that when velocity-tuned neural populations are presented with their preferred stimulus (here rightward), STDP causes their receptive field to shift over time in the direction opposite to their preferred velocity (i.e., here leftward).

3.3. Velocity-dependence

To be able to directly compare how the evolution of level 2 receptive fields depended on velocity, for each velocity we fitted Gaussians to the average receptive field at each time-step (e.g., each row in each panel in Figure 6). We then repeated the entire simulation 15 times to reduce the impact of stochastic noise. Subsequently, we averaged the horizontal centre of the best-fit Gaussians across all 15 iterations. Finally, we plotted this receptive field centre as a function of time, separately for each velocity. The result is illustrated in Figure 6.

Figure 6: The evolution of receptive field centre over time, as a function of velocity. Colours represent

populations tuned to 26 equally-spaced velocities, ranging from zero (red) to 5 cycles/s (blue). For all

non-zero velocities, receptive fields shifted against the direction of motion, with a clear monotonic

relationship between velocity and the asymptotic magnitude of the shift.

We observed two key features. Firstly, the initial rate at which the receptive field centres of the

different velocity-tuned populations shifted increased with velocity. At zero velocity the receptive field

centre stayed in the same position, and as velocity increased, the initial rate of change grew until it

reached an asymptote. This is perhaps unsurprising, because at lower velocities the simulated object

needs more time to traverse the receptive fields of a given number of neurons. As a result, the object

drives fewer individual neurons, and in turn provides fewer opportunities for the network to learn.

Furthermore, the neural populations tuned to different velocities differed not only in their initial rate

of change, but also in the asymptote of that change. In other words, the spatial shift in receptive field

position at which subsequent time-steps produced no further net change in position increased with

increasing velocity. This observation is significant because the asymptote represents the position of

the receptive field after learning has effectively completed, and therefore reflects the stable situation

in visual systems that have had even a short history of exposure to moving stimuli. The rate at which

the asymptote is approached depends upon the STDP learning rate. the Because these velocity-tuned

populations are subpopulations of an overall population coding for position (as illustrated

schematically in Figure 2), the overall population effectively represents a moving object ahead of

where a physically-aligned static stimulus is represented. As a consequence, we might expect the

asymptotic receptive field shift to be similarly reflected in conscious perception as the instantaneous

perceived position of a moving object.

3.4 Behavioural predictions

Our model reveals how STDP-induced shifts in receptive field position depend on velocity. In the final

section of this paper, we evaluate the degree to which these predictions match observed

dependencies on velocities in the localisation of moving objects by healthy human observers.

A much-studied behavioural paradigm used to probe the instantaneous perceived position of a moving

object is the flash-lag paradigm, in which a flashed object is briefly presented alongside a moving

object (Nijhawan, 1994). Observers are then required to report where they perceived the moving

object to be at the moment the flashed object was presented. Strikingly, in this paradigm observers

consistently localise the moving object ahead of the physically aligned flashed object, a phenomenon

known as the Flash-Lag Effect (FLE) (Nijhawan, 1994). Although the mechanisms underlying this effect

have been hotly debated over the last 25 years, convergent evidence supports Nijhawan's original

proposal that it reflects some kind of neural motion extrapolation process (Hogendoorn, 2020;

Nijhawan, 1994). What is particularly relevant to the present context is that the effect has been

observed to scale with velocity (Wojtach et al., 2008): when an object moves faster, its perceived

position at any given instant lies further along the object's trajectory.

In our model, the perceived position of a moving object corresponds to the level 2 neural population

activated by that object. As outlined in the previous section, this is determined by the asymptotic

receptive field position after learning. As a measure of asymptotic receptive field shift after learning,

we averaged the receptive field shift in the final 100 ms of our simulation (e.g., the 100 rightmost

datapoints for each curve in Figure 6), averaged across the 15 iterations of the simulation. We

subsequently fitted a logarithmic function to the data, as has previously been done for behavioural

estimates of perceived position shifts using the FLE (Wojtach et al., 2008). This function explained a

total of 96.8% of the variance, showing that the dependence of final receptive field position on velocity

was very well described by a logarithmic relationship (Figure 7).

Figure 7: Velocity-dependence of asymptotic receptive field shifts. Marker colours correspond to

velocities in Figure 6, and the dashed line represents a logarithmic fit to the data, explaining 96.8% of

variance.

In order to compare the perceptual shifts predicted by our model to those measured in behavioural

experiments with human observers, we compared the velocity dependence of receptive field shifts in

our model to the velocity dependence of the FLE, as previously measured for the full range of

detectable velocities by (Wojtach et al., 2008). We noted that the magnitudes of both RF-shifts in our

model and perceptual shifts in the FLE were very well-described by a logarithmic dependence on

stimulus velocity (Figure 7, Figure 8A). We then directly compared RF-shifts in our model to perceptual

shifts in the FLE by treating the maximum velocities tested in each paradigm to be equal. For the

behavioural paradigm, this was 50 deg/s, the highest velocity at which an FLE could be measured

(Wojtach et al., 2008). For our model, this was 5 cycles/second, at which point the period of the motion

(200 ms) reached the approximate width of the STDP window. The correlation between RF-shifts in

our model and perceived position shifts in the FLE was near perfect (r > 0.99). Note that this pattern

Figure 8: A) Behaviourally measured perceptual shift magnitude as a function of velocity using the

flash-lag effect (reproduced from (Wojtach et al., 2008). The solid line indicates a logarithmic fit to

their data. B) Scatterplot showing the magnitude of receptive field shifts predicted by our model and

the corresponding perceptual shifts reported by (Wojtach et al., 2008) using the FLE. The

correspondence is near-perfect, with a Pearson correlation r > 0.99. C) Absolute magnitudes of RF-

shifts produced by our model and position shifts observed in the FLE, expressed as a time-constant (the

equivalent duration necessary to traverse the shift distance at each velocity). Dashed lines indicate

exponential fits.

of results arose spontaneously as a result of STPD, without requiring any tuning of the model. This

shows that the velocity-dependence of STDP-induced receptive-field shifts in our model very closely

matches the velocity-dependence of perceptual mislocalisation for moving objects as measured using

the FLE.

Finally, we compared the absolute magnitude of the shifts in receptive field position produced by our

model to the absolute magnitude of perceptual shifts observed in the FLE. To do so, we expressed the

magnitude of the shift at each velocity as a time constant, by dividing shift magnitude by velocity. This

is equivalent to the time necessary for an object moving at that velocity to be displaced a distance

equivalent to the receptive field shift (Figure 8C). We observed that for both the FLE and our model,

this time constant tended to decrease exponentially with increasing velocity (exponential fits

explained 96.3% and 98.5% of variance in FLE and model time constants, respectively). Across the

entire range of velocities tested, the time constant produced by our model was roughly 12-20% of the

time constant for the behaviourally-measured FLE, as might be expected given that our model

reflected receptive field shifts in just a single layer of synaptic connections.

3.5 Parameter dependence

Parameters in our model were chosen to be biologically plausible. For some parameters, choosing

different values would be expected to have predictable effects. For example, varying the STDP learning

rate ρ , Eq.(2), would be expected to cause the model to converge to its asymptotic state either more

rapidly or more slowly. However, we would not expect it to change the asymptotic state itself – merely

the simulation time necessary to reach that state. Indeed, we deliberately chose a relatively high

learning rate in order to keep the computation tractable; we would not expect a biological system to

reach its asymptotic state within just 5 seconds of exposure.

For other parameters, it is less obvious how choosing different values would affect the pattern of

results. In particular, we chose a value of 32 for N_p, the number of place fields in each layer. This

parameter corresponds loosely to the size of receptive fields at each layer, and might be expected to

vary for neurons in different areas in retinotopic visual cortex. For example, place field width inevitably

varies as a function of eccentricity, with foveal retinotopic areas showing smaller receptive fields than

peripheral retinotopic areas. To investigate the effect of manipulating this parameter, we ran

additional simulations with higher (64) and lower (16) values of N_p. Although we observed small

differences in the absolute magnitude of predicted receptive field shifts, the overall pattern of results

was very similar (Figure 9). In particular, the pattern of velocity dependence for both absolute

receptive field shifts and the equivalent time constants was highly similar, giving confidence that our

results are not restricted to a small region of parameter space.

Figure 9: Comparison of model prediction for different numbers of place fields. A) Receptive field shift

as a function of velocity for models with 16, 32, or 64 place fields. The magnitude of the receptive field

shift increases slightly with increasing place field width (corresponding to a lower number of place

fields), but the overall pattern of velocity-dependence is comparable for all three simulations. Dashed

lines show logarithmic fits as in the main analysis. B) Comparison of time constants for models with

16, 32, or 64 place fields. Time constants were remarkably similar for models with different numbers

of place fields, showing comparable exponential dependence on velocity. Dashed lines show

exponential fits.

4. Discussion

We investigated a computational problem faced by the brain in processing the visual position of moving objects: the fact that neural transmission takes time, and that the brain therefore only has access to outdated visual information. Motion extrapolation is one way the brain might compensate for these delays: by extrapolating the position of moving stimuli along their trajectory, their perceived position would be closer to the their physical position in the world (Hogendoorn, 2020; Nijhawan, 1994, 2008; Hubbard, 2005). We simulated a possible neural mechanism (spike-timing dependent plasticity - STDP) by which a layered neural network might implement such an extrapolation mechanism. We show that a two-layer hierarchical network comprised of velocity-tuned neural population is not only able to implement motion extrapolation, but actually learns to do so spontaneously without supervision, due only to the temporal contingencies of its connectivity. We go on to show that the velocity-dependence of the resulting receptive field shifts predicts previously reported, behaviourally measured effects on the perceived position of moving objects.

The magnitude of the receptive field shifts we observe for each velocity in our simulations corresponds roughly to the equivalent displacement resulting from 10-20 ms of motion at that velocity. Although this is 5-8 times smaller than the FLE, it is important to note that our model analysed here includes just two layers, and only one stage at which learning takes place. If we were to extend our model to include additional layers, each with comparable properties, then each output layer would constitute the input layer for the synapses at the next stage. As a result, we would expect the same learning process, and therefore the same receptive field shift, to take place at each stage. In this way, receptive field shifts would add up as information ascends the hierarchy. Although it is unknown which cortical areas ultimately determine where we consciously perceive a moving object, it seems highly likely that information from the retina will cross at least a handful of synapses before it is accessed for conscious awareness. The magnitude of the receptive field shifts predicted by our model are therefore of roughly

the same order of magnitude as, and comparable to, those we would expect based on the magnitude

of the perceptual effect.

It is interesting to note that the flash-lag effect is just one of several related motion-position illusions

in which the position of a moving object is biased by motion (e.g., Eagleman & Sejnowski, 2007). In

the Fröhlich effect (Kirschfeld & Kammer, 1999) for example, a moving object suddenly appears, and

the perceived initial position of the object is mislocalised in the direction of motion. The pattern of

receptive fields shifts that we observe in our model can also qualitatively explain the Fröhlich effect.

Subtle differences with the FLE paradigm (such as the likely transient neuronal onset response to the

initial appearance of the moving object in the Fröhlich effect) are not currently captured by our model,

but a direct comparison of predictions for these (and other) illusions could be informative to further

develop the model.

The magnitude of receptive field shifts predicted by our model is also consistent with previous

neurophysiological recordings as well as human neuroimaging. Jancke and colleagues (Jancke et al.,

2004) recorded neurons in cat primary visual cortex, and compared the latencies of responses to

flashes with the latencies of responses to smoothly moving objects. They observed that peak neural

responses to smoothly moving objects were approximately 16 ms further along the motion trajectory

than peak responses to static flashed objects. Almost identical results were found by Subramaniyan

and colleagues (Subramaniyan et al., 2018) who recorded neurons in primary visual cortex of awake

macaques, and observed a latency advantage for moving stimuli compared to flashed stimuli of

between 10-20 ms depending on stimulus velocity. These results from invasive recordings in cats and

macaques are therefore in quantitative agreement with the predictions of our model. In humans, we

recently used an EEG decoding paradigm to investigate the latency of neural responses to predictably

and unpredictably moving objects (Hogendoorn & Burkitt, 2018). Using an apparent motion paradigm,

we showed that when objects move along predictable trajectories, their position is represented with

a lower latency than when they move along unpredictable trajectories. Like the neurophysiology

studies, we observed a latency of 16 ms for the predictably moving object. Our present modelling

result is therefore consistent not only with behavioural measurements of motion perception, but also

with neural recordings in both humans and animals.

The underlying mechanism is the same property of STDP that causes neurons to tune to the earliest

spikes (Song, Miller & Abbott, 2000; Guyonneau, Van Rullen & Thorpe, 2005), which has diverse

manifestations in the brain's neural circuits, including in the context of phase precession in the

hippocampus (Mehta, Quirk & Wilson, 2000) and of localizing a repeating spatio-temporal spike

pattern embedded in a noisy spike train (Masquelier, Guyonneau & Thorpe, 2008). Essentially STDP

leads to potentiation of the weights associated with the leading edge of the receptive field as the

receptive field is entered, and depression of weights on the tailing edge as the motion moves out of

the neuron's receptive field as illustrated in Figure 3. Our study extends the understanding of this

mechanism by examining the velocity dependence of this effect. It is important to note that in our

model, the extrapolation mechanism emerged spontaneously and without supervision, simply as a

result of STDP. By extension, extrapolation would similarly be expected to develop spontaneously in

any hierarchical network of velocity-selective populations when it is exposed to visual motion.

Furthermore, it would be expected to arise between every layer in such a network. This structure of

extrapolation mechanisms at multiple levels of the visual hierarchy is consistent with previous

empirical findings showing extrapolation at both monocular and binocular stages of processing (van

Heusden et al., 2019). It is also consistent with the Real Time Temporal Alignment hypothesis that we

recently proposed (Hogendoorn and Burkitt, 2019) as a theoretical extension of classical predictive

coding (Rao & Ballard, 1999), although this hypothesis also posited feedback projections that are not

included in our current model. Indeed, the network architecture considered here is entirely feed-

forward, which represents a good model for the initial wave of neural activity travelling through the

visual pathway in response to a stimulus, but it neglects the feedback activity from higher visual

centres. This descending feedback activity, which occurs and can persist over a longer timeframe than

the initial wave of neural activity, may play an important role in the understanding of temporal

processing in the brain on these longer timescales. These questions, which lie outside the scope of the

present study, form the focus of ongoing research.

The mechanism here differs from that proposed by Lim & Choe (2006), who show that STDP, together

with facilitating synapses, can provide a neural basis for understanding the orientation flash-lag effect,

a visual illusion involving the perceived misalignment of a rotating bar, which is located between two

aligned flanking bars that briefly flash when the rotating bar is aligned. Importantly, their model

architecture is very different, involving a bilaterally ring-connected network of orientation-tuned

neurons, in which the lateral connections between these neurons are trained using a combination of

STDP and activity-dependent facilitation, rather than the feedforward connections examined in our

study. They also did not examine the velocity dependence of the effect.

The network and learning parameter values chosen for this proof-of-concept study represent values

consistent with cortical neural processing, but without incorporating many of the details of vision

processing in the human visual pathway. The description of the neural activity in terms of a Poisson

process is a widely used approximation for the time distribution of action potentials. Although it

neglects all spike after-effects, such as refractoriness, it nevertheless provides a good description for

the situation examined here in which the visual stimulus moves with a constant velocity and is

modelled as having a spatial intensity distribution that is Gaussian, without any edges or other spatial

discontinuities (Aviel & Gerstner, 2006). The STDP time constants of potentiation and depression, τ_p

and τ_d , are chosen to both have a value of 20ms, which is in the range of that observed in neurons in

the visual cortex (Froemke & Dan, 2002).

In the primate visual system, the processing of motion begins in the retina, where it has long been

known that there are direction-selective retinal ganglion cells (Barlow & Hill, 1963; Barlow et al., 1964).

These neurons are maximally activated by motion in their preferred direction and strongly suppressed

by *motion in the opposite direction.* Within the retina there are a number of mechanisms involving

multilayered retinal circuits that provide reliable motion detection (Frechette et al., 2005; Manookin

et al., 2018), consistent with the broad experimental evidence for velocity tuning neurons in the retina

of mammals and vertebrates (Olveczky et al., 2007; Vaney et al., 2012; Ravello et al., 2019; for a review

see Wei, 2018). This motion selective information is transmitted via the LGN to the primary visual

cortex, V1, where direction selective neurons are concentrated in layer 4B of V1 and project from

there to higher motion processing areas of the visual hierarchy pathway, particularly area MT

(Maunsell & van Essen, 1983).

In the analysis presented here we have made the simplifying assumption that the same principles

apply at each successive level of a layered network. However, in the visual system the receptive fields

of neurons at successive levels become progressively larger as information moves up the visual

hierarchy. For example, a motion selective neuron in area MT which has a receptive field of 10^0

diameter receives its input from neurons in V1 that have receptive fields of 1^0 diameter (Andersen,

1997). The restricted receptive field of neurons in the lower stages of this vision processing hierarchy

can result in ambiguous motion signals as a result of the aperture effect. Consequently, at each

successive stage of the visual hierarchy the motion information is not only transmitted, but it also can

be refined: the information at earlier stages is integrated so that the motion of larger objects can be

more accurately determined and objects moving at different speeds are disambiguated – for a review

see Bradley & Goyal (2008). The larger receptive field sizes in the higher stages of the hierarchy

correspond to broader place field representations, which it would be straightforward to accommodate

in a multi-layer extension of the processing framework presented here. It may also be possible that

the dorsal and ventral pathways of the visual system (i.e., the where and the what pathways) have

very different encoding of velocity. While the dorsal pathway relies upon an accurate representation

of position and the visual motion extrapolation analysed here, it is possible that in the ventral pathway

the velocity coding is so broadly tuned that it is effectively absent.

In the analysis presented here we have for convenience used a discrete coding of the velocity, rather

than allowing it to take a continuum of values from zero up to some maximal value. Consequently, an

object with changing velocity will, in this simplified model, make discrete jumps between velocity sub-

populations. It is, however, also possible to formulate the velocity using such a continuous

representation, for example as a set of overlapping Gaussian distributed velocity fields similar to the

(spatial) place fields. We anticipate that this would give a smoother, possibly more biologically

plausible, transition between velocity sub-populations, but that it would not change the essential

results of this study in any significant way.

In sum, we have implemented spike-time dependent plasticity in a layered network of velocity-

selective neurons, and shown that this results in a pattern of receptive-field shifts that causes the

network to effectively extrapolate the position of a moving object along its trajectory. The magnitude

of this shift is in quantitative agreement with previous findings from both animal neurophysiology and

human neuroimaging experiments, and also qualitatively predicts the perceptual mislocalisation of

moving objects in the well-known FLE. Most strikingly, we show that it emerges spontaneously and

without supervision, suggesting that extrapolation mechanisms are likely to arise in many locations

and at many levels in the visual system.

Finally, the model we present here includes only feed-forward connections, and a natural extension

to the model would be to include lateral and/or feedback connection. Previous modelling work, most

notably by Jancke and Erlhagen (Jancke & Erlhagen, 2010), has proposed an instrumental role for

lateral connections in generating the perceptual mislocalisation that characterises the FLE. It would

be interesting to investigate in more detail what the emergent characteristics would be of a network

implementing both STDP and lateral connectivity, and whether that would explain any other

perceptual phenomenology. In a similar vein, it would be interesting to implement feedback

connections in the model we present here, as we proposed in our previous Real-Time Temporal

Alignment hypothesis (Hogendoorn and Burkitt, 2019). An exciting possibility is that these feedback

connections might function to calibrate receptive field shifts to the relative transmission delays

between layers in the hierarchy, allowing extrapolation mechanisms to accurately compensate for

processing delays. Further research will be necessary to evaluate these possible extensions to the

current model.

6. References

Aitken F, Turner G, Kok P (2020) Prior expectations of motion direction modulate early sensory

processing. bioRxiv:2020.03.06.980672.

Andersen RA (1997) Neural Mechanisms of Visual Motion Perception in Primates. Neuron 18:865–

872.

Aviel Y, Gerstner W (2006) From spiking neurons to rate models: A cascade model as an

approximation to spiking neuron models with refractoriness. Phys Rev E 73:051908.

Barlow HB, Hill RM (1963) Selective Sensitivity to Direction of Movement in Ganglion Cells of the

Rabbit Retina. Science (80-) 139:412–412.

Barlow HB, Hill RM, Levick WR (1964) Retinal ganglion cells responding selectively to direction and

speed of image motion in the rabbit. J Physiol 173:377–407.

Benvenuti G, Chemla S, Boonman A, Perrinet L, Masson G, Chavane F (2020) Anticipatory responses

along motion trajectories in awake monkey area V1. bioRxiv:2020.03.26.010017.

Berry MJ, Brivanlou IH, Jordan TA, Meister M (1999) Anticipation of moving stimuli by the retina.

Nature 398:334–338.

Bi GQ, Poo MM (1998) Synaptic modifications in cultured hippocampal neurons: Dependence on

spike timing, synaptic strength, and postsynaptic cell type. J Neurosci 18:10464–10472.

Blom T, Feuerriegel D, Johnson P, Bode S, Hogendoorn H (2020) Predictions drive neural

representations of visual events ahead of incoming sensory information. Proc Natl Acad Sci

117:7510–7515.

Bradley DC, Goyal MS (2008) Velocity computation in the primate visual system. Nat Rev Neurosci

9:686–695.

Eagleman DM, Sejnowski TJ (2007) Motion signals bias localization judgments: A unified explanation

for the flash-lag, flash-drag, flash-jump, and Frohlich illusions, J Vision 7:1–12.

Frechette E.S, Sher A, Grivich MI, Petrusca D, Litke AM, Chichilnisky EJ (2005) Fidelity of the

ensemble code for visual motion in primate retina, J Neurophysiol 94:119–135.

Froemke RC, Dan Y (2002) Spike-timing-dependent synaptic modification induced by natural spike

trains. Nature 416:433–438.

Gerstner W, Kistler WM, Naud R, Paninski L (2014) Neuronal Dynamics: From Single Neurons to

Networks and Models of Cognition, Cambridge University Press, UK.

Guyonneau R, VanRullen R, Thorpe SJ (2005) Neurons Tune to the Earliest Spikes Through STDP,

Neural Comput 17:859–879.

Hogendoorn H (2020) Motion Extrapolation in Visual Processing: the Resolution to 25 Years of Flash-

Lag Debate. J Neurosci 40:5698–5705.

Hogendoorn H, Burkitt AN (2018) Predictive coding of visual object position ahead of moving objects

revealed by time-resolved EEG decoding. Neuroimage 171:55–61.

Hogendoorn H, Burkitt AN (2019) Predictive Coding with Neural Transmission Delays: A Real-Time

Temporal Alignment Hypothesis. eNeuro 6:e0412-18.2019.

Hosoya T, Baccus SA, Meister M (2005) Dynamic predictive coding by the retina. Nature 436:71–77.

Hubbard TL (2005) Representational momentum and related displacements in spatial memory: A

review of the findings. Psychon Bull Rev 12:822–851.

Inaba N, Miura K, Kawano K (2011) Direction and speed tuning to visual motion in cortical areas MT

and MSTd during smooth pursuit eye movements, J Neurophysiol.105:1531–1545.

Jancke D, Erlhagen W (2010) Bridging the gap: a model of common neural mechanisms underlying

the Fröhlich effect, the flash-lag effect, and the representational momentum effect. In: Space

and Time in Perception and Action (Nijhawan R, Khurana B, eds), pp 422–440. Cambridge:

Cambridge University Press.

Jancke D, Erlhagen W, Schöner G, Dinse HR (2004) Shorter latencies for motion trajectories than for

flashes in population responses of cat primary visual cortex. J Physiol 556:971–982.

Kempter R, Gerstner W, van Hemmen JL, Wagner H (1998) Extracting Oscillations: Neuronal

Coincidence Detection with Noisy Periodic Spike Input. Neural Comput 10:1987–2017.

Khoei MA, Masson GS, Perrinet LU (2013) Motion-based prediction explains the role of tracking in

motion extrapolation. J Physiol Paris 107: 409–420.

Khoei MA, Masson GS, Perrinet LU (2017) The Flash-Lag Effect as a Motion-Based Predictive Shift.

PLOS Comput Biol 13:e1005068.

Koch C, Wang HT, Mathur B (1989) Computing motion in the primate's visual system, J Exp Biol

146:115-139.

Kirchner H, Thorpe SJ (2006) Ultra-rapid object detection with saccadic eye movements: Visual

processing speed revisited. Vision Res 46:1762–1776.

Kirschfeld K, Kammer T (1999) The Fröhlich effect: a consequence of the interaction of visual focal

attention and metacontrast. Vis Res 39:3702 – 3709.

Lamme VAF, Roelfsema PR (2000) The distinct modes of vision offered by feedforward and recurrent

processing. Trends Neurosci 23:571–579.

Lim H, Choe Y (2006) Delay Compensation Through Facilitating Synapses and STDP: A Neural Basis

for Orientation Flash-Lag Effect. Proc. 2006 IEEE Int. Joint Conf. Neural Networks, pp. 4269-

4276.

Manookin MB, Patterson SS, Linehan CM (2018) Neural mechanisms mediating motion sensitivity in

parasol ganglion cells of the primate retina. Neuron 97:1327–1340.

Markram H, Lübke J, Frotscher M, Sakmann B (1997) Regulation of synaptic efficacy by coincidence

of postsynaptic APs and EPSPs. Science 275:213–215.

Masquelier T, Guyonneau R, Thorpe SJ (2008) Spike Timing Dependent Plasticity Finds the Start of

Repeating Patterns in Continuous Spike Trains, PLoS ONE 3(1):e1377.

Maunsell J, van Essen D (1983) The connections of the middle temporal visual area (MT) and their

relationship to a cortical hierarchy in the macaque monkey. J Neurosci 3:2563–2586.

Maunsell JH, Gibson JR (1992) Visual response latencies in striate cortex of the macaque monkey. J

Neurophysiol 68:1332–1344.

Mehta MR, Quirk MC, Wilson MA (2000) Experience-Dependent Asymmetric Shape of Hippocampal

Receptive Fields, Neuron 25:707–715.

Movshon JA, Newsome WT (1996) Visual response properties of striate cortical neurons projecting

to area MT in macaque monkeys. J Neurosci 16:7733–7741.

Nijhawan R (1994) Motion extrapolation in catching. Nature 370:256–257.

Nijhawan R (2008) Visual prediction: Psychophysics and neurophysiology of compensation for time

delays. Behav Brain Sci 31:179–239.

Olveczky BP, Baccus SA, Meister M (2007) Retinal adaptation to object motion. Neuron 56:689–700.

Orban GA, Kennedy H, Bullier J (1986) Velocity sensitivity and direction selectivity of neurons in

areas V1 and V2 of the monkey: influence of eccentricity. J Neurophysiol 56:462–480.

Perrone JA, Thiele A (2001) Speed skills: measuring the visual speed analyzing properties of primate

MT neurons, Nature Neurosci 4:526-532.

Rao RPN, Ballard DH (1999) Predictive coding in the visual cortex: A functional interpretation of

some extra-classical receptive-field effects. Nat Neurosci 2:79–87.

Ravello CR, Perrinet LU, Escobar M-J, Palacios AG (2019) Speed-Selectivity in Retinal Ganglion Cells is

Sharpened by Broad Spatial Frequency, Naturalistic Stimuli. Sci Rep 9:456.

Robinson AK, Grootswagers T, Shatek SM, Gerboni J, Holcombe AO, Carlson TA (2020) Now you see

it, now you don't: Overlapping neural representations for the position of visible and invisible

objects. bioRxiv:2020.03.02.974162.

Schwartz G, Taylor S, Fisher C, Harris R, Berry MJ (2007) Synchronized Firing among Retinal Ganglion

Cells Signals Motion Reversal. Neuron 55:958–969.

Sillito AM, Jones HE, Gerstein GL, West DC (1994) Feature-linked synchronization of thalamic relay

cell firing induced by feedback from the visual cortex. Nature 369:479–482.

Smeets JBJ, Brenner E, De Lussanet MHE (1998) Visuomotor delays when hitting running spiders. In:

EWEP 5 - Advances in perception-action coupling (B B, A L, G D, A R-B, eds), pp 36–40. Paris:

Editions {EDK}.

Song S, Miller KD, Abbott LF (2000) Competitive Hebbian learning through spike-timing-dependent

synaptic plasticity, Nat Neurosci 3:919-926.

Subramaniyan M, Ecker AS, Patel SS, Cotton RJ, Bethge M, Pitkow X, Berens P, Tolias AS (2018) Faster

processing of moving compared with flashed bars in awake macaque V1 provides a neural

correlate of the flash lag illusion. J Neurophysiol 120:2430–2452.

Thorpe S, Fize D, Marlot C (1996) Speed of processing in the human visual system. Nature 381:520–

522.

van Heusden E, Harris AM, Garrido MI, Hogendoorn H (2019) Predictive coding of visual motion in

both monocular and binocular human visual processing. J Vis 19:3.

Vaney DI, Sivyer B, Taylor WR (2012) Direction selectivity in the retina: symmetry and asymmetry in

structure and function. Nature Rev Neurosci 13:194-208.

Wei W (2018) Neural Mechanisms of Motion Processing in the Mammalian Retina. Annu Rev Vis Sci

4:165–192.

Wojtach WT, Sung K, Truong S, Purves D (2008) An empirical explanation of the flash-lag effect. Proc

Natl Acad Sci 105:16338–16343.