1

3   **Emergence of Non-Linear Mixed Selectivity in Prefrontal Cortex after Training**

5   Wenhao Dang*, Russell J. Jaffe*, Xue-Lian Qi, Christos Constantinidis

6   Department of Neurobiology & Anatomy, Wake Forest School of Medicine,

7   Winston-Salem, NC 27157, USA

8   * These authors contributed equally to this work

10   Abbreviated Title: Mixed Selectivity in PFC

Address correspondence to:
Christos Constantinidis, PhD.
Department of Neurobiology and Anatomy,
Wake Forest School of Medicine
Medical Center Blvd, Winston Salem, NC, 27157
E-mail: cconstan@wakehealth.edu

18
19
Number of pages: 46
Number of figures: 10
Number of words for Abstract: 221
Number of words for Significance: 117
Number of words for Introduction: 478
Number of words for Discussion: 1457
26
27   The authors report no conflicts of interest during the research conducted.
28
29

ACKNOWLEDGMENTS
Research reported in this paper was supported by the National Eye Institute of the
National Institutes of Health under award number R01 EY017077. We wish to
acknowledge Kathini Palaninathan and Austin Lodish for technical help; Junda Zhu, and
Sihai Li for helpful comments on the manuscript.

ABSTRACT

Neurons in the prefrontal cortex (PFC) are typically activated by different cognitive

tasks, and also by different stimuli and abstract variables within these tasks. A single

neuron's selectivity for a given stimulus dimension often changes depending on its

context, a phenomenon known as nonlinear mixed selectivity (NMS). It has previously

been hypothesized that NMS emerges as a result of training to perform tasks in different

contexts. We tested this hypothesis directly by examining the neuronal responses of

different PFC areas before and after monkeys were trained to perform different working

memory tasks involving visual stimulus locations and/or shapes. We found that training

induces a modest increase in the proportion of PFC neurons with NMS exclusively for

spatial working memory, but not shape working memory tasks, with area 9/46

undergoing the most significant increase in NMS cell proportion. We also found that

increased working memory task complexity, in the form of simultaneously storing

location and shape combinations, does not increase the degree of NMS for stimulus shape

with other task variables. Lastly, in contrast to the previous studies, we did not find

evidence that NMS is predictive of task performance. Our results thus provide critical

insights on the representation of stimuli and task information in neuronal populations,

which may pave the way to a greater understanding of neural selectivity in working

memory.

SIGNIFICANCE STATEMENT

How multiple types of information are represented in working memory remains a

complex computational problem. It has been hypothesized that nonlinear mixed

selectivity allows neurons to efficiently encode multiple stimuli in different contexts,

after subjects have been trained in complex tasks. Our analysis of prefrontal recordings

obtained before and after training monkeys to perform working memory tasks only

partially agreed with this prediction, in that nonlinear mixed selectivity emerged for

spatial but not shape information, and mostly in mid-dorsal PFC. Nonlinear mixed

selectivity also displayed little modulation across either task complexity or correct

performance. These results point to other mechanisms, in addition to nonlinear mixed

selectivity, to represent complex information about stimulus and context in neuronal

activity.

INTRODUCTION

Working memory (WM) is broadly defined as the ability to encode, maintain, and

manipulate information in the conscious mind over a period of seconds without the

presence of any sensory inputs. As a core component of complex cognitive abilities such

as planning and reasoning, the true importance of WM ultimately depends on whether it

can maintain and manipulate task relevant information in a task relevant manner

(Baddeley, 2012). Multiple variables, including external sensory inputs and internal task

requirements, must be encoded in order to achieve the level of adaptability in WM that is

necessary for complex tasks. The mechanisms that underlie this encoding process across

time and neuronal population is one of the most important questions in current WM

research.

When individuals are required to maintain objects in their WM, neurons from a

network of brain regions may exhibit selective and sustained increases or decreases in

their activity in order to represent the remembered objects through these unique patterns

of activity (Constantinidis and Procyk, 2004). The prefrontal cortex (PFC) plays a leading

role in this network, and by extension, in the use of WM (Riley and Constantinidis,

2016). For example, when the PFC is damaged or degraded, whether through trauma,

illness, or experimental lesions, performance in WM tasks seems to decrease dramatically

(Curtis and D'Esposito, 2004; Morris and Baddeley, 1988; Rossi et al., 2007).

Individual PFC neurons typically encode more than one variables, and the exact

variables encoded are task dependent (Asaad et al., 2000; Machens et al., 2010; Mansouri

et al., 2006; Qi et al., 2015; Warden and Miller, 2010). More interestingly, a portion of

neurons exhibit nonlinear mixed selectivity (NMS) for different variables, which means

that their response to the combination of variables cannot be predicted by the linear

summation of their responses to single variables (Johnston et al., 2020; Parthasarathy et

al., 2017; Rigotti et al., 2013). Theoretical studies have shown that NMS is useful for

linear readouts of flexible, arbitrary combinations of variables (Buonomano and Maass,

2009; Fusi et al., 2016; Rigotti et al., 2010), and may also control the trade-off between

discrimination and generalization (Barak et al., 2013; Johnston et al., 2020).

Despite the proposed importance of NMS on theoretical grounds, some

experimental studies have failed to detect neurons with NMS (Cavanagh et al., 2018). It

is therefore possible that NMS may manifest exclusively in a limited set of PFC

subdivisions or alternatively, that NMS emerges exclusively after training to perform

specific types of cognitive tasks. Moreover, the implications of NMS on other aspects of

neural encoding, such as code stability, have not yet been investigated. We were therefore

motivated to analyze and compare neural data from rhesus macaque monkeys before and

after training. Here we report results of NMS as a function of task training, performance

of different types of working memory tasks, and correct and error trials, across different

prefrontal areas.

METHODS

<u>Animals:</u> Data obtained from six male rhesus monkeys (*Macaca mulatta*), age 5–9 years

old, weighing 5–12 kg, as previously documented (Riley et al., 2018), were analyzed in

this study. None of the animals had any prior experimentation experience at the onset of

our study. Monkeys were either single-housed or pair-housed in communal rooms with

sensory interactions with other monkeys. All experimental procedures followed

guidelines set by the U.S. Public Health Service Policy on Humane Care and Use of

Laboratory Animals and the National Research Council's Guide for the Care and Use of

Laboratory Animals and were reviewed and approved by the Wake Forest University

Institutional Animal Care and Use Committee.

<u>Experimental setup:</u> Monkeys sat with their head fixed in a primate chair while viewing a

monitor positioned 68 cm away from their eyes with dim ambient illumination. Animals

were required to fixate on a 0.2° white square appearing in the center of the screen.

During each trial, the animals maintained fixation on the square while visual stimuli were

presented either at a peripheral location or over the fovea in order to receive a juice

reward. Any break of fixation immediately terminated the trial and no reward was given.

Eye position was monitored throughout the trial using a non-invasive, infrared eye

position scanning system (model RK-716; ISCAN, Burlington, MA). The system

achieved a $< 0.3°$ resolution around the center of vision. Eye position was sampled at 240

Hz, digitized and recorded. Visual stimuli display, monitoring of eye position, and the

synchronization of stimuli with neurophysiological data were performed with in-house

software implemented on the MATLAB environment (Mathworks, Natick, MA), and

utilizing the Psychophysics Toolbox (Meyer and Constantinidis, 2005).

Pre-training task: Following a brief period of fixation training and acclimation to the

stimuli, monkeys were required to fixate on a center position while stimuli were

displayed on the screen. The monkeys were rewarded for maintaining fixation during the

trial with a liquid reward (fruit juice). The stimuli shown in the pre-training passive

spatial task were white 2° squares, presented in one of nine possible locations arranged in

a 3 × 3 grid with 10° distance between adjacent stimuli. The stimuli shown in the pre-

training passive feature task were white 2° circles, diamonds, H-letters, hashtags, plus

signs, squares, triangles, or inverted Y-letters, also presented in one of nine possible

locations arranged in a 3 × 3 grid with 10° distance between adjacent stimuli.

144   Presentation began with a fixation interval of 1 s where only the fixation point

was displayed, followed by a 500 ms of stimulus presentation (referred to hereafter as

cue), followed by a 1.5 s "delay" interval (referred to hereafter as delay1) where, again,

only the fixation point was displayed. A second stimulus (referred to hereafter as sample)

was subsequently shown for 500 ms. In the spatial task, this second stimulus would be

either identical in location to the initial stimulus, or diametrically opposite the first

stimulus. In the feature task, this second stimulus would always be identical in location to

the initial stimulus and would either be an identical shape or the corresponding non-

match shape (each shape was paired with one non-match shape).

In both the spatial and feature task, this second stimulus display was followed by another

"delay" period (referred to hereafter as delay2) of 1.5 s where only the fixation point was

displayed. The location and identity of stimuli was of no behavioral relevance to the

monkeys during the "pre-training" phase, as fixation was the only necessary action for

obtaining reward.

Post-training task: Four of the six monkeys were trained to complete active spatial,

feature and conjunction WM tasks. These active spatial and feature tasks were identical

to the passive spatial and feature tasks that were applied during the "pre-training" phase,

except that these tasks now required the monkeys to remember the spatial location or the

shape feature of the first presented stimulus, and report whether the second stimulus

matched the spatial location or shape feature of the first stimulus, respectively, via

saccading to one of two target stimuli (green for match, blue for non-match). Each target

stimulus could appear at one of two locations orthogonal to the cue/sample stimuli,

pseudo-randomized in each trial.

The conjunction task combined the active spatial and feature tasks, and the stimuli

shown were a white 2° circle, diamond, H-letter, hashtag, plus sign, square, triangle, or

inverted Y-letter shapes, presented in one of nine possible locations arranged in a 3 × 3

grid with 10° distance between adjacent stimuli. Each trial consisted of a fixation interval

of 1 s where only the fixation point was displayed, followed by 500 ms of the first

stimulus presentation, followed by a 1.5 s delay interval where, again, only the fixation

point was displayed. A second stimulus was subsequently shown for 500 ms, and after a

second, 1.5 s delay period the monkeys would report whether the second stimulus

matched both the spatial location and shape feature of the first stimulus, via saccading to

one of the two target stimuli. The conjunction task was therefore the most complex task,

as the monkeys were required to simultaneously store two different items—location and

shape—in their WM.

<u>Surgery and neurophysiology:</u> A 20 mm diameter craniotomy was performed over the

PFC and a recording cylinder was implanted over the site. The location of the cylinder

was visualized through anatomical magnetic resonance imaging (MRI) and stereotaxic

coordinates post-surgery. For two of the four monkeys the recording cylinder was moved

after an initial round of recordings in the post-training phase to sample an additional

surface of the PFC.

<u>Anatomical localization</u>. Each monkey underwent an MRI scan prior to

neurophysiological recordings. Electrode penetrations were mapped onto the cortical

surface. We identified 6 lateral PFC regions: a posterior-dorsal region that included area

8A, a mid-dorsal region that included area 8B and area 9/ 46, an anterior-dorsal region

that included area 9 and area 46, a posterior-ventral region that included area 45, an

anterior-ventral region that included area 47/12, and a frontopolar region that included

area 10. However, the frontopolar region was not sampled sufficiently to be included in

the present analyses.

In addition to comparisons between brain areas segmented in this fashion, other analyses

were performed to account for the position of each neuron along the AP axis. For the

purposes of our analysis, we defined the AP axis as the line connecting the genu of the

arcuate sulcus to the frontal pole. The recording coordinates of each neuron were

projected onto this line, with position expressed as a proportion of the line's length.

Neuronal recordings: Neural recordings were carried out in areas 8, 9, 9/46, 45, 46, and

47/12 of the PFC both before and after training in each WM task. Subsets of the data

presented here were previously used to determine the collective properties of neurons in

the dorsal and ventral PFC, as well as the properties of neurons before and after training

in the posterior-dorsal, mid-dorsal, anterior-dorsal, posterior-ventral, and anterior-ventral

PFC subdivisions. Extracellular recordings were performed with multiple

microelectrodes that were either glass- or epoxylite-coated tungsten, with a 250 μm

diameter and 1–4 MΩ impedance at 1 kHz (Alpha-Omega Engineering, Nazareth, Israel).

A Microdrive system (EPS drive, Alpha- Omega Engineering) advanced arrays of up to

8-microelectrodes, spaced 0.2–1.5 mm apart, through the dura and into the PFC. The

signal from each electrode was amplified and band-pass filtered between 500 Hz and 8

kHz while being recorded with a modular data acquisition system (APM system, FHC,

Bowdoin, ME). Waveforms that exceeded a user-defined threshold were sampled at 25 μs

resolution, digitized, and stored for off-line analysis. Neurons were sampled in an

unbiased fashion, collecting data from all units isolated from our electrodes, with no

regard to the response properties of the isolated neurons. A semi-automated cluster

analysis relied on the KlustaKwik algorithm, which applied principal component analysis

of the waveforms, to sort recorded spike waveforms into separate units. To ensure a

stable firing rate in the analyzed recordings, we identified recordings in which a

significant effect of trial sequence was evident at the baseline firing rate (ANOVA, p <

0.05), e.g., due to a neuron disappearing or appearing during a run, as we were collecting

data from multiple electrodes. Data from these sessions were truncated so that analysis

was only performed on a range of trials with stable firing rate. Less than 10% of neurons

were corrected in this way. Identical data collection procedures, recording equipment,

and spike sorting algorithms were used before and after training in order to prevent any

analytical confounds.

Data analysis: Data analysis was implemented with the MATLAB computational

environment (Mathworks, Natick, MA), with additional statistic tests implemented

through Originlab (OriginLab Corporation, Northampton, MA) and StatsDirect

(StatsDirect Ltd. England). Peristimulus time histograms (PSTHs) for illustrations were

calculated through the moving window average method with a Gaussian window that had

a 200 ms standard deviation, with the shaded area indicating two times standard error

cross trials. For all tasks, only cells with at least 12 correct trials for each cue-sample

location/shape pairs were included in the analysis. To classify neurons of the spatial task

into different categories of selectivity, we performed two-way ANOVAs on the spike

count between either the stimuli location x matching status in for the trial, or between

stimuli location x task epoch (first or second stimulus presentation). Neurons with classic

selectivity (CS) exhibited a main effect of only one factor without significant interaction

term. Neurons with linear mixed selectivity (LMS) exhibited main effects of both factors

without significant interaction term. Neurons with NMS exhibited a significant

interactions term. Finally, non-selective (NS) neurons exhibited no significant term for

both the main effects and the interaction term. Similarly, the two factors for feature task

ANOVA analysis were stimuli shape x matching status, and stimuli shape x task epoch

for the trial.

A method based on singular value decomposition (SVD) and cross validation was

used to calculate population dimensionality for the spatial and feature tasks (Ahlheim and

Love, 2018). The dimensionality of a matrix is defined as its number of non-zero singular

values, identified by SVD. Under this condition however, the noise in the recorded neural

data could potentially inflate the number of non-zero singular values, even if the true

dimensionality were low. Reconstruction from components with cross-validation could

be used to estimate dimensionality with noise, under the assumption that only true

underlying dimensionality can contribute to the reconstruction performance in the cross-

validation dataset. In short, data from j trials were randomly assigned to training the

dataset with j-2 trials, using one of the remaining trials for validation and the other for

testing. SVD was then applied to the averaged training data to obtain all of the possible

low-dimensional reconstructions, which were then correlated through a validation run.

The dimensionality that produced the highest average correlation across j-1 runs was

selected as dimensionality estimate k for this fold, and a k-dimensional reconstruction

was correlated with the held-out test data, resulting in the final reconstruction correlation.

A similar method had also been recently used to estimate the dimensionality over time for

neural data (Cueva et al., 2020). The dimensionality of the sample and delay2 period in

the spatial and feature task was calculated on 50 resamples of a 200-cell pseudo-

population in the corresponding datasets.

Only PFC areas with more than 50 cells in both pre- and post-training time points

were included in the subdivision mixed selectivity comparison analysis. Thus, for the

feature task, only data from the mid-dorsal, posterior-dorsal and posterior-ventral PFC

were analyzed, while the spatial task analyzed data from the mid-dorsal, posterior-dorsal,

posterior-ventral, anterior-dorsal and anterior-ventral PFC.

Neural data from tasks that applied the exact same visual stimuli were used to

compare mixed selectivity between feature/spatial and conjunction tasks. For example, to

compare the feature and the conjunction tasks, we started by selecting a subset of

conjunction trials, in which both visual stimuli appeared at the same location as the

corresponding feature task trials. Then, since all eight shapes were used in a single

recording session for the feature task, a subset of trials, in which same shape pairs were

used as the corresponding conjunction task, could be chosen as the feature dataset. Our

prior methods of ANOVA analyses could thus be applied for comparison across these

datasets.

For comparing mixed selectivity in success and error trials from the spatial task,

we first examined two task variables—the stimuli location and matching status. In this

analysis, we utilized neurons that had at least 3 match and non-match trials for both the

correct and error dataset, in at least 3 stimuli locations. The number of minimum trials

and stimuli locations were chosen to maximize the average trial number for each cell

included into the analysis, while still retaining a sufficiently large sample (i.e. >150

cells). The same number of trials from each stimuli location were randomly chosen in the

correct and error dataset. This randomized trial selection process was repeated 50 times in

order to make the best use of the uneven number of available trials in two datasets. We

also analyzed factors of stimuli location and task epoch. For this analysis, we used neural

data from match trials only, and from neurons that had at least 4 correct and error trials,

in at least 4 stimuli locations.

For decoding analysis, spiking responses from 1 second before cue onset to 5

seconds after cue onset were first binned using a 400 ms wide window and 100 ms steps

to create a spike count vector with a length of 57 elements. A pseudo-population was then

constructed using the spike count vectors from all the available neurons of all the

available animals, thus resulting in a dataset with 96 trials, as if they were recorded

simultaneously. The population response matrix was z-score normalized before being

used to train the decoder. A linear Support Vector Machine (SVM) decoding algorithm

was implemented using the MATLAB fitcecoc function to decode stimuli location,

stimuli shape, or the match/non-match status of trials. A 10-fold cross validation method

was used to estimate the decoder performance and 20 random samplings were

implemented to calculate a 95% confidence interval. For the location and feature task, the

decoding baseline for sensory information was 12.5%, since there were 8 different

options, and 50% for the matching status, since there were only 2 different options. In the

pre-training vs post-training decoding analysis (Fig. 8), linear (CS and LMS) and

nonlinear (NMS) neurons are first defined by their pre and post training responses in the

sample or delay2 period. Each classified population was then applied to decode sensory

information (location and shape) and matching status. A randomization test was used to

determine the time points at which decoding performance was significantly different

between different selectivity categories. In short, we constructed the null distribution by

randomly reassigning the cell selectivity labels under comparison, and re-computing the

maximum absolute difference across all time points of the data in each iteration. This

procedure was repeated for 5000 times. A difference was deemed to be significant if the

true response difference occurred at the extremes of this null distribution (p <0.05, two-

tailed). Since every point in the null distribution is the maximum of all time points, this

method already corrected for the multiple comparisons.

Only informative neurons (CS, LMS, and NMS neurons) in the delay2 period

were used for the cross temporal decoding analysis (Fig. 7), since we wanted to explore

the decoding dynamics during the delay period. The linear SVM decoder was trained on

individual time points and thus had 57 linear decision boundaries. The same dataset was

then classified by every decision boundary in the vector to produce a 57x57 matrix—a

process that was repeated 20 times in order to eliminate the noise. The decoding

performance matrix for each condition was normalized individually to highlight the

coding dynamics rather than absolute performance.

Data availability: All relevant data and code will be available from the

corresponding author on reasonable request. Matlab decoder code for figure 8 and 9 is

available at https://github.com/dwhzlh87/mixed-selectivity

RESULTS

Extracellular neurophysiological recordings were collected from the lateral PFC of six

monkeys before and after they were trained to perform the match/nonmatch WM tasks

(Meyer et al., 2011; Riley et al., 2018). The task required them to view two stimuli

appearing in sequence, with delay periods intervening between them, and to report

whether or not the second stimulus was identical to the first. The two stimuli could differ

in terms of their location (spatial task, Fig. 1A), shape (feature task, Fig. 1B), or both

(conjunction task, Fig. 1C). If the second stimulus matched with the first, monkeys would

saccade towards a green target during a subsequent interval, otherwise to a blue target at

a diametrical location. A total of 1617 cells from six monkeys and 1495 cells from five

monkeys were recorded while the animals were performing the passive spatial and

feature tasks, respectively, which were mutually dubbed "pre-training." A total of 1104

cells from three monkeys and 1116 cells from two monkeys were collected while the

animals were performing the active spatial and feature tasks, respectively, which were

mutually dubbed "post-training". We also collected neural data from 247 neurons for the

passive spatial task from two monkeys after they were trained in the active spatial task.

An additional 975 cells from two monkeys were collected while they were performing

the active "post-training" conjunction task.

Types of selectivity in individual neuronal responses

In our tasks, the context of a given stimulus depends upon the task interval and sequence

in which it is presented. We first considered how selectivity for stimulus location and

shape in the spatial, feature and conjunction WM tasks may vary when the same sample

stimulus appears as a match (it is preceded by a cue at the same location/shape) or a

nonmatch (i.e. is preceded by a cue stimulus of a different location/shape). The neuronal

firing rate is therefore a function of the stimulus location/shape (eight shapes, eight

locations arranged on a 3x3 grid with 10 degrees distance between stimuli, excluding the

center location) and whether this sample stimulus matched the cue stimulus. We used a 2-

way ANOVA with the factors of stimulus location/shape and match/nonmatch status to

classify neurons into four categories of selectivity. CS neurons exhibited a significant

main effect to only one of the factors (stimulus identity or matching status) and had no

significant interaction term. In Fig. 2, the first exemplar displays a cell selective

exclusively for the location of the stimuli, which does not respond differently regardless

of whether the stimulus appeared as a match or nonmatch. The second exemplar of Fig. 2

displays a cell not selective for the location of the stimuli but demonstrates higher mean

response when the stimulus appeared as a non-match. LMS neurons exhibited a

significant main effect for both factors but had no significant interaction term. The third

exemplar of Fig. 2 displays a neuron demonstrating a higher mean response when stimuli

appear as non-match, while simultaneously displaying the same rank order preference for

location. NMS neurons exhibited a significant interaction effect, as shown in the last

exemplar in Fig. 2, a neuron demonstrating different selectivity pattern for locations

under match vs. non-match conditions. Finally, NS indicated the neurons with no

selectivity to any factors or their interaction under consideration. These analyses were

performed using the firing rate recorded during the stimulus presentation period, and

again, using the firing rate recorded during the delay period that followed it.

A second type of NMS was identified in terms of selectivity for stimulus

sequence, that is, whether the same stimulus appeared first (cue) or second (sample). To

avoid the confound of the match or nonmatch status of the second stimulus, we relied

exclusively on match stimuli. This form of NMS was also evaluated through a 2-way

ANOVA model, identifying CS, LMS, NMS, and NS neurons in terms of how the

neurons represented the exact same stimulus when it appeared as a cue and as a match

stimulus.

Effects of training on NMS

When we used the factors of stimulus location/shape and match/nonmatch status for our

two-way ANOVA, we found that training in the spatial WM task increased the proportion

of NMS cells in both the sample period and the delay period that followed the sample

(sample period: pre-training proportion=6.2%, post-training proportion=12.3%, two-

sample proportion test, z=5.31, p=1.13×10^{-7}; delay2 period: pre-training proportion=

2.8%, post-training proportion=6.2%, two-sample proportion test, z=4.62, p=4.86×10^{-5}).

However, this increase in selectivity was not exclusive to NMS cells. The proportion of

CS cells also increased in the delay period following the sample (pre-training proportion=

10.6%, post-training proportion=14.8%, two-sample proportion test, z=3.19, p=0.0014).

The increase in NMS cells was not evident for all types of training. When we

looked at the proportion of change across the pre-training and post-training feature task,

we only found an increase of proportion for CS cells (sample period: pre-training

proportion= 12.0%, post-training proportion=15.7%, two-sample proportion test, z=2.65,

p=0.0081; delay2 period: pre-training proportion= 9.0%, post-training proportion=22.6%,

two-sample proportion test, z=9.37, p=0). No significant increase in the proportion of

NMS cells was observed (sample period: pre-training proportion=5.8%, post-training

proportion=6.7%, two-sample proportion test, z=1.01, p=0.314; delay2 period: pre-

training proportion= 4.2%, post-training proportion=4.6%, two-sample proportion test,

z=0.522, p=0.602) (Fig. 3 A,B).

Similar results were observed when we used the factors of stimulus location/shape

and task epoch (cue vs. match) for the two-way ANOVA instead (Fig. 3 C,D). For the

spatial task, training increased the proportion of NMS cells, at least in the delay period

(stimulus period: pre-training proportion= 5.7%, post-training proportion=7.4%, two-

sample proportion test, z=1.78, p=0.075; delay period: pre-training proportion= 2.1%,

post-training proportion=6.2%, two-sample proportion test, z=5.03, p=4.94×10^{-7}). A

similar increase was observed for the CS cells (stimulus period: pre-training proportion=

21.3%, post-training proportion=25.3%, two-sample proportion test, z=2.37, p=0.018;

delay period: pre-training proportion= 24.2%, post-training proportion=31.1%, two-

sample proportion test, z=3.93, p=8.53×10^{-5}). In the feature task, once again, only the

proportion of CS cells changed (stimulus period: pre-training proportion= 21.9%, post-

training proportion=32.6%, two-sample proportion test, z=6.05, p=1.45×10^{-9}; delay

period: pre-training proportion= 27.6%, post-training proportion=41%, two-sample

proportion test, z=7.16, p=8.32×10^{-13}). The proportion of NMS cells with an effect in the

stimulus period remained relatively unchanged for the cue/match period (pre-training

proportion= 3.4%, post-training proportion=4.7%, two-sample proportion test, z=1.59,

p=0.112), as well as the delay period (pre-training proportion= 2.4%, post-training

proportion=3.8%, two-sample proportion test, z=1.95, p=0.051).

To further validate our proportional measure for NMS and compare our results to

previous research on NMS in the PFC, we plotted the F scores for the interaction term

(i.e. stimulus identity × matching status) in both the spatial and the feature task (Fig. 4

A). We found that this measure of NMS for individual cells increased specifically for the

spatial task, indicated by much higher F score values after training for the spatial task.

We also measured the dimensionality of population responses in the sample and delay2

period for the spatial and feature task. Again, this analysis confirmed the results of our

cell proportion measure (Fig. 4 B). For the spatial task, there was a significant increase of

dimensionality after training (sample period: pre-training dimensionality= 5.72, post-

training dimensionality=10.33, two-sample t test, $t(98)=12.21$, $p=2.18×10^{-21}$; sample

period: pre-training dimensionality= 3.25, post-training dimensionality=6.29, two-sample

t test, $t(98)=9.39$, $p=2.51×10^{-15}$). For the feature task however, no significance increase

was observed in mean F score (Fig. 4A) or dimensionality (Fig 4B; sample period: pre-

training dimensionality= 2.72, post-training dimensionality=2.73, two-sample t test,

$t(98)=0.027$, $p=0.978$; sample period: pre-training dimensionality= 2.43, post-training

dimensionality=2.11, two-sample t test, $t(98)=3.49$, $p=7.29×10^{-4}$).

Regional localization of NMS

To assess whether specific sub-regions of the PFC may be specialized for NMS, we

divided the lateral PFC into six regions (Fig. 5 A) and analyzed the respective

neurophysiological data from five of these regions in order to determine the different

areas' proportional contributions to the observed changes in NMS. We examined NMS

defined by location/shape and match/nonmatch status in the sample period and ultimately

found that the mid-dorsal subdivision underwent the greatest proportional change in

NMS cells for the spatial task after training (Fig. 5 C), without a comparable increase in

the proportion of CS neurons (Mid-dorsal: CS 21.7% pre-training to 19.0% post- training,

NMS 8.2% pre-training to 21.8% post-training). For the feature task however, the

proportional change in NMS cells was relatively small, with moderate increases in CS

and LMS observed in all three analyzed areas (Fig. 5 B).

NMS in task context

Previous theoretical studies linked NMS with more flexible readouts of multiple task

variables, thus leading to the hypothesis that task complexity may modulate NMS. To test

this hypothesis, we compared the neural responses to different shapes at the same

location when the stimuli appeared as match or nonmatch in the conjunction task, to the

same neurons' responses to the same stimuli when they appeared in the feature task. In

the conjunction task, animals needed to simultaneously remember both location and

shape of visual stimuli, while in the feature task, they were only required to remember

shape. Although the hypothesis predicted that the conjunction task would result in greater

NMS than the feature task when the sensory stimuli was the same, this was not what we

observed. No significant differences were observed for either CS cells (feature task

sample proportion= 11.9%, conjunction task sample proportion=9.9%, exact matched

pair sample proportion test ,F=1.229, p=0.197, feature task delay2 proportion= 10.8%,

conjunction task delay2 proportion=11.6%, exact matched pair delay2 proportion test

,F=1.069, p=0.681) or NMS cells (feature task sample proportion= 4.1%, conjunction

task sample proportion=4.4%, exact matched pair sample proportion test, F=1.031,

p=0.901, feature task delay2 proportion= 4.6%, conjunction task delay2

proportion=5.9%, exact matched pair delay2 proportion test, F=1.278, p=0.266) (Fig. 6

A) in the sample period or the delay2 period that followed. We also examined changes in

individual cells' selectivity across the feature and conjunction tasks. Although this

analysis was limited by the relatively low proportion of NMS cells in both tasks, we

found an unstable mapping between tasks in the selectivity categories for both CS and

NMS cells (Fig. 6 B), evidenced by the observation that most cells in CS and NMS

category in the feature task changed their selectivity category in the conjunction task.

There is a possibility, however, that the majority of the informative cells were simply due

to chance (p=0.05 was used as threshold for detecting significant terms), which is

enforced by the observation that NMS cells with larger degree of interaction in one task

tend to also fall into the NMS category in the other task (Fig. 6 C). No significant

difference was observed in the proportion of NMS cells when we performed a similar

comparison between the spatial and conjunction task with a relatively small sample size

(Fig. S1).

The comparison of the naïve and trained conditions allowed us to test the overall

incidence of NMS in different populations of PFC neurons, sampled randomly before and

after training, which was carried out over the course of several months. If NMS were

critical for the representation of task-relevant information, we would expect a difference

in the observed proportion of neurons with NMS, when animals are passively viewing

stimuli vs. when they are actively performing the task and storing representations of the

stimuli in their WM. We therefore applied a two-way ANOVA to compare the neural

responses of neurons between the active and passive spatial tasks after the monkeys had

been trained to perform the active spatial task. We ultimately observed an increase in the

proportion of cells that coded matching status during the sample period, as well as an

increase in the proportion of cells coding sensory information in the delay1 period when

the animal was prompted to report the matching decision. However, the observed

increase in the proportion of NMS cells was not significant (passive proportion= 9.3%,

active proportion=11.7%, exact matched pair proportion test, F=1.385, p=0.362) (Fig. 7

A). Interestingly, a large proportion of cells changed their selectivity category across

tasks, especially for CS cells (Fig. 7 B), and the degree of NMS does not seem to be

predictive of whether a given neuron would fall in the same selectivity category in both

tasks (Fig. 7 C).

Information encoding by NMS neurons

It is known that training leads to increased incorporation of task relevant information in

neural populations, with relatively little change to stimulus information (confirmed in our

dataset, Fig. S2). However, the relative contribution by NMS is not clear. To quantify the

amount of task relevant information contained in linear (CS and LMS) and NMS cells,

we used a linear SVM decoder to decode sensory information (location and shape) and

match/or nonmatch status information. Since the cell selectivity category could be

defined by their response in either sample or delay2 period, we randomly selected equal

numbers of linear and nonlinear cells in both task epochs for each comparison. The

random selecting process was repeated multiple times to obtain a confidence interval. We

ultimately found that linear and nonlinear cells contain comparable amounts of linearly

decodable information in regard to both sensory information and task relevant

information. The only observed difference between the decodable information in the

linear and nonlinear cells occurred in the post training feature task, where linear cells

were observed to contain more stimulus information in the sample period.

We also applied cross temporal decoding to compare classic and linear mixed

cells in regard to population coding dynamics during the delay period. If information

were represented by a stable pattern of activity, the classifier trained at one timepoint

would be expected to work equally effectively at other time points where the information

is present. Conversely, if information were represented by dynamic patterns of activity,

then the decision boundary at one time point would not contribute to decoding

information at other time points. The most prominent result from this analysis is that

NMS cells produced significantly more stable code for matching spatial task information,

compared to CS and LMS cells, as indicated by higher performance off the diagonal

during the delay2 period (Fig. 9).

NMS in correct and error trials

The presence of decodable information in the PFC does not necessarily imply the

presence of information in the conscious mind, and the representation of task relevant

information is ultimately revealed by the ability to conduct the task successfully. In order

to decipher the role of NMS, we examined the F score of the main effects and their

interaction in the ANOVA test in correct vs. error trials for the spatial task (Fig. 10),

which displayed higher NMS levels than the feature or conjunction tasks. Similar to the

pre- vs. post training comparisons, we examined two types of mixed selectivity: stimulus

location vs. matching status and stimulus identity vs. task epoch. The number of trials

and task variables were matched for each cell to avoid confounds in the comparison. The

mean F score in correct trials for the location variable in the stimulus epochs for the

location × epoch comparison was equal to 1.86, while for the error trials the mean was

2.59 (paired t test, t(147)=3.38, p=9.42 × 10^{-4}). The effect extended into the delay

epochs, where the average F score for location in correct trials was 1.43, and that of error

trials was 1.79 (paired t test, t(150)=2.61, p=0.010). However, we did not find any

differences in the F score for the interaction terms in any comparison. The results indicate

that NMS may not be necessary in representing task relevant information in WM.

DISCUSSION

Selectivity for different types of information is critical in representing the plethora of
stimuli and task contexts that can be maintained in WM. NMS is thought to be critical in
that respect, as it allows efficient representation of flexible, arbitrary combinations of
variables (Barak et al., 2013; Buonomano and Maass, 2009; Fusi et al., 2016; Johnston et
al., 2020; Rigotti et al., 2010). Consistent with this idea, increased dimensionality in
NMS has been highlighted as a potential means of increasing the efficiency of WM task
performance (Johnston et al., 2020; Rigotti et al., 2013) and dimensional collapse
characterizes task errors (Rigotti et al., 2013). Moreover, all task relevant information
could be decoded from NMS neurons alone, despite their relative scarcity, with decoder
accuracy actually increasing as the task became more complex (Rigotti et al., 2013).
NMS is assumed to emerge with training in complex tasks that combine multiple types of
information, or in multiple tasks, even without an explicit requirement to combine such
information (Johnston et al., 2020; Lindsay et al., 2017). However, this idea has not been
tested experimentally until now. Our study, by virtue of analyzing neural recordings
before and after training in a series of cognitive tasks, directly tested these postulates. We
found that NMS resulted in a modest increase with training, but only for some tasks, and
furthermore, task complexity was not a predictor of NMS emergence. A causal
relationship between success and dimensionality—and by extension, NMS—was also not
supported by our results, as we did not observe any significant changes in NMS between
error and success trials. These insights refine and qualify the role NMS plays in WM, and
identify a number of open questions.

Effects of training on neural responses

WM is considerably plastic and at least some aspects of it, such as mental processing

speed and the ability to multitask, can be improved with training (Bherer et al., 2008;

Dux et al., 2009; Jaeggi et al., 2008; Klingberg et al., 2005; Klingberg et al., 2002). WM

training has been proven particularly beneficial for clinical populations, e.g. in the case of

traumatic brain injury, attention deficit hyperactivity disorder (ADHD), and

schizophrenia (Klingberg et al., 2002; Subramaniam et al., 2012; Westerberg et al.,

2007). However, the verdict of whether WM training confers tangible benefits on normal

adults and whether these benefits transfer to untrained domains, remains a matter of

heated debate. (Constantinidis and Klingberg, 2016; Cortese et al., 2015; Fukuda et al.,

2010; Owen et al., 2010; Peijnenborgh et al., 2015; Schwaighofer et al., 2015).

This malleability of cognitive performance is thought to be mediated by the

underlying plasticity in neural responses, most importantly within the PFC

(Constantinidis and Klingberg, 2016). In a series of prior studies, we have investigated

changes in PFC responsiveness and selectivity (Meyer et al., 2011; Meyers et al., 2012;

Qi et al., 2011; Riley et al., 2018), as well as other aspects of neuronal discharges such as

trial-to-trial variability and correlation between neurons (Qi and Constantinidis, 2012a,

2012b). This led to our present analysis where—guided by experimental and theoretical

predictions (Rigotti et al., 2013)—we examined NMS as another potential source of

enhanced ability to represent WM information after training.

In agreement with our hypothesis, we found that training increased the proportion

of neurons that exhibit NMS. However, training does not seem to be a prerequisite, as

NMS was also observed in animals that were naïve to any cognitive training. Prior

research has established that the human and primate PFC represent stimuli in memory

even when not prompted to do so (Foster et al., 2017), or without training in WM tasks

(Meyer et al., 2007). Our finding of NMS neurons in naïve monkeys provides another

exemplar of that principle. However, NMS only increased for certain types of task

information and not for others, thus suggesting that its role in the simultaneous

representation of multiple types of information is not universal across tasks. These results

stand in agreement with some prior studies, which have failed to uncover substantial

NMS in the tasks they employed (Cavanagh et al., 2018). Examining where NMS failed

to appear—and where WM representations fail to spontaneously appear— will be an

important area of future investigation for NMS.

The greatest increase in the proportion of neurons that exhibited NMS for the

spatial working memory task was observed at the mid-dorsal region during the sample

presentation period (Fig. 5). This disproportionate increase in NMS neurons was

associated with a modest decrease in neurons that exhibit CS, as predicted by theoretical

studies (Lindsay et al., 2017). However, this finding, too did not generalize across

conditions. During the delay periods of the spatial task, we saw an across-the-board

increase in neurons with CS, which were much more abundant in the trained than the

naïve PFC (Fig. 3). In fact, the increase in feature selectivity after training was driven

almost exclusively by CS cells, suggesting a potential division of labor between NMS

and CS in the PFC for different types of information.

Task Complexity and Difficulty

Another potential factor that determines the emergence of NMS is task complexity. NMS

may arise exclusively in highly complex tasks that require subjects to maintain and

combine multiple types of information in their WM, simplifying the involved neural

circuits to achieve greater efficiency (Rigotti et al., 2013). We thus tested this concept by

applying a dataset that relied on three tasks which differed in complexity (and overall

difficulty). The spatial and feature tasks each required maintenance of a single stimulus

property in memory (location or shape). The conjunction task required both. Surprisingly

however, we did not observe a higher incidence of NMS in the conjunction task when

compared to the feature task. Moreover, we observed a much lower incidence of NMS in

the feature task compared to the spatial task despite the fact that the latter was no more

complex or difficult for the monkeys to perform (Meyer et al., 2011; Riley et al., 2018).

This implies that NMS in the PFC may not be necessary for certain types of information,

like object shape, even when the task complexity is high. Alternatively, we may also

consider the possibility that the increased NMS that results from training may be

sufficient for the majority of behavioral requirements, without any additional increases

required. Future research is therefore necessary to assess and examine all of these

possibilities, and more.

Regional Specialization

Different types of information are represented across the dorso-ventral and anterior-

posterior axes of the PFC (Constantinidis and Qi, 2018), and examining the regional

distribution of NMS neurons within the PFC therefore bears a clear importance. We

found that NMS was most strongly demonstrated in the mid-dorsal area for the spatial

task and the posterior dorsal area for the feature task. This pattern was generally

consistent with the known distribution of neuronal selectivity for stimuli in the PFC

(Riley et al., 2018).

Information Content and Task Performance

A critical issue regarding the role of Mixed Selectivity is whether nonlinear match

selectivity, by virtue of representing information more efficiently, is also more necessary

for effective task performance (Rigotti et al., 2013). We relied on a linear SVM decoder

to decipher the specific information that may be represented by NMS cells, compared to

CS cells. In the current study, we found similar quantities of information could be

decoded from (equal-sized) populations of CS and NMS neurons, though the coding

dynamics for some types of information were significantly different between CS and

NMS cells. Similarly, when we compared the NMS levels of successful and failed task

trials, we were surprised to find that there was no appreciable difference. This suggests

that loss of information encoded in a nonlinear manner is not the primary factor of

successfully maintaining information in the conscious mind. An important caveat for this

conclusion is that the combination of small and unbalanced number of error trials in

match vs. nonmatch conditions make the detection power for the interaction fairly small

in our analysis. Moreover, with very little NMS presented even in success trials after

matching the trial number for the error condition, a floor effect may have prevented a

further decline from becoming apparent. Nonetheless, our result reinforces the idea that

NMS is not necessary in all tasks, without which performance fails. An interesting

observation in this analysis was that the proportion of cells tuned to spatial location was

elevated in error trials. The result may imply that task success also depends on the task

relevance of the represented information, with error trials incorporating greater quantities

of task irrelevant spatial information and therefore unnecessarily drawing away WM

resources without benefit. Ultimately, by comparing and evaluating the conditions in

which NMS emerges, we may decipher its true role in WM and other cognitive functions.

REFERENCES

Ahlheim, C., Love, B.C. (2018). Estimating the functional dimensionality of neural representations. Neuroimage.*179*, 51-62.

Asaad WF, Rainer G, Miller EK. 2000. Task-specific neural activity in the primate prefrontal cortex. J Neurophysiol. 84: 451-459.

Baddeley A. 2012. Working memory: theories, models, and controversies. Annu Rev Psychol. 63: 1-29.

Barak O, Rigotti M, Fusi S. 2013. The sparseness of mixed selectivity neurons controls the generalization-discrimination trade-off. J Neurosci. 33: 3844-3856.

Bherer L, Kramer AF, Peterson MS, Colcombe S, Erickson K, Becic E. 2008. Transfer effects in task-set cost and dual-task cost after dual-task training in older and younger adults: further evidence for cognitive plasticity in attentional control in late adulthood. Exp Aging Res. 34: 188-219.

Buonomano DV, Maass W. 2009. State-dependent computations: spatiotemporal processing in cortical networks. Nat Rev Neurosci. 10: 113-125.

Cavanagh SE, Towers JP, Wallis JD, Hunt LT, Kennerley SW. 2018. Reconciling persistent and dynamic hypotheses of working memory coding in prefrontal cortex. Nature communications. 9: 3498.

Constantinidis C, Klingberg T. 2016. The neuroscience of working memory capacity and training. Nat Rev Neurosci. 17: 438-449.

Constantinidis C, Procyk E. 2004. The primate working memory networks. Cogn Affect Behav Neurosci. 4: 444-465.

Constantinidis C, Qi XL. 2018. Representation of Spatial and Feature Information in the Monkey Dorsal and Ventral Prefrontal Cortex. Front Integr Neurosci. 12: 31.

Cortese S, Ferrin M, Brandeis D, Buitelaar J, Daley D, Dittmann RW, Holtmann M, Santosh P, Stevenson J, Stringaris A, Zuddas A, Sonuga-Barke EJ, European AGG. 2015. Cognitive training for attention-deficit/hyperactivity disorder: meta-analysis of clinical and neuropsychological outcomes from randomized controlled trials. J Am Acad Child Adolesc Psychiatry. 54: 164-174.

Cueva CJ, Saez A, Marcos E, Genovesio A, Jazayeri M, Romo R, Salzman CD, Shadlen MN, Fusi S. 2020. Low-dimensional dynamics for working memory and time encoding. Proc Natl Acad Sci U S A. 117: 23021-23032.

Curtis CE, D'Esposito M. 2004. The effects of prefrontal lesions on working memory performance and theory. Cogn Affect Behav Neurosci. 4: 528-539.

Dux PE, Tombu MN, Harrison S, Rogers BP, Tong F, Marois R. 2009. Training improves multitasking performance by increasing the speed of information processing in human prefrontal cortex. Neuron. 63: 127-138.

Foster JJ, Bsales EM, Jaffe RJ, Awh E. 2017. Alpha-Band Activity Reveals Spontaneous Representations of Spatial Position in Visual Working Memory. Curr Biol. 27: 3216-3223 e3216.

Fukuda K, Awh E, Vogel EK. 2010. Discrete capacity limits in visual working memory. Curr Opin Neurobiol. 20: 177-182.

Fusi S, Miller EK, Rigotti M. 2016. Why neurons mix: high dimensionality for higher cognition. Curr Opin Neurobiol. 37: 66-74.

Jaeggi SM, Buschkuehl M, Jonides J, Perrig WJ. 2008. Improving fluid intelligence with
training on working memory. ProcNatlAcadSciUSA. 105: 6829-6833.
Johnston WJ, Palmer SE, Freedman DJ. 2020. Nonlinear mixed selectivity supports
reliable neural computation. PLoS Comput Biol. 16: e1007544.
Klingberg T, Fernell E, Olesen P, Johnson M, Gustafsson P, Dahlstrîm K, Gillberg CG,
Forssberg H, Westerberg H. 2005. Computerized Training of Working Memory in
Children with ADHD - a Randomized, Controlled Trial. J Am Acad Child
Adolesc Psychiatry. 44: 177-186.
Klingberg T, Forssberg H, Westerberg H. 2002. Training of working memory in children
with ADHD. J Clin Exp Neuropsychol. 24: 781-791.
Lindsay GW, Rigotti M, Warden MR, Miller EK, Fusi S. 2017. Hebbian Learning in a
Random Network Captures Selectivity Properties of the Prefrontal Cortex. J
Neurosci. 37: 11021-11036.
Machens CK, Romo R, Brody CD. 2010. Functional, but not anatomical, separation of
"what" and "when" in prefrontal cortex. J Neurosci. 30: 350-360.
Mansouri FA, Matsumoto K, Tanaka K. 2006. Prefrontal cell activities related to
monkeys' success and failure in adapting to rule changes in a Wisconsin Card
Sorting Test analog. J Neurosci. 26: 2745-2756.
Meyer T, Constantinidis C. 2005. A software solution for the control of visual behavioral
experimentation. J Neurosci Methods. 142: 27-34.
Meyer T, Qi XL, Constantinidis C. 2007. Persistent discharges in the prefrontal cortex of
monkeys naive to working memory tasks. Cereb Cortex. 17 Suppl 1: i70-76.
Meyer T, Qi XL, Stanford TR, Constantinidis C. 2011. Stimulus selectivity in dorsal and
ventral prefrontal cortex after training in working memory tasks. J Neurosci. 31:
6266-6276.
Meyers EM, Qi XL, Constantinidis C. 2012. Incorporation of new information into
prefrontal cortical activity after learning working memory tasks. Proc Natl Acad
Sci U S A. 109: 4651-4656.
Morris RG, Baddeley AD. 1988. Primary and working memory functioning in
Alzheimer-type dementia. J Clin Exp Neuropsychol. 10: 279-296.
Owen AM, Hampshire A, Grahn JA, Stenton R, Dajani S, Burns AS, Howard RJ, Ballard
CG. 2010. Putting brain training to the test. Nature. 465: 775-778.
Parthasarathy A, Herikstad R, Bong JH, Medina FS, Libedinsky C, Yen SC. 2017. Mixed
selectivity morphs population codes in prefrontal cortex. Nat Neurosci. 20: 1770-
1779.
Peijnenborgh JC, Hurks PM, Aldenkamp AP, Vles JS, Hendriksen JG. 2015. Efficacy of
working memory training in children and adolescents with learning disabilities: A
review study and meta-analysis. Neuropsychol Rehabil. 1-28.
Qi XL, Constantinidis C. 2012a. Correlated discharges in the primate prefrontal cortex
before and after working memory training Eur J Neurosci. 36: 3538-3548.
Qi XL, Constantinidis C. 2012b. Variability of prefrontal neuronal discharges before and
after training in a working memory task. PLoS ONE. 7: e41053.
Qi XL, Elworthy AC, Lambert BC, Constantinidis C. 2015. Representation of
remembered stimuli and task information in the monkey dorsolateral prefrontal
and posterior parietal cortex. J Neurophysiol. 113: 44-57.

Qi XL, Meyer T, Stanford TR, Constantinidis C. 2011. Changes in Prefrontal Neuronal
Activity after Learning to Perform a Spatial Working Memory Task. Cereb
Cortex. 21: 2722-2732.
Rigotti M, Barak O, Warden MR, Wang XJ, Daw ND, Miller EK, Fusi S. 2013. The
importance of mixed selectivity in complex cognitive tasks. Nature. 497: 585-590.
Rigotti M, Ben Dayan Rubin D, Wang XJ, Fusi S. 2010. Internal representation of task
rules by recurrent dynamics: the importance of the diversity of neural responses.
Front Comput Neurosci. 4: 24.
Riley MR, Constantinidis C. 2016. Role of prefrontal persistent activity in working
memory. Front Syst Neurosci. 9: 181.
Riley MR, Qi XL, Zhou X, Constantinidis C. 2018. Anterior-posterior gradient of
plasticity in primate prefrontal cortex. Nature communications. 9: 3790.
Rossi AF, Bichot NP, Desimone R, Ungerleider LG. 2007. Top down attentional deficits
in macaques with lesions of lateral prefrontal cortex. J Neurosci. 27: 11306-
11314.
Schwaighofer M, Fischer F, Buhner M. 2015. Does Working Memory Training Transfer?
A Meta-Analysis Including Training Conditions as Moderators. Educ Psychol. 50:
138-166.
Subramaniam K, Luks TL, Fisher M, Simpson GV, Nagarajan S, Vinogradov S. 2012.
Computerized cognitive training restores neural activity within the reality
monitoring network in schizophrenia. Neuron. 73: 842-853.
Warden MR, Miller EK. 2010. Task-dependent changes in short-term memory in the
prefrontal cortex. J Neurosci. 30: 15801-15810.
Westerberg H, Jacobaeus H, Hirvikoski T, Clevberger P, Ostensson ML, Bartfai A,
Klingberg T. 2007. Computerized working memory training after stroke--a pilot
study. Brain Inj. 21: 21-29.
781

783
Figure 1. Task structure and stimuli used. The animals were required to maintain
center fixation throughout both active and passive task trials. At the end of active
tasks trials however, monkeys were required to make a saccade to a green
target if the stimuli matched or to a blue target if the stimuli did not match. (A)
Spatial location match-to-sample task, nine possible cue locations in a session
shown in the inset. (B) Shape feature match-to-sample task, 8 possible shapes in
a session shown in the inset. (C) Spatial-shape conjunction task, up to two
locations and two stimuli shapes were used for any single particular session.
Stimuli in all tasks extended 2 degree of visual angle.

793
Figure 2. Exemplar neural responses from the spatial task for CS (classical
selective), LMS (linear mixed selective) and NMS (nonlinear mixed selective)
cells, defined by the task variables of stimulus location and match status.
Selectivity classification were based on the spike responses of the 500 ms
sample period. The locations the stimuli were color coded, with a solid line/bar
representing when the stimulus was a match with the cue, and a dash line/bar
representing when the stimulus was a nonmatch with the cue. Shaded regions
and error bar indicate 2 times SE of firing rate.
802

803
Figure 3. Training increased mixed selectivity preferentially in the spatial task. (A)
Bar graphs show the proportions of cells tuned to stimuli identities
(Location/Shape), matching status and their interaction (i.e. NMS) in different
stages of the task trials, both before and after the animals were trained for the
active tasks. Pie charts show the proportion of different selectivity categories
(NS, CS, LMS and NMS) in the sample and delay2 periods of both tasks, both
before and after the animals were trained for the active tasks. (B) Plots of
corresponding proportion changes. (C) Same as (A) but examining the
interaction between stimuli identities (Location/Shape), and task epoch
(cue/delay1 vs sample/delay2 period), instead of trials matching status. (D) Plots
of corresponding proportion changes.
815

817
818
Figure 4. (A) Analysis of F score for the interaction term (stimuli identity ×
match/nonmatch) shows that the degree of mixed selectivity increased after
training for the spatial task only. Black dots in the box represent mean, box
boundaries indicate 25%-75% range, and whiskers represents 1.5 IQR. (B)
Dimensionality measure of neural responses in the spatial (left) and feature
(right) task, before and after training in the active tasks.

Figure 5. Cell selectivity changes by brain regions. (A) PFC subdivisions that
were utilized for recording in the current study. (B) The effects of training in for
the active feature task on the proportion of different selectivity categories (NS,
CS, LMS and NMS) in the sample period. There were significant increases in the
proportion of LMS cells in all three PFC regions included for analysis, but
relatively low increases in the proportion of NMS cells. (C) The effects of training
in the active spatial task on the proportion of different selectivity categories (NS,
CS, LMS and NMS) in the sample period. The greatest increase in NMS
occurred at the mid-dorsal region.

Figure 6. Cell selectivity in tasks with different task complexity. Analyses were performed on neural data from the same population of cells, with matching numbers of trials in the feature and the conjunction tasks. Only trials with the same stimuli were included in this analysis. (A) Examining interaction (NMS) across stimulus preference and matching status. Bar graphs show the proportions of cells tuned to stimuli shape, trials matching status and their interaction in different stages of both the feature and conjunction tasks. Pie charts display the proportion of different selectivity categories (NS, CS, LMS and NMS) in corresponding sample and delay2 periods. (B) Cell selectivity category mapping cross tasks. (C) F scores of the interaction term in the ANOVA were compared between cells that were classified as NMS cell in both tasks (overlapping cells), and those only classified as NMS in one of the tasks (non-overlapping cells).

850
Figure 7. Cell selectivity in tasks with the same sensory input but different
behavioral requirements. Analyses were performed on neural data from the same
population of cells, with matching number of trials in the passive and active
spatial tasks. Only trials that had the exact same stimuli pairs in both tasks were
included in this analysis. (A) Examining interaction (NMS) between stimulus
preference and matching status. Bar graphs show the proportions of cells tuned
to stimuli location, trials matching status and their interaction in different stages of
the tasks. Pie charts display the proportion of different selectivity categories (NS,
CS, LMS and NMS) in the sample period. (B) Cell selectivity category mapping
across tasks in the sample period. (C) F scores of the interaction term in the
ANOVA were compared between cells that were classified as NMS cell in both
tasks (overlapping cells), and those only classified as NMS in one of the tasks
(non-overlapping cells).

864
Figure 8. Comparing linear SVM decoder information between pure and linear
selective cells (CS and LMS) vs. NMS cells. An equal number of linear (CS and
LMS) and NMS cells were randomly selected from the sample or delay2 period.
The selectivity categories were defined by spiking count in corresponding periods
with reference to stimuli identity (stimuli location or shape) and matching status.
The decoders were trained to classify either stimuli identity or match/nonmatch
status with z-score normalized pseudo-population response. (A) Decoding
performance in the spatial task before and after training. (B) Decoding
performance in the feature task before and after training. Red and blue bars
indicate time points when the performance for NMS and linear cells differs
significantly within the shaded regions.

Figure 9. Coding dynamics of pure and linearly selective cells (CS and LMS) vs. NMS cells. Linear kernel SVM decoders were trained to perform cross-temporal decoding with different selectivity populations in the delay2 period, for both spatial and feature tasks, as indicated by the Y-axis. The decoder was then required to predict whether a match or non-match occurred at each time point based on a different test set of data, as indicated by the X-axis. Normalized decoding accuracy is indicated in the color bar, demonstrating how spatial and feature WM representations can be decoded from specific patterns of neural activity. Coding of matching information for NMS cells is more stable across time for the spatial task.

888
Figure 10. Comparison of cell selectivity in correct and error trials in the same
population for the spatial task, after controlling for trial number and location pairs
used. Two forms of mixed selectivity were examined (location x matching,
location x task epoch). No change (location x matching delay2 period) or
increase (location x task stim and delay epochs, location x matching sample
period) in the F score of the interaction term of ANOVA results were observed in
cells with significant interaction (NMS) terms. Higher F score for the variable of
stimuli location was observed in error trials in the location x Epoch comparison.
Box boundaries represent 25%-75% data range, whiskers indicate 1.5 IQR and
squares indicates means across cells.

899
Fig. S1. Plotting conventions same as Fig. 6. Comparing cell selectivity between
the spatial and the conjunction tasks. The analysis was performed on neural data
from the same population of cells, with matching numbers of trials in the spatial
and the conjunction tasks. Only trials with the same stimuli were included in this
analysis. (A) Examining interaction (NMS) between stimulus preference and
matching status. Bar graphs display the proportions of cells tuned to stimuli
shape, trials matching status and their interaction in different stages of the tasks,
in both the feature and conjunction task. Pie charts display the proportion of
different selectivity categories (NS, CS, LMS and NMS) in corresponding sample
and delay2 periods. (B) Cell selectivity category mapping across tasks. (C) F
scores of the interaction term in the ANOVA were compared between cells that
were classified as NMS cells in both tasks (overlapping cells), and those only
classified as NMS in one of the tasks (non-overlapping cells). Similar to the
comparison between the feature and the conjunction tasks, no change in the
proportion of NMS was observed.

915
Fig. S2. Incorporation of new information after training for the spatial and the
feature tasks. Linear SVM decoders were trained to classify either stimuli identity
or match/nonmatch status with z-score normalized pseudo-population response.
Color bars indicate time points when the performance for NMS and linear cells
differs significantly. Red bar for decoding matching status, blue bar for decoding
stimuli identity.