Increasing temperature threatens an already endangered coastal dune plant

Contact information of authors: Aldo Compagnoni[1,2], Eleanor Pardini[3], & Tiffany M. Knight[1,2,4]

[1]Institute of Biology, Martin Luther University Halle-Wittenberg, Am Kirchtor 1,06108, Halle

(Saale), Germany, aldo.compagnoni@gmail.com

[2]German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz

5e, 04103, Leipzig, Germany

[3]Environmental Studies Program, Washington University in St. Louis, 1 Brookings Drive, Box

1165, St. Louis, MO 63130, USA

[4]Department of Community Ecology, Helmholtz Centre for Environmental Research – UFZ,

Theodor-Lieser-Straße 4, 06120, Halle (Saale), Germany

Person to whom correspondence should be sent: Aldo Compagnoni, German Centre for

Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103, Leipzig,

Germany, aldo.compagnoni@gmail.com

ABSTRACT

Climate change has the potential to reduce the abundance and distribution of species and threaten

global biodiversity, but it is typically not listed as a threat in classifying species conservation

status. This likely occurs because demonstrating climate change as a threat requires data-

intensive demographic information. Moreover, the threat from climate change is often studied in

specific biomes, such as polar or arid ones. Other biomes, such as coastal ones, have received

little attention, despite being currently exposed to substantial climate change effects. We forecast

the effect of climate change on the demography and population size of a federally endangered

coastal dune plant (*Lupinus tidestromii*). We use data from a 14-year demographic study across

seven extant populations of this endangered plant. Using model selection, we found that survival

and fertility measures responded negatively to temperature anomalies. We then produced

forecasts based on stochastic individual based population models that account for uncertainty in

demographic outcomes. Despite large uncertainties, we predict that all populations will decline if

temperatures increase by 1° Celsius. Considering the total number of individuals across all seven

populations, the most likely outcome is a population decline of 90%. Moreover, we predict

extinction is certain for one of our seven populations. These results demonstrate that climate

change will profoundly decrease the current and future population growth rates of this plant, and

its chance of persistence. Thus, our study provides the first evidence that climate change is an

extinction threat for a plant species classified as endangered under the USA Endangered Species

Act.

Keywords: climate change, forecast, integral projection model, individual based model, life

table response experiment, population dynamics.

INTRODUCTION

Year to year variation in weather influences demographic processes such as survival, growth and reproduction, and ultimately population persistence (Sæther et al. 2000). For many natural populations, climate change will increase the frequency of weather conditions outside the physiological tolerances of individuals, and is thus expected to reduce the number, size and distribution of populations (IPCC 2014). As a result, conservation planning (Akçakaya et al., 2014) and projections of biodiversity change (Pereira et al. 2010) will benefit from understanding how climate change will threaten population viability.

Researchers typically use species distribution models (SDMs) to assess the effects of climate change on species (Elith and Leathwick 2009). However, species distributions might include declining populations of the species because of a currently unsuitable climate (Schurr et al. 2012). Subsequently, while SDMs describe current ranges well, their prediction of future ranges are highly uncertain (Keenan et al. 2011). Hence, forecasting the effect of climate change on species persistence should be more reliable using process-based models that require long-term demographic data across the entire life cycle of the organism (Ehrlén and Morris 2015, Paniw et al. 2019). Despite this, our knowledge on how climate drives demography is relatively limited, most studies coming from animals in cold and arid biomes. For example, changing temperature has been linked to reductions in population growth rates of alpine plants (Doak and Morris 2010, Campbell 2019, Iler et al. 2019) and polar animals (Jenouvrier et al. 2009, Hunter et al. 2010). Changing precipitation has been linked to changes in population growth rates of organisms in arid environments (Tews and Jeltsch 2004, Jonzén et al. 2010). However, the effect of climate

change in other biomes, regardless of their extent or importance to humans, has received less

attention.

Coastal ecosystems are a biome for which the extinction threat from climate change has received

little attention. This occurs despite the importance of species conservation for human well-being

(Díaz et al. 2019), and despite the importance of coastal ecosystems and their susceptibility to

climate change. Coastal ecosystems occupy a small portion of terrestrial surface, but are home to

41% of the human population (Martínez et al. 2007) and to a large portion of the earth's

economy; for example, 31% of the USA gross domestic product in 1985 (Luger 1991). Coastal

biomes protect inland ecosystems during storms by attenuating or resisting wave action (Barbier

2015). Current and future climate change in coastal ecosystems is altering precipitation patterns,

increasing the frequency of high temperature extremes (USGCRP 2018), and causing habitat loss

and salt water inundation (Feagin et al. 2005). Thus, it is timely to consider the effects of climate

on the demography of coastal species.

Understanding the effect of climatic drivers on future population size requires quantifying how

the link between climate and demography translates into population dynamics. Studies

establishing such link are the foundation to prioritize conservation efforts. For example, the polar

bear was listed as a threatened species under the endangered species act thanks to demographic

studies (USFWS 2015). Demographic analysis linked warming-induced reductions in sea ice

extent to reductions in survival and breeding probabilities, and increased risk of extinction

(Hunter et al. 2010). As a result of this work, the US Fish and Wildlife Service indicated curbing

Arctic warming as the most important conservation action for this species (USFWS 2015). Such

demographic studies are most useful when carried out across multiple populations, in order to

account for the substantial variation in populations sizes and vital rates (Glenn et al. 2010).

Here, we examine the effect of changing climate on the vital rates and population growth of a

perennial coastal dune plant. We first link the effect of climate on separate vital rates, and we use

an integral projection model to estimate their contribution to population growth rate. Then, we

use an individual based model to forecast the effect of projected climate change on future

population numbers in the next 30 years. These numerical simulations account for the effect of

initial population numbers, year to year stochastic weather, and demographic stochasticity. We

parameterize these models using a 14-year data set that comprehensively sampled 7 of the 15

known remaining populations of this species. The populations we included range from the

smallest to the largest known (17 and 180000 individuals, respectively). Our focal species,

Lupinus tidestromii, is listed as endangered under the USA endangered species act, but to date,

climate change is not considered a threat to this or any plant species.

MATERIALS AND METHODS

Study Species and Site

Lupinus tidestromii (Fabaceae) is a perennial, herbaceous plant that is endemic to coastal dunes

in northern California, where it is found in 15 extant populations. These populations are located

in Sonoma, Monterey, and Marin counties, seven of these are contained within Point Reyes

National Seashore (Appendix S1). In this protected area, the population located at Abbotts

Lagoon contains more than 50% of the remaining individuals of the entire species (USFWS

2009). *L. tidestromii* was listed as federally endangered in 1992 and among its primary threats

are habitat loss, trampling by visitors and cattle, hybridization, direct competition with invasive

plants, and elevated levels of seed predation in the presence of the invasive grass *Ammophila*

arenaria (USFWS 2009, Dangremond et al. 2010). Our study was conducted at Point Reyes

National Seashore (latitude: 33.1120, longitude: -122.9579). This site has a Mediterranean

climate, with wet winters and dry summers (Evens 2008).

L. tidestromii is a herbaceous perennial plant that typically lives three years. It produces prostrate

stems, and small, upright inflorescences with a whorl of flowers that produce leguminous fruits

(Baldwin et al. 2012). Seeds are dispersed locally by explosive dehiscence (USFWS 1998).

Seeds have a tough coat that likely allows survival for many years in the seed bank. However,

we assume that if they do not germinate within the first two years, seeds become buried by sand.

This is a reasonable assumption in dunes that are highly stabilized by introduced invasive plants.

Flowers experience occasional insect herbivory, and fruits and seeds experience pre- and post-

dispersal seed predation by the native *Peromyscus maniculatus* (deer mouse). We have never

observed pre-dispersal insect seed predation.

Monitoring

We began demographic censuses in 2005 at three sites (AL, ATT, NB) in 2005, and expanded

monitoring at an additional four sites (Pop9, DR, BR, and BS) in 2008 (Appendix S1: Fig. S1).

Populations NB, BS, and DR are small enough that we monitored every individual plant in the

population each year. In AL and BR, and ATT and Pop9 starting in 2013 and 2015, we stratified

sampling, censusing plants located in clusters across the extent of the population. We

haphazardly chose plants across a range of sizes and microhabitats. Annually, we also tagged at

least 50 new seedlings across each site. We know total population counts for ATT, Pop9, NB,

BS, and DR for every year. We inferred population counts for NB, BS, and DR, and we counted

the total number of individuals in the populations after starting stratified sampling in ATT and

Pop9.

Vital rate data

To quantify the vital rates, in 2005, we started censusing tagged plants in June when fruits on

most plants were dehiscing. These censuses quantified the transition rates from one year (t) to the

next ($t+1$). To track individuals, we attached numbered aluminum tags to the basal stem of

plants, or staked these tags near plants. We relocated plants each year using a GPS and a metal

detector. For each plant, every year we recorded size, survival, flowering status, and number of

flowering inflorescences. Our way to estimate size changed after 2008. Before 2008 we

quantified size counting the number of branches on a plant. After 2008, we quantified the size of

each individual approximating it using as an ellipse. The major axis of this ellipse was the

longest segment covered by each individual, the minor axis the width perpendicular to the major

axis. For data before 2008, we inferred the area of the ellipse by developing a regression that

used a sample of plants for which we counted branches and measured major and minor axis

length. In total, this regression contained 234 plants (82 measured in 2008, 152 in 2018). We

predicted area as a quadratic function of the number of branches (Appendix S1: Fig. S2).

For reproductive plants, we recorded the total number of inflorescences and of those that were

aborted, consumed, or intact. Flowers on inflorescences can abort or produce fruits; fruits can be

clipped by mice or remain intact, dehisce and disperse seeds. We used the fate of inflorescences

to calculate abortion, and consumption (pre-dispersal seed predation), for each population and

year.

To calculate abortion, we summed the number of aborted inflorescences and divided by the sum

of the total number of observed inflorescences. To calculate consumption, we summed the

number of clipped inflorescences, and divided by the sum of the number of not aborted

inflorescences. The resulting abortion and consumption are population-level means, because we

performed the sums to calculate these metrics across all the plants in each population. We do not

have observations on abortion and consumption for all populations and all years. We have

abortion rates for all populations in 2010 and 2011, and from 2013 to 2017 (Appendix S2: Fig.

S5) and consumption rates for all years except 2012 (Appendix S2: Fig. S6). In years and

locations for which we did not have an abortion or consumption rate, we used the population-

specific mean taken across censuses.

We assumed the number of fruits per inflorescence, and the number of seeds per fruit are

constant across years and populations. We calculated fruits per inflorescence using data from

2011, averaging the number of fruits across all inflorescences for each plant, and then averaging

across plants to produce a population-level value. We quantified the average number of seeds per

fruit by counting seeds in 213 randomly collected fruits from multiple years and populations:

AL: 2005 (23), 2008 (3), 2009 (15), 2010 (39); ATT: 2011 (33); BR: 2010 (23); DR: 2010 (10),

2011 (17); NB: 2010 (16) Pop9: 2010 (15).

We estimated germination rates through a field germination trial and calculated a recruitment

adjustment factor to account for post-dispersal seed predation, and for other sources of

unaccounted seed or seedling mortality. We installed caged seed baskets in 2008 at the AL

population, and recording the proportion of seedlings that germinated in 2009 (g_1), 2010 (g_2),

and 2011 (g_3) (Pardini et al. 2015). Here g_2, and g_3 represent emergence out of seed bank stages.

We then calculated a recruitment adjustment factor to account for post-dispersal seed predation

and for other sources of unaccounted seed or seedling mortality. To account for these other

sources of seed and seedling loss, we used our known population counts to estimate a

recruitment adjustment factor, δ_p, for each p population (Appendix S3). We modified

germination rates g_1, g_2, and g_3 employed in the IPM using δ_p as $g_t\left(1-\delta_p\right)$. We could only

calculate δ_p for the five populations with known population counts (see "monitoring"). For

population BR and ATT, we applied the mean of δ_p from BS and DR, and from ATT and Pop9,

respectively the closest populations to BR and ATT.

Modeling of size-dependent vital rates and climate effects

We modeled four vital rates as size dependent processes for each year and site: survival, growth,

the probability of flowering, and the number of inflorescences. We modeled these vital rates

based on generalized mixed linear models. We fit models in R using the lme4 package (Bates et

al. 2015), and we performed a preliminary model selection based on Akaike Information

Criterion (Burnham and Anderson 2002). In this preliminary model selection, we compared

models predicting vital rates using linear, quadratic, and cubic predictors. Our baseline model

structure included a random intercept and slope (the effect of size on the vital rate) for both years

and populations. We modeled survival and probability of flowering data as Bernoulli distributed,

the number of inflorescences as Poisson distributed, and growth data as normally distributed. For

survival, the best model included a linear, squared, and cubic predictor of size. For the remaining

three vital rates, the best model included only a linear predictor (Appendix S4).

We selected which climatic predictor, if any, affected vital rates by performing a leave-one-year-

out cross-validation. We selected whether climatic drivers increased the predictive power of our

linear models (Appendix S4) and, if so, what climatic drivers improved predictive power the

most. The models in this cross-validation simply added the climatic predictor to the vital rate

models (Appendix S4). We considered four climatic predictors: temperature, precipitation,

Oceanic Niño Index (ONI) (CPC n.d.), and the standardized precipitation evapotranspiration

index (SPEI, Vicente-Serrano et al. 2009). We obtained temperature and precipitation data

starting in 1990 from the PRISM dataset (Daly et al. 1994), the ONI data from the internet site of

the Climate Prediction Center of the National Oceanic and Atmospheric Administration (CPC

n.d.), and we derived the SPEI using our precipitation and temperature data using the R package

SPEI (Beguería and Vicente-Serrano 2017). We calculated the yearly climate anomalies for each

one of these four drivers. We computed anomalies using the mean and standard deviation

observed between 1990 and 2018. Here, we define "year" as the time elapsing between

demographic censuses: from June to May of the following year.

For each of the four climatic drivers, we used two yearly anomalies: one referred to the year

leading up to the current census (*t*), and one to the year leading up to the preceding census (*t-1*).

For example, for year 2012 (t=2012) we considered the climate anomaly observed between June

2011 and May 2012, and between June 2010 and May 2011 to predict the flowering probability

and number of inflorescences observed in June 2012, and to predict the survival and growth

occurred from June 2012 to June 2013. We thus tested for a lag in the effect of climatic drivers

on plant vital rates (see also Teller et al. 2016; Tenhumberg et al. 2018).

We measured model performance summing the negative log-likelihood of each out-of-sample

prediction, with the best model having the lowest score. We used negative log-likelihood as it is

a proper scoring rule (Gneiting and Raftery 2007) across our generalized linear models. We also

ranked models based on root mean squared error which, despite being not proper, yielded

qualitatively identical results (Appendix S5: Table S1-2). We fit the final models using the best

climatic predictor (if any).

Population projections

We quantified the effect of a changing climate on the population dynamics of *L. tidestromii*

looking at its projected long-term population growth rates and population sizes. We projected

long-term population growth rates using an integral projection model (IPM, Ellner et al. 2016),

and future population sizes using an individual based model (IBM). We used IBMs to add

demographic stochasticity, which is a key component to predict population sizes, particularly for

smaller populations (Lande 1993, Caswell 2001).

We used an IPM because it is a computationally inexpensive way to perform stochastic

population projections of species whose dynamics depend on a continuous state variable: in this

case, the size of individuals. We constructed the IPM using the parameters of the vital rates

models described above. Most of these parameters changed based on year and population

(Appendix S6).

We first used the IPMs to compute how climate affects λ_s, the long-term stochastic population

growth rate. We calculated λ_s across the whole range of climate anomalies observed during our

study period (Appendix S6). Our best climatic predictor was always temperatures. Therefore, we

calculated λ_s from an average annual temperature of about 11° Celsius to an average annual

temperature of 13°.

To quantify the contribution of each vital rate in changing λ_s under a different climate, we used

the IPM to perform a Life Table Response Experiment (LTRE) and a sensitivity analysis. The

LTRE quantified the difference in λ_s observed when only one of the vital rates responded to

deviations from the mean temperature. For example, we calculated λ_s for an IPM where only

survival experienced the full range of temperature anomalies. In these simulations, all other vital

rates experienced average climate (about 12° Celsius). We then calculated the contribution of

each vital rate to the effect of climate. This contribution is the difference between the λ_s

computed by these stochastic simulations, and the λ_s where all vital rates experienced a 12°

Celsius climate.

To further interpret the LTRE results, we calculated vital rate elasticities. The LTRE results by

combination of two factors. First, the effect size of climate on each vital rate, which is provided

by parameter β_c in the vital rate models (Appendix S6). Second, the sensitivity of stochastic

population growth rate (λ_s) to each vital rate. We approximated λ_s using the asymptotic

population growth rate (λ). We then calculated the elasticity of each vital rate, which is the

proportional response of λ to a proportional increase in vital rate parameter.

Forecast

We forecasted population numbers through an IBM that incorporates probabilistic events of the

life cycle by projecting the dynamics of each individual. Given our life cycle (Appendix S7),

these probabilistic events were: survival, growth, probability of flowering, production of

inflorescences, the abortion of inflorescences, the consumption of inflorescences, and the

germination of seeds. We simulated each of these processes using the appropriate statistical

distributions (Appendix S7).

We used this IBM to produce a forecast of population sizes in the next 30 years that included

uncertainty arising from process and demographic stochasticity. We projected each population

for 30 time steps, and we replicated each run 1000 times. Each run included a different sequence

of years, and a different realization of each vital rate process (Appendix S7). We started each

simulation with the number of individuals observed in June 2018, assuming the stable stage

distribution suggested by the population-specific IPM model. We ran two simulations: one

assuming the current mean climate (average annual temperature of 12° Celsius), and one

assuming the largest weather anomaly we observed during our 2005-2018 study period (which

was a year with an average temperature of 13° Celsius). A mean annual temperature of 13°

Celsius is a conservative forecast, because projections suggest that in 2055, temperature will be

on average 2.1 Celsius higher than the 1990-2018 mean (Vose et al. 2017). We used the 1000

simulations for each population and climate scenario to calculate 95% confidence intervals of

population abundances during the 30-year projections.

RESULTS

Observed and future climate variation

Across our 14-year study period, we observe variation in weather on par with that across longer

time periods (Appendix S1: Fig. S4), but not within the range of weather projected with climate

change by the middle of the century (Wang et al. 2016, n.d.). For example, mean annual

temperature was 11.5° C across the study period, 12° C from 1990-2018. This temperature is

projected to increase to about 14° C (range: 13.8-14.6° C) by 2055 according to the

Representative Concentration Pathway of 4.5 W/m^2 (Wang et al. 2016, n.d., Vose et al. 2017).

The range of annual temperatures observed from 1990-2018 was 11-13° C, and thus we only

consider conservative scenarios of a 1° C increase in temperature.

Vital rates response to climate

Survival, growth, probability of flowering, number of inflorescences, rates of abortion, and

consumption rates of fruits vary widely across populations and years (Appendix S2: Fig. S5-10).

This variation drives fluctuations in population growth, including a remarkable population crash

between 2014 and 2015 (Appendix S2: Fig. S11). Growth did not respond to climate, but the

remaining three size-dependent vital rates correlated negatively with average annual temperature

(Fig. 1, Appendix S2: Fig. S7, Fig. S9-10).

Population responses to climate

There is a sharp decline in population growth rate with increasing mean annual temperature (Fig.

2A). While the slope of the relationship between population growth rate and annual temperature

is similar for all populations, these vary in their mean vital rates (i.e., their intercepts and slopes,

Appendix S2: Fig. S7-10). If the temperatures remained at the 1990-2018 mean (12° C), three of

the four smallest populations would be expected to decline. At these temperatures, the remaining

four populations, including the three largest, are projected to grow. On the other hand, in a

scenario where mean annual temperature increases by 1° C, all populations are projected to

decline (Fig. 2A).

Perturbation analyses indicate that the decline in population growth rate with increasing

temperature is primarily due to the effects of temperature on plant survival (Fig. 2B). Two

factors explain the high contribution of plant survival on the change in population growth rate.

First, population growth rate is very sensitive to changes in survival (Appendix S8: Fig. S12),

and second, survival responds dramatically to increasing temperature (Fig. 2A).

Forecast

Population trajectories indicate that if temperature remained at its 1990-2018 average, this

endangered plant species would double its population number in the next 30 years. However,

with a 1° C increase in temperature, there would be a 90% average reduction in individuals (Fig.

3). This scenario is conservative, as more dramatic increases in temperature are projected for this

region in the next 30 years (Wang et al. 2016, n.d., Vose et al. 2017).

DISCUSSION

Our results demonstrate that climate change is having profound negative effects on the current

and future population growth rates of *L. tidestromii*. Within the next 30 years, climate change

will result in high probabilities of extinction for four out of the seven populations we analyzed.

These high extinction probabilities add to the substantial decrease in total population size. As our

analysis considers 7 of the 15 extant populations of the species, including the largest population,

it is clear that climate change will substantially deteriorate the conservation status of *L.*

tidestromii.

We hypothesize that our results could reflect a general pattern among species evolved in coastal

dune habitats experiencing low variation in annual temperatures. Low variation in climatic

variables, both temporally and spatially (e.g. across topographic gradients), should decrease the

ability of species to respond to climatic changes. This expectation is justified because climatic

variation in time and space selects for high genetic variability and high phenotypic plasticity

(Nadeau et al. 2017). Dune plants of the pacific coast such as *L. tidestromii* experience low

variability in annual temperature, and do not occur across large topographic gradients. The

standard deviation in annual temperatures at our study populations is only 0.43° Celsius, which

is below the 5th percentile of those found across the conterminous United States. Low variation

in annual temperatures should be common among Western temperate coasts as a result of

Westerly currents. Moreover, dune plants occur only at sea level, and therefore they are not

adapted to relatively small topographic gradients. Given these considerations, it might be fruitful

to carry out comparative tests on the climatic sensitivity of taxa adapted to dune habitats of mid-

latitude Western coasts.

Alternatively, our results for *L. tidestromii* could reflect an isolated case, because climate change

can threaten species in any biome as long as their physiological tolerances are exceeded. For

example, *L. tidestromii* could be vulnerable because it thrives only within a narrow range of

temperatures (Jenouvrier, 2013). One of the reasons for a narrow temperature niche is that the

geographic distribution of rare species like *L. tidestromii* could be much smaller than its potential

(Svenning and Skov 2004). If *L. tidestromii* did not fill its potential range, its populations could

fall, by sheer luck, at sites whose average climate is close to the upper limits of its thermal

physiological tolerance. An additional hypothesis for such high climate sensitivity is that the

phylogeny of *L. tidestromii* might make this species prone to physiological stress from heat. The

productivity of legume crops is very sensitive to heat stress (Liu et al. 2019), a trait which might

be shared with wild legumes.

Our results show three advantages of demographic approaches compared to species distribution

modeling. First the temporally explicit projections of our demographic models provide highly

relevant information to conservation planning and prioritization (Fig. 3). Second, demographic

analysis holds mechanistic insights. In our case, we found that the population growth rate of *L.*

tidestromii responds to climate mainly through survival. Hence, management actions that

increase average individual survival might ameliorate the risks imposed by climate change.

Finally, it would be hard or impossible to construct an SDM for *L. tidestromii*. This species

occurs in just three locations which differ in average temperature by just 0.12 degrees Celsius.

Moreover, the minimum number of sites needed to build an SDM free of statistical artifacts is

above 10 (Proosdij et al. 2016). Hence, long-term population or demographic data might be a

preferable way to devise climate change forecasts for species that have a geographically

restricted range. Some authors suggest that establishing a link between demographic rates and

climatic drivers might require 20 to 25 years of data (Teller et al. 2016, Tenhumberg et al. 2018).

Instead, our results show that such long data collection might not be needed when a study period

spans a large range of climate anomalies (Appendix S1: Fig. S4).

To our knowledge, the evidence we present here is the first example suggesting that climate

change should be included as a threat in the status for a species listed under the Endangered

Species Act. To date, climate change has been considered in listing actions for four animal

species occurring in arctic, alpine, or arid biomes. These animals are the polar bear, the

American pika, the American wolverine, and the Gunnison sage-grouse (Blumm and Marienfeld

2013, USFWS 2015). While our results on *L. tidestromii* could be an isolated case, they suggest

that the extinction threat posed by climate change might be overlooked in temperate biomes.

Thus, we hope that our study serves as a model for other similar forecasting efforts using long-

term monitoring data.

ACKNOWLEDGMENTS

Funding for this research was provided by the NSF (DEB-0743731), the Alexander von

Humboldt foundation, and Washington University in St. Louis. We thank Valentin Ştefan for

support with data processing. We thank Dr. Ben Becker at Point Reyes National Seashore for

lodging and extensive logistical support, and Lorraine Parsons at Point Reyes National Seashore

for long-term collaboration on the monitoring of this species, and sharing site-based knowledge.

We thank numerous students for assisting with field data collection, in particular Emily

Dangremond and Melissa Patten who worked at least three summers on this project.

REFERENCES

Baldwin, B. G., D. H. Goldman, D. J. Keil, R. Patterson, T. J. Rosatti, and D. H. Wilken. 2012.
The Jepson Manual: Vascular Plants of California. University of California Press.

Barbier, E. B. 2015. Valuing the storm protection service of estuarine and coastal ecosystems.
Ecosystem Services 11:32–38.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models
Using {lme4}. Journal of Statistical Software 67.

Beguería, S., and S. Vicente-Serrano. 2017. SPEI: Calculation of the Standardised Precipitation-
Evapotranspiration Index.

Blumm, M. C., and K. B. Marienfeld. 2013. Endangered Species Act Listings and Climate
Change: Avoiding the Elephant in the Room Symposium. Animal Law:277–310.

Burnham, K. P., and D. R. Anderson. 2002. Model Selection and Multimodel Inference: A
Practical Information-Theoretic Approach. Second edition. Springer-Verlag, New York.

Campbell, D. R. 2019. Early snowmelt projected to cause population decline in a subalpine
plant. Proceedings of the National Academy of Sciences 116:12901–12906.

Caswell, H. 2001. Matrix population models. Massachusetts: Sinauer Associates.

CPC. (n.d.).
https://origin.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ONI_v5.php.htt
ps://origin.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ONI_v5.php.

Daly, C., R. P. Neilson, and D. L. Phillips. 1994. A Statistical-Topographic Model for Mapping
Climatological Precipitation over Mountainous Terrain. Journal of Applied Meteorology
33:140–158.

Dangremond, E. M., E. A. Pardini, and T. M. Knight. 2010. Apparent competition with an

invasive plant hastens the extinction of an endangered lupine 91:11.

Díaz, S., J. Settele, E. Brondízio, H. T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, K.

Brauman, R. T. Watson, I. A. Baste, A. Larigauderie, P. Leadley, U. Pascual, B. Baptiste,

S. Demissew, L. Dziba, G. Erpul, A. Fazel, M. Fischer, A. María, M. Karki, V. Mathur,

T. Pataridze, I. S. Pinto, M. Stenseke, K. Török, and B. Vilá. 2019. Summary for

policymakers of the global assessment report on biodiversity and ecosystem services of

the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem

Services:45.

Doak, D. F., and W. F. Morris. 2010. Demographic compensation and tipping points in climate-

induced range shifts. Nature 467:959–962.

Ehrlén, J., and W. F. Morris. 2015. Predicting changes in the distribution and abundance of

species under environmental change. Ecology Letters 18:303–314.

Elith, J., and J. R. Leathwick. 2009. Species Distribution Models: Ecological Explanation and

Prediction Across Space and Time. Annual Review of Ecology, Evolution, and

Systematics 40:677–697.

Ellner, S. P., D. Z. Childs, and M. Rees. 2016. Data-driven Modelling of Structured Populations:

A Practical Guide to the Integral Projection Model. Springer International Publishing.

Evens, J. G. 2008. natural history of the Point Reyes Peninsula. University of California Press.

Feagin, R. A., D. J. Sherman, and W. E. Grant. 2005. Coastal erosion, global sea-level rise, and

the loss of sand dune plant habitats. Frontiers in Ecology and the Environment 3:359–

364.

Glenn, E. M., R. G. Anthony, and E. D. Forsman. 2010. Population trends in northern spotted

owls: Associations with climate in the Pacific Northwest. Biological Conservation

143:2543–2552.

Gneiting, T., and A. E. Raftery. 2007. Strictly Proper Scoring Rules, Prediction, and Estimation.

Journal of the American Statistical Association 102:359–378.

Hunter, C. M., H. Caswell, M. C. Runge, E. V. Regehr, S. C. Amstrup, and I. Stirling. 2010.

Climate change threatens polar bear populations: a stochastic demographic analysis.

Ecology 91:2883–2897.

Iler, A. M., A. Compagnoni, D. W. Inouye, J. L. Williams, P. J. CaraDonna, A. Anderson, and T.

E. X. Miller. 2019. Reproductive losses due to climate change-induced earlier flowering

are not the primary threat to plant population viability in a perennial herb. Journal of

Ecology 107:1931–1943.

IPCC. 2014. Summary for policymakers. In: Climate Change 2014: Impacts, Adaptation, and

Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to

the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.

Cambridge University Press, Cambridge, UK and New York, NY, USA.

Jenouvrier, S. 2013. Impacts of climate change on avian populations. Global Change Biology

19:2036–2057.

Jenouvrier, S., H. Caswell, C. Barbraud, M. Holland, J. Strœve, and H. Weimerskirch. 2009.

Demographic models and IPCC climate projections predict the decline of an emperor

penguin population. Proceedings of the National Academy of Sciences 106:1844–1847.

Jonzén, N., T. Pople, J. Knape, and M. Sköld. 2010. Stochastic demography and population

dynamics in the red kangaroo Macropus rufus. Journal of Animal Ecology 79:109–116.

Keenan, T., J. M. Serra, F. Lloret, M. Ninyerola, and S. Sabate. 2011. Predicting the future of

forests in the Mediterranean under climate change, with niche- and process-based

models: CO_2 matters! Global Change Biology 17:565–579.

Lande, R. 1993. Risks of Population Extinction from Demographic and Environmental

Stochasticity and Random Catastrophes. The American Naturalist 142:911–927.

Liu, Y., J. Li, Y. Zhu, A. Jones, R. J. Rose, and Y. Song. 2019. Heat Stress in Legume Seed

Setting: Effects, Causes, and Future Prospects. Frontiers in Plant Science 10.

Luger, M. I. 1991. The economic value of the coastal zone:23.

Martínez, M. L., A. Intralawan, G. Vázquez, O. Pérez-Maqueo, P. Sutton, and R. Landgrave.

2007. The coasts of our world: Ecological, economic and social importance. Ecological

Economics 63:254–272.

Paniw, M., N. Maag, G. Cozzi, T. Clutton-Brock, and A. Ozgul. 2019. Life history responses of

meerkats to seasonal changes in extreme environments. Science 363:631–635.

Pardini, E. A., K. E. Vickstrom, and T. M. Knight. 2015. Early Successional Microhabitats

Allow the Persistence of Endangered Plants in Coastal Sand Dunes. PLOS ONE

10:e0119567.

Pereira, H. M., P. W. Leadley, V. Proença, R. Alkemade, J. P. W. Scharlemann, J. F. Fernandez-

Manjarrés, M. B. Araújo, P. Balvanera, R. Biggs, W. W. L. Cheung, L. Chini, H. D.

Cooper, E. L. Gilman, S. Guénette, G. C. Hurtt, H. P. Huntington, G. M. Mace, T.

Oberdorff, C. Revenga, P. Rodrigues, R. J. Scholes, U. R. Sumaila, and M. Walpole.

2010. Scenarios for Global Biodiversity in the 21st Century. Science 330:1496–1501.

Proosdij, A. S. J. van, M. S. M. Sosef, J. J. Wieringa, and N. Raes. 2016. Minimum required

number of specimen records to develop accurate species distribution models. Ecography

39:542–552.

Sæther, B.-E., J. Tufto, S. Engen, K. Jerstad, O. W. Røstad, and J. E. Skåtan. 2000. Population

Dynamical Consequences of Climate Change for a Small Temperate Songbird. Science

287:854–856.

Schurr, F. M., J. Pagel, J. S. Cabral, J. Groeneveld, O. Bykova, R. B. O'Hara, F. Hartig, W. D.

Kissling, H. P. Linder, G. F. Midgley, B. Schröder, A. Singer, and N. E. Zimmermann.

2012. How to understand species' niches and range dynamics: a demographic research

agenda for biogeography. Journal of Biogeography 39:2146–2162.

Svenning, J.-C., and F. Skov. 2004. Limited filling of the potential range in European tree

species. Ecology Letters 7:565–573.

Teller, B. J., P. B. Adler, C. B. Edwards, G. Hooker, and S. P. Ellner. 2016. Linking demography

with drivers: climate and competition. Methods in Ecology and Evolution 7:171–183.

Tenhumberg, B., E. E. Crone, S. Ramula, and A. J. Tyre. 2018. Time-lagged effects of weather

on plant demography: drought and Astragalus scaphoides. Ecology 99:915–925.

Tews, J., and F. Jeltsch. 2004. Modelling the impact of climate change on woody plant

population dynamics in South African savanna. BMC Ecology 4:17.

USFWS. 1998. Seven coastal plants and the Myrtle's silverspot butterfl y recovery plan.

USFWS, Portland, Oregon, USA.

USFWS. 2009. Lupinus tidestromii (Clover lupine) 5-Year Review: Summary and Evaluation.

Sacramento Fish and Wildlife Office, Sacramento, California, USA.

USFWS. 2015. Polar Bear (Ursus maritimus) Conservation Management Plan, Draft. U.S. Fish

and Wildlife, Region 7, Anchorage, Alaska.

USGCRP. 2018. Fourth National Climate Assessment. U.S. Global Change Research Program,

Washington, DC, USA.

Vicente-Serrano, S. M., S. Beguería, and J. I. López-Moreno. 2009. A Multiscalar Drought Index

Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index.

Journal of Climate 23:1696–1718.

Vose, R. S., D. R. Easterling, K. E. Kunkel, A. N. LeGrande, and M. F. Wehner. 2017.

Temperature Changes in the United States. Climate Science Special Report: Fourth

National Climate Assessment, Volume I. U.S. Global Change Research Program.

Wang, T., A. Hamann, D. Spittlehouse, and C. Carroll. 2016. Locally Downscaled and Spatially

Customizable Climate Data for Historical and Future Periods for North America. PLOS

ONE 11:e0156720.

Wang, T., A. Hamann, D. Spittlehouse, and C. Carroll. (n.d.). ClimateNA_MAP -- An

Interactive Plantform for Visualization and Data Access. http://www.climatewna.com/.

Figure 1. The effect of plant size and mean annual temperature (when relevant) on survival (A),

growth (B), the probability of flowering (C), and the production of racemes (D). The dots in the

plots showing survival (A) and probability of flowering (C) represent the average proportion of

individuals surviving and flowering, respectively, within 10 equally spaced intervals of

individual log sizes. Shaded areas show the 95% confidence intervals of the mean model

predictions. For the three climate-dependent vital rates, we show credible intervals of vital rates

responding to the average yearly temperature observed between 1990 and 2018 (12° Celsius, in

blue), and the highest average yearly temperature observed during the same period (13° Celsius,

in orange).

Figure 2. Long-term stochastic population growth rate (λ_s) as a function of mean annual

temperature for each of the seven *L. tidestromii* populations (A), and the partial effect of each

climate-dependent vital rate on the λ_s (B). The average yearly temperature observed between

1990 and 2018 was 12° Celsius; the average temperature during the study period was 11.5°

Celsius.

Figure 3. Projected population sizes of *L. tidestromii* assuming historical (12° Celsius) and

future average annual temperatures (13° Celsius). Polygons show the lower 2.5th and upper 97.5th

percentile of 1000 separate projections. The upper left panel shows the population sizes of all

sites at Point Reyes National Seashore; the remaining panels show the population sizes at each

one of the seven sites separately. Each simulation was initiated using the population sizes

observed in 2018.

538   **APPENDICES**
539
540   **Appendix S1: Model system figures**
541

543   **Figure S1**. Map of the seven study populations of *L. tidestromii* at Point Reyes National
544   Seashore, Marin County, California, USA. The entire range of this species includes two
545   additional clusters of populations located in Sonoma and Monterey county.
546

547
548 **Figure S2**. Relationship between the number of branches counted on *L. tidestromii* plants, and
549 their area. The black line represents the mean model prediction of the quadratic linear model that
550 describes the relationship between area and the number of branches. The grey area represents the
551 95% confidence interval of the mean model prediction. The color of dots refers to data from six
552 separate populations.
553

554
Figure S3. Life cycle model for *L. tidestromii*. This life cycle refers to the IPM described in
Appendix S6 (Eq. 7-9). The IPM separates the population in three discrete stages: continuously
sized plants (*plants*), seeds that germinate after one winter (*B1*), and seeds that germinate after
two winters (*B2*). These stages are modeled, respectively, by equation 9, 7, and 8 in Appendix
S6. To minimize the use of space, in this figure π corresponds to $(1-a)(1-c)\iota\kappa(1-\delta)$ in
Equations 7-9.
561

562
563 **Figure S4**. Annual climate values of temperature, precipitation, Oceanic Niño Index (ONI), and
564 standardized aridity index (SPEI) between 1990 and 2018. The dashed lines denote the beginning
565 of *L. tidestromii* demographic censuses.
566
567

568 **Appendix S2: Vital rates figures**
569

570
571 **Figure S5**. Average abortion rate of inflorescences by year and population.
572

573
574 **Figure S6**. Average consumption rate by year and population.
575

576
577 **Figure S7**. Survival to time t+1 of individuals of size log(Z) at time *t*, by transition year
578 (columns) and population (rows). Red dots show the observed proportion of surviving
579 individuals in ten equally spaced size log(Z) intervals at time *t*. Grey lines show the average size-
580 dependent survival probability predicted by the generalized linear mixed model.
581

582
583 **Figure S8**. Individual log size (Z) at time t+1 as a function of individual log size (Z) at time t, by
584 transition year (columns) and population (rows). Red dots show individual data points, and grey
585 lines show the prediction of the linear mixed model.
586

587
588 **Figure S9**. Flowering probability at time *t* based on the log size of individuals (Z) at time *t*, by
589 year (columns) and population (rows). Red dots show the observed proportion of flowering
590 individuals in ten equally spaced intervals of log sizes at time *t*. Grey lines show the average
591 size-dependent flowering probability predicted by the generalized linear mixed model.
592

593
594 **Figure S10**. Production of inflorescences by reproductive individuals at time t as a function of
595 individual log size (Z) at time t, by year (columns) and population (rows). Red dots show
596 individual data points, and grey lines show the predictions of the generalized linear mixed model.
597

598
599 **Figure S11**. Asymptotic population growth rates (λ) based on transition year and population. λ
600 are calculated using year- and population- specific integral projection models. λ values prior to
601 the 2008-2009 transition are available for two sites only (AL and ATT).
602

603 **Appendix S3: Calculation of the germination adjustment factor**
604
We estimated a germination adjustment factor, δ, by combining information on germination
trials, on seed production, and on the number of new seedlings. We used data from "known
population counts", which refers to the populations and transition years for which we have
counted the total number of individuals and seedlings. First, we estimated the expected number
of seeds produced at each population, p, and year, t (*estimated seeds*$_{pt}$). To estimate the number
of seeds, we used year- and population-specific consumption and abortion, and constant values
of fruits per inflorescence and seeds per fruit data. We then used the ratio of observed
seedlings$_{pt+1}$ to the estimated seeds$_{pt}$ to calculate the recruitment adjustment factor δ:
613

614
$$\frac{observed\ seedlings_{pt+1}}{estimated\ seeds_{pt}} = g_1(1 - \delta_p)$$

615
616 Rearranging,

617
$$\delta_p = \frac{g_1 - (\frac{observed\ seedlings_{pt+1}}{estimated\ seeds_{pt}})}{g_1}$$

618 Where g_1 is the proportion of seeds germinating in the first year (after one winter) in the
619 germination experiment. In our population models, we apply δ_p to g_2 and g_3 because calculating
620 δ_p using g_1 provides a conservative estimate of unobserved seed and seedling mortality. Our δ_p
621 is conservative because the observed seedlings at $t+1$ emerge from dispersed seeds (g_1) and from
622 the seed bank (g_2, g_3). We modified germination rates g_1, g_2, and g_3 used in the IPM using δ_p as:

623
$$g_t\left(1 - \delta_p\right).$$

We could only calculate δ_p for the five populations with known population counts. For
population BR, we applied the mean of δ_p from BS and DR; for AL, we applied the mean of δ_p
from ATT and Pop9. We used these estimates due to the geographic proximity of BR and AL to
these smaller populations for which δ could be quantified. We could only calculate δ_p for the
five populations with known population counts. For population BR, we applied the mean of δ_p
from BS and DR; for AL, we applied the mean of δ_p from ATT and Pop9. We used these
estimates due to the geographic proximity of BR and AL to these smaller populations for which
δ could be quantified.
632
633

Appendix S4: Vital rate models

635 The baseline model structure for our models of size-dependent vital rates was

636
$$\text{link}\left(\widehat{\psi_{itp}}\right) = \gamma_t + \gamma_p + \beta_t \, log(Z_i) + \beta_p \, log(Z_i), \qquad (1)$$

where $\hat{\psi}$ is the mean prediction of the vital rate model for individual i, in year t, at population p,
Z is the size of individual i, the γ coefficients are normally distributed random intercepts, the β
coefficients are normally distributed random slopes of log size, and *link* is the link function for
the vital rate being modelled. The link function depends on the type of response variable. The
models on survival (S) and flowering (F) used a logit link, because the response variable is
Bernoulli distributed:

643
$$S_{itp} \sim Bernoulli(\hat{S}_{itp}), \qquad (2)$$

644
$$F_{itp} \sim Bernoulli(\hat{F}_{itp}). \qquad (3)$$

645 The best survival model was a cubic function of log(size), so that:

646
$$\text{logit}\left(\widehat{S_{itp}}\right) = \gamma_t + \gamma_p + \beta_t \, log(Z_i) + \beta_p \, log(Z_i) + \beta_2 \, log(Z_i)^2 + \beta_3 \, log(Z_i)^3. \quad (4)$$

647 The number of inflorescences (R) used a log link, because this response variable follows a
648 Poisson distribution:

649
$$R_{itp} \sim Poisson(\hat{R}_{itp}). \qquad (5)$$

650 The growth model used an identity link: that is, it did not use a link function:

651
$$G_{itp} \sim Normal(\hat{G}_{itp}, \sigma). \qquad (6)$$

Finally, we used these models to select the best climatic predictor through a leave-one-year-out
cross-validation. We tested four climatic predictors: temperature, precipitation, Oceanic Niño
Index (ONI), and the standardized precipitation evapotranspiration index (SPEI, see main text).
In these cross-validations, Equation 1 and 4 included the term $\beta_c C_t$, where C was one of the
climatic predictors, and β_c was its slope. In such model, the linear predictor changes based on the
year t only, because climatic drivers are the same across populations p and individuals i.

Appendix S5: Model selection tables

Table S1. Cross-validation results based on log-likelihood. The models with higher support appear first, those with the least support are last. Model "null" does not contain a climatic predictor. We compare models with four annual climatic anomalies: temperature (tmp), precipitation (ppt), Oceanic Niño Index (oni), and the standardized precipitation evapotranspiration index (spei). Moreover, we compare models with climatic predictors referred to two years: the year of the demographic census (t) and that preceding it ($t-1$).

Survival		Growth		Flowering		Racemes	
Model	log_lik_sum	Model	log_lik_sum	Model	log_lik_sum	Model	log_lik_sum
tmp_t	8320.713	null	9643.271	tmp_t	4860.269	tmp_t	20538.58
null	8339.689	ppt_{t-1}	9683.304	null	5594.391	null	20863.18
$spei_{t-1}$	8378.673	$spei_t$	9690.057	oni_t	5794.024	ppt_t	20886.95
oni_t	8382.177	ppt_t	9696.37	tmp_{t-1}	5804.951	$spei\_to$	21036.85
tmp_{t-1}	8385.776	$spei_{t-1}$	9697.36	$spei_{t-1}$	5874.919	tmp_{t-1}	21236.65
ppt_t	8408.783	oni_{t-1}	9697.919	ppt_{t-1}	5876.703	oni_t	21281.96
oni_{t-1}	8439.169	tmp_{t-1}	9700.742	oni_{t-1}	5882.694	ppt_{t-1}	21292.39
$spei_t$	8442.731	oni_t	9730.185	$spei_t$	5888.286	$spei_{t-1}$	21326.18
ppt_{t-1}	8494.641	tmp_t	9808.604	ppt_t	5890.526	oni_{t-1}	21336.45

Table S2. Cross-validation results based on root mean squared error. The models tested are the same as those described in Table S1.

Survival		Growth		Flowering		Racemes	
Model	rmse	Model	rmse	Model	rmse	Model	rmse
tmp_t	0.44768	null	0.979553	tmp_t	0.354357	tmp_t	7.82832
null	0.448003	ppt_{t-1}	0.984409	null	0.374067	null	7.918863
$spei_{t-1}$	0.449193	$spei_t$	0.985296	oni_t	0.381321	ppt_t	7.954278
tmp_{t-1}	0.449645	ppt_t	0.986036	ppt_{t-1}	0.381385	tmp_{t-1}	7.961484
oni_t	0.449673	$spei_{t-1}$	0.986127	tmp_{t-1}	0.381954	oni_t	8.001534
ppt_t	0.4504	oni_{t-1}	0.986247	$spei_{t-1}$	0.382086	$spei_t$	8.014872
$spei_t$	0.451516	tmp_{t-1}	0.986565	$spei_t$	0.382173	oni_{t-1}	8.052406
oni_{t-1}	0.451523	oni_t	0.990133	ppt_t	0.382333	ppt_{t-1}	8.230619
ppt_{t-1}	0.453314	tmp_t	0.999683	oni_{t-1}	0.384076	$spei_{t-1}$	8.258191

Appendix S6: Integral projection model

Our IPM projects the number of z-sized *L. tidestromii* individuals, represented by vector $n(z)$, to the number of z-sized individuals next year, represented by vector $n(z')$. This population is also composed of seeds that germinate one year after the projection interval, represented by scalar B_1, and two years after, represented by scalar B_2. The dynamics of the population are given by

$$B_{1,t+1} = (1 - a)(1 - c)\iota\kappa g_2(1 - \delta)\int_L^U p_b(z; \beta_c)b(z; \beta_c)n(z,t)dz + B_{2,t} \qquad (7)$$

$$B_{2,t+1} = (1 - a)(1 - c)\iota\kappa g_3(1 - \delta)\int_L^U p_b(z; \beta_c)b(z; \beta_c)n(z,t)dz \qquad (8)$$

$$n(z', t + 1) = \eta(z')B_{1,t} + \eta(z')(1 - a)(1 - c)\iota\kappa g_1(1 - \delta)\int_L^U p_b(z; \beta_c)b(z; \beta_c)n(z,t)dz +$$

$$\int_L^U s(z; \beta_c)G(z', z)n(z,t)dz. \qquad (9)$$

We show a schematic representation that links these equations to the life cycle of *L. tidestromii* in Appendix S1, Fig. S3. These equations contain four vital rates that depend directly on the size of individuals, z, three of which depend on temperature anomaly, β_c, while the remaining demographic rates are size- and climate-independent. Three vital rates are size- and climate-dependent: $p_b(z;\beta_c)$ is the size- and climate-dependent probability of flowering, $b(z;\beta_c)$ is the size- and climate-dependent production of inflorescences, $s(z;\beta_c)$ is the size- and climate-dependent probability of survival, and $G(z',z)$ is size-dependent growth. The size- and climate-independent vital rates mostly refer to the transition from the number of inflorescences to the number of establishing seedlings. In particular, a is the population- and year-specific abortion rate, c is the population- and year-specific consumption rate (or pre-dispersal predation), ι is the average number of fruits produced per inflorescence, κ is the average number of seeds produced per fruit, δ is the recruitment adjustment factor (Appendix S3), and g values refer to germination rates. g_0 is the fraction of seeds that germinate before the end of the first transition, g_1 is the fraction of seeds that germinate before the end of the second transition, and g_2 is the fraction of seeds that germinate before the end of the third transition. Finally, $\eta(z')$ is the size distribution of seedlings emerging by the end of the transition.

We used the IPMs to compute how climate affects λ_s, the long-term stochastic population growth rate, and how each climate-dependent vital rate contributes to these changes in λ_s. We computed the stochastic population growth rate, λ_s, by projecting the population numbers of each one of our seven population 50000 times. We started with a population of only one individual, discarding the first 1000 projections to allow populations to reach a stable size distribution. These stochastic simulations kept a constant average annual temperature, and randomly varied the year-specific conditions (parameters γ_t and β_t in equation 1). We calculated λ_s as:

$$\log(\lambda_s) = E\left[\log\left(\frac{N_{t+1}}{N_t}\right)\right] \qquad (10)$$

Where E calculates the mean, and N is the total population size at the end ($t+1$) and at the beginning (t) of a transition. We repeated these projections across the whole range of observed climate anomalies we observed during our study period. These temperatures go from an average annual temperature of about 11° Celsius to an average annual temperature of 13°.

Appendix S7: Individual based model

Our individual based model simulated each probabilistic event in the life cycle of *L. tidestromii* (Appendix S1: Fig. S3) using an appropriate statistical distribution. Our probabilistic events were linked to survival, growth, probability of flowering, production of inflorescences, the abortion of inflorescences, the consumption of inflorescences, and the germination of seeds. Starting from the production of flowers, each individual flowered based on a Bernoulli process depending on its individual size, so that:

$$Flower \sim Bernoulli(\ p_b(z; \beta_c)\ ) \qquad (11)$$

Where *Flower* is either a 0, if an individual failed to flower, or 1, if an individual flowered. $p_b(z;\beta_c)$ shows that the probability of reproduction depends on individual size, *z,* and on the climate anomaly. If an individual reproduces, its production of inflorescences depends on a negative binomial distribution, so that

$$RacemesN \sim Poisson(\ b(z; \beta_c), \theta\ ) \qquad (12)$$

Which is also a size- and climate- dependent process. The number of inflorescences would decrease because of abortion following a binomial process

$$RacemesN \sim Binomial(RacemesN, (1 - a)\ ) \qquad (13)$$

where 1-*a* is the probability of an inflorescence not aborting. The remaining inflorescences could still be consumed by predators,

$$RacemesN \sim Binomial(RacemesN, (1 - c)\ ) \qquad (14)$$

where 1-*c* is the probability of not being consumed (the probability of experiencing pre-dispersal predation). The seeds produced by the surviving inflorescences on each individual could then germinate based on a binomial process, so that:

$$SeedlingN \sim Binomial(seeds, g_t(1 - \delta)) \qquad (15)$$

Where SeedlingN is the number of seedlings emerging in each population, and g_t is one of the germination rates associated with seeds that germinate in the year of the transition (g_0), the year after (g_1), and two years after (g_2). Finally, individuals could survive based on a size- and climate-dependent Bernoulli process

$$Surv \sim Bernoulli(\ s(z; \beta_c)\ ) \qquad (16)$$

and, in case they survived, they could grow based on a normal process:

$$z' \sim Normal(\ G(z', z), \sigma\ ). \qquad (17)$$

 Appendix S8: Elasticity analysis

744
Figure S12. Elasticity of asymptotic population growth rate (λ) to the parameters of the IPM.
Parameters refer to the average values across years and populations. The elasticity is a
proportional sensitivity, meaning that it quantifies by what percentage λ will change after a
percent change in each parameter. Thus, elasticity facilitates the comparison among parameters.
The parameter names for size-dependent processes show the effect of annual temperature
anomaly (_clim), the intercept of linear models (_b0), and the linear (_b1), quadratic (_b2), and
cubic (_b3) effects of size. Parameters "recr_size" and "recr_sd" refers to the mean size and
standard deviation of recruits. The remaining parameter names are identical to those in equations
7-9.