Title Page

Title: An extension of Shannon's entropy to explain taxa diversity and human

diseases

Author list and full affiliations:

Farzin Kamari[*1,2], MD, MPH; Sina Dadmand[2,3], PharmD.

[1]Neurosciences Research Centre, Tabriz University of Medical Sciences, Tabriz, Iran.

[2]Synaptic ProteoLab, Synaptic ApS, Skt Knuds Gade 20, 5000 Odense C, Denmark.

[3]Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

[*]**Corresponding author:** Farzin Kamari   10

Email: kamari.farzin@gmail.com

Number of words in the main text (excluding Abstract, Methods, References, and

Figure/Table Legends): 3,934

Number of words in the Abstract: 148

Keywords: origin of diseases/ protein-protein interaction/ Shannon's entropy/ taxonomic

classification/ tree of life

Abstract

In this study, with the use of the information theory, we have proposed and proved a mathematical theorem by which we argue the reason for the existence of human diseases. To introduce our theoretical frame of reference, first, we put forward a modification of Shannon's entropy, computed for all available proteomes, as a tool to compare systems complexity and distinguish between the several levels of biological organizations. We establish a new approach to differentiate between several taxa and corroborate our findings through the latest tree of life. Furthermore, we found that human proteins with higher mutual information, derived from our theorem, are more prone to be involved in human diseases. We further discuss the dynamics of protein network stability and offer probable scenarios for the existence of human diseases and their varying occurrence rates. Moreover, we account for the reasoning behind our mathematical theorem and its biological inferences.

Introduction

The term 'entropy' was originally introduced by Rudolf Clausius in thermodynamics more than one and a half centuries ago (Clausius, 1864). Entropy is predominantly known as a measure of the disorder and uncertainty in dynamic systems (Bailey, 2009; Ghahramani, 2006). In information theory, entropy, also known as Shannon's entropy, is defined as the average minimum rate at which information is produced or predicted in an uncertain stochastic setting (Shannon, 1948). In recent decades, information theory has been vastly applied in many fields of science (Andrews et al., 2015). Biology is of no exception, but compared to other areas, the applications of information theory in biological sciences have been indeed limited (Battail, 2013). More importantly, medical sciences lack any use of information theory in daily practice. The applications of information theory in molecular biology have been mostly focused on genome sequence analysis (Vinga, 2013). To date, no study has investigated the evolutionary nature of human diseases using information theory.

The backbone of evolution is random genetic mutations being selected according to the natural environment. So what has been encountered in nature after some 3.5 billion years of life history is a 'selected randomness'. This is the reason why we believe information theory can be a perfect language to understand life – i.e., this selected randomness. In the literature, single nucleotide polymorphisms (SNPs) accounting for the main portion of this randomness have been associated with inherited disease susceptibility (Bodmer & Bonilla, 2008; Z. Wang & Moult, 2001). However, such approaches have only focused on the genome investigation and, in most part, neglected the human proteome and the protein-protein interactions (PPIs). PPIs are the leading cause of cellular metabolic processes. They are induced-fit physical contacts between macromolecules of proteins allowing the cellular function (Changeux & Edelstein, 2011; Keskin et al., 2008; Koshland Jr, 1995). In order to employ information

theory in medical sciences, it would be necessary to investigate diseases in detail considering

their molecular networks and PPIs. We believe evolutionary evidence interpreted by

stochastic information analysis can provide substantial help in understanding diseases of

living organisms.

In this study, to understand the nature of human diseases, we have focused on human

interactome and available proteomes of living organisms. To avoid confusion, by 'human

diseases' we only refer to non-communicable diseases in human with at least one reported

genetic basis. Also, as the term 'proteome' has sometimes referred to all proteins of a cell or

a tissue in the literature, it is to be noted that in this article, the term 'proteome' will refer to

the complete set of proteins that *can be* expressed by an *organism*. Because proteomes are

functional representatives of the 'expressed genome' of organisms, we have used them as the

means of our investigation. We have used Shannon's entropy as a retrograde approach to

trace ~180 million proteins with more than 61 billion amino acids through the tree of life and

investigated the trends of complexity among organisms. We have shown that this

methodology agrees with the classification of phyla and may be used as a new tool in

taxonomy. Also, using our new mathematical theorem presented in the Methods section, we

have focused on *Homo sapiens'* PPI network and discussed potential clinical applications in

the practice of medicine. We argue why there are only the diseases we know, and not others,

and discuss why some diseases are more prevalent. We also elaborate on the reasonable links

between our mathematical theory, Shannon's entropy, the evolution of taxa, and human

diseases.

Results

CAIR comparisons among taxonomic groups

Calculated Average Information per Residue (CAIR) was calculated (see Methods) for all proteomes available at the UniProt database until April 2020. Nearly 180 million proteins with more than 61 billion amino acids were analysed to classify ~29,000 organisms in 92 phyla. Table 1 shows CAIRs of the most popular proteomes and model organisms (for all ~29k proteomes, see Supplementary Table 1). The minimum CAIR of an organism is that of *Zinderia insecticola* (0.8247) and the maximum is of *Ciona savignyi* (0.9449). The mean ± standard deviation considering all organisms is 0.9210 ± 0.0129 with a median (interquartile range) of 0.9250 (0.0160). Having performed a literature review of articles published no later than April 2020, we have drawn the most updated tree of life for UniProt taxonomic lineage data (Figure 1A). For each bifurcation point on the tree, we tested if two sides of the bifurcation have developed divergent CAIRs. Figure 1B illustrates how CAIR divergence is present through the different lineages of taxonomy. On all bifurcation points of Figure 1A, a number is written whose respective statistical test results are demonstrated in Figure 1B with two half violin plots for upper and lower sides of the bifurcation and their box-and-whisker plots. Since the groups were negatively skewed, unbalanced, and heteroscedastic, their difference was investigated via the two-sided Brunner-Munzel statistical test (Neuhäuser & Ruxton, 2009). It is noteworthy that groups with ten or fewer organisms were excluded from comparisons, as the Brunner-Munzel test is statistically imprecise even with a permutation. Among 56 performed tests, 48 tests demonstrated a significant difference at the point of bifurcation. Interestingly, the bifurcation points of eight insignificant tests are mostly known to be a matter of controversy in the scientific literature (Evans et al., 2019; Spang et al.,

2017). Along with the *p*-value significance, estimated effect sizes (ES) and 95% confidence

intervals (CI) are also reported. For the exact *p*-values of each test, please see Table 2.

CAIR as a means of understanding the behaviour of natural selection

The extent of natural selection's capacity to show bias in favour of selecting a spectrum of

organisms is open to question. Since genetic mutations are known to be generally random, the

CAIR density plot of such a random condition without a selection bias shall turn out to have a

uniform distribution. In a simulation of various random protein systems, we obtained a

similar distribution to the actual CAIR density plot taking into account a negative skewness

of -0.90 (Figure 2). Figure 2A shows the density plot of life and Figure 2B depicts that of our

simulation. Figure 2C, also, shows how these two distributions are similar given a tiny

bandwidth. To test for their similarity of distributions, we also performed two-sample

Kolmogorov-Smirnov tests whose mean *p*-value was 0.40 on 1000 iterations. It is

noteworthy, not unexpected though, that the natural selection is biased toward the organisms

with higher CAIRs. This might have stemmed from the random mutations over the course of

ages as discussed in the next section. Also, natural selection favours more complex and more

unpredictable protein systems, as they can accommodate superior functionalities. Besides, the

density plot of CAIRs possesses one other interesting property. Since there are significant

differences among taxonomic groups of the second hierarchy as shown in Figure 1B, it can

further be expected that all organisms are noticeable on the density plot. Figure 3 shows how

different taxonomic hierarchies are manifested in the density plot of organism CAIRs. On the

'wave of life', members of the succeeding taxonomic ranks are revealed by zooming in on the

preceding taxonomic group. This property of CAIR density suggests an original methodology

to help classifying organisms into various taxa.

Human proteome analysis and the estimation of mutual information for a protein (EMIP)

According to the theorem presented in the Methods section and its biological inferences,

EMIP has been calculated for each human protein entry using its PPI network. Each Swiss-

Prot, i.e. reviewed, entry has been categorized into three groups using the Orphanet database

of diseases. In case of a reported disease(s) related to an entry, all disease point-

prevalence/incidences of the entry (from Orphanet database) are summed up to obtain the

total occurrence of the disease, that is to say, the protein's overall malfunction. Table 3 shows

the narrative data of the human disease categories. As seen in the table, the groups are

unbalanced in size and heteroscedastic which makes conventional statistical analyses

unfavourable. Herein, our results demonstrate how well indicators of diseases can be

correlated to the disease occurrence categories. In the Methods section, we have explained

why such independent variables were candidates of correlation and further statistical

analyses. Figure 4 shows the results of comparisons between disease occurrence categories in

four disease indicators. In each comparison, we have also included gene age categories to test

our hypotheses and biological inferences. The Dunnett-Tukey-Kramer pairwise multiple

comparison test adjusted for unequal variances and unequal sample sizes (Dunnett, 1980) was

performed to test overall comparisons. Results of comparisons show that disease indicators

correlate significantly with disease occurrence categories. Among four indicators, EMIP is

revealed to have the most significant differences between categories, while CAIR was

incongruous. The inconsistency seen in CAIR confirms that the use of Shannon's entropy

alone is not a good enough indicator of disease occurrence categories.

Additionally, since gene ages are presented as ranked data in the literature (Liebeskind et al.,

2016), ranked analysis with an equal number of ranks has been performed to make all five

indicators comparable with one another. Figure 5 shows the Likert plots and rank

comparisons. As illustrated, natural Logarithm of EMIP (LEMIP) is by far better than other

indicators in correlating disease occurrence categories which might be stemmed from its bell-

shaped histogram. Unlike other indicators, the distribution of LEMIP allows it to be ranked

with mean and standard deviations which have been reported in the Table 4. Please refer to

Table 4 for detailed information about the ranks in disease indicators.

As results suggest, EMIP seems to be a superior indicator of human diseases. Calculated

values of disease indicators for all reported diseases (Supplementary Table 2) and all human

proteins (Supplementary Table 3) are available in the Supplementary Material. In a nutshell,

we have presented 16 human proteins with the highest EMIPs in Table 5. It is noteworthy

that high-EMIP proteins are more susceptible to have diseased networks and are clinically

crucial for human health. Figure 6 shows the network topology of the same proteins (for its R

code, see the Data Availability section).

Discussion

Proteome evolution during ages

As shown in the Methods section, relative frequencies of residues in a protein decide on its CAIR. Obviously, biochemical properties of amino acids play a central role in determining their primary relative frequencies in *de novo* proteins. Such dissimilarity of chemical properties would encourage unbalanced primary relative frequencies, thus lesser CAIRs, as shown in our results for younger proteins in Figure 4D. This finding follows the theory of *de novo* gene birth from non-coding DNA (Neme et al., 2017; Wilson et al., 2017). Nevertheless, during the course of evolution, residues are subjected to random mutation which equalizes their relative frequencies. This, in turn, increases the CAIR of proteins as they age which also agrees with the trend of CAIRs in Figure 4D. This can be a corroborating rationale for the study carried out with a different methodology in which it is shown that intrinsic disorder of proteins negatively correlates with gene age (Banerjee & Chakraborty, 2017). Random mutations aside, natural selection's bias in favour of more complex proteins may have also contributed to increasing the CAIR in older proteins. This is also noticeable from the results seen in Figure 2 verifying the identical behaviour of natural selection toward all living organizations. Accordingly, it is not surprising that the human proteome includes a negatively-skewed distribution whose lesser CAIRs are mostly associated with proteins expressed by younger genes. Unfortunately, the literature lacks any thorough investigation on linguistic complexity of proteins and gene ages.

Moreover, it is evident from Figure 4C that the younger eukaryotic proteins are shorter than their older counterparts. A significant decline is noticed, however, in the length, interactions, and mutual information of proteins during the old ages. This refers to the different evolutionary rates of prokaryotic and eukaryotic genes and is compatible with the findings of

previous studies (Alba & Castresana, 2005; Wolf et al., 2009). Even for proteins of

prokaryotic ages, the trend of increase in protein length and interactions is observed;

nonetheless, the trend line is discrete from that of eukaryotic proteins. Interestingly, this

decline is not seen in Figure 4D, as the CAIR, in both eukaryotic and prokaryotic settings, is

identically affected by directional selection. The dome shape increase of disease indicators

from younger to older proteins refutes the justification of the study by Elhaik et al claiming

that the slower rate of evolution in older genes is 'an artefact of increased genetic distance

(Elhaik et al., 2006)'.

Furthermore, it is demonstrated in Figure 4B that interactions increase as genes age which

agrees with the literature (Saeed & Deane, 2006). However, the rate of increase seems to be

slower in a prokaryotic setting. That means the rate of interaction turnover in eukaryotic

proteomes may be comparatively higher. This might be due to the denser networks of

eukaryotes and the gene duplication (Wagner, 2003). It is also noteworthy that the rate of

interaction turnover seems to be non-decreasing as eukaryotic genes get older. Lastly, mutual

information might be considered as an assembly of protein information and interactions. So

as seen in Figure 4A, the trend of EMIP is also increasing by time for both eukaryotic and

prokaryotic genes.

Dynamics of protein networks

A protein network is considered 'stable' when the odds of network malfunction is tiny.

Among various protein networks, particular ones malfunction often, and they generally are

responsible for the networks of non-communicable diseases. According to the theorem

presented in the Methods section, the number of interactions is negatively correlated to

network stability. This deduction is contrary to what is seen in Figures 4B, 5B, and the

literature (Jonsson & Bates, 2006; Oti et al., 2006; Xu & Li, 2006). According to the

literature, the PPI networks of disease genes are different in topology containing more

interactions, as it has been similarly shown in the mentioned figure panels. Previously, it was

clarified that the interactions increase with a stationary rate as a gene ages. As a matter of

fact, there are also many genes encoding for proteins of the cellular organisms that have not

established any interactions. Hence, a confounding factor that might be overlooked would be

the primary stability of the protein itself. In other words, proteins that are prone to

malfunction need substantial interactions to compensate for their malfunctions within their

network. That is the reason for increasing interactions along with the disease occurrences. In

the literature, the finding of the increased mutation rate of disease genes may reason their

increasing interactions (Smith & Eyre-Walker, 2003). Natural selection might be the other

reason which would favour interactions merely for faulty networks rejecting in others.

As mentioned previously, an inaccurate estimate of a protein's stability would be its CAIR.

Figure 4D illustrates that CAIRs of disease proteins are significantly higher than those of

non-disease proteins. However, the incongruous decrease of CAIRs in rare disease comparing

with extremely rare diseases had not expected in the inferences of our theorem. This might

be, in part, the influence of abundant younger proteins responsible for rare diseases that have

not developed interactions yet. Proteins with less CAIR are more stable, yet they are

negatively selected as they do not contain sufficient information. Complexity allows a protein

to have the potential capacity to carry more intricate functions (Babu, 2016). Thus, in order to

obtain higher functionalities, proteins grow both in their sizes and CAIRs. Consequently, this

necessitates new interactions to arrive. However, new protein interactors may take millions of

years to appear and reform the instability of the network (Fraser et al., 2002). These cycles

begin all over again as the new proteins come into existence. All these aside, the essentiality

of a protein's function is an argument that should not be dismissed. Generally, old proteins

are more crucial to forming life than younger ones (Chen et al., 2012). The less crucial roles

of younger proteins render their diseases to be less severe. As human ages, numerous

interactors in various networks are cancelled out causing these networks to malfunction. The

coincidental rush of diseases at the late ages of human life may be caused by the

accumulative effect of interactors removal and the loss of proteostasis (Kaushik & Cuervo,

2015; Labbadia & Morimoto, 2015). So, even substantial interactions cannot fully guarantee

networks of complex unstable proteins.

Gene age, hubs, and human diseases

Our results presented in Figure 5C illustrates the critical role of proteins encoded by the older

genes to be responsible for a broad spectrum of diseases. This finding is in total agreement

with a previous study in the literature highlighting the importance of ancient genes in human

genetic disorders (Domazet-Lošo & Tautz, 2008). We also discussed that the older genes

might take the leading role in creating the necessary fundamentals of life. A series of papers

by Barabási et al dedicatedly demonstrate the distinction between disease genes and the

essential genes. According to their work, disease genes are in most cases non-essentials being

located at the periphery of the network, rather than being a central hub (Barabási et al., 2011;

Domazet-Lošo & Tautz, 2008; Goh et al., 2007). In Figure 5D, although the trend of the

increase in CAIR is parallel to more odds of disease to occur, the proportion of rare to

extremely rare diseases suggests that more prevalent diseases are not associated with the

proteins with the highest CAIRs. This is also evident from the overall comparison of CAIRs

in disease occurrence categories in Figure 4D. This finding puts forward an argument that the

most complex proteins located at the centre of networks are encoded by the essential old

genes that in case of their malfunction, the condition would be fatal or cause an extremely

rare and severe disease. However, the more prevalent diseases are caused by comparably less

complex proteins that are indeed younger than central hubs. So, the proportion of rare to

extremely rare diseases in Figure 5 is of great importance and should be noticed as they agree

with the scenarios presented by Barabási et al.

EMIP and human diseases

A disease may occur when the process of network compensation works inaccurately. For any

network, the removal of the nodes would weaken its stability. This fact is evidently derived

from our theorem. As we discussed, it is the reason why many of the unstable networks have

urged to have many interactions. Integrating the effect of interactions with the protein

information is what mutual information tries to demonstrate. In a sense, the mutual

information of a protein is the information that is identical between the protein and its

network. Thus, in the case of a malfunction, it would not result in a disease, as the network

already carries the identical information. Estimation of mutual information, mathematically,

is equivalent to the difference between the information of the network as a whole and the

scalar summation of information that interactors carry when they are not interacting within a

network. This, of course, will show how much capacity the network carries to compensate for

its malfunction. In Figures 4A and 5A, the superior relation between EMIP and disease

occurrence categories stems from this fact.

On the existence of diseases

Based on what is discussed above, the scenario of a gene to cause a disease or not is being

summarized as the following (Figure 7). When the evolution was in its initial periods, the

involved proteins for life network was mainly the crucial ones. The metabolic pillars of life

owe the fundamental and critical components to the first formation of these networks. There

are two types of proteins at this stage, i.e. either 'robust' or 'weak'. To define, robust proteins

are those which are structurally not very susceptible to malfunction. These proteins have

evolved without many interactions comparing to others. Hence, during the path of evolution,

and still, we cannot detect as many interactions for them. On the other hand, weak proteins

would have caused vital errors that result in severe diseases. It would be reasonable to assume

that the incidence of such diseases would have been higher at ancient ages. So we may expect

to observe many interactions of them till now. According to our theorem, it can be reasoned

that these interactions would bypass the hub protein in case it malfunctions. So the incidence

of such diseases would have been drastically lessened by now. As a general point, for an

unstable weak protein that initiates a network, there are two prospects to be considered, i.e.

being able to develop a mature network at the time of the investigation, or still getting

involved in an immature one. By definition, mature networks would be those which have

totally cancelled out the adverse outcomes of the malfunctioning hub protein. Based on the

theorem, maturity is an ideal that no network can satisfy it in a biology setting.

Having mentioned all these about the archaic proteins, it should also be noticed that the

proteins which have come into existence in more chronologically proximal periods would

have by far lower chances of being a vital hub. The functioning network of life, in one piece,

has less to do with a mammalian protein than an archaic cellular respiration protein. So, the

other category of proteins is that of the contemporary period on which the logical assumption

would be that they are either non-hubs or if hubs do not function in places crucial to life. The

contemporary category, i.e. the young proteins, again, is subdivided into robust and weak

proteins. Robust young proteins are those without interactions which are not very much

susceptible to cause diseases, because if otherwise, evolution would bring interactions for

them. It is noteworthy that very young proteins are mainly very stable, simple, and proteins

with minor functional capacities. Stable young proteins may by time change to unstable

complex proteins because of the random mutations and the natural selection's bias toward

more complex proteins. In the way of transformation, most of the weak young proteins arise

which are primarily responsible for the prevalent metabolic diseases of the current

evolutionary era. Since the average evolutionary rate interaction turnover has not satisfied the

optimum number of network nodes for them, they are apprentices susceptible to malfunction.

It would be easy to infer that these proteins cause less severe diseases comparing to archaic

proteins, as their functions are still not vital. Nevertheless, they cause diseases with higher

incidences.

Applicability of the method

The methodology we have presented in the next section is not sensitive to the level of

taxonomy, i.e. whether the calculation is for a species, a genus, an order, a kingdom, or the

complete tree of life. The reason for this fact is that calculating the amino acid frequencies is

the same considering a species proteome or any other more extensive proteomic combination

of taxonomic levels. Also, we can calculate the Shannon entropy for a single protein, or a

peptide. This insensitivity to the size of the network in calculation enables a homogenous

analysis through the whole tree of life.

A perspective of future studies

Future studies may focus on each of the non-communicable diseases to elaborate more on the

speculations made in this study. Communicable diseases may also be the focus of further

studies to investigate the CAIRs of organisms and probable relations to their pathogenicity.

Treatments that are targeting networks with high-CAIR interacting protein crowds would be

an option to be explored. Moreover, better estimations for mutual information would be of

great interest. Besides, the notation of the wave of life and CAIR comparisons may bring

further arguments to taxonomists. Lastly, the theorem may be used in various fields of

science which are shaped by networks.

Methods

Introducing the Calculated Average Information per Residue (CAIR) and the Protein

Information (PI)

Proteins, form a mathematical point of view, are once randomly-occurred sequences of

residues that have gone through a process of selection in nature. Considering this fact, a

protein structure can be defined using a random sequence that carries mathematical

information. The information that a protein carries may be defined as to be equivalent to the

amount of uncertainty in predicting its residues. It is to be highlighted that regardless of the

protein conformation, the information of a protein is determined merely by its primary

structure. The average information carried by a residue in a protein is calculated by

Shannon's entropy (H) equation as below:

$$H = - \sum_{i=1}^{s} p_i \, \log_2 p_i \tag{1}$$

337 where p_i is the probability of state i, and s is the total number of possible states. In the

338 current context, the CAIR notion is introduced to be the same as Shannon's entropy except

339 for the logarithm base which is 22 in the former and 2 in the latter. In other words, CAIR is

340 the 22-ary of Shannon's entropy and is formulated as:

$$\text{CAIR} = - \sum_{r=1}^{t} p_r \, \log_{22} p_r \tag{2}$$

341 in which r is a numeral given to each residue, t is the total number of residues, p_r is the

342 relative frequency of r^{th} residue in the protein. More simply, the CAIR could be written as:

$$\text{CAIR} = kH \tag{3}$$

343 in which H is Shannon's entropy in equation (1), and k is a constant equivalent to:

$$k = \log_{22} 2 \cong 0.224243$$

344 As it is evident from equation (3), the patterns in the results obtained in the current article

345 were independent of the base of the logarithm; however, the scale of entropy would be much

346 more tangible considering the base of 22, as the ideal proteinogenic alphabet contains 22

347 letters. Deriving from CAIR, protein information (PI) is the amount of information carried by

348 all the residues of a protein as of the following equation:

$$\text{PI} = -l \sum_{r=1}^{t} p_r \, \log_{22} p_r \tag{4}$$

in which the variables are the same as those in equation (2), and l is the length of the protein.

The notations of PI and CAIR are proposed, instead of the conventional H, for the fact that

they are more expressive and pertinent for the field of proteomics. It would also be humbly

proposed – with an analogy to Shannon's bit – to use 'pit', i.e. protein unit, for the unit of

CAIR, PI, and EMIP in order to be readily comprehensible. As an example, one kilo-pit

would be equivalent to the PI of a 1000-lengthed protein whose residues have equal

frequencies.

Mathematical Theorem

Suppose $\{X\}, \{Y_1\}, \{Y_2\} \dots, \{Y_n\}$ are sets of sequences from which $\{Y_1\}, \{Y_2\} \dots, \{Y_n\}$ are all

dependent to $\{X\}$, but are pairwise independent from each other, not necessarily identically

distributed random variables having characteristic functions of $\varphi_1, \varphi_2, \dots, \varphi_n$, distribution

functions of $f_1, f_2, \dots, f_n$, and entropies of $H_1, H_2, \dots, H_n$. Let Φ be the characteristic function

of $\{X\}$, $F_{\tilde{x}}$ be its distribution function, and $H_{\tilde{x}}$ be its entropy. Then:

$$\lim_{n \to \infty} H(\Phi | \varphi_1, \varphi_2, \ldots, \varphi_n) = 0$$

364   ***Proof.*** According to the definition of mutual information, we can write the following

365   relations (Cover & Thomas, 2012):

$$I(X; Y_i) = I(Y_i; X)$$

$$I(Y_i; X) = H(X) - H(X|Y_i)$$

$$\sum_{i=1}^{\infty} H(X|Y_i) = H(X) - \sum_{i=1}^{\infty} I(X; Y_i) \tag{5}$$

366   Corollary. Non-negativity of mutual information(Cover & Thomas, 2012):

$$I(X; Y) \geq 0$$

367   with equality i□ X and Y are independent.

368   Based on the corollary of non-negativity of mutual information, and because Y_i are all

369   dependent on X, the mutual information of X with respect to all Y_i is always positive:

$$I(X; Y_i) > 0 \tag{6}$$

370   from which it can be inferred that infinite sum of positive values yields not to infinite, but to

371   the maximum mutual information possible, i.e., the entropy of X:

$$\sum_{i=1}^{\infty} I(X; Y_i) = H(X) \tag{7}$$

372   So, substituting equation (7) in equation (5):

$$\sum_{i=1}^{\infty} H(X|Y_i) = H(X) - H(X)$$

$$\sum_{i=1}^{\infty} H(X|Y_i) = 0 \tag{8}$$

373   Also, since $\sum_{i=1}^{\infty} H(X|Y_i)$ is the same as $\lim\limits_{n \to \infty} H(\Phi|\varphi_1, \varphi_2, \dots, \varphi_n)$, thus:

$$\lim_{n \to \infty} H(\Phi|\varphi_1, \varphi_2, \dots, \varphi_n) = 0 \tag{9}$$

Biological inferences and hypotheses

In the above theorem, supposing Φ be a protein with the amino acid sequence of $\{X\}$,

interacting with n number of other proteins, namely $\varphi_1, \varphi_2, \dots, \varphi_n$, with sequences of

$\{Y_1\}, \{Y_2\} \dots, \{Y_n\}$:

1)   Said interactions are mathematically interpreted as the dependency of $F_{\mathfrak{x}}$ on

$f_1, f_2, \dots, f_n$. To elucidate, the probability distribution functions of proteins

correspond to their Boltzmann distributions. Because of the induced-fit nature of

biochemical interactions, it might be plausible to consider the distribution

functions to be dependent on one another. For that reason, each interaction is

deduced as the dependency of two distributions in the theorem.

2)   As the theorem suggests, $H_{\mathfrak{x}}$ indicates Shannon's entropy of the protein Φ with an

amino acid sequence of $\{X\}$. This might be interpreted as the extent of probable

variations that could lay in the primary structure of a protein affecting its function.

This measure would be an inaccurate estimate of a protein's malfunction as no

biochemical conditions have been taken into account. Despite its inaccuracy, we

have included the CAIR as an indicator of diseases in our analysis. The intuitive

hypothesis would maintain that the CAIR is significantly more in disease proteins

than non-disease ones for their surplus odds of having potential disadvantageous

variations in the primary structure leading to malfunction and disease.

3) According to the assumptions of the theorem, the notation of 'information' could

also apply to any system containing proteins, i.e. a particular metabolic pathway,

an interactome, diseasome, or the entire living organisms. Measuring the

information is independent of the size of the network. This property allows us to

calculate the information among the different taxonomic hierarchies and to utilize

the results in practice. Also, in a single organism like *Homo sapiens*, it would

allow us to compare different disease pathways, track hub proteins, and discern

potential disease proteins from non-disease ones.

4) The length of a protein sequence determines the PI, as shown in equation (4). We

have not included the PI itself as an indicator in the study, but have included both

the CAIR and the protein length separately. Proteins with longer sequences carry

more information and are more prone to malfunction as the overall odds of a

faulty residue in their sequence is higher compared to a protein with a shorter

sequence.

5) Considering equation (9), it is understandable that the conditional entropy of Φ

with respect to the knowledge of *n* number of interactors, i.e. $\varphi_1, \varphi_2, \ldots, \varphi_n$, would

equal zero when *n* approaches infinity. The conditional entropy designates the

new information carried by the Φ protein when functioning in its network with

other proteins of φ_1 to φ_n. So, as a preliminary and naive inference, it could be

easily inferred from the theorem that networks with more interactions are more

stable. Therefore, we have included the number of interactions as an indicator of

human diseases to test our hypothesis.

6) Although equation (9) is a relatively straightforward approach to calculate the

stability of a network, it can be shown that its exact quantitative calculation is not

possible in proteomic analysis. An equivalent measure of network stability would

be to use equation (6) in which mutual information is shown to be always positive.

Unlike the conditional entropy which is negatively correlated to the network

stability, the mutual information is positively correlated. To quantitate mutual

information in the current context, we propose the following estimation,

henceforth referred to as EMIP (μ):

$$\mu_\Phi = -[\sum_{\varphi=1}^{n} l_\varphi + l_\Phi] \sum_{\varphi=1}^{n} \sum_{r=1}^{t} \frac{p_{(r,\varphi)} l_\varphi + p_{(r,\Phi)} l_\Phi}{l_\varphi + l_\Phi} \log_{22} \frac{p_{(r,\varphi)} l_\varphi + p_{(r,\Phi)} l_\Phi}{l_\varphi + l_\Phi}$$

$$+ \sum_{\varphi=1}^{n} \sum_{r=1}^{t} l_\varphi \, p_{(r,\varphi)} \log_{22} p_{(r,\varphi)} \tag{10}$$

in which μ_Φ is the mutual information for the Φ protein, n is the number of

interactions, l_φ is the length of the φ^{th} interactor, l_Φ is the length of protein Φ, r is

a numeral given to each residue in proteins, t is the total number of residues,

$p_{(r,\varphi)}$ is the relative frequency of r^{th} residue in φ^{th} interactor, and $p_{(r,\Phi)}$ is the

relative frequency of r^{th} residue in Φ protein. EMIP has also been included as an

indicator of diseases in our analysis.

7) An essential element in the course of life evolution and human disease analysis is

to consider the gene age of proteins. It is conceivable to hypothesize that the

chronological data of genes can be very much associated with the indicators of

human diseases. One reason is the additive effect of gene ages to introduce new

interactions. The second reason is based on the assumption that natural selection

can show bias towards proteins with a specific range of CAIRs. Also, the third

reason is the possibility that the older proteins can grow to have longer sequences

during evolution. Therefore, we have also included the gene ages as a covariate of

human diseases in our analysis.

8)　According to the theorem, it is also inferred that in an ideal condition with an

infinite number of interactors for a protein, the functionality of such a hub protein

reaches an infallible state, i.e. no disease would ever happen. This could also be

applied to other fields of science to reason why there is an order in complex

systems with innumerable components. The rate of decline in error in our theorem

as shown in equation (9) under random conditions is generally consistent with the

simple statistical rule of $\sqrt{n}$ as proposed in physicist Schrödinger's 'what is life'

(Schrödinger, 1944).

Protein database

For taxonomic comparisons, all complete proteomes were extracted from the UniProt

(Consortium, 2019) FTP server, freely available at

ftp://ftp.uniprot.org/pub/databases/uniprot/current_release/knowledgebase/complete/. Both

SwissProt and TrEMBL files were downloaded in FASTA format. We calculated protein

entropies separately for each entry in both files including a total of more 61 billion amino

acids and merged them with every individual organism. Then, we used the calculations for

further statistical analyses.

Besides, for human proteins analyses, the complete list of *Homo sapiens* proteins was

downloaded directly from the UniProt website in a tab-separated (.tab) format containing the

following columns: 'Entry', 'Length', 'Sequence', 'Orphanet', and 'Involvement in disease'.

Data were updated on 22nd April 2020 with UniProt release 2020_02.

Protein-protein interactions database

Protein-protein interactions in human proteome were obtained from Protein InteraCtion

KnowLedgebasE (Gioutlakis et al., 2017) (PICKLE) meta-database, the release of 2.5.

PICKLE is a cross-checked integration of all available human PPIs included in BioGRID,

IntAct, HPRD, MINT, DIP databases. The default filter mode was selected to download

191,113 binary interactions among 16,418 UniProtKB/SwissProt entries. All interactions

were included in our study for further analysis.

Taxonomy database

Taxonomy data of organisms were also extracted from the UniProt FTP, the release of

2020_02. All organisms were included and matched according to their 'OX', i.e. organism

number used by UniProt and other databases. The evolutionary tree was plotted based on a

landmark study (Hug et al., 2016) by Hug et al, published in 2016, with a review of updates

(Carnevali et al., 2019; Carr et al., 2019; Cavalier-Smith et al., 2014, 2018; Dombrowski et

al., 2019; Eloe-Fadrosh et al., 2016; Eme et al., 2017; Hahnke et al., 2016; Hamilton et al.,

2016; Jay et al., 2018; Jungbluth et al., 2017; Kevbrin et al., 2020; Kirkegaard et al., 2016;

Martinez et al., 2019; L. Momper et al., 2017; L. M. Momper et al., 2018; Munoz et al., 2016;

Pavan et al., 2018; Y. Wang et al., 2019; Ward et al., 2019; Youssef et al., 2019; Zhou et al.,

2020) since then until April 2020. All updates were added to the tree and were matched with

UniProt taxonomy data.

Diseases database

According to cross-references between UniProt and Orphanet, related epidemiological data

were downloaded and extracted from the Orphanet database (Weinreich et al., 2008) in XML

format. The file was then used in Python code for further analysis. Only diseases with at least

one reported worldwide occurrence were included. Normally, in the Orphanet database,

occurrences are reported from one or more of the following categories: annual incidence,

cases/families, lifetime prevalence, point prevalence, and prevalence at birth. In the case of

more than one reported occurrence, the priority of selection was for incidence, prevalence at

birth, point prevalence, respectively. Accumulative occurrence data, i.e. cases/families and

lifetime prevalence, were excluded from the analysis.

Analysis of taxonomic hierarchies

The following steps were carried out to implement the theorem over the taxonomic

hierarchies (Figure 8A):

1) TrEMBL and SwissProt FASTA files of complete proteomes were downloaded from

UniProt KnowledgeBase. A total number of ~180 million protein entries were

included.

2) The frequencies of all amino acid residues were calculated with regards to all protein

entries.

3) All proteins were grouped according to their organisms using organism IDs.

4) Organisms that are not proteomes were excluded. Duplicates are also removed.

5) Viruses are excluded from the study.

6) Shannon's entropy was calculated according to the residue frequencies for all

included non-virus proteomes.

7) Taxonomic data were downloaded directly from the UniProt website.

8) The most updated tree of life was drawn after a thorough review of the literature until

April 2020.

9) Organisms were grouped with respect to taxonomic hierarchies.

10) Organisms with unknown taxonomic lineage were excluded from the analysis.

11) Brunner-Munzel test was performed for every bifurcation node through the tree of

life. The level of significance was 0.05. Preparation of data and protein information

calculations were all executed in Python 3.8.0 (Van Rossum & Drake, 2009) using

NumPy (Oliphant, 2006), Pandas (McKinney & others, 2010), and Biopython (Cock

et al., 2009) libraries. Statistical analysis and violin plots were carried out using R-

3.6.0 (R Core Team, 2019) with brunnermunzel (Ara, 2020) and Plotly (Sievert,

2018) packages.

Analysis of human disease proteins

The following steps were carried out to implement EMIP and analyse disease occurrence

categories (Figure 8B):

1) The human proteome was downloaded from the UniProt database with the organism

522   ID of 9606. 20,350 of reviewed and 54,473 of unreviewed proteins were included in

523   the study.

524   2) CAIR was calculated for all human entries using the sequences of residues.

525   3) Protein-protein interactions were extracted from the PICKLE database as the default

526   UniProt normalized file with a total number of ~190k interactions.

527   4) Interactions were altered in order to match the UniProt 'interactions with' column.

528   This was done to keep the homogeneity of the data.

529   5) EMIP was then calculated for all entries with the help of the PPIs.

530   6) The unreviewed proteins were excluded after the calculations of disease indicators

531   because they have not been reported to cause any diseases.

532   7) Ordinal age categories of genes were merged to the file using the consensus data

533   article (Liebeskind et al., 2016).

8) Disease categories are ranked using R into three groups of no diseases, extremely rare

diseases, and rare diseases.

9) Disease indicators were also categorized into eight groups to draw Likert plots.

10) Statistical differences between groups were tested among variables with the DTK test.

Significance levels of the DTK test were added to the comparison graphs with stars.

The highest significance level was set to 0.05. Graphs of comparisons were plotted in

R with DTK (Lau, 2013) and ggpubr (Kassambara, 2019) packages. Likert plots were

plotted with HH (Heiberger, 2019) package. Networks were plotted using igraph

(Csardi & Nepusz, 2006) package.

Data availability

Data used for analysis is available as the following files. FASTA files of all Swiss-Prot and

TrEMBL entries are publicly available from UniProt's FTP server at

https://ftp.expasy.org/databases/uniprot/current_release/knowledgebase/complete/. Also, all

non-redundant proteomes could be downloaded from UniProt website:

https://www.uniprot.org/proteomes/?query=redundant:no&format=tab&force=true&columns

=id,name,organism-id,lineage&compress=yes. Tab-separated format of human proteome data

used in our analysis is achievable from

https://www.uniprot.org/uniprot/?query=proteome:UP000005640&format=tab&force=true&

columns=id,reviewed,genes(PREFERRED),protein%20names,sequence,database(Orphanet),

comment(INVOLVEMENT%20IN%20DISEASE),interactor&compress=yes. Additionally,

PICKLE interactions are freely available from

http://www.pickle.gr/Data/2.5/PICKLE2_5_UniProtNormalizedTabular-default.zip. Orphanet

data is also freely available from http://www.orphadata.org/data/xml/en_product9_prev.xml.

Data of gene ages are adopted from the Gene-Ages GitHub repository at

https://github.com/marcottelab/Gene-Ages/raw/master/Main/main_HUMAN.csv.

Supplementary information is available in the online version.

All Python and R codes necessary to reproduce all parts of the analysis and for the illustration

of the figures are available under the MIT license on our GitHub repository at

https://github.com/synaptic-proteolab/CAIR_EMIP, or the Zenodo link at

https://zenodo.org/record/3975465. For executing codes online on cloud servers, Google

Colab links are also available on the GitHub page. Python

(https://www.python.org/downloads/), Jupyter Notebook (https://jupyter.org/install), R

(https://cran.r-project.org/), and RStudio (https://rstudio.com/products/rstudio/download/)

are all freely available for the public.

Acknowledgments

Authors would like to thank Dr Saeed Sadigh-Eteghad, PhD in neurosciences, Tabriz

University of Medical Sciences, and Dr Pedram Dindari, a PhD candidate in computer

sciences, University of Tabriz, for their assistance during the preparation of the manuscript.

No funding to declare.

Author contributions

FK proposed the mathematical theorem and its proof. SD gathered and handled the large data

and prepared the data for analysis using Python. FK reviewed the literature to illustrate the

tree of life. FK analysed the data using R. SD and FK prepared the codes for open-source

publishing. FK wrote the manuscript, and SD agreed with all sections. FK designed the

figures and SD prepared the tables of the manuscript.

Conflict of interest

The authors have no conflict of interest to disclose.

References

Alba, M. M., & Castresana, J. (2005). Inverse relationship between evolutionary rate and age of mammalian genes. *Molecular Biology and Evolution*, *22*(3), 598–606.

Andrews, J. G., Dimakis, A., Dolecek, L., Effros, M., Medard, M., Milenkovic, O., Montanari, A., Vishwanath, S., Yeh, E., & Berry, R. (2015). A perspective on future research directions in information theory. *ArXiv Preprint ArXiv:1507.05941*.

Ara, T. (2020). *brunnermunzel: (Permuted) Brunner-Munzel Test*. https://cran.r-project.org/package=brunnermunzel

Babu, M. M. (2016). The contribution of intrinsically disordered regions to protein function, cellular complexity, and human disease. *Biochemical Society Transactions*, *44*(5), 1185–1200.

Bailey, K. D. (2009). Entropy systems theory. *Systems Science and Cybernetics, Eolss Publishers, Oxford, UK*, 152–169.

Banerjee, S., & Chakraborty, S. (2017). Protein intrinsic disorder negatively associates with gene age in different eukaryotic lineages. *Molecular BioSystems*, *13*(10), 2044–2055.

Barabási, A.-L., Gulbahce, N., & Loscalzo, J. (2011). Network medicine: a network-based approach to human disease. *Nature Reviews Genetics*, *12*(1), 56–68.

Battail, G. (2013). Biology needs information theory. *Biosemiotics*, *6*(1), 77–103.

Bodmer, W., & Bonilla, C. (2008). Common and rare variants in multifactorial susceptibility to common diseases. *Nature Genetics*, *40*(6), 695.

Carnevali, P. B. M., Schulz, F., Castelle, C. J., Kantor, R. S., Shih, P. M., Sharon, I., Santini, J. M., Olm, M. R., Amano, Y., Thomas, B. C., & others. (2019). Hydrogen-based

metabolism as an ancestral trait in lineages sibling to the Cyanobacteria. *Nature*

Communications, *10*(1), 1–15.

Carr, S. A., Jungbluth, S. P., Eloe-Fadrosh, E. A., Stepanauskas, R., Woyke, T., Rappé, M.

S., & Orcutt, B. N. (2019). Carboxydotrophy potential of uncultivated

Hydrothermarchaeota from the subseafloor crustal biosphere. *The ISME Journal*, *13*(6),

1457–1468.

Cavalier-Smith, T., Chao, E. E., & Lewis, R. (2018). Multigene phylogeny and cell evolution

of chromist infrakingdom Rhizaria: contrasting cell organisation of sister phyla

Cercozoa and Retaria. *Protoplasma*, *255*(5), 1517–1574.

Cavalier-Smith, T., Chao, E. E., Snell, E. A., Berney, C., Fiore-Donno, A. M., & Lewis, R.

(2014). Multigene eukaryote phylogeny reveals the likely protozoan ancestors of

opisthokonts (animals, fungi, choanozoans) and Amoebozoa. *Molecular Phylogenetics*

and Evolution, *81*, 71–85.

Changeux, J.-P., & Edelstein, S. (2011). Conformational selection or induced fit? 50 years of

debate resolved. *F1000 Biology Reports*, *3*.

Chen, W.-H., Trachana, K., Lercher, M. J., & Bork, P. (2012). Younger genes are less likely

to be essential than older genes, and duplicates are less likely to be essential than

singletons of the same age. *Molecular Biology and Evolution*, *29*(7), 1703–1706.

Clausius, R. (1864). *Abhandlungen über die mechanische Wärmetheorie* (Vol. 1). F. Vieweg.

Cock, P. J. A., Antao, T., Chang, J. T., Chapman, B. A., Cox, C. J., Dalke, A., Friedberg, I.,

Hamelryck, T., Kauff, F., Wilczynski, B., & others. (2009). Biopython: freely available

Python tools for computational molecular biology and bioinformatics. *Bioinformatics*,

25(11), 1422–1423.

Consortium, U. (2019). UniProt: a worldwide hub of protein knowledge. *Nucleic Acids*

Research, *47*(D1), D506–D515.

Cover, T. M., & Thomas, J. A. (2012). *Elements of information theory*. John Wiley & Sons.

Csardi, G., & Nepusz, T. (2006). The igraph software package for complex network research.

InterJournal, *Complex Sy*, 1695. http://igraph.org

Domazet-Lošo, T., & Tautz, D. (2008). An ancient evolutionary origin of genes associated

with human genetic diseases. *Molecular Biology and Evolution*, *25*(12), 2699–2707.

Dombrowski, N., Lee, J.-H., Williams, T. A., Offre, P., & Spang, A. (2019). Genomic

diversity, lifestyles and evolutionary origins of DPANN archaea. *FEMS Microbiology*

Letters, *366*(2), fnz008.

Dunnett, C. W. (1980). Pairwise multiple comparisons in the unequal variance case. *Journal*

of the American Statistical Association, *75*(372), 796–800.

Elhaik, E., Sabath, N., & Graur, D. (2006). The "inverse relationship between evolutionary

rate and age of mammalian genes" is an artifact of increased genetic distance with rate

of evolution and time of divergence. *Molecular Biology and Evolution*, *23*(1), 1–3.

Eloe-Fadrosh, E. A., Paez-Espino, D., Jarett, J., Dunfield, P. F., Hedlund, B. P., Dekas, A. E.,

Grasby, S. E., Brady, A. L., Dong, H., & Briggs, B. R. (2016). Global metagenomic

survey reveals a new bacterial candidate phylum in geothermal springs. *Nature*

Communications, *7*, 10476.

Eme, L., Spang, A., Lombard, J., Stairs, C. W., & Ettema, T. J. G. (2017). Archaea and the

origin of eukaryotes. *Nature Reviews Microbiology*, *15*(12), 711.

Evans, P. N., Boyd, J. A., Leu, A. O., Woodcroft, B. J., Parks, D. H., Hugenholtz, P., &

Tyson, G. W. (2019). An evolving view of methane metabolism in the Archaea. *Nat.*

Rev. Microbiol, *17*(4), 219–232.

Fraser, H. B., Hirsh, A. E., Steinmetz, L. M., Scharfe, C., & Feldman, M. W. (2002).

Evolutionary rate in the protein interaction network. *Science*, *296*(5568), 750–752.

Ghahramani, Z. (2006). Information theory. *Encyclopedia of Cognitive Science*.

Gioutlakis, A., Klapa, M. I., & Moschonas, N. K. (2017). PICKLE 2.0: A human protein-

protein interaction meta-database employing data integration via genetic information

ontology. *PloS One*, *12*(10).

Goh, K.-I., Cusick, M. E., Valle, D., Childs, B., Vidal, M., & Barabási, A.-L. (2007). The

human disease network. *Proceedings of the National Academy of Sciences*, *104*(21),

8685–8690.

Hahnke, R. L., Meier-Kolthoff, J. P., García-López, M., Mukherjee, S., Huntemann, M.,

Ivanova, N. N., Woyke, T., Kyrpides, N. C., Klenk, H.-P., & Göker, M. (2016).

Genome-based taxonomic classification of Bacteroidetes. *Frontiers in Microbiology*, *7*,

2003.

Hamilton, T. L., Bovee, R. J., Sattin, S. R., Mohr, W., Gilhooly III, W. P., Lyons, T. W.,

Pearson, A., & Macalady, J. L. (2016). Carbon and sulfur cycling below the chemocline

in a meromictic lake and the identification of a novel taxonomic lineage in the FCB

superphylum, Candidatus Aegiribacteria. *Frontiers in Microbiology*, *7*, 598.

Heiberger, R. M. (2019). *HH: Statistical Analysis and Data Display: Heiberger and Holland.*

https://cran.r-project.org/package=HH

Hug, L. A., Baker, B. J., Anantharaman, K., Brown, C. T., Probst, A. J., Castelle, C. J.,

Butterfield, C. N., Hernsdorf, A. W., Amano, Y., & Ise, K. (2016). A new view of the

tree of life. *Nature Microbiology*, *1*(5), 16048.

Jay, Z. J., Beam, J. P., Dlakić, M., Rusch, D. B., Kozubal, M. A., & Inskeep, W. P. (2018). Marsarchaeota are an aerobic archaeal lineage abundant in geothermal iron oxide microbial mats. *Nature Microbiology*, *3*(6), 732.

Jonsson, P. F., & Bates, P. A. (2006). Global topological features of cancer proteins in the human interactome. *Bioinformatics*, *22*(18), 2291–2297.

Jungbluth, S. P., Amend, J. P., & Rappé, M. S. (2017). Metagenome sequencing and 98 microbial genomes from Juan de Fuca Ridge flank subsurface fluids. *Scientific Data*, *4*(1), 1–11.

Kassambara, A. (2019). *ggpubr: "ggplot2" Based Publication Ready Plots*. https://cran.r-project.org/package=ggpubr

Kaushik, S., & Cuervo, A. M. (2015). Proteostasis and aging. *Nature Medicine*, *21*(12), 1406.

Keskin, O., Gursoy, A., Ma, B., & Nussinov, R. (2008). Principles of protein– protein interactions: What are the preferred ways for proteins to interact? *Chemical Reviews*, *108*(4), 1225–1244.

Kevbrin, V., Boltyanskaya, Y., Grouzdev, D., Koziaeva, V., Park, M., & Cho, J.-C. (2020). Natronospirillum operosum gen. nov., sp. nov., a haloalkaliphilic satellite isolated from decaying biomass of a laboratory culture of cyanobacterium Geitlerinema sp. and proposal of Natronospirillaceae fam. nov., Saccharospirillaceae fam. nov. and Gynuell. *International Journal of Systematic and Evolutionary Microbiology*, *70*(1), 511–521.

Kirkegaard, R. H., Dueholm, M. S., McIlroy, S. J., Nierychlo, M., Karst, S. M., Albertsen, M., & Nielsen, P. H. (2016). Genomic insights into members of the candidate phylum Hyd24-12 common in mesophilic anaerobic digesters. *The ISME Journal*, *10*(10), 2352.

Koshland Jr, D. E. (1995). The key–lock theory and the induced fit theory. *Angewandte*

Chemie International Edition in English, *33*(23□24), 2375–2378.

Labbadia, J., & Morimoto, R. I. (2015). The biology of proteostasis in aging and disease.

Annual Review of Biochemistry, *84*, 435–464.

Lau, M. K. (2013). *DTK: Dunnett-Tukey-Kramer Pairwise Multiple Comparison Test*

Adjusted for Unequal Variances and Unequal Sample Sizes. https://cran.r-

project.org/package=DTK

Liebeskind, B. J., McWhite, C. D., & Marcotte, E. M. (2016). Towards consensus gene ages.

Genome Biology and Evolution, *8*(6), 1812–1823.

Martinez, M. A., Woodcroft, B. J., Espinoza, J. C. I., Zayed, A. A., Singleton, C. M., Boyd, J.

A., Li, Y.-F., Purvine, S., Maughan, H., Hodgkins, S. B., & others. (2019). Discovery

and ecogenomic context of a global Caldiserica-related phylum active in thawing

permafrost, Candidatus Cryosericota phylum nov., Ca. Cryosericia class nov., Ca.

Cryosericales ord. nov., Ca. Cryosericaceae fam. nov., comprising the four species C.

Systematic and Applied Microbiology, *42*(1), 54–66.

McKinney, W., & others. (2010). Data structures for statistical computing in python.

Proceedings of the 9th Python in Science Conference, *445*, 51–56.

Momper, L., Jungbluth, S. P., Lee, M. D., & Amend, J. P. (2017). Energy and carbon

metabolisms in a deep terrestrial subsurface fluid microbial community. *The ISME*

Journal, *11*(10), 2319–2333.

Momper, L. M., Aronson, H., & Amend, J. P. (2018). Genomic description of 'Candidatus

Abyssubacteria,'a novel subsurface lineage within the candidate phylum

Hydrogenedentes. *Frontiers in Microbiology*, *9*, 1993.

Munoz, R., Rosselló-Móra, R., & Amann, R. (2016). Revised phylogeny of Bacteroidetes and

proposal of sixteen new taxa and two new combinations including Rhodothermaeota

phyl. nov. *Systematic and Applied Microbiology*, *39*(5), 281–296.

Neme, R., Amador, C., Yildirim, B., McConnell, E., & Tautz, D. (2017). Random sequences

are an abundant source of bioactive RNAs or peptides. *Nature Ecology & Evolution*,

1(6), 1–7.

Neuhäuser, M., & Ruxton, G. D. (2009). Distribution-free two-sample comparisons in the

case of heterogeneous variances. *Behavioral Ecology and Sociobiology*, *63*(4), 617–623.

Oliphant, T. E. (2006). *A guide to NumPy* (Vol. 1). Trelgol Publishing USA.

Oti, M., Snel, B., Huynen, M. A., & Brunner, H. G. (2006). Predicting disease genes using

protein–protein interactions. *Journal of Medical Genetics*, *43*(8), 691–698.

Pavan, M. E., Pavan, E. E., Glaeser, S. P., Etchebehere, C., Kämpfer, P., Pettinari, M. J., &

López, N. I. (2018). Proposal for a new classification of a deep branching bacterial

phylogenetic lineage: transfer of Coprothermobacter proteolyticus and

Coprothermobacter platensis to Coprothermobacteraceae fam. nov., within

Coprothermobacterales ord. nov., Coprothermobacte. *International Journal of*

Systematic and Evolutionary Microbiology, *68*(5), 1627–1632.

R Core Team. (2019). *R: A Language and Environment for Statistical Computing*.

https://www.r-project.org

Saeed, R., & Deane, C. M. (2006). Protein protein interactions, evolutionary rate, abundance

and age. *BMC Bioinformatics*, *7*(1), 128.

Schrödinger, E. (1944). *What is life? The physical aspect of the living cell and mind*.

Cambridge University Press Cambridge.

Shannon, C. E. (1948). A mathematical theory of communication. *Bell System Technical Journal*, *27*(3), 379–423.

Sievert, C. (2018). *plotly for R*. https://plotly-r.com

Smith, N. G. C., & Eyre-Walker, A. (2003). Human disease genes: patterns and predictions. *Gene*, *318*, 169–175.

Spang, A., Caceres, E. F., & Ettema, T. J. G. (2017). Genomic exploration of the diversity, ecology, and evolution of the archaeal domain of life. *Science*, *357*(6351).

Van Rossum, G., & Drake, F. L. (2009). *Python 3 Reference Manual*. CreateSpace.

Vinga, S. (2013). Information theory applications for biological sequence analysis. *Briefings in Bioinformatics*, *15*(3), 376–389.

Wagner, A. (2003). How the global structure of protein interaction networks evolves. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *270*(1514), 457–466.

Wang, Y., Wegener, G., Hou, J., Wang, F., & Xiao, X. (2019). Expanding anaerobic alkane metabolism in the domain of Archaea. *Nature Microbiology*, *4*(4), 595–602.

Wang, Z., & Moult, J. (2001). SNPs, protein structure, and disease. *Human Mutation*, *17*(4), 263–270.

Ward, L. M., Cardona, T., & Holland-Moritz, H. (2019). Evolutionary Implications of Anoxygenic Phototrophy in the Bacterial Phylum Candidatus Palusbacterota (WPS-2). *BioRxiv*, 534180.

Weinreich, S. S., Mangon, R., Sikkens, J. J., Teeuw, M. E., & Cornel, M. C. (2008). Orphanet: a European database for rare diseases. *Nederlands Tijdschrift Voor*

765 *Geneeskunde*, *152*(9), 518–519.

766 Wilson, B. A., Foy, S. G., Neme, R., & Masel, J. (2017). Young genes are highly disordered

767  as predicted by the preadaptation hypothesis of de novo gene birth. *Nature Ecology &*

768  *Evolution*, *1*(6), 1–6.

769 Wolf, Y. I., Novichkov, P. S., Karev, G. P., Koonin, E. V, & Lipman, D. J. (2009). The

770  universal distribution of evolutionary rates of genes and distinct characteristics of

771  eukaryotic genes of different apparent ages. *Proceedings of the National Academy of*

772  *Sciences*, *106*(18), 7273–7280.

773 Xu, J., & Li, Y. (2006). Discovering disease-genes by topological features in human protein–

774  protein interaction network. *Bioinformatics*, *22*(22), 2800–2805.

775 Youssef, N. H., Farag, I. F., Hahn, C. R., Jarett, J., Becraft, E., Eloe-Fadrosh, E., Lightfoot,

776  J., Bourgeois, A., Cole, T., Ferrante, S., & others. (2019). Genomic characterization of

777  candidate division LCP-89 reveals an atypical cell wall structure, microcompartment

778  production, and dual respiratory and fermentative capacities. *Appl. Environ. Microbiol.*,

779  *85*(10), e00110--19.

780 Zhou, Z., Liu, Y., Xu, W., Pan, J., Luo, Z.-H., & Li, M. (2020). Genome-and Community-

781  Level Interaction Insights into Carbon Utilization and Element Cycling Functions of

782  Hydrothermarchaeota in Hydrothermal Sediment. *MSystems*, *5*(1).

 Figure legends

 Figure 1. CAIR comparisons through the tree of life.

(A) The most updated tree of life comprising all second hierarchies stemming from cellular organisms. For bacteria superkingdom, the second hierarchy includes all 56 bacterial phyla; Candidate Phyla Radiation (CPR) is included as one separate phylum. For Archaea and Eukaryota superkingdoms, the second hierarchy encompasses 20 archaeal phyla and 16 eukaryote supergroups and divisions. On each bifurcation point of the tree, the arbitrary number corresponds to the test number and the associated plots in panel B. The blue and light green colours indicate the superior and inferior arm of the bifurcation point, respectively, which correspond to the left and right sides of the violin- and box-and-whisker plots. The red colour indicates a bifurcation point with both arms from which at least one arm contains less than ten organisms and thus the Brunner-Munzel test would not be reliable. The numbers in parentheses designate the total number of available complete proteomes in each group. **(B)** Violin plots of each bifurcation point in the tree of life, except for those in red. The vertical axes refer to the CAIR in all plots. The box-and-whiskers are overlaid within each violin plot, and the white dashed line in each box indicates the CAIR mean in the corresponding group of organisms. None of the outliers were excluded from the analysis. Asterisks after each test number indicate the significance level of tests. ES; estimated effect size. CI; 95% confidence interval.

Figure 2. A simulation of natural selection shows the selection's bias towards higher CAIRs.

(A) CAIR density plot of the organisms considering the interquartile range (IQR) of phyla sizes through the tree of life after removing the tiny phyla with less than 10 organisms (Q_1-Q_3 16.5 – 171). From the 27 phyla within the IQR, random sampling was performed with a size of $Q_1 \sim 17$. IQR inclusion and the subsequent sampling was done to remove the size effects of populated phyla on the density plot. The mean of phyla skewness is -0.82 in Q_0-Q_4 and -0.74 in Q_1-Q_3. **(B)** CAIR simulation of the tree of life with 27 negatively skewed normal distributions. The means of these simulated random distributions equal to the respective medians of the 27 phyla (marginal rugs) explained in panel A. The function was iterated 1000 times to find the skewness in which the Kolmogorov-Smirnov (KS) test has the maximum *p*-value. The simulation revealed a skewness of -0.90. **(C)** Both panels of A and B are overlaid with a lesser bandwidth to show the details of the distributions. KS test reveals a *p*-value of 0.40 not rejecting the null hypothesis that distributions are identical. Negative skewness of the red wave (depicting phyla data) suggests that the natural selection is biased in favour of organisms with higher CAIRs. Q; quartile.

Figure 3. From the whole life to *E. coli* as an exemplary organism illustrating the CAIR density of proteomes in several ranks of taxonomy.

(**A**) The 'wave of life' denotes the CAIR density of proteomes through the tree of life. As it is argued, the wave of life is proposed to be a summation of skewed distributions. As a result, the wave of life holds an interesting property; namely, zooming in on the wave of life by narrowing the range on the horizontal axis reveals the members in the next rank of taxonomy. In all panels, the peaks of the plots are named according to the most abundant phyla with the closest median to the peak. (**B**) CAIR density of the proteomes in the Proteobacteria phylum. (**C**) Zooming in further on panel B and narrowing the range of horizontal axis to the distribution of organisms under Gammaproteobacteria class, i.e. CAIRs of 0.900 – 0.936, reveals the taxonomic orders. (**D-G**) Proceeding further to zoom in on the peaks of the previous plot shows a perfect agreement with all taxa lineage. Lastly, several strains of *E. Coli* form a bell-shaped distribution, as shown in panel G.

Figure 4. Comparisons of disease indicators in three occurrence groups across gene age categories.

(A) EMIP increases significantly as the occurrence of diseases increases. Generally, EMIP has the highest level of significance among disease indicators. Not surprisingly, the trend of EMIP is also increasing as the genes age. The decline of EMIP in primordial gene age eras are due to the eukaryotic branching and the nucleic genetic material. **(B)** The number of interactions is the second-best indicator of the disease occurrence category. Similarly, as expected, the trend of interactions is increasing as genes grow older. Evolution brings new interactions and adds new nodes to the network. Similarly, there is a decline in the number of interactions in the last two gene age eras. **(C)** The bigger the size of a protein, the more likely it is to be involved in a disease. Also, it is noticeable that the gene ages correlates positively with the protein size in both eukaryotic and prokaryotic settings. However, the disparity between these two settings is easily discernible. **(D)** Unlike what is expected, extremely rare diseases account for the proteins with the highest CAIRs. This observation has been further elucidated in the Discussion section. Nonetheless, the trend of gene ages agrees with the expectation as the complexity of proteins increases with age. Error bars illustrate mean ± 95% confidence intervals, and the significance test is the Dunnett-Tukey-Kramer pairwise multiple comparison test adjusted for unequal variances and unequal sample sizes.

Figure 5. Likert plots of disease indicators (all classified into 8 ranks) with regard to occurrence categories.

(A) Ranked data of EMIP show a robust relationship with the disease categories. Note that the occurrence of diseases increases as EMIP increases. Because the log-transformed of EMIP (LEMIP) forms a bell-shaped curve, the ranking has been done by the mean and three standard deviations of LEMIP. **(B)** Ranked data of interactions increase with the occurrences, maintaining the order of ranks. However, the number of interactions is not as well correlated to occurrences as EMIP. It is noteworthy in B, D, and E panels, since indicator distributions are inconsistent with a Gaussian distribution, rankings have been accomplished by median and equal percentile intervals. **(C)** The link between gene age ranks and disease categories is satisfactory considering the first and last ranks; however, the overall order of ranks does not match with the order of occurrence categories. Gene age ranks are the same as the eight age groups presented previously in the literature (Liebeskind et al., 2016). **(D)** Among Likert plots, CAIR ranks have shown the least correlation with the disease categories which is in line with Figure 4D. **(E)** The ranks of protein length are associated with disease occurrence categories maintaining the order of ranks, except for the sixth rank.

Figure 6. Protein networks of the proteins with the highest EMIPs.

(A-P) The illustrated networks represent the proteins in Table 5 as the 16 proteins with the

highest EMIPs in the human proteome. The size of each circle represents the EMIP of the

protein. According to the UniProt database, networks illustrated in C, D, I, K, and L panels

are non-disease networks; yet, the rest are involved in at least one disease. The main proteins

are demonstrated in colours other than sky blue, and all other interactors are coloured in sky

blue. The figure has been drawn with the interactions data available at the UniProt website up

to the second-degree interactions. It is noteworthy that the illustrated proteins with the highest

EMIP values are markedly present in various disease networks.

Figure 7. The cone of life and the evolution of diseases.

The cone of life summarizes the probable scenarios for proteins put forward in the discussion

section. It is to be noted that the shape of the cone is schematic and an exemplary instance of

every possible occasion has been schematically illustrated. Also, for the clearness of the

drawing, the dark blue cylinder has been cut in half in order not to block other elements. The

diameter of the cone in any cross section shows the amount of existing protein material in

that given time period. The proteins are born from the circumference of any cone base. As

shown in the figure, a general rule would be that diseases are a result of weak proteins with

immature networks. Details have been depicted in the figure for every protein scenario. It is

to be highlighted that the eukaryotic and prokaryotic proteomes have not been discerned in

the figure. s, size; w/, with.

Figure 8. Detailed steps needed to carry out the presented methodology.

(A) Flowchart shows how the proteins were included, grouped according to their respective

organisms, and the exclusion criteria. **(B)** A similar flowchart explaining the steps used to

integrate human proteome data, PICKLE interactions, and Orphanet diseases. Green

rectangles, steps; purple rectangles, executions; red flags, exclusions.

Tables and their legends

Table 1. Proteome CAIRs of organisms mainly used in biological models and studies.

Organism	CAIR	# of proteins	Organism	CAIR	# of proteins
Arabidopsis thaliana	0.9366	39,364	*Mus musculus* (Mouse)	0.9376	55,398
Caenorhabditis elegans	0.9419	26,850	*Neurospora crassa* [74A]	0.9323	10,257
Chlamydomonas reinhardtii	0.8887	18,829	*Rattus norvegicus* (Rat)	0.9378	29,951
Ciona savignyi	0.9444	20,004	*Saccharomyces cerevisiae* [S288c]	0.9336	6,049
Danio rerio (Zebrafish)	0.9398	46,848	*Schizosaccharomyces pombe*	0.9347	5,141
Drosophila melanogaster (Fruit fly)	0.9390	21,973	*Tetrahymena thermophila*	0.9119	26,976
Escherichia coli [K12]	0.9328	4,391	*Xenopus tropicalis* (Western clawed frog)	0.9410	55,258
Homo sapiens (Human)	0.9392	74,823	*Zea mays* (Maize)	0.9341	99,254
Medicago truncatula (Barrel medic)	0.9392	57,065	*Zinderia insecticola*	0.8247	206

The table has been sorted in alphabetical order. Where there were different proteomes of a single organism, the number of proteins refers to that of the most popular proteome used in the literature. Terms enclosed in parentheses are the common names used colloquially, and those enclosed in brackets are the strain names of organisms that have different indexed proteomes for their strains in the UniProt database. Please note that the prevalent model organisms of humans, like rats, mice, fruit flies, and zebrafish are very similar to us in terms of CAIR. #; number.

Table 2. Exact *p*-values of the two-sided Brunner-Munzel tests.

Test no.	*p*-value	Test no.	*p*-value	Test no.	*p*-value	Test no.	*p*-value
T1	5×10^{-137}	T18	2.72×10^{-10}	T42	0.000965	T66	4.29×10^{-08}
T2	3.42×10^{-17}	T19	0.14649	T44	0.35037	T67	0
T3	6.91×10^{-31}	T20	6.98×10^{-07}	T46	0	T68	0
T4	7.42×10^{-131}	T21	0.000417	T47	2.88×10^{-28}	T69	5.64×10^{-32}
T5	2.92×10^{-76}	T26	2.74×10^{-42}	T49	2.51×10^{-07}	T70	0.008022
T6	0	T28	1.66×10^{-05}	T51	7.23×10^{-205}	T73	0.004432
T7	6.41×10^{-47}	T29	0.99445	T54	0.000363	T75	2.71×10^{-263}
T8	0.32303	T32	8.86×10^{-17}	T55	5.04×10^{-07}	T76	7.62×10^{-08}
T9	0.019045	T33	0.14933	T56	5.13×10^{-05}	T77	9.92×10^{-05}
T10	6.98×10^{-08}	T36	3.83×10^{-09}	T57	8.48×10^{-56}	T79	1.30×10^{-08}
T14	7.70×10^{-08}	T38	0.012529	T59	0.87909	T80	7.16×10^{-05}
T15	0.001135	T39	0.030999	T60	0.46556	T81	0.000134
T16	0.010167	T40	5.65×10^{-16}	T62	0.55498	T83	6.78×10^{-89}
T17	2.45×10^{-17}	T41	2.73×10^{-51}	T64	0.002841	T88	1.35×10^{-34}

Test no. refer to the test labels illustrated in Figure 1. Because there was limited space in the illustration, the exact *p*-values are reported herein. Please note that no test was performed when either one or both groups contained less than 10 organisms. The numbers of available organisms in each phylum are enclosed by parentheses in Figure 1A. no.; number.

905 **Table 3. Narrative data of protein groups in three human disease categories.**

Disease category	Occurrence	Group size	# of interactions median (IQR)	Protein length median (IQR)
Rare diseases	> 1:1,000,000	819	16.0 (36.0)	599.0 (668.0)
Extremely rare diseases	≤1:1,000,000	1,356	11.0 (23.0)	527.0 (487.5)
No diseases	-	16,296	5.0 (15.0)	386.0 (393.0)

Swiss-Prot entries have been matched with their Orphanet cross-references to obtain the total

occurrences classifying diseases in three categories. Please note that extremely rare diseases

are a vast number of diseases whose names are not even familiar to general practitioners as

they mainly consist of case reports from around the world. Rare diseases are a group of

diseases that have been mainly the focus of attention in scientific literature and medical

books. IQR; interquartile range. #; number.

Disease indicator	Rank 1	Rank 2	Rank 3	Rank 4	Rank 5	Rank 6	Rank 7	Rank 8
LEMIP[a]	[-9, 36.45]	(36.44, 44.94]	(44.94, 53.44]	(68.38, 61.94]	(61.94, 70.44]	(70.44, 78.93]	(78.93, 87.43]	(87.43, 104.00]
Number of interactions[b]	[0]	[1]	(1, 3]	(3, 6]	(6, 11]	(11, 19]	(19, 39]	(39, 2136]
Protein length[b]	[2, 161]	(161, 250]	(250, 328]	(328, 415]	(415, 518]	(518, 671]	(671, 968]	(968, 34350]
CAIR[b]	[0.0217, 0.8803]	(0.8803, 0.8982]	(0.8982, 0.9088]	(0.9088, 0.9164]	(0.9164, 0.9230]	(0.9230, 0.9290]	(0.9290, 0.9351]	(0.9351, 0.9567]
Gene age[c]	Mammalia	Vertebrata	Eumetazoa	Opisthokonta	Eukaryota	Euk_Archaea	Euk+Bac	Cellular_organisms

Table 4. Eight ranks of disease indicators and their quantitative intervals.

a. Because of the bell-shaped curve distribution, mean and standard deviations have been used to classify ranks. b. Because of the non-normal distribution, median and percentiles have been used to classify ranks. c. Ranks are based on scientific literature. LEMIP; Log_e of Estimation of Mutual Information of Proteins. CAIR; Calculated Average Information per Residue.

Entry	Gene	Protein name	Length	CAIR	# of ints	EMIP	Disease
Q8WZ42	TTN	Titin (EC 2.7.11.1) (Connectin)	34350	0.9190	119	33839	Yes
P05067	APP	Amyloid-beta precursor protein (APP)	770	0.9309	2136	23441	Yes
P0CG48	UBC	Polyubiquitin-C [Cleaved into: Ubiquitin]	685	0.8785	916	14231	No
Q8WXI7	MUC16	Mucin-16 (MUC-16) (Ovarian cancer-related tumor marker CA125)	14507	0.8487	3	12440	No
Q9NRI5	DISC1	Disrupted in schizophrenia 1 protein	854	0.9002	650	11312	Yes
P04637	TP53	Cellular tumor antigen p53 (Tumor suppressor p53)	393	0.9250	787	10222	Yes
Q09472	EP300	Histone acetyltransferase p300 (p300 HAT)	2414	0.9145	541	9730	Yes
P00533	EGFR	Epidermal growth factor receptor (EC 2.7.10.1)	1210	0.9439	679	9697	Yes
P62993	GRB2	Growth factor receptor-bound protein 2 (Adapter protein GRB2)	217	0.9387	615	8704	No
Q8NF91	SYNE1	Nesprin-1 (Enaptin)	8797	0.9058	31	8583	Yes
P63104	YWHAZ	14-3-3 protein zeta/delta (Protein kinase C inhibitor protein 1)	245	0.9018	517	8282	No
P78362	SRPK2	SRSF protein kinase 2 (EC 2.7.11.1) (SFRS protein kinase 2)	688	0.9278	441	8161	No
Q03001	DST	Dystonin (230 kDa bullous pemphigoid antigen)	7570	0.9152	58	7806	Yes
Q5VST9	OBSCN	Obscurin (EC 2.7.11.1)	7968	0.9118	14	7769	Yes
P38398	BRCA1	Breast cancer type 1 susceptibility protein (EC 2.3.2.27)	1863	0.9160	456	7655	Yes
Q15149	PLEC	Plectin (PCN) (PLTN)	4684	0.8815	138	7611	Yes

916   **Table 5. Proteins with the highest EMIP values.**

917   Entry is the UniProt entry of the protein; Gene is the preferred gene names used in the literature; Length denotes to the length of the protein

918   sequence; # of ints is the number of interactions adapted from PICKLE database; Disease is represented as a dichotomous variable adapted from

919   UniProt's 'Involvement in disease' column.

A **Connectin**

B **Amyloid-beta precursor protein**

C **Polyubiquitin-C**

D **Ovarian carcinoma antigen CA125**

E **Disrupted in schizophrenia 1 protein**

F **Cellular tumor antigen p53**

G **Histone acetyltransferase p300**

H **Epidermal growth factor receptor**

I **Growth factor receptor-bound protein 2**

J **Nesprin-1**

K **Protein kinase C inhibitor protein 1**

L **SRSF protein kinase 2**

M **Dystonin**

N **Obscurin**

O **Breast cancer type 1 susceptibility protein**

P **Plectin**

● Connectin	● Amyloid-beta precursor protein	● Polyubiquitin-C	● Ovarian carcinoma antigen CA125
● Disrupted in schizophrenia 1 protein	● Cellular tumor antigen p53	● Histone acetyltransferase p300	● Epidermal growth factor receptor
● Growth factor receptor-bound protein 2	● Nesprin-1	● Protein kinase C inhibitor protein 1	● SRSF protein kinase 2
● Dystonin	● Obscurin	● Breast cancer type 1 susceptibility protein	● Plectin
● Other proteins	— Indicative of a protein-protein interaction		

Provided that the network is ideally mature, the brown protein is a non-disease

Current era

Always a non-disease

After years, the same protein is the cause of the same disease; however, w/ an extremely rare incidence

Potentially a rare disease protein

Non-disease, if ideally matured

Always a non-disease

Three exemplary non-essential proteins are born while adding to cone radius

An exemplary extinction era

The non-essential protein is extinct

During the illustrated cross section era, the red protein is likely to cause a rare disease

An exemplary cross section of an evolutionary era

A non-essential protein is born

Time

Light blue area encompasses proteins that are not essential for life essence

s

Dark blue cylinder encompasses the essential proteins for life essence

By time, proteins grow in size and CAIR, leading to new interactions, in case the protein is not ideally robust

Essential proteins meet the threshold for the essence of life. Life begins here.

s

Pre-life building blocks

In general, proteins are born with a small size, less CAIR, and few interactions

Essential proteins lacked for the beginning of life

Old weak protein w/ mature network	Old weak protein w/ immature network	Old robust protein scenario
Young weak protein w/ mature network	Young weak protein w/ immature network	Young robust protein scenario
Extinct protein scenario	A single protein	Protein-protein interaction

A

B

Diseases were filtered out based on the below criteria:

1) A geodistribution other than "Worldwide", *or*
2) An occurrence value of N/A, "Unknown", or "Not yet documented", *or*
3) An occurrence type of "Lifetime prevalence", or "Case/families"

15,459 diseases from Orphanet database were included

54,473 TrEMBL and 20,350 Swiss-Prot human entries were included

191,113 binary interactions were included from PICKLE database

3,336 remaining Orphanumber were integrated with the data

EMIP and CAIR for all organisms were calculated

Total occurences for all entries were calculated

54,473 TrEMBL entries were excluded

Age-independent analyses were executed

20,350 Swiss-Prot entries remained

Entries without a gene age category were excluded

19,305 proteins were further analyzed