1

A computational model of homeostatic cerebellar compensation of ageing in vestibulo-ocular reflex adaptation

Niceto R. Luque [1, 2, *], Francisco Naveros[2, *], Eduardo Ros[2], Angelo Arleo[1]

[1] Sorbonne Université, INSERM, CNRS, Institut de la Vision, 17 rue Moreau, F-75012 Paris, France

[2] Department of Computer Architecture and Technology, University of Granada (CITIC), Granada, Spain

* NL and FN share the first authorship

Corresponding author: Niceto R. Luque, nluque@ugr.es

Declaration of interest: The authors hereby declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Abstract

The vestibulo-ocular reflex (VOR) stabilises vision during head motion. Age-related

structural changes predict a linear VOR decay, whereas epidemiological data show a

non-linear temporal profile. Here, we model cerebellar-dependent VOR adaptation to link

structural and functional changes throughout ageing. We posit that three neurosynaptic

factors codetermine VOR ageing patterns: electrical coupling between inferior olive

neurons, intrinsic plasticity at Purkinje cell synapses, and long-term spike timing

dependent plasticity at parallel fibre - Purkinje cell synapses as well as mossy fibre -

medial vestibular nuclei synapses. Our cross-sectional simulations show that long-term

plasticity acts as a global homeostatic mechanism mediating the non-linear temporal

profile of VOR. Our results also suggest that intrinsic plasticity at Purkinje cells acts as a

local homeostatic mechanism sustaining VOR at old ages. Importantly, longitudinal

simulations show that residual fibres coding for the peak and trough of the VOR cycle

constitute a predictive hallmark of VOR ageing trajectories.

Keywords: vestibulo-ocular reflex (VOR), VOR ageing, cerebellar adaptation, spiking

neural networks, spike timing dependent plasticity, intrinsic plasticity, electrical

synapses.

1. Introduction

Healthy ageing progressively degrades postural control, balance, and spatial orientation (Anson and Jeka, 2016, Brandt et al., 2005, Zalewski, 2015). The consequent loss of static and dynamic balance hinders older adults' autonomy and it increases their risk of fall (Piirtola and Era, 2006, Desai et al., 2010, Tinetti, 2003). Postural control is an adaptive process that involves biomechanical reflexes, sensorimotor functions, as well as attentional and cognitive faculties (Horak, 2006). The sensory modalities that concurrently describe the spatiotemporal dynamics of body's position and orientation in space, include visual, vestibular, and somesthetic afferents (Lackner and DiZio, 2005, Arleo and Rondi-Reig, 2007, Cullen, 2012). The integration of these multisensory signals mediates postural control, which relies upon both body and gaze stability (Mergner and Rosemeier, 1998). Here, we focus on the vestibular control of eye movements, ensuring gaze stability during head motion (Grossman and Leigh, 1990). In particular, the vestibulo-ocular reflex (VOR) generates rapid contralateral eye movements that stabilise images on the retinal fovea during head displacements (Fig. 1A). The VOR plays a key role in maintaining balance and spatial orientation. Age-related deficits in VOR can impair visual acuity and they can lead to oscillopsia (i.e., a perturbing illusory oscillation of the visual scene) during locomotion (Demer et al., 1994).

There is evidence that healthy ageing decreases the VOR gain, i.e. the ratio between the antagonist changes of eye and head angles during head displacements (Baloh et al., 1993, Demer et al., 1994, Demer et al., 1993, Baloh et al., 2001, Agrawal et al., 2013, Li et al., 2015, Anson et al., 2016). Studies in the early 90s shed light on the impact of ageing on rotatory VOR (r-VOR) (Baloh et al., 1993, Peterka et al., 1990). A decline in the r-VOR function was observed in older adults in response to low frequency sinusoidal

rotations (<1 Hz; Peterka et al., 1990) as well as to high-amplitude and high-velocity

sinusoidal rotations (Paige, 1992).Subsequent epidemiological studies reported

discordant patterns of results on age-related VOR deficits, due to differences in VOR

measures, tested head motion frequencies, age range, and sample size (Peterka et al.,

1990, Paige, 1994, Furman and Redfern, 2001, McGarvie et al., 2015, Anson and Jeka,

2016, Li et al., 2015, Matiño-Soler et al., 2015). For instance, (Li et al., 2015) conducted

a cross-sectional VOR evaluation on a study population of 110 community-dwelling

adults spanning a broad age range (26-92 years, from the Baltimore Longitudinal Study

of Aging cohort). They tested the VOR function by employing the video head-impulse

testing (vHIT), which provides a specific clinical assessment of the peripheral vestibular

system. They found is that the r-VOR gain remained stable across participants aged up

to about 79-80 years, whilst it significantly declined with age afterwards (Li et al., 2015).

In contrast, McGarvie et al. (2015) reported an unchanged VOR gain in older adults of

80-89 years of age, by using the vHIT for all six semi-circular canals. Similarly, Matiño-

Soler et al. (2015) observed a stable r-VOR until 90 years of age and a decline

afterwards. Thus, despite their discrepancies about when the VOR impairment actually

occurs in older adults, all these epidemiological studies consistently found a non-linear

VOR profile as a function of age, with the VOR gain remaining unchanged until 80,

maximum 90 years of age, and declining thereafter. That is, the temporal dynamics of r-

VOR modulation across the lifetime involve a first phase of steady, quasi linear

persistence of the gain, and a second phase characterised by an abrupt drop in

performances (with the cut-off occurring at 80-90 years of age).

Age-related VOR loss depends on numerous neuroanatomical properties of the

vestibular system that degenerate with age (Allen et al., 2017, Anson and Jeka, 2016).

The number of vestibular receptors (i.e., hair cells) decreases at a rate of about 6% per

decade, tending to degenerate from middle age on, independently from pathology (Baloh

et al., 1989, Bergström, 1973). Also, the number of neurons in the vestibular nuclei

undergoes a 3% loss per decade, starting at approximately 40 years of age (Alvarez et

al., 2000, Lopez et al., 1996). As a consequence, throughout ageing, fewer primary

vestibular afferents reach the brain, in particular the downstream control structures

responsible for VOR adaptation, such as the cerebellum (Allen et al., 2017). Thus, ageing

gradually hinder the detection and encoding of head displacements (in particular head

rotations; Hirvonen et al., 1997). It has been suggested that neural compensatory

factors (including increased sensitivity to afferent nerve fibres) may counterbalance age-

related vestibular losses, thus preserving, to a certain extent, VOR in older adults (Jahn

et al., 2003, Li et al., 2015, McGarvie et al., 2015). However, to the best of our knowledge,

there are neither experimental nor theoretical explicit hypotheses about the

compensatory mechanisms that synergistically determine the observed non-linear VOR

profile across lifetime.

Here, we propose a model of cerebellar-dependent VOR adaptation to cross-link

neuroanatomical and functional aspects of gaze stabilisation during head rotation. The

model aimed at both reproducing epidemiological data and making testable predictions

about the interplay of neuronal and plasticity mechanisms at stake during ageing. We

hypothesised that three neuro-synaptic factors are critical during VOR ageing: *(i)*

Electrical synaptic coupling between inferior olive neurons (through gap junctions)

(Llinas et al., 1974, Sotelo et al., 1974). Electrical synapses determine the synchronicity

level of the inferior olive network, thus shaping its oscillatory dynamics(Lefler et al.,

2020). We considered the impact of age-related changes in the electrical coupling of

inferior neurons, due to degrading GABAergic inputs from medial vestibular nuclei(Lefler

et al., 2014, Best and Regehr, 2009). Because inferior olive cells are assumed to encode

retina slips during gaze stabilisation (i.e., error signalling), we evaluated to what extent

age-related changes in the electrical coupling of inferior olive neurons could impact the

VOR function. *(ii)* Intrinsic plasticity occurring at Purkinje cell synapses (Jang et al., 2020,

Shim et al., 2018), which regulates the excitability of Purkinje neurons (by modulating

their membrane capacitance) to adapt their response to synaptic morphological changes

(Zhang et al., 2010, Andersen et al., 2003). We investigated the possible role of intrinsic

plasticity at Purkinje cell synapses as a *local* homeostatic process, i.e., compensating

for decreasing levels of vestibular afferent signals (which reach Purkinje cells through

cerebellar granule cells) as well as for electro-responsiveness changes induced by

ageing in Purkinje cells themselves (Zhang et al., 2010, Andersen et al., 2003). *(iii)* Long-

term plasticity operating at different synaptic sites of the cerebellar circuit (Luque et al.,

2019, Gao et al., 2012) Long-term potentiation (LTP) and depression (LTD) mechanisms

are instrumental to cerebellar-dependent sensorimotor adaptation (D'Angelo et al.,

2016). We explored their role as a *global* homeostatic compensatory process, enhancing

neural sensitivity to the remaining sensory inputs during ageing.

First, we examined these tree neuro-synaptic processes independently from each other,

assessing their individual impact on VOR adaptation as a function of age. Second, we

simulated cross-sectional and longitudinal studies to explore how these factors may

codetermine the non-linear VOR temporal pattern observed throughout ageing. Third,

we sought to identify the factors responsible for the interindividual variability observed

during ageing (i.e., underlying different VOR ageing trajectories across study

populations). We tested the hypothesis that the variability in terms of adaptive

compensation to residual fibres/connections might partially explain the apparent

discrepancies between different epidemiological outcomes.

133   ***** *Fig 1 about here* *****

134 .

2. Results

Cerebellar-dependent VOR adaptation

We framed cerebellar-dependent VOR adaptation within a forward control scheme (Fig. 1B; Lorente de Nó, 1933, Santina et al., 2001, Luque et al., 2019). Computationally, the model reproduced the main properties of the cerebellar circuit, and it consisted of five neural networks (Fig. 1C). A population of 100 mossy fibres (MFs) conveyed primary vestibular inputs (signalling head angular accelerations) onto the cerebellar network. MFs projected excitatory afferents onto both 200 medial vestibular nuclei (MVN) and 2000 granular cells (GCs). GCs generated a sparse representation of the MF inputs and they transmitted the encoded sensory information to 200 Purkinje cells (PCs), through excitatory projections. An intrinsic plasticity mechanism (Jang et al., 2020, Shim et al., 2018) regulated the excitability of model PCs, consistently with electrophysiological recordings (Shim et al., 2017, Turrigiano et al., 1994) (see Methods). PCs integrated the afferent signals from PFs (i.e., the axons of GCs), which elicited Purkinje simple spikes (i.e., tonic firing mode, see Luque et al., 2019). PCs also integrated the error-related signal from climbing fibres (CFs), i.e., the axons of inferior olive (IO) cells. CFs, assumed to code for retina slips, (Luque et al., 2019, Ito, 2013, Naveros et al., 2019) elicited Purkinje complex spikes (i.e., bursting mode). The responses of PCs (either simple or complex spiking) inhibited MVN cells, which also integrated inputs from MFs and IOs to generate the cerebellar output controlling eye movements (Fig. 1C). The CF-PC-MVN subcircuit comprised two symmetric microcomplexes, controlling leftward and rightward eye compensatory rotations, respectively (see Methods). Long-term plasticity (LTP and

LTD) modulated PF-PC and MF-MVN synapses (Clopath et al., 2014, Badura et al.,

2016), whereas the remaining synaptic connections in the model were either non plastic

or electrical, as between IO cells (Fig. 1C; see Methods).

We assessed cerebellar-dependent r-VOR adaptation by means of a 1 Hz sinusoidal

head rotation protocol (i.e., within the natural head rotation range [0.05-5 Hz], Leigh and

Zee, 2015). During 2500 s of simulation (Fig. 2), LTP/LTD mechanisms shaped PF-PC

and MF-MVN synaptic efficacies (which were randomly initialised) to adapt the r-VOR

gain such as to reduce the error signalled by IO cells. After about 1000 s, the r-VOR gain

(averaged over 40 simulated individuals) plateaued at 0.95 (Fig. 2A), a value consistent

with experimental VOR data in humans under 1 Hz sinusoidal head rotations (Dits et al.,

2013). Retina slip errors, as encoded by firing of IO cells, decreased as VOR accuracy

improved (Fig. 2B).

170   ***** *Fig 2 about here* *****

Impact of age-related vestibular loss on the electrical coupling of Inferior

Olive neurons

Since IO neurons conveyed error-related signals mediating cerebellar sensorimotor

learning, we tested the impact of age-related vestibular loss on IO activity. In the model,

IO neurons formed an electrically coupled network, whose recurrent dynamics was

regulated by the PC-MVN-IO cerebellar loop (Fig. 1C). In particular, the inhibitory action

of MVN modulated the strength of IO electrical coupling, which in turn determined the

synchronicity of IO firing (Najac and Raman, 2015, Lefler et al., 2014, Best and Regehr,

2009). A strong IO electrical coupling (i.e., a highly synchronised IO network) would allow

large errors to be transmitted to PCs, eliciting fast VOR learning (Tokuda et al., 2013,

Schweighofer et al., 2013). A reduced IO electrical coupling would lead to slower but

more accurate VOR adaptation (e.g., during late learning).

We sought to understand how a progressive age-related decrease of the MVN

GABAergic input to IO neurons (due to vestibular primary afferent loss) would impact

their network activity (Lefler et al., 2014, Best and Regehr, 2009). We simulated two age

groups (20 young subjects: 20 yo; 20 older subjects: 100 yo) and we linearly decreased

the inhibitory MVN input to IO as a function of age (from a maximum at 20 years to zero

at 100 years, see Methods). We compared the dynamics of IO spatiotemporal firing

patterns in a 5x5 lattice configuration (Nobukawa and Nishimura, 2016), when an error-

related pulse activated the central IO neuron of the network (neuron 1 in Supp. Fig. 1A).

The electrical coupling between IO neurons produced a rapid transient propagation

within the network, eliciting a sequential bursting of IO cells along the radial outward

direction of the lattice (Supp. Figs. 1A, B). When comparing IO network propagation

patterns across age, we found that the central stimulation did elicit more rapid and

pronounced membrane potential variations in the IO lattices of older individuals, which

resulted in simpler on/off network dynamics as compared to young individuals (Fig. 3A

and Supp. Figs. 1B, C). Also, this transient on/off patterns produced a higher mean

activation frequency in older IO networks (Fig. 3A). We quantified the complexity of IO

spatiotemporal patterns by using the Discrete Wavelet Transform (DWT) (Latorre et al.,

2013). DWT considered these patterns as sequences of images (obtained every ms) and

it estimated the compression rate of each image by calculating the DWT coefficients.

High (low) DWT values corresponded to complex (simple) spatial structures of IO

network patterns at a given time. We found that the electrical coupling among IO neurons

in older individuals gave rise to significantly simpler spatiotemporal network activations,

as compared to young individuals (Fig. 3B; ANOVA $F_{(294,16)}=18$, $p < 10^{-7}$). This was

consistent with a more uniform and synchronised activity of older IO neurons, and with

a higher mean frequency (Fig. 3A). The simpler spatiotemporal dynamics of older IO

networks were likely to induce a poorer capacity to encode retina slips. Therefore, we

subsequently tested the impact of this less effective error signalling on VOR

performance.

***** *Fig 3 about here* *****

Impact of age-related vestibular loss on r-VOR performance

First, we investigated the consequences of age-related vestibular degradations on the

VOR function without any compensatory mechanism in the downstream cerebellar

network. To do so, we blocked intrinsic plasticity at PC synapses as well as LTP/LTD at

MF-MVN and PF-PC synapses. We simulated a cross-sectional study over a large-scale

study population of 2400 individuals aged from 40 to 100 years, by taking a group of 40

individuals per each of year of age (i.e., uniform distribution). Each individual underwent

an independent 1 Hz head rotation protocol (during 2500 s, as above). At the beginning

of the ageing simulation, the cerebellar synaptic weights of each 40-year-old individual

were those obtained at the end of r-VOR learning (Fig. 2). Then, a loss of primary

vestibular afferents took place as a function of age, based on a degeneration rate of 3%

MVN neurons per decade (Lopez et al., 1996, Alvarez et al., 2000). The loss of MVN

neurons induced a change in the MVN-IO inhibitory action, which in turn gradually

increased the electrical coupling within the IO network (Najac and Raman, 2015, Lefler

et al., 2014, Best and Regehr, 2009). The age-related degradation of vestibular primary

afferents also translated into a loss of 0.3% MF-MVN connections per year, as well as

0.3% MF-GC projections per year (starting at 40 yo). In addition, the simulated ageing

accounted for a loss of approximately 6% of GCs per decade (Baloh et al., 1993,

Bergström, 1973, Renovell et al., 2001, Viswasom et al., 2013), which engendered a

degradation of 0.6% of PF-PC connections per year. Each of the 2400 individuals

independently lost a number of *randomly selected* fibres and neurons as a function of

age, based on the above degeneration rates.The results of the ageing simulation showed

a steady decline of the VOR function (Figs. 3C-E), with the accuracy of the r-VOR gain

significantly impaired. Across the study population, the VOR gain declined quasi-linearly

as a function of age (Fig. 3E), in contrast to actual epidemiological data (e.g., Li et al.,

2015).

Intrinsic plasticity at Purkinje cell synapses as a local homeostatic

mechanism

The detailed Purkinje cell (PC) model reproduced the three characteristics spiking

patterns observed experimentally (Fig. 4A): simple spiking response (i.e., tonic firing at

10-250 Hz), complex spiking response (i.e., bursting activity up to 600 Hz), and post-

complex spike pauses. We previously showed that Purkinje spike burst-pause dynamics

are likely to play a key role in VOR adaptation and reversal learning (Luque et al., 2019).

Here, we investigated the consequences of age-dependent changes of PC excitability

on the VOR function. With ageing, the number and the surface of PC synapses decrease

significantly (Zhang et al., 2010). We reasoned that intrinsic plasticity could adapt the

response of PCs during ageing (thus acting as a local homeostatic mechanism). The

membrane capacitance of model PCs decreased as a function of age (Li and Li, 2013)

(Fig. 4B; see Methods). This led to an increase of tonic firing rates in older PCs (Fig. 4C),

consistently with electrophysiological data (Zhang et al., 2010). We also assessed the

relation between the duration of post-complex spike pauses and the duration of pre-

complex spike ISIs in model PCs. We realised this measure by incrementally increasing

the PF inputs whilst maintaining the CF stimulation constant (i.e., only ISIs immediately

following complex spikes were considered for this analysis, as in experimental data,

Grasselli et al., 2016). The PC model with the intrinsic plasticity mechanism predicted

that the linear relation between the duration of post-complex spike pauses and the

duration of pre-complex spike ISIs would be preserved during ageing (Fig. 4D; R^2 =

0.9932; $p < 10^{-4}$).

We then ran a second ageing simulation to test to what extent PC intrinsic plasticity may

operate in the presence of vestibular loss. We mimicked again a cross-sectional study

by taking a sample of 2400 individuals (age range: 40-100 yo; 40 individuals per each

year of age). Each individual underwent the same VOR adaptation protocol (1 Hz

sinusoidal rotation during 2500 s). The initial conditions (in terms of cerebellar synaptic

weights) corresponded to those obtained after r-VOR learning per each independent

individual (Fig. 2A). Age-dependent vestibular (and MVN) degeneration translated into a

loss of 0.3% and 0.6% MFs and PFs per year, respectively. As a consequence of MVN

loss, the IO electrical coupling progressively increased with age (Najac and Raman,

2015, Lefler et al., 2014, Best and Regehr, 2009). All LTP/LTD mechanisms at MF-MVN

and PF-PC synapses were blocked, such as to isolate the effect of the local homeostatic

mechanism provided by intrinsic plasticity at PCs. We found that the increasing

excitability of PCs could partially counterbalance the decreased depolarising currents

elicited by PFs throughout ageing. This resulted in a quasi-linear decrease of the r-VOR

gain across lifetime, along with an increasing interindividual variability (Fig. 4E)

275   ***** *Fig 4 about here* *****

Cerebellar spike-timing dependent plasticity as a global homeostatic compensatory mechanism

Spike-based plasticity mediated LTP/LTD at PF-PC and MF-MVN synapses during r-VOR learning (Fig. 2). We tested whether this adaptation mechanism could enhance the sensitivity of PCs and MVN to degraded input signals during ageing. First, we analysed the weight distributions at PF-PC and MF-MVN synapses after r-VOR learning as a function of age. We compared the synaptic weights of simulated young and older individuals (20 and 100 yo, respectively). In both age groups, cerebellar learning led to anti-symmetric weight distributions at both PF-PC and MF-MVN synapses (Fig. 5), corresponding to the two microcomplexes that controlled rightward and leftward eye movements. Expectedly, the inhibitory action of PCs onto MVN generated opposite weight patterns at PF-PC as compared to MF-MVN synapses (Figs. 5A, B *vs.* 5C, D). In older individuals, an increase of the weights of the remaining fibres compensated the loss of vestibular afferents (Figs. 5B, D). When comparing the distributions obtained by the normalised sums of synaptic weights across PFs (i.e., to estimate the input drive received by PCs), we found platykurtic-like distributions in older individuals (Figs. 5B,D) as compared to more leptokurtic profiles in young individuals (Figs. 5A,C). The ratio between the number of saturated synaptic weights and the number of active afferents increased significantly with age: 28% in young vs. 64% in older PF-PC synapses (Fig. 5A vs. 5B); and 21% in young vs. 31% in older MF-MVN synapses (Fig. 5C vs 5D). Consequently, the neural drive (defined as the area obtained by convolving a unitary impulse signal with the weight distributions) increased significantly with age: it was 2.64 times larger in the older PF-PC synaptic distribution, and 1.64 times larger in the older MF-MVN synaptic distribution, as compared to younger individuals, respectively.

***** *Fig 5 about here* *****

We then ran a third cross-sectional ageing simulation to isolate the role of spike-based

cerebellar LTP/LTD in preserving VOR accuracy across lifetime (i.e., by blocking intrinsic

plasticity at PC synapses). Again, we considered a study population of 2400 individuals

(age range: 40-100 years; 40 individuals per each year of age) and we applied the same

VOR protocol (1 Hz head rotation during 2500 s). We found that compensatory LTP/LTD

at PF-PC synapses increased the input drive to PCs from 40 to 60 years (Fig. 6A).

However, from 60 to 75 years, increasing the neural sensitivity of PCs only could not

maintain the same level of neural drive. Thus, a rapid boost of neural sensitivity at MVN

synapses sustained the neural drive (Fig. 6B). Beyond 80 years of age, the neural inputs

driving both PC and MVN responses could no longer be maintained against age-related

vestibular losses (Figs. 6A,B), which impaired the learning of sensorimotor associations

underpinning VOR accuracy. Accordingly, the r-VOR gain remained unchanged until

about 80 years and it declined significantly afterwards, accompanied by higher

performance variations across individuals (Fig. 6C).

***** *Fig 6 about here* *****

Impact of ageing on the VOR function: cross-sectional and longitudinal

analyses

We combined all age-related factors and all compensatory mechanisms examined so far

to assess their synergistic impact on r-VOR adaptation. First, we ran a fourth cross-

sectional ageing simulation by taking again a study population of 2400 individuals (age

range: 40 - 100 yo), with each individual undertaking the same r-VOR adaptation protocol

(1 Hz sinusoidal head rotation during 2500 s). Intrinsic plasticity at PC synapses and

LTP/LTD at MF-MVN and PF-PC synapses provided local and global homeostatic

adaptation, respectively, to age-related alterations. Simulation results suggested that the

above factors interacted to shape the r-VOR function across the lifespan. In agreement

with cross-sectional analyses in humans, the r-VOR performances remained quasi-

stable until 85-90 years of age, after which it declined sharply at a rate of 0.8% per year

(Fig. 7A). The contribution of PCs' intrinsic plasticity allowed the VOR function to be

sustained further in age as compared to when only LTP/LTD adaptation was present

(Fig. 7B). The variability across individuals increased smoothly until 80 years and it then

augmented significantly (up to 4 times larger at 100 years of age, Fig. 7C), consistently

with the tendency observed experimentally (e.g., Li et al., 2015).

333   ***** *Fig 7 about here* *****

Second, we ran a longitudinal ageing simulation. We took a study population of 40

individuals and we emulated a 60-year follow-up for each of them (i.e., from 40 to 100

yo). Again, we considered a loss of 0.3% and 0.6% of MFs and PFs loss per year,

respectively. Also, age-related changes of MVN GABAergic inputs to IO shaped the

electrical coupling within the IO network as before. However, in contrast to the previous

cross-sectional scenario, these neural losses accumulated across the lifetime of each

individual. Expectedly, the VOR performance across the study population remained

quasi-stable until about 85 yo and it declined sharply afterwards (Fig. 7D). The

interindividual variability increased significantly during the last 15 years of age and it

became almost 5 times larger at 100 yo as compared to 85 yo (Fig. 7C).

Finally, we sought to understand the underlying factors determining the difference

between steady and declining VOR trajectories (e.g., thick green curve vs. thick red

curve in Fig. 7D, respectively). Knowing that: *(i)* all individuals of same age had the same

probability of losing vestibular primary afferents, MVN, MFs, and PFs; *(ii)* the

degeneration process affected fibres and neurons based on a random selection; *(iii)* the

local and global homeostatic mechanisms (i.e., intrinsic plasticity and LTP/LTD) operated

equally across all individuals, we reasoned that a possible determinant of the VOR

ageing trajectory could lie in the distribution of the remaining (i.e., post age-related loss)

fibres/synaptic connections. We postulated that the activity of some subsets of the

remaining connections may be more critical than others in terms of information content

for the encoding of sensorimotor associations and then to maintain the VOR function. To

test this hypothesis, we first sorted all individuals on the basis of their VOR gain

performance at 100 years (Fig. 8A; the red and green dots corresponded to the worst

and best aging trajectories of Fig. 7D, respectively). Then, we compared the subsets of

residual connections that were active at specific moments of the VOR cycle across

individuals. We found that the number of remaining connections responsible for the

encoding of the peak and the trough of the eye velocity function (Fig. 8B) correlated

significantly with the VOR performances of 100-year-old individuals (Fig. 8C, right

column; Supp. Fig. S2). That is, the sorting of 100yo individuals based on their residual

VOR performance matched perfectly the sorting of the same individuals based on the

number of residual PFs and MF-MVN projections coding for the sinusoid's peak and

trough (Fig. 8C, right column). Strikingly, this strong correlation held already at 85 years

of age (i.e., the cut-off age between steady and declining VOR trajectories; Fig. 8C,

centre) and even much earlier, at 60 years of age (i.e., numerous years before the

discontinuity time point between "good" and "bad" ageing trajectories; Fig. 8C, left).

Given that the overall number of lost connections was the same across all the study

population, this implied that those individuals that *by chance* had most of the remaining

connections involved in the encoding of those two critical moments in the VOR period

(i.e., between 200-300 ms and 700-800 ms) had the best chances to have stable VOR

performances throughout ageing.

 ***** *Fig 8 about here* *****

3. Discussion

We modelled cerebellar-dependent sensorimotor adaptation to study how ageing alters

the VOR function. The spiking neural model captured the main computations made by

the cerebellum and it reproduced the biophysical properties of PCs, which are core to

sensorimotor learning. The model offered a transparent tool to assess the impact of age-

related vestibular losses on downstream, cerebellar adaptive coding of gaze stability

(during head rotations). The proposed model complements previous qualitative VOR

models that addressed age-related issues (Anson et al., 2016, Baloh et al., 1993, Li et

al., 2015, Paige, 1992, Peterka et al., 1990) by providing a more mechanistic insight into

the factors underpinning VOR ageing. We hypothesised that three neurosynaptic factors

are key to the understanding of the link between age-related anatomical and functional

VOR changes: the electrical coupling between inferior olive neurons, the intrinsic

plasticity at PC synapses, and LTP/LTD at PF - PC as well as MF - MVN synapses.

To test this hypothesis, we ran a series of ageing simulations to single out the role of

these three factors in determining VOR changes throughout ageing. First, we found that

age-related vestibular loss caused the spatiotemporal patterns of the IO network to

become simpler, similar to an on/off ensemble dynamics. This reduced the accuracy of

retina slip (i.e., error) coding, which in turn impaired VOR adaptation. As a consequence,

our first cross-sectional ageing simulation (which isolated the effect of vestibular loss and

IO coupling alteration, with no compensatory mechanism in the downstream cerebellar

network) showed a linear VOR decline as a function of age (i.e., following the steady

vestibular degeneration). This result contrasted epidemiological data in humans, which

rather show a non-linear VOR temporal profile (with VOR remaining stable until 85-90

years of age and declining abruptly afterwards, Li et al., 2015, Matiño-Soler et al., 2015,

McGarvie et al., 2015). Second, we assessed the local homeostatic action provided by

PC intrinsic plasticity, which adaptively increased PC excitability throughout ageing

(countering the decreasing levels of PF afferents, due to the loss of vestibular signals

integrated by cerebellar granule cells). At the level of single PCs, the tonic firing rates

increased with age according to experimental data (Zhang et al., 2010), whereas the

linear relation between the duration of post-complex spike pauses and the duration of

pre-complex spike ISIs did not change during ageing (testable prediction). At the level of

the VOR function, our second cross-sectional simulation (which accounted for age-

related vestibular loss, IO coupling changes, and PC intrinsic plasticity) showed that the

adaptively increasing excitability of PCs could only moderately counter the impaired

encoding of sensory (vestibular) - motor (eye movement) associations, resulting in a

linear decline in VOR accuracy over the years (again, in contrast with actual

epidemiological data). Third, we isolated the impact of LTP/LTD at PF - PC and MF -

MVN connections and we quantified the increase of synaptic weights to adapt the neural

drive of PCs and MVN to degrading input signals. The results from another cross-

sectional ageing simulation (accounting for age-related vestibular loss, IO coupling

changes, and LTP/LTD) captured the two regimes of human VOR performances during

ageing. That is, LTP/LTD sustained the sensorimotor associations underlying the VOR

function by enhancing the neural sensitivity to residual afferent signals throughout ageing

(i.e., it allowed the full synaptic range to be exploited in order to maintain constant the

neuronal drives). However, the compensatory action by LTP/LTD became ineffective in

the presence of significant levels of vestibular losses (i.e., beyond 80 yo), because

synaptic weights saturated in PF-PC and MV-MVN connections.

We ran a fourth cross-sectional simulation to assess how the three neuro-synaptic

factors would concurrently work during VOR adaptation. The results confirmed that the

global homeostatic compensation mediated by cerebellar LTP/LTD was primarily

responsible for the non-linear temporal profile observed in VOR epidemiological data

(e.g., Li et al., 2015). This prediction is consistent with the saturation hypothesis by

(Nguyen-Vu et al., 2017), stating that an intense change of the synaptic strength shall

temporarily prevent further adaptation. They showed that a specific type of pre-training

that desaturates synapses can improve the ability of mutant mice to learn an eye-

movement task. Conversely, they found that a specific procedure that saturates

synapses can impair the learning ability. In our model, the progressive saturation of PF-

PC and MF-MVN synapses limited VOR adaptation, thus impairing the compensatory

action of LTP/LTD in oldest individuals and leading to the non-linear VOR dynamics

throughout ageing. Our results also showed that the local homeostasis implemented by

intrinsic plasticity at PC synapses played a role in further sustaining VOR between 80

and 90 years of age and in attenuating its decline rate afterwards. A compensatory action

related to intrinsic plasticity at PCs was recently reported during long-term VOR

consolidation in mice (Jang et al., 2020). Interestingly, the slope attenuation due to

intrinsic plasticity while compensating VOR decline in the model was within the same

range as that observed in mice (Jang et al., 2020).

Finally, we further exploited the model vantage point to run a longitudinal ageing

simulation. This allowed us to follow individual ageing trajectories over 60 years, in the

attempt to better understand the factors determining interindividual differences across

ageing (i.e., differentiating steady vs. declining VOR trajectories). Strikingly, we found

that the number of remaining PFs and MF-MVN projections coding for the peak and the

trough of the VOR cycle provided a predictive hallmark for the VOR ageing trajectory on

a single-subject basis. That is, those individuals lacking active PF and MF-MVN afferents

in those precise moments could robustly be expected to have more difficulties in VOR

adaptation throughout ageing. This prediction could possibly be tested in animal models

by seeking for those fibres that are most active when the ocular velocity is maximal during

VOR. For instance, the identification of specific cerebellar GCs (and of their PFs) that

are active upon induction of specific stimulation is possible in vivo mice experiments

(Ishikawa et al., 2015).

The model presented here assumed that the GC layer univocally encoded vestibular

(head motion related) signals through the temporal activation of non-overlapping cell

populations during cerebellar VOR adaptation. GCs are thought to encode vestibular

signals into sparse representations allowing interferences across tasks to be minimised

and neuronal resources to be optimised by reducing redundancy (D'Angelo and De

Zeeuw, 2009). The recurrent inhibitory Golgi cells - GC connections suggest that the

granular layer may act as a recurrent dynamic network (Yamazaki and Tanaka, 2005)

Thus, GCs are likely to generate a randomly repetitive network response characterised

by active/inactive state transitions with no repetition of active cell populations (Yamazaki

and Tanaka, 2007). The model also assumed a progressive degradation of vestibular

afferents integrated by the GC layer with ageing (Baloh et al., 1989, Bergström, 1973),

which led to a degradation of PFs, impairing, in turn, long-term PF-PC synaptic

adaptation. Interestingly, neural regeneration can occur at PF-PC synapses thanks to

the Gluδ2 receptor (Ichikawa et al., 2016), whereas Gluδ2 deficits lead to disruption of

LTD at PC synapses and motor impairment in VOR tasks (Pernice et al., 2019, Yuzaki,

2013). Some evidence suggests that neural loss can be related to the absence of Gluδ2

receptor, since the deletion of GluRδ2 expression in mutant mice (GluRδ2ho/ho) induces

PC and GC reduction over lifetime (Zanjani et al., 2016). A gradual decrease on Gluδ2

with ageing would compromise Gluδ2-dependent processes that, in turn, would reduce

intrinsic PC excitability and eventually impair LTD at PCs.

The model also assumed a compromised IO electrical coupling due to degraded

GABAergic afferents from MVN during ageing. The strength of the gap junctions amongst

modelled olivary neurons was asymmetric (Lefler et al., 2014, De Zeeuw et al., 1998).

The level and the direction of this asymmetry was regulated by emulating the GABAergic

feedback (Lefler et al., 2014). The coupling asymmetry allowed for the creation of

different spatial configurations of PC's complex spike patterns. The GABAergic inputs

from MVN could directly cause a transient decrement in electrical coupling amongst IO

cells (Lefler et al., 2014). GABAergic feedback not only temporarily blocked the

transmission of signals through the olivary system but it could also isolate IO neurons

from the network by shunting the junction current (Loewenstein, 2002). In the absence

of GABAergic feedback, electrical coupling was not counteracted and IO network

oscillations were not mitigated but rather increased. There is only indirect evidence for

an age-related degeneration of the GABAergic MVN inputs throughout aging. The r–

aminobutyric acid, GABA, inhibits the formation of lipoxidation end products (Deng et al.,

2010). The presence and accumulation of lipofuscin with ageing, a lipoxidation product,

is an essential part of the traditional theory of ageing (Sulzer et al., 2008). Lipofuscin

accumulates in postmitotic cells with age, impairing their functioning. Its presence is

caused by unbalanced cell metabolic and waste-degradation functions. IO neurons are

relatively immune to apoptosis (Lasn et al., 2001) and they preserve their function with

ageing, although they tend to accumulate significant amount of lipofuscin with

age (Brizzee et al., 1975). It is unclear whether the presence of a large amount of

lipofuscin is due to higher lipofuscin generation and/or decelerated removal (Fonseca et

al., 2005). Since lipofuscin aggregates are unavoidable reactions in biological systems,

the lack of a cycle involving lipofuscin elimination is more plausible than the absence of

lipofuscin generation (Yin, 1996). The r-aminobutyric acid scavenging effects proposed

by Deng et al. (2010) over advanced lipoxidation end products (ALEs) may be

instrumental for lipofuscin clearance in the olivary system. A gradual decline of r–

aminobutyric acid presence with age, may explain the accumulation of lipoxidation

products in IO neurons. MVN GABAergic afferents are the main source of r–aminobutyric

acid for the olivary cells but they also mediate the electrical coupling amongst them. The

gradual degeneration of these GABAergic afferents may explain the gradual presence

of IO lipofuscin as well as the altered activations of IO ensembles with ageing.

4. Methods

Vestibulo-Ocular Reflex (VOR) Model

The VOR was defined as a continuous-time mathematical model with two poles (Eq. 1),

whose parameters were adjusted recursively to fit experimental and clinical data

(Skavenski and Robinson, 1973, Robinson, 1981, Gordon et al., 1989):

511
$$VOR(s) = \frac{E(s)}{H(s)} = \frac{K \cdot T_{c1} \cdot s}{(T_{c1} \cdot s + 1) \cdot (T_{c2} \cdot s + 1)} \cdot e^{-s\tau_{delay}} \qquad (1)$$

Where $e(t), E(s): eye\ motion\ (output),$ and $h(t), H(s): head\ motion\ (input).$

There were 4 parameters in the model: $Q = [K, T_{C1}, T_{C2}, \tau_{delay}]$. The delay parameter

τ_{delay} captured the delay in communicating the signals from the inner ear to the brain and

the eyes. This delay is the consequence of the time needed for neurotransmitters to

traverse the synaptic clefts between nerve cells. Based on the number of synapses

involved in the VOR, the estimate of this delay is of 5 ms (Skavenski and Robinson,

1973, Robinson, 1981). The gain parameter *K,* assumed to be between 0.6 and 1,

modelled the fact that the eyes do not perfectly cope with the movement of the

head(Skavenski and Robinson, 1973, Robinson, 1981). The T_{c1} parameter represented

the dynamics associated with the semicircular canals as well as some additional neural

processing. The canals are high-pass filters, as the neural active membranes in the

canals slowly relax back to their resting position after rotational experimentation (the

canals stop sensing motion). Based on the mechanical characteristics of the canals,

combined with additional neural processing which prolongs this time constant to improve

the VOR accuracy, the T_{c1} parameter was estimated to be around 15 sec, in agreement

with the biologically range which is 10-30 sec (Skavenski and Robinson, 1973, Robinson,

1981). Finally, the T_{c2} parameter captured the dynamics of the oculomotor plant, i.e. the

eye and the muscles and tissues attached to it. Its value was between 0.005 and 0.05

sec.

To find the temporal response for the VOR transfer function, we needed to calculate the

inverse Laplace transform (Eq. 2). The outcome of the inverse Laplace transform

consisted in a differential equation system defined in the same time domain as the

spiking cerebellar network (see below; note that we modelled the delay and we inserted

within the sensorimotor delay).

$$
\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -a_0 & -a_1 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} 0 \\ h(t) \end{bmatrix}
$$

$$
e(t) = \begin{bmatrix} b_0 & b_1 \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}
$$

(2)

537   Where:

$$a_0 = \frac{1}{T_{c1} \cdot T_{c2}}, \quad a_1 = \frac{T_{c1} + T_{c2}}{T_{c1} \cdot T_{c2}}$$

$$b_0 = 0, \quad b_1 = \frac{K \cdot T_{c1}}{T_{c1} \cdot T_{c2}}$$

539 *VOR analysis and assessment.* The periodic functions representing eye and head

540 velocities were analysed through a discrete-time Fourier transform:

$$FFT\, X(k) = \frac{1}{N} \sum_{n=0}^{N-1} x(n) \cdot e^{-j\frac{2\cdot\pi\cdot k\cdot n}{N}}$$

541 $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad$ (3)

$$Forward\, FFT\, x(n) = \frac{1}{N} \sum_{k=0}^{N-1} X(k) \cdot e^{j\frac{2\cdot\pi\cdot k\cdot n}{N}}$$

542 where $x(n)$ indicates the periodic function, and N the number of samples within the

543 considered time window. For each k, the term constituted a harmonic component (the

544 complex version) with amplitude and frequency defined as:

$$Harmonic\; amplitude\; A_k = \frac{X(k)}{N}$$

545 $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad$ (4)

$$Harmonic\; frequency\; f_k = \frac{F_s}{N}$$

546 with F_s denoting the sampling frequency (0.5 KHz). The harmonic distortion values, which

547 indicated the harmonic content of a waveform compared to its fundamental, were

548 negligible. We calculated the **VOR gain** as the ratio between the first harmonic

549 amplitudes of the forward Fourier eye- and head-velocity transforms

$$VOR\; GAIN\; G = \frac{A_1^{eye-velocity}}{A_1^{head-velocity}}$$

$\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad$ (5)

VOR protocols. In rotational chair testing, the subject (mouse, monkey, human) is seated

on a rotatory table (Dumas et al., 2016). Speed and velocity of rotation are controlled

and measured. The subject's head is restrained, assuming that the movement of the

table equals to the subject's head movement. During normal VOR adaptation, a visual

target is given in anti-phase with vestibular stimulation. The eyes must follow the visual

target thus minimising the retinal slip. In the model, the eye output function was defined

as:

558
$$Vestibular \quad stimulation = sin(2 \cdot \pi \cdot t)$$
$$Eye\ output\ function = A_E \cdot sin(2 \cdot \pi \cdot t + \pi \cdot \phi_E)$$
(6)

where the ideal VOR experiment values corresponded to $A_E = -1, \phi_E = 0$ (visual field

fixed).

Cerebellar Network Model

The cerebellar network model consisted of five neural populations (Fig. 1C).

Mossy fibres (MFs). 100 MFs constituted the input to the cerebellar network. Mossy

fibres (MFs) conveyed the sensory signals from the vestibular organ and the eye muscles

onto granule cells (GCs) and medial vestibular nuclei (MVN). MF activity evolved based

on a sinusoidal function (1Hz with a step size of 0.002 ms) to encode head movements

consistently with the functional principles of VOR control (Badura et al., 2016, Clopath et

al., 2014, Arenz et al., 2008, Lisberger and Fuchs, 1978). MF responses consisted of

non-overlapping activations of equally sized neural subpopulations, which maintained a

constant overall firing rate (Luque et al., 2016).

Granular cells (GCs). 2000 GCs operated as a state generator (Yamazaki and Tanaka,

2007, Yamazaki and Tanaka, 2005, Yamazaki and Tanaka, 2009). In the presence of a

constant MF input, the granular layer generated a sequence of non-overlapping

spatiotemporal patterns (i.e., states, Fujita, 1982). The same sequence of 500 states

(each consisting of 4 active GCs per time step of 2 ms) repeatedly activated every 1-sec

during learning (see below).

Purkinje cells (PCs). We modelled a population of 200 PCs, divided into 2 groups of 100

cells to control agonist and antagonist eye muscles, respectively. PCs integrated the

excitatory input from the parallel fibres (PFs), i.e. the axons of GCs, as well as the input

from the climbing fibres (CFs), i.e. the axons of inferior olive (IO) cells. PCs projected

inhibitory connections onto MVN cells, to close the cerebellar loop and generate the VOR

output.

Inferior olive (IO) and climbing fibres (CFs). We modelled 200 IO cells, divided in 2

groups of 100 IO cells for agonist/antagonist muscles, respectively. Each IO cell

projected a CF onto one PC and one MVN cell. IO cells were interconnected via

excitatory gap junctions, whose electrical coupling followed preferred directions (Devor

and Yarom, 2002). The preferred paths were disposed radially from the centre of 5x5 IO

cell subpopulations, as in a square regular lattice network (Nobukawa and Nishimura,

2016). The strength of the electrical coupling, which drove the recurrent dynamics of the

olivary population, was equal between all IO cells of the lattice network (see Table 1). In

terms of external inputs, the IO population received excitatory afferents coding for retina

slips (Clopath et al., 2014).This input reached the centre of each lattice network and it

was generated by a Poisson spiking process process (Boucheny et al., 2005, Luque et

al., 2011b). The IO population also received an inhibitory external input from MVN cells

(Fig. 1C) whose action regulated the IO network synchronisation via electrical coupling

modulation (Lefler et al., 2014, Best and Regehr, 2009). We assumed a progressive age-

related decrease of this inhibitory action based on the progressive age-loss of MVN

neurons (Torvik et al., 1986), which modulated the MVN-IO inhibitory synaptic weight

distribution of each 5x5 IO cell subpopulation. The variance of the Gaussian MVN-IO

weight distribution varied linearly from 0.4 to 1.75 causing a more homogeneous

electrical coupling along each 5x5 IO cell subpopulation whilst ageing.

The error-related inputs (coding for retina slips), combined with the recurrent electrical

coupling modulated by inhibitory MVN inputs, determined the overall activity of the IO

population, which generated the CF bursting output. The probabilistic spike sampling of

retina slips ensured an exploration of the whole error space over trials, whilst maintaining

the CF activity below 10 Hz per fibre (in agreement with electrophysiological data,

Kuroda et al., 2001). The evolution of the error could be sampled accurately even at

such a low frequency (Carrillo et al., 2008, Luque et al., 2011b). A graded representation

of the error signal (Najafi and Medina, 2013) led to a correlation between the intensity of

the sampled instantaneous error and the number of the spikes within the CF burst (Eq.

7):

$$s_{spikes} : [0,1] \subseteq \mathbb{R} \to \mathbb{R}$$
$$\varepsilon \to y = s_{spikes}(\varepsilon)$$

612
$$s_{spikes}(\varepsilon) = \begin{cases} 2 & if \quad 0.25 \le \varepsilon \le 0.50 \\ 3 & if \quad 0.50 \le \varepsilon \le 0.75 \\ 4 & if \quad 0.75 \le \varepsilon \le 0.85 \\ 5 & if \quad 0.85 \le \varepsilon \le 0.95 \\ 6 & if \quad 0.95 \le \varepsilon \le 1.0 \end{cases} \tag{7}$$

We assumed a perfect transmission of bursts from CFs to target PCs, i.e. the number of

spikes in a PC complex spike linearly depended on the number of spikes in the CF burst

(Mathy et al., 2009). The IO transmitted from 2 to 6 CF stimuli, delivered at inter-stimulus

intervals of 2 ms, a range representative of inter-spike intervals recorded in olivary axons

during bursts (Mathy et al., 2009, Davie et al., 2008), depending on the retina slips to be

compensated.

Medial Vestibular Nuclei (MVN) cells. We modelled a population of 200 MVN cells, with

again 2 groups of 100 cells for agonist/antagonist muscles, respectively. Each MVN cell

received an inhibitory afferent from a PC and an excitatory afferent from the IO cell that

was also contacting that PC (Luque et al., 2014, Uusisaari and De Schutter, 2011). MVN

cells also received excitatory projections from all MFs. The subcircuit IO-PC-MVN was

then organised in a single microcomplex. This circuitry arrangement rested upon the

principles of circuit integrity and uniformity on the olivo-cortico-nucleo-olivary loop

(Uusisaari and De Schutter, 2011).

Translation of MVN spike trains into analogue eye motor commands. The MVN output

was translated into analogue output signals by averaging the spiking activity of each

MVN subpopulation (one for each agonist/antagonist group of muscles) (Eqs. 8, 9):

630
$$MVN_i\left(t\right) = \int\limits_{t}^{t+T_{step}} \delta_{MVN_{spike}}\left(t\right) \cdot dt \qquad (8)$$

631
$$MVN_{output}\left(t\right) = \alpha\left(\sum_{i=1}^{100} MVN_{ag,i}\left(t\right) - \sum_{j=1}^{100} MVN_{ant,j}\left(t\right) \right) \qquad (9)$$

where α is the kernel amplitude that normalised the contribution of each MVN cell spike

to the cerebellar output correction (the MVN_{ag} output controlled the agonist muscle,

whilst the MVN_{ant} output controlled the antagonist muscle).

Table 1 summarises the parameters of the cerebellar topology used in the model.

***** *Table 1 about here* *****

Neuronal Models

<u>MVN cell model</u>. We modelled MVN cells as LIF neurons with excitatory (AMPA and

NMDA) and inhibitory (GABA) chemical synapses (Eqs. 10-16).

640
$$C_m \cdot \frac{dV}{dt} = I_{leaky} + I_{external} \tag{10}$$

641
$$I_{leaky} = -g_L \cdot (V + E_L) \tag{11}$$

642
$$I_{external} = -\left(g_{AMPA}(t) + g_{NMDA}(t) \cdot g_{NMDA\_INF}\right) \cdot (V - E_{AMPA}) - \\ -g_{GABA}(t) \cdot (V - E_{GABA}) \tag{12}$$

643
$$g_{AMPA}(t) = g_{AMPA}(t_0) \cdot e^{\frac{t-t_0}{\tau_{AMPA}}} + \sum_{i=1}^{N} \delta_{AMPA,i}(t) \cdot w_i \tag{13}$$

644
$$g_{NMDA}(t) = g_{NMDA}(t_0) \cdot e^{\frac{t-t_0}{\tau_{NMDA}}} + \sum_{i=1}^{N} \delta_{NMDA,i}(t) \cdot w_i \tag{14}$$

645
$$g_{GABA}(t) = g_{GABA}(t_0) \cdot e^{\frac{t-t_0}{\tau_{GABA}}} + \sum_{i=1}^{N} \delta_{GABA,i}(t) \cdot w_i \tag{15}$$

646
$$g_{NMDA\_INF} = \frac{1}{1 + e^{-62 \cdot V} \cdot \frac{1.2}{3.57}} \tag{16}$$

where: C_m denoted de membrane capacitance; V the membrane potential; I_{leaky} the leak

current; $I_{external}$ the external currents; E_L the resting potential; g_L the conductance

responsible for the passive decay term towards the resting potential; w_i the synaptic

weight of the synapses between the neuron i and the target neuron. Conductances g_{AMPA},

g_{NMDA} and g_{GABA} integrated all the contributions received by each receptor (AMPA,

NMDA, GABA) through individual synapses. These conductances were defined as

decaying exponential functions, which were proportionally incremented via w_i upon each

presynaptic spike arrival (Dirac delta function). Finally, g_{NMDA_INF} stand for the NMDA

activation channel. Note that we set the neuron membrane potential to E_L during the

refractory period (T_{ref}), just after reaching V_{thr} (votage firing threshold) (Gerstner and

Kistler, 2002, Gerstner et al., 2014). All the parameters of the neuronal models are shown

in Table 2.

***** **Table 2 about here** *****

Inferior olive (IO) neuronal model. We modelled IO cells as LIF neurons with excitatory

(AMPA) and inhibitory (GABA) chemical synapses as well as with electronic gap

junctions (Llinas et al., 1974, Sotelo et al., 1974) (Eqs. 17-22):

663
$$C_m \cdot \frac{dV}{dt} = I_{leaky} + I_{external} \tag{17}$$

664
$$I_{leaky} = -g_L \cdot (V + E_L) \tag{18}$$

665
$$I_{external} = -g_{AMPA}(t) \cdot (V - E_{AMPA}) - \\ -g_{GABA}(t) \cdot (V - E_{GABA}) - I_{EC} \tag{19}$$

$$g_{AMPA}(t) = g_{AMPA}(t_0) \cdot e^{\frac{t-t_0}{\tau_{AMPA}}} + \sum_{i=1}^{N} \delta_{AMPA,i}(t) \cdot w_i \qquad (20)$$

$$g_{GABA}(t) = g_{GABA}(t_0) \cdot e^{\frac{t-t_0}{\tau_{GABA}}} + \sum_{i=1}^{N} \delta_{GABA,i}(t) \cdot w_i \qquad (21)$$

$$I_{EC} = \sum_{i=1}^{N} w_i \cdot (V - V_i) \cdot \left(0.6 \cdot e^{-\frac{(V-V_i)^2}{50^2}} + 0.4 \right) \qquad (22)$$

where: C_m denotes de membrane capacitance; V the membrane potential; I_{leaky} the leak

current; $I_{external}$ the external currents; E_L the resting potential; g_L the conductance

responsible for the passive decay term toward the resting potential; w_i the synaptic

weight of the synapses between the neuron i and the target neuron. Conductances g_{AMPA}

and g_{GABA} integrated all the contributions received by each chemical receptor (AMPA,

GABA) through individual synapses. These conductances were defined as decaying

exponential functions, which were proportionally incremented via w_i upon each

presynaptic spike arrival (Dirac delta function) (Gerstner and Kistler, 2002, Ros et al.,

2006). I_{EC} represented the total current injected through the electrical synapses

(Schweighofer et al., 1999). V was the membrane potential of the target neuron, V_i the

membrane potential of the neuron i, and N was the total number of input synapses of the

target neuron. Finally, for a correct operation of the electrical synapses, this model

emulated the depolarisation and hyperpolarisation phases of an action potential. The LIF

neuron incorporated a simple threshold process that enabled the generation of a

triangular voltage function (maximum/minimum value V_{peak} /E_L respectively) each time

the neuron fired (Bezzi et al., 2004). All the parameters of the IO neuronal model are

shown in Table 2.

Purkinje cell model. The PC model was the same as in (Miyasho et al., 2001, Luque et

al., 2019, Middleton et al., 2008). It reproduced the three spiking modes of Purkinje cells,

namely tonic, bursting, and spike pauses (Forrest, 2008). The PC model consisted of a

single compartment with five ionic currents and two excitatory (AMPA) and inhibitory

(GABA) chemical synapses (Eqs. 23-27):

691
$$C_m \cdot \frac{dV}{dt} = I_{internal} + I_{external} \tag{23}$$

692
$$I_{internal} = -g_K \cdot n^4 \cdot (V + 95) - g_{Na} \cdot m_0 [V]^3 \cdot h \cdot (V - 50) - \\ -g_{Ca} \cdot c^2 \cdot (V - 125) - g_L \cdot (V + 70) - g_M \cdot M \cdot (V + 95) \tag{24}$$

693
$$I_{external} = -g_{AMPA}(t) \cdot (V - E_{AMPA}) - g_{GABA}(t) \cdot (V - E_{GABA}) \tag{25}$$

694
$$g_{AMPA}(t) = g_{AMPA}(t_0) \cdot e^{\frac{t - t_0}{\tau_{AMPA}}} + \sum_{i=1}^{N} \delta_{AMPA,i}(t) \cdot w_i \tag{26}$$

695
$$g_{GABA}(t) = g_{GABA}(t_0) \cdot e^{\frac{t - t_0}{\tau_{GABA}}} + \sum_{i=1}^{N} \delta_{GABA,i}(t) \cdot w_i \tag{27}$$

where C_m denotes de membrane capacitance, V the membrane potential, I the internal

currents, $I_{external}$ the external currents, and w_i the synaptic weight of the synapses

between the neuron i and the target neuron. Conductances g_{AMPA} and g_{GABA} integrated

all the contributions received by each chemical receptor type (AMPA, GABA) through

individual synapses. These conductances were decaying exponential functions that were

proportionally incremented via w_i upon each presynaptic spike arrival (Dirac delta

function) (Ros et al., 2006, Gerstner and Kistler, 2002) . Finally, g_K was the delayed

rectifier potassium current, g_{Na} the transient inactivating sodium current, g_{Ca} the high-

threshold non-inactivating calcium current, g_L the leak current, and g_M the muscarinic

receptor suppressed potassium current (see Table 3).

The dynamics of each gating variable (n, h, c, and M) followed the Eq. 28:

$$\dot{x} = \frac{x_0[V] - x}{\tau_x[V]}$$ (28)

where x corresponds to variables n, h, c, and M. The equilibrium function was given by

the term $x_0[V]$ and the time constant $\tau_x[V]$ (see Table 3).

The sodium activation variable was replaced and approximated by its equilibrium

function $m_0[V]$. The M current presented a temporal evolution significantly slower than

the rest of variables. Each spike in the neuron generated a fast increase of the M current

that took several milliseconds to return to its stable state. A high M current prevented the

PC from entering in its tonic mode (when the neuron generated spikes due to PFs

activity). A complex spike caused a rapid increase of the M current that depended, in

turn, on the size of the spikelet within the burst. PC tonic mode resumed when the M

current decreased.

We first validated the PC model in the NEURON simulator and then we reduced it to

make it compatible with EDLUT (Luque et al., 2019). In the reduced PC model, we

implemented the I_K and I_{Na} currents through a simple threshold process that triggered the

generation of a triangular voltage function each time the neuron fired (Bezzi et al., 2004).

This triangular voltage depolarisation drove the state of ion channels similarly to the

original voltage depolarisation during the spike generation. The final internal current was

given by Eq. 29:

725
$$I_{internal} = -g_{Ca} \cdot c^2 \cdot (V - 125) - g_L \cdot (V + 70) - g_M \cdot M \cdot (V + 95) \quad (29)$$

726   All the parameters are shown in Table 3.

727   ***** **Table 3 about here** *****

728   *Mossy fibres (MF) & granule cells (GC) models.* MFs and GC neurons were simulated

729   as leaky integrate–and–fire (LIF) neurons, with the same excitatory (AMPA) and

730   inhibitory (GABA) chemical synapses and parameters as in (Luque et al., 2019).

Synaptic Plasticity Models

732   *PC Intrinsic Plasticity.* We equipped the PC model with a mechanism to update the value

733   of the membrane capacitance (Cm) according to Eq. 30:

734
$$\frac{dC_m}{dt} = \frac{-\left(C_m - \frac{\beta}{2}\right)}{\beta \cdot \tau_{IP} \cdot (1 + I_{external})} \quad (30)$$

where τ_{IP} denotes the intrinsic plasticity time constant set to 12 10^3 sec (this large time

constant prevented interferences between intrinsic plasticity and other STDP

mechanisms during the learning process (Garrido et al., 2016); β controls the shape of

the firing rate distribution and it is equal to 1 (see Garrido et al. (2016) for details about

all intrinsic plasticity mechanism parameters). Whenever a spike was elicited, the C_m

variable was updated according to the following equation:

741
$$\Delta C_m = \frac{\varepsilon_{Cm}}{\tau_{IP}} \quad if \ PC \ is \ active \quad (31)$$

where ε_{Cm} = 0.0475 (Garrido et al., 2016) determined the influence of each spike on C_m

. Note that the membrane capacitance of PCs could not diminish below a lower limit (0.77

± 0.17 µF cm−2 where mean ± s.d.; range, 0.64-1.00 µF cm−2; Roth and Häusser,

2001).

PF-PC synaptic plasticity. The model of long-term depression (LTD) and long-term

potentiation (LTP) at PF–PC synapses was the same as in (Luque et al., 2019) and it

followed the Eqs. 32 and 33:

749
$$LTD\ \Delta w_{PF_j-PC_i}(t) = \int_{-\infty}^{IO_{spike}} k\left(\frac{t - t_{IO_{spike}}}{\tau_{LTD}}\right) \cdot \delta_{GrC_{spike}}(t) \cdot dt \quad if\ PF_j\ is\ active\ at\ t$$

750　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　　(32)

751
$$LTP\ \Delta w_{PF_j-PC_i}(t) = \alpha \quad const \tag{33}$$

752　where $\Delta w_{PF_j-PC_i}(t)$ denotes the weight change between the j^{th} PF and the target i^{th}

753　PC; τ_{LTD} is the time constant that compensates for the sensory motor delay (i.e., about

754　100 ms, Sargolzaei et al., 2016); δ_{GrC} is the Dirac delta function corresponding to an

755　afferent spike from a PF; and the kernel function *k(x)* is defined as in Eq. 34:

756
$$k(x) = e^{-x} \cdot sin(x)^{20} \tag{34}$$

757　With this parametric configuration, the effect on presynaptic spikes arriving through PFs

758　is maximal over the 100 ms time window before CF spike arrival, thus accounting for the

759　sensorimotor pathway delay. For the sake of computational efficiency, note that the

kernel $k(x)$ combines exponential and trigonometric functions that allow for recursive computation suitable for an event-driven simulation scheme as EDLUT (Naveros et al., 2017, Naveros et al., 2015, Ros et al., 2006). Computational recursion avoids integrating the whole kernel upon each new spike arrival.

Finally, as shown in Eq. 33, the amount of LTP at PFs was fixed (Kawato and Gomi, 1992, Luque et al., 2016, Luque et al., 2011a), with an increase of synaptic efficacy equal to α each time a spike arrived through a PF to the targeted PC. This STDP mechanism correlated the activity patterns coming through the PFs to PCs with the instructive signals coming from CFs to PCs (producing LTD in the activated PF–PC synapses). The correlation process at PC level identified certain PF activity patterns and it consequently reduced the PC output activity. A decrease of PC activations caused a subsequence reduction on the PC inhibitory action over the target MVN. Since the MVN received an almost constant gross MF activation, a lack of PC inhibitory action caused increasing levels of MVN activation. Conversely, the STDP mechanism increased the PC inhibitory activity by potentiating PF-PC synapses in the absence of instructive signal, thus causing decreasing levels of MVN activations. Consequently, PC axon activity governed MVN activation by shaping their inhibitory action produced onto MVN. This spike-timing-dependent plasticity (STDP) mechanism, which regulated the LTP/LTD ration at PF-PC synapses, shaped the inhibitory action of PCs onto MVN cells.

MF-MVN synaptic plasticity. The LTD/LTP dynamics at MF-MVN synapses were the same as in Luque et al. (2019), i.e., they were based on the following rules:

781
$$LTD \; \Delta w_{MF_j - MVN_i}(t) = \int_{-\infty}^{\infty} k\left(\frac{t - t_{PC_{spike}}}{\sigma_{MF-MVN}}\right) \cdot \delta_{MF_{spike}}(t) \cdot dt \quad if \, MF_j \, is \, active \, at \, t$$

782
(35)

783
$$LTP \; \Delta w_{MF_j - MVN_i}(t) = \alpha \quad const \tag{36}$$

784   with $\Delta w_{MF_j - MVN_i}(t)$ denotes the weight change between the j^{th} MF and the target i^{th}

785   MVN; σ_{MF-MVN} the temporal width of the kernel; and δ_{MF} the Dirac delta function that

786   defined a MF spike. The integrative kernel function $k(x)$ was taken as:

787
$$k(x) = e^{-|x|} \cdot cos(x)^2 \tag{37}$$

Note that there is no need for sensorimotor delay compensation thanks to the previous

learning rule (τ_{LTD} in Eq. 32). This second STDP mechanism accounted for learning

consolidation at MVN (see Luque et al., 2016). The PC output operated as an

instructive signal and correlated the activity patterns coming from MFs to MVN

(producing LTD in the activated MF–MVN synapses upon the arrival of the instructive

signal and LTP otherwise). Well-timed sequences of increasing/decreasing levels of

MVN activation ultimately shaped the cerebellar output during VOR adaptation.

The EDLUT source code is available at the following URL:

www.ugr.es/~nluque/restringido/CODE_Cerebellar_Ageing_Vestibulo_Ocular_Adaptati

on.rar

User: REVIEWER, password: REVIEWER.

5.   Figure Captions

Figure 1| Cerebellum-dependent adaptation of Vestibulo-Ocular Reflex (VOR). (A)
VOR stabilises the visual field during head motion, in this example during horizontal head
rotations x(t), by producing contralateral eye movements y(t). (B) Cerebellar VOR
adaptation was theoretically modelled in terms of a classic feedforward control loop. The
control system compared a known reference, or input variable x(t) to the actual output
y(t), to quantify an error signal ε(t) driving adaptation. Cerebellar learning compensated
for the difference between eye and head velocity (i.e., it minimised the error function ε(t)).
The cerebellar model presented here had to learn contralateral eye control in the
presence of a 1-Hz sinusoidal head velocity function (iteratively presented as in classical
horizontal rotational VOR (i.e., r-VOR) (Leigh and Zee, 2015). (C) Schematic
representation of the main cerebellar layers, cells, and synaptic connections considered
in the model. Mossy fibres (MFs) conveyed vestibular information onto granular cells
(GCs) and medial vestibular nuclei (MVN). GCs, in turn, projected onto Purkinje cells
(PCs) through parallel fibres (PFs). PCs also received excitatory inputs from the inferior
olivary (IO) system. IO cells were electrically coupled and delivered an error signal
through the climbing fibres (CFs). Finally, MVN were inhibited by PCs and provide the
cerebellar output driving oculomotor neurons. Two spike-dependent plasticity
mechanisms operated at PF - PC and MF – MVN synapses.

Figure 2| Time course of VOR gain and error during adaptation. (A) Evolution of
VOR gain averaged across 40 individuals with stochastically initialised weights at PF-PC
and MF-MVN synapses, under a sinusoidal vestibular stimulus of 1 Hz. The gain
plateaued at 0.95 in agreement with human records (Dits et al., 2013). (B) Red curve:
convergence of the mean absolute VOR error during adaptation averaged over 40

individuals. Green squares: error-related mean frequency of IO neurons throughout VOR

adaptation (diminishing from 8-9 Hz to 2-3 Hz).

Figure 3| Impact of age-related vestibular loss on IO electrical coupling and

cerebellar-dependent VOR adaptation. (A) *Mean frequency, averaged over 20 young*

(20 yo, top) and 20 older (100 yo, bottom) individuals, across a cluster of 5 × 5 IO cells

(lattice configuration Nobukawa and Nishimura, 2016) during the first second of VOR

adaptation under a sinusoidal vestibular stimulus of 1 Hz. Ageing engendered an on/off

like transient pattern of IO network activity, leading to a higher average activation

*frequency in older IO cells. **(B)** Discrete Wavelet Transformation (DWT) applied to the*

snapshot sequence of the IO membrane potentials (Supp. Fig. 1A) obtained during the

first second of VOR adaptation for a young (20 yo) and an older (100 yo) adult. In the

presence of equivalent input stimulation to the IO network, the mean DWT coefficients

were significantly higher in young adults (ANOVA $F_{(294,16)}=18$, $p < 10^{-7}$), indicating a more

complex evolution of the spatial-temporal patterns, higher variability in membrane

*potentials, and lower overall frequencies in IO network activity. **(C)** Increase of the mean*

absolute VOR error (red curve) throughout ageing and corresponding increase of the

average IO frequency (green squares), caused by altered electrical coupling (simulated

*from 40 to 100 years of age). **(D)** Age-related impact on VOR velocity compensatory eye*

*velocity functions (with young and older IO network, red and black, respectively). **(E)***

Cross-sectional ageing simulation over a study population of 2400 individuals aged from

40 to 100 years (40 individuals per each of year of age). Only the effect of age-related

vestibular loss and IO coupling alteration was considered in this ageing simulation, with

no compensatory mechanism operating in the downstream cerebellar network. Each

individual underwent an independent 1 Hz r-VOR protocol (during 2500 s). On average,

the VOR performance declined quasi-linearly throughout ageing, with an increasing

interindividual variability.

Figure 4| Intrinsic plasticity at PC synapses as a local homeostatic mechanism.

(A) Trimodal spiking patterns of model PCs: tonic firing, corresponding to simple spikes

elicited by PF inputs; bursting mode, during which complex spikes (bursts of spikes) are

elicited by CFs (~500 synapses Palay and Chan-Palay, 2012) wrapping around PC

dendrites that can even suppress simple spiking; and silent mode, corresponding to an

extended hyperpolarisation period called the post-complex spike pause. (B) Intrinsic

plasticity modified PC excitability through capacitance adaptation to age-dependent

changes in the synaptic inputs (see Methods). The model PC capacitance decreased

with ageing to facilitate neural excitability (Zhang et al., 2010) (C) As a consequence,

PCs' firing rates increased significantly in older simulated individuals as compared to

young ones (Zhang et al., 2010). This measure was realised with PCs operating in

spiking tonic mode (10-250Hz). (D) The linear correlation between pause duration and

pre-complex spike ISI duration (Grasselli et al., 2016) in older PCs was preserved ($R^2 =$

0.9932; $p < 10^{-4}$). (E) Cross-sectional ageing simulation accounting for vestibular loss,

IO electrical coupling chances, and intrinsic plasticity at PC synapses. The VOR gain

decayed linearly across the lifespan as in the previous ageing simulation. However, the

adaptive excitability of PCs (induced by intrinsic plasticity) partially counterbalanced the

vestibular degeneration (as compared to Fig. 3E).

Figure 5| Spike-based plasticity mediating LTP/LTD at PF-PC and MF-MVN

***synapses. (A, B)** Synaptic weight distributions obtained at PF-PC connections by*

averaging over 20 young individuals (20 yo) and 20 older ones (100 yo), respectively.

Each individual underwent an independent 1 Hz r-VOR adaptation (during 2500 s). The

two cerebellar microcomplexes devoted to the control of rightward and leftward eye

*movements are visible. **(C, D)** Synaptic weight distributions at MF-MVN connections by*

averaging over 20 young individuals (20 yo) and 20 older ones (100 yo), respectively.

The antisymmetric distributions with respect to (A, B) are caused by the inhibitory PC

projections onto MVN. To counter age-related afferent loss, LTP/LTD increased the

weight of remaining synapses both at PF-PC and MF-MVN projections, significantly

increasing the ratio between the number of saturated vs. active afferents. As a

consequence, the neural drive increased significantly as a function of age.

Figure 6| Cerebellar LTP/LTD as a global homeostatic compensatory mechanism.

***(A)** LTP/LTD sustained the input drive to PCs (computed by convolving unitary pulses*

and synaptic weight distributions) until about 80 years of age and it dropped afterwards.

***(B)** A rapid boost of the neural drive at MVN synapses around 60 years of age helped*

maintaining the overall responses. Beyond 80 years of age, the neural drive could no

*longer be maintained against age-related vestibular losses. **(C)** Cross-sectional ageing*

simulation accounting for vestibular loss, IO electrical coupling alterations and LTP/LTD

at PF-PC and MF-MVN synapses. The VOR performances remained quasi stable until

about 80 years and it began to decline afterwards, with an incrementally increasing

interindividual variability.

Figure 7| Impact of ageing on VOR: cross-sectional and longitudinal studies. (A)

Cross-sectional ageing simulation accounting for all factors (vestibular loss, IO electrical

coupling changes, PC intrinsic plasticity, and LTP/LTD at PF-PC and MF-MVN

synapses) to evaluate their synergistic contribution to r-VOR adaptation. Again, a study

population of 2400 individuals (age range: 40-100 years; 40 individuals per each year of

age) underwent the same r-VOR protocol (1 Hz head rotation during 2500 s). The VOR

*gain remained quasi-stable until 80-85 yo, and it declined sharply afterwards. **(B)** The*

local homeostatic mechanism (PC intrinsic plasticity) helped sustaining the VOR gain

*further in age, complementing the global homeostatic action of LTP/LTD. **(C)** The*

interindividual variability increased smoothly until 80-85 yo and it augmented rapidly

*afterwards, consistently with the tendency observed experimentally (Li et al., 2015). **(D)***

Longitudinal ageing simulation considering all age-related and compensatory factors. A

study population of 40 individuals were followed-up individually during 60 years (from 40

to 100 years of age) using the same r-VOR protocol (1 Hz head rotation during 2500 s).

Age-related losses accumulated longitudinally for each individual. VOR performance

remained quasi-stable until 80-85 yo and it declined sharply afterwards. Interindividual

variability increased significantly during the last 15 years of age becoming larger than

the one predicted by to the cross-sectional simulation (C). The thick green and red curves

correspond to the 100 yo individuals with best and worst VOR performance, respectively.

Figure 8| Distribution of remaining fibres/synaptic connections in the longitudinal

***ageing simulation. (A)** VOR gain at 100 years of age sorted incrementally across the*

study population. The red and green dots correspond to the worst and best individuals,

*respectively (i.e., red and green thick curves in Fig. 7D, respectively). **(B)** Eye velocity*

function, with a focus on time windows corresponding to the peak and the trough of the

*sinusoidal profile (i.e., between 200-300 ms and 700-800 ms). **(C)** Residual PFs (top)*

and MF-MVN projections (bottom) active at the peak and the trough of the eye velocity

profile across all individuals sorted on the basis of their VOR performance (left: 60 yo;

centre: at 85 yo; right: 100 yo). The larger the number of residual PFs and MV-MVN

projections encoding the peak and the trough of the eye velocity function, the higher the

VOR performance throughout ageing (see correlations in Supp. Fig. S2).

Supplementary S1| Spatial-temporal evolution of the IO network activity patterns

***in young and older individuals. (A)** Spike propagation through electrical coupling along*

the diagonal of an IO cell cluster (5x5 lattice configuration) induced by an external

*stimulus delivered at the centre of the network. **(B)** Time course of the membrane*

potentials of IO cells in 5x5 clusters caused by a large retinal slip (input stimulus received

*at the centre of the network at 1 ms) in young (top) and older (bottom) individuals. **(C)***

Ageing favours a faster and higher electrical coupling in old adults.

Supplementary S2| Correlation matrix between the VOR gain and the number of

residual PFs and MF-MVN projections active at the peak and the trough of the eye

velocity function. *The diagonal represents the histograms of the VOR performance*

values.

6.   Tables

Table 1. *Cerebellar network topology parameters*

Neurons		Synapses			
Pre-synaptic cells (number)	*Post-synaptic cells (number)*	*Number*	*Type*	*Initial weight*	*Weight range*
2000 GCs	*200 PCs*	*400000*	*AMPA*	*rand*	*[0, 3.65]*
200 IO	*200 PCs*	*200*	*AMPA*	*40*	*–*
100 MFs	*200 MVN*	*20000*	*AMPA*	*0*	*[0, 1]*
200 PCs	*200 MVN*	*200*	*GABA*	*1.5*	*–*
200 IO	*200 MVN*	*200*	*NMDA*	*7*	*–*
IO to IO (lattice configuration)		*320*	*EC*	*5*	*–*

933   **Table 2.** *Neuronal model parameters.*

Parameters	*MVN*	*IO*	*PC*
C_m *(pF)*	*2*	*10*	*7.16*
G_L *(nS)*	*0.2*	*0.15*	*0.15*
E_L *(mV)*	*-70*	*-70*	*-70*
E_{AMPA} *(mV)*	*0*	*0*	*0*
E_{GABA} *(mV)*	*-80*	*-80*	*-80*
τ_{AMPA} *(ms)*	*0.5*	*1*	*1*
τ_{NMDA} *(ms)*	*14*		
τ_{GABA} *(ms)*	*10*	*2*	*2*
V_{thr} *(mV)*	*-40*	*-50*	*-35*
T_{ref} *(ms)*	*1*	*1.35*	*1.35*

		31	31
V_{peak} (mV)		31	31
gc_a (mS)			0.0075
g_M (mS)			5.65

935 **Table 3.** *Ionic conductance kinetic parameters.*

Conductance type	Steady–state Activation/Inactivation	Time constant (ms)
g_K delayed rectifier potassium current	$x_0[V] = \dfrac{1}{1 + e^{\frac{-V-29.5}{10}}}$	$\tau_x[V] = 0.25 + 4.35 \cdot e^{\frac{-\lvert V+10 \rvert}{10}}$
g_{Na} transient inactivating sodium current	$x_0[V] = \dfrac{1}{1 + e^{\frac{V-59.4}{10.7}}}$	$\tau_x[V] = 0.15 + \dfrac{1.15}{1 + e^{\frac{V+33.5}{15}}}$
$m_0[V]$	$m_0[V] = \dfrac{1}{1 + e^{\frac{-V-48}{10}}} \cdot m$	
	Forward Rate Function (α)	**Backward Rate Function** (β)
g_{Ca} high threshold	$\alpha = \dfrac{1.6}{1 + e^{-0.0072 \cdot (V-5)}}$	$\beta = \dfrac{0.02 \cdot (V + 8.9)}{e^{\frac{V+8.9}{5}}}$
g_M muscarinic receptor suppressed potassium current	$\alpha = \dfrac{0.3}{1 + e^{\frac{-V-2}{5}}}$	$\beta = 0.001 \cdot e^{\frac{-V-70}{18}}$
	Steady–state Activation/Inactivation	**Time constant(ms)**
	$x_0[V] = \dfrac{\alpha}{\alpha + \beta}$	$\tau_x[V] = \dfrac{1}{\alpha + \beta}$

7. References

938 AGRAWAL, Y., DAVALOS-BICHARA, M., ZUNIGA, M. G. & CAREY, J. P. 2013. Head
939 impulse test abnormalities and influence on gait speed and falls in older
940 individuals. *Otology & Neurotology,* 34**,** 1729-1735.

ALVAREZ, J. C., DíAZ, C., SUAREZ, C., FERNANDEZ, J. A., GONZALEZ DEL REY, C., NAVARRO, A. & TOLIVIA, J. 2000. Aging and the human vestibular nuclei: morphometric analysis. *Mechanisms of ageing and development,* 114**,** 149-72.

ALLEN, D., RIBEIRO, L., ARSHAD, Q. & SEEMUNGAL, B. M. 2017. Age-related vestibular loss: current understanding and future research directions. *Frontiers in neurology,* 7**,** 231.

ANDERSEN, B. B., GUNDERSEN, H. J. G. & PAKKENBERG, B. 2003. Aging of the human cerebellum: a stereological study. *Journal of Comparative Neurology,* 466**,** 356-365.

ANSON, E. & JEKA, J. 2016. Perspectives on aging vestibular function. *Frontiers in neurology,* 6**,** 269.

ANSON, E. R., BIGELOW, R. T., CAREY, J. P., XUE, Q.-L., STUDENSKI, S., SCHUBERT, M. C. & AGRAWAL, Y. 2016. VOR gain is related to compensatory saccades in healthy older adults. *Frontiers in aging neuroscience,* 8**,** 150.

ARENZ, A., SILVER, R. A., SCHAEFER, A. T. & MARGRIE, T. W. 2008. The Contribution of Single Synapses to Sensory Representation in Vivo. *Science* 321**,** 977-80.

ARLEO, A. & RONDI-REIG, L. 2007. Multimodal sensory integration and concurrent navigation strategies for spatial cognition in real and artificial organisms. *Journal of integrative neuroscience,* 6**,** 327-366.

BADURA, A., CLOPATH, C., SCHONEWILLE, M. & DE ZEEUW, C. I. 2016. Modeled changes of cerebellar activity in mutant mice are predictive of their learning impairments. *Sci. Rep.,* 6**,** 36131.

BALOH, R. W., ENRIETTO, J., JACOBSON, K. M. & LIN, A. 2001. Age-related changes in vestibular function: a longitudinal study. *Annals of the New York Academy of Sciences,* 942**,** 210-219.

BALOH, R. W., JACOBSON, K. M. & SOCOTCH, T. M. 1993. The effect of aging on visual-vestibuloocular responses. *Exp. Brain Res.,* 95**,** 509-16.

BALOH, R. W., SLOANE, P. D. & HONRUBIA, V. 1989. Quantitative vestibular function testing in elderly patients with dizziness. *Ear, nose, & throat journal,* 68**,** 935-9.

BERGSTRÖM, B. 1973. Morphology of the vestibular nerve: III. Analysis of the calibers of the myelinated vestibular nerve fibers in man at various ages. *Acta oto-laryngologica,* 76**,** 331-8.

BEST, A. R. & REGEHR, W. G. 2009. Inhibitory regulation of electrically coupled neurons in the inferior olive is mediated by asynchronous release of GABA. *Neuron,* 62**,** 555-565.

BEZZI, M., NIEUS, T., COENEN, O. J.-M. D. & D'ANGELO, E. 2004. An I&F model of a cerebellar granule cell. *Neurocomp.,* 58**,** 593-8.

BOUCHENY, C., CARRILLO, R. R., ROS, E. & COENEN, O. J.-M. D. 2005. Real-time spiking neural network: an adaptive cerebellar model. *LNCS,* 3512**,** 136–44.

BRANDT, T., SCHAUTZER, F., HAMILTON, D. A., BRÜNING, R., MARKOWITSCH, H. J., KALLA, R., DARLINGTON, C., SMITH, P. & STRUPP, M. 2005. Vestibular loss causes hippocampal atrophy and impaired spatial memory in humans. *Brain,* 128**,** 2732-2741.

BRIZZEE, K. R., KAACK, B. & KLARA, P. 1975. Lipofuscin: intra-and extraneuronal accumulation and regional distribution. *Neurobiology of Aging.* Springer.

CARRILLO, R. R., E. ROS, E., BOUCHENY, C. & COENEN, O. 2008. A real-time spiking cerebellum model for learning robot control. *Biosystems,* 94**,** 18-27.

CLOPATH, C., BADURA, A., DE ZEEUW, C. I. & BRUNEL, N. 2014. A cerebellar learning model of VOR adaptation in wild-type and mutant mice. *J. Neurosci.,* 34**,** 7203-15.

CULLEN, K. E. 2012. The vestibular system: multimodal integration and encoding of self-
motion for motor control. *Trends in neurosciences,* 35**,** 185-196.
D'ANGELO, E. & DE ZEEUW, C. I. 2009. Timing and plasticity in the cerebellum: focus
on the granular layer. *Trends Neurosci.,* 32**,** 10.
D'ANGELO, E., MAPELLI, L., CASELLATO, C., GARRIDO, J. A., LUQUE, N.,
MONACO, J., PRESTORI, F., PEDROCCHI, A. & ROS, E. 2016. Distributed
circuit plasticity: new clues for the cerebellar mechanisms of learning. *The*
Cerebellum, 15**,** 139-151.
DAVIE, J. T., CLARK, B. A. & HÄUSSER, M. 2008. The origin of the complex spike in
cerebellar Purkinje cells. *J. Neurosci.,* 28**,** 7599-7609.
DE ZEEUW, C. I., HOOGENRAAD, C. C., KOEKKOEK, S. K. E., RUIGROK, T. J.,
GALJART, N. & SIMPSON, J. I. 1998. Microcircuitry and function of the inferior
olive. *Trends Neurosci.,* 21**,** 391-400.
DEMER, J. L., HONRUBIA, V. & BALOH, R. W. 1994. Dynamic visual acuity: a test for
oscillopsia and vestibulo-ocular reflex function. *The American journal of otology,*
15**,** 340-347.
DEMER, J. L., OAS, J. G. & BALOH, R. W. 1993. Visual-vestibular interaction in humans
during active and passive, vertical head movement. *J. Vestib. Res.,* 3**,** 101-14.
DENG, Y., XU, L., ZENG, X., LI, Z., QIN, B. & HE, N. 2010. New perspective of GABA
as an inhibitor of formation of advanced lipoxidation end-products: it's interaction
with malondiadehyde. *J Biomed Nanotechnol,* 6**,** 318-24.
DESAI, A., GOODMAN, V., KAPADIA, N., SHAY, B. L. & SZTURM, T. 2010.
Relationship between dynamic balance measures and functional performance in
community-dwelling elderly people. *Physical therapy,* 90**,** 748-760.
DEVOR, A. & YAROM, Y. 2002. Generation and propagation of subthreshold waves in
a network of inferior olivary neurons. *Journal of neurophysiology,* 87**,** 3059-3069.
DITS, J., HOUBEN, M. M. & VAN DER STEEN, J. 2013. Three dimensional Vestibular
ocular reflex testing using a Six degrees of freedom motion platform. *JoVE*
*(Journal of Visualized Experiments)***,** e4144.
DUMAS, G., PERRIN, P., OUEDRAOGO, E. & SCHMERBER, S. 2016. How to perform
the skull vibration-induced nystagmus test (SVINT). *European annals of*
otorhinolaryngology, head and neck diseases, 133**,** 343-348.
FONSECA, D. B., SHEEHY, M. R. J., BLACKMAN, N., SHELTON, P. M. J. & PRIOR, A.
E. 2005. Reversal of a hallmark of brain ageing: lipofuscin accumulation.
Neurobiol. Aging, 26**,** 69-76.
FORREST, M. 2008. *Biophysics of Purkinje computation.* University of Warwick.
FUJITA, M. 1982. Adaptive filter model of the cerebellum. *Biological cybernetics,* 45**,**
195-206.
FURMAN, J. M. & REDFERN, M. S. 2001. Effect of aging on the otolith-ocular reflex.
Journal of Vestibular Research, 11**,** 91-103.
GAO, Z., VANBEUGEN, B. J. & DE ZEEUW, C. I. 2012. Distributed Synergistic Plasticity
and Cerebellar Learning. *Nat. Rev. Neurosci.,* 13**,** 1-17.
GARRIDO, J. A., LUQUE, N. R., TOLU, S. & D'ANGELO, E. 2016. Oscillation-driven
spike-timing dependent plasticity allows multiple overlapping pattern recognition
in inhibitory interneuron networks. *Int. J. Neural Systems,* 26**,** 1650020.
GERSTNER, W. & KISTLER, W. M. 2002. *Spiking neuron models: Single neurons,*
populations, plasticity, Cambridge university press.
GERSTNER, W., KISTLER, W. M., NAUD, R. & PANINSKI, L. 2014. *Neuronal dynamics:*
From single neurons to networks and models of cognition, Cambridge University
Press.

GORDON, J. L., FURMAN, J. M. R. & KAMEN, E. W. System identification of the
vestibulo-ocular reflex: application of the recursive least-squares algorithm.
Bioengineering Conference, 1989., Proceedings of the 1989 Fifteenth Annual
Northeast, 1989. IEEE, 199-200.

GRASSELLI, G., HE, Q., WAN, V., ADELMAN, J. P., OHTSUKI, G. & HANSEL, C. 2016.
Activity-Dependent Plasticity of Spike Pauses in Cerebellar Purkinje Cells. *Cell
Reports,* 14**,** 2546-53.

GROSSMAN, G. E. & LEIGH, R. J. 1990. Instability of gaze during locomotion in patients
with deficient vestibular function. *Annals of Neurology: Official Journal of the
American Neurological Association and the Child Neurology Society,* 27**,** 528-
532.

HIRVONEN, T., AALTO, H., PYYKKÖ, I., JUHOLA, M. & JÄNTTIL, P. 1997. Changes in
vestibulo-ocular reflex of elderly people. *Acta Oto-Laryngologica,* 117**,** 108-110.

HORAK, F. B. 2006. Postural orientation and equilibrium: what do we need to know about
neural control of balance to prevent falls? *Age and ageing,* 35**,** ii7-ii11.

ICHIKAWA, R., SAKIMURA, K. & WATANABE, M. 2016. GluD2 endows parallel fiber–
Purkinje cell synapses with a high regenerative capacity. *Journal of
neuroscience,* 36**,** 4846-4858.

ISHIKAWA, T., SHIMUTA, M. & HÄUSSER, M. 2015. Multimodal sensory integration in
single cerebellar granule cells in vivo. *Elife,* 4**,** e12916.

ITO, M. 2013. Error Detection and Representation in the Olivo-Cerebellar System. *Front.
Neural Circuits***,** 1-8.

JAHN, K., NAEßL, A., SCHNEIDER, E., STRUPP, M., BRANDT, T. & DIETERICH, M.
2003. Inverse U-shaped curve for age dependency of torsional eye movement
responses to galvanic vestibular stimulation. *Brain,* 126**,** 1579-1589.

JANG, D. C., SHIM, H. G. & KIM, S. J. 2020. Intrinsic plasticity of cerebellar purkinje
cells contributes to motor memory consolidation. *Journal of Neuroscience,* 40**,**
4145-4157.

KAWATO, M. & GOMI, H. 1992. A computational model of four regions of the cerebellum
based on FEL. *Biol. Cybern.,* 68**,** 95-103.

KURODA, S., YAMAMOTO, K., MIYAMOTO, H., DOYA, K. & KAWATO, M. 2001.
Statistical characteristics of climbing fiber spikes necessary for efficient
cerebellar learning. *Biol. Cybern.,* 84**,** 183-92.

LACKNER, J. R. & DIZIO, P. 2005. Vestibular, proprioceptive, and haptic contributions
to spatial orientation. *Annu. Rev. Psychol.,* 56**,** 115-147.

LASN, H., WINBLAD, B. & BOGDANOVIC, N. 2001. The number of neurons in the
inferior olivary nucleus in Alzheimer's disease and normal aging: A stereological
study using the optical fractionator. *J. Alzheimer's Disease,* 3**,** 159-68.

LATORRE, R., AGUIRRE, C., RABINOVICH, M. & VARONA, P. 2013. Transient
dynamics and rhythm coordination of inferior olive spatio-temporal patterns.
Frontiers in Neural Circuits, 7.

LEFLER, Y., AMSALEM, O., VRIELER, N., SEGEV, I. & YAROM, Y. 2020. Using
subthreshold events to characterize the functional architecture of the electrically
coupled inferior olive network. *Elife,* 9**,** e43560.

LEFLER, Y., YAROM, Y. & UUSISAARI, M. Y. 2014. Cerebellar inhibitory input to the
inferior olive decreases electrical coupling and blocks subthreshold oscillations.
Neuron, 81**,** 1389-400.

LEIGH, R. J. & ZEE, D. S. 2015. *The neurology of eye movements*, Oxford University
Press.

LI, C., LAYMAN, A. J., GEARY, R., ANSON, E., CAREY, J. P., FERRUCCI, L. & AGRAWAL, Y. 2015. Epidemiology of vestibulo-ocular reflex function: data from the Baltimore Longitudinal Study of Aging. *Otol Neurotol,* 36**,** 267-72.

LI, C. & LI, Y. 2013. A spike-based model of neuronal intrinsic plasticity. *IEEE Trans. Autonomous Mental Develop.,* 5**,** 62-73.

LISBERGER, S. G. & FUCHS, A. F. 1978. Role of primate flocculus during rapid behavioral modification of VOR. II. Mossy fiber firing patterns during horizontal head rotation and eye movement. *J. Neurophysiol.,* 41**,** 764-77.

LOEWENSTEIN, Y. 2002. A possible role of olivary gap-junctions in the generation of physiological and pathological tremors. *Mol. Psychiatry,* 7**,** 129.

LOPEZ, I., HONRUBIA, V. & BALOH, R. W. 1996. Aging and the human vestibular nucleus. *J. Vestibular research: equilibrium & orientation,* 7**,** 77-85.

LORENTE DE NÓ, R. 1933. Vestibulo-ocular reflex arc. *Archiv Neurol & Psychiatry.*

LUQUE, N. R., GARRIDO, J. A., CARRILLO, R. R., COENEN, O. J. M. D. & ROS, E. 2011a. Cerebellar Input Configuration Toward Object Model Abstraction in Manipulation Tasks. *IEEE Trans. Neural. Netw.,* 22**,** 1321-8.

LUQUE, N. R., GARRIDO, J. A., CARRILLO, R. R., COENEN, O. J. M. D. & ROS, E. 2011b. Cerebellarlike Corrective Model Inference Engine for Manipulation Tasks. *IEEE Trans. Syst. Man. Cybern.,* 41**,** 1299-312.

LUQUE, N. R., GARRIDO, J. A., CARRILLO, R. R., D'ANGELO, E. & ROS, E. 2014. Fast convergence of learning requires plasticity between inferior olive and deep cerebellar nuclei in a manipulation task: a closed-loop robotic simulation. *Front. Comput. Neurosci.,* 8.

LUQUE, N. R., GARRIDO, J. A., NAVEROS, F., CARRILLO, R. R., D'ANGELO, E. & ROS, E. 2016. Distributed Cerebellar Motor Learning; a STDP Model. *Front. Comp. Neurosci.,* 10.

LUQUE, N. R., NAVEROS, F., CARRILLO, R. R., ROS, E. & ARLEO, A. 2019. Spike burst-pause dynamics of Purkinje cells regulate sensorimotor adaptation. *PLOS Computational Biology,* 15**,** e1006298.

LLINAS, R., BAKER, R. & SOTELO, C. 1974. Electrotonic coupling between neurons in cat inferior olive. *J. Neurophysiol.,* 37**,** 560-71.

MATHY, A., HO, S. S., DAVIE, J. T., DUGUID, I. C., CLARK, B. A. & HAUSSER, M. 2009. Encoding of oscillations by axonal bursts in inferior olive neurons. *Neuron,* 62**,** 388-99.

MATIÑO-SOLER, E., ESTELLER-MORE, E., MARTIN-SANCHEZ, J.-C., MARTINEZ-SANCHEZ, J.-M. & PEREZ-FERNANDEZ, N. 2015. Normative data on angular vestibulo-ocular responses in the yaw axis measured using the video head impulse test. *Otology & Neurotology,* 36**,** 466-471.

MCGARVIE, L. A., MACDOUGALL, H. G., HALMAGYI, G. M., BURGESS, A. M., WEBER, K. P. & CURTHOYS, I. S. 2015. The video head impulse test (vHIT) of semicircular canal function–age-dependent normative values of VOR gain in healthy subjects. *Frontiers in neurology,* 6**,** 154.

MERGNER, T. & ROSEMEIER, T. 1998. Interaction of vestibular, somatosensory and visual signals for postural control and motion perception under terrestrial and microgravity conditions—a conceptual model. *Brain research reviews,* 28**,** 118-135.

MIDDLETON, S. J., RACCA, C., CUNNINGHAM, M. O., TRAUB, R. D., MONYER, H., KNOPFEL, T., SCHOFIELD, I. S., JENKINS, A. & WHITTINGTON, M. A. 2008. High-frequency network oscillations in cerebellar cortex. *Neuron,* 58**,** 763-74.

MIYASHO, T., TAKAGI, H., SUZUKI, H., WATANABE, S., INOUE, M., KUDO, Y. & MIYAKAWA, H. 2001. Low-threshold potassium channels and a low-threshold

calcium channel regulate Ca2+ spike firing in the dendrites of cerebellar Purkinje
neurons: a modeling study. *Brain Res.,* 891**,** 106-15.
NAJAC, M. & RAMAN, I. M. 2015. Integration of Purkinje Cell Inhibition by Cerebellar
Nucleo-Olivary Neurons. *J. Neurosci.,* 35**,** 544-9.
NAJAFI, F. & MEDINA, J. F. 2013. Beyond "all-or-nothing" climbing fibers: graded
representation of teaching signals in Purkinje cells. *Front. Neural Circuits,* 7**,** 1-
15.
NAVEROS, F., GARRIDO, J. A., CARRILLO, R. R., ROS, E. & LUQUE, N. R. 2017.
Event-and Time-Driven Techniques Using Parallel CPU-GPU Co-processing for
Spiking Neural Networks. *Front. Neuroinformatics,* 11.
NAVEROS, F., LUQUE, N. R., GARRIDO, J. A., CARRILLO, R. R., ANGUITA, M. &
ROS, E. 2015. A Spiking Neural Simulator Integrating Event-Driven and Time-
Driven Computation Schemes Using Parallel CPU-GPU Co-Processing: A Case
Study. *IEEE Trans. Neural Netw. Learn. Syst.,* 26**,** 1567-74.
NAVEROS, F., LUQUE, N. R., ROS, E. & ARLEO, A. 2019. VOR Adaptation on a
Humanoid iCub Robot Using a Spiking Cerebellar Model. *IEEE transactions on*
cybernetics.
NGUYEN-VU, T. B., ZHAO, G. Q., LAHIRI, S., KIMPO, R. R., LEE, H., GANGULI, S.,
SHATZ, C. J. & RAYMOND, J. L. 2017. A saturation hypothesis to explain both
enhanced and impaired learning with enhanced plasticity. *elife,* 6**,** e20147.
NOBUKAWA, S. & NISHIMURA, H. 2016. Chaotic resonance in coupled inferior olive
neurons with the Llinás approach neuron model. *Neural computation,* 28**,** 2505-
2532.
PAIGE, G. 1992. Senescence of human visual-vestibular interactions. 1. Vestibulo-
ocular reflex and adaptive plasticity with aging. *Journal of vestibular research:*
equilibrium & orientation, 2**,** 133-151.
PAIGE, G. D. 1994. Senescence of human visual-vestibular interactions: smooth pursuit,
optokinetic, and vestibular control of eye movements with aging. *Experimental*
brain research, 98**,** 355-372.
PALAY, S. L. & CHAN-PALAY, V. 2012. *Cerebellar cortex: cytology and organization*,
Springer Science & Business Media.
PERNICE, H. F., SCHIEWECK, R., JAFARI, M., STRAUB, T., BILBAN, M., KIEBLER,
M. A. & POPPER, B. 2019. Altered Glutamate Receptor Ionotropic Delta Subunit
2 Expression in Stau2-Deficient Cerebellar Purkinje Cells in the Adult Brain.
International journal of molecular sciences, 20**,** 1797.
PETERKA, R. J., BLACK, F. O. & SCHOENHOFF, M. B. 1990. Age-related changes in
human vestibulo-ocular reflexes: sinusoidal rotation and caloric tests. *J Vestib*
Res, 1**,** 49-59.
PIIRTOLA, M. & ERA, P. 2006. Force platform measurements as predictors of falls
among older people–a review. *Gerontology,* 52**,** 1-16.
RENOVELL, A., GINER, J. & PORTOLES, M. 2001. Loss of granule neurons in the aging
human cerebellar cortex. *International Journal of Developmental Biology,* 40**,**
S193-S194.
ROBINSON, D. A. 1981. The use of control systems analysis in the neurophysiology of
eye movements. *Annual review of neuroscience,* 4**,** 463-503.
ROS, E., CARRILLO, R. R., ORTIGOSA, E. M., BARBOUR, B. & AGÍS, R. 2006. Event-
driven simulation scheme for spiking neural networks using lookup tables to
characterize neuronal dynamics. *Neural Comput.,* 18**,** 2959-93.
ROTH, A. & HÄUSSER, M. 2001. Compartmental models of rat cerebellar Purkinje cells
based on simultaneous somatic and dendritic patch-clamp recordings. *J.*
Physiol., 535**,** 445-72.

SANTINA, C. C., CREMER, P. D., CAREY, J. P. & MINOR, L. B. 2001. The Vestibulo-
Ocular Reflex during Self-Generated Head Movements by Human Subjects with
Unilateral Vestibular Hypofunction. *Annals New York Academy of Sciences,* 942**,**
465-6.
SARGOLZAEI, A., ABDELGHANI, M., YEN, K. K. & SARGOLZAEI, S. 2016.
Sensorimotor control: computing the immediate future from the delayed present.
BMC bioinformatics, 17**,** 245.
SCHWEIGHOFER, N., DOYA, K. & KAWATO, M. 1999. Electrophysiological properties
of inferior olive neurons: a compartmental model. *Journal of neurophysiology,* 82**,**
804-817.
SCHWEIGHOFER, N., LANG, E. J. & KAWATO, M. 2013. Role of the olivo-cerebellar
complex in motor learning and control. *Frontiers in neural circuits,* 7**,** 94.
SHIM, H. G., JANG, D. C., LEE, J., CHUNG, G., LEE, S., KIM, Y. G., JEON, D. E. & KIM,
S. J. 2017. Long-term depression of intrinsic excitability accompanied by the
synaptic depression in the cerebellar Purkinje cells. *J. Neurosci.***,** 3464-16.
SHIM, H. G., LEE, Y.-S. & KIM, S. J. 2018. The emerging concept of intrinsic plasticity:
activity-dependent modulation of intrinsic excitability in cerebellar Purkinje cells
and motor learning. *Experimental neurobiology,* 27**,** 139.
SKAVENSKI, A. A. & ROBINSON, D. A. 1973. Role of abducens neurons in
vestibuloocular reflex. *Journal of Neurophysiology,* 36**,** 724-738.
SOTELO, C., LLINAS, R. & BAKER, R. 1974. Structural study of inferior olivary nucleus
of the cat: morphological correlates of electrotonic coupling. *J. Neurophysiol.,* 37**,**
541-59.
SULZER, D., MOSHAROV, E., TALLOCZY, Z., ZUCCA, F. A., SIMON, J. D. & ZECCA,
L. 2008. Neuronal pigmented autophagic vacuoles: lipofuscin, neuromelanin, and
ceroid as macroautophagic responses during aging and disease. *J. Neurochem.,*
106**,** 24-36.
TINETTI, M. E. 2003. Preventing falls in elderly persons. *New England journal of*
medicine, 348**,** 42-49.
TOKUDA, I. T., HOANG, H., SCHWEIGHOFER, N. & KAWATO, M. 2013. Adaptive
coupling of inferior olive neurons in cerebellar learning. *Neural Networks,* 47**,** 42-
50.
TORVIK, A., TORP, S. & LINDBOE, C. 1986. Atrophy of the cerebellar vermis in ageing:
a morphometric and histologic study. *Journal of the neurological sciences,* 76**,**
283-294.
TURRIGIANO, G., ABBOTT, L. F. & MARDER, E. 1994. Activity-dependent changes in
the intrinsic properties of cultured neurons. *Science,* 264**,** 974-6.
UUSISAARI, M. & DE SCHUTTER, E. 2011. The mysterious microcircuitry of the
cerebellar nuclei. *The Journal of physiology,* 589**,** 3441-3457.
VISWASOM, A. A., SIVAN, S. & JOBBY, A. 2013. Age related changes in the granule
cell number in the human cerebellar cortex. *Journal of Evolution of Medical and*
Dental Sciences, 2**,** 2698-2705.
YAMAZAKI, T. & TANAKA, S. 2005. Neural modeling of an internal clock. *Neural*
Comput, 17**,** 1032-58.
YAMAZAKI, T. & TANAKA, S. 2007. The cerebellum as a liquid state machine. *Neural*
Netw., 20**,** 290-7.
YAMAZAKI, T. & TANAKA, S. 2009. Computational models of timing mechanisms in the
cerebellar granular layer. *Cerebellum,* 8**,** 423-32.
YIN, D. 1996. Biochemical basis of lipofuscin, ceroid, and age pigment-like fluorophores.
Free Radical Biology and Medicine, 21**,** 871-888.

YUZAKI, M. 2013. Cerebellar LTD vs. motor learning—lessons learned from studying
GluD2. *Neural networks,* 47**,** 36-41.
ZALEWSKI, C. K. Aging of the human vestibular system. Seminars in hearing, 2015.
Thieme Medical Publishers, 175-196.
ZANJANI, H. S., VOGEL, M. W. & MARIANI, J. 2016. Deletion of the GluRδ2 receptor in
the hotfoot mouse mutant causes granule cell loss, delayed Purkinje cell death,
and reductions in Purkinje cell dendritic tree area. *The Cerebellum,* 15**,** 755-766.
ZHANG, C., ZHU, Q. & HUA, T. 2010. Aging of cerebellar Purkinje cells. *Cell and tissue*
research, 341**,** 341-7.

1252

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

B

C

VOR gain/year = -0.05%   VOR gain/year = -0.9%

VOR gain

Age (years)

piece-wise linear regression

Figure 6

Figure 7

A

B

Right

Eye velocity (deg/s)

0.2 0.3

0.7 0.8   t(s)

Left

200 ms

Active connections:

during the period 200-300 ms

during the period 700-800 ms

C

Figure 8

Supplementary Figure 1

Correlation Matrix

Figure S2