Sequential onset and concurrent expression of miR-9 genomic loci in single cells contributes to the temporal increase of mature miR-9 in zebrafish neurogenesis

Soto, X.[1], Minchington, T.[1], Lea, R.[1], Lee, J.[2] and Papalopulu, N[1]*

*correspondence: nancy.papalopulu@manchester.ac.uk

[1]Division of Developmental Biology and Medicine, School of Medical Sciences, Faculty of Biology Medicine and Health, The University of Manchester, Oxford Road, Manchester, M13 9PT, UK;

[2]Medicines Discovery Catapult, Block 35, Mereside, Alderley Park, Alderley Edge, Cheshire, SK10 4TG, UK

Abstract

Gene expression oscillations of the Hes/Her family members of transcriptional repressors are important for cell state transitions during neural development. The input of miR-9 is necessary to constrain gene expression noise, allowing oscillations to occur and to be decoded by downstream genes. Hes/Her dynamics are sensitive to the amount of miR-9 present in the cell but the mechanism by which miR-9 is quantitatively controlled is not known. In vertebrates, there are several distinct genomic loci that produce the same mature miR-9, leading to a number of possibilities of how the production of mature miR-9 may be regulated.

Here, we show that the expression of miR-9 increases spatially and temporarily over zebrafish development. A detailed time course of the expression of 7 pri-miR-9 genomic loci shows that they have distinct temporal and spatial profiles, which may be brought about by different numbers of E/N-boxes in their regulatory regions. Focusing on pairs that are expressed in the same area of the hindbrain region, namely pri-miR-9-1/pri-miR-9-4 and pri-miR-9-1/pri-miR-9-5, we find that they sequentially activated during neurogenesis, concurrent with a known change in Her6 dynamics from noisy to oscillatory. Analysis of expression at the single-cell level shows that although they are sequentially activated, early and late pri-miRs are mostly concurrently transcriptionally active in the same cells. This finding supports the idea that increased mature miR-9 is contributed, at least in part, by overlapping activation of distinct loci. We propose that this may lead to an additive, sharp, increase of mature miR-9 which in turn may lead to rapid changes in Her6 dynamics.

Introduction

MicroRNAs are a class of small (~22 nucleotides) regulatory non-coding RNAs, which regulate gene expression at the post-transcriptional level. These small RNAs are processed from large microRNA primary transcripts (pri-miRs) into 70~90nt precursors (pre-miRs) before further splicing into ~22nt mature microRNA. miR-9 is a highly conserved microRNA which is expressed predominantly in the central nervous system (CNS) of vertebrates and plays a crucial role during CNS development.

Previous work in xenopus, zebrafish and mice has shown that miR-9 is essential for the correct timing of cell fate transitions during neurogenesis (Coolen et al., 2013) (Bonev et al., 2011; Bonev et al., 2012). miR-9 post-transcriptionally targets many transcription factors that are involved in neural development such as FoxG1 (Shibata et al., 2008), Tlx (Zhao et al., 2009) and members of the Hes/Her helix-loop-helix family of transcription factors, including Hes1 in mouse and Xenopus (Bonev et al., 2011; Bonev et al., 2012) and Her6 in zebrafish (Coolen et al., 2012; Galant et al., 2016; Soto et al., 2020).

The Hes/Her family of proteins are expressed dynamically in an oscillatory manner at the ultradian timescale (Hirata et al., 2002; Shimojo et al., 2008). Hes/Her oscillations are achieved by a negative feedback loop, whereby Hes/Her proteins inhibit their own transcription coupled with a rapid turnover of protein and mRNA. Instability of both protein and mRNA allows for levels of the protein to fall, de-repression to occur and expression to resume, generating a cyclic pattern (Hirata et al., 2002; Novak and Tyson, 2008). Indeed, both mRNA and protein of Hes family

genes are unstable, for example, in mice, the half-life of *Hes1* mRNA is ~24 minutes and the HES1 protein half-life is in the order of 22 minutes (Hirata et al., 2002) and the Her6 (Zebrafish orthologue) protein half-life is around 10 minutes (Soto et al., 2020).

The instability of the mRNA, as well as the translation of the protein, are controlled by microRNAs. Indeed, our previous work revealed that miR-9 regulation is important for allowing the oscillatory pattern of expression of HES1 to emerge. However, the amount of miR-9 is important as too much or too little miR-9 can lead to dampening of HES1 oscillations (Bonev et al., 2012; Goodfellow et al., 2014). We have also recently shown that in zebrafish, the dynamics of Her6 protein expression switch from noisy to oscillatory and this coincides temporally with the onset of miR-9 expression in the hindbrain (Soto et al., 2020). Furthermore, when the influence of miR-9 on *her6* is removed experimentally, Her6 expression does not evolve away from the "noisy" regime with a consequent reduction in progenitor differentiation. We have interpreted this to mean that the miR-9 input is necessary to constrain gene expression noise, enabling oscillations to occur and to be decoded by downstream genes (Soto et al., 2020). Together these findings support that Hes/Her dynamics are sensitive to the amount of mature miR-9 present in the cell, however, the mechanism by which this is controlled is not known.

The question is complicated by the observation that vertebrates (and some invertebrates) possess multiple copies of the miR-9 gene at distinct loci but all capable of producing the same mature microRNA. For example, both human and mouse contain 3 copies of miR-9 (Rodriguez-Otero et al., 2011; Shibata et al., 2011)

and frogs have 4 (Walker and Harland, 2008). Due to an additional round of whole-

genome duplications in teleost fish (Amores et al., 1998; Jaillon et al., 2004),

zebrafish have 7 paralogues of miR-9 (miR-9-1 to miR-9-7) (Chen et al., 2005).

One possibility is that different genomic loci contribute to miR-9 regulation in a

qualitative way, that is due to differential temporal and spatial specificity of mature

miR-9 expression. Indeed, there is some limited evidence that these discrete copies

of miR-9 are expressed differentially during development both temporally and

spatially (Nepal et al., 2016; Tambalo et al., 2020). Another, and yet unexplored,

possibility is that transcription from different loci may serve to control miR-9

quantitatively, that is to increase the amount of miR-9 in the cell and perhaps do so

in a temporally controlled manner.

Here, we undertake a systematic study of pri-miR-9 expression in zebrafish that aims

to address the likelihood of these distinct scenarios, with special attention to the

possibility of a quantitative control mechanism. We show by *in situ* hybridization that

the amount of mature miR-9 increases over zebrafish development, spreading from

the forebrain to the hindbrain between 24 and 48hpf. A detailed time course of the

expression of all 7 pri-miR-9 paralogues shows that they are all transcriptionally

active, but exhibit subtle, yet distinct, temporal and spatial profiles which could be

contributed by differences in their regulatory regions, for example, correlating with

the number of E/N-boxes present in the proximal promotor of each pri-miR-9.

Focusing on 2 pairs of early and late expressed pri-miR-9s, (pri-miR-9-1 and pri-miR-

9-4, or pri-miR-9-5) by fluorescent in situ hybridization at single cell level, we find

that, surprisingly, in some cells early and late pri-miR-9s were concurrently

transcriptionally active. Taken together, we find evidence for a qualitative mechanism

in the deployment of pri-miR-9s, as well as a previously un-appreciated quantitative

component, both of which may contribute to the decoding function of mature miR-9s.

RESULTS

All pri-mir-9s are expressed but with differed temporal onset

To investigate the expression of the mature miR-9 in zebrafish embryos, we first

performed a whole mount in situ hybridization (WM-ISH) for the mature miR-9 using

an LNA (locked nucleic acid) probe. Mature miR-9 was detected only in forebrain at

24hpf (**Fig. 1a**) while at 30hpf miR-9 was observed in the midbrain and in

rhombomere 1 (r1) of the hindbrain (hb), maintaining high expression in forebrain

(**Fig. 1a**, 30hpf-red head arrow). Later in development miR-9 expression was seen in

the posterior hindbrain (**Fig. 1a**, 34-36hpf-blue arrow). As development progressed,

increased levels of expression throughout the hindbrain were observed while

expression in the forebrain decreased (**Fig. 1a**, 48hpf, blue arrow: hindbrain, green

arrow: forebrain). These results show a temporally controlled antero-posterior wave

of miR-9 expression along the brain/hindbrain axis.

In zebrafish, the mature miR-9 can be produced from seven paralogous of miR-9.

The miR-9 paralogues occupy 7 unique loci across the genome (GRCz11; Genome

Reference Consortium Zebrafish Build 11) (Yates et al., 2020). With the exception of

miR-9-3 which is located upstream of a lincRNA, all miR-9 genes are intragenic,

overlapping annotations of lincRNAs or proteins(Yates et al., 2020) (**Figure S1a,b**).

Our *in silico* analysis of previously published RNA-seq data confirms the differential

temporal expression of six of the seven miR-9 paralogues host genes (White et al.,

2017). It is also clear that expression of miR-9 host genes coincides with a gradual

decline in the expression of Her/Hes family genes expression, consistent with the

idea that *hes/her* genes are major targets of miR-9 (**Figure S1c**) (Bonev et al.,

2011).

Previous work has revealed that the 7 miR-9 zebrafish paralogues are expressed in

forebrain at early stages of neurogenesis while toward the end are also expressed in

hindbrain (Nepal et al., 2016). However, little is known about the period spanning the

peak of neurogenesis, when miR-9 controls downstream targets such as the

ultradian oscillator Her6. To characterize the expression in greater spatiotemporal

detail, particularly over the hindbrain area where Hes/Her target genes are

expressed, we investigated the nascent expression of all 7 precursors over a time

period spanning the peak of neurogenesis which occurs at 33hpf (Lyons et al., 2003)

using specific probes for each pri-miR-9 (**Figure S2; Methods**, Molecular cloning).

We observed that all pri-miR-9s were first expressed in the forebrain (24hpf), in a

regional specific manner, which was not further characterised here. At 48hpf they are

all also expressed in hindbrain (**Fig. 1b,d**, 24 and 48hpf and **Figure S3a-c**)

consistent with what has been described before (Nepal et al., 2016). Differential

expression was evident in the intermediate stages. Specifically, pri-miR-9-3, 9-4 and

9-5 were expressed ahead of the others in the hindbrain. Here, pri-miR-9-4 was

evenly and highly expressed while pri-miR-9-3 and pri-miR-9-5 were expressed in a

gradient from anterior to posterior (**Fig. 1b,c**; 30hpf; blue arrowhead). At the peak of

hindbrain neurogenesis (34-36hpf), pri-miR-9-2 and pri-miR-9-7 were upregulated,

joining most of the pri-miR-9s that were highly expressed at this stage (**Fig. 1b**, 34-

36hpf). Pri-miR-9-1 and miR-9-6 were temporally delayed until 48hpf, at which point

all pri-miR-9 were fully expressed.

Thus, overall, every pri-miR-9 was expressed in the CNS and exhibited similar but

unique expression patterns and a temporal progression. Characterising the temporal

order of the onset of expression in intermediate stages was a first step in asking

whether the expression of different pri-miR-9s is distinct or additive in the hindbrain,

which we address below.

Partially overlapping spatial expression of early pri-miR-9s

We selected three different precursors based on the onset of their hindbrain

temporal expression during development, earliest or latest, and the phylogenetic

analysis of precursors sequences in clades/subgroups performed by Alwin Prem

Anan et al., 2018. Pre-miR-9-1 was selected as the latest to express in the hindbrain

belonging to clade I/subgroup II, pre-miR-9-5 as the earliest to express that belongs

to clade I/subgroup I and pre-miR-9-4 as the earliest, belonging to clade II (Alwin

Prem Anand et al., 2018) (**Figure S3d**).

Double WM-FISH for pri-miR-9-1 and pri-miR-9-4 performed on stage 32hpf

embryos revealed expression of pri-miR-9-4 along the antero-posterior hindbrain

axis, while pri-miR-9-1 expression was limited to the region of the anterior hindbrain

corresponding to rhombomere 1 (**Fig. 2a**, red arrows). The transversal view of

rhombomere 4 reveals expression of pri-miR-9-4 within the ventricular zone, while

pri-miR-9-1 was almost completely absent (**Fig. 2d, f**). These data indicate that pri-

miR-9-s are mainly expressed in the region where most of the dorsal progenitors,

medial progenitors and ventral progenitors are found (**Fig. 2f**) (Lyons et al., 2003;

Tambalo et al., 2020). The double WM-FISH experiment was repeated with pri-miR-

9-1 and pri-miR-9-5. At 32hpf, pri-miR-9-5 is evenly distributed antero-posteriorly in

the hindbrain whilst pri-miR-9-1 was mostly expressed in rhombomere 1 as one

would expect due to the delayed onset of expression for pri-miR-9-1 (**Fig. 2b**, red

arrows). Fluorescent intensity quantification of the region of interest (**Fig. 2a,b**

merge, yellow rectangle) show variability in the levels of expression, pri-miR-9-4 and

pri-miR-9-5 are significantly high expressed ($P<0.05$ and $P<0.0001$, respectively) to

pri-miR-9-1 (**Fig. 2c),** supporting the hypothesis of pri-miR-9-4 and pri-miR-9-5 early

expression. Transversal imaging of pri-miR-9-1/pri-miR-9-5 revealed similar pattern

expression to pri-miR-9-1/pri-miR-9-4 analysis. Interestingly, subtle differences were

observed in pri-miR-9-4 which had a broader pattern of expression, extending further

in the dorsal progenitor area than pri-miR-9-5 indicating some differences in spatial

expression between pri-miR-9s of the same temporal profile (**Fig. 2d-e**, compare

middle panels, **Fig. 2f**).

Progressive additive expression of pri-miR-9s

In order to examine whether the late expression of pri-miR-1 is cumulative with pri-

miR-4 and pri-miR-5, or whether it is spatially distinct, we analysed expression at

later stages. Coronal and transversal views of pri-miR-9-1 double WM-FISH with pri-

miR-9-4 and pri-miR-9-5 revealed both overlapping and distinct expression of the

primary transcripts (**Fig. 3a-d**). Distinct expression was most clearly observed in

transversal views that show pri-miR-9-5 is more restricted in its expression to the

middle-ventral progenitor region of the ventricular zone, while pri-miR-9-1 was more

broadly expressed toward the dorsal progenitor region (**Fig. 3d, e**). In order for the

overlapping expression to contribute to the total levels of mature miR-9 in a cell,

early and late pri-miR-9s would need to be expressed in the same cells, which was

examined next.

Total mature miR-9 is contributed by overlapping activation of distinct miR-9

loci

To investigate the existence of a spatiotemporal overlap in pri-miR-9 expression at

the single cell level, we performed double WM-FISH for pri-miR-9-1 with pri-miR-9-4

and pri-miR-9-5. Cell boundaries were visualized with the cell boundary marker

BODIPY-texas red (BP-TR). At 48hpf we observed cells that have only one miR-9

precursor present (**Fig. 4c,d,g,h**) from which, cells expressing only pri-miR-9-1 were

often found in the dorsal progenitor region of the ventricular zone (**Fig. 4a,c,e,g**)

while cells expressing only pri-miR-9-4 or pri-miR-9-5 and not pri-miR-9-1 were

mostly observed in the middle-ventral progenitor region of the ventricular zone (**Fig.**

4a,d,e,h) suggesting differential spatial expression. However, we also found many

cells co-expressing two pri-miR-9s (**Fig. 4a,b,e,f**), suggesting that different miR-9

paralogues are concurrently transcriptionally active in the same cells.

Diverse regulation of miR-9 promoters

To have a better understanding of the origin of these spatial and temporal

expression differences, we focused on the transcriptional

activation/repression/regulation of the different miR-9 precursors. Transcriptional

repressors such as Hes/Her family can inhibit expression through the binding to N/E-

boxes (Akazawa et al., 1992; Sasai et al., 1992). An E-box (enhancer box) is a DNA

response element that can act as a protein-binding site of an activator or repressor,

while an N-box is a protein-binding for a repressor such as Hes1(Sasai et al., 1992).

We first performed an *in-silico* analysis to identify putative canonical DNA binding

motif of the bHLH transcription factors (E-boxes), with consensus sequence

CANNTG, in a 2kb region upstream and 1kb region downstream of the

transcriptional start site of the transcripts of the miR-9 paralogues as previously

defined in Chirag et al. 2016. We found miR-9-2 and miR-9-4 promotor/proximal

enhancer region contained the highest number of putative E-boxes, 21 and 18,

respectively, followed by miR-9-3, 5, 7, 1 and 6 (**Fig. 5a, b**). In concordance, the

bHLH transcription factor Neurogenin1 (Ngn1) is a pro-neural gene known to induce

miR-9 thus is a potential candidate to promote differential temporal expression of the

seven miR-9 paralogues through E-box binding (Zhao et al., 2015)

We next performed *in silico* analysis of the DNA binding motif sequence CACNAG

called, N-box. As previously mentioned, the transcription of miR-9 precursors is

repressed by Hes1 protein through this binding site (Bonev et al., 2012; Sasai et al.,

1992). We identified variability within host gene regulatory sequences; 9 putative N-

boxes were found in miR-9-5, 5 N-boxes in miR-9-2, 3 N-boxes in miR-6 while only

1 in miR-9-7, in miR-9-4 and miR-9-1 and none in miR-9-3 (**Fig. 5a, b**). These results

may reflect variability of Hes1 strength regulation over the miR-9s proximal promotor

dependent on the number of N-boxes that each miR-9 promoter/proximal enhancer

region contains. Indeed, we observed that expression levels of pri-miR-9-5 is

consistently reduced (**Fig. 5e**. dashed line) in the regions where the zebrafish Hes1

orthologues, Her6/Her9, are highly expressed (**Fig. 5c**. her6/her9 merge-white

underlined), corresponding to the medial-dorsal progenitor region of the ventricular

zone and pri-miR-9-1 is expressed more broadly, with expression in the dorsal

progenitor region of the ventricular zone. We also observed expression of pri-miR-9-

4 narrowed to the ventral progenitor region of the ventricular zone (**Fig. 5d**, dashed

line), this can be explained by its higher number of E-boxes when compared to pri-

miR-9-1. Combined, these data outline the dynamical spatial expression of the highly

conserved microRNA, miR-9, based on the ability of repressors/activators to

modulate the different promoters of the seven different mir-9 precursors.

Discussion

miR-9 is expressed from several genomic loci which, after transcription and

processing, produce the same 5' mature form of miR-9 which is particularly

interesting because it targets the key neural progenitors Her/Hes TFs. How common

is this multi-locus organisation? In humans 6.3% of mature microRNA arms are

identical across two or more loci (Kozomara et al., 2019), thus, it is not very

common, but it is not unique to miR-9. In zebrafish this number rises to around

32.3% (Kozomara et al., 2019). The higher number of microRNA expressed from

multiple loci is possibly due to the teleost-specific whole-genome duplication (WGD).

Evidence from rainbow trout also shows that following the salmonid-specific extra

round of WGD, microRNAs appear to be retained at higher levels than protein-

coding genes (Berhelot et al 2014). This may suggest that extra copies of microRNA

are evolutionary advantageous but why this is the case was not understood. Here,

we propose that retention of multiple microRNA loci could have specific functional

advantages for regulatory control of target gene expression of an organism.

By examining in detail the temporal and spatial expression at single cell level of 3

select early and late pri-miR-9s, we offer two possible explanations for this multi-site

organisation or primary transcripts.

The first explanation involves a qualitative mechanism. In this scenario, distinct pri-

miR-9s have different spatial expression, which allows them to target different, i.e.

region-specific, genes. Some of the differences in the spatial expression of pri-miR-

9s are easily discernible macroscopically (e.g. differential expression in the

forebrain) while others are subtle and require post-hybridisation sectioning to

document, as we have done here. An example of the latter is the expression of pri-

miR-1 which extents more dorsally in the hindbrain than pri-miR-4. This correlates

well with the expression of her6 and her9, which are both miR-9 targets but are

expressed adjacent to each other along the D-V axis (Soto et al. 2020).

The second explanation favours a quantitative mechanism. In this scenario, the

differential temporal expression, where some primary transcripts commence their

expression early while others are only expressed late, results in the simultaneous

expression of both (or more) transcriptional loci in the same cells at a particular time

in development. In support of this scenario, we have shown by FISH that pri-miR-9-

1, a late onset pri-miR-9, is co-expressed in the same cells as the earlier onset pri-

miR-9-4 or -5. Interestingly, we have not seen evidence for a mechanism where one

pri-miR-9 switches off and another one comes on, arguing against a mechanism

where an early pri-miR-9 "passes on" the task of repression to a later expressed one;

rather all loci seem to remain active at least for the duration of our observations

which covers the period of embryonic neurogenesis. This means that both early and

late pri-miR-9s are concurrently transcribed at late stages, and assuming they are

both processed the amount of mature miR-9 in a cell could sharply increase. Given

that transcription saturates easily in many systems (Hafner et al., 2020), this co-

expression may be a strategy to increase the amount of miR-9 available to the cell

than what is possible with transcription from one locus alone. Indeed, we have

shown that the dynamical regime of Hes1 (i.e. oscillatory expression to stable

expression at different levels) as well as the amount of time that Hes1 oscillates for,

depends on the amount of miR-9 in the cell (Bonev et al., 2012; Goodfellow et al.,

2014). More recently, we have shown by in vivo manipulations, that the input of miR-

9 changes the dynamic expression of her6 from noisy to oscillatory (Soto et al.,

2020).

Like many other miRs, miR-9 has been shown to be quite stable in xenopus and was

thus presumed to accumulate gradually over time (Bonev et al., 2012; Goodfellow et

al., 2014). In turn, changing the levels of miR-9 in the cell would drive the dynamics

of Hes1 from one dynamical regime to another, as described above. While more

recent studies have questioned the long stability of miRs by using different methods

and showing that the stability of miR-9 varies between tissues (Kim et al., 2020;

Marzi et al., 2016; Ruegger and Grosshans, 2012), the onset of additive transcription

from two loci is nevertheless likely to exert a sharp increase in the amount of miR-9

present in the cell. It is possible that some genes/networks do not respond to slow

increases of miR-9 (a form of adaptation observed in signalling pathways (Dessaud

et al., 2007)), and a sharp, non-linear, increase may be needed to push a dynamical

system into a new state associated with a cell fate change. In fact, non-linearity of

reactions is a key feature of systems that can generate oscillatory gene expression

(Novak and Tyson, 2008).

The qualitative and quantitative mechanisms suggested above are not mutually

exclusive and may be combined and also take place at the same time. A common

element is that both mechanisms would rely on distinct control of expression of

different miR-9 loci. In support of this, we have found that characteristics of the

regulatory regions and the organisation of pri-miR-9 in their host transcriptional units

show some distinct features. Interestingly, some of the regulatory region differences

may also be quantitative rather than simply qualitative; Indeed, we have found a

difference in the number of repressive N-boxes but also differences in the number of

activators E-boxes, which may correlate with the differences in the extent of medio-

lateral (ontogenetically ventral-dorsal) expression of some pri-miR-9 pairs. Another

common element is that in both scenarios, a sequential temporal order of activation

can be involved, and indeed, we have observed a temporal sequence of activation

for most pri-miR-9s, roughly starting anteriorly and spreading posteriorly. This

temporal order was also observed when considering the entire miR-9 family and the

Her gene targets which were expressed on the whole earlier and were

downregulated when pri-miR-9s were upregulated.

Where multiple paralogues of a microRNA are present, differential qualitative and

quantitative regulation may be a more common feature than is currently appreciated.

In this respect, it is interesting that a recent study found that miR-196 paralogues

show both unique and overlapping expression. Single KOs showed some

redundancy but importantly, they also showed unique phenotypes and combinatorial

KOs showed better penetrance together with additional defects, suggesting an

additive role of miR-196 paralogues in establishing vertebral number (Wong et al.,

2015).

In conclusion, by providing evidence for both a quantitative and qualitative

mechanism, we have made conceptual advances on the possible roles of organising

pri-miR-9s in several distinct genomic loci, which may have led to their evolutionary

conservation. In addition, we highlight here some practical benefits of our work for

the experimenter; once mature miR-9 has been produced, it is not possible to tell

which genomic locus it was transcribed from. Thus, with multiple such loci being

potentially involved, it is very difficult for the experimenter to select the correct one to

tag, mutagenize or otherwise manipulate by CRISPR/Cas9. Therefore, an added

benefit of our work is that the detailed characterisation we have described here will

enable the selection of the correct genomic locus for genetic manipulation of miR-9

production, depending on the precise research question.

Acknowledgements

We are grateful to Dr Laure Bally-Cuif for sharing plasmids. The authors would also

like to thank the Biological Services Facility, Bioimaging and Systems Microscopy

Facilities of the University of Manchester for technical support. This work was

supported by a Wellcome Trust Senior Research Fellowship to NP (090868/Z/09/Z).

The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

Author Contributions

Conceptualization, X.S. and N.P.; Methodology, X.S. and N.P; Software, T.M. and
X.S.; Validation, X.S., T.M and R. L.; Formal Analysis, X.S. and T. M. ; Investigation,
X.S., R.L, J. L. and T.M.; Resources, N.P. and X.S.; Data Curation, X.S. and T.M..;
Writing - Original Draft, X.S.; Writing – Review & Editing, N.P., X.S. and T.M..;
Visualization, X.S., T.M and R.L.; Supervision, N.P. and X.S; Project Administration,
N.P and X.S.; Funding Acquisition, N.P.
382
383 **Declaration of Interests**
384
385 The authors declare no competing interests.
386
387 **Methods**
388 **Research Animals**
389 Animal experiments were performed under UK Home Office project licenses
390 (PFDA14F2D) within the conditions of the Animal (Scientific Procedures) Act 1986.
391 Animals were only handled by personal license holders.
392
mRNA extraction and Quantitative real-time PCR (qRT-PCR)
mRNA was extracted using Trizol from a pool of 10 zebrafish embryos. SuperScript
II (Invitrogen) was used to do reverse cDNA synthesis using hexamer primers. For
quantitative real-time PCR (qRT-PCR) cDNA samples were diluted to a
concentration of 50 ng/µl with dH_2O. Each qPCR reaction was prepared in triplicate in
a 96-well plate with each well consisting of 0.2 µM each forward and reverse primer,
50ng cDNA and SYBR Green Mastermix (ThermoFisher). Zebrafish embryo
Reactions were run on Step One Plus Real-time PCR System (Applied Biosystems)
alongside negative controls. The data for each sample was normalized to the
expression level of *b-actin* and analysed by the $2^{-\Delta\Delta Ct}$ method, normalized to miR-9-1
expression. For each primer pair, the PCR product was examined by gel

electrophoresis and its melting curve to ensure a single fragment of the predicted

molecular weight.

Molecular cloning

RNA probes for pri-miR-9-1, pri-miR-9-2, pri-miR-9-4, pri-miR-9-5 and pri-miR-9-7

were PCR amplified and cloned into pCRII vector using primers described in **Table**

S1. Except for pri-miR-9-2 probe, they were designed to distinguish the primary

transcripts by including sequences, intron and exon, before and after each

microRNA processing, while also covering the sequence corresponding to mature

miR-9 **(Figure S2)**. Since the mature miR-9 sequence is conserved between

paralogs, to avoid any cross-binding of probes to this sequence we mutated it on

each probe by using QuikChange II XL Site-Directed Mutagenesis assay. This

allowed us introduce deletions and single nucleotide exchange in specific regions of

the mature miR-9 sequence (**Table S2**; **Figure S2**; sequence highlighted in red).

pri-miR-9-3 and pri-miR-9-6 probes were generated from plasmids kindly gifted by

Laure Bally-Cuif (Nepal et al., 2016).

Whole mount chromogenic and fluorescence _in situ_ hybridization and

sectioning

Chromogenic _in situ_ hybridisation was performed as previously described by

Christine Thisse (Thisse and Thisse, 2008). Multicolour fluorescence _in situ_

hybridisation was modified from Hoppler and Vize (Lea et al., 2012) by developing

with tyramide amplification (Perkin Elmer) after addition of antisense RNA probes

and antibodies conjugated to horseradish peroxidase (Lea et al., 2012).

Sections were obtained as described in Dubaissi (Dubaissi et al., 2012) with

modifications. Embryos were embedded in 25% fish gelatine and 30% sucrose for a

minimum of 24 hrs. 18µm thickness hindbrain sections were collected and

transferred onto superfrost glass slides. The slides were air dried overnight under the

fume hood and stained with 5µM BODYPI-TR (Thermo-Fisher Scientific) before

mounting with Prolong Diamond Antifade.

Imaging

Chromogenic *in situs* were imaged using a Leica M165FC with a DFC7000T camera.

Fluorescent *in situ* sections were imaged using Leica TCS SP5 upright confocal with

HCX PL APO CS 40.0x1.25 OIL UV lens or Olympus FLUOVIEW FV1000 confocal

with UPLSAPO 20X NA:0.75 lens or UPLSAPO 20X NA:0.75 lens.

Expression analysis of hes/her genes and microRNA hosts

For the *in silico* analysis of the microRNA host gene expression we downloaded the

time course RNA-seq data (TPM) from White et al. 2017 supplemental file 3. Here

we used the overlapping host genes as a proxy for the expression of the microRNA.

MicroRNA would not show up in standard RNA-seq analysis and there is no current

microRNA time course data. Host genes were identified as those with overlapping

annotations with the miR-9 genes. The host genes for each microRNA are in **Table**

S4. MiR-9-7 has no overlapping annotation at this time and is thus not reported on in

these data.

We filtered the RNA-seq data removing genes which were neither the host genes of

the microRNA or members of the Her family. 3 repeats for each stage of

development are included in the data and we averaged the expression across the 3

repeats for each stage. The stages reported in the data are based on standard

embryonic stages in zebrafish development. However, we wanted to visualize the

expression in terms of hours and the stages were converted accordingly. Finally,

before plotting these data were z-scored to normalize the expression of each of the

genes so that we could compare changes in expression over time rather than

absolute levels. These data were then plotted using the heatmap.3 package in R.

N-box / E-box analysis

To assess the number of N and E -box elements in the promoter/proximal enhancer

region of the transcriptional start site (TSS) of the microRNA host genes/pri-miRs

(Table S4) we downloaded the *Danio rerio* genome from Ensembl. Using the

genome coordinates from (Nepal et al., 2016) (supplementary Table-S4) for the miR-

9 TSS we took a region of 2kb upstream to 1kb downstream of the TSS. The N and

E-boxes in each region were then counted. The N and E-box sequences can be

found in **Table S5**. These data were then plotted using R (R Core, 2019) and the

ggplot2 library (Wickham, 2016)

Figure Legends

Figure 1. Pri-miR-9 paralogues are expressed with different temporal onset. (a)

Representative example of chromogenic whole mount in situ hybridization (WM-ISH)

of miR-9 using miR-9 LNA 5'-Dig observed at different stages during development;

longitudinal view, anterior to the left. Green arrow: forebrain (fb) expression. Blue

arrow: hindbrain (hb) expression. **(b)** Chromogenic WM-ISH (Cro-WMISH) of

different pri-miR-9s using specific probes for each paralogue observed at different

stages during development; longitudinal view, anterior to the left. Blue arrowhead:

expression in hb at 30-31hpf, light blue arrowhead: expression in hb at 34-36hpf,

blue arrow: expression in hb at 48hpf. **(c)** High magnification of hb region from

respective paralogue at 30-31hpf. (**d**) Heatmap plot representing expression levels in

hindbrain; grade 0= no expression in hindbrain, grade 1= weak expression in

hindbrain (rhombomere 1 or 2), grade 2= distinct expression in hb and grade 3=

strong expression in hb.

Figure 2. Partially overlapping expression of pri-mIR-9s. (a, b) Representative

example of double Fluorescent WM-ISH (WM-FISH) labelling of pri-miR-9-1/pri-miR-

9-4 **(a)** and pri-miR-9-1/pri-miR-9-5 **(b)** in hindbrain, coronal view, from wild-type

embryo observed at 32hpf; merged images (**a, b;** bottom panel) indicate pri-miR-9-

4(**a**; magenta**)** and pri-miR-9-5 (**b**; magenta**)** are expressed along the antero-

posterior axis while pri-miR-9-1 is limited to anterior hindbrain (**a, b;** green, red

arrows). Annotation denotate hindbrain (HB). **(c)** Pixel intensity mean of pri-miR-9-1,

pri-miR-9-4 and pri-miR-9-5 observed in ROI (region of interest), delineated by the

yellow rectangle in **(a)** and **(b).** Error bar in Scatter dot plot represent mean and SEM

of 32-34hpf pri-miR-9-1 (28 slices, 6 embryos), 32-34hpf pri-miR-9-4 (8 slices, 3

embryos) and 32-34hpf pri-miR-9-5 (18 slices, 3 embryos). Kruskal-Wallis with

Dunn's multiple comparison test, significance: $p<0.05^*$, $p<0.0001^{****}$. **(d, e)**

Representative example of double WM-FISH labelling of pri-miR-9-1/pri-miR-9-4 **(d)**

and pri-miR-9-1/pri-miR-9-5 **(e)** in hindbrain rhombomere 4 (r4), transversal view,

from wild-type embryo observed at 32hpf; merged images (**d, e;** right panel) indicate

pri-miR-9-4(**d**; magenta**)** and pri-miR-9-5 (**e**; magenta**)** are expressed in the

ventricular zone region while pri-miR-9-1 is almost absent (**d, e;** green). Scale bar 30

μm. **(f)** Schematic representation of transverse section from 32hpf zebrafish

hindbrain at the level of the otic vesicle. VZ: ventricular zone, region where most of

progenitor cells are located. MZ: mantle zone, region of ongoing neurogenesis.

Within the VZ there are dorsal progenitors (DP), medial progenitors (MP) and ventral

progenitors (VP).

Figure 3. Progressive additive expression of pri-miR-9s. (a, b) Representative

example of double WM-FISH labelling of pri-miR-9-1 (left panel) and pri-miR-9-4

(middle panel) in hindbrain from wild-type embryo observed at 48hpf, coronal view

(a) and transversal view **(b). (c, d)** Representative example of double WM-FISH

labelling of pri-miR-9-1 (left panel) and pri-miR-9-5 (middle panel) in hindbrain from

wild-type embryo observed at 48hpf, coronal view **(c)** and transversal view **(d)**;

merged images in coronal view **(a, c; right)** shows overlapping expression of the

primary transcripts yet in transversal view **(b, d; right)** reveals distinct expression,

pri-miR-9-1 is expressed more broadly toward the dorsal progenitor region. Scale bar

30μm. hb: hindbrain **(e)** Schematic representation of transverse section from 48hpf

zebrafish hindbrain at the level of the otic vesicle. VZ: ventricular zone, region where

most of progenitor cells are located. MZ: mantle zone/marginal zone, region of

ongoing neurogenesis and mature neurons. Within the VZ there are dorsal

progenitors (DP), medial progenitors (MP) and ventral progenitors (VP).

Figure 4. mature miR-9 produced by overlapping activation of distinct miR-9

loci. (a-d) Representative example of double WM-FISH labelling pri-miR-9-1 and pri-

miR-9-4 combined with cell boundary staining, BODIPY-texas red (BP-TR). **(b)**

Indicates cells expressing both, pri-miR-9-1 (green) and pri-miR-9-4 (magenta). **(c)**

Indicates cells expressing only pri-miR-9-1 (green). **(d)** Indicates cells expressing

only pri-miR-9-4 (magenta). **(e-h)** Representative example of double WM-FISH labelling pri-miR-9-1 and pri-miR-9-5 combined with BP-TR. **(f)** Indicates cells expressing both, pri-miR-9-1 (green) and pri-miR-9-5 (magenta). **(g)** Indicates cells expressing only pri-miR-9-1 (green). **(h)** Indicates cells expressing only pri-miR-9-5 (magenta). Images were acquired from hindbrain r4, transversal sections, from 48hpf wild-type embryos. Scale bars 30μm and 5μm.

Figure 5. Diverse regulation of miR-9 promoters. (a) Bar plot showing the number of N (green) and E (blue) boxes around the TSS (2kb upstream, 1kb downstream) of the miR-9 genes. **(b)** Scatter plot showing the relationship between the number of N boxes and E boxes around the TSS (2kb upstream, 1kb downstream) of the miR-9 genes**. (c)** Representative example of double WM-FISH labelling *her9* (green) and *her6* (magenta) in hindbrain, transversal view, from wild-type embryo observed at 44hpf, white line indicates dorsal and medial progenitor region. Scale bar 30μm. **(d,e)** Representative example of double WM-FISH labelling pri-miR-9-1 (green)/pri-miR-9-4 (magenta) **(d)** and pri-miR-9-1 (green)/pri-miR-9-5 (magenta) **(e)** in hindbrain, transversal view, from wild-type embryos observed at 48hpf. Dashed white line indicates ventral progenitor region. Scale bar 30μm.

REFERENCE

Akazawa, C., Sasai, Y., Nakanishi, S., and Kageyama, R. (1992). Molecular characterization of a rat negative regulator with a basic helix-loop-helix structure predominantly expressed in the developing nervous system. J Biol Chem *267*, 21879-21885.

Alwin Prem Anand, A., Huber, C., Asnet Mary, J., Gallus, N., Leucht, C., Klafke, R., Hirt, B., and Wizenmann, A. (2018). Expression and function of microRNA-9 in the mid-hindbrain area of embryonic chick. BMC Dev Biol *18*, 3.

Amores, A., Force, A., Yan, Y.L., Joly, L., Amemiya, C., Fritz, A., Ho, R.K., Langeland, J., Prince, V., Wang, Y.L.*, et al.* (1998). Zebrafish hox clusters and vertebrate genome evolution. Science *282*, 1711-1714.

Bonev, B., Pisco, A., and Papalopulu, N. (2011). MicroRNA-9 reveals regional diversity of neural progenitors along the anterior-posterior axis. Dev Cell *20*, 19-32.

Bonev, B., Stanley, P., and Papalopulu, N. (2012). MicroRNA-9 Modulates Hes1 ultradian oscillations by forming a double-negative feedback loop. Cell Rep *2*, 10-18.

Chen, P.Y., Manninga, H., Slanchev, K., Chien, M., Russo, J.J., Ju, J., Sheridan, R., John, B., Marks, D.S., Gaidatzis, D.*, et al.* (2005). The developmental miRNA profiles of zebrafish as determined by small RNA cloning. Genes Dev *19*, 1288-1293.

Coolen, M., Katz, S., and Bally-Cuif, L. (2013). miR-9: a versatile regulator of neurogenesis. Frontiers in cellular neuroscience *7*, 220.

Coolen, M., Thieffry, D., Drivenes, O., Becker, T.S., and Bally-Cuif, L. (2012). miR-9 controls the timing of neurogenesis through the direct inhibition of antagonistic factors. Dev Cell *22*, 1052-1064.

Dessaud, E., Yang, L.L., Hill, K., Cox, B., Ulloa, F., Ribeiro, A., Mynett, A., Novitch, B.G., and Briscoe, J. (2007). Interpretation of the sonic hedgehog morphogen gradient by a temporal adaptation mechanism. Nature *450*, 717-720.

Dubaissi, E., Panagiotaki, N., Papalopulu, N., and Vize, P.D. (2012). Antibody development and use in chromogenic and fluorescent immunostaining. Methods Mol Biol *917*, 411-429.

Galant, S., Furlan, G., Coolen, M., Dirian, L., Foucher, I., and Bally-Cuif, L. (2016).

Embryonic origin and lineage hierarchies of the neural progenitor subtypes building

the zebrafish adult midbrain. Dev Biol *420*, 120-135.

Goodfellow, M., Phillips, N.E., Manning, C., Galla, T., and Papalopulu, N. (2014).

microRNA input into a neural ultradian oscillator controls emergence and timing of

alternative cell states. Nature communications *5*, 3399.

Hafner, A., Reyes, J., Stewart-Ornstein, J., Tsabar, M., Jambhekar, A., and Lahav,

G. (2020). Quantifying the Central Dogma in the p53 Pathway in Live Single Cells.

Cell Syst *10*, 495-505 e494.

Hirata, H., Yoshiura, S., Ohtsuka, T., Bessho, Y., Harada, T., Yoshikawa, K., and

Kageyama, R. (2002). Oscillatory expression of the bHLH factor Hes1 regulated by a

negative feedback loop. Science *298*, 840-843.

Jaillon, O., Aury, J.M., Brunet, F., Petit, J.L., Stange-Thomann, N., Mauceli, E.,

Bouneau, L., Fischer, C., Ozouf-Costaz, C., Bernot, A*., et al.* (2004). Genome

duplication in the teleost fish Tetraodon nigroviridis reveals the early vertebrate

proto-karyotype. Nature *431*, 946-957.

Kim, C.K., Asimes, A., Zhang, M., Son, B.T., Kirk, J.A., and Pak, T.R. (2020).

Differential Stability of miR-9-5p and miR-9-3p in the Brain Is Determined by Their

Unique Cis- and Trans-Acting Elements. eNeuro *7*.

Kozomara, A., Birgaoanu, M., and Griffiths-Jones, S. (2019). miRBase: from

microRNA sequences to function. Nucleic Acids Res *47*, D155-D162.

Lea, R., Bonev, B., Dubaissi, E., Vize, P.D., and Papalopulu, N. (2012). Multicolor

fluorescent in situ mRNA hybridization (FISH) on whole mounts and sections.

Methods Mol Biol *917*, 431-444.

Lyons, D.A., Guy, A.T., and Clarke, J.D. (2003). Monitoring neural progenitor fate through multiple rounds of division in an intact vertebrate brain. Development *130*, 3427-3436.

Marzi, M.J., Ghini, F., Cerruti, B., de Pretis, S., Bonetti, P., Giacomelli, C., Gorski, M.M., Kress, T., Pelizzola, M., Muller, H*., et al.* (2016). Degradation dynamics of microRNAs revealed by a novel pulse-chase approach. Genome Res *26*, 554-565.

Nepal, C., Coolen, M., Hadzhiev, Y., Cussigh, D., Mydel, P., Steen, V.M., Carninci, P., Andersen, J.B., Bally-Cuif, L., Muller, F*., et al.* (2016). Transcriptional, post-transcriptional and chromatin-associated regulation of pri-miRNAs, pre-miRNAs and moRNAs. Nucleic Acids Res *44*, 3070-3081.

Novak, B., and Tyson, J.J. (2008). Design principles of biochemical oscillators. Nature reviews Molecular cell biology *9*, 981-991.

R Core, T. (2019). R: A language and environment for statistical computing.

Rodriguez-Otero, P., Roman-Gomez, J., Vilas-Zornoza, A., Jose-Eneriz, E.S., Martin-Palanco, V., Rifon, J., Torres, A., Calasanz, M.J., Agirre, X., and Prosper, F. (2011). Deregulation of FGFR1 and CDK6 oncogenic pathways in acute lymphoblastic leukaemia harbouring epigenetic modifications of the MIR9 family. Br J Haematol *155*, 73-83.

Ruegger, S., and Grosshans, H. (2012). MicroRNA turnover: when, how, and why. Trends Biochem Sci *37*, 436-446.

Sasai, Y., Kageyama, R., Tagawa, Y., Shigemoto, R., and Nakanishi, S. (1992). Two mammalian helix-loop-helix factors structurally related to Drosophila hairy and Enhancer of split. Genes Dev *6*, 2620-2634.

Shibata, M., Kurokawa, D., Nakao, H., Ohmura, T., and Aizawa, S. (2008).

MicroRNA-9 modulates Cajal-Retzius cell differentiation by suppressing Foxg1

expression in mouse medial pallium. J Neurosci *28*, 10415-10421.

Shibata, M., Nakao, H., Kiyonari, H., Abe, T., and Aizawa, S. (2011). MicroRNA-9

regulates neurogenesis in mouse telencephalon by targeting multiple transcription

factors. J Neurosci *31*, 3407-3422.

Shimojo, H., Ohtsuka, T., and Kageyama, R. (2008). Oscillations in notch signaling

regulate maintenance of neural progenitors. Neuron *58*, 52-64.

Soto, X., Biga, V., Kursawe, J., Lea, R., Doostdar, P., Thomas, R., and Papalopulu,

N. (2020). Dynamic properties of noise and Her6 levels are optimized by miR-9,

allowing the decoding of the Her6 oscillator. EMBO J *39*, e103558.

Tambalo, M., Mitter, R., and Wilkinson, D.G. (2020). A single cell transcriptome atlas

of the developing zebrafish hindbrain. Development *147*.

Thisse, C., and Thisse, B. (2008). High-resolution in situ hybridization to whole-

mount zebrafish embryos. Nature protocols *3*, 59-69.

Walker, J.C., and Harland, R.M. (2008). Expression of microRNAs during embryonic

development of Xenopus tropicalis. Gene Expr Patterns *8*, 452-456.

White, R.J., Collins, J.E., Sealy, I.M., Wali, N., Dooley, C.M., Digby, Z., Stemple,

D.L., Murphy, D.N., Billis, K., Hourlier, T*., et al.* (2017). A high-resolution mRNA

expression time course of embryonic development in zebrafish. eLife *6*.

Wickham, H. (2016). ggplot2, Elegant Graphics for Data Analysis (Springer

International Publishing).

Wong, S.F., Agarwal, V., Mansfield, J.H., Denans, N., Schwartz, M.G., Prosser,

H.M., Pourquie, O., Bartel, D.P., Tabin, C.J., and McGlinn, E. (2015). Independent

regulation of vertebral number and vertebral identity by microRNA-196 paralogs.

Proc Natl Acad Sci U S A *112*, E4884-4893.

Yates, A.D., Achuthan, P., Akanni, W., Allen, J., Allen, J., Alvarez-Jarreta, J.,

Amode, M.R., Armean, I.M., Azov, A.G., Bennett, R.*, et al.* (2020). Ensembl 2020.

Nucleic Acids Res *48*, D682-D688.

Zhao, C., Sun, G., Li, S., and Shi, Y. (2009). A feedback regulatory loop involving

microRNA-9 and nuclear receptor TLX in neural stem cell fate determination. Nat

Struct Mol Biol *16*, 365-371.

Zhao, J., Lin, Q., Kim, K.J., Dardashti, F.D., Kim, J., He, F., and Sun, Y. (2015).

Ngn1 inhibits astrogliogenesis through induction of miR-9 during neuronal fate

specification. eLife *4*, e06885.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5