Equal performance but distinct behaviors: *Astatotilapia burtoni* sex differences in

a novel object recognition task and spatial maze

Kelly J. Wallace[1] & Hans A. Hofmann[1]

[1]Department of Integrative Biology, University of Texas, Austin, TX, 78712, USA

Article Type: Research Article

Kelly J. Wallace ORCID: 0000-0002-2361-1213

Hans A. Hofmann ORCID: 0000-0002-3335-330X

Author for correspondence: Kelly J. Wallace

kwallace@utexas.edu

University of Texas at Austin, Department of Integrative Biology

1 University Station C0990

Austin, Texas 78712

Highlights

- To ask if sexual dimorphism in cognition is reduced in a species where both sexes navigate highly fluctuating social environments, we assessed performance and behavior of the cichlid fish *Astatotilapia burtoni* in a novel object recognition task and a spatial maze.
- Both sexes exhibited a preference for the familiar object during a novel object recognition task, with males exhibiting the preference early in the task, whereas females exhibiting the preference late in the task.
- In the spatial task, females reached the learning criterion more often than expected by chance, whereas males did not. Females exhibited significantly longer decision latencies and quicker error correction in the spatial task, suggesting a sex-specific speed-accuracy tradeoff.
- The sexes differ in behaviors related to neophobia, space use, and decision latencies.
- A model selection analysis to predict sex finds that novel object recognition task preference, approach, and interaction duration are the most important terms on average.

Abstract

Sex differences in behavior and cognition can be driven by differential selection pressures from the environment and in the underlying neuromolecular mechanisms of decision-making. The highly social cichlid fish *Astatotilapia burtoni* exhibits dynamic and complex social hierarchies, yet explicit cognitive testing (outside of social contexts) and investigations of sex differences in cognition have yet to be fully explored. Here we assessed male and female *A. burtoni* in two cognitive tasks: a novel object recognition task and a spatial task. We hypothesized that males outperform females in a spatial learning task and exhibit more neophilic/exploratory behavior in across both tasks. In the present study we find that both sexes prefer the familiar object in a novel object recognition task, but the time at which they exhibit this preference differs between the sexes. Females more frequently learned the spatial task, exhibiting longer decision latencies and quicker error correction, suggesting a potential speed-accuracy tradeoff.

Furthermore, the sexes differ in space use in both tasks and in a principal component analysis of the spatial task. A model selection analysis finds that preference, approach, and interaction duration in the novel object recognition task that reach a threshold of importance averaged across all models. This work highlights the need to explicitly test for sex differences in cognition to better understand how individuals navigate dynamic social environments.

Keywords

sex differences, spatial learning, novel object recognition, social behavior, *Astatotilapia burtoni*

Introduction

For humans and non-human animals alike, navigating dynamic social environments is a constant challenge (O'Connell & Hofmann 2011, Taborsky & Oliveira 2012). Individuals must attend to changing signals and cues, remember stimuli and their valence, assess risk, make inferences from often incomplete or ambiguous information, use and update prior associations, and display socially appropriate behaviors in varied contexts including mate choice (Phelps & Ophir 2009, White & Galef 1999, Hebets & Sullivan-Beckers 2010), territorial/dominance aggression (Reichert & Quinn 2017, Bshary & Brown 2014), social foraging (Rapaport & Brown 2008, Thornton & McAuliffe 2006), and nest building (Keagy et al. 2011).

Males and females frequently make different decisions even within the same social environment. Sex differences in cognition, "the mechanisms by which animals acquire, process, store, and act on information from the environment" (Shettleworth 2010) have been described across taxa (rodents: Orsini & Setlow 2017, Dalla & Shors 2009, Rice et al. 2017, songbirds: Titulaer et al. 2012, lizards: Carazo et al. 2014, poeciliid fish: Lucon-Xiccato & Bisazza 2017a,b, Wallace et al. 2020), although males and females do not always differ in this regard (Lucon-Xiccato & Bisazza 2014 and 2016; Healy 1999, Guillette et al. 2009, Etheredge et al. 2018). When present, sex differences in cognition often arise from varying ecological demands and fitness benefits of cognitive abilities

such as sexually divergent pressures related to dispersal, predation, and mate choice

(Watson & Platt 2008). For example, males and females can differ in foraging patterns

(mice: Maille & Schradin 2016, cormorants: Ishikawa & Watanuki 2002, sandperch

Sano 1993) and response to predation (poeciliid fish: Magurran & Seghers 1994,

Magurran & Nowak 1991). Cognition can additionally impact fitness via reproductive

performance directly, such as sexual display quality (e.g. aspects of song quality in

songbirds), (Smith et al 2015, Boogert et al. 2011, Farrell et al. 2011) mate preference,

(Shohet & Watt 2009, Branch et al. 2019) or responses to sexual conflict (Cummings

2018, Buechel et al. 2016). Furthermore, behavior and decision-making can vary along

many axes including time (seasonality: Clayton & Krebs 1995, Yaskin 2011), social

status (Stowe et al. 2007, Layton & Fulton 2014), satiety (Caraco 1981), reproductive

status (Lynch et al 2005), and social context (Rosati & Hare 2012).

Burton's mouthbrooder cichlid, *Astatotilapia burtoni,* is a highly social east African

cichlid fish that provides a unique system in which to explore sex differences in

cognition in a temporally dynamic social system (Hofmann & Fernald 2001, Maruska &

Fernald 2018). In this species, both sexes have been shown to navigate socio-

ecological demands that fluctuate across time: males bidirectionally shift between a

dominant phenotype (colorful, aggressive, territorial) and a subordinate phenotype (drab

coloration, submissive, non-territorial) (Fernald & Hirata 1977, Burmeister et al 2005,

Maruska & Fernald 2010, Huffman et al 2012), and females can also form complex

social hierarchies and display remarkable behavioral plasticity across reproductive

states (Fernald & Hirata 1977, Kidd et al. 2013a, White et al. 1996, Renn et al. 2012,

Kidd et al. 2013 a,b). Successfully navigating these complex and ever-changing social

surroundings requires cognitive abilities such as transitive inference and cognitive

flexibility, (Fernald 2014, Alcazar et al. 2014, Grosenick et al. 2007, Desjardins. et al

2012, Weitekamp et al. 2017, Rodriguez-Santiago et al 2020). Furthermore, this species

has emerged as a model system for understanding the overlapping neurobiological

mechanisms of social behavior and decision-making (Hofmann 2003, O'Connell &

Hofmann 2012, Maruska & Fernald 2018). While cognition has been examined in

naturalistic communities of *A. burtoni* (see Wood et al. 2011, Rodriguez-Santiago et al

2020), studies that assess cognitive performance independent of social context and

across the sexes are lacking. In the present study we compare the sexes in two

cognitive tasks to examine whether and how the social environment might influence

cognition *outside* of the fast-paced social lives of this species. Assessing cognition

outside of social context is critical for understanding how socioecological pressures

influence decision-making and behavior more generally independent of ongoing social

interactions.

In the present study, we investigated whether *A. burtoni* males and females differed in

cognition by assessing individuals in a novel object recognition task (NOR) and a spatial

task (SPA). Both tasks require learning - a change in cognitive state or behavior

resulting from prior experience (Shettleworth 2010, Staddon 1983, Domjan 2015)- and

test abilities that are relevant to *A. burtoni*'s natural history, as males typically establish

territories under specific objects (such as plant debris or rocks), to which they reliably

return after extensive excursions in search of females found in shoals on the periphery

of the lek (Fernald & Hirata 1977). We hypothesized that if we were to find sex

differences in cognition in this species they would manifest as greater male

performance in the spatial task. We employed here a spatial task, where individuals

were tasked with learning a simple route (behind a left barrier or right barrier to reach a

social reward). This paradigm is appropriate given that in the natural environment both

sexes are often found in shoals, while dominant males remember the location of their

territories. Sex differences in spatial learning have been previously described in fish,

with most studies reporting males outperforming females (Carazo et al. 2014, Lucon-

Xiccato & Bisazza 2017a,b, Fabre et al. 2014, Saucier et al. 2008, Jonasson 2005, but

see Healy et al. 1999 which finds no sex difference and Jones et al. 2003 which

emphasizes that home range may be a better predictor of spatial performance than

sex). Importantly, behavior exhibited during a task can vary by sex independently of

cognitive performance (Burns & Rodd 2008, Mamuneas et al. 2015; Lucon-Xiccato &

Bisazza 2016; Etheredge et al. 2018; Titulaer 2012, Dougherty & Guillette 2018), for

instance due to where males are typically identified as the "bolder" or "more exploratory"

sex, often explained by sex-specific variation in predation pressures and dispersal

behavior as well as life history tradeoffs (Lucon-Xiccato & Dadda 2016, Han et al. 2015,

Harris et al. 2010, Overli et al. 2006, Gagnon et al. 2016, but see Hughes 1968 &

Swanson 1966). In concordance with this literature, we hypothesized that across both

tasks males would display more neophilic and exploratory behaviors than females.

Lastly, we assessed whether behavior and cognition in the two tasks was correlated to

the other (Lucon-Xiccato & Bisazza 2017b) via multidimensional analyses. We

hypothesized that, since both memory and spatial learning are needed to successfully

navigate the *A. burtoni* social environment, both sexes would successfully learn both

tasks and additionally would exhibit correlated individual performance across tasks.

Materials and Methods

(a) Housing and Initial Community Formation

128 adult *A. burtoni* (64M/64F) were used from a lab population descended from wild-

caught stock. Prior to testing, the fish were maintained in naturalistic communities (114-

liter aquaria), which allowed for individual variation in dominance and reproductive

status. Both prior to and during testing, fish were kept in reverse osmosis water treated

with Seachem® Lake Tanganyika salt and bicarbonate buffer. Lights were on a 12:12

light-dark cycle and fish were fed once daily with Omega One Natural Protein Formula

Cichlid Flakes. All individuals were tagged with colored plastic beads for individual

identification. One week prior to testing, fish were placed as a community of 8M/8F (n=

8 replicate communities) in a circular pool (1m diameter, 25cm depth, lined with a PEVA

curtain liner and Spectrastone White aquarium gravel) containing four halved terra-cotta

pots to be used as territorial shelters (Supplemental Figure 1), and at this time fish were

weighed and measured for standard length (average standard length: males = 4.78cm,

females = 4.76cm). One week after completion of the experiment, individuals were

again weighed and subsequently dissected for gonadal weight to calculate

gonadosomatic index (GSI, i.e., the ratio of gonad mass to body mass), a proxy for

reproductive status.

(b) Novel Object Recognition Task (NOR)

To assess memory, we utilized a novel object recognition task (also referred to as the spontaneous object recognition task). The novel object recognition task, at its simplest, requires an individual to discriminate between a previously encountered object and a novel one. This paradigm is commonly used in the rodent literature to assess memory (Ennaceur & Delacour 1988, Broadbent et al. 2010), but additionally has the capacity to characterize recognition, neophobia, and exploration (Antunes & Biala 2012).

In the novel object recognition task conducted here, individuals were assessed for relative preference (physical proximity) to a familiarized object versus a novel object in a 114 liter acrylic aquarium filled to a depth of 20cm with a central alley way and two object regions (Figure 1A, B). Four objects were used as stimuli in the task: a pink centrifuge tube, a purple falcon tube cap, a blue weight boat (quartered), and a yellow table tennis ball filled with sand (Supplemental Figure 2). Object pairs and their roles (familiar vs novel) were counter-balanced across communities. Prior to testing, the entire community was habituated at once to two identical objects (coined the "familiar object") for one hour (e.g. A + A). We chose this method of habituation for two reasons: First, to maintain daily consistency in the timing of the experimental design; and second, to avoid disruption of the social hierarchy, which frequently occurs when only one individual is removed from the community at a time. Approximately 24 hours later, each fish was individually presented with a familiar object and a novel object at the opposite ends of the aquarium (the side of the familiar object, left or right, was counter-balanced across communities) (e.g. A + B). Previous studies in zebrafish have shown that the delay between habituation and testing can range between 2 and 24 hours without impacts on resulting performance (Lucon-Xiccato & Dadda 2014). During the trial, individuals were acclimated in an opaque PVC tube for one minute then allowed to swim freely in the apparatus for twelve minutes. Following the trial, individuals were kept in a holding area until all community members were tested, then simultaneously returned to the community to prevent social hierarchy destabilization. We recorded fish as they transited between the following zones (Figure 1): the central alley, the interaction area (where the object was located), and the corners associated with each object. For each zone we calculated the proportion of time spent in the zone, the number of entries into the zone, and the time of first entry. Performance was calculated

as the relative preference for the familiar object ("NOR familiar preference"): the relative

proportion of time spent at the familiar object to time spent at the novel object, where a

score of 1 means the entirety of time spent at an object was at the familiar object, a

score of 0.5 means equal preference, and of 0 means the entirety of time spent at an

object was at the novel object.

(c) Spatial Task (SPA)

Spatial learning involves following a route composed of a sequence of stimuli that must

occur in a specific order (Dodson 1988). There has been a long and extensive usage of

spatial tasks ("mazes") to assess spatial learning in fish: see Salas et al. 1996

(goldfish), Lucon-Xiccato & Bisazza 2017a (guppy), Burns & Rodd 2008 (guppy), Jun et

al. 2016 (electric fish) Wood et al. 2011 (cichlid). In the spatial task conducted here,

individuals in a 57 liter acrylic aquarium filled to a depth of 20cm were required to detour

around an opaque barrier in one of two directions (left or right) to reach a social reward

consisting of a female in a clear plexiglass tube behind one side of the barrier (Figure

1C, D, average standard length of female conspecific reward = 4.49cm). In *A. burtoni,*

both sexes are highly social, and the use of a female reward (from a separate stimulus

community and novel to all subjects) for both sexes was employed to avoid potentially

confounding effects of male-male aggression. Behind the other side of the barrier, an

empty plexiglass tube was placed as a control stimulus. Transit between the rewarded

and unrewarded area required an individual to navigate back out in front of the barrier,

thus creating a penalty for errors. For all trials, individuals were initially placed in an

opaque PVC tube for one minute to acclimate to the aquarium, then allowed to swim

freely for twelve minutes. Each individual underwent an initial apparatus habituation trial

(using two control stimuli) the day prior to the reward trials. Following this habituation

trial, individuals were assayed over four rewarded trials over two days, with the reward

stimulus present behind either the left or right barrier, counter-balanced across

individuals. Across these trials the same reward female stimuli were used per individual

(excluding rare experimental limitations). As with the NOR task, following the trial,

individuals were kept in a holding area until all community members were tested then

simultaneously returned to the community. We recorded fish as they transited between

the following zones (Figure 1): the central neutral area, within 2cm of the various walls,

within 2cm of the barrier, the rewarded and unrewarded corners, and the rewarded and

unrewarded platforms (which contained the plexiglass tube). For each zone we

calculated the proportion of time spent in the zone, the number of entries into the zone,

and the time of first entry. To assess performance, we established a learning criterion in

which successful learners first approached the rewarded stimulus in all rewarded trials

after the first trial (i.e. trials 2-4). Relative preference for the rewarded stimuli ("SPA

reward preference") was calculated as the relative proportion of time spent at the

rewarded stimulus to time spent at the unrewarded stimulus, where a score of 1 means

the entirety of time spent near a stimulus was at the rewarded stimulus, a score of 0.5

means equal preference, and of 0 means the entirety of time spent at an object was at

the unrewarded stimulus.

(d) Video Scoring

Videos were recorded overhead of the experimental apparat via the Alibi Security

Camera system and scored by three human observers (who had not conducted the

behavioral assays) in VLC media player using CowLog video scoring software version

3.0.2 (Hänninen & Pastell 2009), a software which has been validated to be highly

consistent across multiple observers (Wallace et al. 2020). Focal fish location was

recorded though the duration of the test trial of the novel object recognition task and the

fourth (test) trial of the spatial task. Additionally, for trials one through three of the spatial

task the first approach (rewarded or unrewarded) was recorded.

(e) Statistics

Data analysis and visualization were performed in R (R Core Team 2013) version 3.6.1

(2019). We conducted an unpaired t-test or unpaired two-sided Wilcoxon Rank Sum test

with continuity correction for continuous data split into 2 categories, or a One-way

ANOVA or Kruskal-Wallis test for continuous data split into >2 categories (determined

by a Shapiro-Wilk normality test). For entirely categorical data we conducted Chi-

squared tests. We conducted standard multiple linear regressions to determine

correlations and effect sizes of relationships between continuous independent variables

using the formula lm(x1 ~ x2) using the R function "lm()". To identify interaction effects

of sex, we conducted linear models with interaction terms using the formula lm(x1 ~ x2 *

x3). Effect sizes were calculated as either Cohen's D using the R function "cohen.d()"in

the R package "effsize" (Torchiano 2020) for categorical data or as the Pearson

correlation coefficient using the R function "cor()" for continuous data. We conducted

principal component analyses (PCA) on z-scored data via the R function "prcomp".

Hierarchical clustering analyses were conducted using the R package "pvclust" (Suzuki

et al 2019) with distance metric: "correlation", linkage metric: "average", and bootstrap

values set to 100. Logistic regression models were constructed and selected using the

R package "glmulti" (Calcagno 2020). Variables included in data analysis (including in

the principal component analyses) were first assessed via a covariance matrix: any two

variables with a Pearson's correlation coefficient above 0.8 were considered redundant,

thus for subsequent analysis the authors used the variable more commonly used in

previous literature. Raw data and analysis code can be found at

https://github.com/kellyjwallace/Wallace_Hofmann_2020_sex_differences, or via the

Texas Data Repository (https://data.tdl.org/).

Results

(a) Novel Object Recognition Task (NOR)

Before quantifying performance (first object choice and relative preference for the

familiar object) on the NOR task, we first assessed if any objects were preferred

independent of familiarity. No object was significantly preferred or avoided (average

preferences: cap 55%, weigh boat 45%, ball 43%, tube 57%, Kruskal-Wallis test of

preference by object: $p = 0.080$, df = 3, $\eta^2 = 0.041$, Wilcoxon test of preference between

the two pairs: $W_{41,41} = 389$, $p = 0.229$ and $W_{55,55} = 497$, $p = 0.054$, respectively).

However, because the ANOVA across object and the ball/tube pair showed a near

significant bias, we additionally tested whether individuals' first choice (familiar or novel)

differed by the pair of objects they received and found no significant difference (chi^2

test: $p = 0.108$, $n = 97$, $X^2 = 2.577$, df = 1).

We then asked whether males and females differed in task performance. We quantified

the proportion of individuals that first approached the familiar object in each sex and

found that neither sex first approached the familiar object more than would be expected

by chance: 63 individuals (38M/25F) first approached the familiar object, and 47

individuals (20M/27F) first approached the novel object. Consistent with this result, we

also found that a continuous measure of performance across the length of the trial, the

"NOR familiar preference," did not differ by sex (average NOR familiar preference

across both sexes = 0.52, Wilcoxon test: $p = 0.386$, $W_{64,64} = 1268$, Cohen's D = 0.188)

and did not diverge from a random 50% expectation in either sex. This preference score

did not differ by round (rounds differed in the object pair presented) (Kruskal-Wallis test:

$p = 0.174$, df = 7, $\eta^2 = 0.038$), and was not correlated with GSI in either sex (female

linear regression: $R^2 = -0.021$, $F_{37} = 6.321$, $p = 0.647$, male linear regression: $R^2 = -$

0.020, $F_{21} = 0.565$, $p = 0.461$).

Given that neither sex displayed a significant preference when averaged across the

entire task (contrary to our hypothesis that both sexes would learn the task), we further

asked whether a preference for either the familiar or novel object could be observed at

any time during the task. We initially observed that males exhibited a significant

preference for the familiar object in the first minute of the task (Wilcoxon test: $p = 0.040$,

V = 742, n = 49, Figure 2A), and upon a more in-depth analysis found that preference

was significantly correlated with time for both males (linear regression: $R^2 = 0.372$, $F_8 =$

6.321, $p = 0.036$) and females (linear regression: $R^2 = 0.570$, $F_8 = 12.93$, $p = 0.007$),

with a significant interaction effect of sex: male's familiar object preference decreased

across time whereas female's increased (linear regression: $R^2 = 0.575$, $F_{16} = 9.55$, $p =$

0.0009) (Figure 2B).

To assess whether males were the more neophilic sex, we compared latencies to

approach the objects and the proportion of time spent with objects: latency to approach

either object did not differ by sex (Wilcoxon test: $p = 0.463$, $W_{64,64} = 1275$, Cohen's D =

0.198), nor did the proportion of time spent interacting with either object (Wilcoxon test:

$p = 0.180$, $W_{64,64} = 987$, Cohen's D = 0.331) though a large proportion of time was spent

in the corners near the objects ("association"), significantly more so in females with a

small effect (average of females = 85%, average of males = 78%, Wilcoxon test: p =

0.022, $W_{64,64}$ = 1490, Cohen's D = 0.482) (Figure 2C). Activity (total entries) during the

first minute of the task did not significantly differ between the sexes (Wilcoxon test: p =

0.539, $W_{49,45}$ = 1021.5, Cohen's D = 0.051). Visualizations of focal location across time

in the NOR task are available in (Supplemental Figure 3).

We then conducted a multivariate analysis of variables measured in the NOR task

(PCA) to further assess relationships between performance and other behaviors

observed during the task. The analysis identified 9 PC axes, three of which explained

>10% each (and together 74%) of the total variance in the dataset (Supplemental Figure

6, Supplemental Table 1). None of these PC's differed significantly between the sexes.

To identify clusters of behaviors that shared similar loading patterns, we performed a

hierarchical clustering analysis of behavior using the PCA eigenvalues. The analysis

yielded one highly supported cluster (AU > = 95%): NOR non-engagement (a

neophobia metric) and NOR familiar preference.

(b) Spatial Task (SPA)

To first confirm that no cues (e.g. olfactory) from the reward fish could be detected by

the focal fish, we compared the proportion of individuals who first chose the rewarded

side during the first trial to a random expectation, and found that individuals first

approached the rewarded and unrewarded sides randomly (chi^2 test: p = 0.313, n =

111, X^2 = 1.0182, df = 1). To test our hypothesis that the sexes would not differ in

spatial learning or, alternatively, that males would outperform females due to

territoriality-related cognitive demands, we assessed performance on the final trial of the

task (trial 4) and surprisingly found that 31 of 51 females (61%) first chose the rewarded

side, which was significantly more than the 25 of 59 males (42%) who first chose the

rewarded side (chi^2 test: p = 0.043, n = 111, X^2 = 4.064, df = 1, Supplemental Figure 4).

We then assessed performance based on a criterion of learning where we classified

learners as those who first approached the rewarded stimulus in all rewarded trials after

the first trial (trials 2-4). The probability for this to occur by random choice is 12.5%.

Interestingly, 7 of 58 males (12.1%) reached this criterion (suggesting random choice),

whereas 12 of 51 females (23.5%) reach this criterion, representing a significant

cumulative binomial probability in which the females learn the task more than would be

expected by chance (p = 0.021, Figure 3A). However, this is not a significant sex

difference in performance, as the proportion of males that reached the learning criterion

did not differ from proportion of females that reached the learning criterion (chi^2 test: p =

0.186, n = 109, X^2 = 1.744, df = 1).

Next, we asked whether males were the more neophilic sex and whether behavior was

related to performance in the task. To this end, we assessed several behavioral and

performance metrics in the fourth trial of the assay. Interestingly, we detect a small

effect where *males* spent significantly more time spent near the walls of the aquarium

during the task (thigmotaxis), which is commonly considered an anxiety behavior

(Maximino et al. 2010) (Wilcoxon test: p = 0.045, $W_{53,50}$= 1023, Cohen's D = 0.435). We

found a small effect where females exhibited a significantly higher SPA reward

preference across the duration of the task (female average = 0.6, male average = 0.4,

Wilcoxon test: p = 0.032, $W_{64,64}$ = 1346, Cohen's D = 0.447) (Figure 3B). This

preference was not correlated with GSI in either sex (female linear regression: R^2 = -

0.006, F_{38} = 0.770, p = 0.386, male linear regression: R^2 = -0.026, F_{24} = 6.321, p =

0.549). Additionally, females took significantly *more* time to decide than males with a

moderate effect size (Wilcoxon test: p = 0.0028, $W_{64,64}$= 1381, Cohen's D = 0.538)

(Figure 3C). Also, their latency to solve the task (enter the rewarded area) was highly

correlated with this latency to decide, a relationship that was not present in males (linear

regression: R^2 = 0.851, F_{60} = 121.3, p = 1.296x10^{-14}) (Figure 3D), suggesting that the

sexes might differ in error correction. To test this idea, we subsetted the results by

individuals who first approached the unrewarded side and subsequently entered the

rewarded side and measured the time between the first unrewarded entry and the

subsequent rewarded entry. We found that, indeed, females correct these first approach

"errors" significantly faster than males with a very large effect (female average = 101

seconds, male average = 309 seconds, Wilcoxon test: p = 0.029, $W_{7,8}$ = 9, n = 15,

Cohen's D = 1.193) (Figure 3E). Latency to solve the task was not correlated with GSI

in either sex (female linear regression: $R^2 = 0.057$, $F_{33} = 3.038$, p = 0.091, male linear

regression: $R^2 = -0.10$, $F_{21} = 0.0002$, p = 0.990). Visualizations of focal location across

time in the SPA task are available in (Supplemental Figure 5).

As with the NOR task, we conducted a principal component analysis of variables

measured in the spatial task to understand the relationships between the behaviors

observed during the task. Of the 10 PC axes identified, we explored further the four PCs

that explained >10% each (and together 85%) of the total variance in the dataset

(Supplemental Figure 7A, Supplemental Table 2). Interaction, association, decision

latency, and interaction latency loaded most strongly onto PC2 (20% of the variance)

and significantly differed by sex with a large effect (Wilcoxon test with FDR correction, p

=0.008, $W_{26,24} = 481$, Cohen's D = 0.884) (Supplemental Figure 7B). None of the other

PCs varied across sexes, and in a hierarchical clustering analysis of the eigenvalue

patterns did not yield any well supported clusters (AU > = 95%).

(d) Relationships across the tasks

Completing the NOR task requires learning and memory, and both abilities are likely

important for successfully navigating the cognitive demands of *A. burtoni* social life. We

therefore asked whether performance across the two tasks was correlated. When first

simply using the categorical metric of "first choice" (rewarded or unrewarded in SPA,

familiar or novel in NOR) the two tasks are not correlated. However, performance in the

two tasks can be compared appropriately only for those individuals who entered both

the familiar and novel object recognition task areas *and* the rewarded and unrewarded

areas of the spatial task. 13 males and 7 females met this criterion. When we then

correlated the continuous preference metrics across the tasks (relative preference for

the familiar object in the novel object recognition task and relative preference for the

rewarded stimuli in the spatial task), we found that they were indeed significantly

correlated in a sex-invariant manner (linear regression: $R^2 = 0.229$, $F_{18} = 5.359$, p =

0.032) (Figure 4A). Because we had previously discovered that decision latencies

differed by sex in the SPA tasks (Figure 3C), we next asked whether decision latencies

related to choices across tasks, and observed a modest effect where individuals who

first approached the novel object in NOR task showed longer decision latencies in SPA

(Wilcoxon test: p = 0.033, $W_{58,39}$= 427, Cohen's D = 0.333), and conversely, individuals

who first approach the rewarded stimulus in SPA exhibit longer decision latencies in

NOR (Wilcoxon test: p = 0.010, $W_{53,50}$ = 1062, Cohen's D = 0.029) (Figure 4B,C).

To better understand the relationships between variables across both tasks, we

conducted a principal component analysis that included the variables from both the SPA

and NOR tasks (Figure 5, Supplemental Table 3), and identified four principal

components (of ten total) that explained at least 10% each (and together 61%) of the

total variance. The first two principal components were largely defined by task: PC1

(accounting for 24% of the variance) loaded most strongly with measures of activity and

reward preference in the spatial task, whereas PC2 (accounting for 16% of the

variance) loaded primarily by object preference and approaches, though it also included

latency variables across both tasks. PC3 (11%) loaded most strongly by variables

associated with stimulus interaction across both tasks, and PC4 (10%) loaded primarily

by a single variable: SPA reward latency. No PC axes in this combined analysis differed

significantly by sex. When clustering behaviors by their eigenvalue patterns in this PCA,

we identified five highly supported clusters (AU > = 95%), some of which were with a

task and some which spanned across tasks (Figure 5, highlighted in grey boxes).

Finally, to identify which variables across both tasks were best able to distinguish males

and females, we conducted three logistic regression model selection analyses to predict

sex via an exhaustive screening of all possible models and selecting based on AICc:

one with variables from the NOR task, one with variables from the SPA task, and a

combined model. Using variables from the NOR task, the selected model (of 511

possible models, McFadden's Pseudo-R^2 value of 0.032) included only NOR novel

object interaction bout duration (Supplemental Table 4). Using variables from only the

SPA task, the selected model (of 1023 possible models, McFadden's Pseudo-R^2 value

of 0.213) included the SPA preference measure and % association (Supplemental

Table 5). For each of analysis we additionally conducted a model-averaged importance

of terms analysis. In a model-averaged importance of terms analysis, the importance of

a term is determined by the number of models the term appears in and their weights (a

metric of the model's likelihood to be the "best" model). Neither of these task-specific

models identified a term that reached the importance threshold of 0.8 (Supplemental

Tables 4,5). Using variables from both tasks, the selected model (of 524,287 possible

models, McFadden's Pseudo-R^2 value of 1.00) included the following variables from

both tasks: NOR novel object interaction bout duration, NOR familiar preference, NOR

novel object approaches, NOR association, NOR interaction, NOR object approach

latency, SPA decision latency, SPA % association, SPA reward latency, NOR non-

engagement, SPA reward preference, SPA reward preference following solution, and

SPA thigmotaxis (Table 1, bolded). The terms in a model-averaged importance of

terms analysis that reached the importance threshold were three NOR variables: NOR

novel object interaction bout duration (0.994), NOR familiar preference (0.938), and

NOR novel object approaches (0.927). (Table 1, highlighted with an asterisk)

<u>Discussion</u>

The present study investigated sex differences in cognition (outside of social contexts)

in a species in which both sexes navigate temporally dynamic socio-ecological

pressures would show less sex differences in cognition. We used Burton's

mouthbrooder cichlid, *Astatotilapia burtoni,* a species well-characterized for its fast-

paced and complex social system (Hofmann & Fernald 2001). We assessed male and

female *A. burtoni* (across 8 replicate communities, a total of 128 individuals) in two

cognitive tasks: a novel object recognition task and a spatial task. In the novel object

recognition task, both sexes exhibited a preference for the familiar object but at different

times of the task, and males spent less time associating with the objects than females.

In the spatial task, females reached a learning criterion above chance levels and

exhibited stronger preferences for the reward stimulus. Females exhibited longer

decision latencies and quicker error correction speeds. These differences were also

captured in a principal component analysis of the spatial task, which identified a PC axis

that significantly varied by sex and loaded most strongly by stimulus interaction and

latency. Broadening our analysis to look at relationships across both tasks, we found

that in a sex-invariant manner preference scores across the two tasks significantly

correlated to each other (preference for the familiar object in the novel object

recognition task, and preference for the rewarded stimulus in the spatial task), and

decision latencies predicted first choice behaviors across tasks. The first axis of a

principal component analysis using variables from both tasks was primarily loaded by

SPA preference and activity. A hierarchical clustering analysis of eigenvalue patterns

validated behavioral relationships both within and across tasks, and additionally

identified a potential neophobia relationship across the tasks. Finally, using a logistic

regression model selection approach, we found that the behavioral variables that best

predicted sex spanned both tasks. However, when assessing the importance of

variables across all models, only NOR measures (novel object interaction bout duration,

familiar preference, and number of novel object approaches) reached an importance

threshold.

(a) Sex differences in the Novel Object Recognition Task

In the novel object recognition task, both sexes exhibited a significant preference for the

familiar object, confirming our hypothesis that both sexes would learn the task. While

rodents (of both sexes) typically prefer the *novel* object, studies in fish have yielded

conflicting results (May et al. 2016, guppies: Miletto Petrazzini et al. 2012, Lucon-

Xiccato & Dadda 2016). Where we did identify sex differences was in the *timing* of the

NOR preference for the familiar object: males exhibited the familiar preference early on

in the task, whereas females exhibited the familiar preference later in the task. Studies

in fish have found cases where preferences are only exhibited during for the first few

minutes of the task (Lucon-Xiccato & Dadda 2016, Miletto Petrazzini et al. 2012). We do

not believe this sex difference in the timing of the familiar preference in our experiment

to be a reflection of motivation or initial anxiety during the task, as latency to approach

either object did not differ by sex. Furthermore, both sexes are equally active in the first

minute of the task (our closest proxy to freezing behavior, Blaser et al. 2010). Instead,

we suggest that this timing difference reflects variation in space use between the sexes.

In a NOR experiment conducted by Lucon-Xiccato & Dadda (2016), zebrafish males

exhibited similar object preferences to females, but more quickly approached the novel

object. The authors emphasize, as we do, that the differences found likely do not reflect

differences in memory but rather space use (as zebrafish and *A. burtoni* are both small,

social fish that may respond similarly to novel isolated environments).

One additional consideration that may explain the sex difference in NOR familiar

preference is that, in our experimental design, the animals were familiarized to the task

(all individuals from a given community were familiarized together). A similar study in

juvenile guppies also used a NOR task while minimizing social isolation (i.e., habituation

to the apparatus occurred in pairs) and found a preference for the familiar object during

the subsequent test (Miletto Petrazzini et al. 2012). In ravens, dominance status

influenced approach latencies in a novel object task, and this relationship between

dominance status and approach varied by social context (e.g. same-sex vs mixed-sex

pair) (Stowe et al. 2006). Dominance status has been shown to relate to exploratory

behavior (Dingemanse & de Goede 2004, Stapley & Keogh 2004, Carazo et al. 2014)

and cognitive performance across multiple tasks and taxa (pheasants, Madden et al.

2018, starlings, Boogert et al. 2006, dogs, Pongracz et al. 2008). In *A. burtoni*, males

vary in dominance status, with dominant males aggressively establishing and defending

territories (Fernald & Hirata 1977). Because the habituation phase of the NOR task was

conducted in a social setting, the preferences exhibited during the test may be the result

of social associations (e.g. males more readily seeking the familiar object as a territory).

While we failed to discover a relationship between GSI (as a proxy for reproductive

state), further experiments assessing within-sex variation in *A. burtoni* and a possible

role of dominance and reproductive status are needed to identify potential social factors

driving NOR preferences.

(b) Sex differences in the Spatial Task

Across both sexes only a small proportion of individuals successfully reached the

learning criterion in the spatial task. This is not unexpected, given that our experimental

paradigm has few trials and a strict learning criterion ("learners" had to learn after only

one training trial and not make a mistake in the following three trials). A similar design

has been used in another cichlid species, the Nile tilapia *Oreochromis niloticus*

(Lombardi Brandão et al. 2015). However, this study also used a T-shaped barrier in

which a focal individual had to navigate around by making either a left turn or right turn

to reach a reward and after 30 trials (2x/day) with a criterion of 9/10 correct trials in a

row 50% of their focal individual successfully learned the task. In contrast, our design

lacked landmark cues and used a much shorter training procedure. It is thus not

surprising that only a small proportion of individuals reached the learning criterion.

It has been suggested that in territorial species competition for mates drives increased

spatial cognitive performance (Heske & Ostfeld 1990), and several studies have indeed

shown that the more territorial sex (which is often male) exhibits increased spatial

cognition (Gaulin & Fitzgerald 1989, Carazo et al. 2014, Rice et al. 2017, Carazo et al.

2014, Lucon-Xiccato & Bisazza 2017, Fabre et al. 2014, Saucier et al. 2008, Jonasson

2005, but see Jones et al. 2003, Healy et al. 1999, Lucon-Xiccato & Bisazza 2017a).

We therefore hypothesized that because some *A. burtoni* males defend territories (and

territory defense is the only scenario in which members of this species exhibit clearly

delineated home ranges), males would exhibit greater spatial learning performance.

Contrary to our expectations, we found that in the spatial task it was *females* who

reached the learning criterion more than would be expected by chance and thus

successfully learned the task, whereas males did not. However, males approached

either stimulus more quickly than females, suggesting that the sex difference in spatial

learning performance was not due to a difference in motivation. Despite females

reaching the learning criterion more than expected by chance, the sexes did not

significantly differ in performance when compared to each other. Female *A burtoni* must

also attend to spatial locales for spawning in male territories, and furthermore females

are capable of exhibiting male-typical behavior (including territory defense) in the

absence of males (Renn et al. 2012), suggesting that despite rarely exhibiting territory

defense under normal conditions the cognitive abilities needed for territory defense are

present in both sexes (White et al. 1996). Further research in *A. burtoni* females on

foraging ecology, predation pressure, risk assessment, and shoaling behavior is

necessary to validate these explanations. And even though reproductive status (GSI)

was not significantly correlated with performance in either sex in the present study,

future research will need to examine within-sex variation in cognition in this species.

In the PCA of the spatial task, PC2 (explaining 20% of the variance) significantly differed by sex. This PC2 loaded most strongly by variables of stimuli interaction, association, and latencies. Additionally, females exhibited longer latencies to decide in the spatial task with medium effect and more frequently reached the learning criterion. Taken together these results suggest a possible speed-accuracy tradeoff, such that females are the slower but more accurate sex in this task. Speed accuracy tradeoffs have been previously identified in the literature (chickadees: Guillette et al. 2011, voles: Mazza et al. 2018, see Chittka et al. 2009 for review). This interpretation is further supported by the finding that females exhibit their NOR familiar preference later in the task, suggesting that females may be the "slower" sex in the NOR as well. Additionally, females corrected errors much faster than males, suggesting higher accuracy as well as increased cognitive flexibility. Cognitive flexibility is the ability to modify behavior in response to variations in consequences or context of the environment (Bond et al. 2007; Easton 2005). While we did not specifically assess cognitive flexibility in this task, our observations of quicker female error correction (a term used to assess cognitive flexibility in other tasks, see Mazza et al. 2018) complements previous findings that female guppies have been shown to outperform males in tests of cognitive flexibility (Miletto Petrazzini et al. 2017, Lucon-Xiccato & Bisazza 2017). Given the large effect size of the sex difference in error correction found in this study, further experiments in this species quantifying sex differences in cognitive flexibility are likely to yield fruitful insights.

(c) Relationships across tasks

A growing body of literature has demonstrated a high correlation in performance across cognitive tasks (Chiappe & McDonald 2005, Kolata et al. 2008, Banerjee et al. 2009). At this current state of research on sex differences in (animal) cognition, relatively few studies have specifically investigated sex differences across multiple cognitive tasks. Those that do often find sex differences in specific tasks rather than general greater performance by one sex across all tasks (e.g. Dalla & Shors 2009, Lucon-Xiccato & Bisazza 2017a, Wallace et al. 2020, Miletto Petrazzini et al. 2017; Lucon-Xiccato &

Bisazza 2014; Brust et al. 2013; Carazo et al. 2014; Laland & Reader 1999). Because

both memory and spatial learning are likely needed by *A. burtoni* males and females to

successfully navigate the dynamic and fast-paced social environment typical for this

species, we predicted performance would be correlated across tasks for both sexes. In

a PCA of both tasks, SPA variables largely loaded on PC1 (explaining 24% of variance)

and NOR variables load on PC2, suggesting that these tasks are not strongly

correlated. However, preference for the familiar object in the NOR task correlates

significantly preference for the reward stimulus in the SPA task at an individual level.

Furthermore, when clustering variables by their eigenvalue patterns in this PCA, a

majority of clusters contain variables from both tasks. NOR novel object approaches

clustered with SPA interaction, both potential measures reflecting boldness/neophilia.

SPA thigmotaxis (wall-hugging) clustered with SPA activity (number of entries during

the task), which is unsurprising given both these behaviors reflect anxiety-like

behaviors. NOR familiar preference clustered with the number of familiar object

approaches, consistent with what we would expect. The two SPA latencies (decision

and interaction) clustered with NOR non-engagement, reflecting another possible

boldness/anxiety/neophobia cluster. Lastly, NOR novel object interaction bout duration,

SPA decision (time spent in both the association and interaction areas), and SPA

preference (both over and preference following solution) clustered together. It is

important to note that these hierarchical clusters of eigenvalues not only help identify

potential dimensions where the sexes might differ; more generally, they can validate the

interpretation of our behavioral measurements. For example, the clustering of SPA

thigmotaxis and activity confirm our expectation that both these behaviors relate to

anxiety (Maximino et al 2010). Additionally, these clusters elucidate some behavioral

patterns that may not be as expected: the clustering of NOR novel object approaches

and SPA interaction suggests that close contact with the conspecific reward in the SPA

should perhaps not be considered a measure of "social affiliation," rather it may

additionally relate to boldness.

Finally, we hypothesized that the spatial task would capture greater sexual dimorphism

in cognition than the novel object recognition task, and indeed a PCA of variables in the

SPA task identified an axis (PC2) that significantly differs by sex, whereas no axis in the

NOR PCA differed significantly. Furthermore, the only sex differences with large effect

sizes resulted from the SPA task, suggesting that the sexes diverge in this task more

than the NOR task. But counter to this hypothesis, only variables in the NOR task (novel

object bout duration, familiar preference, and number of novel object approaches)

reached an average importance threshold across all models. These three variables are

all related to exploration of a novel object, indicating that variation in neophobia may be

at the root of the observed sex differences in *A. burtoni*. Additional experiments that

directly quantify exploratory behavior, stress response, and neophobia in this species

will help disentangle these behavioral factors from cognitive performance and will

facilitate improved cognitive assay design in the future.

(d) Conclusion

In the present study, we asked to which extent the sexes differ in cognition in a species

that lives in a highly dynamic social environment. This question reflects a larger

theoretical framework on the reduction or exaggeration of sex differences depending on

the social environment. This framework has been successfully applied as it relates to

mating system. In voles, monogamous species show a lower degree of sexual

dimorphism in morphology, behavior, and cognition relative to their non-monogamous

congeners (Kleiman 1977, Jacobs 1995, Heske & Ostfeld 1990, Jacobs et al. 1990,

Gaulin & Fitzgerald 1989). And in two species of polygynandrous fish (the shanny and

the Azorean rock-pool blenny), females have larger home ranges and a larger

dorsolateral telencephalon (hippocampus) than males, though this has yet to be

correlated to their learning and memory abilities (Costa et al. 2011). In the present study

we hypothesized that in more dynamic or fluctuating social environments we would find

fewer sex differences in cognition performance, and indeed found no significant

differences in *performance* in either task (though the greater female performance on the

spatial task warrants additional work). We did, however, identify several behavioral

differences between the sexes primarily related to neophobia, space use, and action

latencies. Without a rigorous, phylogenetically controlled species comparison and

without a more detailed understanding of the variation in ecologically relevant behaviors

(e.g. shoaling, foraging, anti-predation behavior) between the sexes in *A. burtoni,* we

cannot definitively know whether the sex differences we have observed here are greater

or less than "expected", but we contribute to this question by uncovering previously

undescribed differences in cognition and behavior across the sexes. Our findings were

facilitated by an experimental design employing two distinct cognitive tasks followed by

multivariate analyses to describe the complex relationships observed in the data. We

argue that pairing a more "reductionist" design of discrete suites of cognitive tasks with

a strong foundation of more "naturalistic" observations of behavior is a powerful

approach to understanding the relationship between the social environment and

cognition. Future research across suites of cognitive tasks specifically designed to test

variation within each sex and between other social phenotypes (e.g. mating systems,

reproductive status, dominance status) will allow us to further relate cognitive

performance to ecologically-relevant behaviors. Ultimately, these research endeavors

will allow us to better understand the evolution of cognition and decision-making in a

dynamic social world.

Data Accessibility

Raw data (Microsoft excel) as well as analysis and visualization R code can be found at

https://github.com/kellyjwallace/Wallace_Hofmann_2020_sex_differences, via the

Texas Data Repository (https://data.tdl.org/), or via request to the corresponding author.

Acknowledgements

The authors would like to thank Kavyaa Choudhary, Layla Kutty, Don Le, Matthew Lee,

and Karleen Wu for assistance in data collection, and to Mike Ryan, Tessa Solomon-

Lane, Felicity Muth, and all members of the Hofmann Lab for discussion and

assistance.

Funding

This work was supported by the National Science Foundation (NSF) Bio/computational

Evolution in Action Consortium (BEACON) Center for the Study of Evolution in Action

and an NSF Grant IOS1354942 (to HAH), a Ford Foundation Predoctoral Fellowship

(National Academies of Sciences, Engineering, & Medicine), a UT Austin Graduate

School Continuing Fellowship, The Zoology Scholarship Endowment for Excellence

(Graduate School at the University of Texas at UT Austin), and a Department of

Integrative Biology Doctoral Dissertation Improvement grant (to KJW).

Competing interests

700 The authors have no competing interests.

Ethical approval

702 The authors certify that this work followed ethical treatment of animals outlined in their

703 IACUC protocol (AUP-2018-00236).

Author's Contributions

705 HAH and KJW conceived of the study. KJW designed and constructed the experimental

706 apparatus and data collection procedures, collected behavioral data, and performed

707 statistical analyses. HAH and KJW interpreted the results. KJW wrote the initial draft of

708 the manuscript, and HAH provided feedback and comments during manuscript writing.

709 All authors give final approval for publication.

Figure Legends

Figure 1. Experimental Designs. The novel object recognition task (A,B) consisted of a 1-hour habituation (community) (A) then a 12-minute test (individual) ~ 24 hours later (B). The spatial task (C,D) consisted of a 1 trial unrewarded habituation (C) then 4 rewarded trials over two days (D).

Figure 2. Sex Differences in the Novel Object Recognition Task. Males, but not females, exhibited a significant preference for the familiar object during the first minute of the task (A), and when assessing preference across time male preference decreased over the length of the task whereas female preference increased, with a significant interaction effect of sex (B). Females spent proportionally more time in the object association zone (C).

Figure 3. Sex differences in the Spatial Task. More females reached the spatial task learning criterion than would be expected by chance, whereas males did not (A). Females exhibited a greater preference for the reward stimuli during the task (B). Females also demonstrated longer decision latencies (C) and their decision latencies significantly predicted their latency to solve the task (enter the reward area) whereas this relationship was not seen in males (D). When subsetting individuals who first entered the unrewarded side and then corrected and entered the rewarded side, females' error correction speed was faster than males' (E).

Figure 4. Individual Variables are Related Across Tasks. When subsetting by individuals who had an intermediate level of preference, relative preference scores are correlated across tasks (A). Decision latency in the NOR task differed by the first choice made in the SPA task (rewarded or unrewarded) (B), and decision latency in the SPA task differed by the first choice made in the NOR task (familiar or novel) (C).

Figure 5. Principal Component Analysis of Variables from Both Tasks. A PCA of variables assessed in both tasks identifies four axes that explain >10% of the variance. Shown here are the proportion of variance for each PC axis (top) and a heatmap of the

eigenvalues of each behavioral variable for each PC axis (bottom) (A). The first PC axis

(24%) can be described by measurements of activity and preference in the spatial task.

The second PC axis (16%) load most strongly by preference in the novel object

recognition task as well as measurements of latency across both tasks. The third PC

axis (11%) loads most strongly with interaction variables across both tasks. The fourth

PC axis (10%) loads very strongly by a single variables: SPA latency to solve the task.

A hierarchical clustering of variables by loading patterns (dendrogram) identified five

highly supported clusters (AU > = 95%, grey boxes), three of which contain variables

form both tasks.

Table 1. Model Selection and Model-Averaged Importance of Terms Analysis.

Through an exhaustive "all-possible models" selection analysis, we identified the model

with the lowest AICc value, which included 13 variables spanning both tasks (variables

in the resulting model in bold). In a model-averaged importance of terms analysis (in

which each term's importance score is related to the number of models it appears in

and the weight of those models), three NOR variables reach the importance threshold

(highlighted with asterisks).

References

Alcazar RM, Hilliard AT, Becker L, Bernaba M, Fernald RD (2014) Brains over brawn: Experience overcomes a size disadvantage in fish social hierarchies. J. Exp. Biol. 217: 1462–1466. https://doi.org/10.1242/jeb.097527

Antunes M, Biala G (2012) The novel object recognition memory: Neurobiology, test procedure, and its modifications. Cogn Process 13: 93–110. https://doi.org/10.1007/s10339-011-0430-z

Banerjee K, Chabris CF, Johnson VE, Lee JJ, Tsao F, Hauser MD (2009) General intelligence in another primate: Individual differences across cognitive task performance in a new world monkey (Saguinus oedipus). PLoS One 4(6): e5883. https://doi.org/10.1371/journal.pone.0005883

Blaser RE, Chadwick L, McGinnis GC (2010). Behavioral measures of anxiety in zebrafish (*Danio rerio*). Behav Brain Res 208:56-62. doi:10.1016/j.bbr.2009.11.009

Bond AB, Kamil AC, Balda RP (2007) Serial reversal learning and the evolution of behavioral flexibility in three species of North American corvids (Gymnorhinus cyanocephalus, Nucifraga columbiana, Aphelocoma californica). J Comp Psychol 121: 372-379. https://doi.org/10.1037/0735-7036.121.4.372

Boogert NJ, Reader SM, Laland KN (2006) The relation between social rank, neophobia and individual learning in starlings. Anim Behav 72(6):1229-39. https://doi.org/10.1016/j.anbehav.2006.02.021

Boogert NJ, Giraldeau LA, Lefebvre L (2008) Song complexity correlates with learning ability in zebra finch males. Anim Behav 76: 1735–1741. https://doi.org/10.1016/j.anbehav.2008.08.009

Branch CL, Pitera AM, Kozlovsky DY, Bridge ES, Pravosudov VV (2019) Smart is the new sexy: female mountain chickadees increase reproductive investment when mated to males with better spatial cognition. Ecol Lett 22: 897–903. https://doi.org/10.1111/ele.13249

Broadbent NJ, Gaskin S, Squire LR, Clark RE (2004) Object recognition memory and the rodent hippocampus. Learn Memory 17:5-11. https://doi.org/10.1101/lm.1650110

Brust V, Wuerz Y, Krüger O (2013) Behavioural Flexibility and Personality in Zebra Finches. Ethology 119: 559–569. https://doi.org/10.1111/eth.12095

Bshary R, Brown C (2014) Fish cognition. Curr Biol 24(19): R947–R950. https://doi.org/10.1016/j.cub.2014.08.043

Buechel SD, Booksmythe I, Kotrschal A, Jennions MD, Kolm N (2016) Artificial selection on male genitalia length alters female brain size. Proc R Soc B Biol Sci 283: 20161796. https://doi.org/10.1098/rspb.2016.1796

Burmeister SS, Jarvis ED, Fernald RD (2005) Rapid behavioral and genomic responses to social opportunity. PLoS Biol 3:e363. https://doi.org/10.1371/journal.pbio.0030363

Burns JG, Rodd FH (2008) Hastiness, brain size and predation regime affect the performance of wild guppies in a spatial memory task. Anim Behav 76: 911–922. https://doi.org/10.1016/j.anbehav.2008.02.017

Calcagno V (2020) glmulti: Model Selection and Multimodel Inference Made Easy. R package
version 1.0.8. https://CRAN.R-project.org/package=glmulti

Caraco T (1981) Energy budgets, risk and foraging preferences in dark-eyed juncos (Junco
hyemalis). Behav Ecol Sociobiol 8: 213–217. https://doi.org/10.1007/BF00299833

Carazo P, Noble DWA, Chandrasoma D, Whiting MJ (2014) Sex and boldness explain
individual differences in spatial learning in a lizard. Proc R Soc B Biol Sci 281: 20133275.
https://doi.org/10.1098/rspb.2013.3275

Chiappe D, MacDonald K (2005) The evolution of domain-general mechanisms in intelligence
and learning. J Gen Psychol 132: 5–40. https://doi.org/10.3200/GENP.132.1.5-40

Chittka L, Skorupski P, Raine NE (2009) Speed-accuracy tradeoffs in animal decision making.
Trends Ecol Evol 24: 400–407. https://doi.org/10.1016/j.tree.2009.02.010

Clayton NS, Krebs JR (1995) Memory in food-storing birds: from behaviour to brain. Curr Opin
Neurobiol 5: 149–154. https://doi.org/10.1016/0959-4388(95)80020-4

Costa SS, Carneiro LA, Gonçalves J (2011) Sex Differences in the Dorsolateral Telencephalon
Correlate with Home Range Size in Blenniid Fish. Brain Behav Evolut 77(1): 55–64.
https://doi.org/10.1159/000323668

Cummings M (2018) Sexual conflict and sexually dimorphic cognition — reviewing their
relationship in poeciliid fishes Behavioral Ecology and Sociobiology. Behav Ecol Sociobiol 72:
73. https://doi.org/10.1007/s00265-018-2483-9

Dalla C, Shors TJ (2009) Sex differences in learning processes of classical and operant
conditioning. Physiol Behav 97: 229–238. https://doi.org/10.1016/j.physbeh.2009.02.035

Desjardins JK, Hofmann HA, Fernald RD (2012) Social context influences aggressive and
courtship behavior in a cichlid fish. PLoS One 7:e32781.
https://doi.org/10.1371/journal.pone.0032781

Dingemanse NJ, De Goede P (2004) The relation between dominance and exploratory behavior
is context-dependent in wild great tits. Behav Ecol 15:1023–1030.
https://doi.org/10.1093/beheco/arh115

Dodson JJ (1988) The nature and role of learning in the orientation and migratory behavior of
fishes. Environ Biol Fishes 23:161-182.

Domjan MP (2015) Principles of Learning and Behavior, 7th Edition. Cengage Learning,
Stamford, CT.

Dougherty LR, Guillette LM (2018) Linking personality and cognition: a meta-analysis. Philos
Trans R Soc B Biol Sci 373: 20170282. https://doi.org/10.1098/rstb.2017.0282

Easton A (2004) Behavioural flexibility, social learning, and the frontal cortex. The cognitive
neuroscience of social behaviour. Psychology Press, Routledge, pp 71–92

Ennaceur A, Delacour J (1988) A new one-trial test for neurobiological studies of memory in
rats. 1: Behavioral data. Behav Brain Res 31: 47–59. https://doi.org/10.1016/0166-
4328(88)90157-X

Etheredge RI, Avenas C, Armstrong MJ, Cummings ME (2018) Sex-specific cognitive–
behavioural profiles emerging from individual variation in numerosity discrimination in Gambusia
affinis. Anim Cogn 21: 37–53. https://doi.org/10.1007/s10071-017-1134-2

Fabre N, García-Galea E, Vinyoles D (2014) Spatial learning based on visual landmarks in the
freshwater blenny Salaria fluviatilis (Asso, 1801). Learn Motiv 48:47–54.
https://doi.org/10.1016/j.lmot.2014.10.002

Farrell TM, Weaver K, An YS, MacDougall-Shackleton SA (2011) Song bout length is indicative
of spatial learning in European starlings. Behav Ecol 23: 101–111.
https://doi.org/10.1093/beheco/arr162

Fernald RD (2014) Cognitive skills needed for social hierarchies. Cold Spring Harb Symp Quant
Biol 79: 229–236. https://doi.org/10.1101/sqb.2014.79.024752

Fernald RD, Hirata NR (1977) Field study of Haplochromis burtoni: quantitative behavioural
observations. Anim Behav 25(4):964-975. https://doi.org/10.1016/0003-3472(77)90048-3

Gagnon KT, Cashdan EA, Stefanucci JK, Creem-regehr SH (2016) Sex Differences in
Exploration Behavior and the Relationship to Harm Avoidance. Hum Nat 27:82–97.
https://doi.org/10.1007/s12110-015-9248-1

Gaulin SJC, Fitzgerald RW (1989) Sexual selection for spatial learning ability. Anim Behav 37:
322–331. https://doi.org/10.1016/0003-3472(89)90121-8

Grosenick L, Clement TS, Fernald RD (2007) Fish can infer social rank by observation alone.
Nature 445:429–432. https://doi.org/10.1038/nature05511

Guillette LM, Reddon AR, Hurd PL, Sturdy CB (2009) Exploration of a novel space is associated
with individual differences in learning speed in black-capped chickadees, Poecile atricapillus.
Behav Processes 82:265–270. https://doi.org/10.1016/j.beproc.2009.07.005

Guillette LM, Reddon AR, Hoeschele M, Sturdy CB (2011) Sometimes slower is better: slow-
exploring birds are more sensitive to changes in a vocal discrimination task. Proc R Soc B Biol
Sci 278:767–773. https://doi.org/10.1098/rspb.2010.1669

Han CS, Jablonski PG, Brooks RC (2015) Intimidating courtship and sex differences in
predation risk lead to sex-specific behavioural syndromes. Anim Behav 109:177–185.
https://doi.org/10.1016/j.anbehav.2015.08.017

Hänninen L, Pastell M (2009) CowLog: open-source software for coding behaviors from digital
video. Behav Res Methods 41(2):472-476. doi:10.3758/BRM.41.2.472

Harris S, Ramnarine IW, Smith HG, Pettersson LB (2010) Picking personalities apart: estimating
the influence of predation, sex and body size on boldness in the guppy Poecilia reticulata. Okios
119(11):1711–1718. https://doi.org/10.1111/j.1600-0706.2010.18028.x

Healy SD, Braham SR, Braithwaite VA (1999) Spatial working memory in rats: No differences
between the sexes. Proc R Soc B Biol Sci 266: 2303–2308.
https://doi.org/10.1098/rspb.1999.0923

Hebets E, Sullivan-Beckers L (2010) Mate Choice and Learning. *Encyclopedia of Animal*
Behavior. Elsevier pp 389-393

Heske EJ, Ostfeld RS (1990) Sexual Dimorphism in Size, Relative Size of Testes, and Mating
Systems in North American Voles. J Mammal 71: 510–519. https://doi.org/10.2307/1381789

Hofmann HA (2003) Functional genomics of neural and behavioral plasticity. J Neurobiol
54:272–282. https://doi.org/10.1002/neu.10172

Hofmann HA, Fernald RD (2001) What cichlids tell us about the social regulation of brain and
behaviour. Journal of Aquaculture & Aquatic Sciences 9:17-31.

Huffman LS, Mitchell MM, O'Connell LA, Hofmann HA (2012) Rising StARs: Behavioral,
hormonal, and molecular responses to social challenge and opportunity. Horm Behav 61:631-
641. https://doi.org/10.1016/j.yhbeh.2012.02.016

Hughes RN (1968) Behaviour of male and female rats with free choice of two environments
differing in novelty. Anim Behav 16:92–96. https://doi.org/10.1016/0003-3472(68)90116-4

Ishikawa K, Watanuki Y (2002) Sex and individual differences in foraging behavior of Japanese
cormorants in years of different prey availability. J Ethol 20:49–54.
https://doi.org/10.1007/s10164-002-0053-z

Jacobs LF, Gaulin SJC, Sherry DF, Hoffman GE (1990) Evolution of spatial cognition: Sex-
specific patterns of spatial behavior predict hippocampal size. PNAS 87:6349–6352.
https://doi.org/10.1073/pnas.87.16.6349

Jacobs LF (1995) The Ecology of Spatial Cognition. Adaptive Patterns of Space Use and
Hippocampal Size in Wild Rodents. Behavioural Brain Research in Naturalistic and Semi-
Naturalistic Settings. Nato Science Series D 82:301–322.Springer

Jonasson Z (2005) Meta-analysis of sex differences in rodent models of learning and memory:
A review of behavioral and biological data. Neurosci Biobehav Rev 28:811–825.
https://doi.org/10.1016/j.neubiorev.2004.10.006

Jones CM, Braithwaite VA, Healy SD (2003) The evolution of sex differences in spatial ability.
Behav Neurosci 117:403–411. https://doi.org/10.1037/0735-7044.117.3.403

Jun JJ, Longtin A, Maler L (2016) Active sensing associated with spatial learning reveals
memory-based attention in an electric fish. J Neurophysiol 115:2577–2592.
https://doi.org/10.1152/jn.00979.2015

Kidd MR, Dijkstra PD, Alcott C, Lavee D, Ma J, O'Connell LA, Hofmann HA (2013a)
Prostaglandin F2alpha facilitates female mating behavior based on male performance.
Behavioral Ecology and Sociobiology 67:1307 - 1315. https://doi.org/10.1007/s00265-013-1559-
9

Kidd MR, O'Connell LA, Kidd CE, Chen CW, Fontenot MR, Williams SJ, Hofmann HA (2013b)
Female preference for males depends on reproductive physiology in the African cichlid fish
Astatotilapia burtoni. General and Comparative Endocrinology 180:56 - 63.
https://doi.org/10.1016/j.ygcen.2012.10.014

Kleiman DG (1977) Monogamy in Mammals. Q Rev Biol 52(1):39-69

Kolata S, Light K, Matzel LD (2008) Domain-specific and domain-general learning factors are
expressed in genetically heterogeneous CD-1 mice. Intelligence 36:619–629.
https://doi.org/10.1016/j.intell.2007.12.001

Laland KN, Reader SM (1999) Foraging innovation in the guppy. Anim Behav 57:331–340.
https://doi.org/10.1006/anbe.1998.0967

Layton C, Fulton CJ (2014) Status-dependent foraging behaviour in coral reef wrasses. Coral
Reefs 33:345–349. https://doi.org/10.1007/s00338-014-1138-1

Lombardi Brandão M, Braithwaite VA, Gonçalves-de-Freitas E (2015) Isolation impairs cognition
in a social fish. Appl Anim Behav Sci 171:204–210.
https://doi.org/10.1016/j.applanim.2015.08.026

Lucon-Xiccato T, Bisazza A (2014) Discrimination reversal learning reveals greater female
behavioural flexibility in guppies. Biol Lett 10:20140206. https://doi.org/10.1098/rsbl.2014.0206

Lucon-Xiccato T, Dadda M, Bisazza A (2016) Sex Differences in Discrimination of Shoal Size in
the Guppy (Poecilia reticulata). Ethology 122:481–491. https://doi.org/10.1111/eth.12498

Lucon-Xiccato T, Bisazza A (2017a) Sex differences in spatial abilities and cognitive flexibility in
the guppy. Anim Behav 123:53–60. https://doi.org/10.1016/j.anbehav.2016.10.026

Lucon-Xiccato T, Bisazza A (2017b) Individual differences in cognition among teleost fishes.
Behav Processes 141:185-195. http://dx.doi.org/10.1016/j.beproc.2017.01.015

Lynch KS, Rand AS, Ryan MJ, Wilczynski W (2005) Plasticity in female mate choice associated
with changing reproductive states. Anim Behav 69:689–699.
https://doi.org/10.1016/j.anbehav.2004.05.016

Madden JR, Langley EJG, Whiteside MA, Beardsworth CE, Van Horik JO (2018) The quick are
the dead: Pheasants that are slow to reverse a learned association survive for longer in the wild.
Philos Trans R Soc B Biol Sci 373: 20170297. https://doi.org/10.1098/rstb.2017.0297

Magurran AE, Nowak MA (1991) Another battle of the sexes: The consequences of sexual
assymmetry in mating costs and predation risk in the guppy, Poecilia reticulata. Proc R Soc B
Biol Sci 246: 31–38. https://doi.org/10.1098/rspb.1991.0121

Magurran AE, Seghers BH (1994) A Cost of Sexual Harassment in the Guppy, Poecilia
reticulata. Proc R Soc B Biol Sci 258:52–89. https://doi.org/10.1098/rspb.1994.0147

Maille A, Schradin C (2016) Survival is linked with reaction time and spatial memory in African
striped mice. Biol Lett 12: 20160346. https://doi.org/10.1098/rsbl.2016.0346

Mamuneas D, Spence AJ, Manica A, King AJ (2015) Bolder stickleback fish make faster
decisions, but they are not less accurate. Behav Ecol 26: 91–96.
https://doi.org/10.1093/beheco/aru160

Maruska KP, Fernald RD (2010) Behavioral and physiological plasticity: rapid changes during
social ascent in an African cichlid fish. Horm Behav 58:230–240.
https://doi.org/10.1016/j.yhbeh.2010.03.011

Maruska KP, Fernald RD (2018) Astatotilapia burtoni: A Model System for Analyzing the
Neurobiology of Behavior. ACS Chem Neurosci 9:1951–1962.
https://doi.org/10.1021/acschemneuro.7b00496

Maximino C, Marques de Brito T, de Mattos Dias CAG, Gouveia Jr A, Morato S (2010)
Scototaxis as anxiety-like behavior in fish. Nat Protoc 5(2): 221-228.
https://doi.org/10.1038/nprot.2009.225

May Z, Morrill A, Holcombe A, Johnston T, Gallup J, Fouad K, Schalomon M, Hamilton TJ
(2016) Object recognition memory in zebrafish. Behav Brain Res 296:199–210.
https://doi.org/10.1016/j.bbr.2015.09.016

Mazza V, Eccard JA, Zaccaroni M, Jacob J, Dammhahn M (2018) The fast and the flexible:
cognitive style drives individual variation in cognition in a small mammal. Anim Behav 137: 119–
132. https://doi.org/10.1016/j.anbehav.2018.01.011

Miletto Petrazzini ME, Agrillo C, Piffer L, Dadda M, Bisazza A (2012) Development and
application of a new method to investigate cognition in newborn guppies. Behav Brain Res
233:443–449. https://doi.org/10.1016/j.bbr.2012.05.044

Miletto Petrazzini ME, Bisazza A, Agrillo C, Lucon-Xiccato T (2017) Sex differences in
discrimination reversal learning in the guppy. Anim Cogn 20:1081–1091.
https://doi.org/10.1007/s10071-017-1124-4

O'Connell LA, Hofmann HA (2011) Genes, hormones, and circuits: an integrative approach to
study the evolution of social behavior. Front Neuroendocrinol 32:320-35.
https://doi.org/10.1016/j.yfrne.2010.12.004

O'Connell LA, Hofmann HA (2012) Evolution of a Vertebrate Social Decision-Making Network.
Science 336:1154–1157. https://doi.org/10.1126/science.1218889

Orsini CA, Setlow B (2017) Sex differences in animal models of decision making. J Neurosci
Res 95:260–269. https://doi.org/10.1002/jnr.23810

Øverli Ø, Sørensen C (2006) Behavioral indicators of stress-coping style in rainbow trout: Do
males and females react differently to novelty? Physiol Behav 87:506–512.
https://doi.org/10.1016/j.physbeh.2005.11.012

Phelps SM, Ophir AG (2009) Monogamous Brains and Alternative Tactics: Neuronal V1aR,
Space Use, and Sexual Infidelity among Male Prairie Voles. Cognitive Ecology II. University of
Chicago Press. pp 156-176.

Pongrácz P, Vida V, Bánhegyi P, Miklósi Á (2008) How does dominance rank status affect
individual and social learning performance in the dog (Canis familiaris)? Anim Cogn 11:75–82.
https://doi.org/10.1007/s10071-007-0090-7

R Core Team (2013) R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. http://www.R-project.org/

Rapaport LG, Brown GR (2008) Social Influences on Foraging Behavior in Young Nonhuman
Primates: Learning What, Where, and How to Eat. Evol Anthropol 17:189–201.
https://doi.org/10.1002/evan.20180

Reichert MS, Quinn JL (2017) Cognition in Contests: Mechanisms, Ecology, and Evolution.
Trends Ecol Evol 32:773–785. https://doi.org/10.1016/j.tree.2017.07.003

Rice MA, Hobbs LE, Wallace KJ, Ophir AG (2017) Hormones and Behavior Cryptic sexual
dimorphism in spatial memory and hippocampal oxytocin receptors in prairie voles (Microtus
ochrogaster). Horm Behav 95: 94–102. https://doi.org/10.1016/j.yhbeh.2017.08.003

Renn SCP, Fraser EJ, Aubin-Horth N, Trainor BC, Hofmann HA (2012) Females of an African
cichlid fish display male-typical social dominance behavior and elevated androgens in the
absence of males. Horm Behav 61:496–503. https://doi.org/10.1016/j.yhbeh.2012.01.006

Rodriguez-Santiago M*, Nührenberg P*, Derry J, Deussen O, Francisco FA, Garrison LK, Garza
SF, Hofmann HA, Jordan A (2020) Behavioral traits that define social dominance are the same
that reduce social influence in a consensus task. PNAS.
https://doi.org/10.1073/pnas.2000158117

Rosati AG, Hare B (2012) Decision making across social contexts: Competition increases
preferences for risk in chimpanzees and bonobos. Anim Behav 84:869–879.
https://doi.org/10.1016/j.anbehav.2012.07.010

Salas C, Rodríguez F, Vargas JP, Durán E, Torres B (1996) Spatial learning and memory
deficits after telencephalic ablation in goldfish trained in place and turn maze procedures. Behav
Neurosci 110(5):965–980. https://doi.org/10.1037/0735-7044.110.5.965

Sano M (1993) Foraging activities and diets of males and females in a haremic sandperch
(Pisces: Pinguipedidae). Mar Ecol Prog Ser 98: 55–59. https://doi.org/10.3354/meps098055

Saucier DM, Shultz SR, Keller AJ, Cook CM, Binsted G (2008) Sex differences in object location
memory and spatial navigation in Long-Evans rats. Anim Cogn 11:129–137.
https://doi.org/10.1007/s10071-007-0096-1

Shettleworth SJ (2010) Cognition, evolution, and behaviour, 2nd edn. Oxford University Press,
Oxford

Shohet AJ, Watt PJ (2009) Female guppies Poecilia reticulata prefer males that can learn fast. J
Fish Biol 75:1323–1330. https://doi.org/10.1111/j.1095-8649.2009.02366.x

Smith C, Phillips A, Reichard M (2015) Cognitive ability is heritable and predicts the success of
an alternative mating tactic. Proc R Soc B Biol Sci 282: 20151046.
http://dx.doi.org/10.1098/rspb.2015.1046

Staddon, JER (1983) Adaptive behavior and learning. Second Edition. New York: Cambridge
University Press

Stapley J, Keogh JS (2004) Exploratory and antipredator behaviours differ between territorial
and nonterritorial male lizards. Anim Behav 68:841–846.
https://doi.org/10.1016/j.anbehav.2004.02.008

Stowe M, Kotrschal K (2007) Behavioural phenotypes may determine whether social context
facilitates or delays novel object exploration in ravens (Corvus corax). J Ornithol 148(Supple 2):
179–184. https://doi.org/10.1007/s10336-007-0145-1

Suzuki R, Terada Y, Shimodaira H (2019) pvclust: Hierarchical Clustering with P-Values via
Multiscale Bootstrap Resampling. R package version 2.2-0. https://CRAN.R-
project.org/package=pvclust

Swasnson HH (1966) Sex differences in behaviour of hamsters in open field and emergence
tests: effects of pre- and post-pubertal gonadectomy. Anim Behav 14(4): 522-529
https://doi.org/10.1016/S0003-3472(66)80054-4

Taborsky B, Oliveira RF (2012) Social competence: an evolutionary approach. TREE 27:679-
688. https://doi.org/10.1016/j.tree.2012.09.003

Thornton A, McAuliffe K (2006) Teaching in Wild Meerkats. Science 313:227–229.
https://doi.org/10.1126/science.1128727

Titulaer M, van Oers K, Naguib M (2012) Personality affects learning performance in difficult
tasks in a sex-dependent way. Anim Behav 83:723–730.
https://doi.org/10.1016/j.anbehav.2011.12.020

Torchiano M (2020) effsize: Efficient Effect Size Computation. doi: 10.5281/zenodo.1480624
(https://doi.org/10.5281/zenodo.1480624), R package version 0.8.0. https://CRAN.R-
project.org/package=effsize

Wallace KJ, Rausch RT, Ramsey ME, Cummings ME (2020) Sex differences in cognitive
performance and style across domains in mosquitofish (Gambusia affinis). Anim Cogn 23:655-
669. https://doi.org/10.1007/s10071-020-01367-2

Watson KK, Platt ML (2008) Neuroethology of reward and decision making. Philos Trans R Soc
B Biol Sci 363:3825–3835. https://doi.org/10.1098/rstb.2008.0159

Weitekamp CA, Hofmann HA (2017) Neuromolecular correlates of cooperation and conflict
during territory defense in a cichlid fish. Horm Behav 89: 145–156.
https://doi.org/10.1016/j.yhbeh.2017.01.001

Galef Jr BG, White DJ (1998) Mate choice copying and conspecific cueing in Japanese quail,
Coturnix coturnix japonica. Anim Behav 55: 545-552. https://doi.org/10.1006/anbe.1997.0616

White SA, Nguyen T, Fernald RD (2002) Social regulation of gonadotropin-releasing hormone. J
Exp Biol 205:2567–2581.

Wood LS, Desjardins JK, Fernald RD (2011) Effects of stress and motivation on performing a
spatial task. Neurobiol Learn Mem 95: 277–285. https://doi.org/10.1016/j.nlm.2010.12.002

Yaskin VA (2011) Seasonal changes in hippocampus size and spatial behavior in mammals and
birds. Biol Bull Rev 1: 279. https://doi.org//10.1134/S2079086411030108

A   NOR Habituation

50cm

53cm

n = **16** individuals
(free-swimming for habituation)

15cm

75cm

B   NOR Test

familiar
object

novel
object

opaque tube
for focal
acclimation

C   SPA Habituation

transparent
tube

11cm

7cm

37.5cm

31cm

opaque tube
for focal
acclimation

7cm

50cm

D   SPA Test

female
conspecific
reward

Term Name	Importance Score
NOR novel object interaction bout duration	0.994 *
NOR familiar preference	0.938 *
NOR novel object approaches	0.927 *
SPA decision latency	0.800
SPA % association	0.769
NOR % interaction	0.758
SPA reward latency	0.622
NOR non-engagement	0.548
NOR % association	0.520
SPA reward preference	0.394
SPA % decision (association + interaction)	0.365
SPA reward preference following solution	0.358
SPA % interaction	0.352
NOR familiar object interaction bout duration	0.281
SPA thigmotaxis	0.239
NOR object approach latency	0.183
SPA interaction latency	0.161
SPA activity	0.096
NOR familiar object approaches	0.033