Plant carbohydrate depletion impairs water relations and spreads via

ectomycorrhizal networks.

Running title: Fungal networks spread plant carbon stress

Length: Abstract: 238 words. Main text (excluding methods): 3,224 words. Methods: 2,013 words.

References: 52. Figures: 6. Tables: 0. Text boxes: 0.

Authors: Gerard Sapes[1, 2]*, Patrick Demaree[1], Ylva Lekberg[3,4], & Anna Sala[1].

[1] Division of Biological Sciences, University of Montana, Missoula, MT 59812, USA.

[2] Department of Ecology, Evolution, and Behavior, University of Minnesota, Saint Paul,

Minnesota 55108, USA

[3] MPG Ranch, Missoula, MT 59801, USA

[4] W.A. Franke College of Forestry & Conservation, University of Montana, Missoula, MT 59812,

USA.

Corresponding Author:

Gerard Sapes (gsapes@umn.edu)

Phone: (406) 207-7955

Department of Ecology, Evolution, and Behavior

University of Minnesota

1987 Upper Buford Circle 55108, Saint Paul, Minnesota, USA

Classification: Biological Sciences/Plant Biology

Keywords: Carbon starvation, carbon transfer, drought, ectomycorrhizae, osmotic potential, *Pinus*

ponderosa, seedlings, turgor, tradeoff, fungal networks.

Author contributions:

G.S., P.D., Y.L. and A.S. designed the experiment. G.S. and P.D. collected data and G.S. analyzed

the data with contributions from A.S. and Y.L. G.S., P.D., and A.S. wrote the manuscript with

contributions from Y.L.

Competing Interests: The authors declare no competing interests.

This PDF file includes:

Main Text

Figures 1 to 6

Abstract

Carbon and water relations are fundamental to plant life and strongly interact. Under drought, the

ability of plants to assimilate carbon is reduced, which increases their consumption of stored labile

carbon in the form of non-structural carbohydrates (NSC). Stored NSC depletion may impair plant

water relations, but mechanisms are not clear, and we do not know if its effects

are independent of water deficit. If so, carbon costs of fungal symbionts could also indirectly

influence plant drought tolerance through stored NSC depletion. We connected well-watered

Pinus ponderosa seedling pairs via ectomycorrhizal (EM) networks where one seedling was shaded

and the other experienced full light and compared responses to seedling pairs in the light. We

measured plant water relations and traced carbon movements using $^{13}CO_2$ to explore the

mechanisms linking stored NSC to water relations, and to identify potential tradeoffs between the

ability to endure low water potentials and maintaining EM fungi under carbon-limiting conditions.

Even in the absence of drought, mild NSC depletion impaired osmoregulation capacity and turgor

maintenance, a critical strategy to tolerate drought. This demonstrates that NSC storage

influences plant water relations independently of plant water status. We also found that EM

networks propagated NSC depletion and its negative effects on water relations from carbon

stressed hosts to non-stressed hosts. These results highlight carbon allocation tradeoffs between

supporting fungal symbionts and retaining water via stored NSC and have implications for biotic

interactions and forest drought responses.

Significance Statement

The potential effects of future drought on global carbon cycles, vegetation-climate feedbacks,

species distributions and their ecological impacts, urgently call for a clear understanding of factors

influencing vegetation tolerance to drought. Key to this is the understanding of mechanisms and

processes by which plants tolerate drought and how prevalent plant-fungal interactions may

influence these processes.

We demonstrate that even mild depletion of plant non-structural carbohydrate (NSC) storage

readily decreases plant water retention capacity, therefore decreasing tolerance to drought.

Because plant-fungal interactions depend on NSC exchange, plants face carbon-allocation

tradeoffs between maintaining drought tolerance and feeding fungal symbionts. The impacts of

these tradeoffs extend across plants connected via ectomycorrhizal networks as fungi propagate

NSC depletion from NSC-limited plants to non-stressed individuals.

Main Text

Introduction

Under future climate, more intense and frequent droughts are expected to increase forest

mortality and to impact forest ecosystems and global carbon cycles [1]. Predicting the extent of

these impacts requires a full understanding of the mechanisms and processes associated with

drought-induced mortality (DIM) [2,3]. Under drought, plants ultimately die from desiccation

because water supply does not keep pace with water loss. Desiccation and death are generally

associated with failure of the vascular system to transport water due to embolism formation

(hydraulic failure, Adams *et al.* 2017). Reduced photosynthesis under prolonged drought

eventually leads to the depletion of stored non-structural carbohydrates (NSC) and carbon

limitation, a process thought to contribute to desiccation by decreasing the plant's ability to retain

and move water [5–8]. Hydraulic failure is thought to be the main driver of DIM with NSC depletion

playing a secondary, interacting role. However, it is not known whether NSC depletion *per se*

impairs plant water relations because its effects have not been experimentally isolated from those

caused by drought (e.g., reduced water transport) (Hartmann and Trumbore 2016, Adams et al.

2017). Yet, NSC depletion may be more relevant than currently thought as the wide array of

physiological processes that demand NSC may limit carbon allocation to storage and its positive

effects on water relations. For instance, in addition to metabolic costs, many trees incur significant

carbon costs to maintain symbiotic partnerships with ectomycorrhizal (EM) fungi [11]. Consequently,

plants may face carbon allocation tradeoffs between sustaining symbionts and maintaining water

relations via NSC storage under carbon-limiting conditions. Whether these tradeoffs exist has not

been tested despite their potential implications for forest mortality under global change. In this

study, we examine the independent effects of stored NSC for plant water relations and assess

potential tradeoffs between maintaining water relations and sustaining EM symbionts when

carbon becomes limiting.

Stored NSC have been classically understood as a carbon safeguard for growth and energy.

However, NSC may play an important role in the movement and retention of water [5] including

lowering tissue osmotic potential to increase tolerance to low water potentials under drought [12].

While the osmotically active soluble fraction of NSC plays a direct osmotic role, starch could also

play an indirect role as a source of osmotically active organic compounds and energy [13]. Under

carbon-limiting conditions of any kind, NSC-consuming processes such as respiration, defense

production, and maintenance of ectomycorrhizae rely on the same pool of stored NSC potentially

involved in osmoregulation [14]. Consistently, complete consumption of stored NSC is rare under

natural and experimental conditions, and plants rarely deplete their stored NSC below ca. 40% of

their seasonal maximums [4,15]. While NSC have been associated with maintenance of hydraulic

conductivity, capacitance, and turgor [9,16,17], it is unclear how much of the total stored NSC pool

plants must conserve to maintain these functions, even under non-drought conditions [18]. If a

significant fraction of stored NSC must remain unconsumed to maintain the capacity to move and

retain water, we may have underestimated the role of NSC depletion in DIM, which could have

important implications for modelling and prediction of tree mortality.

In temperate coniferous and boreal ecosystems, virtually all trees and seedlings form

associations with EM fungi and establish underground EM networks through which resources can

be exchanged [19–21]. These networks have been shown to enhance seedling establishment and

increase ecosystem-level nutrient cycling [22,23]. Plant hosts allocate up to 30% of their assimilated

carbon to EM fungi [11,24] in exchange for nutrients. As such, EM fungi could indirectly influence

tolerance to drought via network-mediated carbon redistribution and host NSC depletion under

carbon-limiting conditions. For instance, if stochastic or microsite effects (e.g., herbivory, shading,

microsite low moisture) cause carbon deficit and NSC depletion on some plant hosts but not

others within the network, NSC-depleted hosts could decrease carbon supply to their associated

EM fungi and trigger fungal carbon deficit. Non-NSC-depleted hosts may then offset the deficit of

the EM network with their own stored NSC, a response that could indirectly compromise their own

water relations especially if they cannot regulate carbon flow to EM fungi. There is debate on

whether carbon can be transferred among connected hosts via EM networks [20,25,26] or whether

NSC simply stays within the EM fungi [27]. In either case, network-mediated carbon redistribution

may influence how individual plants and entire forest stands respond to drought. Whether and

under what circumstances fungal carbon demand and redistribution compromise allocation to

plant NSC storage and water relations is unknown but deserves attention.

Here, we used an experimental design that depletes plant NSC pools by combining shading

treatments and EM networks among well-watered ponderosa pine seedlings (*Pinus ponderosa*

Douglas ex C. Lawson). This design allowed us to unveil the mechanisms (and their importance)

linking stored NSC and plant water relations without the confounding effects of water deficit, as

well as the carbon dynamics between ectomycorrhizal fungi and their plant hosts under carbon-

limiting conditions. Our experiment involved seedlings as they often determine forest distributions

through establishment [28]. Also, because of their small size and little storage capacity, seedlings

may be especially susceptible to carbon allocation tradeoffs between maintaining water relations

and EM fungi. Specifically, we asked 1) does stored NSC depletion affect host water relations even

without water deficit?; and 2) do ectomycorrhizal networks extend the effects of NSC depletion on

plant water relations to non-stressed neighbors? We found that, even in the absence of drought,

mild NSC depletion indirectly decreased plant tolerance to drought by impairing osmoregulation

capacity and turgor maintenance. These results unambiguously link NSC storage to plant water

relations. We also found that EM networks propagated NSC depletion and impaired critical

drought tolerance strategies in non-stressed hosts. This demonstrates the existence of resource-

allocation tradeoffs between supporting fungal symbionts and retaining water via stored NSC and

suggests that EM networks can affect tree and possibly entire forest tolerance to drought.

Results

<u>NSC decreased in LD and D seedlings</u>

Shading caused a 72% reduction in whole-plant NSC concentrations in dark (D) relative to

light (LL) seedlings ($p < 0.001$). Light seedlings paired with dark (LD) plants also experienced NSC

depletion (45% reduction) and showed intermediate whole-plant NSC pools relative to LL and D

seedlings ($p < 0.001$, and $p = 0.009$, respectively) (Fig. 2). When broken down to each NSC

compound, we found that starch was higher in LL seedlings ($p < 0.001$) relative to LD and D

seedlings. LD seedlings had intermediate starch levels although only marginally higher than D

seedlings (p = 0.079). Both sucrose and glucose + fructose were lower in D seedlings than in LL and

LD seedlings (sucrose: LL: p < 0.001, LD: p = 0.040; glucose and fructose: LL: p < 0.001, LD: p =

0.018). However, no differences were detected between LL and LD seedlings. Patterns within each

organ were consistent with those observed at the whole-plant level (Figure S6).

<u>Labeled carbon stayed within ectomycorrhizal fungi</u>

Labeled LL and LD seedlings showed significant increases in $^{13}C/^{12}C$ ratios ($\delta^{13}C$) across all

organs relative to pre-labeling base-line values (Fig. 3, Figure S7) while non-labeled seedlings

stayed at base-line levels. This indicates that only labeled seedlings incorporated the C-isotope and

that there was no contamination during the labeling process. In labeled seedlings, ^{13}C progressively

declined from needles to roots (p < 0.001 in all pairwise contrasts across seedling types and

organs, Fig. 3, Figure S7: Panels a-c). High levels of ^{13}C were also present in fungal tissue associated

with labeled LL and LD seedlings (Fig. 3, Figure S8: Panel a). To a lesser extent, ^{13}C was also present

in fungal tissue associated with non-labeled LL plants paired with labeled LL plants (Fig. 3, Figure

S8: Panel b). ^{13}C in fungi associated with non-labeled LL plants confirmed the existence of a

mycorrhizal network between seedlings at harvesting (Figure S1 and S2). However, ^{13}C was not

detected in colonized root tips of non-labeled D plants paired with labeled LD seedlings (Fig. 3,

Figure S8: Panel b). In all cases, ^{13}C levels within non-labeled seedlings were similar to base-line

values regardless of organ or seedling type (Fig. 3, Figure S7: Panels d-f).

<u>LD and D seedlings exhibited lower turgor</u>

Hydraulic conductivity, which describes the capacity to transport water, did not

significantly differ among seedling types in any organ (Figure S9). However, osmotic potentials,

which describe the ability to retain cell water at low water potentials, as is typical under drought,

were up to ca. 1.5 MPa lower (i.e., higher water retention capacity) in LL seedlings than in LD and

D seedlings (Fig. 4a-c). These differences were observed in needles (p < 0.001 in all pairwise

contrasts), stems (D: p = 0.002, LD: p < 0.001), and roots (D: p < 0.001, LD: p = 0.001). High osmotic

potentials were associated with low pressure potentials (i.e., turgor) in D and LD needles (D: p <

0.001, LD: p < 0.001, Fig. 4d) and stems (D: p = 0.012, LD: p < 0.001, Fig. 4e). Pressure potentials in

LD and D stems were low enough to bring stems to turgor loss (i.e., pressure potential lower than

0). As a result, we observed lower stem relative water content in NSC-depleted seedlings than in

non-depleted plants (D: p < 0.001, LD: p = 0.047, Figure S10).

NSC depletion impaired water relations

NSC depletion in needles was associated with less negative (higher) needle osmotic

potentials (p < 0.001, R = -0.83, Table S2, Fig. 5a). This relationship was linear when NSC depletion

was large and plateaued when NSC were ca. 20% depleted relative to non-depleted seedlings. The

pressure potential (necessary for positive turgor) mirrored this pattern (p < 0.001, R = 0.64, Table

S2, Fig. 5b) and plateaued at 1.5 MPa when NSC was ca. 20% NSC depleted.

Stored NSC depletion also influenced the relationship between needle pressure potential

and needle water potential (WP) (p < 0.001, R = 0.78, Fig. 6; Table S2). Plants with greater NSC

depletion showed lower pressure at any given WP and higher WP at turgor loss (NSC deviation

from controls: p < 0.001, Fig. 6). Additionally, plants with greater NSC depletion lost more turgor

per unit decline in WP than less NSC-depleted plants with (Leaf WP x NSC deviation from controls:

p = 0.006, Fig. 6).

Discussion

It has long been known that plants exchange water (lost through stomata) for carbon

(assimilated from atmospheric CO_2). Our results demonstrate that plants also need stored labile

carbon in the form of non-structural carbohydrates (NSC) to retain water. We asked 1) whether

stored NSC depletion affect host water relations even without water deficit and 2) whether

ectomycorrhizal networks extend the effects of NSC depletion on plant water relations to non-

stressed neighbors. We found that NSC depletion impaired turgor maintenance -a critical drought

tolerance strategy- even in well-watered plants. We also found that carbon-limiting conditions can

result in a tradeoff between supporting symbionts and retaining water via stored NSC, and that

this effect can be propagated by the EM network to otherwise non-stressed plants. During periods

such as drought, when carbon supply via photosynthesis is limited, carbon allocation tradeoffs

may be exacerbated by EM fungi and influence plant survival.

<u>Depletion of stored NSC impairs a critical drought tolerance strategy</u>

Despite being well-watered, seedlings that experienced stored NSC depletion below 20%

of non-NSC-depleted plants experienced declines in turgor and osmoregulation capacity (Figs. 4-6).

Leaf osmotic potentials shifted from values typical of drought-tolerant temperate conifers (-2.3

MPa) to values greater than those found in drought-vulnerable croplands (-1.3 MPa) (Bartlett *et al.*

2012) (Fig. 4), which led to turgor loss (Fig. 6). These shifts were consistent with declines in total

NSC concentrations but were not always reflected by declines in the osmotically active fraction of

NSC (glucose, fructose, and sucrose). In fact, shifts in osmotic potential were mainly associated

with declines in osmotically inactive starch. These results suggest that the effects of NSC depletion

on osmoregulation are largely indirect, perhaps via the synthesis of osmotically active molecules

other than soluble sugars (e.g., sugar alcohols, proteins, amino acids) [29–31]. Regardless of the

specific mechanisms, however, our results show that stored NSC depletion impairs a critical

drought tolerance strategy in plants. The link between stored NSC and maintenance of osmotic

and pressure potentials could explain previous findings that NSC-depleted plants experience lower

phloem turgor under well-watered conditions [9], lower water content at any degree of water deficit

[6], and reach lower stem WP and die at faster rates than NSC-enriched plants when exposed to

similar water deficit [16]. Based on our results, and given that osmotic potentials decrease from

moist to dry biomes to compensate for low water potentials typical of dry regions [12], drought-

tolerant plants from dry biomes might require greater NSC pools to maintain lower osmotic

potentials than those from moist biomes and be more prone to experience carbon-allocation

tradeoffs.

212   Drought impairs plant water transport and depletes NSC, and both have negative effects

on plant water relations. Therefore, it is not surprising that both hydraulic failure and NSC

depletion are often associated with drought-induced mortality (DIM) [4,16]. However, despite an

increasing interest in the potential role of stored NSC on plant water relations [5,9,10], evidence that

NSC affects plant water relations independent of drought has been lacking. The negative impact of

NSC depletion on osmoregulation capacity and turgor maintenance we show here provides such

evidence. Our findings are consistent with the observation that plants prevent NSC depletion

below certain thresholds at all costs [15,32], as well as observed increases in stomatal conductance in

response to increases in osmotic potential [33]. Together, these findings suggest that NSC depletion

may further decrease drought tolerance by preventing total stomatal closure under drought.

While still hypothetical, this could explain sudden increases in stomatal conductance at low water

potentials [32], accelerated water loss [6] and accelerated declines in WP observed in drought-

stressed, NSC-depleted plants [16] leading to early hydraulic failure and death. Our results provide

support for a greater focus on plant water balance (turgor maintenance) in our efforts to

understand interactions between water and carbon to better monitor and predict DIM [6,7,34].

Mechanistic models that consider turgor loss may achieve higher species-specific predictive

capacity because turgor loss marks the onset of many processes that lead to plant death.

Additionally, turgor loss integrates many processes, including regulation of water transport, water

loss and water retention. Additional research is needed to understand the specific mechanisms by

which NSC depletion impairs osmoregulation across plant types, drought strategies, and

environments and how species-specific carbon allocation strategies, including to storage and

osmoregulation, influence tolerance to drought.

<u>Ectomycorrhizal networks can exacerbate NSC depletion and carbon allocation tradeoffs</u>

We hypothesized that shaded seedlings (D) would indirectly increase carbon demand on

connected, non-shaded seedlings (LD) due to a disruption of carbon allocation to EM fungi from D

seedlings. Our results supported this hypothesis: LD seedlings became NSC-depleted relative to LL

seedlings (Fig. 2), indicating that carbon demand in LD seedlings exceeded supply. Additional

measurements confirmed that assimilation did not decrease in LD relative to LL seedlings (i.e.

supply did not decrease; Figure S5). Likewise, there were no differences in growth and respiration

between LD and LL seedlings (Figure S5). Clearly, stored carbon was lost in LD seedlings during the

three weeks of shading, but it did not go to any other sink we could measure, leaving the EM

network as the most likely sink. However, we did not find more ^{13}C in the EM network in the LD

pair than the LL pair, and the fungal tissue closest to the D seedling was not enriched, which is

opposite to what we expected if the LD seedling bore a greater cost for supporting the network.

This is not because the EM network connecting the seedlings was lacking. We confirmed the

presence of the network both visually (Figures S1 and S2) and isotopically as ^{13}C travelled from

labeled seedlings to fungal tissue connected to neighboring, unlabeled seedlings (Fig 3). We

believe that the most likely explanation is that, by the time labeling occurred, the LD seedling had

either reduced carbon allocation to the EM network as costs exceeded benefits [24,35,36], or the EM

fungi closest to the starved D seedling had died or become dormant, reducing the overall carbon

demand on the LD seedling. While we cannot unambiguously attribute the missing carbon in LD

seedlings to increased allocation to EM fungi (and possibly D seedlings [20]), it does not change the

fact that the EM network propagated NSC depletion to a non-stressed plant and that this had

consequences for drought tolerance strategies. Thus, although EM fungi may increase survival by

hydraulic lift [19], they may also reduce plant tolerance to drought [37] by contributing to the carbon

allocation tradeoff between supporting symbionts and retaining water via stored NSC.

Carbon costs of sustaining EM may increase when plant or fungal carbon demand

increases [38] and during periods of carbon limitation such as prolonged drought [39]. Costs of

sustaining EM networks might also increase substantially when some individuals connected to the

network die (e.g. fire, drought, disease, and insect induced mortality) and the surviving trees must

subsidize carbon supply to EM until fungal biomass re-adjusts to new levels of carbon supply. In

these situations, EM networks might increase NSC depletion rates in hosts, hinder their

osmoregulation capacity, and hasten desiccation rates. It is possible that our pairwise approach

overestimates costs to individual seedlings and that EM network-driven NSC depletion is less

substantial in complex networks of highly interconnected plants where increased costs are

subsidized by several hosts [40,41]. As such, EM networks may even enhance community resilience to

drought, although this will likely depend on the proportion of stressed hosts within the network.

The carbon costs of EM-networks under carbon-limiting conditions could also be offset over time

by positive effects on overall performance or fitness that occur at times when carbon is not

limiting (e.g., enhanced growth, photosynthetic rate, nutrient acquisition) [42,43]. Studying all the

factors that influence costs and benefits of EM networks at different spatiotemporal scales is

critical to predict when and where they might have positive or negative effects under future

climates [44]. While studies often assess costs and benefits of mycorrhizal associations based on

whether they increase or decrease plant growth [45], growth is only one of several functions

requiring NSC and may not reflect costs and benefits for other vital plant functions, especially

under stressful conditions that inhibit growth. Using a carbon balance approach, we can integrate

all the physiological processes that require carbon (e.g., growth, defense, osmoregulation,

reproduction) and provide a more comprehensive cost-benefit assessment.

While our novel results highlight a critical role of stored NSC on plant water relations and a

potentially delicate balance between costs and benefits of EM networks under carbon stress,

much research is still needed. For instance, we need to elucidate the specific mechanisms

(including the involved organic solutes) by which stored NSC depletion decreases osmoregulation,

and how these mechanisms vary across species and climate gradients. We also need a better

understanding of the potential effect of plant-fungal interactions on NSC depletion and plant

water relations in complex EM networks under climate change. For instance, does individual-level

(as opposed to stand level) mortality caused by low intensity fire, drought or insects cause rippling

effects of similar magnitude to the ones we document when multiple trees are interconnected?

We hope that our research motivates such work.

Methods

Experimental Design

Ponderosa pine is one of the most widely distributed species in North America and has

been extensively used as a representative gymnosperm in ecological, physiological, and forestry

studies. In the spring of 2016, we planted eighty, one-year-old ponderosa pine seedlings (source:

Zone IV-V block of the Missoula Ponderosa Pine Seed Orchard, Department of Natural Resources)

in forty 19 L pots at the University of Montana greenhouse. Each pot contained two seedlings. We

used a soil mixture consisting of 40% sand, 30% topsoil, and 30% peat moss, which was kept at

field capacity throughout the duration of the experiment to avoid water deficit. Seedlings came

already inoculated with ectomycorrhizal Pezizales fungi from source. However, we also inoculated

the rhizosphere of each seedling with a mixture of *Rhizopogon* spores (Mycorrhizal Applications.

Grants Pass, OR) to ensure colonization. Both *Rhizopogon* and Pezizales form ectomycorrhizal

structures that potentially redistribute resources among plants [20,46]. Half of the pots contained a

stainless-steel mesh barrier that separated seedlings (139.7 μm wire diameter and 177.8 μm pore

diameter) which was penetrable by hyphae but not roots. The remaining pots contained a

Plexiglass barrier to prevent both root and ectomycorrhizal connections between seedlings to

create a control treatment and observe patterns in the absence of ectomycorrhizal networks. Both

barrier types were secured using silicone sealant. After inoculation, seedlings grew unperturbed

for 47 weeks and established mature ectomycorrhizal associations (Figure S1 and S2). Despite our

efforts, roots in pots with plexiglass barriers pierced the silicon sealant to form ectomycorrhizal

networks. While this precluded a treatment without EM networks, our main goal was to assess

how NSC depletion of neighbor seedlings influenced NSC pools and water relations in non-

stressed, connected hosts. Because all key variables measured did not differ between pots with

and without root connections (Table S1), the two barrier treatments were pooled for statistical

analyses. All subsequent treatments described below were applied uniformly in each of the two

underground barrier treatments.

In the fall of 2017, we split pots from each treatment into two groups (NSC-depleted and

non-depleted) and applied a 3-week NSC depletion treatment to one group using light-blocking

covers that reduced photosynthetically photon flux density below compensation point (Fig. 1,

Figure S3). In each pot within the NSC-depleted treatment, we placed a cover over one of the two

seedlings (dark, D) to inhibit photosynthesis while the other seedling (light paired with dark, LD)

was left undisturbed. Each cover consisted of a wire scaffolding (20 x 40 cm) overlaid with

aluminum foil that blocked incoming light while keeping air temperatures similar to those in non-

covered neighbor seedlings (ca. 24 ºC). We pierced 5 mm diameter holes evenly across the cover

walls which facilitated air flow, kept the atmosphere around the plant unsaturated at less than

50%, and allowed canopy transpiration. Relative humidity was measured using a Vaisala HMP35C

sensor (Vantaa, Finland). We maximized ventilation inside the covers while maintaining PAR below

0.40 µmol quanta m^{-2} s^{-1} which is far below compensation point (Figure S4). We measured PAR

with a Licor LI-190SA Quantum sensor (Lincoln, NE, USA). Combining shading with ectomycorrhizal

networks allowed us to alter the overall carbon budget of a networked system and indirectly

modify NSC pools in non-shaded plants. This procedure generated non-depleted networks with

both connected seedlings in the light (light, LL) and NSC-depleted networks with one seedling in

the light (LD) connected to one seedling in the dark (D). After three weeks of light deprivation, we

randomly selected plants from ten pots (five from each underground barrier treatment) in both

NSC-depleted and non-depleted treatments to assess physiological status (see below) of LL, LD,

and D seedlings and harvested them to establish base-line $^{13}C/^{12}C$ ratios ($\delta^{13}C$) in all organs across

seedling types. We then labeled with ^{13}C (see below) the remaining LD seedlings from NSC-

depleted networks and one of the two LL seedlings (chosen at random) from the non-depleted

NSC networks to assess potential carbon transfer between individuals in the presence and

absence, respectively, of carbon-limiting conditions seven days later. Comparing carbon transfer

between non-depleted and NSC-depleted networks let us assess how carbon transfer responds to

carbon limitation. One week after the labeling, we re-assessed physiological status and harvested

the remaining seedlings in both treatments to measure changes in $\delta^{13}C$ across all organs (Fig. 1).

<u>Isotopic Labeling and Carbon Transfer Variables</u>

Labeling was performed following similar methods to Song *et al.* (2015). We enclosed the

entire canopy of each labeled plant in a clear 2 L plastic bag. Each bag was injected with 20 mL of

Carbon-^{13}C dioxide (99 atom % 13C, <3 atom % 18O, Sigma Aldrich, (St. Louis, MO, USA)) for a ratio

of 10 mL of $^{13}C \cdot L^{-1}$ of air and was left undisturbed for 2 hours. ^{13}C injections were performed at

midday over two consecutive days. After each labeling event, enriched air was removed from bags

and directed outside the greenhouse facilities using an industrial vacuum, thus preventing

contamination of neighboring seedlings. Additionally, extra seedlings were randomly interspersed

throughout the treatments and their needles were used to assess possible contamination during

labeling. No signs of increased δ^{13}C were observed in these seedlings. Ground samples of dry

needles, stems, and roots of all remaining seedlings (labeled or not) were sent to the Stable

Isotope Facility at University of California, Davis and analyzed to obtain δ^{13}C values. We also

collected a representative amount of fungal material from ectomycorrhizal root tips of LL, LD, and

D seedlings (5 samples) before the labeling event to obtain base-line δ^{13}C fungal values as well as

several labelled LL (4) and LD (3) seedlings, and non-labelled LL (9), and D (10) seedlings after

labeling and analyzed their δ^{13}C. Fungal samples were collected to ensure that the applied ^{13}C label

traveled from needles to fungal tissue on the labeled side as well as to fungal tissue of the paired,

non-labeled plants connected through the ectomycorrhizal network. These descriptive

measurements served as an extra validation of the existence of an ectomycorrhizal network

between seedling pairs already observed at the time of harvest (Figure S1).

<u>Measurements of water relations</u>

After re-acclimating D seedlings to light for 2 hours (see Figure S5, panel F), we measured

assimilation, stomatal conductance, and respiration in each seedling with a Licor 6400XT (Lincoln,

NE, USA). We then measured midday leaf and stem water potential (WP) (i.e., xylem tension due

to water deficit) immediately before harvesting (Fig. 1) using a pressure chamber (PMS Instrument

Company, (Corvallis, OR, USA)) following methods in Kaufmann (1968). Stem WP was estimated by

equilibrating the WP of a needle bundle with that of the stem following methods from Begg &

Turner (1970) and measuring the equilibrated bundle as above. Subsequently, we harvested

seedlings and collected tissue samples (needles, stem, and roots) for osmotic potential

measurements (i.e., water retention capacity via solutes potentially derived from NSC). Tissue

samples were wrapped in aluminum foil to prevent artificial changes in osmotic potential due to

water loss or photosynthetic activity, placed in small Ziploc bags, frozen in dry ice and transported

to the lab within 2 hours. The rest of the seedling was placed in a Ziploc bag containing a wet

paper towel partially covered in tin foil to prevent desiccation without introducing external

moisture into the tissues in contact with the towel [32], placed in a cooler, and transported to the

lab. There, we measured relative water content (RWC) and hydraulic conductivity (i.e., capacity to

transport water [49]). We describe the methods used to measure water content and hydraulic

conductivity in Methods S1. We also used the rest of the seedling to measure plant NSC pools (see

below). Upon arrival to the lab, samples collected for osmotic potential were pressed to extract

cellular liquid contents and the expressed solution was used to saturate 28 mm^2 filter paper disks

following methods from Grange (1983) [50]. This method avoids potential osmotic artifacts due to

starch mobilization or low sample water content. Disks were then placed in a C-52 sample

chamber attached to a Psypro data logger (Wescor, Inc. Logan, UT, USA) to measure osmotic

potential. Finally, pressure potential (i.e., positive pressure exerted by cells against cell walls when

turgid [12]) was calculated in stems and needles as the difference between their respective water

and osmotic potential. Pressure potential was not calculated in roots because we lacked root WP

measurements.

<u>Non-structural carbohydrates</u>

Non-structural carbohydrates were analyzed in all tissue types (needles, stems, and roots)

collected at harvest. A sample of each tissue was microwaved for 180 seconds at 900 Watts in

three cycles of 60 seconds to stop metabolic consumption of NSC pools, and oven-dried at 70 $^\circ$C.

Samples were dried to a constant mass and finely ground into a homogenous powder.

Approximately 11 mg of needle tissue and 13 mg of stem or root tissue were used to analyze NSC

dry mass content following the procedures and enzymatic digestion method from Hoch, Popp &

Körner (2002) and Galiano, Martínez-Vilalta, Sabaté & Lloret (2012). NSC, starch, sucrose, and

glucose + fructose tissue concentrations were then multiplied by their respective tissue fraction

(the dry mass of a tissue relative to whole-plant dry mass) to obtain whole-plant concentrations.

Whole-plant dry mass for each seedling was calculated by combining the dry mass of all samples

and the remaining biomass. These methods are described in Methods S1.

<u>Statistical Analyses</u>

We observed no differences between pots with (Plexiglas barrier) and without (mesh

barrier) root connections for all key variables measured (Table S1), and data from the two barrier

treatments were pooled. We tested differences among seedling types (LL, LD, D) in all variables

using ANOVA and Tukey's Honestly Significant Difference tests. Note that sample size of LL

seedlings is twice that of LD and D seedlings because both plants in non-depleted networks lack of

covers and are LL seedlings (Fig. 1). However, because treatment comparisons excluding a random

half of the LL seedlings yielded the same results, we decided not to exclude any data. To test

differences in NSC depletion, we compared total NSC concentration and concentrations of each

NSC component (i.e., starch, sucrose, and glucose and fructose together) among seedling types at

both organ and whole-plant levels. To test the effectiveness of the labeling process and assess

carbon translocation among organs, for each organ and treatment, we compared δ^{13}C values after

labelling to those prior to labeling. Note that D seedlings were excluded because they were never

labeled. We only analyzed a few fungal samples for δ^{13}C (see above) as an additional check that

the ^{13}C label could travel from the labelled seedling to the fungal network of the non-labelled

seedling, thus ensuring that plants were indeed connected via fungal networks. Likewise, to assess

potential carbon transfer to non-labeled seedlings, for each organ and treatment, we compared

δ^{13}C values after labeling the paired seedling to those before labelling. LD seedlings were excluded

because they were all labeled. We also tested differences in hydraulic conductivity, WP, osmotic

potential, pressure potential and relative water content in each organ for which we collected data

among seedling types.

To assess both direct and indirect effects of NSC depletion on water relations, we assessed

whether osmotic potential and pressure potential changed in response to NSC depletion. We used

two linear regressions with needle % NSC deviation from control as predictor and either leaf

osmotic or pressure potential as response variables. We focused on needles because they are the

organ with most living cells exposed to dry conditions and, therefore, most susceptible to turgor

loss. Finally, we assessed whether NSC depletion influences turgor loss as WP decreases using

linear regressions. Leaf pressure potential was the response variable and the interaction between

leaf WP and whole-plant % NSC deviation from control as predictor. In this analysis, we focus only

on differences in intercepts among NSC levels (i.e. NSC effect on leaf WP at turgor loss point, a

broadly used indicator of drought tolerance) and among their slopes (i.e. rate of turgor loss per

unit change in water potential) because a significant relationship between leaf pressure potential

and leaf WP is expected given that the former is calculated from the later.

Acknowledgements

This work was partially supported by a National Science Foundation grant to AS (BCS

1461576). GS received funding from the NSF Experimental Program to Stimulate Competitive

Research (EPSCoR) Track-1 EPS-1101342 (INSTEP 3). PD received funding from NSF EPSCoR RII

Track 1 award number IIA-1443108. YL is grateful to MPG Ranch for funding. The authors thank

Laura Thornton and Maria Pilar de Moreta for their help collecting data, and Mauri Valett and Lila

Fishman for letting us use their facilities. The authors also thank D. Stanton, R. Montgomery, R.

Koide, and D. Ulrich for their comments in early versions of this manuscript.

References

1.   Stocker, T. F. *et al.* Summary for Policymakers. In: Climate Change 2013: The Physical

Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the

Intergovernmental Panel on Climate Change. *CEUR Workshop Proc.* **1542**, 33–36 (2015).

2.   Meir, P., Mencuccini, M. & Dewar, R. C. Drought-related tree mortality: addressing the gaps

in understanding and prediction. *New Phytol.* **207**, 28–33 (2015).

3.   Hartmann, H. *et al.* Research frontiers for improving our understanding of drought-induced

tree and forest mortality. *New Phytol.* **218**, 15–28 (2018).

4.   Adams, H. D. *et al.* A multi-species synthesis of physiological mechanisms in drought-

induced tree mortality. *Nat. Ecol. Evol.* **1**, 1285–1291 (2017).

5.   Sala, A., Woodruff, D. R. & Meinzer, F. C. Carbon dynamics in trees: feast or famine? *Tree*

Physiol. **32**, 764–75 (2012).

6.   Sapes, G. *et al.* Plant water content integrates hydraulics and carbon depletion to predict

drought-induced seedling mortality. *Tree Physiol.* (2019). doi:10.1093/treephys/tpz062

7.   Martinez-Vilalta, J., Anderegg, W. R. L., Sapes, G. & Sala, A. Greater focus on water pools

may improve our ability to understand and anticipate drought-induced mortality in plants.

New Phytol. **223**, 22–32 (2019).

8.   Marshall, J. D. Drought and shade interact to cause fine-root mortality in Douglas-fir

seedlings. *Plant Soil* **91**, 51–60 (1986).

9.   Sevanto, S., McDowell, N. G., Dickman, L. T., Pangle, R. & Pockman, W. T. How do trees die?

A test of the hydraulic failure and carbon starvation hypotheses. *Plant. Cell Environ.* **37**,

153–61 (2014).

10.   Hartmann, H. & Trumbore, S. Understanding the roles of nonstructural carbohydrates in

forest trees - from what we can measure to what we want to know. *New Phytol.* (2016).

doi:10.1111/nph.13955

11.   Nehls, U., Grunze, N., Willmann, M., Reich, M. & Küster, H. Sugar for my honey:

Carbohydrate partitioning in ectomycorrhizal symbiosis. *Phytochemistry* **68**, 82–91 (2007).

12.   Bartlett, M. K., Scoffoni, C. & Sack, L. The determinants of leaf turgor loss point and

prediction of drought tolerance of species and biomes: a global meta-analysis. *Ecol. Lett.*

15, 393–405 (2012).

13.   Salleo, S., Trifilò, P., Esposito, S., Nardini, A. & Lo Gullo, M. A. Starch-to-sugar conversion in

wood parenchyma of field-growing Laurus nobilis plants: a component of the signal

pathway for embolism repair? *Funct. Plant Biol.* **36**, 815–825 (2009).

14.   Dietze, M. C. *et al.* Nonstructural carbon in woody plants. *Annu. Rev. Plant Biol.* **65**, 667–

687 (2014).

15.   Martinez-Vilalta, J. *et al.* Dynamics of non-structural carbohydrates in terrestrial plants: A

global synthesis. *Ecol. Monogr.* **86**, 495–516 (2016).

16.   O'Brien, M. J., Leuzinger, S., Philipson, C. D., Tay, J. & Hector, A. Drought survival of tropical

tree seedlings enhanced by non-structural carbohydrate levels. *Nat. Clim. Chang.* **4**, 1–5

(2014).

17.   Secchi, F. & Zwieniecki, M. A. Accumulation of sugars in the xylem apoplast observed under

water stress conditions is controlled by xylem pH. *Plant Cell Environ.* **39**, 2350–2360 (2016).

18.   Sala, A. & Mencuccini, M. Plump trees win under drought. *Nat. Clim. Chang.* **4**, 666–667

(2014).

19.   Warren, J. M., Brooks, J. R., Meinzer, F. C. & Eberhart, J. L. Hydraulic redistribution of water

488   from Pinus ponderosa trees to seedlings: evidence for an ectomycorrhizal pathway. *New*

489   *Phytol.* **178**, 382–394 (2008).

490 20. Song, Y. Y., Simard, S. W., Carroll, A., Mohn, W. W. & Zeng, R. Sen. Defoliation of interior

491   Douglas-fir elicits carbon transfer and stress signalling to ponderosa pine neighbors through

492   ectomycorrhizal networks. *Sci. Rep.* **5**, 8495 (2015).

493 21. He, X., Xu, M., Qiu, G. Y. & Zhou, J. Use of15N stable isotope to quantify nitrogen transfer

494   between mycorrhizal plants. *J. Plant Ecol.* **2**, 107–118 (2009).

495 22. van der Heijden, M. G. & Horton, T. R. Socialism in Soil? the Importance of Mycorrhizal

496   Fungal Networks for Facilitation in Natural Ecosystems. *J. Ecol.* **97**, 1139–1150 (2009).

497 23. Tedersoo, L., May, T. W. & Smith, M. E. Ectomycorrhizal lifestyle in fungi : global diversity ,

498   distribution , and evolution of phylogenetic lineages. 217–263 (2010). doi:10.1007/s00572-

499   009-0274-x

500 24. Selosse, M.-A., Richard, F., He, X. & Simard, S. W. Mycorrhizal networks: des liaisons

501   dangereuses? *Trends Ecol. Evol.* **21**, 621–628 (2006).

502 25. Simard, S. W. & Perry, D. a. Net transfer of carbon between ectomycorrhizal tree species in

503   the field. *Nature* **388**, 579 (1997).

504 26. Klein, T., Siegwolf, R. T. W. & Korner, C. Belowground carbon trade among tall trees in a

505   temperate forest. *Science (80-.).* **352**, 342–344 (2016).

506 27. Koide, R. T. & Fernandez, C. W. The continuing relevance of 'older' mycorrhiza literature:

507   insights from the work of John Laker Harley (1911–1990). *Mycorrhiza* **28**, 577–586 (2018).

508 28. Simeone, C. *et al. Coupled ecohydrology and plant hydraulics modeling predicts ponderosa*

509   *pine seedling mortality and lower treeline in the US Northern Rocky Mountains*. *New*

510   *Phytologist* (2018). doi:10.1111/nph.15499

511   29.   Chaves, M. Effects of Water Deficits on Carbon Assimilation. *J. Exp. Bot.* **42**, 1–16 (1991).

512   30.   Thalmann, M. *et al.* Regulation of Leaf Starch Degradation by Abscisic Acid Is Important for

513   Osmotic Stress Tolerance in Plants. *Plant Cell* **28**, 1860–1878 (2016).

514   31.   Wang, Z., Quebedeaux, B. & Stutte, W. Osmotic Adjustment: Effect of Water Stress on

515   Carbohydrates in Leaves, Stems and Roots of Apple. *Aust. J. Plant Physiol.* **22**, 747–54

516   (1995).

517   32.   Garcia-Forner, N., Sala, A., Biel, C., Savé, R. & Martínez-Vilalta, J. Individual traits as

518   determinants of time to death under extreme drought in Pinus sylvestris L. *Tree Physiol.* **36**,

519   1196–1209 (2016).

520   33.   Salmon, Y. *et al.* Leaf carbon and water status control stomatal and non-stomatal

521   limitations of photosynthesis in trees. *New Phytol.* (2019). doi:10.1111/nph.16436

522   34.   Trueba, S. *et al.* Thresholds for leaf damage due to dehydration: declines of hydraulic

523   function, stomatal conductance and cellular integrity precede those for photochemistry.

524   *New Phytol.* **223**, 134–149 (2019).

525   35.   Grman, E. Plant species differ in their ability to reduce allocation to non-beneficial

526   arbuscular mycorrhizal fungi. *Ecology* **93**, 711–718 (2012).

527   36.   Nehls, U. Mastering ectomycorrhizal symbiosis: the impact of carbohydrates. *J. Exp. Biol.*

528   **59**, 1097–1108 (2008).

529   37.   Ulrich, D. E. M. *et al.* Plant-microbe interactions before drought influence plant

530   physiological responses to subsequent severe drought. *Sci. Rep.* **9**, 1–10 (2019).

38. Franklin, O., Näsholm, T., Högberg, P. & Högberg, M. Forests trapped in nitrogen limitation

– an ecological market perspective on ectomycorrhizal symbiosis. *New Phytol.* **203**, 657–

666 (2014).

39. Schellenbaum, L., Muller, J., Boller, T., Wiemken, A. & Schuepp, H. Effects of drought on

non-mycorrhizal and mycorrhizal maize: changes in the pools of non-structural

carbohydrates, in the activities of invertase and trehalase, and in the pools of amino acids

and imino acids. *New Phytol.* **138**, 59–66 (1998).

40. Bingham, M. a. & Simard, S. Ectomycorrhizal Networks of Pseudotsuga menziesii var. glauca

Trees Facilitate Establishment of Conspecific Seedlings Under Drought. *Ecosystems* **15**, 188–

199 (2012).

41. Beiler, K. J., Simard, S. W. & Durall, D. M. Topology of tree–mycorrhizal fungus interaction

networks in xeric and mesic Douglas-fir forests. *J. Ecol.* **103**, 616–628 (2015).

42. Thomson, B. D., Grove, T. S., Malajczuk, N. & Hardy, G. The Effectiveness of Ectomycorrhizal

Fungi in Increasing the Growth of Eucalyptus-Globulus Labill in Relation To Root

Colonization and Hyphal Development in Soil. *New Phytol.* **126**, 517–524 (1994).

43. Nardini, A., Salleo, S., Tyree, M. T. & Vertovec, M. Influence of the ectomycorrhizas formed

by Tuber melanosporum Vitt . on hydraulic conductance and water relations of Quercus

ilex L . seedlings. *Ann. For. Sci.* **57**, 305–312 (2000).

44. Hasselquist, N. *et al.* Greater carbon allocation to mycorrhizal fungi reduces tree nitrogen

uptake in a boreal forest. *Ecology* **97**, 1012–1022 (2016).

45. Morgan, J. A. W., Bending, G. D. & White, P. J. Biological costs and benefits to plant–

microbe interactions in the rhizosphere. *J. Exp. Bot.* **56**, 1729–1739 (2005).

46.   Beiler, K. J., Durall, D. M., Simard, S. W., Maxwell, S. A. & Kretzer, A. M. Architecture of the

wood-wide web: Rhizopogon spp. genets link multiple Douglas-fir cohorts. *New Phytol.* **185**,

543–553 (2010).

47.   Kaufmann, M. Evaluation of the Pressure Chamber Technique for Estimating Plant Water

Potential of Forest Tree Species. *For. Sci.* **14**, 369–374 (1968).

48.   Begg, J. E. & Turner, N. C. Water Potential Gradients in Field Tobacco. *PLANT Physiol.*

(1970). doi:10.1104/pp.46.2.343

49.   Tyree, M. T. & Sperry, J. S. Vulnerability of Xylem to Cavitation and Embolism. *Ann. Rev.*

Plant. Phys. Mol. Bio. **40**, 19–38 (1989).

50.   Grange, R. I. Solute Production during the Measurement of Solute Potential on Disrupted

Tissue. *J. Exp. Bot.* **34**, 757–764 (1983).

51.   Galiano, L., Martínez-Vilalta, J., Sabaté, S. & Lloret, F. Determinants of drought effects on

crown condition and their relationship with depletion of carbon reserves in a

Mediterranean holm oak forest. *Tree Physiol.* **32**, 478–489 (2012).

52.   Hoch, G., Popp, M. & Körner, C. Altitudinal increase of mobile carbon pools in Pinus cembra

suggests sink limitation of growth at the Swiss treeline. *Oikos* (2002). doi:10.1034/j.1600-

0706.2002.980301.x

Figure legends

Fig. 1. Experimental design. Circles represent pots with barriers and a seedling on each side.

Colors represent LL seedlings exposed to natural light (golden), D seedlings with light-blocking

covers (black), and LD seedlings exposed to natural light paired with plants with covers (teal

border). Numbers on the side of each circle indicate the sample size of the seedling type

represented in that side of the pot at a given point on time. Red 13C annotations indicate pots

labeled with ^{13}C. Arrows indicate harvesting events.

Fig. 2. Stored NSC decreased in seedlings of the NSC-depletion treatment. Panels correspond to

whole-plant concentrations in percent dry weight. Colors represent LL seedlings exposed to

natural light (golden), D seedlings with light-blocking covers (black), and LD seedlings exposed to

natural light paired with plants with covers (teal). Lines within boxes represent the median and top

and bottom hinges represent 25th and 75th percentiles. Whiskers indicate highest and lowest

value no further than 1.5 times the inter-quartile range represented by the hinges. Dots represent

the distribution of the data. Asterisks indicate the degree of significance between groups (* = 0.05,

** = 0.01, *** = < 0.001).

 Fig. 3. ^{13}C isotope reached all organs in labeled plants and the ectomycorrhizal network but was

not transferred to NSC-depleted hosts. The area of the yellow circles represents the average

change in ^{13}C/^{12}C ratios in each tissue as a result from labeling (mean tissue δ^{13}C after labeling –

mean tissue δ^{13}C before labeling) in parts per thousand (‰; also, annotated besides each circle).

Positive values indicate an increase in δ^{13}C. See Figure S7 and S8 for raw δ^{13}C values for each

tissue and labeling time, confidence intervals, and statistical significances.

Fig. 4. All organs in NSC-depleted seedlings exhibited an increase of osmotic potential and a decrease in turgor. Panels on the left correspond to osmotic potentials of A) needles, B) stems, and C) roots. Panels on the right correspond to pressure potentials of D) needles and E) stems. Colors represent LL seedlings exposed to natural light (golden), D seedlings with light-blocking covers (black), and LD seedlings exposed to natural light paired with plants with covers (teal). Lines within boxes represent the median and top and bottom hinges represent 25th and 75th percentiles. Whiskers indicate highest and lowest value no further than 1.5 times the inter-quartile

605   range represented by the hinges. Dots represent the distribution of the data. Asterisks indicate the

606   degree of significance between groups (* = 0.05, ** = 0.01, *** = < 0.001).

Fig. 5. Water relations are impaired when NSC drop below values corresponding to non-stressed

plants. Panels correspond to A) osmotic potential and B) turgor pressure. All seedling types are

merged for this analysis. Vertical golden lines indicate average leaf NSC content in non-depleted

pots (i.e., light plants). A loess function was fit to the data to best represent the relationship

between variables. Dashed lines indicate 95% confidence interval of the regression lines.

Fig. 6. NSC depletion is associated with turgor loss at high water potentials. Colors represent LL

seedlings exposed to natural light (golden), D seedlings with light-blocking covers (black), and LD

seedlings exposed to natural light paired with plants with covers (teal). Vertical red line and points

indicate differences in turgor at a given water potential. Horizontal line indicates turgor loss point.

Dashed lines indicate 95% confidence interval of the regression lines.