Title
CSDE1 is a Post-Transcriptional Regulator of the LDL Receptor

Authors
Geoffrey A. Smith, MD PhD[†,1,2]; Arun Padmanabhan, MD PhD[†,3,4,5]; Bryan H. Lau, BS[6]; Akhil Pampana, MS[7,8]; Li Li, BS[9]; Y. Clara Lee, BA[5]; Angelo Pelonero, MS[5]; Tomohiro Nishino, MD PhD[5]; Nandhini Sadagopan, MS[3,4,5]; Rajan Jain, MD[9,10]; Pradeep Natarajan, MD[7,8,11]; Roland S. Wu, MD[3,4,6,12]; Brian L. Black, PhD[6]; Deepak Srivastava, MD[5,13,14]; Kevan M. Shokat, PhD[1,15]; John S. Chorba, MD*[,4,16]

[†]Drs. Smith and Padmanabhan contributed equally.
*Corresponding Author

Author Affiliations
[1]Department of Cellular and Molecular Pharmacology, University of California, San Francisco CA
[2]Present Affiliation: Boston Children's Hospital, Boston MA
[3]Division of Cardiology, UCSF Health, San Francisco CA
[4]Department of Medicine, University of California, San Francisco CA
[5]Gladstone Institute of Cardiovascular Disease, San Francisco CA
[6]Cardiovascular Research Institute, University of California San Francisco
[7]Cardiovascular Research Center, Massachusetts General Hospital, Boston MA
[8]Program in Medical and Population Genetics, Broad Institute of Harvard and MIT, Cambridge MA
[9]Department of Medicine and Penn Cardiovascular Institute, University of Pennsylvania, Philadelphia PA
[10]Department of Cell and Developmental Biology, Institute of Regenerative Medicine, and Penn Epigenetics Institute, University of Pennsylvania, Philadelphia PA
[11]Department of Medicine, Harvard Medical School, Boston MA
[12]Present Affiliation: Amgen, South San Francisco CA
[13]Departments of Pediatrics, and Biochemistry and Biophysics, University of California, San Francisco CA
[14]Roddenberry Center for Stem Cell Biology and Medicine at Gladstone, San Francisco CA
[15]Howard Hughes Medical Institute, University of California, San Francisco CA
[16]Division of Cardiology, Zuckerberg San Francisco General Hospital, San Francisco CA

Corresponding Author
John S. Chorba MD
600 16th St
Genentech Hall N514, MC2280
San Francisco CA 94143
phone: 415-601-7361
email: john.chorba@ucsf.edu

Abstract

The low-density lipoprotein receptor (LDLR) controls cellular delivery of cholesterol and clears LDL from the bloodstream, protecting against atherosclerotic heart disease, the leading cause of death in the United States. We therefore sought to identify regulators of the LDLR beyond the targets of current clinical therapies and known causes of familial hypercholesterolemia. We show that Cold Shock Domain-Containing Protein E1 (CSDE1) enhances hepatic *LDLR* mRNA decay via its 3' untranslated region to regulate atherogenic lipoproteins *in vivo*. Using parallel phenotypic genome-wide CRISPR interference screens, we found 40 specific regulators of the LDLR left unidentified by observational human genetics. Among these, we show that CSDE1 regulates the LDLR at least as strongly as the mechanistically distinct pathways exploited by the best available clinical therapies: statins and PCSK9 inhibitors. Additionally, we show that hepatic gene silencing of *Csde1* treats diet-induced dyslipidemia in mice better than that of *Pcsk9*. Our results reveal the therapeutic potential of manipulating a newly identified key factor in the post-transcriptional regulation of the *LDLR* mRNA for the prevention of cardiovascular disease. We anticipate that our approach of modelling a clinically relevant phenotype in a forward genetic screen, followed by mechanistic pharmacologic dissection and *in vivo* validation, will serve as a generalizable template for the identification of therapeutic targets in other human disease states.

Keywords

Atherosclerosis, cholesterol metabolism, CRISPR interference, CSDE1, genome-wide CRISPR screen, post-transcriptional regulation, LDL receptor, mRNA decay, pharmacogenetics, transferrin receptor

One Sentence Summary

A genome-wide CRISPR screen identifies CSDE1 as a key regulator of hepatic *LDLR* mRNA decay *in vivo*, making it a promising target for heart disease.

73   **Graphical Abstract**
74

75
76
77

Introduction

Introduction
The low-density lipoprotein receptor (LDLR) delivers cholesterol from low-density lipoprotein
(LDL) to cells to maintain membrane homeostasis *(1)*. By clearing atherogenic LDL from the
bloodstream, the hepatic LDLR protects against atherosclerotic heart disease *(2, 3)*. Despite
successful therapies that upregulate the hepatic LDLR and reduce heart attacks *(4)*,
cardiovascular disease remains the leading cause of death in Western countries *(5)*. Lowering
LDL beyond the levels achieved by HMG-CoA reductase inhibitors (statins) improves clinical
outcomes without adverse effects *(6, 7)*. Though there is a theoretical level at which LDL could
get too low *(8)*, this has yet to be discovered in large randomized trials *(9, 10)*. Whether other
LDLR regulatory mechanisms could be leveraged to further treat heart disease remains
unknown.
89
The genetics of familial hypercholesterolemia (FH), which manifests as an isolated elevation in
serum LDL, underlies the clinical success of LDLR upregulation by statins and PCSK9
inhibitors. Estimates suggest that 20-40% of FH phenotypes remain unexplained outside of the
four major causes: *LDLR*, *APOB*, *PCSK9*, and *LDLRAP1 (11, 12)*. This implies that additional
regulators of the LDLR exist. Advances in forward genetics *(13–15)* can now enable searches
for tissue and disease-specific effects across the entire genome that may elude the sporadic
natural variants found in observational studies, which themselves require compatibility
throughout the entire lifespan and in all cell types. Moreover, hepatic delivery of gene silencing
agents is effective in the clinic *(16)*, providing a therapeutic modality against hits whose
phenotypes are driven by expression in the liver. We therefore employed a genome-wide
CRISPR interference screen for factors involved in hepatic LDLR regulation, both to understand
the biology of this important receptor and to uncover potential therapeutic targets in
cardiovascular disease.
103

Results

Results
A Genome-Wide CRISPR Interference Screen for LDL Receptor Regulation
We engineered the HepG2 cell line, which models the regulation of the LDLR *(17)*, to
constitutively express a dCas9-KRAB fusion protein, enabling the knockdown of any given gene
with an appropriate sgRNA (Fig. 1A) *(13, 14, 18)*. Because statins *(19, 20)* and PCSK9
inhibitors *(21–23)* increase cell surface LDLR, we scored surface LDLR levels. To focus on
factors that preferentially affect the LDLR over other receptors, we performed a parallel screen
for regulators of the transferrin receptor (TFR). This critical player in iron metabolism shares a
clathrin-mediated intake mechanism, but is otherwise orthogonally regulated from the LDLR *(24,
25)*. Prior to our screen, we confirmed both dCas9-KRAB activity and an appropriate dynamic
range for both LDLR and TFR regulation by transduction with sgRNAs expected to alter
receptor levels in either direction (Fig. S1) *(26, 27)*.
116
We next performed our pooled screens in parallel by transducing a library encoding sgRNAs
with 5-fold coverage of the entire protein-coding human genome *(14)*. We then selected the
cells at the upper and lower third of receptor abundance by FACS and quantified the sgRNAs
for each population via deep sequencing (Fig. S2, Tables S1-S4). We compared the degree of
enrichment of LDLR or TFR surface levels in the high abundance to the low abundance cells
(defined as ρ, Fig. 1B). We also compared the high and low receptor abundance cells to the
unsorted population (defined as τ or γ, respectively) and included these results in our final hit
count. This resulted in 130 total hits for the LDLR and 186 hits for the TFR (Tables S5-S6). We
hypothesized that hits with shared phenotypes would likely have global effects on surface
receptors, leaving us with 117 hits specific for LDLR regulation (Fig. 1C, Table 1). Gene
ontology (GO) analysis *(28)* revealed a 15-fold enrichment for cholesterol metabolism as a
biologic process (11 total hits, $p = 5.7 \times 10^{-10}$), providing confidence that we recapitulated our

target biology. The hits also included 48 members of potentially druggable protein classes,
including 29 with proposed enzymatic activity, and 22 hits were unclassified in GO databases
(Fig. S3A).
132
Cross-referencing Human Genetic Datasets Identifies LDLR Regulators in vivo
We next compared genes associated with serum LDL cholesterol (LDL-C) from published
genome-wide association studies (GWAS) *(29–31)* to our list of hits. Intriguingly, only 13 of
these genes overlapped with our results (Fig. 1D), even when we relaxed our threshold for hit
selection. To improve power for multiple hypothesis testing across the entire genome, we turned
to 390,375 UK Biobank participants with genome-wide genotypes and known plasma lipids
(Table S7) to search for variants associated with LDL-C amongst only our hits *(32)*. We filtered
to nonsynonymous protein coding variants in these hits by a threshold minor allele frequency
(>0.001) and minimum statistical significance (Table 2). For *BCAM*, we found both an
association between higher LDL-C and a nonsense variant, along with bidirectional associations
between LDL-C and missense variants, suggesting that this pathway may be tunable. We also
found associations between elevated LDL-C and variants in *MSMO1*, *C6orf132*, *HNF4A*, and
TIMELESS, suggesting that these hits may be functional in the human and warrant further
evaluation. The results also suggest that the accessible "genomic space" of the CRISPRi and
GWAS strategies is only partially overlapping.
148
Regulators of Surface LDL Receptor Abundance Affect Functional Uptake of LDL
To validate our screen results, we generated CRISPRi HepG2 cells harboring either of the two
top-scoring sgRNAs for 77 of our hits as well as established controls. We preferentially tested
hits with an increase in surface LDLR upon inhibition, as well as those with potentially druggable
functions or lacking associated GO terms. Since surface receptor levels might not necessarily
correlate to increased function, we evaluated both LDLR and TFR surface phenotypes
alongside a functional assay of LDL uptake *(33)*. Lastly, as knockdowns could also cause
growth phenotypes, we assayed the number of cells surviving to FACS analysis as a proxy for
viability.
158
We recapitulated the phenotypes for receptor abundance for at least one of the guides in the
majority of the hits (55 genes, 71% of those tested, Table S8). Moreover, for 40 of these genes,
both sgRNAs independently validated, suggesting against an off-target effect. We visualized
these hits based on their effects, at single cell resolution, on LDLR and TFR levels, the
LDLR/TFR ratio, functional LDL uptake and number of cells surviving to analysis (Figs. 2, S3B).
Notably, most knockdowns had independently validated effects on LDLR abundance and LDL
uptake of similar or greater magnitude than our *HMGCR* or *PCSK9* controls.
166
Knockdown of hits expected to alter cellular cholesterol balance or transcriptionally regulate the
LDLR showed directionally consistent effects between LDLR abundance and function (Fig. 2).
For genes in the enzymatic pathway of cholesterol metabolism *(34)* (*HMGCS1* and *MSMO1*),
this is consistent with activation of SREBP2-mediated *LDLR* transcription. For genes encoding
certain transcription factors (*HNF1A (35)*, *HNF4A (36)*, *ONECUT1 (37)*, and *ZEB1 (38)*), this is
consistent with an effect on *LDLR* transcription itself. Knockdowns of *SLC25A27*, which
encodes a mitochondrial uncoupling protein *(39)*, and *ABCA4*, encoding a known lipid
transporter *(40)*, both exhibited reductions in LDLR abundance and function (Fig. 2). These
genes could plausibly induce a negative lipid balance, increasing LDL uptake via both LDLR
dependent and independent mechanisms.
177
Targeting of hits that either affect multiple transcriptional pathways or regulate endocytosis
showed opposite effects on LDLR abundance and function. Knockdown of *TRIB1*, a GWAS hit

(29) encoding a pseudokinase that regulates the COP1 E3 ligase *(41, 42)* and affects multiple
transcription factors *(43)*, showed this phenotype. In the mouse, *TRIB1* overexpression lowers
serum cholesterol, while the knockout has the opposite effect *(44, 45)*, consistent with our
results. Knockdown of *AP2M1*, a TFR screen hit that encodes an adaptor protein required for
endocytosis *(46)*, was similar, consistent with an accumulation of non-functional receptors at the
cell surface. This phenotype, though specific to the LDLR, was also seen with knockdown of
BCAM, which encodes a membrane cell adhesion molecule *(47)* identified by GWAS *(31)*, and
TMEM217, which encodes an uncharacterized transmembrane protein (Figs. 2, S4). This
suggests that these proteins could have a similar endocytosis adaptor function specific for the
LDLR, akin to *LDLRAP1 (48)*, in which mutations cause a recessive form of FH.
190
Pharmacologic Inhibition of Clinically Relevant Pathways Provides Mechanistic Insight into
Putative LDLR Regulators
We next turned to pharmacologic approaches to perturb specific pathways of LDLR regulation.
We hypothesized that hits might alter cholesterol metabolism, LDLR recycling, or a yet
unspecified pathway. By combining CRISPRi knockdown with either a statin, to inhibit
endogenous cholesterol biosynthesis *(49, 50)*, or a PCSK9 inhibitor, to arrest LDLR lysosomal
degradation *(23)*, and assessing the combined effect, we inferred mechanistic information about
the target gene. Furthermore, we hypothesized that either additive or potentiating effects
between a clinically validated therapy and a hit gene might suggest promising therapeutic
targets.
201
We evaluated the receptor abundance and function phenotypes for 29 of our validated hits in
the presence or absence of a statin *(51)* or PF-846, a selective inhibitor of PCSK9 translation
(52, 53) (Fig. 3, Table S9). We calculated a synergy score by subtracting the differential effects
of CRISPRi knockdown, compared to the control, in the presence of compound from that with
the DMSO vehicle. A more positive value indicated synergy, and a more negative value
indicated antagonism.
208
Upon knockdown, regulators of cholesterol biosynthesis (*SREBF2*, *HMGCR*, *HMGCS1*,
MSMO1, and *PMVK*) showed antagonism with the statin, but mild synergy with PCSK9
inhibition (Fig. 3). This is consistent with the SREBP2-mediated *PCSK9* transcription that
underlies the clinical synergy between statins and PCSK9 inhibitors. The synergy phenotypes
for knockdown of *MRPL16*, which encodes a structural component of the mitochondrial
ribosome *(54)*, mirrored these cholesterol biosynthetic genes (Fig. 3), suggesting that MRP-L16
may play a role in the mitochondrial generation of metabolic precursors to sterol biogenesis. In
contrast, *C6orf132* knockdown showed the opposite phenotype: mild synergy with a statin, and
mild antagonism with PF-846 (Fig. 3). Given that C6orf132 localizes to the Golgi *(55)*, this
suggests it may function by facilitating LDLR delivery to the cell surface, prior to any interaction
with extracellular PCSK9. For some transcription factors, the synergy phenotypes can point to
their downstream targets. For example, synergy of *HNF1A* knockdown with a statin (Fig. 3) is
consistent with disruption of HNF1-α-mediated *PCSK9* transcription *(56)*.
222
CSDE1 Regulates the Stability of LDLR mRNA
One of our strongest hits, *CSDE1*, also known as upstream of N-ras (UNR), encodes an RNA
binding protein with varied regulatory functions *(57–59)*, including mRNA decay *(60)*. As the
LDLR 3' UTR consists of adenylate-uridylate (AU)-rich elements (AREs) implicated in mRNA
stability *(61)*, we hypothesized that CSDE1 could mediate the degradation of the *LDLR*
transcript, thereby explaining its observed receptor abundance, function, and synergy
phenotypes.
230

In the setting of *CSDE1* knockdown, we observed progressively higher LDLR abundance with
sterol depletion and a concomitant statin (Figs. 4A, S5A-C). This suggests the mechanism of
CSDE1 disruption is at least additive with SREBP2-mediated *LDLR* transcription and statin
therapy. We also observed that overexpression of isoform 1 of CSDE1, but not isoforms 2
through 4, reduced surface LDLR in HepG2 cells (Figs. 4B, S6A-D). Overexpression of all four
isoforms of CSDE1 downregulated LDLR levels in the *CSDE1* knockdown cells, though isoform
1 showed the strongest effect (Fig. S6E-I). The opposing directional effects of *CSDE1*
knockdown and overexpression suggest that, under physiologic expression conditions, isoform
1 of CSDE1 is a rate-limiting regulator of the LDLR.
240
Consistent with our mechanistic hypothesis, we noted over a 2-fold increase in steady-state
mRNA levels of *LDLR* (Fig. 4C), as well as depleted CSDE1 (Figs. 4C, S7), in the *CSDE1*
knockdown cells. Among control mRNA targets, we also observed significant increases in
MYLIP and *KHSRP* mRNA. These gene products downregulate the LDLR *(26, 62)*, which is the
opposite of our observed phenotype, suggesting that the direct effect of *CSDE1* knockdown on
the *LDLR* mRNA predominates. To specifically evaluate transcriptional decay, we treated cells
with actinomycin D and measured *LDLR* transcript levels over time. We observed significantly
higher *LDLR* mRNA in the *CSDE1* knockdown cells at all subsequent timepoints (Fig. 4D). The
mRNA half-life, modeled by a single-phase decay equation, was nearly 1.5-fold longer for the
CSDE1 knockdowns compared to controls (p = 0.0021, Fig. 4D). Notably, *CSDE1* knockdown
had no significant effect on *HMGCR*, *SREBF2*, *PCSK9*, or *TFRC* mRNA levels over time (Fig.
S8), suggesting that the effect on mRNA stability was isolated to *LDLR* among our tested
transcripts.
254
To probe the relationship of CSDE1 to the *LDLR* 3' UTR, we transiently expressed luciferase
constructs (Fig. 4E) under control of the native *LDLR* promoter in the *CSDE1* knockdown cells.
The luciferase-only constructs showed appropriate physiologic upregulation by sterols,
regardless of *CSDE1* knockdown (Fig. S9). Constructs fused to the *LDLR* 3' UTR, but not those
fused to the *LDLR* coding sequence alone, exhibited increased reporter activity with *CSDE1*
knockdown (Fig. 4F). Notably, this increase in activity was attenuated by removing the first of
four AREs *(61, 63)* from the 3' UTR (Fig. 4F). Activity of the 3' UTR-fused construct increased
further with statin coadministration (Fig. 4G), suggesting that *CSDE1* knockdown may be
synergistic with statins, consistent with our prior results (Fig. 3). Taken together, we conclude
that under physiologic conditions, CSDE1 mediates decay of the *LDLR* mRNA through its 3'
UTR, with the first ARE of the UTR required for its full effect.
266
Disruption of CSDE1 Upregulates the LDLR in vivo
We then turned to an *in vivo* model in zebrafish, as the 3' UTR of its ortholog *ldlra*
(XM_005163870.4) is highly AU-rich and contains at least two canonical ARE sequences for
mRNA regulation *(64)*. We employed yolk microinjection of a Cas9-ribonucleoprotein (RNP)
complex containing redundant guides to achieve near-saturation gene disruption *(65)*, followed
with dietary cholesterol supplementation, and evaluated total cholesterol in the larvae *(66)*.
Targeting of *csde1* protected against cholesterol accumulation, with a modest (12%) but
significant reduction in total cholesterol in 8-day post fertilization (dpf) zebrafish, without any
obvious phenotypic abnormalities (Figs. 5A, S10). By contrast, targeting of *ldlra* showed the
expected 1.4-fold increase in total cholesterol (Fig. 5A) *(66)*.
277
We next probed the effect of *Csde1* gene silencing in the mouse as a therapeutic proof-of-
principle, given even greater homology between the 3' UTRs of the murine and human *LDLR*
orthologs *(67)*. Using C57BL/6 mice on an atherogenic diet *(68)*, we delivered shRNA against
Csde1 (59), *Pcsk9*, or scramble control via low-dose adeno-associated virus 8 (AAV8). Two

weeks later, we observed a 25% reduction in fasting plasma cholesterol in the *Csde1*
knockdown mice, which exceeded the effect of *Pcsk9* knockdown (Fig. 5B). We then re-dosed
the *Csde1* and scramble AAV8-shRNA and, 2 weeks later, observed an even stronger
phenotype (Fig. S11A). Lipoprotein fractionation of the mouse plasma showed that *Csde1*
knockdown mostly affected the VLDL-containing fractions (Fig. 5C), consistent with upregulation
of the murine LDLR on our dietary background *(69, 70)*. Accordingly, we observed an increase
in *Ldlr* expression in the liver tissue of the *Csde1* knockdown mice (Fig. S11B). Notably, we
observed no differences in appearance or behavior of the mice, nor in plasma levels of alanine
aminotransferase activity (Fig. S11C), arguing against hepatic or systemic toxicity of either the
Csde1 shRNA or the gene knockdown.
292
To gain further insight, we performed bulk RNA sequencing on the liver tissue (Figs. 5D, S11D-
F, Table S10). We compared the *Csde1* knockdown to control (scramble) mice, using the mice
with the highest transcript counts of a vector-delivered eGFP reporter to control for variations in
transduction efficiency. We then filtered our results for the differentially expressed transcripts in
the control mice at the extremes of eGFP expression, to control for the effects of viral
transduction alone. As expected, we found higher *Ldlr* expression in the *Csde1* knockdown mice
($\log_2$FC = 0.43, p = 0.0029, Table S10). Consistent with our mechanistic hypothesis, GO
enrichment analysis of the differentially expressed genes revealed that mRNA processing was
the most significantly downregulated biological process in the *Csde1* knockdown mice (OR =
2.7, adj. p = 0.0085, Figs. 5D, S11F, Table S11). Spliceosome complex assembly was also
significantly downregulated (OR = 7.4, adj. p = 0.0085, Figs. 5D, S11F, Table S11).
304
Discussion
The powerful biology of the LDLR is unquestioned in cardiovascular medicine *(71)*. Since their
introduction, statins, which upregulate the LDLR, have become a major public health success,
and with the discovery of PCSK9 and the therapeutic antibodies targeting it, patients can safely
reach much lower LDL levels than is achievable by statins alone *(10)*. Together, this suggests
that we can push further on this LDL-LDLR axis and still achieve a clinical benefit.
311
In this study, we modeled a clinically relevant phenotype of LDLR abundance and function,
complementing the independent investigations of other groups *(72, 73)*. When synthesizing our
screening and validation data together with large-scale genomics and additional pharmacologic
perturbations, we produce an exploratory map of potential regulatory mechanisms for the LDLR
(Fig. 6). These data represent not just promising targets but also pathways likely to be impacted
by therapies already in use in the clinic.
318
We have shown that *CSDE1*, one of our strongest hits, regulates LDLR levels in HepG2 cells by
promoting *LDLR* mRNA decay via its 3' UTR. These data lay in concert with CSDE1's
destabilizing effects on other transcripts, such as c-Fos *(60)*. We have also shown that *in vivo*
knockdown of *Csde1* upregulates the hepatic LDLR and improves atherogenic lipid profiles in
mice. This mimics the effect of deleting the 3' UTR *in vivo (74)* and illustrates the promise of
targeting CSDE1 to lower LDL and protect against atherosclerosis. It is notable that several
small molecules, including triciribine *(63)* and berberine *(67, 75)*, have stabilizing effects on
LDLR mRNA, though whether their mechanisms directly involve CSDE1 remain to be
elucidated. The magnitude of LDLR upregulation imparted by *CSDE1* knockdown mirrors or
exceeds that of *HMGCR* and *PCSK9* in both tissue culture and mouse models, suggesting that
a high-fidelity approach targeting CSDE1-mediated *LDLR* mRNA decay in the clinic could have
similarly impressive effects. Additionally, our mechanistic data suggest that targeting CSDE1
would be at least additive with the use of statins.
332

The degree to which CSDE1 inhibition affects other transcripts, or other tissues *(59, 76)*,
remains an important question. As an RNA chaperone, CSDE1 can have a variety of effects,
from mRNA stabilization *(58)* to promotion or inhibition of translation *(77–80)*, dependent on the
identity of the RNA it binds and the cofactors with which it interacts. Intriguingly, though CSDE1
was found to bind biotinylated *LDLR* 3' UTR transcripts in HepG2 cell lysates *(62)*, cross-linking
immunoprecipitation approaches in both mouse brain and melanoma cells failed to identify
LDLR mRNA as a CSDE1 binding partner *(81, 82)*. This suggests that the CSDE1-*LDLR*
interaction is context dependent. Advances in liver-specific delivery of gene-silencing agents
(16, 83), novel gene editing technologies *(84)*, and small molecules *(85)* offer the possibility that
selectively targeting hepatic CSDE1 for cholesterol lowering could avoid systemic toxicities.
343
Though we observed no toxicity, transcriptional profiling suggests that disrupting hepatic
CSDE1 upregulates both apoptosis-related and GTPase-mediated signaling pathways (Figs.
5D, S11F, Table S11). Notably, CSDE1 appears to protect from apoptosis in both colorectal
cancer *(86)* and Huh7 cells *(87)*. Though biologically plausible, we hesitate to make definitive
conclusions from these data since the cholate-rich diet we used to obtain hyperlipidemia also
causes liver inflammation and hepatic steatosis *(88)*. Despite this confounder, we expect that
our transcriptomic analysis will guide further investigations of potential toxicities from hepatic
CSDE1 disruption. Given that CSDE1 has such varied effects on other transcripts, future
mechanistic dissection of the hepatic CSDE1-*LDLR* interaction could identify what makes this
relationship unique and guide a potential therapeutic strategy. Combination therapies targeting
interconnected pathways to disease can provide increased benefits without inducing extreme
side effects, with angiotensin receptor blockade and neprilysin inhibition in heart failure a
prominent clinical example *(89)*. Though speculative, we are intrigued by the effects of CSDE1
on mRNA splicing (Figs. 5D, S11F, Table S11), and we note in particular that the spliceosome
helicase DDX39B was both validated as a regulator of the LDLR from our CRISPRi screen (Fig.
2, Table S8) and downregulated by *Csde1* knockdown *in vivo* ($\log_2$FC = -0.32, p = 0.011, Table
S10). To the extent hepatic CSDE1 utilizes specific factors to downregulate *LDLR* mRNA,
simultaneous tissue-specific drugging of both CSDE1 and these factors could widen the overall
therapeutic window. We anticipate that our study will serve as the seed for these and other
further investigations.
364
Methods
Study Design
We designed the study as a discovery biology experiment to identify new regulators of the LDL
receptor. We used an established tissue culture model, HepG2 cells, to evaluate for LDL
receptor regulation. We used wild-type zebrafish (Ekwill) and wild-type mice (C57BL/6) to
validate the contribution of our top hit, CSDE1, to LDL receptor regulation *in vivo*. We evaluated
sufficient cells for the LDLR and TFR screens to provide adequate coverage for transduction
and downstream sequencing of each sgRNA in the genome-wide library. Sample sizes for
animal experiments were estimated to provide 80% power (two-tailed α = 0.05) for a 25% effect
in cholesterol levels compared to controls, based on effects in these models in the existing
literature. The numbers of animals used in each experiment are noted in the figures and
manuscript. Unless otherwise noted, all *in vitro* data are representative of multiple (≥ 3)
experimental outcomes to ensure robust outcomes. Experiments were not performed in a
blinded fashion. All animal studies were performed in accordance with IACUC approved
protocols at the University of California, San Francisco.
380
Plasmids and Cloning
SFFV-dCas9-BFP-KRAB (Addgene 46911), CRISPRi/a v2 (Addgene 84832), pMD2.G, dR8.91,
and the hCRISPRi v2 top5 sgRNA library (Addgene 83969) were gifts from L. Gilbert and J.

Weissman. Oligonucleotides of the protospacers of validated sgRNA sequences *(14)*, as well as
those for PCR amplification and isothermal assembly, were obtained from Elim
Biopharmaceuticals (Hayward CA). Protospacers were cloned into the CRISPRi/a v2 vector
using restriction enzyme digest (BlpI and BstXI, ThermoFisher, Waltham MA) and ligation with
10× T4 ligase (NEB, Ipswich MA). CSDE1 overexpression constructs were created by PCR
expansion of *CSDE1* (HsCD00949797, DNASU, Tempe AZ) or AcGFP1 (vector control,
pIRES2-AcGFP1, Clontech, Mountain View CA) and isothermal assembly *(90)* into the
pcDNA5/FRT/TO backbone (ThermoFisher), followed by site-directed mutagenesis to generate
the four CSDE1 isoforms (UniprotKB O75534 1 through 4). pLuc2-Prom$_{LDLR}$ was created by
PCR expansion of the target luciferase from pGL4Luc-RLuc (Addgene 64034), custom gene
synthesis of the LDLR promoter (NCBI Reference Sequence NG_009060.1, from -687 bp to the
LDLR start codon, Twist Biosciences, South San Francisco CA) and isothermal assembly into
the pcDNA5/FRT/TO backbone. pSS-NLuc was created by PCR expansion of the target
luciferase from pNL1.1 (Promega, Madison WI) into a vector containing the PCSK9 signal
sequence from the same backbone *(91)*. The remaining pLuc2-Prom$_{LDLR}$ constructs were
created by PCR expansion of the coding region of *LDLR* (HsCD00004643, DNASU), custom
gene synthesis of the 3' UTR of the *LDLR* mRNA (NCBI Reference Sequence
NM_001252658.1, Twist Biosciences), or custom oligonucleotides to add the P2A ribosomal
skipping linker *(92, 93)* and isothermal assembly into pLuc2-Prom$_{LDLR}$, as appropriate for each
construct. All plasmids were confirmed by Sanger sequencing. Expansion of the top5 sgRNA
library was as previously described *(13)*.
405
Cell Culture and Lentiviral Production
HepG2 (ATCC HB-8065) and derivatives were cultured in low-glucose DMEM (1 g/L,
ThermoFisher) with 10% FBS (Axenia BioLogix, Dixon CA), GlutaMax (ThermoFisher) and 1×
penicillin-streptomycin (ThermoFisher), and sent thrice through a 21g needle during passaging
to minimize cell clumping. HEK-293T (ATCC CRL-3216) were cultured in standard DMEM
(ThermoFisher) with 10% FBS. All cell lines were cultured at 37 °C at 5% CO_2, seeded for
approximately 50% confluency at the time of experiment, and were confirmed free of
Mycoplasma contamination by the MycoAlert PLUS Mycoplasma Detection Kit (Lonza,
Switzerland). Lentivirus was produced in 293T cells by transfection of dR8.91, pMD2.G, and the
appropriate pLKO-derived vector (at ratios of 8 µg, 1 µg, and 8 µg, respectively, per 15 cm dish)
with Trans-LT1 (Mirus Bio, Madison WI), according to the manufacturer's instructions. Viral
harvest media was supplemented with Viralboost (Alstem, Richmond CA), collected 2-3 days
after transfection, and filtered through 0.44 µm polyvinylidene difluoride filters and either frozen
for storage at -80 °C or used immediately for transduction.
420
Generation of CRISPRi Cell Lines
All cell lines were transduced using virus-containing supernatant in the presence of 8 µg/ml
polybrene (Millipore-Sigma, St. Louis MO). HepG2 expressing dCas9-KRAB were derived by
transduction with lentivirus harboring SFFV-dCas9-BFP-KRAB, followed by two rounds of FACS
for BFP-positive cells on a BD FACSAria II. dCas9-KRAB HepG2 with individual targeting
sgRNAs were derived by transduction with lentivirus harboring the desired sgRNA, followed by
48 hrs of puromycin selection (2 µg/ml, InvivoGen, San Diego CA), prior to experiments.
428
Quantitative Real-Time PCR
dCas9-KRAB HepG2 stably expressing an appropriate sgRNA were harvested, lysed, and total
RNA was extracted via the RNeasy Mini Kit (Qiagen, Germantown MD). RNA was converted
into cDNA using qScript cDNA SuperMix (QuantaBio, Beverly MA) following the manufacturer's
instructions. RT-qPCR was performed against indicated targets with PrimeTime qPCR primers
(IDT, Coralville IA) using the SYBR Select Master Mix (ThermoFisher) according to the

manufacturer's instructions on a CFX96 Touch Real-Time PCR Detection System (BioRad,
Hercules CA). Fold changes were calculated using ΔΔCt analysis, normalizing each sample to
B2M controls, using CFX Maestro software (BioRad).
438
Receptor Abundance Analysis
1-2 days prior to analysis, dCas9-HepG2 cells and derivatives were cultured in low-glucose
DMEM with 5% lipoprotein deficient serum (Kalen Biomedical, Germantown MD). Prior to
analysis, cells were dissociated with Accutase (Innovative Cell Technologies, San Diego CA),
collected, washed in PBS (ThermoFisher), live-dead stained with Ghost Dye Red 780 (1:1000
dilution, Tonbo Biosciences, San Diego CA), washed, and then stained with the indicated
antibody in FACS buffer (PBS with 1% FBS, 10 U/ml DNAse I, GoldBio, St. Louis MO) for 30
minutes on ice with gentle agitation. Cells were washed, resuspended in FACS buffer, filtered
through 90 µm mesh (Elko Filtering, Miami FL) to give a single cell suspension, and placed on
ice. Cells were then analyzed on either a BD Fortessa, BD LSRII, BD FACSAria II, or Beckman
Coulter CytoFLEX, or sorted on a BD FACSAria II, depending on the experiment. In general,
gating excluded cells positive for live-dead staining and included only the cells positive for the
level of BFP expression induced by the CRISPRi/a v2 vector. FACS analysis and figure
preparation was performed with FlowJo v10 (BD, Ashland OR).
453
Genome-Wide CRISPRi Screen
The screen was conducted similarly to prior descriptions *(13–15)*. Approximately 200×10^6
dCas9-KRAB HepG2 were transduced with hCRISPRi-v2 top 5 sgRNAs/gene lentivirus at an
MOI of ~0.5, and with polybrene at 8 µg/ml, on day 1. Cells were grown on 15-cm dishes,
subdivided into four replicates immediately upon transduction (biological duplicate for each
screen), and reseeded every 3-4 days as necessary to avoid overconfluence. Cells were
selected with puromycin (2 mg/ml) from day 2 through day 6. On day 5, cells for the LDLR sort
were placed in DMEM with lipoprotein depleted serum (5%). On day 7, approximately 50×10^6
cells from 2 replicates were live-dead stained and stained for LDLR as described above, and
then two-way sorted on a BD FACSAria II for the top and bottom 33% by LDLR abundance.
Cells were spun down, washed in PBS and frozen at -80 °C. On day 8, the sort was repeated
except in one replicate, cells were stained for TFR instead of LDLR and then sorted as per
above. Genomic DNA was isolated using a NucleoSpin Blood DNA extraction kit (Macherey-
Nagel, Bethlehem PA). The sgRNA-containing region was PCR-amplified with NEBNext Ultra II
Q5 MasterMix (NEB), acrylamide gel-purified, and size-selected by SPRI beads (Beckman
Coulter, Indianapolis IN), all as previously described, prior to sequencing on an Illumina HiSeq
4000.
471
Screen Processing
Sequencing data were aligned to the top5 library, counted, and quantified using the
ScreenProcessing pipeline (accessed from https://github.com/mhorlbeck/ScreenProcessing*(14)*
4/25/2019). Phenotypes and Mann-Whitney *P* values were determined as previously described
(14), with the phenotypes defined as follows: ρ indicated the comparison in high-abundance vs.
low abundance cells, τ indicated the comparison in high-abundance vs. unsorted cells, and γ
indicated the comparison in low-abundance vs. unsorted cells. Counts from 4 guides were
removed from the final analysis as there was evidence of contamination from individually cloned
plasmids (PCSK9_+_55505255.23-P1P2, HMGCR_+_74633053.23-P1P2, TFRC_-
195808987.23-P1P2, ACO1-_32384733.23-P1P2). A hit threshold of 7 (normalized
phenotype z score $\times$ -log10(p-value) $\geq$ 7) *(13)* was used to identify hits from ρ, τ, and γ
phenotypes, which were then compiled. Identical analysis of the TFR screen was used to
prioritize hits unique to LDLR regulation. Gene ontology analysis was performed using the

PANTHER Classification System database (v15) *(28, 94)*. For relaxation of the hit threshold for
comparison to GWAS studies, a score of 6 was used. Cellular localization of hits was imputed
by manual curation from UniProt *(95)* and the Human Protein Atlas *(55)*.
488
Human Genomic Analysis
Protein coding variants for hits validated at the individual sgRNA level were assayed in the UK
Biobank *(96)* for associations with LDL-C. In the setting of a statin medication, LDL-C was
divided by 0.7 as before *(30)*. Genotyping and imputation was performed in the UK Biobank as
previously described *(32)*, and nonsynonymous protein coding variants with minor allele
frequencies greater than 0.001 were considered. Efficient linear mixed models adjusting for age,
sex, genotyping array, and principal components of ancestry were employed, using BOLT-LMM
(97). Statistical significance was assigned at α = 0.05/117 = 0.000427 to account for multiple
hypothesis testing.
498
499 *Validation Experiments of Individual sgRNAs*
500 Cloning of protospacers, as described above, was performed in 96-well plate format until
501 selecting individual colonies. Lentiviral production in 293T, transduction of dCas9-KRAB HepG2
502 with lentiviral sgRNA vectors, and receptor abundance and LDL uptake assays were similarly
503 performed in 96-well plate format to maximize throughput.
504
LDL Uptake Assays
Assays were performed as previously described*(33)* with the following modifications. dCas9-
HepG2 cells harboring individual sgRNAs were treated similarly to receptor abundance analysis,
except that prior to harvest, cells were washed and then treated with 5 µg/ml 1,1'-dioctadecyl-
3,3,3',3'-tetramethylindocarbocyanine perchlorate (DiI) labeled LDL (Kalen Biomedical) in low-
glucose DMEM with 0.5% BSA (MilliporeSigma) for 1 hr at 37 $^{\circ}$C. Cells were then washed,
collected, and prepared for FACS analysis, as described above, but without antibody labelling.
512
513 *Pharmacologic Synergy Experiments*
514 Receptor and LDL uptake assays were performed as described, with cells treated overnight with
515 either simvastatin (6 µM, MilliporeSigma), PF-6846 (10 µM, MilliporeSigma), or DMSO vehicle
516 (final concentration of 0.5%) overnight prior to analysis. Synergy scores were calculated by the
517 following equation:
518
519 $$\left(LDLR_{sgRNA+compound} - LDLR_{NegCtrl+compound} \right) - \left(LDLR_{sgRNA+vehicle} - LDLR_{NegCtrl+vehicle} \right)$$
520
521 with $LDLR$ obtained as the mean fluorescence, background subtracted from an unstained
522 control and subsequently normalized to $LDLR_{NegCtrl+vehicle}$ within a given experiment.
523
Overexpression Experiments
HepG2 or engineered dCas9-HepG2 cell lines were seeded into 96 well plates at 5×10^4 cells
per well in HepG2 growth medium. After 24 hrs, cells were washed and changed into low-
glucose DMEM with 5% lipoprotein-deficient serum. Each well was transfected with 100 ng of
the appropriate CSDE1 overexpression construct, or vector control, in a total of 10 µL OptiMEM
(ThermoFisher) using Lipofectamine 3000 (ThermoFisher) according to the manufacturer's
instructions. Cells were incubated at 37 $^{\circ}$C with 5% CO_2 for 72 hrs, and then harvested for LDL
receptor expression analysis as above.
532
mRNA Decay Experiments

Engineered dCas9-HepG2 cell lines harboring sgRNAs against CSDE1
(CSDE1_+_115300577.23-P1P2) or a negative control (Unassigned=negZNF335_-
_44601297.24-all) were seeded into 12 well plates at 5×10^5 cells per well in HepG2 growth
medium. After 24 hrs, cells were washed and changed into sterol-depleting media (low-glucose
DMEM with 5% lipoprotein-deficient serum) supplemented with 6 µM simvastatin. After an
additional 24 hrs, actinomycin D (MilliporeSigma) was added at 5 µg/ml, and cells were
harvested as described at the indicated timepoints.
541
Immunoblots
Engineered dCas9-HepG2 cell lines harboring appropriate sgRNAs were grown in growth
medium and harvested with 0.25% trypsin digestion. Cells were washed and lysed in lysis buffer
on ice (50 mM Tris-HCl pH 7.4, 150 mM NaCl, 0.1% NP-40). Lysates were clarified at $21,000 \times$
g for 10 min, and the supernatant was recovered. Equivalent amounts of lysates, as measured
by BCA assay (ThermoFisher), were resolved on 4-12% Bis-Tris NuPAGE gels (ThermoFisher),
transferred to nitrocellulose, probed with primary and secondary antibodies as noted (see Table)
in 5% BSA in TBS-T, and visualized on an Odyssey imaging system (LI-COR, Lincoln NE).
550
Dual-Luciferase Assays
Engineered dCas9-HepG2 cells were seeded into opaque white 96 well plates, at 2.2×10^4 cells
per well, in 100 µL growth medium the day prior to transfection. On day of transfection, medium
was replaced or changed to sterol-depleted medium (low-glucose DMEM with 5% lipoprotein-
deficient serum) with or without 6 µM simvastatin as appropriate. Each well was transfected with
100 ng of Luc2-Prom$_{LDLR}$ based construct and 1 ng of secreted nanoluciferase control construct
(pSS-NLuc) in a total of 10 µL OptiMEM using Lipofectamine 3000 according to the
manufacturer's instructions. 6 replicates were transfected per construct per experiment. After 48
hours at 37°C and 5% CO_2, 10 µL of medium was removed from each plate and aliquoted into a
separate 384 well plate. Firefly luciferase activity was evaluated in the plates containing the
cells by adding an equal volume of a $2\times$ firefly lytic assay buffer (100 mM Tris-HCl pH 7.7, 50
mM NaCl, 2 mM $MgCl_2$, 0.17% Triton X-100, 10 mM DTT, 0.4 mM coenzyme A, 0.3 mM ATP,
and 0.28 mg/ml luciferin (Goldbio)) *(98)*. Nanoluciferase activity was evaluated from the
conditioned medium using a non-lytic $2\times$ coelenterazine (Goldbio) reagent as previously
described *(91)*. Raw luminescence was obtained on a SPARK plate reader (Tecan, San Jose
CA) with 1 second integration time. Readout of firefly luciferase in each well was normalized to
the corresponding secreted nanoluciferase control and data were visually inspected and
cleaned to remove values from poorly transfected wells (formally defined by ROUT = 1%) during
analysis.
570
Zebrafish Handling, Maintenance, and Cas9-Ribonucleoprotein Knockdowns
All zebrafish studies were performed as previously described*(65)* with minor modifications.
Briefly, wild type zebrafish embryos were injected at the one-cell stage with Cas9-RNP
complexes and raised at 28 °C. Cas9-RNP complexes were prepared as previously
described*(65)* using custom oligonucleotides against the indicated targets (Elim
Biopharmaceuticals). Targeting of tyrosinase, which results in larval albinism, was used as an
injection control. Larvae were fed a diet of Golden Pearls (5-50 micron, Brine Shrimp Direct,
Ogden UT) $3\times$ daily from 4 days post fertilization (dpf), fasted on 7 dpf to clear intestinal
cholesterol, and harvested at 8 dpf. Larvae were collected, extensively washed, anesthetized in
tricaine, and collected in groups of 10 per sample prior to storage at -80 °C. All zebrafish
experiments were performed in accordance with IACUC-approved protocols at the University of
California, San Francisco.
583

Cholesterol Analysis of Zebrafish Homogenates
Total cholesterol levels were analyzed as previously described *(66)* with minor modifications.
Briefly, frozen larvae were homogenized in PBS with a plastic pestle, and then clarified at
18,000 × *g* for 15 min. Supernatants were recovered and total protein content was analyzed by
BCA assay. Homogenates were then analyzed, in duplicate, at the appropriate dilution (typically
1:12 in PBS) for total cholesterol content using a commercial fluorometric assay (Cayman
Chemical, Ann Arbor MI). Fluorescence outputs were measured on a Tecan SPARK plate
reader, and cholesterol concentrations were interpolated from a regression line calculated from
a standard curve. Cholesterol levels were normalized to total protein content for analysis and
subsequently to the scramble control for comparison between experiments.
594
Mouse Handling, Maintenance, and shRNA Knockdowns
All mouse manipulations were performed in accordance with IACUC approved protocols at the
University of California, San Francisco following guidelines described in the US National
Institutes of Health Guide for the Care and Use of Laboratory Animals. 8-10 week old male
C57BL/6 mice (The Jackson Laboratory, Bar Harbor ME) were placed on an atherogenic diet
(1.25% cholesterol, 15% fat, 0.5% cholate, D12336i, Research Diets, New Brunswick NJ)*(68)* at
the beginning of the experiment (week 0). At week 2, AAV8-packaged shRNA against mouse
Csde1 (NM_144901), *Pcsk9* (NM_153565), or scramble control (Vector Biolabs, Malvern PA)
were diluted in sterile PBS to a concentration of 2×10^{11} genomes/ml. 100 µL of diluted AAV8 (2
$\times 10^{10}$ genomes/mouse) harboring the appropriate shRNA was administered to each mouse via
tail vein injection. At week 4, and again at week 6, mice were fasted overnight and then
underwent blood sampling via submandibular vein puncture. Approximately 50 µL of blood was
collected into an EDTA-coated tube, centrifuged at 2000 × g for 10 min at 4 °C, and the plasma
recovered and stored at -20 °C until further analysis. Total cholesterol of the plasma, after
approximately 1:200 to 1:400 dilution in assay buffer, was evaluated by commercial fluorometric
cholesterol assay (Cayman) as described above. At week 8, mice from the same exposure arm
were re-dosed with AAV8 targeting either *Csde1* or scramble control. At week 10, the mice were
again fasted overnight and then euthanized after CO_2 narcosis followed by cervical dislocation.
The abdominal cavity was opened with a ventral midline incision, the IVC was cannulated, and
plasma was collected as described above. The liver and vasculature were perfused with PBS,
and the samples of the liver were harvested. Tissue samples for RNA evaluation were placed in
TRIzol (ThermoFisher) and those for protein analysis were flash frozen in liquid N_2 and stored at
-80 °C.
618
Lipoprotein Fractionation
Plasma samples were thawed and centrifuged at 2000 × g for 10 min at 4 °C, and the
supernatant recovered. 100 µL of individual mouse plasma was loaded onto a Superose 6
Increase 10/300 GL column (Cytiva, Marlborough MA) and eluted with PBS with 1 mM EDTA at
0.5 ml/min on an AKTA Pure chromatography system (Cytiva). Fixed 0.5 ml fractions were
collected from 0.2 to 1 column volumes along the isocratic elution. Fractions were subjected to
total cholesterol analysis and immunoblots as described above.
626
RNA-seq Library Preparation
Total RNA was extracted from frozen liver samples using the Qiagen RNeasy Plus Universal
mini kit followed by Manufacturer's instructions (Qiagen). RNA samples were quantified using
Qubit 2.0 Fluorometer (ThermoFisher) and RNA integrity was checked using Agilent
TapeStation 4200 (Agilent Technologies, Palo Alto CA). Purified RNA was used for mouse
qPCR experiments as described above. RNA sequencing libraries were prepared via polyA
selection using the NEBNext Ultra RNA Library Prep Kit for Illumina using manufacturer's

instructions (NEB). Briefly, mRNAs were initially enriched with Oligod(T) beads. Enriched
mRNAs were fragmented for 15 minutes at 94 °C. First strand and second strand cDNA were
subsequently synthesized. cDNA fragments were end repaired and adenylated at 3'ends, and
universal adapters were ligated to cDNA fragments, followed by index addition and library
enrichment by PCR with limited cycles. The sequencing library was validated on the Agilent
TapeStation (Agilent) and quantified by using Qubit 2.0 Fluorometer (Invitrogen) as well as by
quantitative PCR (KAPA Biosystems, Wilmington, MA, USA). The sequencing libraries were
clustered on a single lane of a flowcell. After clustering, the flowcell was loaded on the Illumina
HiSeq instrument (4000 or equivalent) according to manufacturer's instructions. The samples
were sequenced using a 2x150bp Paired End (PE) configuration. Image analysis and base
calling were conducted by the HiSeq Control Software (HCS). Raw sequence data (.bcl files)
generated from Illumina HiSeq was converted into fastq files and de-multiplexed using Illumina's
bcl2fastq 2.17 software. One mismatch was allowed for index sequence identification. RNA
library preparation and sequencing were conducted by GENEWIZ, LLC (South Plainfield, NJ).
648
RNA-seq Analysis
All raw sequencing data underwent quality control checks with FastQC (v 0.11.8). Reads were
mapped to the mm10 mouse reference genome using Rsubread (v 2.4.3) and assigned to
Ensembl gene IDs. Ensemble gene IDs were then mapped to gene symbols using
AnnotationDBI (v1.52.0). Gene expression was quantified using raw counts and differential
expression gene testing was performed on the scramble-shRNA samples comparing the groups
(n=3 in each group) at the highest and lowest levels of raw eGFP expression with EdgeR *(99,*
100) (v.3.32.1) using the glmQLFit method, default settings *(101)*. Statistical significance was
set at 5% false discovery rate (FDR; Benjamini-Hochberg). Differential expression gene testing
was then performed on the *Csde1*-shRNA and scramble-shRNA at the highest levels of eGFP
expression with the overlap of differentially expressed genes identified between these two
analyses subsequently removed. Functional enrichment gene-set analysis for GO (Gene
Ontology) terms was performed using Enrichr *(102)* via the enrichR R package (v.3.0).
Heatmaps were generated using the Bioconductor package ComplexHeatmap *(103)* (v.2.6.2)
using log2-transformed CPM values (counts-per-million; values shown are log2-transformed and
row-normalized). Volcano plots were generated using the Bioconductor package
EnhancedVolcano (v.1.2.0).
666
Statistical Analysis
Fluorescence values from gated populations in flow cytometry experiments were background
corrected by unstained controls and were normalized to the values of the cell line harboring
negative control sgRNA. Normalized data were then grouped by the Cochrane method *(104)*,
and values for cell lines transduced with individual sgRNAs were compared those of the
negative control by T-test with Holm-Sidak correction. For comparison of one-phase decay
regression curves in mRNA decay experiments, the extra sum-of-squares F test was used.
Pairwise testing to controls was performed in all other experiments using Welch's T-test with
Holm-Sidak correction unless otherwise noted. One or two-way ANOVA with Tukey's or Sidak's
multiple comparisons tests was used when comparing across all groups in a particular
experiment. Adjusted p values < 0.05 were considered significant. Statistical analysis was
performed using Prism 7 (GraphPad Software, San Diego CA). All experiments were replicated
thrice unless otherwise noted.
680
681 **Key Resources Tables**

Antibodies				
Target	**Fluorophore**	**Clone**	**Source**	**Dilution and Final Conc.**

Human LDL Receptor	Alexa Fluor 647	472413	R&D Systems	1:100, 2 µg/ml (FACS)
Human Transferrin Receptor	Alexa Fluor 647	29806	R&D Systems	1:100, 2 µg/ml (FACS)
Human Transferrin Receptor	Alexa Fluor 488	29806	R&D Systems	1:100, 2 µg/ml (FACS)
Human CSDE1	None	62328	Cell Signaling Technology	1:1000 (WB)
Human beta-Actin	None	8H10D10	Cell Signaling Technology	1:2000 (WB)
Mouse ApoA-I	None	2G4	Santa Cruz Biotechnology	1:1000 (WB)
Mouse ApoB	None	2G11	Millipore Sigma	1:1000 (WB)
Rabbit IgG	IRDye 800CW	926-32211	LI-COR	1:5000, 0.1 µg/ml (WB)
Mouse IgG	IRDye 800CW	926-32210	LI-COR	1:5000, 0.1 µg/ml (WB)

qPCR Primers				
Target	Ref. Sequence	Assay ID	Source	Conc.
B2M	NM_004048(1)	Hs.PT.58v.18759587	IDT	300 nM
LDLR	NM_000527(6)	Hs.PT.58.2004261	IDT	300 nM
HMGCR	NM_000859(2)	Hs.PT.58.41105492	IDT	300 nM
CSDE1	NM_001007553(6)	Hs.PT.58.40309152	IDT	300 nM
PCSK9	NM_174936(1)	Hs.PT.58.20317141	IDT	300 nM
MYLIP	NM_013262(1)	Hs.PT.58.39124976	IDT	300 nM
TFRC	NM_003234(1)	Hs.PT.39a.22214826	IDT	300 nM
HNRNPD	NM_031369(4)	Hs.PT.58.3757916	IDT	300 nM
KHSRP	NM_003685(1)	Hs.PT.58.555216	IDT	300 nM
B2m	NM_009735(1)	Mm.PT.39a.22214835	IDT	300 nM
Ldlr	NM_001252658(3)	Mm.PT.58.9930556	IDT	300 nM
Csde1	NM_001161854(2)	Mm.PT.58.8160050	IDT	300 nM
eGFP	n/a	Custom Primers Fwd: GAACCGCATCGAGCTGAA Rev: TGCTTGTCGGCCATGATATAG	IDT	300 nM

Acknowledgements

We thank Max Horlbeck for guidance with the CRISPRi system, Jonathan D. Brown for helpful discussion on *in vivo* experimental designs, and Paul Cheng and Richard Baylis for helpful discussion on histologic insights and *in vivo* experimental planning. Plasmids for the CRISPRi system were a generous gift from Luke Gilbert and Jonathan Weissman. We thank the Gladstone Institutes Flow Cytometry core facility for their assistance with flow cytometry experiments. UK Biobank analyses were conducted using the UK Biobank resource under application 7089.

Funding Sources

Ar. P. receives support from the Tobacco-Related Disease Research Program (578649), the A.P. Giannini Foundation (P0527061), NIH/NHLBI (K08 HL157700), the Michael Antonov Charitable Foundation Inc., and the Sarnoff Cardiovascular Research Foundation. T.N. is supported by the Japan Society for the Promotion of Science Overseas Research Fellowship. P.N. is supported by a Hassenfeld Scholar Award from the Massachusetts General Hospital, the NIH/NHLBI (R01 HL142711, R01 HL148565, and R01 HL148050), and the Fondation Leducq (TNE-18CVD04). B.L.B. is supported by NIH grants R01 DK119621 and P01 HL146366. D.S. is supported by NIH grants P01 HL098707, P01 HL146366, R01 HL057181 and R01 HL127240, the Roddenberry Foundation, the L.K. Whittier Foundation, the Younger Family Fund, and the NIH/National Center for Research Resources grant C06 RR018928 to the Gladstone Institute. K.M.S. is supported by the Howard Hughes Medical Institute. J.S.C. is supported by the NIH/NHLBI (K08 HL124068 and R03 HL145259), a Pfizer ASPIRE Cardiovascular Award and the Harris Fund and Research Evaluation and Allocation Committee of the UCSF School of Medicine.

Author Contributions

Overall study design: J.S.C. Execution of *in vitro* screen and data processing: G.A.S. Genomic analyses: Ak.P., P.N. *In vitro* validation and synergy experiments, and analysis of mouse blood samples: J.S.C. Execution of zebrafish gene knockdowns: B.H.L., R.S.W. Oversight of zebrafish husbandry: R.S.W., B.L.B. Planning and execution of *in vivo* mouse experiments: Ar.P. Mouse handling, blood, and tissue collection: Ar.P., Y.C.L., T.N., N.S. Processing of RNA-seq data: Ar.P., An.P. Analysis of mouse samples: L.L., R.J. Critical data review and analysis: B.L.B, D.S., K.M.S. Preparation of manuscript: J.S.C. Critical review and revision of manuscript: All authors.

Declaration of Interests

P.N. reports investigator-initiated grant support from Amgen, Apple, and Boston Scientific, and personal fees from Apple, Blackstone Life Sciences, and Novartis, all unrelated to the present work. R.S.W. is an employee of Amgen, Inc. D.S. is the scientific cofounder, shareholder, and director of Tenaya Therapeutics, unrelated to the present work. K.M.S. has consulting agreements for the following companies involving cash and/or stock compensation: Black Diamond Therapeutics, BridGene Biosciences, Denali Therapeutics, Dice Molecules, eFFECTOR Therapeutics, Erasca, Genentech/Roche, Janssen Pharmaceuticals, Kumquat Biosciences, Kura Oncology, Merck, Mitokinin, Petra Pharma, Revolution Medicines, Type6 Therapeutics, Venthera, Wellspring Biosciences (Araxes Pharma). J.S.C. has received consulting fees from Gilde Healthcare and is an unpaid scientific advisor to Eko, both unrelated to this work.

Resource and Data Availability Statement

Additional supporting data are available upon request from the corresponding author. All requests for raw and analyzed data, and materials, including plasmids or cell lines, generated in

735 this study will be responded to promptly. UK Biobank data is available by application to the UK
736 Biobank.
737
738 **Supplementary Materials**
739
Figure S1: Validation of dCas9-KRAB-HepG2 Cells.
Figure S2: Recovered sgRNAs from Screening Phenotypes.
Figure S3: Gene Ontology and Localization Analysis.
Figure S4: Selective LDLR Effect of Transmembrane Proteins.
Figure S5: Effect of Sterol Conditions on CSDE1 Knockdown.
Figure S6: Effect of CSDE1 Overexpression.
Figure S7: CSDE1 Knockdown at Protein Level.
Figure S8: Effect of *CSDE1* Knockdown on Decay of Non-*LDLR* Transcripts.
Figure S9: Physiologic Response of Luciferase Reporter System.
Figure S10: Visual Phenotypes of Zebrafish Cas9-sgRNA Saturation Gene Disruption.
Figure S11: Effects of *in vivo Csde1* Disruption in Mice.
751
Table S1: LDLR Screen Data by Gene *(provided as an Excel file)*
Table S2: LDLR Screen Data by Guide *(provided as an Excel file)*
Table S3: TFR Screen Data by Gene *(provided as an Excel file)*
Table S4: TFR Screen Data by Guide *(provided as an Excel file)*
Table S5: LDLR Screen Hits *(provided as an Excel file)*
Table S6: TFR Screen Hits *(provided as an Excel file)*
Table S7: Baseline Characteristics of UK Biobank Participants in Genomic Association Analyses
(included below)
Table S8: Validation Data by Guide *(provided as an Excel file)*
Table S9: Pharmacology Synergy Data by Guide *(provided as an Excel file)*
Table S10: Differentially Expressed Genes by *in vivo* RNA Seq *(provided as an Excel file)*
Table S11: Enriched GO Terms by *in vivo* RNA Seq *(provided as an Excel file)*
764

References

1. M. S. Brown, J. L. Goldstein, A receptor-mediated pathway for cholesterol homeostasis,
Science (80-.). (1986), doi:10.1126/science.3513311.

2. J. L. Goldstein, M. S. Brown, A Century of Cholesterol and Coronaries: From Plaques to
Genes to Statins, *Cell* **161**, 161–172 (2015).

3. J. L. Goldstein, M. S. Brown, The LDL receptor*Arterioscler. Thromb. Vasc. Biol.* (2009),
doi:10.1161/ATVBAHA.108.179564.

4. M. G. Silverman, B. A. Ference, K. Im, S. D. Wiviott, R. P. Giugliano, S. M. Grundy, E.
Braunwald, M. S. Sabatine, Association Between Lowering LDL-C and Cardiovascular Risk
Reduction Among Different Therapeutic Interventions, *JAMA* **316**, 1289 (2016).

5. K. D. Kochanek, J. Xu, E. Arias, Mortality in the United States, 2019., *NCHS Data Brief* , 1–8
(2020).

6. G. G. Schwartz, P. G. Steg, M. Szarek, D. L. Bhatt, V. A. Bittner, R. Diaz, J. M. Edelberg, S.
G. Goodman, C. Hanotin, R. A. Harrington, J. W. Jukema, G. Lecorps, K. W. Mahaffey, A.
Moryusef, R. Pordy, K. Quintero, M. T. Roe, W. J. Sasiela, J. F. Tamby, P. Tricoci, H. D. White,
A. M. Zeiher, L. B. Schiavi, M. Garrido, A. F. Alvarisqueta, S. A. Sassone, A. P. Bordonava, A.
E. Alves De Lima, J. M. Schmidberg, E. A. Duronto, O. C. Caruso, L. P. Novaretto, M. A.
Hominal, O. R. Montaña, A. Caccavo, O. A. Gomez Vilamajo, A. J. Lorenzatti, L. R. Cartasegna,
G. A. Paterlini, I. J. Mackinnon, G. D. Caime, M. Amuchastegui, O. R. Codutti, H. O. Jure, J. O.
E. Bono, A. D. Hrabar, J. A. Vallejos, R. A. Ahuad Guerrero, F. Novoa, C. A. Patocchi, C. J.
Zaidman, M. E. Giuliano, R. D. Dran, M. L. Vico, G. S. Carnero, P. N. Guzman, J. C. Medrano
Allende, D. F. Garcia Brasca, M. H. Bustamante Labarta, S. Nani, E. D. S. Blumberg, H. R.
Colombo, A. Liberman, H. L. Luciardi, G. D. Waisman, M. A. Berli, R. O. Garcia Duran, H. G.
Cestari, H. A. Luquez, J. A. Giordano, S. S. Saavedra, J. H. Waites, N. Collins, A. Soward, P. E.
Aylward, C. L. S. Hii, J. Shaw, M. A. Arstall, J. Horowitz, J. F. Rogers, D. Colquhoun, R. E.
Oqueli Flores, P. Roberts-Thomson, O. Raffel, S. J. Lehman, S. G. M. Coverdale, P. J. Garrahy,
G. Starmer, M. Sader, P. A. Carroll, R. Zweiker, U. Hoppe, H. Drexel, K. Huber, R. Berger, F.
Weidinger, D. Faes, K. Hermans, B. Pirenne, A. Leone, E. Hoffer, P. Sinnaeve, M. C. M. Vrolix,
L. De Wolf, B. Wollaert, M. Castadot, K. Dujardin, C. Beauloye, G. Vervoort, H. Striekwold, C.
Convens, J. Roosen, E. Barbato, M. Claeys, F. Cools, I. Terzic, F. Barakovic, Z. Midzic, B.
Pojskic, E. Fazlibegovic, M. Dilic, A. Durak-Nalbantic, D. Vulic, A. Muslibegovic, G. Reis, L.
Sousa, J. C. Nicolau, F. E. Giorgeto, R. P. Silva, L. Nigro Maia, R. Rech, P. R. F. Rossi, M.
Cerqueira, N. Duda, R. Kalil, A. Kormann, J. A. M. Abrantes, P. Pimentel Filho, A. P. Soggia, M.
O. N. de Santos, F. Neuenschwander, L. C. Bodanese, Y. L. Michalaros, F. G. Eliaschewitz, M.
H. Vidotti, P. E. Leaes, R. V. Botelho, S. Kaiser, E. R. F. Fernandes Manenti, D. B. Precoma, J.
C. Moura Jorge, P. G. M. de B Silva, J. A. Silveira, W. Saporito, J. A. Marin Neto, G. S. Feitosa,
L. E. F Ritt, J. A. de Souza, F. Costa, W. Souza, H. J. L. Reis, R. D. Lopes, L. Machado, J. C.
Aidar Ayoub, G. V. Todorov, F. P. Nikolov, E. S. Velcheva, M. L. Tzekova, H. O. Benov, S. L.
Petranov, H. S. Tumbev, N. S. Shehova-Yankova, D. T. Markov, D. H. Raev, M. N. Mollov, K. N.
Kichukov, K. A. Ilieva-Pandeva, N. N. Gotcheva, R. Ivanova, V. M. Mincheva, P. V. Lazov, B. I.
Dimov, M. Senaratne, J. Stone, J. Kornder, S. Pearce, D. Dion, D. Savard, Y. Pesant, A.
Pandey, S. Robinson, G. Gosselin, S. Vizel, G. Hoag, R. Bourgeois, A. Morisset, E. Sabbah, B.
Sussex, S. Kouz, P. MacDonald, N. Michaud, D. Fell, R. Leung, T. Vuurmans, C. Lai, F. Nigro,
R. Davies, G. Nogareda, R. Vijayaraghavan, J. Ducas, S. Lepage, S. Mehta, J. Cha, R. Dupuis,
P. Fong, J. Rodes-Cabau, H. Fadlallah, D. Cleveland, T. Huynh, I. Bata, A. Hameed, C. Pincetti,
S. Potthoff, J. C. Prieto, M. Acevedo, A. Aguirre, M. Vejar, M. Yañez, G. Araneda, M.
Fernandez, L. Perez, P. Varleta, F. Florenzano, L. Huidobro, C. A. Raffo, C. Olivares, J. Chen,
Y. Dong, W. Huang, J. Wang, S. Huang, Z. Yao, L. Cui, W. Lin, Y. Sun, J. Li, X. Zhang, H. Zhu,
D. Chen, L. Huang, S. Dong, G. Su, B. Xu, X. Su, X. Cheng, J. Lin, W. Zong, H. Li, Y. Feng, D.
Xu, X. Yang, Y. Ke, X. Lin, Z. Zhang, Z. Zheng, Z. Luo, Y. Chen, C. Ding, Y. Zheng, X. Li, D.
Peng, Y. Li, M. Wei, S. Liu, Y. Yu, B. Qu, W. Jiang, Y. Zhou, X. Zhao, Z. Yuan, Y. Guo, X. Xu,

X. Shi, J. Ge, G. Fu, F. Bai, W. Fang, X. Shou, N. Jaramillo, G. Sanchez Vallejo, D. C. Luna
Botia, R. Botero Lopez, D. I. Molina De Salazar, A. J. Cadena Bonfanti, J. Diego Higuera, S. I.
Barrera Silva, H. J. Garcia Lozada, J. A. Coronel Arroyo, J. L. Accini Mendoza, R. L. Fernandez
Ruiz, A. M. Fernandez, F. G. Manzur Jatin, A. Sotomayor Herazo, J. Castellanos Parada, M. A.
Urina Triana, M. Strozzi, S. Car, D. Miličić, M. L. Benčić, H. Pintarić, D. Prvulović, J. Šikić, V.
Peršić, D. Mileta, K. Štambuk, Z. Babić, J. Spinar, D. Horak, J. Stasek, D. Alan, V. Machova, A.
Linhart, V. Novotny, V. Kaucak, R. Rokyta, R. Naplava, Z. Coufal, V. Adamkova, I. Podpera, J.
Zizka, Z. Motovska, I. Marusincova, P. Svab, P. Ostadal, P. Heinc, J. Kuchar, P. Povolny, S. H.
Poulsen, B. Raungaard, P. Clemmensen, L. E. Bang, O. May, M. Bøttcher, J. D. Hove, L. Frost,
G. Gislason, J. Larsen, P. B. Johansen, F. Hald, J. Jeppesen, T. Nielsen, K. S. Kristensen, P.
M. Walichiewicz, J. D. Lomholdt, I. C. Klausen, P. K. Nielsen, F. Davidsen, L. Videbaek, M.
Viigimaa, M. Soots, V. Vahula, A. Hedman, U. Soopõld, K. Märtsin, M. R. Taskinen, K. Porthan,
J. K. Airaksinen, M. Juonala, T. Kiviniemi, S. Vikman, P. Posio, J. Taurio, H. Huikuri, K.
Kaikkonen, P. Coste, E. Ferrari, N. Danchin, O. Morel, G. Montalescot, G. Barone-Rochette, J.
Mansourati, Y. Cottin, F. Leclercq, A. Belhassane, N. Delarche, F. Boccara, F. Paganelli, J.
Clerc, F. Schiele, V. Aboyans, V. Probst, J. Berland, T. Lefèvre, V. Chumburidze, I. Khintibidze,
T. Shaburishvili, Z. Pagava, R. Ghlonti, Z. Lominadze, G. Khabeishvili, R. Hemetsberger, U.
Rauch-Kröhnert, M. Stratmann, K. F. Appel, E. Schmidt, H. Omran, C. Stellbrink, T. Dorsel, E.
Lianopoulos, R. Marx, A. Zirlik, D. Schellenberg, T. Heitzer, U. Laufs, N. Marx, S. Gielen, B.
Winkelmann, S. Behrens, K. Sydow, G. Simonis, T. Muenzel, N. Werner, S. Leggewie, D.
Böcker, R. Braun-Dullaeus, N. Toursarkissian, M. Jeserich, M. Weißbrodt, T. Schaeufele, J.
Weil, H. Völler, J. Waltenberger, M. Natour, S. Steiner, L. Heidenreich, U. Gremmler, H. Killat,
S. Patsilinakos, A. Kartalis, A. Manolis, D. Sionis, E. Liberopoulos, I. Skoumas, V. Athyros, P.
Vardas, F. Parthenakis, D. Alexopoulos, G. Hahalis, J. Lekakis, A. Xatzitolios, S. R. Fausto
Ovando, P. C. Montenegro Valdovinos, J. L. Arango Benecke, E. R. Rodriguez De Leon, B. P.
Y. Yan, D. C. W. Siu, T. Turi, B. Merkely, R. G. Kiss, I. Ungi, G. Lupkovics, L. Nagy, A. Katona,
I. Édes, G. Müller, I. Horvath, T. Kapin, J. Faluközy, M. Kumbla, M. Sandhu, S. Annam, N. R.
Proddutur, R. K. Premchand, A. Mahajan, A. D. Abhyanakar, P. Kerkar, R. A. Govinda, A.
Oomman, D. Sinha, S. N. Patil, D. Kahali, J. Sawhney, A. B. Joshi, S. Chaudhary, P. Harkut, S.
Guha, S. Porwal, S. Jujjuru, R. B. Pothineni, M. R. Monteiro, A. Khan, S. S. Iyengar, J. S.
Grewal, M. Chopda, M. C. Fulwani, A. Patange, V. K. Chopra, N. K. Goyal, R. Shinde, G. V.
Manakshe, N. Patki, S. Sethi, V. Munusamy, S. Karna, S. Adhyapak, U. Pandurangi, R. Mathur,
S. Kalashetti, A. Bhagwat, B. Raghuraman, S. K. Yerra, P. Bhansali, R. Borse, S. Das, J.
Abdullakutty, S. Saathe, P. Palimkar, S. Atar, M. Shechter, M. Mosseri, Y. Arbel, C. Lotan, U.
Rosenschein, A. Katz, Y. Henkin, A. Francis, M. Klutstein, E. Nikolsky, Y. Turgeman, M. Halabi,
R. Kornowski, M. Jonas, O. Amir, Y. Rozenman, S. Fuchs, O. Hussein, D. Gavish, Z. Vered, Y.
Caraco, M. Elias, N. Tov, G. Piovaccari, A. De Pellegrin, G. Guardigli, G. Licciardello, C.
Auguadro, C. Cuccia, A. Salvioni, G. Musumeci, P. Calabrò, S. Novo, P. Faggiano, N. B. De
Cesare, S. Berti, C. Cavallini, E. Puccioni, M. Galvani, M. Tespili, P. Piatti, M. Palvarini, G. De
Luca, R. Violini, A. De Leo, P. Perrone Filardi, M. Ferratini, K. Dai, H. Kamiya, K. Ando, Y.
Takeda, Y. Morino, Y. Hata, K. Kimura, K. Kishi, I. Michishita, H. Uehara, T. Higashikata, A.
Hirayama, K. Hirooka, S. Sakagami, S. Taguchi, A. Koike, H. Fujinaga, S. Koba, K. Kozuma, T.
Kawasaki, Y. Ono, M. Shimizu, Y. Katsuda, A. Wada, T. Shinke, T. Kimura, J. Ako, K. Fujii, T.
Takahashi, T. Sakamoto, Y. Furukawa, H. Sugino, T. Mano, N. Utsu, K. Ito, T. Haraguchi, Y.
Ueda, A. Nishibe, K. Fujimoto, J. H. Yoon, S. H. Kim, H. S. Park, I. H. Chae, M. H. Kim, M. H.
Jeong, S. Rha, C. Kim, H. S. Kim, T. Hong, A. Busmane, N. Pontaga, A. Strelnieks, I. Mintale, I.
Sime, Z. Petrulioniene, R. Kavaliauskiene, R. Jurgaitiene, G. Sakalyte, R. Slapikas, S. Norkiene,
N. Misonis, A. Kibarskis, R. Kubilius, S. Bojovski, S. Kedev, N. Lozance, A. Kjovkaroski, S.
Doncovska, T. K. Ong, S. Kasim, O. Maskon, B. Kandasamy, K. Yusoff, H. B. Liew, W. M. I.
Wan Mohamed, A. García Castillo, G. A. Ramos López, J. Carrillo Calvillo, P. Fajardo Campos,
J. C. Núñez Fragoso, E. A. Bayram Llamas, M. A. Alcocer Gamba, J. Carranza Madrigal, L. G.

González Salas, E. López Rosas, B. González Díaz, E. Salcido Vázquez, A. Nacoud Ackar, G.
A. Llamas Esperón, C. R. Martínez Sánchez, M. Guerrero De Leon, R. Suarez Otero, G.
Fanghänel Salmón, J. A. Pérez Ríos, J. A. Garza Ruíz, M. Alings, R. W. Breedveld, P. A. M.
Hoogslag, H. Suryapranata, A. Oomen, J. J. Wiersma, R. M. A. Van Der Wal, P. S. Hooft Van
Huysduynen-Monraats, I. Karalis, G. J. E. Verdel, B. R. G. Brueren, R. P. T. Troquay, E. P.
Viergever, N. Y. Y. Al-Windy, G. L. Bartels, J. H. Cornel, W. R. M. Hermans, J. P. R. Herrman,
R. J. Bos, R. Groutars, C. C. Van Der Zwaan, R. Kaplan, E. Ronner, B. E. Groenemeijer, P. N.
A. Bronzwaer, A. A. H. Liem, B. Rensing, M. Bokern, R. Nijmeijer, F. Hersbach, F. F. Willems,
A. T. M. Gosselink, J. Elliott, G. Wilkins, R. Fisher, D. Scott, H. Hart, R. Stewart, S. Harding, I.
Ternouth, N. Fisher, D. Aitken, R. Anscombe, T. Tomala, O. Nygård, J. A. Sparby, K. Andersen,
L. Gullestad, J. Jortveit, P. S. Munk, S. Halvorsen, U. Hurtig, R. M. Correa Flores, J. R.
Calderon Ticona, J. R. Durand Velasquez, S. A. Negron Miguel, E. S. Sanabria Perez, J. M.
Carrion Chambilla, C. A. Chavez Ayala, R. P. Castillo Leon, R. J. Vargas Gonzales, J. D.
Hernandez Zuniga, L. A. Camacho Cosavalente, J. E. Bravo Mannucci, N. C. Llerena Navarro,
Y. M. Roldan Concha, V. E. Rodriguez Chavez, H. A. Anchante Hernandez, C. A. Zea Nunez,
A. Ferrolino, R. A. G. Sy, L. Tirador, R. G. Sy, G. Matiga, R. M. Coching, A. Bernan, G. Rogelio,
D. D. Morales, E. Tan, A. Wlodarczak, K. Jaworska, G. Skonieczny, L. Pawlowicz, P.
Wojewoda, B. Busz-Papiez, J. Bednarski, A. Goch, P. Staneta, E. Dulak, A. Budaj, K. Saminski,
W. Krasowski, W. Sudnik, A. Zurakowski, M. Skorski, R. Lysek, B. Miklaszewicz, J. Kubica, J. A.
Lipko, E. Kostarska-Srokosz, M. Piepiorka, A. Drzewiecka, R. Sciborski, A. Stasiewski, T.
Blicharski, L. Bystryk, M. Szpajer, M. Korol, T. Czerski, E. Mirek-Bryniarska, J. Gniot, A.
Lubinski, J. Gorny, E. Franek, P. Monteiro, J. Mesquita Bastos, H. H. Pereira, D. Martins, J.
Morais, F. Seixo, C. Mendonça, A. Botelho, B. Minescu, O. Istratoaie, D. N. Tesloianu, M.
Dorobantu, G. Cristian, C. G. C. Podoleanu, M. C. A. Constantinescu, C. M. Bengus, C. Militaru,
D. Rosu, I. R. Parepa, A. V. Matei, T. M. Alexandru, Y. Shvarts, O. Orlikova, Z. Kobalava, O. L.
Barbarash, V. Markov, N. Lyamina, A. Gordienko, K. Zrazhevsky, A. Y. Vishnevsky, V.
Gurevich, R. Stryuk, N. V. Lomakin, I. Bokarev, S. Shalaev, L. Khaisheva, P. Chizhov, I.
Viktorova, N. Osokina, E. Akatova, G. Chumakova, I. Libov, M. I. Voevoda, T. V. Tretyakova, E.
Baranov, S. Shustov, S. Yakushin, I. Gordeev, N. Khasanov, O. Reshetko, T. Sotnikova, O.
Molchanova, K. Y. Nikolaev, L. Gapon, E. Baranova, Z. Shogenov, E. Kosmachova, Y. Karpov,
A. Povzun, L. Egorova, V. V. Tyrenko, I. G. Ivanov, D. Simic, N. Ivanovic, G. Davidovic, N.
Tasic, M. R. Asanin, S. Stojic, S. R. Apostolovic, S. Ilic, B. Putnikovic, A. Stankovic, A.
Arandjelovic, S. Radovanovic, A. D. Ristic, J. Balinovac, D. V. Dincic, P. Seferovic, S. Dodic, S.
Dimkovic, T. Chua, K. K. Poh, H. Y. Ong, K. Micko, J. Nociar, D. Pella, P. Fulop, M. Hranai, J.
Palka, J. Mazur, I. Majercák, A. Dzupina, F. Fazekas, J. Gonsorcik, V. Bugan, J. Murin, J.
Selecky, G. Kamensky, J. Strbova, R. Smik, A. Dukat, I. Žuran, J. Oklukar, N. C. Šuligoj, M.
Cevc, L. Lipar, H. P. Cyster, N. Ranjith, C. Corbett, J. Bayat, E. M. Makotoko, I. E. Kapp, M. M.
V. Basson, H. Lottering, L. J. Van Zyl, P. J. Sebastian, T. Pillay, J. A. Saaiman, P. J.
Commerford, S. Cassimjee, I. O. Ebrahim, M. Sarvan, J. H. Mynhardt, A. J. Dalby, H. Reuter, R.
Moodley, M. Vida, A. R. Cequier Fillat, V. Bodí Peris, F. Fuentes Jimenez, F. Marín, J. M. Cruz
Fernández, B. Gil-Extremera, F. W. Diz, D. Garcia-Dorado, A. Iñiguez, J. Tuñón Fernández, J.
R. Gonzalez-Juanatey, J. Fernandez Portales, F. Civeira Murillo, L. Matas Pericas, J. L.
Zamorano, M. De Mora Martin, J. Bruguera Cortada, J. J. Alonso Martin, J. R. De Berrazueta
Fernández, J. F. Díaz Fernández, J. A. García Lledó, J. Cosín Sales, J. Botas Rodriguez, G.
Gusi Tragant, A. Benedicto, C. Gonzalez-Juanatey, M. Camprubí Potau, I. Plaza Perez, C. M.
De La Tassa, P. Loma-Osorio Rincon, J. Balaguer Recena, J. M. Escudier, G. Constantine, R.
Haniffa, N. Tissera, S. Amarasekera, N. Fernando, J. Jayawardena, W. Santharaj, R.
Ekanayaka, S. Mendis, V. Senaratne, G. Mayurathan, T. Sirisena, A. Rajapaksha, J. I. Herath,
N. Amarasena, S. Berglund, G. Rasmanis, E. Hagström, N. Witt, G. Mourtzinis, P. Nicol, O.
Hansen, S. Romeo, S. A. Jensen, I. Torstensson, U. Ahremark, T. Sundelin, T. Moccetti, C.
Müller, F. Mach, R. Binder, C. E. Chiang, W. C. Tsai, K. C. Ueng, W. T. Lai, M. E. Liu, J. J.

Hwang, W. H. Yin, I. C. Hsieh, W. H. Lin, J. Y. Kuo, T. Y. Huang, C. Y. Fang, P. Kaewsuwanna, W. Soonfuang, W. Jintapakorn, A. Sukonthasarn, P. Sritara, N. Wongpraparut, K. Sastravaha, N. Sansanayudh, W. Kehasukcharoen, D. Piyayotai, A. Camsari, H. Kultursay, S. Guneri, B. Mutlu, M. Ersanli, M. Demirtas, C. Kirma, E. Ural, L. Koldas, O. Karpenko, A. Prokhorov, I. Vakaluyk, H. Myshanych, D. Reshotko, V. Batushkin, L. Rudenko, I. Kovalskyi, M. Kushnir, V. Tseluyko, Y. Mostovoy, M. Stanislavchuk, Y. Kyiak, Y. Karpenko, Y. Malynovsky, A. Klantsa, O. Kutniy, E. Amosova, V. Tashchuk, O. Leshchuk, A. Parkhomenko, M. Rishko, M. Kopytsya, A. Yagensky, M. Vatutin, A. Bagriy, O. M. Barna, O. Ushakov, G. Dzyak, B. Goloborodko, A. Rudenko, J. Trevelyan, A. Zaman, K. Lee, A. Moriarty, R. K. Aggarwal, P. Clifford, Y. K. Wong, S. M. R. Iqbal, E. Subkovas, D. Braganza, D. Sarkar, R. Storey, H. Griffiths, S. Mcclure, R. Muthusamy, J. Kurian, T. Levy, C. Barr, H. Kadr, R. Gerber, A. Simaitis, H. Soran, A. Mathur, A. Brodison, R. Oliver, T. Mudawi, T. Reynolds, D. Sharman, R. Butler, P. Wilkinson, G. Y. H. Lip, J. Halcox, G. Vardi, D. Baldari, D. Brabham, C. Treasure, C. Dahl, B. Palmer, A. Wiseman, S. Puri, A. E. Mohart, C. Ince, E. Flores, S. Wright, S. C. Cheng, M. Rosenberg, W. Rogers, E. Kosinski, L. Forgosh, J. Waltman, M. Khan, M. Shoukfeh, G. Dagher, I. Lieber, P. Kumar, C. East, P. Krichmar, L. White, T. Knickelbine, T. Haldis, E. Gillespie, D. Suh, I. Arif, F. Akhter, E. Carlson, M. D'Urso, F. El-Ahdab, W. Nelson, B. Harris, S. Cohen, L. Carter, K. Sabatino, T. Haddad, A. Malik, S. Rao, A. Mulkay, I. Jovin, K. Klancke, V. Malhotra, S. K. Devarapalli, M. Koren, H. Chandna, G. Dodds, M. Janik, J. Moran, A. Sumner, J. Kobayashi, W. Davis, S. Yazdani, J. Pasquini, M. Thakkar, A. Vedere, W. Leimbach, J. Rider, N. Singh, A. V. Shah, P. M. Moriarty, D. Janosik, C. Pepine, B. Berman, J. Gelormini, C. Daniels, F. Keating, N. I. Kondo, S. Shetty, W. Waider, T. Takata, M. Abu-Fadel, V. Shah, R. Aggarwal, M. Izzo, A. Kumar, B. Hattler, C. Link, A. Bortnick, G. Kinzfogl, A. Ghitis, J. Larry, E. Teufel, P. Kuhlman, B. Mclaurin, W. Zhang, S. Thew, J. Abbas, M. White, N. Ranadive, C. Gring, D. Henderson, T. Schuchard, N. Farhat, G. Kline, S. Mahal, J. Whitaker, S. Speirs, R. Andersen, N. Daboul, P. Horwitz, Z. Jafar, J. Mcgarvey, V. Panchal, S. Voyce, T. Blok, W. Sheldon, M. M. Azizad, C. Schmalfuss, M. Picone, W. Herzog, J. Lindsey, R. Nowins, N. Lepor, M. El Shahawy, H. Weintraub, A. Irimpen, W. May, T. Galski, A. Chu, F. Mody, Z. Hodes, J. Fairlamb, C. Lambert, A. Raisinghani, A. Abbate, M. King, C. Carey, J. Gerber, L. Younis, H. T. Park, M. Vidovich, T. Knutson, D. Friedman, F. Chaleff, A. Loussararian, C. Kimmelstiel, K. Silver, M. Foster, G. Tonnessen, M. Amlani, A. Wali, C. Malozzi, K. Wattanakit, P. J. O'Donnell, D. Singal, N. Jaffrani, S. Banuru, D. Fisher, M. Xenakis, N. Perlmutter, R. Bhagwat, J. Strader, A. Akyea-Djamson, A. Labroo, H. J. Marais, E. Claxton, M. Berk, P. Rossi, P. Joshi, A. S. Khaira, G. Kumkumian, S. Lupovitch, J. Purow, S. Welka, D. Hoffman, S. Fischer, E. Soroka, D. Eagerton, S. Pancholy, M. Ray, M. Farrar, S. Pollock, W. J. French, S. Diamantis, L. Gimple, S. Schwartz, E. Pereira, D. Spriggs, J. Strain, A. Vo, M. Chane, J. Hall, N. Vijay, K. Lotun, F. M. Lester, A. Nahhas, T. Pope, P. Nager, R. Vohra, R. Bashir, H. Ahmed, M. Berlowitz, R. Fishberg, R. Barrucco, E. Yang, M. Radin, D. Sporn, S. Eisenberg, J. Landzberg, M. Mcgough, S. Turk, M. Schwartz, P. S. Sundram, D. Jain, M. Zainea, C. Bayron, R. Karlsberg, H. Lui, W. Keen, D. Westerhausen, S. Khurana, H. Agarwal, J. Birchem, W. Penny, M. Chang, J. M. Gilbert, G. Chalavarya, C. Eaton, J. F. Schmedtje, S. Christenson, D. Denham, A. Macdonell, P. Gibson, A. Rahman, T. Al Joundi, G. Conrad, P. Kotha, M. Love, G. Giesler, H. Rubenstein, L. Akright, B. Schifferdecker, J. Krawczyk, T. Wells, J. Welker, R. Foster, R. Gilmore, J. Anderson, D. Jacoby, G. Gardner, R. Dandillaya, K. Vora, J. Kostis, J. Hunter, D. Laxson, E. Ball, Alirocumab and cardiovascular outcomes after acute coronary syndrome, *N. Engl. J. Med.* (2018), doi:10.1056/NEJMoa1801174.

7. M. S. Sabatine, R. P. Giugliano, A. C. Keech, N. Honarpour, S. D. Wiviott, S. A. Murphy, J. F. Kuder, H. Wang, T. Liu, S. M. Wasserman, P. S. Sever, T. R. Pedersen, FOURIER Steering Committee and Investigators, Evolocumab and Clinical Outcomes in Patients with Cardiovascular Disease, *N. Engl. J. Med.* **376**, 1713–1722 (2017).

8. C. Ma, M. E. Gurol, Z. Huang, A. H. Lichtenstein, X. Wang, Y. Wang, S. Neumann, S. Wu, X.

Gao, Low-density lipoprotein cholesterol and risk of intracerebral hemorrhage, *Neurology* (2019), doi:10.1212/WNL.0000000000007853.

9. M. S. Sabatine, S. D. Wiviott, K. Im, S. A. Murphy, R. P. Giugliano, Efficacy and safety of further lowering of low-density lipoprotein cholesterol in patients starting with very low levels: A meta-analysis, *JAMA Cardiol.* (2018), doi:10.1001/jamacardio.2018.2258.

10. R. P. Giugliano, T. R. Pedersen, J.-G. Park, G. M. De Ferrari, Z. A. Gaciong, R. Ceska, K. Toth, I. Gouni-Berthold, J. Lopez-Miranda, F. Schiele, F. Mach, B. R. Ott, E. Kanevsky, A. L. Pineda, R. Somaratne, S. M. Wasserman, A. C. Keech, P. S. Sever, M. S. Sabatine, FOURIER Investigators, Clinical efficacy and safety of achieving very low LDL-cholesterol concentrations with the PCSK9 inhibitor evolocumab: a prespecified secondary analysis of the FOURIER trial, *Lancet* **390**, 1962–1971 (2017).

11. A. Taylor, D. Wang, K. Patel, R. Whittall, G. Wood, M. Farrer, R. D. Neely, S. Fairgrieve, D. Nair, M. Barbir, J. L. Jones, S. Egan, R. Everdale, Y. Lolin, E. Hughes, J. A. Cooper, S. G. Hadfield, G. Norbury, S. E. Humphries, Mutation detection rate and spectrum in familial hypercholesterolaemia patients in the UK pilot cascade project, *Clin. Genet.* **77**, 572–580 (2010).

12. A. Garg, S. Fazio, P. B. Duell, A. Baass, C. Udata, T. Joh, T. Riel, M. Sirota, D. Dettling, H. Liang, P. D. Garzone, B. Gumbiner, H. Wan, Molecular Characterization of Familial Hypercholesterolemia in a North American Cohort, *J. Endocr. Soc.* (2019), doi:10.1210/jendso/bvz015.

13. L. A. Gilbert, M. A. Horlbeck, B. Adamson, J. E. Villalta, Y. Chen, E. H. Whitehead, C. Guimaraes, B. Panning, H. L. Ploegh, M. C. Bassik, L. S. Qi, M. Kampmann, J. S. Weissman, Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation, *Cell* **159**, 647–661 (2014).

14. M. A. Horlbeck, L. A. Gilbert, J. E. Villalta, B. Adamson, R. A. Pak, Y. Chen, A. P. Fields, C. Y. Park, J. E. Corn, M. Kampmann, J. S. Weissman, Compact and highly active next-generation libraries for CRISPR-mediated gene repression and activation., *Elife* **5** (2016), doi:10.7554/eLife.19760.

15. B. Adamson, T. M. Norman, M. Jost, M. Y. Cho, J. K. Nuñez, Y. Chen, J. E. Villalta, L. A. Gilbert, M. A. Horlbeck, M. Y. Hein, R. A. Pak, A. N. Gray, C. A. Gross, A. Dixit, O. Parnas, A. Regev, J. S. Weissman, A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response., *Cell* **167**, 1867-1882.e21 (2016).

16. K. K. Ray, R. S. Wright, D. Kallend, W. Koenig, L. A. Leiter, F. J. Raal, J. A. Bisch, T. Richardson, M. Jaros, P. L. J. Wijngaard, J. J. P. Kastelein, Two phase 3 trials of inclisiran in patients with elevated LDL cholesterol, *N. Engl. J. Med.* (2020), doi:10.1056/NEJMoa1912387.

17. B. B. Knowles, C. C. Howe, D. P. Aden, Human hepatocellular carcinoma cell lines secrete the major plasma proteins and hepatitis B surface antigen, *Science (80-.).* (1980), doi:10.1126/science.6248960.

18. M. A. Mandegar, N. Huebsch, E. B. Frolov, E. Shin, A. Truong, M. P. Olvera, A. H. Chan, Y. Miyaoka, K. Holmes, C. I. Spencer, L. M. Judge, D. E. Gordon, T. V. Eskildsen, J. E. Villalta, M. A. Horlbeck, L. A. Gilbert, N. J. Krogan, S. P. Sheikh, J. S. Weissman, L. S. Qi, P. L. So, B. R. Conklin, CRISPR Interference Efficiently Induces Specific and Reversible Gene Silencing in Human iPSCs, *Cell Stem Cell* (2016), doi:10.1016/j.stem.2016.01.022.

19. A. W. Alberts, J. Chen, G. Kuron, V. Hunt, J. Huff, C. Hoffman, J. Rothrock, M. Lopez, H. Joshua, E. Harris, A. Patchett, R. Monaghan, S. Currie, E. Stapley, G. Albers-Schonberg, O. Hensens, J. Hirshfield, K. Hoogsteen, J. Liesch, J. Springer, Mevinolin: A highly potent competitive inhibitor of hydroxymethylglutaryl-coenzyme A reductase and a cholesterol-lowering agent, *Proc. Natl. Acad. Sci. U. S. A.* (1980), doi:10.1073/pnas.77.7.3957.

20. M. S. Brown, J. R. Faust, J. L. Goldstein, I. Kaneko, A. Endo, Induction of 3-hydroxy-3-methylglutaryl coenzyme A reductase activity in human fibroblasts incubated with compactin (ML-236B), a competitive inhibitor of the reductase, *J. Biol. Chem.* (1978).

21. S. Benjannet, D. Rhainds, R. Essalmani, J. Mayne, L. Wickham, W. Jin, M.-C. Asselin, J.
Hamelin, M. Varret, D. Allard, M. Trillard, M. Abifadel, A. Tebon, A. D. Attie, D. J. Rader, C.
Boileau, L. Brissette, M. Chrétien, A. Prat, N. G. Seidah, NARC-1/PCSK9 and its natural
mutants: zymogen cleavage and effects on the low density lipoprotein (LDL) receptor and LDL
cholesterol., *J. Biol. Chem.* **279**, 48865–75 (2004).
22. S. Rashid, D. E. Curtis, R. Garuti, N. N. Anderson, Y. Bashmakov, Y. K. Ho, R. E. Hammer,
Y. A. Moon, J. D. Horton, Decreased plasma cholesterol and hypersensitivity to statins in mice
lacking Pcsk9, *Proc. Natl. Acad. Sci. U. S. A.* **102**, 5374–5379 (2005).
23. J. C. Chan, D. E. Piper, Q. Cao, D. Liu, C. King, W. Wang, J. Tang, Q. Liu, J. Higbee, Z. Xia,
Y. Di, S. Shetterly, Z. Arimura, H. Salomonis, W. G. Romanow, S. T. Thibault, R. Zhang, P.
Cao, X. P. Yang, T. Yu, M. Lu, M. W. Retter, G. Kwon, K. Henne, O. Pan, M. M. Tsai, B.
Fuchslocher, E. Yang, L. Zhou, K. J. Lee, M. Daris, J. Sheng, Y. Wang, W. D. Shen, W. C. Yeh,
M. Emery, N. P. Walker, B. Shan, M. Schwarz, S. M. Jackson, A proprotein convertase
subtilisin/kexin type 9 neutralizing antibody reduces serum cholesterol in mice and nonhuman
primates, *Proc. Natl. Acad. Sci. U. S. A.* **106**, 9820–9825 (2009).
24. P. Aisen, Transferrin receptor 1., *Int. J. Biochem. Cell Biol.* **36**, 2137–43 (2004).
25. R. Coffey, T. Ganz, Iron homeostasis: An anthropocentric perspective, *J. Biol. Chem.* **292**,
12727–12734 (2017).
26. N. Zelcer, C. Hong, R. Boyadjian, P. Tontonoz, LXR regulates cholesterol uptake through
Idol-dependent ubiquitination of the LDL receptor, *Science* **325**, 100–104 (2009).
27. M. Yoshinaga, Y. Nakatsuka, A. Vandenbon, D. Ori, T. Uehata, T. Tsujimura, Y. Suzuki, T.
Mino, O. Takeuchi, Regnase-1 Maintains Iron Homeostasis via the Degradation of Transferrin
Receptor 1 and Prolyl-Hydroxylase-Domain-Containing Protein 3 mRNAs, *Cell Rep.* (2017),
doi:10.1016/j.celrep.2017.05.009.
28. H. Mi, A. Muruganujan, D. Ebert, X. Huang, P. D. Thomas, PANTHER version 14: more
genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools, *Nucleic*
Acids Res. **47**, D419–D426 (2019).
29. T. M. Teslovich, K. Musunuru, A. V. Smith, A. C. Edmondson, I. M. Stylianou, M. Koseki, J.
P. Pirruccello, S. Ripatti, D. I. Chasman, C. J. Willer, C. T. Johansen, S. W. Fouchier, A. Isaacs,
G. M. Peloso, M. Barbalic, S. L. Ricketts, J. C. Bis, Y. S. Aulchenko, G. Thorleifsson, M. F.
Feitosa, J. Chambers, M. Orho-Melander, O. Melander, T. Johnson, X. Li, X. Guo, M. Li, Y. Shin
Cho, M. Jin Go, Y. Jin Kim, J.-Y. Lee, T. Park, K. Kim, X. Sim, R. Twee-Hee Ong, D. C.
Croteau-Chonka, L. A. Lange, J. D. Smith, K. Song, J. Hua Zhao, X. Yuan, J. Luan, C. Lamina,
A. Ziegler, W. Zhang, R. Y. L. Zee, A. F. Wright, J. C. M. Witteman, J. F. Wilson, G. Willemsen,
H.-E. Wichmann, J. B. Whitfield, D. M. Waterworth, N. J. Wareham, G. Waeber, P.
Vollenweider, B. F. Voight, V. Vitart, A. G. Uitterlinden, M. Uda, J. Tuomilehto, J. R. Thompson,
T. Tanaka, I. Surakka, H. M. Stringham, T. D. Spector, N. Soranzo, J. H. Smit, J. Sinisalo, K.
Silander, E. J. G. Sijbrands, A. Scuteri, J. Scott, D. Schlessinger, S. Sanna, V. Salomaa, J.
Saharinen, C. Sabatti, A. Ruokonen, I. Rudan, L. M. Rose, R. Roberts, M. Rieder, B. M. Psaty,
P. P. Pramstaller, I. Pichler, M. Perola, B. W. J. H. Penninx, N. L. Pedersen, C. Pattaro, A. N.
Parker, G. Pare, B. A. Oostra, C. J. O'Donnell, M. S. Nieminen, D. A. Nickerson, G. W.
Montgomery, T. Meitinger, R. McPherson, M. I. McCarthy, W. McArdle, D. Masson, N. G.
Martin, F. Marroni, M. Mangino, P. K. E. Magnusson, G. Lucas, R. Luben, R. J. F. Loos, M.-L.
Lokki, G. Lettre, C. Langenberg, L. J. Launer, E. G. Lakatta, R. Laaksonen, K. O. Kyvik, F.
Kronenberg, I. R. König, K.-T. Khaw, J. Kaprio, L. M. Kaplan, Å. Johansson, M.-R. Jarvelin, A.
Cecile J. W. Janssens, E. Ingelsson, W. Igl, G. Kees Hovingh, J.-J. Hottenga, A. Hofman, A. A.
Hicks, C. Hengstenberg, I. M. Heid, C. Hayward, A. S. Havulinna, N. D. Hastie, T. B. Harris, T.
Haritunians, A. S. Hall, U. Gyllensten, C. Guiducci, L. C. Groop, E. Gonzalez, C. Gieger, N. B.
Freimer, L. Ferrucci, J. Erdmann, P. Elliott, K. G. Ejebe, A. Döring, A. F. Dominiczak, S.
Demissie, P. Deloukas, E. J. C. de Geus, U. de Faire, G. Crawford, F. S. Collins, Y. I. Chen, M.
J. Caulfield, H. Campbell, N. P. Burtt, L. L. Bonnycastle, D. I. Boomsma, S. M. Boekholdt, R. N.

Bergman, I. Barroso, S. Bandinelli, C. M. Ballantyne, T. L. Assimes, T. Quertermous, D. Altshuler, M. Seielstad, T. Y. Wong, E.-S. Tai, A. B. Feranil, C. W. Kuzawa, L. S. Adair, H. A. Taylor Jr, I. B. Borecki, S. B. Gabriel, J. G. Wilson, H. Holm, U. Thorsteinsdottir, V. Gudnason, R. M. Krauss, K. L. Mohlke, J. M. Ordovas, P. B. Munroe, J. S. Kooner, A. R. Tall, R. A. Hegele, J. J. P. Kastelein, E. E. Schadt, J. I. Rotter, E. Boerwinkle, D. P. Strachan, V. Mooser, K. Stefansson, M. P. Reilly, N. J. Samani, H. Schunkert, L. A. Cupples, M. S. Sandhu, P. M. Ridker, D. J. Rader, C. M. van Duijn, L. Peltonen, G. R. Abecasis, M. Boehnke, S. Kathiresan, Biological, clinical and population relevance of 95 loci for blood lipids, *Nature* **466**, 707–713 (2010).

30. P. Natarajan, G. M. Peloso, S. M. Zekavat, M. Montasser, A. Ganna, M. Chaffin, A. V. Khera, W. Zhou, J. M. Bloom, J. M. Engreitz, J. Ernst, J. R. O'Connell, S. E. Ruotsalainen, M. Alver, A. Manichaikul, W. C. Johnson, J. A. Perry, T. Poterba, C. Seed, I. L. Surakka, T. Esko, S. Ripatti, V. Salomaa, A. Correa, R. S. Vasan, M. Kellis, B. M. Neale, E. S. Lander, G. Abecasis, B. Mitchell, S. S. Rich, J. G. Wilson, L. A. Cupples, J. I. Rotter, C. J. Willer, S. Kathiresan, NHLBI TOPMed Lipids Working Group, Deep-coverage whole genome sequences and blood lipids among 16,324 individuals, *Nat. Commun.* **9**, 3391 (2018).

31. D. J. Liu, G. M. Peloso, H. Yu, A. S. Butterworth, X. Wang, A. Mahajan, D. Saleheen, C. Emdin, D. Alam, A. C. Alves, P. Amouyel, E. Di Angelantonio, D. Arveiler, T. L. Assimes, P. L. Auer, U. Baber, C. M. Ballantyne, L. E. Bang, M. Benn, J. C. Bis, M. Boehnke, E. Boerwinkle, J. Bork-Jensen, E. P. Bottinger, I. Brandslund, M. Brown, F. Busonero, M. J. Caulfield, J. C. Chambers, D. I. Chasman, Y. E. Chen, Y.-D. I. Chen, R. Chowdhury, C. Christensen, A. Y. Chu, J. M. Connell, F. Cucca, L. A. Cupples, S. M. Damrauer, G. Davies, I. J. Deary, G. Dedoussis, J. C. Denny, A. Dominiczak, M.-P. Dubé, T. Ebeling, G. Eiriksdottir, T. Esko, A.-E. Farmaki, M. F. Feitosa, M. Ferrario, J. Ferrieres, I. Ford, M. Fornage, P. W. Franks, T. M. Frayling, R. Frikke-Schmidt, L. G. Fritsche, P. Frossard, V. Fuster, S. K. Ganesh, W. Gao, M. E. Garcia, C. Gieger, F. Giulianini, M. O. Goodarzi, H. Grallert, N. Grarup, L. Groop, M. L. Grove, V. Gudnason, T. Hansen, T. B. Harris, C. Hayward, J. N. Hirschhorn, O. L. Holmen, J. Huffman, Y. Huo, K. Hveem, S. Jabeen, A. U. Jackson, J. Jakobsdottir, M.-R. Jarvelin, G. B. Jensen, M. E. Jørgensen, J. W. Jukema, J. M. Justesen, P. R. Kamstrup, S. Kanoni, F. Karpe, F. Kee, A. V Khera, D. Klarin, H. A. Koistinen, J. S. Kooner, C. Kooperberg, K. Kuulasmaa, J. Kuusisto, M. Laakso, T. Lakka, C. Langenberg, A. Langsted, L. J. Launer, T. Lauritzen, D. C. M. Liewald, L. A. Lin, A. Linneberg, R. J. F. Loos, Y. Lu, X. Lu, R. Mägi, A. Malarstig, A. Manichaikul, A. K. Manning, P. Mäntyselkä, E. Marouli, N. G. D. Masca, A. Maschio, J. B. Meigs, O. Melander, A. Metspalu, A. P. Morris, A. C. Morrison, A. Mulas, M. Müller-Nurasyid, P. B. Munroe, M. J. Neville, J. B. Nielsen, S. F. Nielsen, B. G. Nordestgaard, J. M. Ordovas, R. Mehran, C. J. O'Donnell, M. Orho-Melander, C. M. Molony, P. Muntendam, S. Padmanabhan, C. N. A. Palmer, D. Pasko, A. P. Patel, O. Pedersen, M. Perola, A. Peters, C. Pisinger, G. Pistis, O. Polasek, N. Poulter, B. M. Psaty, D. J. Rader, A. Rasheed, R. Rauramaa, D. F. Reilly, A. P. Reiner, F. Renström, S. S. Rich, P. M. Ridker, J. D. Rioux, N. R. Robertson, D. M. Roden, J. I. Rotter, I. Rudan, V. Salomaa, N. J. Samani, S. Sanna, N. Sattar, E. M. Schmidt, R. A. Scott, P. Sever, R. S. Sevilla, C. M. Shaffer, X. Sim, S. Sivapalaratnam, K. S. Small, A. V Smith, B. H. Smith, S. Somayajula, L. Southam, T. D. Spector, E. K. Speliotes, J. M. Starr, K. E. Stirrups, N. Stitziel, K. Strauch, H. M. Stringham, P. Surendran, H. Tada, A. R. Tall, H. Tang, J.-C. Tardif, K. D. Taylor, S. Trompet, P. S. Tsao, J. Tuomilehto, A. Tybjaerg-Hansen, N. R. van Zuydam, A. Varbo, T. V Varga, J. Virtamo, M. Waldenberger, N. Wang, N. J. Wareham, H. R. Warren, P. E. Weeke, J. Weinstock, J. Wessel, J. G. Wilson, P. W. F. Wilson, M. Xu, H. Yaghootkar, R. Young, E. Zeggini, H. Zhang, N. S. Zheng, W. Zhang, Y. Zhang, W. Zhou, Y. Zhou, M. Zoledziewska, J. M. M. Howson, J. Danesh, M. I. McCarthy, C. A. Cowan, G. Abecasis, P. Deloukas, K. Musunuru, C. J. Willer, S. Kathiresan, G. Abecasis, P. Deloukas, K. Musunuru, C. J. Willer, S. Kathiresan, Exome-wide association study of plasma lipids in >300,000 individuals, *Nat. Genet.* **49**, 1758–1766 (2017).

32. C. Bycroft, C. Freeman, D. Petkova, G. Band, L. T. Elliott, K. Sharp, A. Motyer, D. Vukcevic,
O. Delaneau, J. O'Connell, A. Cortes, S. Welsh, A. Young, M. Effingham, G. McVean, S. Leslie,
N. Allen, P. Donnelly, J. Marchini, The UK Biobank resource with deep phenotyping and
genomic data, *Nature* (2018), doi:10.1038/s41586-018-0579-z.
33. A. Loregger, J. K. Nelson, N. Zelcer, in *Methods in molecular biology (Clifton, N.J.)*, (2017),
vol. 1583, pp. 53–63.
34. A. Mazein, S. Watterson, W. Y. Hsieh, W. J. Griffiths, P. Ghazal, A comprehensive machine-
readable view of the mammalian cholesterol biosynthesis pathway, *Biochem. Pharmacol.*
(2013), doi:10.1016/j.bcp.2013.03.021.
35. R. B. Rose, J. H. Bayle, J. A. Endrizzi, J. D. Cronk, G. R. Crabtree, T. Alber, Structural basis
of dimerization, coactivator recognition and MODY3 mutations in HNF-1α, *Nat. Struct. Biol.*
(2000), doi:10.1038/78966.
36. A. Delaforest, F. Di Furio, R. Jing, A. Ludwig-Kubinski, K. Twaroski, A. Urick, K. Pulakanti,
S. Rao, S. A. Duncan, HNF4A regulates the formation of hepatic progenitor cells from human
iPSC-derived endoderm by facilitating efficient recruitment of RNA pol II, *Genes (Basel).* (2019),
doi:10.3390/genes10010021.
37. K. Wang, A. X. Holterman, Pathophysiologic role of hepatocyte nuclear factor 6*Cell. Signal.*
(2012), doi:10.1016/j.cellsig.2011.08.009.
38. U. Wellner, J. Schubert, U. C. Burk, O. Schmalhofer, F. Zhu, A. Sonntag, B. Waldvogel, C.
Vannier, D. Darling, A. Zur Hausen, V. G. Brunton, J. Morton, O. Sansom, J. Schüler, M. P.
Stemmler, C. Herzberger, U. Hopt, T. Keck, S. Brabletz, T. Brabletz, The EMT-activator ZEB1
promotes tumorigenicity by repressing stemness-inhibiting microRNAs, *Nat. Cell Biol.* (2009),
doi:10.1038/ncb1998.
39. C. L. Gao, J. G. Zhu, Y. P. Zhao, X. H. Chen, C. B. Ji, C. M. Zhang, C. Zhu, Z. K. Xia, Y. Z.
Peng, X. R. Guo, Mitochondrial dysfunction is induced by the overexpression of UCP4 in 3T3-L1
adipocytes, *Int. J. Mol. Med.* (2010), doi:10.3892/ijmm_00000315.
40. F. Quazi, R. S. Molday, Differential phospholipid substrates and directional transport by
ATP-binding cassette proteins ABCA1, ABCA7, and ABCA4 and disease-causing mutants, *J.*
Biol. Chem. (2013), doi:10.1074/jbc.M113.508812.
41. J. E. Kung, N. Jura, The pseudokinase TRIB 1 toggles an intramolecular switch to regulate
COP 1 nuclear export , *EMBO J.* (2019), doi:10.15252/embj.201899708.
42. J. M. Murphy, Y. Nakatani, S. A. Jamieson, W. Dai, I. S. Lucet, P. D. Mace, Molecular
Mechanism of CCAAT-Enhancer Binding Protein Recruitment by the TRIB1 Pseudokinase,
Structure (2015), doi:10.1016/j.str.2015.08.017.
43. S. Soubeyrand, A. Martinuk, R. McPherson, TRIB1 is a positive regulator of hepatocyte
nuclear factor 4-Alpha, *Sci. Rep.* (2017), doi:10.1038/s41598-017-05768-1.
44. R. C. Bauer, M. Sasaki, D. M. Cohen, J. Cui, M. A. Smith, B. O. Yenilmez, D. J. Steger, D. J.
Rader, Tribbles-1 regulates hepatic lipogenesis through posttranscriptional regulation of
C/EBPα, *J. Clin. Invest.* (2015), doi:10.1172/JCI77095.
45. R. Burkhardt, S. A. Toh, W. R. Lagor, A. Birkeland, M. Levin, X. Li, M. Robblee, V. D.
Fedorov, M. Yamamoto, T. Satoh, S. Akira, S. Kathiresan, J. L. Breslow, D. J. Rader, Trib1 is a
lipid- and myocardial infarction-associated gene that regulates hepatic lipogenesis and VLDL
production in mice, *J. Clin. Invest.* **120**, 4410–4414 (2010).
46. A. Motley, N. A. Bright, M. N. J. Seaman, M. S. Robinson, Clathrin-mediated endocytosis in
AP-2-depleted cells, *J. Cell Biol.* (2003), doi:10.1083/jcb.200305145.
47. S. F. Parsons, G. Mallinson, C. H. Holmes, J. M. Houlihan, K. L. Simpson, W. J. Mawby, N.
K. Spurr, D. Warne, A. N. Barclay, D. J. Anstee, The Lutheran blood group glycoprotein, another
member of the immunoglobulin superfamily, is widely expressed in human tissues and is
developmentally regulated in human liver, *Proc. Natl. Acad. Sci. U. S. A.* (1995),
doi:10.1073/pnas.92.12.5496.
48. S. K. Mishra, P. A. Keyel, M. A. Edeling, A. L. Dupin, D. J. Owen, L. M. Traub, Functional

dissection of an AP-2 β2 appendage-binding sequence within the autosomal recessive hypercholesterolemia protein, *J. Biol. Chem.* (2005), doi:10.1074/jbc.M501029200.

49. A. Radhakrishnan, J. L. Goldstein, J. G. McDonald, M. S. Brown, Switch-like Control of SREBP-2 Transport Triggered by Small Changes in ER Cholesterol: A Delicate Balance, *Cell Metab.* (2008), doi:10.1016/j.cmet.2008.10.008.

50. J. D. Horton, N. A. Shah, J. A. Warrington, N. N. Anderson, S. W. Park, M. S. Brown, J. L. Goldstein, Combined analysis of oligonucleotide microarray data from transgenic and knockout mice identifies direct SREBP target genes., *Proc. Natl. Acad. Sci. U. S. A.* **100**, 12027–32 (2003).

51. Scandinavian Simvastatin Survival Study Group, Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S), *Lancet* (1994), doi:10.1016/S0140-6736(94)90566-5.

52. N. G. Lintner, K. F. McClure, D. Petersen, A. T. Londregan, D. W. Piotrowski, L. Wei, J. Xiao, M. Bolt, P. M. Loria, B. Maguire, K. F. Geoghegan, A. Huang, T. Rolph, S. Liras, J. A. Doudna, R. G. Dullea, J. H. D. Cate, C. Khosla, Ed. Selective stalling of human translation through small-molecule engagement of the ribosome nascent chain., *PLoS Biol.* **15**, e2001882 (2017).

53. W. Li, F. R. Ward, K. F. McClure, S. T.-L. Chang, E. Montabana, S. Liras, R. G. Dullea, J. H. D. Cate, Structural basis for selective stalling of human ribosome nascent chain complexes by a drug-like molecule, *Nat. Struct. Mol. Biol.* **26**, 501–509 (2019).

54. T. Suzuki, M. Terasaki, C. Takemoto-Hori, T. Hanada, T. Ueda, A. Wada, K. Watanabe, Structural compensation for the deficit of rRNA with proteins in the mammalian mitochondrial ribosome. Systematic analysis of protein components of the large ribosomal subunit from mammalian mitochondria, *J. Biol. Chem.* (2001), doi:10.1074/jbc.M100432200.

55. P. J. Thul, L. Akesson, M. Wiking, D. Mahdessian, A. Geladaki, H. Ait Blal, T. Alm, A. Asplund, L. Björk, L. M. Breckels, A. Bäckström, F. Danielsson, L. Fagerberg, J. Fall, L. Gatto, C. Gnann, S. Hober, M. Hjelmare, F. Johansson, S. Lee, C. Lindskog, J. Mulder, C. M. Mulvey, P. Nilsson, P. Oksvold, J. Rockberg, R. Schutten, J. M. Schwenk, A. Sivertsson, E. Sjöstedt, M. Skogs, C. Stadler, D. P. Sullivan, H. Tegel, C. Winsnes, C. Zhang, M. Zwahlen, A. Mardinoglu, F. Pontén, K. Von Feilitzen, K. S. Lilley, M. Uhlén, E. Lundberg, A subcellular map of the human proteome, *Science (80-.).* (2017), doi:10.1126/science.aal3321.

56. H. Li, B. Dong, S. W. Park, H.-S. Lee, W. Chen, J. Liu, Hepatocyte nuclear factor 1alpha plays a critical role in PCSK9 gene transcription and regulation by the natural hypocholesterolemic compound berberine., *J. Biol. Chem.* **284**, 28885–95 (2009).

57. A.-X. Guo, J.-J. Cui, L.-Y. Wang, J.-Y. Yin, The role of CSDE1 in translational reprogramming and human diseases, *Cell Commun. Signal.* **18**, 14 (2020).

58. M. Dinur, R. Kilav, A. Sela-Brown, H. Jacquemin-Sablon, T. Naveh-Many, In vitro evidence that upstream of N-ras participates in the regulation of parathyroid hormone messenger ribonucleic acid stability, *Mol. Endocrinol.* (2006), doi:10.1210/me.2005-0333.

59. K. S. Moore, N. Yagci, F. Van Alphen, N. A. Paolini, R. Horos, N. M. Held, R. H. Houtkooper, E. Van Den Akker, A. B. Meijer, P. A. C. T'Hoen, M. Von Lindern, Csde1 binds transcripts involved in protein homeostasis and controls their expression in an erythroid cell line, *Sci. Rep.* (2018), doi:10.1038/s41598-018-20518-7.

60. T. C. Chang, A. Yamashita, C. Y. A. Chen, Y. Yamashita, W. Zhu, S. Durdan, A. Kahvejian, N. Sonenberg, A. Bin Shyu, UNR, a new partner of poly(A)-binding protein, plays a key role in translationally coupled mRNA turnover mediated by the c-fos major coding-region determinant, *Genes Dev.* (2004), doi:10.1101/gad.1219104.

61. G. M. Wilson, M. Z. Vasa, R. G. Deeley, Stabilization and cytoskeletal-association of LDL receptor mRNA are mediated by distinct domains in its 3' untranslated region, *J. Lipid Res.* (1998).

62. H. Li, W. Chen, Y. Zhou, P. Abidi, O. Sharpe, W. H. Robinson, F. B. Kraemer, J. Liu,

Identification of mRNA binding proteins that regulate the stability of LDL receptor mRNA through AU-rich elements, *J. Lipid Res.* (2009), doi:10.1194/jlr.M800375-JLR200.

63. K. Bjune, L. Wierød, S. Naderi, Triciribine increases LDLR expression and LDL uptake through stabilization of LDLR mRNA, *Sci. Rep.* (2018), doi:10.1038/s41598-018-34237-6.

64. T. Bakheet, M. Frevel, B. R. G. Williams, W. Greer, K. S. A. Khabar, ARED: Human AU-rich element-containing mRNA database reveals an unexpectedly diverse functional repertoire of encoded proteins, *Nucleic Acids Res.* (2001), doi:10.1093/nar/29.1.246.

65. R. S. Wu, I. I. Lam, H. Clay, D. N. Duong, R. C. Deo, S. R. Coughlin, A Rapid Method for Directed Gene Knockout for Screening in G0 Zebrafish., *Dev. Cell* **46**, 112-125.e4 (2018).

66. C. Liu, Y. S. Kim, J. Kim, J. Pattison, A. Kamaid, Y. I. Miller, Modeling hypercholesterolemia and vascular lipid accumulation in LDL receptor mutant zebrafish, *J. Lipid Res.* **59**, 391–399 (2018).

67. A. B. Singh, H. Li, C. F. K. Kan, B. Dong, M. R. Nicolls, J. Liu, The critical role of mRNA destabilizing protein heterogeneous nuclear ribonucleoprotein D in 3′ untranslated region-mediated decay of low-density lipoprotein receptor mRNA in liver tissue, *Arterioscler. Thromb. Vasc. Biol.* (2014), doi:10.1161/ATVBAHA.112.301131.

68. B. Paigen, A. Morrow, C. Brandon, D. Mitchell, P. Holmes, Variation in susceptibility to atherosclerosis among inbred strains of mice, *Atherosclerosis* (1985), doi:10.1016/0021-9150(85)90138-8.

69. G. S. Getz, C. A. Reardon, Diet and murine atherosclerosis*Arterioscler. Thromb. Vasc. Biol.* (2006), doi:10.1161/01.ATV.0000201071.49029.17.

70. M. von Scheidt, Y. Zhao, Z. Kurt, C. Pan, L. Zeng, X. Yang, H. Schunkert, A. J. Lusis, Applications and Limitations of Mouse Models for Understanding Human Atherosclerosis*Cell Metab.* (2017), doi:10.1016/j.cmet.2016.11.001.

71. R. Collins, C. Reith, J. Emberson, J. Armitage, C. Baigent, L. Blackwell, R. Blumenthal, J. Danesh, G. D. Smith, D. DeMets, S. Evans, M. Law, S. MacMahon, S. Martin, B. Neal, N. Poulter, D. Preiss, P. Ridker, I. Roberts, A. Rodgers, P. Sandercock, K. Schulz, P. Sever, J. Simes, L. Smeeth, N. Wald, S. Yusuf, R. Peto, Interpretation of the evidence for the efficacy and safety of statin therapy, *Lancet* **388**, 2532–2561 (2016).

72. B. T. Emmer, E. J. Sherman, P. J. Lascuna, S. E. Graham, C. J. Willer, D. Ginsburg, Genome-scale CRISPR screening for modifiers of cellular LDL uptake, *PLOS Genet.* (2021), doi:10.1371/journal.pgen.1009285.

73. M. N. Trinh, M. S. Brown, J. L. Goldstein, J. Han, G. Vale, J. G. McDonald, J. Seemann, J. T. Mendell, F. Lu, Last step in the path of LDL cholesterol from lysosome to plasma membrane to ER is governed by phosphatidylserine, *Proc. Natl. Acad. Sci. U. S. A.* (2020), doi:10.1073/pnas.2010682117.

74. C. Knouff, S. Malloy, J. Wilder, M. K. Altenburg, N. Maeda, Doubling Expression of the Low Density Lipoprotein Receptor by Truncation of the 3′-Untranslated Region Sequence Ameliorates Type III Hyperlipoproteinemia in Mice Expressing the Human ApoE2 Isoform, *J. Biol. Chem.* (2001), doi:10.1074/jbc.M009423200.

75. W. Kong, J. Wei, P. Abidi, M. Lin, S. Inaba, C. Li, Y. Wang, Z. Wang, S. Si, H. Pan, S. Wang, J. Wu, Y. Wang, Z. Li, J. Liu, J. D. Jiang, Berberine is a novel cholesterol-lowering drug working through a unique mechanism distinct from statins, *Nat. Med.* (2004), doi:10.1038/nm1135.

76. H. Ju Lee, D. Bartsch, C. Xiao, S. Guerrero, G. Ahuja, C. Schindler, J. J. Moresco, J. R. Yates, F. Gebauer, H. Bazzi, C. Dieterich, L. Kurian, D. Vilchez, A post-transcriptional program coordinated by CSDE1 prevents intrinsic neural differentiation of human embryonic stem cells, *Nat. Commun.* (2017), doi:10.1038/s41467-017-01744-5.

77. O. Boussadia, M. Niepmann, L. Créancier, A.-C. Prats, F. Dautry, H. Jacquemin-Sablon, Unr Is Required In Vivo for Efficient Initiation of Translation from the Internal Ribosome Entry Sites of both Rhinovirus and Poliovirus, *J. Virol.* (2003), doi:10.1128/jvi.77.6.3353-3359.2003.

78. V. Dormoy-Raclet, J. Markovits, A. Jacquemin-Sablon, H. Jacquemin-Sablon, Regulation of
Unr expression by 5′- and 3′-untranslated regions of its mRNA through modulation of stability
and IRES mediated translation, *RNA Biol.* (2005), doi:10.4161/rna.2.3.2203.
79. K. E. Duncan, C. Strein, M. W. Hentze, The SXL-UNR Corepressor Complex Uses a PABP-
Mediated Mechanism to Inhibit Ribosome Recruitment to msl-2 mRNA, *Mol. Cell* (2009),
doi:10.1016/j.molcel.2009.09.042.
80. S. Ray, E. C. Anderson, Stimulation of translation by human Unr requires cold shock
domains 2 and 4, and correlates with poly(A) binding protein interaction, *Sci. Rep.* (2016),
doi:10.1038/srep22461.
81. H. Guo, Y. Li, L. Shen, T. Wang, X. Jia, L. Liu, T. Xu, M. Ou, K. Hoekzema, H. Wu, M. A.
Gillentine, C. Liu, H. Ni, P. Peng, R. Zhao, Y. Zhang, C. Phornphutkul, A. P. A. Stegmann, C. E.
Prada, R. J. Hopkin, J. T. Shieh, K. McWalter, K. G. Monaghan, P. M. van Hasselt, K. van
Gassen, T. Bai, M. Long, L. Han, Y. Quan, M. Chen, Y. Zhang, K. Li, Q. Zhang, J. Tan, T. Zhu,
Y. Liu, N. Pang, J. Peng, D. A. Scott, S. R. Lalani, M. Azamian, G. M. S. Mancini, D. J. Adams,
M. Kvarnung, A. Lindstrand, A. Nordgren, J. Pevsner, I. A. Osei-Owusu, C. Romano, G.
Calabrese, O. Galesi, J. Gecz, E. Haan, J. Ranells, M. Racobaldo, M. Nordenskjold, S. Madan-
Khetarpal, J. Sebastian, S. Ball, X. Zou, J. Zhao, Z. Hu, F. Xia, P. Liu, J. A. Rosenfeld, B. B. A.
de Vries, R. A. Bernier, Z. Q. D. Xu, H. Li, W. Xie, R. B. Hufnagel, E. E. Eichler, K. Xia,
Disruptive variants of CSDE1 associate with autism and interfere with neuronal development
and synaptic transmission, *Sci. Adv.* (2019), doi:10.1126/sciadv.aax2166.
82. L. Wurth, P. Papasaikas, D. Olmeda, N. Bley, G. T. Calvo, S. Guerrero, D. Cerezo-Wallis, J.
Martinez-Useros, M. García-Fernández, S. Hüttelmaier, M. S. Soengas, F. Gebauer,
UNR/CSDE1 Drives a Post-transcriptional Program to Promote Melanoma Invasion and
Metastasis, *Cancer Cell* (2016), doi:10.1016/j.ccell.2016.10.004.
83. P. Gennemark, K. Walter, N. Clemmensen, D. Rekić, C. A. M. Nilsson, J. Knöchel, M.
Hölttä, L. Wernevik, B. Rosengren, D. Kakol-Palm, Y. Wang, R. Z. Yu, R. S. Geary, S. J. Riney,
B. P. Monia, R. Isaksson, R. Jansson-Löfmark, C. S. J. Rocha, D. Lindén, E. Hurt-Camejo, R.
Crooke, L. Tillman, T. Rydén-Bergsten, B. Carlsson, U. Andersson, M. Elebring, A. Tivesten, N.
Davies, An oral antisense oligonucleotide for PCSK9 inhibition, *Sci. Transl. Med.* **13**, eabe9117
(2021).
84. K. Musunuru, A. C. Chadwick, T. Mizoguchi, S. P. Garcia, J. E. DeNizio, C. W. Reiss, K.
Wang, S. Iyer, C. Dutta, V. Clendaniel, M. Amaonye, A. Beach, K. Berth, S. Biswas, M. C.
Braun, H.-M. Chen, T. V Colace, J. D. Ganey, S. A. Gangopadhyay, R. Garrity, L. N. Kasiewicz,
J. Lavoie, J. A. Madsen, Y. Matsumoto, A. M. Mazzola, Y. S. Nasrullah, J. Nneji, H. Ren, A.
Sanjeev, M. Shay, M. R. Stahley, S. H. Y. Fan, Y. K. Tam, N. M. Gaudelli, G. Ciaramella, L. E.
Stolz, P. Malyala, C. J. Cheng, K. G. Rajeev, E. Rohde, A. M. Bellinger, S. Kathiresan, In vivo
CRISPR base editing of PCSK9 durably lowers cholesterol in primates, *Nature* **593**, 429–434
(2021).
85. K. F. McClure, D. W. Piotrowski, D. Petersen, L. Wei, J. Xiao, A. T. Londregan, A. S.
Kamlet, A.-M. Dechert-Schmitt, B. Raymer, R. B. Ruggeri, D. Canterbury, C. Limberakis, S.
Liras, P. DaSilva-Jardine, R. G. Dullea, P. M. Loria, B. Reidich, C. T. Salatto, H. Eng, E. Kimoto,
K. Atkinson, A. King-Ahmad, D. Scott, K. Beaumont, J. R. Chabot, M. W. Bolt, K. Maresca, K.
Dahl, R. Arakawa, A. Takano, C. Halldin, Liver-Targeted Small-Molecule Inhibitors of Proprotein
Convertase Subtilisin/Kexin Type 9 Synthesis, *Angew. Chemie Int. Ed.* **56**, 16218–16222
(2017).
86. J. Martinez-Useros, N. Garcia-Carbonero, W. Li, M. J. Fernandez-Aceñero, I. Cristobal, R.
Rincon, M. Rodriguez-Remirez, A. Borrero-Palacios, J. Garcia-Foncillas, UNR/CSDE1
Expression Is Critical to Maintain Invasive Phenotype of Colorectal Cancer through Regulation
of c-MYC and Epithelial-to-Mesenchymal Transition, *J. Clin. Med.* (2019),
doi:10.3390/jcm8040560.
87. V. Dormoy-Raclet, J. Markovits, Y. Malato, S. Huet, P. Lagarde, D. Montaudon, A.

Jacquemin-Sablon, H. Jacquemin-Sablon, Unr, a cytoplasmic RNA-binding protein with cold-shock domains, is involved in control of apoptosis in ES and HuH7 cells, *Oncogene* (2007), doi:10.1038/sj.onc.1210068.

88. N. Matsuzawa, T. Takamura, S. Kurita, H. Misu, T. Ota, H. Ando, M. Yokoyama, M. Honda, Y. Zen, Y. Nakanuma, K. I. Miyamoto, S. Kaneko, Lipid-induced oxidative stress causes steatohepatitis in mice fed an atherogenic diet, *Hepatology* (2007), doi:10.1002/hep.21874.

89. J. J. V McMurray, M. Packer, A. S. Desai, J. Gong, M. P. Lefkowitz, A. R. Rizkala, J. L. Rouleau, V. C. Shi, S. D. Solomon, K. Swedberg, M. R. Zile, Angiotensin–Neprilysin Inhibition versus Enalapril in Heart Failure, *N. Engl. J. Med.* **371**, 993–1004 (2014).

90. D. G. Gibson, L. Young, R.-Y. Chuang, J. C. Venter, C. A. Hutchison, H. O. Smith, Enzymatic assembly of DNA molecules up to several hundred kilobases., *Nat. Methods* **6**, 343–5 (2009).

91. J. S. Chorba, A. M. Galvan, K. M. Shokat, Stepwise processing analyses of the single-turnover PCSK9 protease reveal its substrate sequence specificity and link clinical genotype to lipid phenotype, *J. Biol. Chem.* **293**, 1875–1886 (2018).

92. J. H. Kim, S.-R. Lee, L.-H. Li, H.-J. Park, J.-H. Park, K. Y. Lee, M.-K. Kim, B. A. Shin, S.-Y. Choi, V. Thiel, Ed. High Cleavage Efficiency of a 2A Peptide Derived from Porcine Teschovirus-1 in Human Cell Lines, Zebrafish and Mice, *PLoS One* **6**, e18556 (2011).

93. D. G. Gibson, H. O. Smith, C. A. Hutchison, J. C. Venter, C. Merryman, Chemical synthesis of the mouse mitochondrial genome, *Nat. Methods* (2010), doi:10.1038/nmeth.1515.

94. H. Mi, A. Muruganujan, X. Huang, D. Ebert, C. Mills, X. Guo, P. D. Thomas, Protocol Update for large-scale genome and gene function analysis with the PANTHER classification system (v.14.0), *Nat. Protoc.* (2019), doi:10.1038/s41596-019-0128-8.

95. A. Bateman, UniProt: A worldwide hub of protein knowledge, *Nucleic Acids Res.* (2019), doi:10.1093/nar/gky1049.

96. C. Sudlow, J. Gallacher, N. Allen, V. Beral, P. Burton, J. Danesh, P. Downey, P. Elliott, J. Green, M. Landray, B. Liu, P. Matthews, G. Ong, J. Pell, A. Silman, A. Young, T. Sprosen, T. Peakman, R. Collins, UK Biobank: An Open Access Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age, *PLoS Med.* (2015), doi:10.1371/journal.pmed.1001779.

97. P. R. Loh, G. Tucker, B. K. Bulik-Sullivan, B. J. Vilhjálmsson, H. K. Finucane, R. M. Salem, D. I. Chasman, P. M. Ridker, B. M. Neale, B. Berger, N. Patterson, A. L. Price, Efficient Bayesian mixed-model analysis increases association power in large cohorts, *Nat. Genet.* (2015), doi:10.1038/ng.3190.

98. J. M. Baker, F. M. Boyce, High-throughput functional screening using a homemade dual-glow luciferase assay, *J. Vis. Exp.* (2014), doi:10.3791/50282.

99. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2, *Genome Biol.* (2014), doi:10.1186/s13059-014-0550-8.

100. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for differential expression analysis of digital gene expression data., *Bioinformatics* **26**, 139–140 (2010).

101. A. T. L. Lun, Y. Chen, G. K. Smyth, It's DE-licious: A Recipe for Differential Expression Analyses of RNA-seq Experiments Using Quasi-Likelihood Methods in edgeR., *Methods Mol. Biol.* **1418**, 391–416 (2016).

102. M. V. Kuleshov, M. R. Jones, A. D. Rouillard, N. F. Fernandez, Q. Duan, Z. Wang, S. Koplev, S. L. Jenkins, K. M. Jagodnik, A. Lachmann, M. G. McDermott, C. D. Monteiro, G. W. Gundersen, A. Ma'ayan, Enrichr: a comprehensive gene set enrichment analysis web server 2016 update, *Nucleic Acids Res.* (2016), doi:10.1093/nar/gkw377.

103. Z. Gu, R. Eils, M. Schlesner, Complex heatmaps reveal patterns and correlations in multidimensional genomic data., *Bioinformatics* **32**, 2847–2849 (2016).

104. G. S. (editors). Higgins JPT, Cochrane Handbook for Systematic Reviews of Interventions

Version 5.1.0 *Cochrane Collab.* (2011).

105. M. Roederer, W. Moore, A. Treister, R. R. Hardy, L. A. Herzenberg, Probability binning

comparison: A metric for quantitating multivariate distribution differences, *Cytometry* (2001),

doi:10.1002/1097-0320(20010901)45:1<47::AID-CYTO1143>3.0.CO;2-A.

106. M. Roederer, A. Treister, W. Moore, L. A. Herzenberg, Probability binning comparison: A

metric for quantitating univariate distribution differences, *Cytometry* (2001), doi:10.1002/1097-

0320(20010901)45:1<37::AID-CYTO1142>3.0.CO;2-E.

107. M. Roederer, R. R. Hardy, Frequency difference gating: A multivariate method for

identifying subsets that differ between samples, *Cytometry* (2001), doi:10.1002/1097-

0320(20010901)45:1<56::AID-CYTO1144>3.0.CO;2-9.

1387

1388

1389 **Figures and Figure Legends**
1390

1391
1392
Figure 1: Genome-Wide CRISPR Interference Screen. *A)* Overall schematic of selection. See
text for details. *B)* Volcano plot showing the statistical significance (Mann-Whitney test) of the
guides recovered for each gene against the mean ρ phenotype of the three guides with the
strongest effect. ρ is defined as the $\log_2$-fold enrichment for high LDLR expressing cells to the
low LDLR expressing cells. Guides targeting known regulators of the LDLR are noted. *C)* Venn
diagram showing the overlap between parallel LDLR and TFR screens. 6 guides common to
both had opposing expression phenotypes in the respective screens and were included as
specific hits. *D)* Venn diagram of hits between the LDLR screen (GWCS) and putative genes
correlated with serum LDL from GWAS. The dotted line indicates a relaxed threshold for hit
selection from LDLR screen, with only an additional 3 genes in the overlap. Overlap genes
shown at right.
1404

1405
1406
Figure 2: Validation of LDLR CRISPRi Hits. Heatmap showing receptor abundance (LDLR,
TFR, and LDLR/TFR ratio) and function (LDL uptake) for dCas9-KRAB HepG2 cells transduced
with sgRNA targeting the indicated gene, analyzed by flow cytometry. Hits are grouped
according to directional effect on LDLR abundance, and then within groups, by effect on LDL
uptake (with uptake from *FOXL3-OT1*, *CIT*, and *DHX15* sgRNAs not significantly different from
negative control sgRNA). *CSDE1* highlighted in blue. Control sgRNAs shown at right. Readouts
show $\log_2$ fold change compared to transduction with negative control sgRNA and represent the
weighted average of the effects from both sgRNAs targeting each gene. Viability indicates the
relative number of cells surviving to flow cytometry in the experiments. Functional classification
of genes is shown visually in Supp. Fig. 3. Note that LDLR/TFR is a separately ascertained
value from individual cells, and not a derived parameter from aggregate data. Only the hits for
which two separate sgRNAs independently validated for receptor expression are shown, defined
as $p < 0.05$ via Holm-Sidak corrected T-test. Data represent summary information from 3 to 4
independent experiments.
1421

Figure 3: CRISPRi Knockdown Synergy with Statin and PF-846. Heatmap showing synergy score with statin (top) or PF-846 (bottom) for knockdowns of indicated validated genes with a single sgRNA for separate LDLR abundance and function experiments. Hits are grouped first according to overall effect on LDLR abundance, and secondarily by effect on LDL uptake, as in Fig. 2. *CSDE1* highlighted in blue. Data represent summary information from 4 independent experiments.

Figure 4: CSDE1 Mediates *LDLR* mRNA Decay. *A)* Relative LDLR abundance, by normalized mean fluorescence, in engineered dCas9-KRAB HepG2 cells transduced with indicated sgRNAs (CRISPRi cells) and grown in the indicated media conditions. *B)* Relative LDLR abundance, by normalized median fluorescence, in HepG2 cells overexpressing indicated CSDE1 isoforms. One-way ANOVA with Tukey's multiple comparisons test shown. *C)* Relative expression, by qPCR, of indicated mRNA targets in CRISPRi cells under sterol-depleted conditions. *D)* Relative expression, by qPCR, of *LDLR* mRNA in CRISPRi cells after arrest of transcription with actinomycin D. Data are normalized to results at T=0 within the sgRNA evaluated to illustrate the change in time. $t_{1/2}$ shown indicates the best fit data to a one-stage exponential decay equation. Unpaired t-tests with Holm-Sidak correction shown for pairwise comparisons. Extra sum-of-squares F test shown for decay equation. *E)* Schematics of Luc2-Prom$_{LDLR}$ reporter constructs, illustrating *LDLR* promoter, start site (arrowhead), P2A ribosomal skipping sequence (if present), AREs in 3' UTR, stop codon (red octagon), and indicated regions of the *LDLR* gene. ARE = Adenylate-uridylate (AU) rich element. Schematics not to scale. *F)* Ratiometric luciferase outputs, normalized to negative control, of CRISPRi cells transfected with indicated luciferase constructs. *G)* Ratiometric luciferase outputs of 3' UTR addended luciferase constructs in CRISPRi cells. *All panels)* Error bars indicate 95% confidence intervals. Data represent summary information from 3-4 independent experiments. n.s. = not significant, * = p < 0.05, ** = p < 0.01, *** = p < 0.001, and **** = p < 0.0001. Unless otherwise indicated, two-way ANOVA with Sidak's multiple comparisons test was used for statistical analysis.

Figure 5: CSDE1 Disruption Upregulates the LDLR *in vivo*. A) Total cholesterol, in µg cholesterol per mg of total protein, of homogenates of 8 days post fertilization (dpf) zebrafish larvae fed a high-cholesterol diet and subjected to Cas9 mediated gene disruption of indicated target. Data are normalized to the scramble control of a particular experiment. Each point represents a homogenate consisting of 10 larvae. Data represent summary information from 4 independent experiments. One-way ANOVA with Holm-Sidak's multiple comparisons test shown. **B)** Total fasting plasma cholesterol of C57BL/6 mice on an atherogenic diet, 2 weeks after transduction with AAV8-packaged shRNA against indicated target. Each point represents an individual mouse. One-way ANOVA with Tukey's multiple comparisons test shown. Error bars = 95% confidence intervals. **C)** Cholesterol levels of fractions collected from gel filtration of plasma from individual mice, harvested 2 weeks after transduction with second dose of AAV8-packaged shRNA. Note that fractions shown begin with the elution front from the size-exclusion column. Each dot represents the mean cholesterol level from mice in the same intervention arm. Immunoblots from representative fractions against mouse ApoB-100 and ApoA-I shown below. Two-way ANOVA with Sidak's multiple comparisons test shown to illustrate comparison between treatment arm within a given fraction. Error bars = standard error of the mean. **D)** Volcano plot showing differentially expressed genes between *Csde1* and scramble shRNA treatment arms, filtered for effects of viral transduction. Statistical significance is shown on the y

axis and strength of effect is shown on the x axis. Comparison made among the 3 mice in each
arm with the highest *eGFP* transcript expression, as a proxy for transduction efficiency. Genes
reaching threshold significance for p-value (blue), $\log_2$ fold change (green), both (red), or neither
(grey) annotated accordingly. Genes associated with lipid and sterol metabolic GO terms
highlighted in yellow (0019216 = lipid metabolic process, 000695 = cholesterol biosynthetic
process, and 0045540 = regulation of cholesterol biosynthetic process). Leading upregulated
and downregulated GO terms of all statistically significant differentially expressed genes noted
in the boxes at right. *All panels)* n.s. = non-significant, * = $p < 0.05$, ** = $p < 0.01$, *** = $p <$
0.001, and **** = $p < 0.0001$.
1483

Figure 6: An Exploratory Map of Potential LDLR Regulatory Targets. Genes identified and validated in the screen are mapped by cellular localization and possible mechanisms of effect. Known downregulators are shown in cyan (including CSDE1 given the results presented in the current study) and known upregulators shown in magenta. Validated hits with observed effects on cell proliferation or viability are excluded.

Tables and Table Legends

Hit	Hit (NCBI Name)
ABCA1	ABCA1
ABCA4	ABCA4
ACAN	ACAN
ALKBH5	ALKBH5
AMMECR1L	AMMECR1L
ANO8	ANO8
BCAR1	BCAR1
C12orf45	C12orf45
C14orf79	CLBA1
C1orf210	C1orf210
C5orf34	C5orf34
C6orf132	C6orf132
C9orf40	C9orf40
C9orf92	C9orf92
CD164L2	CD164L2
CD276	CD276
CD96	CD96
CIT	CIT
CSDE1	CSDE1
CXCR2	CXCR2
CXXC11	FBXL19
CYB5R3	CYB5R3
DDX39B	DDX39B
DESI1	DESI1
DHX15	DHX15
DUOX1	DUOX1
EIF3D	EIF3D
ENG	ENG
ENTPD1	ENTPD1
ESRRG	ESRRG
EVA1B	EVA1B
FAM126A	FAM126A
FAM178B	FAM178B
FAM57A	TLCD3A
FBXW11	FBXW11
FDPS	FDPS

GXYLT1	GXYLT1
HMGCR	HMGCR
HMGCS1	HMGCS1
HNF1A	HNF1A
HNF4A	HNF4A
HPGDS	HPGDS
HRK	HRK
ICAM4	ICAM4
INTS8	INTS8
ITGA11	ITGA11
ITGA7	ITGA7
ITGAV	ITGAV
LDLR	LDLR
LGALS14	LGALS14
LOC100288524	FOXL3-OT1
LOC729159	NPAP1L
LYZ	LYZ
MARK2	MARK2
MATN1	MATN1
MFHAS1	MFHAS1
MRAP2	MRAP2
MRPL16	MRPL16
MRPL22	MRPL22
MSMO1	MSMO1
MYLIP	MYLIP
NCR2	NCR2
NDUFB5	NDUFB5
NDUFS8	NDUFS8
NINJ1	NINJ1
NLRP6	NLRP6
ONECUT1	ONECUT1
OR52A1	OR52A1
PCDH7	PCDH7
PCDHB4	PCDHB4
PCSK9	PCSK9
PHGR1	PHGR1
PIANP	PIANP
PID1	PID1
PLAC1L	OOSP2

PMVK	PMVK
POLD2	POLD2
POLD3	POLD3
PRIM1	PRIM1
PROL1	OPRPN
PTGDR2	PTGDR2
RARRES3	PLAAT4
REPS1	REPS1
RNF151	RNF151
RSG1	CPLANE2
SCUBE1	SCUBE1
SEC61G	SEC61G
SERPINA9	SERPINA9
SF3A2	SF3A2
SLC25A27	SLC25A27
SLC2A7	SLC2A7
SLC6A19	SLC6A19
SLURP1	SLURP1
SMURF1	SMURF1
SON	SON
SREBF2	SREBF2
SSR2	SSR2
SSUH2	SSUH2
ST6GALNAC4	ST6GALNAC4
STAC	STAC
STAG2	STAG2
TIMELESS	TIMELESS
TMEM217	TMEM217
TMEM86A	TMEM86A
TPRG1	TPRG1
TRMT10C	TRMT10C
TRPM1	TRPM1
TRPM7	TRPM7
TTC14	TTC14
TXNDC8	TXNDC8
WDR5	WDR5
WDR75	WDR75
ZBED6CL	ZBED6CL
ZBTB42	ZBTB42

ZC3H12A	ZC3H12A
ZEB1	ZEB1
ZNF595	ZNF595

1495
1496 **Table 1: LDLR Specific CRISPRi Screen Hits.** Hits are listed both by gene name in the
1497 genome-wide library*(14)* as well as NCBI name.
1498

GENE	Variant rsID	BETA	P_BOLT_LMM	Consequence	IMPACT
HNF4A	rs1800961	0.0564144	0	missense_variant	MODERATE
BCAM	rs28399659	-0.0174111	7.70E-29	missense_variant	MODERATE
BCAM	rs200398713	-0.0803165	1.80E-28	splice_region_variant,intron_variant	LOW
BCAM	rs199922856	-0.342179	6.20E-28	missense_variant	MODERATE
BCAM	rs28399654	0.220592	6.10E-10	missense_variant	MODERATE
BCAM	rs3810141	0.020077	5.50E-07	stop_gained	HIGH
TIMELESS	rs2291738	0.00388393	0.00014	splice_region_variant,intron_variant	LOW
BCAM	rs149302547	-0.147327	0.005	missense_variant	MODERATE
BCAM	rs1135062	-0.0213642	0.0074	missense_variant	MODERATE
C6orf132	rs55772414	0.0116856	0.013	missense_variant	MODERATE
MSMO1	rs142496142	0.0432195	0.015	missense_variant	MODERATE

1499

1500 **Table 2: Association of Nonsynonymous Variants in CRISPRi Screen Hits with Serum**
1501 **LDL-C in the UK Biobank.** BETA indicates the linear regression standardized effect size, and
1502 P_BOLT_LMM indicates the linear mixed model p value using BOLT-LMM*(97)*.

1503 **Supplementary Figures and Figure Legends**
1504

1505
1506
Supplementary Figure 1: Validation of dCas9-KRAB-HepG2 Cells. *A)* Relative expression,
by qPCR, of *LDLR* and *HMGCR* in engineered dCas9-KRAB HepG2 cells transduced with
sgRNAs targeting the indicated genes. *B2M* used as qPCR control. Error bars indicate 95%
confidence intervals. *** = p < 0.001 by Holm-Sidak corrected T-test, comparing to negative
control sgRNA of the same target. *B)* Flow cytometry analysis, by surface labelling with anti-
LDLR-AF647, of dCas9-KRAB HepG2 cells transduced with sgRNAs targeting the indicated
genes. Mean fluorescence shown in inset. *MYLIP* (IDOL) is an E3 ligase which ubiquitinates the
LDLR, leading to lysosomal degradation*(26)*. *C)* Flow cytometry analysis as in *B* but transduced
with indicated sgRNAs and labelled with anti-TFR-AF647. *ZC3H12A* (REG1) is an
endoribonuclease that accelerates the degradation of TFR mRNA*(27)*.
1517

1518
1519
Supplementary Figure 2: Recovered sgRNAs from Screening Phenotypes. Distribution of
number of guide RNAs recovered by phenotype in both LDLR (*A-C*) and TFR (*D-F*) screens. ρ
(*A,D*) indicates $\log_2$ fold enrichment for sgRNA in high receptor abundance cells compared to
low receptor abundance cells. γ (*B,E*) indicates $\log_2$ enrichment for sgRNA in low receptor
abundance cells compared to unsorted population. τ (*C,F*) indicates $\log_2$ enrichment for sgRNA
in high receptor abundance cells compared to unsorted population. Mean results are reported
for the 3 replicates of the LDLR screen (*A-C*).
1527

1528
1529
1530 **Supplementary Figure 3: Gene Ontology and Localization Analysis.** *A)* Characterization of
1531 hits from the LDLR screen based on gene ontology (GO) and localization, along with results
1532 from GO enrichment analysis (yellow center). Note that multiple genes fall into more than one
1533 category. *B)* Primary classification of the 40 LDLR hits independently validated outside of the
1534 pooled screen and displayed, as in Fig. 2, according to the color codes in *A*.
1535

1536

1537

1538 **Supplementary Figure 4: Selective LDLR Effect of Transmembrane Proteins.** Flow
1539 cytometric readout of receptor abundance and LDLR function assays, using CRISPRi
1540 knockdowns against genes thought to be involved in endocytosis. Data, which represents 3 to 4
1541 independent experiments, are normalized to readout of negative control sgRNA within each
1542 experiment. Error bars represent 95% confidence intervals. Note the discontinuous Y axis.

1543

1544

1545
1546
Supplementary Figure 5: Effect of Sterol Conditions on CSDE1 Knockdown. Flow
cytometry histograms showing AF647 conjugated anti-LDLR antibody labelling (as proxy for
LDLR abundance) of engineered dCas9-KRAB HepG2 cells transduced with indicated sgRNAs
and grown in standard growth media (*A*), lipoprotein-deficient media (*B*), or lipoprotein deficient
media with a concomitant statin (*C*). $T\chi$ metric (FlowJo v10)*(105–107)* shown on graph, and
mean fluorescence shown in insets.
1553

Supplementary Figure 6: Effect of CSDE1 Overexpression. Flow cytometry histograms showing AF647 conjugated anti-LDLR antibody labelling (as proxy for LDLR abundance) of HepG2 cells (*A-D*) or engineered dCas9-KRAB HepG2 cells transduced with *CSDE1* targeting sgRNA (*E-H*) and transfected with an overexpression construct encoding the indicated CSDE1 isoform or vector alone. Tχ metric (FlowJo v10)*(105–107)* shown on graphs, and median fluorescence shown above. Quantified relative LDLR abundance of normalized median fluorescence of the data from *e-h* shown in *i*, with one-way ANOVA with Tukey's multiple comparisons test. Error bars indicate 95% confidence intervals. **** = p < 0.0001.

1565
1566
1567 **Supplementary Figure 7: CSDE1 Knockdown at Protein Level.** *A*) Representative
1568 immunoblots of lysates of dCas9-KRAB HepG2 cells harboring indicated guide RNA. CSDE1
1569 shown above, and β-actin (loading control) shown below. *B*) Quantification of relative
1570 abundance of CSDE1 (normalized to loading control) shown in immunoblot in *A*. Data includes 3
1571 independent experiments. * indicates p < 0.05 by Welch's T-test.
1572

1573
1574
Supplementary Figure 8: Effect of *CSDE1* Knockdown on Decay of Non-*LDLR*
Transcripts. Relative expression, by qPCR, of *HMGCR (A), TFRC (B), SREBF2 (C), or PCSK9*
(D) mRNA in dCas9-KRAB HepG2 cells transduced with indicated sgRNAs and subjected to
arrest of transcription with actinomycin D. Data are normalized to results at T=0 within the
sgRNA evaluated to illustrate the change in time. Data represent summary results from 3
independent experiments. Error bars = 95% confidence intervals. All pairwise comparisons
(unpaired t-tests with Holm-Sidak correction) are nonsignificant.
1582

Reporter	pLuc2-Prom$_{LDLR}$ (Luc2 Only)		
Lipoprotein	+	-	-
Statin	-	-	+

Supplementary Figure 9: Physiologic Response of Luciferase Reporter System. Relative ratiometric luciferase activity of dCas9-KRAB HepG2 (CRISPRi) cells transiently transfected with unmodified Luc2 construct under the *LDLR* promoter (pLuc2-Prom$_{LDLR}$, Fig. 4E) and secreted Nluc reporter under the CMV promoter and subjected to indicated media conditions. Data represent summary results from 3 independent experiments. n.s. = non-significant. Two-way ANOVA with Sidak's multiple comparisons test was used.

1592
1593
Supplementary Figure 10: Visual Phenotypes of Zebrafish Cas9-sgRNA Saturation Gene
Disruption. *A,B*) Representative microscopic images of zebrafish larvae at 1 day post
fertilization without (*A*) or with (*B*) injected Cas9 and redundant guides against tyrosinase
control performed concomitantly with each zebrafish experiment. Albinism is the readout for
successful injections. *C-E)* Representative microscopic images of zebrafish larvae at 8 dpf and
with injected Cas9 and guides against scramble controls (*C*), *ldlra* (*D*), and *csde1* (*E*).
1600

Supplementary Figure 11: Effects of *in vivo Csde1* Disruption in Mice. *A)* Total fasting plasma cholesterol of C57BL/6 mice on an atherogenic diet, 2 weeks after transduction with second dose of AAV8-packaged shRNA against indicated target (8 weeks after first dose). Welch's t-test shown. *B)* Relative expression, by qPCR, of indicated mRNA targets from mouse liver tissue from the indicated treatment arms. Expression normalized to *B2m* as the housekeeping control. Welch's T-test shown. Error bars = standard error of the mean. *C)* Plasma alanine aminotransferase activity (ALT) of mice from *A*. Welch's T-test revealed no significant difference between intervention arms. *D)* Unsupervised cluster analysis of individual mice analyzed for differential gene analysis. *E)* Heatmap of top 100 differentially expressed genes (by p value) of the individual mice analyzed by RNA-seq. Mice transduced with AAV8-scramble-shRNA at the bottom and those transduced with AAV8-*Csde1*-shRNA at the top. *F)* Top 10 biological process GO terms (by uncorrected p value) identified from all statistically

1615   significant differentially expressed genes downregulated (blue) or upregulated (red) by *Csde1*
1616   shRNA treatment. *All)* Each data point represents an individual mouse. * = p < 0.05, ** = p <
1617   0.01, and **** = p < 0.001. Error bars = 95% confidence intervals unless noted otherwise.
1618

Metric	Value
Age (y)	56.9 (7.9)
Sex	179,963 (46.1%)
European ancestry	376,358 (96.4%)
Cholesterol (mg/dl)	
Total	221.1 (44.3)
HDL	56.1 (14.8)
LDL	138.1 (33.7)
Triglycerides (mg/dl)	132.6 [93.6-191.5]
Statin Rx	64,004 (16.4%)
BMI (kg/m^2)	27.4 (4.8)
Systolic blood pressure (mmHg)	140.2 (19.7)
Diastolic blood pressure (mmHg)	82.3 (10.7)
Current smoker	39,736 (10.2%)
Diabetes mellitus type 2	25,349 (6.5%)
Coronary artery disease	18,204 (4.8%)

1619

1620 **Supplementary Table 7: Baseline Characteristics of UK Biobank Participants in Genomic**
1621 **Association Analyses.** Continuous values are presented as mean (standard deviation) except
1622 for triglycerides which is given as median (Q1-Q3) due to the skewness of the triglyceride
1623 distribution. Categorical data are presented as count (percentage). BMI = body-mass index;
1624 HDL = high-density lipoprotein; LDL = low-density lipoprotein.
1625