EEG signals index a global signature of arousal embedded in neuronal population recordings

Richard Johnston[1, 2, 3], Adam C. Snyder[4, 5, 6], Rachel S. Schibler[7], and Matthew A. Smith[1, 2, 3]

[1]Department of Biomedical Engineering, Carnegie Mellon University, Pittsburgh, USA; [2]Neuroscience Institute, Carnegie Mellon University, Pittsburgh, USA; [3]Center for the Neural Basis of Cognition, Carnegie Mellon University, Pittsburgh, USA; [4]Department of Brain and Cognitive Sciences, University of Rochester, Rochester, USA; [5]Department of Neuroscience, University of Rochester, Rochester, USA; [6]Center for Visual Science, University of Rochester, Rochester, USA; [7] Harvey Mudd College, California, USA.

Corresponding author:
Matthew A. Smith, PhD
Carnegie Mellon University
Department of Biomedical Engineering and Neuroscience Institute
Mellon Institute, Room 115
Pittsburgh
Pa, 15213
Email: mattsmith@cmu.edu

Acknowledgements:

M.A.S. was supported by NIH Grants R01 EY022928, R01 MH118929, R01 EB026953, and P30 EY008098; NSF Grant NCS 1734901; a career development grant and an unrestricted award from Research to Prevent Blindness; and the Eye and Ear Foundation of Pittsburgh. A.C.S. was supported by NIH grant K99 EY025768. R.S.S. was supported by NIH Grant R90 DA023426. The authors would like to thank Ms. Samantha Schmitt for assistance with surgery and data collection.

Abstract

Electroencephalography (EEG) has long been used to index brain states, from early studies describing activity during visual stimulation to modern work employing complex perceptual tasks. These studies shed light on brain-wide signals but lacked explanatory power at the single neuron level. Similarly, single neuron studies can suffer from inability to measure brain-wide signals. Here, we combined these techniques while monkeys performed a change detection task and discovered a link between EEG and a signal embedded in spiking responses. This 'slow drift' was associated with arousal: decreases in pre-stimulus α power/increases in P1 amplitude were accompanied by :1) increases in pupil size, false alarm rate and saccade velocity; and 2) decreases in microsaccade rate and reaction time. These results show that brain-wide EEG signals can be used to index modes of activity acquired from direct neural recordings, that in turn reflect global changes in brain state that influence perception and behavior.

Introduction

For decades, researchers have investigated how the spiking responses of single cortical neurons relate to performance on decision-making [1], attention [2] and working memory [3] tasks. Interactions between pairs of neurons have also been studied extensively since technological advances in neural recording systems (e.g. microelectrode arrays and two-photon imaging) made it possible to monitor the activity of neural populations simultaneously [4–6]. At the same time, it is becoming increasingly apparent that major insight about the neurobiological basis of cognition can be gained if one goes beyond studying single and pairwise statistics [7–14]. Furthermore, it has been suggested that low-dimensional neural activity patterns can be used to index global brain states, which influence performance on cognitive tasks. For example, Stringer et al. [15] applied principal component analysis (PCA) to data recorded from more than 10,000 neurons in the mouse and found that fluctuations in the first principal component were associated with a host of arousal-related variables including whisking, pupil size, and running speed. Musall et al. [16] found that uninstructed

movements, which themselves may occur at varying frequency based on arousal, were related to brain-wide activity in the mouse. In our own work in macaque monkeys, we have reported a brain wide 'slow drift' of neural activity [17] which is correlated with a distinctive pattern of eye metrics that is strongly indicative of changes in arousal [18]. However, it is unknown if they are associated with other arousal-related variables that can be measured in a rapid, accurate and non-invasive manner.

Spontaneous (i.e., pre-stimulus) oscillations in the α frequency band (~8 – 12Hz) are associated with global changes in brain state. For example, research in humans using EEG and magnetoencephalography (MEG) has shown that the likelihood of detecting a near-threshold visual stimulus increases when pre-stimulus oscillations in the α band decrease [19–26]. The principles of signal detection theory [27] dictate that changes in behavior on detection tasks can arise due to changes in sensitivity or response criterion [28–30]. To that end, recent work has shown that a decrease in pre-stimulus α power is associated with increased hit rate and false alarm rate [31,32]. These results point to link between pre-stimulus α power and response criterion, a key component of signal detection theory, which is modulated, at least in part, by subcortical structures that control arousal levels [31,33]. In addition, pre-stimulus α power is known to be correlated with eye metrics including pupil size [34,35] and reaction time [35–38] such that decreased pre-stimulus α power is accompanied by increased pupil size and decreased reaction time. These findings raise the possibility that other EEG signals such as the P1 component of the visually evoked potential (VEP) might also be associated with arousal.

Several studies have established a link between early components of the VEP and changes in global brain state. In contrast to pre-stimulus α oscillations, P1 amplitude is significantly larger on trials in which a weak visual stimulus is detected in comparison to when it is not detected [19,25,39,40]. Furthermore, studies that have explored the effects of spatial attention on early components of the VEP have shown that P1 amplitude is negatively correlated with reaction time [41]. In light of these results, and the well-known involvement of oscillatory brain activity in

stimulus-evoked responses [42], one might expect a negative relationship between pre-stimulus α

oscillations and early VEP components. There is some evidence to suggest that this is the case. For

example, Iemi et al. [43] found that the amplitude of the C1 and N150 increased when pre-stimulus

power in the α frequency band decreased. These results suggest that pre-stimulus α power and early

components of the VEP (e.g. P1 amplitude) can be used to index global changes in arousal. However,

it is currently unclear how to link these EEG signals to underlying patterns of neural activity in the

cortex.

In this study, we explored if pre-stimulus oscillations in the α frequency band and P1

amplitude are associated with slow drift in visual cortex. We simultaneously recorded EEG from the

scalp and spiking activity from populations of neurons in V4 of two monkeys while they performed

an orientation-change detection task (Figure 1A). Results showed that neural slow drift was

associated with a pattern that is indicative of changes in the subjects' arousal levels over time. For

example, decreases in pre-stimulus power specifically in the α band (i.e., not other frequencies) and

increases in P1 amplitude were accompanied by 1) increases in pupil size, false alarm rate and saccade

velocity; and 2) decreases in microsaccade rate and reaction time. These findings are important

because they reveal that spontaneous/evoked components of the EEG signal recorded non-invasively

on the scalp index low-dimensional patterns of neural activity acquired from microelectrode array

recordings in the brain. They support previous research [17,18] showing that slow drift is associated

with gradual changes in arousal over time, and provide a strong link between global measurements

made across recording modalities and species.

Results

To determine if spontaneous and evoked components of the EEG signal can provide insight

into the internal brain state associated with slow drift, we trained two macaque monkeys to perform

an orientation change-detection task in which pairs of stimuli were repeatedly presented (Figure 1A).

Spiking responses of populations of neurons in visual cortex (V4) were recorded using 100-channel

"Utah" arrays as well as EEG on the scalp (Figure 1B). On each trial, we calculated: 1) the mean

amplitude of oscillations in different frequency bands during each pre-stimulus period (300-500ms)

using a fast Fourier transform (FFT); and 2) the VEP during each 400ms stimulus period (see

Methods). Previous research has shown that pre-stimulus α power and stimulus-evoked P1 amplitude

across a wide range of frontal, midline and occipital electrode sites are associated with improved

performance on detection tasks [19,22,31]. Hence, the signals from all 8 EEG electrodes (referenced

online to the head post, see Methods) were averaged prior to computing the FFTs and VEPs. We

focused primarily on pre-stimulus power in the α frequency band and P1 amplitude as they capture

spontaneous (i.e., pre-stimulus) and evoked aspects of the EEG signal and have been linked to global

changes in brain state. However, we also measured pre-stimulus beta (β) and gamma (γ) oscillations

to exclude the possibility that slow drift was associated with aperiodic (i.e. broadband) changes in

spectral power over time (1/f noise) [43–46].

Figure 1. Experimental methods. (A) Orientation-change detection task. After an initial fixation period, a sequence of stimuli (orientated Gabor pairs separated by fixation periods) was presented. The subjects' task was to detect an orientation change in one of the stimuli and make a saccade to the changed stimulus. (B) Electrophysiological recordings. We simultaneously recorded: 1) spiking responses of populations of neurons in V4 using 100-channel microelectrode ('Utah') arrays; and 2) EEG from 8 electrodes positioned on the scalp.

Correlation between pre-stimulus α power and P1 amplitude

First, we explored the relationship between pre-stimulus α power and the amplitude of the P1 component of the VEP. As described above, these two signals appear to operate in an antagonistic

manner on detection tasks with decreases in pre-stimulus α power/increases in P1 amplitude being

associated with improved performance [19–26,39,40]. Hence, we hypothesized that these two large-

scale EEG signals would be negatively correlated over time. To investigate the relationship between

pre-stimulus α power and P1 amplitude, EEG data for each session was binned using a 30-minute

sliding window stepped every 6 minutes (Figure 2A and Figure 2B). The width of the window, and

the step size, were chosen to isolate slow changes over time based on previous studies we performed

[17,18]. An example session is shown in Figure 2C. In support of our hypothesis, pre-stimulus power

in the α frequency band was negatively associated with P1 amplitude.

Figure 2. Isolating slow fluctuations in spontaneous and evoked EEG signals. (A) Spontaneous EEG signals. On
each trial, we calculated the mean amplitude of oscillations in distinct frequency bands for each pre-stimulus period
using a FFT. The data was then binned using a 30-minute sliding window stepped every 6 minutes. Each bin is
represented by a shaded grey line. Note that the duration of the fixation period was held constant at 300ms in this
plot for illustration purposes only. In our analysis, we used the full duration of the fixation period, which was
randomized between 300 to 500ms. (B) Evoked EEG signals. On each trial, we calculated a VEP for each stimulus
presentation. The data was then binned using the same 30-minute sliding window so that direct comparisons could
be made between spontaneous and evoked EEG signals. Each bin is represented by a shaded grey line. (C) Example
session from Monkey 1 showing that pre-stimulus α power was negatively correlated with P1 amplitude over time.
Each metric has been z-scored for illustration purposes. (D) Histogram showing distributions of Pearson's r values
(black) across sessions between pre-stimulus α power and P1 amplitude. Actual distributions were compared to
shuffled distributions (grey) using two-sided permutation tests (difference of medians). Median r values are
indicated by dashed lines. $p < 0.05*$, $p < 0.01**$, $p < 0.001***$. See Figure S1 for additional example sessions from
both monkeys.

Next, we explored if a similar pattern was present across sessions. We computed correlations

(Pearson product-moment correlation coefficient) between pre-stimulus power in the α band and the

amplitude of the P1 component for each session and compared the distribution of Pearson's r values

obtained to shuffled distributions using permutation tests (two-sided, difference of medians).

Consistent with the pattern of results observed in several individual sessions for Monkey 1 and

Monkey 2 (Figure S1), we found that pre-stimulus α power was significantly and negatively

correlated with P1 amplitude (Figure 2D, median r = -0.36, p = 0.024). These results motivated us to

ask if pre-stimulus α power and P1 amplitude are correlated with other task performance and eye

metrics that are often taken to be hallmarks of arousal e.g. hit rate, false alarm rate, pupil size,

microsaccade rate, reaction time and saccade velocity. A direct prediction of our results is that an

opposite pattern should emerge when these measures are correlated with pre-stimulus α power and

P1 amplitude.

<u>Correlating pre-stimulus α power/P1 amplitude with other arousal-related variables</u>

In order to directly compare pre-stimulus α power and P1 amplitude with a host of arousal-

related variables we binned the task performance (hit rate and false rate) and eye metrics (pupil size,

microsaccade rate, reaction time, saccade velocity) in the manner described above using a 30-minute

sliding window stepped every six minutes. We computed Pearson product-moment correlations

between these metrics for each session and we compared the subsequent distributions of Pearson's r

values to shuffled distributions using permutation tests (two-sided, difference of medians). First, we

found that the amplitude of pre-stimulus oscillations in the α band was negatively correlated with

false alarm rate (Figure 3A, median r = -0.39, p < 0.001). No significant correlation was found

between pre-stimulus α power and hit rate (median r = 0.04; p = 0.783) but false alarm rate was itself

positively correlated with hit rate across sessions (Figure S2, median r = 0.49, p < 0.001). Hence, our

results support an emerging body of work showing that the amplitude of pre-stimulus α oscillations

is associated with changes in response criterion as opposed to sensitivity [32,43]. In addition, a

negative correlation was found across sessions between pre-stimulus α power and pupil size in

Monkey 1 (median r = -0.32, p = 0.037), but no such correlation was found in Monkey 2 (median r = 0.07 p = 0.623). We found a significant (p<0.05) correlation between pre-stimulus α power and pupil size in 7/20 sessions for Monkey 1 and 2/16 sessions for Monkey 2. Across sessions and monkeys, we found positive correlations of pre-stimulus α power with reaction time (Figure 3A, median r = 0.26, p = 0.012), and P1 amplitude with saccade velocity (Figure 3B, median r = 0.21, p = 0.026). Negative correlations of pre-stimulus α power with saccade velocity (median r = -0.27, p = 0.002), and P1 amplitude with microsaccade rate (median r = -0.38, p = 0.008) were also observed. No significant correlation was found between P1 amplitude and reaction time (median r = 0.08, p = 0.728), but reaction time was itself negatively correlated with saccade velocity across sessions (Figure S2, median r = -0.52, p < 0.001). Taken together, these results demonstrate that slow time scale changes in spontaneous and evoked EEG signals were accompanied by changes in the subjects' arousal levels over time. In particular, decreases in pre-stimulus α power and increases in P1 amplitude were accompanied by increases in false alarm rate and saccade velocity as well as decreases in microsaccade rate and reaction time.

Figure 3. Correlations between EEG signals and other arousal-related variables. (A) Pre-stimulus α power. Each blue point corresponds to the median r value across sessions between pre-stimulus α power and a given metric. In contrast, grey pints represent median r values computed using shuffled data. Actual distributions of r values were compared to shuffled distributions using two-sided permutation tests (difference of medians). Asterisks indicative a significant effect with a p value at least < 0.05. See Figure S2 for histograms showing distributions of r values (black) across sessions between: A) hit rate and false alarm rate; and B) reaction time and saccade velocity.

Correlations between pre-stimulus α power, P1 amplitude and slow drift

        As described above, we and others have shown that pre-stimulus α power and P1 amplitude are related to global changes in arousal. Therefore, we were keen to explore if these indirect and non-invasive measures of neural activity are associated with low-dimensional modes of neural

activity acquired directly from the spiking activity of a population of neurons. Previously, we reported

a co-fluctuation of neural activity in macaque visual and prefrontal cortex [17]. This 'slow drift' was

related to the subject's tendency to make impulsive decisions in a change detection task, ignoring

sensory evidence (false alarms), and multiple eye metrics that together are indicative of global

changes in brain state [18]. Here we sought to determine if pre-stimulus α power and P1 amplitude

are associated with slow drift.

To calculate slow drift, we binned spike counts in V4 using the same 30-minute sliding

window described above and used in previous research (Figure 4A, see Methods). We then applied

principal component analysis (PCA) to the data and estimated slow drift by projecting the binned

residual spike counts along the first principal component (i.e. the loading vector that explained the

most variance in the data). Because the sign of the loadings in PCA is arbitrary [47], the correlation

between slow drift and a given variable can be positive or negative. This would not affect the overall

pattern of results if one were solely interested in total variance explained. However, we were

interested in whether slow drift was associated with a characteristic pattern of EEG activity that we

and others have shown to be indicative of changes in the subjects' arousal levels over time i.e.

decreased pre-stimulus α power and increased P1 amplitude (Figure 2 and Figure 3). Hence, the sign

of the correlation between slow drift and the spontaneous/evoked EEG signals was critical. In order

to establish a common frame of reference across sessions, and preserve the sign of the correlations,

we constrained the slow drift for each session to have the same relationship to the spontaneous and

evoked activity of the neurons. That is, the slow drift was flipped if the mean projection value for

evoked responses recorded during stimulus periods was less than that for spontaneous responses

recorded during pre-stimulus periods i.e., if the relationship that was naturally observed for

unprojected data did not hold true [18]. This served to align the data across sessions such that

relatively large slow drift values were associated with higher firing rates. To put it another way, a

negative correlation between pre-stimulus α power and slow drift would be indicative of a pattern in

which decreased pre-stimulus α power would lead to higher firing rates. Our hypothesis predicts that

the opposite should be true for P1 amplitude.

We computed the slow drift of the neuronal population in each session using the above-

mentioned method, and then compared it to our two spontaneous and evoked EEG measures: pre-

stimulus α power and P1 amplitude. An example session is shown in Figure 4B (same sessions as in

Figure 2C). In support of our hypothesis, a characteristic pattern was found in which slow drift was

negatively associated with pre-stimulus α power and positively associated with P1 amplitude. Next,

we investigated if a similar pattern was found across sessions.

347
348
349

Figure 4. Calculating slow drift. (A) Three example neurons from a single session (Monkey 1). Each point represents

the mean residual spike count during a 400ms stimulus period. The data was then binned using a 30-minute sliding

window stepped every six minutes (solid line) so that direct comparisons could be made with the EEG signals. PCA

was used to reduce the dimensionality of the data and slow drift was calculated by projecting binned residual spike

counts along the first principal component. (B) Example session from Monkey 1 showing that slow drift was

negatively correlated with pre-stimulus α power and positively correlated with P1 amplitude. Each metric has been

z-scored for illustration purposes.

We computed the Pearson product-moment correlation coefficient between pre-stimulus α

power/P1 amplitude and slow drift for each session. We then compared the actual distribution of

Pearson's r values to shuffled distributions using permutation tests (two-sided, difference of medians).

Because the slow drift was aligned across sessions based on the neural activity alone and not the EEG

metrics, the shuffled distributions were centered on a correlation value of zero. We also computed

correlations between slow drift and pre-stimulus oscillations in the beta (β) and gamma (γ) frequency

bands to rule out the possibility that slow drift was associated with aperiodic (i.e. broadband) changes

in spectral power over time (1/f noise) [43–46]. If this was the case, one might expect a significant

correlation between slow drift and the amplitude of pre-stimulus oscillations in the α, β and γ

frequency bands.

368   Consistent with the pattern of results observed in several individual sessions for Monkey 1

and Monkey 2 (Figure S1), we found that slow drift was negatively correlated with pre-stimulus α

power (median r = -0.38, p = 0.004), whereas it was positively correlated with P1 amplitude (median

r = 0.31, p = 0.027). No significant correlation was found between slow drift and the amplitude of

pre-stimulus power in the β (median r = 0.04, p = 0.078) or γ (median r = 0.09, p = 0.621) bands.

Although these findings suggest that slow drift was not associated with changes in 1/f noise over time

[43–46] we carried out an additional analysis in which aperiodic signals were removed prior to

computing mean oscillatory power in different frequency bands. Importantly, the overall pattern of

results remained the same when slow drift was correlated with aperiodic-adjusted power in the α

(median r = -0.18 , p = 0.016), β (median r = - 0.04 , p = 0.145) and γ (median r = 0.07, p = 0.773)

bands (Figure S3). Therefore, we found that spontaneous and evoked EEG signals are associated with

slow drift even after controlling for changes in 1/f noise [43–46].

Figure 5. Correlations between EEG signals and slow drift in visual cortex. Histogram showing distributions of r values (colors) across sessions between slow drift and spontaneous/evoked EEG signals. Actual distributions were compared to shuffled distributions (grey) using two-sided permutation tests (difference of medians). Median r values are indicated by dashed lines. p < 0.05*, p < 0.01**, p < 0.001***. See Figure S3 for histograms showing distributions of r values across sessions between slow drift and aperiodic-adjusted power in different frequency bands.

Discussion

In this study, we investigated if pre-stimulus α oscillations and the amplitude of the P1 component of the VEP could be utilized as an external signature of an internal brain state, a recently discovered neural activity pattern called slow drift [17]. We know from previous work that slow drift in macaque visual and prefrontal cortex is significantly associated with a host of eye metrics that are strongly indicative of changes in arousal [18]. Since pre-stimulus α power and P1 amplitude are also related to arousal, we wondered if a link could be established between a neural measure of an internal

brain state acquired directly from the spiking activity of populations of neurons (i.e., slow drift) and

indirect signals recorded non-invasively using EEG. Results showed that slow drift was significantly

associated with a pattern that is indicative of changes in the subjects' arousal levels over time. When

pre-stimulus α power was low and P1 amplitude was high, pupil size, false alarm rate and saccade

velocity increased, whereas microsaccade rate and reaction time decreased.

A variety of studies have shown that pre-stimulus α power and P1 amplitude are related to a

behavioral pattern that is associated with arousal level. For example, decreases in the amplitude of

pre-stimulus α oscillations are typically accompanied by improved performance on detection tasks

[19–26] and decreased reaction time [35–38], whereas the opposite is true of P1 amplitude [19,25,39–

41]. This motivated us to ask if pre-stimulus α power was negatively associated with P1 amplitude.

In support of previous research [43], we found that decreases in pre-stimulus α power were

accompanied by increases in P1 amplitude. This antagonistic effect might be related to functional

inhibition. Evidence suggests that neuronal excitability is reduced when pre-stimulus α oscillations

are relatively strong [48–53]. Since early VEP components are thought to be generated in an additive

manner on top of spontaneous activity [54–57] inhibition of low-level sensory areas might lead to a

reduction in P1 amplitude. Regardless of the underlying mechanism, a direct prediction of our results

is that pre-stimulus α oscillations and P1 amplitude should be correlated in an opposite manner with

a constellation of arousal-related variables.

In this study, we explored if pre-stimulus α oscillations and P1 amplitude were associated

with various task performance and eye metrics that have been strongly linked to global changes in

brain state. Results showed that pre-stimulus α power was negatively correlated with false alarm rate

and saccade velocity, whereas it was positively correlated with reaction time. In contrast, P1

amplitude was negatively correlated with saccade velocity and microsaccade rate. Taken together,

these findings suggest that the subjects' arousal levels were changing slowly over time. As described

above, decreases in pre-stimulus α power and increases in P1 amplitude are known to be associated

with improved performance on detection tasks [19–26,39,40]. That these metrics were also coupled

with increases in false alarm rate and saccade velocity as well as decreases in microsaccade rate and reaction time suggests that they can be taken as external measures of heightened arousal levels. In addition, our findings extend recent work in humans [31,32] showing that pre-stimulus α power is associated with changes in response criterion as opposed to sensitivity.

The principles of signal detection theory [27] dictate that improved performance on detection tasks can arise due to changes in sensitivity or response criterion [28–30]. Recent work has shown that decreased pre-stimulus α power is associated with increased hit rate and false alarm rate, which suggests that performance improvements linked to decreased pre-stimulus α power might occur due to changes in response criterion [31,32]. Our results support this hypothesis, as pre-stimulus α power was negatively associated with false alarm rate. Furthermore, we found that slow drift was positively correlated with hit rate (median $r = 0.41$, $p < 0.001$) and false alarm rate (median $r = 0.53$, $p < 0.001$) across sessions, consistent with previous research [17]. Luo and Maunsell (2015) found that changes in sensitivity, not response criterion were related to single-neuron and pairwise statistics in V4. In contrast, our results suggest that V4 activity is related to a shift in response criterion that was indexed by decreased pre-stimulus α power. One possible explanation for this discrepancy is that Luo and Maunsell (2015) manipulated response criterion in a spatially selective manner to avoid shifts arising due to global changes in arousal. In our study, response criterion was not manipulated directly, so shifts are likely to have occurred spontaneously due to global changes in the subjects' arousal levels. In addition, we averaged across electrode sites because decreases in pre-stimulus α power/increases in P1 amplitude are associated with improved performance on detection tasks across large swathes of cortex [19,22,31]. These findings suggest that spatially selective and global shifts in response criterion might be governed by distinct neural mechanisms. This is an important issue to address, not just because it has the potential to resolve a discrepancy in the literature, but it also might explain why pre-stimulus α power on spatial attention tasks is associated with a pattern that is more akin to changes in sensitivity i.e. decreased/increased pre-stimulus α power over electrodes contralateral/ipsilateral to the attended hemifield [58].

In summary, we found that two commonly used metrics of cognitive state in human EEG

studies, pre-stimulus α power and P1 amplitude, are associated with gradual shifts in the underlying

population structure of neural activity throughout the brain. Together, these measures at the scalp, and

in the cortex, were predictive of changes in the monkeys' arousal levels over time. These findings

show that indirect measures of neural activity can be used to index a global signature of arousal. They

support recent work showing that pre-stimulus α power is associated with response criterion [59].

Finally, by linking a vast EEG literature in humans with simultaneous scalp/microelectrode array

recordings in macaques our results bridge the gap between large-scale signals and the responses of

single cortical neurons.

Methods

Subjects

Two adult rhesus macaque monkeys (*Macaca mulatta*) were used in this study. A previous

report [60] presented analysis of different aspects of the same experiments described here. Surgical

procedures to chronically implant a titanium head post (to immobilize the subjects' heads during

experiments) and microelectrode arrays were conducted in aseptic conditions under isoflurane

anesthesia, as described in detail by Smith and Sommer (2013). Opiate analgesics were used to

minimize pain and discomfort during the perioperative period. Neural activity was recorded using

100-channel "Utah" arrays (Blackrock Microsystems) in V4 (Monkey 1 = right hemisphere; Monkey

2 = left hemisphere). The arrays comprised a 10x10 grid of silicon microelectrodes (1 mm in length)

spaced 400 μm apart. Experimental procedures were approved by the Institutional Animal Care and

Use Committee of the University of Pittsburgh and were performed in accordance with the United

States National Research Council's Guide for the Care and Use of Laboratory Animals.

Microelectrode array recordings

Signals from each microelectrode in the array were amplified and band-pass filtered (0.3–

7500 Hz) by a Grapevine system (Ripple). Waveform segments crossing a threshold (set as a multiple

of the root mean square noise on each channel) were digitized (30KHz) and stored for offline analysis

and sorting. First, waveforms were automatically sorted using a competitive mixture decomposition

method [62]. They were then manually refined using custom time amplitude window discrimination

software (code available at https://github.com/smithlabvision/spikesort), which takes into account

metrics including (but not limited to) waveform shape and the distribution of interspike intervals [63].

A mixture of single and multiunit activity was recorded, but we refer here to all units as "neurons".

The mean number of V4 neurons across sessions was 70 (SD = 11) for Monkey 1 and 31 (SD = 16)

for Monkey 2.

EEG recordings

We recorded EEG from 8 Ag/AgCl electrodes (Grass Technologies) adhered to the scalp with

electrically conductive paste. The electrodes were positioned roughly at the following locations: Fz,

Iz, CP3, CP4, F5, F6, PO7, and PO8 (see Fig. 1B). Signals were referenced online to a steel screw on

the titanium head post, digitized at 1 kHz and amplified by a Grapevine system (Ripple) and low-

pass filtered online at 250 Hz. Segments of EEG data (a segment is defined as a 400ms stimulus

period or a 300-500ms pre-stimulus period) were considered excessively noisy and removed if any

of the channels had a standard deviation 10 times greater than the mean of the entire session. We also

removed segments of data if any of the channels exhibited a flat signal defined as a standard deviation

less than 300 nanovolts.

Visual stimuli

Visual stimuli were generated using a combination of custom software written in MATLAB

(The MathWorks) and Psychophysics Toolbox extensions [64,65]. They were displayed on a CRT

monitor (resolution = 1024 X 768 pixels; refresh rate = 100Hz), which was viewed at a distance of 36

cm and gamma-corrected to linearize the relationship between input voltage and output luminance

using a photometer and look-up-tables.

Behavioral task

Subjects fixated a central point (diameter = 0.6°) on the monitor to initiate a trial (Figure 1A).

Each trial comprised a sequence of stimulus periods (400ms) separated by fixation periods (duration

drawn at random from a uniform distribution spanning 300-500ms). The 400ms stimulus periods

comprised pairs of drifting full-contrast Gabor stimuli. One stimulus was presented in the aggregate

receptive field (RF) of the recorded V4 neurons, whereas the other stimulus was presented in the

mirror-symmetric location in the opposite hemifield. Although the spatial (Monkey 1 = 0.85cycles/°;

Monkey 2 = 0.85cycles/°) and temporal frequencies (Monkey 1 = 8cycles/s; Monkey 2 = 7cycles/s)

of the stimuli were not optimized for each individual V4 neuron they did evoke a strong response

from the population. The orientation of the stimulus in the aggregate RF was chosen at random to be

45 or 135°, and the stimulus in the opposite hemifield was assigned the other orientation. There was

a fixed probability (Monkey 1 = 30%; Monkey 2 = 40%) that one of the Gabors would change

orientation by ±1, ±3, ±6, or ±15° on each stimulus presentation. The sequence continued until the

subject) made a saccade to the changed stimulus within 400ms ("hit"); 2) made a saccade to an

unchanged stimulus ("false alarm"); or 3) remained fixating for more than 400ms after a change

occurred ("miss"). If the subject correctly detected an orientation change, they received a liquid

reward. In contrast, a time-out occurred if the subject made a saccade to an unchanged stimulus

delaying the beginning of the next trial by 1s. It is important to note that the effects of spatial attention

were also investigated (although not analyzed in this study) by cueing blocks of trials such that the

orientation change was 90% more likely to occur in one hemifield relative to the other hemifield.

<u>Eye tracking</u>

Eye position and pupil diameter were recorded monocularly at a rate of 1000Hz using an

infrared eye tracker (EyeLink 1000, SR Research).

<u>Microsaccade detection</u>

Microsaccades were defined as eye movements that exceeded a velocity threshold of 6 times

the standard deviation of the median velocity for at least 6ms [66–68]. They were required to be

separated in time by at least 100ms. In addition, we removed microsaccades with an amplitude greater

than 1° and a velocity greater than 100°/s. To assess the validity of our microsaccade detection

method, the correlation (Pearson product-moment correlation coefficient) between the amplitude and

peak velocity of detected microsaccades (i.e., the main sequence) was computed for each session.

The mean correlation between these two metrics across sessions was 0.86 (SD = 0.07) indicating that

our detection algorithm was robust [69].

<u>Pre-stimulus power</u>

To calculate the mean amplitude of oscillations in different frequency bands during pre-

stimulus periods the signals from all 8 EEG electrodes were averaged together. We adopted this

approach because previous research has shown that pre-stimulus α power across a wide range of

frontal, midline and occipital locations is associated with performance on detection tasks [19,22,31].

Furthermore, the overall pattern of results did not change when signals were averaged across frontal,

midline and posterior electrodes. We then computed a FFT of the Hanning-windowed segment of data

spanning the entire duration of the pre-stimulus period (300 - 500ms) and computed mean power in

the α, β and γ bands. To isolate slow changes, the data for each trial and power band was binned using

a 30-minute sliding window stepped every 6 minutes (Figure 2A).

<u>P1 amplitude</u>

To calculate the VEP during stimulus periods the signals from all 8 EEG electrodes were

averaged together. We adopted this approach for similar reasons to those described above i.e., because

P1 amplitude across a wide range of frontal, midline and occipital locations is associated with

performance on detection tasks [19]. In addition, the overall pattern of results remained the same

when signals were averaged across frontal, midline and posterior electrodes. We then computed the

VEP for the period spanning 200ms before to 200ms after stimulus onset, filtered the segment

between 1 and 25Hz, and baseline-corrected by subtracting the average voltage for the period

spanning 200ms before stimulus onset. To isolate slow changes over time, we divided the session into

30-minute bins stepped every 6 minutes and computed the mean VEP for each time bin. We were

interested in whether slow drift was associated with the amplitude of the P1 component. Hence, we

calculated the maximum amplitude of the peak occurring ~100ms after stimulus onset from the VEP

for each time bin (Figure 2B). Note that we were only able to reliably compute VEPs in 3 sessions

for Monkey 2, which means the majority of evoked EEG data is from Monkey 1 (17 sessions).

Aperiodic-adjusted pre-stimulus power

We wanted to rule out the possibility that slow drift was associated with gradual changes in

1/f noise [43–46]. Therefore, we carried out a supplementary analysis in which slow drift was

correlated with aperiodic-adjusted power in different frequency bands. As before, signals from all 8

EEG electrodes were averaged together. We then computed a FFT of the Hanning-windowed segment

of data spanning the first 300ms of the pre-stimulus period. To isolate slow changes over time, we

divided the session into 30-minute bins stepped every 6 minutes and computed the mean FFT for each

time bin. We then subtracted the aperiodic signal from the FFT for each time bin, which was computed

by fitting an exponential function [46]. Finally, mean pre-stimulus power in the α, β and γ bands was

calculated from the aperiodic-adjusted FFT for each time bin. Note that the frequency axis of the FFT

for each pre-stimulus interval had to be identical in this analysis so as to reliably estimate the aperiodic

signal. Since accounting for broadband changes in 1/f noise did not change the overall pattern of

results (Figure S3) we decided to correlate slow drift with pre-stimulus power in different frequency

bands using the full duration of the pre-stimulus interval (300-500ms) to enhance frequency

resolution.

Task performance metrics

Hit rate was defined as the number of saccades toward a stimulus that changed orientation

divided by the total number of stimuli presented that changed orientation. False alarm rate was the

number of saccades toward a stimulus that did not change orientation divided by the total number of

stimuli presented that did not change orientation. To isolate slow changes, hit rate and false alarm rate

were calculated in 30-minute bins stepped every 6 minutes. The width of the bins and step size were

chosen based on previous research to ensure reliable estimates over a relatively small number of trials

[17].

 Eye metrics

 Mean pupil diameter was measured during stimulus periods, whereas microsaccade rate was

 measured during fixation periods between the visual stimulus presentations [18]. We did not include

 the initial fixation period when measuring microsaccade rate as there was an increase in eye position

 variability during this period resulting from fixation having been established a short time earlier (300-

 500ms). Such variability was not present in subsequent fixation periods. Reaction time and saccade

 velocity were measured on trials in which the subjects were rewarded for correctly detecting an

 orientation change. Reaction time was defined as the time from when the change occurred to the time

 at which the saccade exceeded a velocity threshold of 100°/s. Saccade velocity was the peak velocity

 of the saccade to the changed stimulus. To isolate slow changes in the eye metrics over time the data

 for each session was binned using a 30-minute sliding window stepped every 6 minutes.

 Calculating slow drift

 The spiking responses of populations of neurons in V4 were measured during a 400ms period

 that began 50ms after stimulus presentation (Figure 4A). To control for the fact that some neurons

 had a preference for one orientation (45 or 135°) over the other residual spike counts were calculated.

 We subtracted the mean response for a given orientation across the entire session from individual

 responses to that orientation. To isolate slow changes in neural activity over time, residual spike

 counts for each V4 neuron were binned using a 30-minute sliding window stepped every 6 minutes

 (Figure 4A). PCA was then performed to reduce the high-dimensional residual data to a smaller

 number of latent variables [70]. Slow drift in V4 was estimated by projecting the binned residual

 spike counts for each neuron along the first principal component.

 Aligning slow drift across sessions

 As described above, slow drift was calculated by projecting binned residual spike counts along

 the first principal component. The weights in a PCA can be positive or negative [47], which meant

 the sign of the correlation between slow drift and a given metric was arbitrary. Preserving the sign of

 the correlations was particularly important in this study because we were interested in whether slow

drift was associated with a pattern that is indicative of changes in the subjects' arousal levels over time i.e., decreased pre-stimulus α power and increased P1 amplitude. The method for aligning slow drift across sessions is described in detail by Johnston et al. [18]. Briefly, we projected evoked responses acquired during stimulus periods and spontaneous responses acquired during pre-stimulus periods onto the first principal component. The sign of the slow drift was flipped if the mean projection value for the evoked responses was less than that for the spontaneous responses i.e., if the relationship that was naturally observed for unprojected data did not hold true.

Figure S1. Relationships between slow drift, pre-stimulus α power and P1 amplitude (A) Example sessions from Monkey 1. Each metric has been z-scored for illustration purposes. (B) Same as (A) but for Monkey 2.

Figure S2. (A) Histogram showing distributions of r values (black) across sessions between hit rate and false alarm rate. (B) Same as (A) but for r values computed between reaction time and saccade velocity. In (A) and (B) actual distributions were compared to shuffled distributions (grey) using two-sided permutation tests (difference of medians). Median r values are indicated by dashed lines. p < 0.05*, p < 0.01**, p < 0.001***.

Figure S3. Correlations between aperiodic-adjusted power and slow drift. Histograms showing distributions of r values (colors) across sessions between slow drift and EEG power in different frequency bands. Actual distributions were compared to shuffled distributions (grey) using two-sided permutation tests (difference of medians). Median r values are indicated by dashed lines. p < 0.05*, p < 0.01**, p < 0.001***.

References

1. Britten, K.H., Newsome, W.T., Shadlen, M.N., Celebrini, S., and Movshon, J.A. (1996). A relationship between behavioral choice and the visual responses of neurons in macaque MT. Vis. Neurosci. *13*, 87–100.

2. Moran, J., and Desimone, R. (1985). Selective attention gates visual information processing in extrastriate cortex. Science (80-.). *229*, 782–784.

3. Fuster, J.M., and Alexander, G.E. (1971). Neuron activity related to short-term memory. Science (80-.). *173*, 652–654.

4. Cohen, M.R., and Maunsell, J.H.R. (2009). Attention improves performance primarily by reducing interneuronal correlations. Nat. Neurosci. *12*, 1594–1600.

5. Leavitt, M.L., Pieper, F., Sachs, A.J., and Martinez-Trujillo, J.C. (2017). Correlated variability modifies working memory fidelity in primate prefrontal neuronal ensembles. Proc. Natl. Acad. Sci. U. S. A. *114*, E2494–E2503.

6. Zohary, E., Shadlen, M.N., and Newsome, W.T. (1994). Correlated neuronal discharge rate and its implications for psychophysical performance. Nature *370*, 140–143.

7. Murray, J.D., Bernacchia, A., Roy, N.A., Constantinidis, C., Romo, R., and Wang, X.J. (2017). Stable population coding for working memory coexists with heterogeneous neural dynamics in prefrontal cortex. Proc. Natl. Acad. Sci. U. S. A. *114*, 394–399.

8. Harvey, C.D., Coen, P., and Tank, D.W. (2012). Choice-specific sequences in parietal cortex during a virtual-navigation decision task. Nature *484*, 62–68.

9. Khanna, S.B., Snyder, A.C., and Smith, M.A. (2019). Distinct sources of variability affect eye movement preparation. J. Neurosci. *39*, 4511–4526.

10. Mante, V., Sussillo, D., Shenoy, K. V., and Newsome, W.T. (2013). Context-dependent computation by recurrent dynamics in prefrontal cortex. Nature *503*, 78–84.

11. Sohn, H., Narain, D., Meirhaeghe, N., and Jazayeri, M. (2018). Bayesian computation through cortical latent dynamics. bioRxiv, 465419. Available at:

791  https://www.biorxiv.org/content/early/2018/11/08/465419.

792 12. Remington, E.D., Narain, D., Hosseini, E.A., and Jazayeri, M. (2018). Flexible Sensorimotor

793   Computations through Rapid Reconfiguration of Cortical Dynamics. Neuron *98*, 1005-

794   1019.e5. Available at: https://doi.org/10.1016/j.neuron.2018.05.020.

795 13. Wang, J., Narain, D., Hosseini, E.A., and Jazayeri, M. (2018). Flexible timing by temporal

796   scaling of cortical responses. Nat. Neurosci. *21*, 102–112. Available at:

797   http://dx.doi.org/10.1038/s41593-017-0028-6.

798 14. Sadtler, P.T., Quick, K.M., Golub, M.D., Chase, S.M., Ryu, S.I., Tyler-Kabara, E.C., Yu,

799   B.M., and Batista, A.P. (2014). Neural constraints on learning. Nature *512*, 423–426.

800   Available at: http://dx.doi.org/10.1038/nature13665.

801 15. Stringer, C., Pachitariu, M., Steinmetz, N., Reddy, C.B., Carandini, M., and Harris, K.D.

802   (2019). Spontaneous behaviors drive multidimensional, brainwide activity. Science (80-.).

803   *364*.

804 16. Musall, S., Kaufman, M.T., Juavinett, A.L., Gluf, S., and Churchland, A.K. (2019). richly

805   varied movements. Nat. Neurosci. *22*. Available at: http://dx.doi.org/10.1038/s41593-019-

806   0502-4.

807 17. Cowley, B., Snyder, A., Acar, K., Williamson, R., Yu, B., and Smith, M. (2020). Slow drift of

808   neural activity as a signature of impulsivity in macaque visual and prefrontal cortex. Neuron,

809   1–17. Available at: https://doi.org/10.1016/j.neuron.2020.07.021.

810 18. Johnston, R., Snyder, A.C., Khanna, S., Issar, D., and Smith, M.A. (2020). The eyes reflect

811   an internal cognitive state embedded in the population activity of cortical neurons. bioRxiv,

812   2020.06.29.178251. Available at:

813   http://biorxiv.org/content/early/2020/06/30/2020.06.29.178251.abstract.

814 19. Ergenoglu, T., Demiralp, T., Bayraktaroglu, Z., Ergen, M., Beydagi, H., and Uresin, Y.

815   (2004). Alpha rhythm of the EEG modulates visual detection performance in humans. Cogn.

816   Brain Res. *20*, 376–383.

20.   Babiloni, C., Vecchio, F., Bultrini, A., Romani, G.L., and Rossini, P.M. (2006). Pre- and

poststimulus alpha rhythms are related to conscious visual perception: A high-resolution EEG

study. Cereb. Cortex *16*, 1690–1700.

21.   Van Dijk, H., Schoffelen, J.M., Oostenveld, R., and Jensen, O. (2008). Prestimulus

oscillatory activity in the alpha band predicts visual discrimination ability. J. Neurosci. *28*,

1816–1823.

22.   Busch, N.A., Dubois, J., and VanRullen, R. (2009). The phase of ongoing EEG oscillations

predicts visual perception. J. Neurosci. *29*, 7869–7876.

23.   Thut, G., Nietzel, A., Brandt, S.A., and Pascual-Leone, A. (2006). α-Band

electroencephalographic activity over occipital cortex indexes visuospatial attention bias and

predicts visual target detection. J. Neurosci. *26*, 9494–9502.

24.   Hanslmayr, S., Aslan, A., Staudigl, T., Klimesch, W., Herrmann, C.S., and Bäuml, K.H.

(2007). Prestimulus oscillations predict visual perception performance between and within

subjects. Neuroimage *37*, 1465–1473.

25.   Mathewson, K.E., Gratton, G., Fabiani, M., Beck, D.M., and Ro, T. (2009). To see or not to

see: Prestimulus α phase predicts visual awareness. J. Neurosci. *29*, 2725–2732.

26.   Romei, V., Gross, J., and Thut, G. (2010). On the role of prestimulus alpha rhythms over

occipito-parietal areas in visual input regulation: Correlation or causation? J. Neurosci. *30*,

8692–8697.

27.   Green, D.M., and Swets, J.A. (1966). Signal detection theory and psychophysics (Wiley).

28.   Luo, T.Z., and Maunsell, J.H.R. (2015). Neuronal Modulations in Visual Cortex Are

Associated with Only One of Multiple Components of Attention. Neuron *86*, 1182–1188.

Available at: http://dx.doi.org/10.1016/j.neuron.2015.05.007.

29.   Luo, T.Z., and Maunsell, J.H.R. (2018). Attentional Changes in Either Criterion or Sensitivity

Are Associated with Robust Modulations in Lateral Prefrontal Cortex. Neuron *97*, 1382-

1393.e7. Available at: https://doi.org/10.1016/j.neuron.2018.02.007.

30.   Crapse, T.B., Lau, H., and Basso, M.A. (2018). A Role for the Superior Colliculus in

Decision Criteria. Neuron *97*, 181-194.e6. Available at:

https://doi.org/10.1016/j.neuron.2017.12.006.

31.   Iemi, L., Chaumon, M., Crouzet, S.M., and Busch, N.A. (2017). Spontaneous neural

oscillations bias perception by modulating baseline excitability. J. Neurosci. *37*, 807–819.

32.   Limbach, K., and Corballis, P.M. (2016). Prestimulus alpha power influences response

criterion in a detection task. Psychophysiology *53*, 1154–1164.

33.   de Gee, J.W., Colizoli, O., Kloosterman, N.A., Knapen, T., Nieuwenhuis, S., and Donner,

T.H. (2017). Dynamic modulation of decision biases by brainstem arousal systems. Elife *6*,

1–36.

34.   Hong, L., Walz, J.M., and Sajda, P. (2014). Your eyes give you away: Prestimulus changes in

pupil diameter correlate with poststimulus task-related EEG dynamics. PLoS One *9*.

35.   Van Kempen, J., Loughnane, G.M., Newman, D.P., Kelly, S.P., Thiele, A., O'Connell, R.G.,

and Bellgrove, M.A. (2019). Behavioural and neural signatures of perceptual decision-

making are modulated by pupil-linked arousal. Elife *8*, 1–27.

36.   Kelly, S.P., and O'Connell, R.G. (2013). Internal and external influences on the rate of

sensory evidence accumulation in the human brain. J. Neurosci. *33*, 19434–19441.

37.   Zhang, Y., Wang, X., Bressler, S.L., Chen, Y., and Ding, M. (2008). Prestimulus cortical

activity is correlated with speed of visuomotor processing. J. Cogn. Neurosci. *20*, 1915–

1925.

38.   Bompas, A., Sumner, P., Muthumumaraswamy, S.D., Singh, K.D., and Gilchrist, I.D. (2015).

The contribution of pre-stimulus neural oscillatory activity to spontaneous response time

variability. Neuroimage *107*, 34–45.

39.   Del Cul, A., Baillet, S., and Dehaene, S. (2007). Brain dynamics underlying the nonlinear

threshold for access to consciousness. PLoS Biol. *5*, 2408–2423.

40.   Pourtois, G., De Pretto, M., and Hauert, C. (2006). Time course of brain activity during

change blindness and change awareness: performance …. J. Cogn. Neurosci., 2108–2129.

Available at:

http://www.mitpressjournals.org/doi/abs/10.1162/jocn.2006.18.12.2108%5Cnfile:///Users/Me

gan/Documents/Papers2/Articles/2006/Pourtois/Time course of brain activity during change

blindness and change awareness performance … 2006 Pourtois.pdf%5Cnpapers2://pu.

41.   Mangun, G.R., and Hillyard, S.A. (1991). Modulations of Sensory-Evoked Brain Potentials

Indicate Changes in Perceptual Processing During Visual-Spatial Priming. J. Exp. Psychol.

Hum. Percept. Perform. *17*, 1057–1074.

42.   Zoefel, B., ten Oever, S., and Sack, A.T. (2018). The involvement of endogenous neural

oscillations in the processing of rhythmic input: More than a regular repetition of evoked

neural responses. Front. Neurosci. *12*, 1–13.

43.   Iemi, L., Busch, N.A., Laudini, A., Haegens, S., Samaha, J., Villringer, A., and Nikulin, V. V.

(2019). Multiple mechanisms link prestimulus neural oscillations to sensory responses. Elife

8, 1–34.

44.   Voytek, B., Kramer, M.A., Case, J., Lepage, K.Q., Tempesta, Z.R., Knight, R.T., and

Gazzaley, A. (2015). Age-related changes in 1/f neural electrophysiological noise. J.

Neurosci. *35*, 13257–13265.

45.   Peterson, E.J., Rosen, B.Q., Campbell, A.M., Belger, A., and Voytek, B. (2017). 1/F Neural

Noise Is a Better Predictor of Schizophrenia Than Neural Oscillations. Doi.Org, 113449.

Available at: https://www.biorxiv.org/content/early/2017/03/03/113449.

46.   Haller, M., Donoghue, T., Peterson, E., Varma, P., Sebastian, P., Gao, R., Noto, T., Knight,

R.T., Shestyuk, A., and Voytek, B. (2018). Parameterizing neural power spectra. bioRxiv,

299859. Available at:

https://www.biorxiv.org/content/biorxiv/early/2018/04/11/299859.full.pdf.

47.   Jollife, I.T., and Cadima, J. (2016). Principal component analysis: A review and recent

developments. Philos. Trans. R. Soc. A Math. Phys. Eng. Sci. *374*.

48.   Haegens, S., Nácher, V., Luna, R., Romo, R., and Jensen, O. (2011). α-Oscillations in the

896   monkey sensorimotor network influence discrimination performance by rhythmical inhibition

897   of neuronal spiking. Proc. Natl. Acad. Sci. U. S. A. *108*, 19377–19382.

898   49.   Jensen, O. (2002). Oscillations in the Alpha Band (9-12 Hz) Increase with Memory Load

899   during Retention in a Short-term Memory Task. Cereb. Cortex *12*, 877–882.

900   50.   Klimesch, W., Doppelmayr, M., Schwaiger, J., Auinger, P., and Winkler, T. (1999).

901   "Paradoxical" alpha synchronization in a memory task. Cogn. Brain Res. *7*, 493–501.

902   51.   Mayhew, S.D., Ostwald, D., Porcaro, C., and Bagshaw, A.P. (2013). Spontaneous EEG alpha

903   oscillation interacts with positive and negative BOLD responses in the visual-auditory

904   cortices and default-mode network. Neuroimage *76*, 362–372. Available at:

905   http://dx.doi.org/10.1016/j.neuroimage.2013.02.070.

906   52.   Harvey, B.M., Vansteensel, M.J., Ferrier, C.H., Petridou, N., Zuiderbaan, W., Aarnoutse, E.J.,

907   Bleichner, M.G., Dijkerman, H.C., van Zandvoort, M.J.E., Leijten, F.S.S., *et al.* (2013).

908   Frequency specific spatial interactions in human electrocorticography: V1 alpha oscillations

909   reflect surround suppression. Neuroimage *65*, 424–432. Available at:

910   http://dx.doi.org/10.1016/j.neuroimage.2012.10.020.

911   53.   Van Kerkoerle, T., Self, M.W., Dagnino, B., Gariel-Mathis, M.A., Poort, J., Van Der Togt, C.,

912   and Roelfsema, P.R. (2014). Alpha and gamma oscillations characterize feedback and

913   feedforward processing in monkey visual cortex. Proc. Natl. Acad. Sci. U. S. A. *111*, 14332–

914   14341.

915   54.   Mazaheri, A., and Jensen, O. (2006). Posterior α activity is not phase-reset by visual stimuli.

916   Proc. Natl. Acad. Sci. U. S. A. *103*, 2948–2952.

917   55.   Shah, A.S., Bressler, S.L., Knuth, K.H., Ding, M., Mehta, A.D., Ulbert, I., and Schroeder,

918   C.E. (2004). Neural Dynamics and the Fundamental Mechanisms of Event-related Brain

919   Potentials. Cereb. Cortex *14*, 476–483.

56.   Mäkinen, V., Tiitinen, H., and May, P. (2005). Auditory event-related responses are generated

independently of ongoing brain activity. Neuroimage *24*, 961–968.

57.   Bijma, F., De Munck, J.C., Huizenga, H.M., and Heethaar, R.M. (2003). A mathematical

approach to the temporal stationarity of background noise in MEG/EEG measurements.

Neuroimage *20*, 233–243.

58.   Foxe, J.J., and Snyder, A.C. (2011). The role of alpha-band brain oscillations as a sensory

suppression mechanism during selective attention. Front. Psychol. *2*, 1–13.

59.   Samaha, J., Iemi, L., Haegens, S., and Busch, N.A. (2020). Spontaneous Brain Oscillations

and Perceptual Decision-Making. Trends Cogn. Sci., 1–15. Available at:

https://doi.org/10.1016/j.tics.2020.05.004.

60.   Snyder, A.C., Yu, B.M., and Smith, M.A. (2018). Distinct population codes for attention in

the absence and presence of visual stimulation. Nat. Commun. *9*, 4382. Available at:

http://www.nature.com/articles/s41467-018-06754-5.

61.   Smith, M.A., and Sommer, M.A. (2013). Spatial and Temporal Scales of Neuronal

Correlation in Visual Area V4. J. Neurosci. *33*, 5422–5432. Available at:

http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.4782-12.2013.

62.   Shoham, S., Fellows, M.R., and Normann, R.A. (2003). Robust, automatic spike sorting

using mixtures of multivariate t-distributions. J. Neurosci. Methods *127*, 111–122.

63.   Kelly, R.C., Smith, M.A., Samonds, J.M., Kohn, A., Bonds, A.B., Movshon, J.A., and Sing

Lee, T. (2007). Comparison of Recordings from Microelectrode Arrays and Single Electrodes

in the Visual Cortex. J. Neurosci. *27*, 261–264.

64.   Brainard, D.H. (1997). The Psychophysics Toolbox. Spat. Vis. *10*, 433–436.

65.   Pelli, D.G. (1997). The VideoToolbox software for visual psychophysics: Transforming

numbers into movies. Spat. Vis. *10*, 437–442.

66.   Scholes, C., McGraw, P. V., Nyström, M., and Roach, N.W. (2015). Fixational eye

movements predict visual sensitivity. Proc. R. Soc. B Biol. Sci. *282*.

67.   Engbert, R., and Kliegl, R. (2003). Microsaccades uncover the orientation of covert attention.

Vision Res. *43*, 1035–1045.

68.   Loughnane, G.M., Newman, D.P., Tamang, S., Kelly, S.P., and O'Connell, R.G. (2018).

Antagonistic interactions between microsaccades and evidence accumulation processes

during decision formation. J. Neurosci. *38*, 2163–2176.

69.   Zuber, B.L., Stark, L., and Cook, G. (1965). Microsaccades and the velocity-amplitude

relationship for saccadic eye movements. Science (80-.). *150*, 1459–1460.

70.   Cunningham, J.P., and Yu, B.M. (2014). Dimensionality reduction for large-scale neural

recordings. Nat. Neurosci. *17*, 1500–1509.