Seasonal climate variations promote bacterial α-diversity in soil

Xin-Feng Zhao, Wen-Sheng Shu, Yi-Qi Hao[*]

Authors' affiliation

School of Life Sciences, South China Normal University, Guangzhou 510631, China

* Corresponding author: Yi-Qi Hao

E-mail address: yiqihao@m.scnu.edu.cn

Telephone number: (86) 020 - 85211850

Postal address: No.55 Zhongshan Avenue West, Tianhe District, Guangzhou 510631, China

Keywords: climate seasonality, coexistence, diversity, fluctuation-dependent mechanisms, soil bacteria

Abstract

Ecological theory suggests that temporal environmental fluctuations can contribute

greatly to diversity maintenance. Given bacteria's short generation time and rapid

responses to environmental change, seasonal climate fluctuations are very likely to

play an important role in maintaining the extremely high α-diversity of soil bacterial

community, which has been unfortunately neglected in previous studies. Here, with

in-depth analyses of two previously published soil bacterial datasets at global scale,

we found that soil bacterial α-diversity was positively correlated with both seasonal

variations of temperature and precipitation. Furthermore, piecewise structural

equation models showed that seasonal variations of temperature or precipitation

directly promoted soil bacterial α-diversity in each dataset. However, it is noteworthy

that the importance of seasonal climate variations might be underestimated in the

above analyses, due to the high level of environmental variations irrelevant to climate

seasonality among sampling sites and the lack of sampling across seasons.

Supplementary analyses of a previously published wheat cropland dataset with

samples collected in both summer and winter across North China Plain was conducted.

Similarly, we found that bacterial α-diversity was positively correlated with seasonal

climate variations, but much stronger, though the range of seasonal climate variations

was much smaller compared to the two global datasets. Collectively, these findings

implied that fluctuation-dependent mechanisms of diversity maintenance presumably

operate in soil bacterial communities. Based on existing evidence, we speculated that

the storage effect may be the main mechanism responsible for diversity maintenance

in soil bacterial community, but rigorous experimental tests are needed in the future.

Main text

The extraordinarily high α-diversity of soil bacterial communities has fascinated and puzzled microbial ecologists [1, 2]. In contrast to the considerable advances in characterizing soil bacterial biogeographic patterns across spatial and environmental gradients (e.g. [3, 4]), little attempt has been made to infer the underlying mechanisms of diversity maintenance. The high spatial heterogeneity of soil was generally thought as an important factor in maintaining bacterial α-diversity [5]; surprisingly, temporal environmental fluctuations promoting diversity maintenance firstly suggested by Hutchinson was rarely mentioned, which is one of the most influential ideas in community ecology [6, 7]. The fluctuation-dependent coexistence mechanisms are particularly important for micro-organisms due to their short generation time and rapid responses to environmental changes [8, 9]. Temporal changes in temperature and precipitation can significantly alter soil physiochemical and nutritional status, which jointly enhance temporal environmental fluctuations in soil, and drive the seasonal species turnover in bacterial community [10]. However, the majority of biogeographic studies only chose the mean annual temperature (MAT) and annual precipitation (AP) as climatic predictors of soil bacterial diversity and composition, whereas seasonal climate variations were unfortunately neglected. Here we tested the prediction that seasonal climate variations promote soil bacterial α-diversity derived from the fluctuation-dependent coexistence theory, with two global datasets covering wide geographic and climate gradients: global topsoil microbiome (hereafter "topsoil data") [3] and global soil atlas (hereafter "atlas data") [11], and one regional dataset of wheat cropland soil across North China Plain (hereafter "farmland data") [12]. Details of data acquisition, processing, and statistical analyses were provided in Supplementary Methods.

The climate data were obtained from WorldClim2 database [13], whereby the

standard deviation of monthly mean temperature and the coefficient of variation of

monthly precipitation were used to represent temperature seasonality (TS) and

precipitation seasonality (PS), respectively. Soil bacterial richness was measured by

the number of observed exact sequence variants (ESVs). As predicted, soil bacterial

richness significantly increased with TS and PS in both global datasets (Fig. 1).

Piecewise structural equation models [14] were then fitted to address the direct and

indirect effects of geographic (absolute latitude), climatic (MAT, AP, TS, PS)

variables and soil physiochemical properties (pH and C/N ratio, the most important

two soil physiochemical factors in determining bacterial α-diversity and composition)

on soil bacterial richness. The modeling results demonstrated significant direct effect

of PS (Fig. 2a, Table S3; standardized coefficient $\beta = 0.195$, $p = 0.0019$) in promoting

soil bacterial richness in topsoil data, and TS (Fig. 2b, Table S5; standardized

coefficient $\beta = 0.191$, $p = 0.0006$) in atlas data after controlling for the effects of MAT,

AP, soil pH and C/N ratio. Given that richness estimation is sensitive to detection of

rare species, above analyses were repeated with Shannon diversity, which showed

similar results as richness (Fig. S3, S4, Table S4, S6).

Of note, the above two global datasets were consisted of soil samples from

various types of ecosystem, where each location was represented by one sample [3,

11]. The high level of environmental variations between samples and lack of sampling

across seasons may result in underestimating the effects of climate seasonality on

maintaining soil bacterial α-diversity. As a supplement, we analyzed farmland data

with soil samples collected in both summer and winter across North China Plain [12].

We found that soil bacterial α-diversity in summer was positively correlated with the

collective α-diversity of two seasons (Fig. 3a, Fig. S5a). Besides, both the summer

α-diversity and whole year α-diversity of soil bacterial communities were positively

correlated with TS and PS (Fig. 3b, c, Fig. S5b, c). Such results enhanced our

confidence in the global analyses with cross-sectional data. Despite the narrow range

of climate seasonality in farmland data compared to the two global datasets, climate

seasonality explained a much larger portion of variations of the whole year soil

bacterial α-diversity within one type of ecosystem (Fig. 3b, c; Fig. S5b, c), where soil

samples shared similar properties. This finding suggested that the importance of

seasonal climate variations in maintaining soil bacterial α-diversity was probably

underestimated in the global analyses. Unfortunately, the scarcity of studies

incorporating seasonal variations in soil bacterial α-diversity across climate gradients

impeded further meta-analysis.

Theoretically, there are two classes of fluctuation-dependent coexistence

mechanisms: the storage effect and relative nonlinearity of competition [6, 9]. The

storage effect mediates coexistence via temporal niche partitioning [6], and its

operation requires that (1) species differ in their responses to environments, which

results in (2) the relative strength of interspecific and intraspecific competition

varying with fluctuating environments; and that (3) there are mechanisms buffering

population from extinction under unfavorable conditions [6], such as a variety of

physiological stress-resistance and dormancy mechanisms of soil bacteria [15]. There

has been much experimental evidence that soil bacteria show diverse responses to

moisture and temperature gradients. For example, a collection of soil bacterial isolates

exhibited a wide range of responses to moisture gradient, and the derived niche

parameters suggested a potential of coexistence via partitioning the moisture niche

axis [16]. In addition to dry- and wet-adapted bacteria, stress-tolerant and

opportunistic strategies of soil bacteria in response to drought or raining events were

also discovered, as well as sensitive taxa [15, 17]. Likewise, both warm- and

cold-responsive taxa were identified [18], and soil bacterial communities can rapidly

diverge under contrasting experimental temperature treatments [19]. Besides evidence

strongly supporting differential responses to moisture and temperature (requirement 1)

discussed above, repeated reports of seasonal turnover in soil bacterial community

structure (e.g. [10]) imply altered competitive intensities among soil bacteria under

fluctuating environments (requirement 2); although distinguishing whether it is due to

the fluctuating temperature and moisture *per se*, and/or the accompanying seasonal

changes in availability of various resources is difficult. So far, rigorous tests of the

storage effect have been restricted to simplistic microcosms [20, 21]; although being

presumably important, its role in maintaining soil bacterial α-diversity in natural

system remains unexamined yet.

By contrast, relative nonlinearity of competition might not be the major

mechanism responsible for the promoting effect of seasonal climate variations on the

soil bacterial α-diversity. Coexistence via relative nonlinearity of competition relies

on that the per capita growth rates of species are different nonlinear functions of

limiting resource [6, 8, 22]. Such nonlinearity can result in the fluctuation of limiting

resource, and the stable coexistence can achieve when each species is disadvantaged

relative to the others in the fluctuation environment caused by itself, but not

necessarily requires fluctuation of external conditions [6, 8, 22]. On the other hand,

previous modeling investigations emphasizing the importance of relative nonlinearity

of competition merely focused on microorganisms in homogenous aquatic systems [8,

9], and their implications cannot be directly generalized to the soil environment with

high spatial heterogeneity, where resources are not simultaneously accessible to all

species.

In conclusion, our findings supported the idea that seasonal temperature and

precipitation variations promote soil bacterial α-diversity via fluctuation-dependent

mechanisms of diversity maintenance. Our study highlighted the importance of

considering both the average values and temporal variations of abiotic environmental

factors to understand the soil bacterial community composition. Further filed

investigations of spatial-temporal turnover of soil bacterial diversities and

experimental studies designed to investigate specific coexistence mechanisms will

shed new light on the diversity maintenance of soil bacterial communities in the

future.

Acknowledgements

We thank Mohammad Bahram for generously sharing the metadata with us.

References

1. Shade A (2017) Diversity is the question, not the answer. ISME J 11:1-6.

https://doi:10.1038/ismej.2016.118

2. Fierer N (2017) Embracing the unknown: disentangling the complexities of the soil

microbiome. Nat Rev Microbiol 15:579-590. https://doi:10.1038/nrmicro.2017.87

3. Bahram M, Hildebrand F, Forslund SK, Anderson JL, Soudzilovskaia NA,

Bodegom PM, et al. (2018) Structure and function of the global topsoil microbiome.

Nature 560:233-237. https://doi:10.1038/s41586-018-0386-6

4. Martiny JB, Bohannan BJ, Brown JH, Colwell RK, Fuhrman JA, Green JL, et al.

(2006) Microbial biogeography: putting microorganisms on the map. Nat Rev

Microbiol 4:102-112. https://doi:10.1038/nrmicro1341

5. Vos M, Wolf AB, Jennings SJ, Kowalchuk GA (2013) Micro-scale determinants of

bacterial diversity in soil. FEMS Microbiol Rev 37:936-954.

https://doi:10.1111/1574-6976.12023

6. Chesson P (2000) Mechanisms of maintenance of species diversity. Annu Rev Ecol

Syst 31:343-366. https://doi:10.1146/annurev.ecolsys.31.1.343

7. Hutchinson GE (1961) The paradox of the plankton. Am Nat 95:137-145.

https://doi.org/10.1086/282171

8. Huisman J, Weissing FJ (1999) Biodiversity of plankton by species oscillations and

chaos. Nature 402:407-410. https://doi:10.1038/46540

9. Letten AD, Dhami MK, Ke P-J, Fukami T (2018) Species coexistence through

simultaneous fluctuation-dependent mechanisms. Proc Natl Acad Sci USA

115:6745-6750. https://doi:10.1073/pnas.1801846115

10. Rasche F, Knapp D, Kaiser C, Koranda M, Kitzler B, Zechmeister-Boltenstern S,

Richter A, Sessitsch A (2011) Seasonality and resource availability control bacterial

and archaeal communities in soils of a temperate beech forest. ISME J 5:389-402.

https://doi:10.1038/ismej.2010.138

11. Delgado-Baquerizo M, Oliverio AM, Brewer TE, Benavent-Gonzalez A, Eldridge

DJ, Bardgett RD, Maestre FT, Singh BK, Fierer N (2018) A global atlas of the

dominant bacteria found in soil. Science 359:320-325.

https://doi:10.1126/science.aap9516

12. Zhang K, Delgado-Baquerizo M, Zhu Y-G, Chu H (2020) Space is more

important than season when shaping soil microbial communities at a large spatial

scale. mSystems 5:e00783-19. https://doi:10.1128/mSystems.00783-19

13. Fick SE, Hijmans RJ (2017) WorldClim 2: new 1-km spatial resolution climate

surfaces for global land areas. Int J Climatol 37:4302-4315.

https://doi:10.1002/joc.5086

14. Lefcheck JS, Freckleton R (2015) piecewiseSEM : Piecewise structural

equation modelling in R for ecology, evolution, and systematics. Methods Ecol Evol

7:573-579. https://doi:10.1111/2041-210x.12512

15. Schimel J, Balser TC, Wallenstein M (2007) Microbial stress-response physiology

and its implications for ecosystem function. Ecology 88:1386-1394.

https://doi:10.1890/06-0219

16. Lennon JT, Aanderud ZT, Lehmkuhl B, Schoolmaster Jr DR (2012) Mapping the

niche space of soil microorganisms using taxonomy and traits. Ecology 93:1867-1879.

https://doi.org/10.1890/11-1745.1

17. Evans SE, Wallenstein MD (2014) Climate change alters ecological strategies of

soil bacteria. Ecol Lett 17:155-164. https://doi:10.1111/ele.12206

18. Oliverio AM, Bradford MA, Fierer N (2017) Identifying the microbial taxa that

consistently respond to soil warming across time and space. Global Change Biol

23:2117-2129. https://doi:10.1111/gcb.13557

19. Wu J, Xiong J, Hu C, Shi Y, Wang K, Zhang D (2015) Temperature sensitivity of

soil bacterial community along contrasting warming gradient. Appl Soil Ecol

94:40-48. https://doi:10.1016/j.apsoil.2015.04.018

20. Descamps-Julien B, Gonzalez A (2005) Stable coexistence in a fluctuating

environment: an experimental demonstration. Ecology 86:2815-2824.

https://doi:10.1890/04-1700

21. Jiang L, Morin PJ (2007) Temperature fluctuation facilitates coexistence of

competing species in experimental microbial communities. J Anim Ecol 76:660-668.

https://doi:10.1111/j.1365-2656.2007.01252.x

22. Armstrong RA, McGehee R (1980) Competitive exclusion. Am Nat 115:151-170.

https://doi:10.1086/283553

 Figure legends

Fig. 1 The relationships between soil bacterial richness (the number of observed ESVs)

and temperature (a, c) or precipitation (b, d) seasonality in two global datasets (topsoil

data: red squares, atlas data: blue circles). Significant relationships are depicted by

regression lines

Fig. 2 Structural equation models (piecewise SEM) to address direct and indirect

effects of geographic (absolute latitude), climatic (mean annual temperature, annual

precipitation, temperature seasonality, precipitation seasonality) variables, and soil

physiochemical properties (pH, C/N ratio) on soil bacterial richness for topsoil data

(169 soil samples) (a) and atlas data (186 soil samples) (b). Red arrows represent

positive paths, and black arrows represent negative paths; only significant paths are

shown ($p < 0.05$). Standardized effect sizes of path coefficients are reported and

indicated by path thickness. R^2 for component models are given in the boxes of

corresponding response variables. Shipley's test of directed separation: Fisher's *C*

statistic (if $p > 0.05$, then there are no missing associations and the model reproduces

the data well) and AIC are used to evaluate the overall fit of the model. Details of

model evaluation, simplification and coefficients estimation are provided in

supplementary methods and Table S3, S5

Fig. 3 The relationship between soil bacterial richness in summer and of the whole

year (a), and the relationships between soil bacterial richness (summer: red squares;

whole year: blue circles) and temperature (b) or precipitation (c) seasonality. In total

of 90 soil samples were collected from 45 soil plots (geographic distances up to 873

km) across North China Plain in winter and the following summer. Richness of the

whole year is calculated by merging community compositions in summer and winter

for each plot. Significant relationships are depicted by regression lines

(a) Topsoil data

Fisher's C = 6.588, p = 0.884
AIC = 78.588

(b) Atlas data

Fisher's C = 9.032, p = 0.912
AIC = 77.032