1

Concerted evolution reveals co-adapted amino acid substitutions in frogs that
prey on toxic toads

Authors: Shabnam Mohammadi[1]*, Lu Yang[2]*, Arbel Harpak[3]*, Santiago Herrera-Álvarez[4, ψ],
María del Pilar Rodríguez-Ordoñez[4, #], Julie Peng, Karen Zhang[2], Jay F. Storz[1], Susanne Dobler[6],
Andrew J. Crawford[4]** and Peter Andolfatto[3]**

Affiliations:

[1] School of Biological Sciences, University of Nebraska, Lincoln, NE, USA

[2] Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA

[3] Department of Biological Sciences, Columbia University, New York, NY, USA

[4] Department of Biological Sciences, Universidad de los Andes, Bogotá, 111711, Colombia

[5] Lewis-Sigler Institute, Princeton University, Princeton, NJ, USA

[6] Molecular Evolutionary Biology, Zoological Institute, Universität Hamburg, Hamburg,
Germany

*Co-first authorship

**Correspondence to: andrew@dna.ac, pa2543@columbia.edu

ψ Current address: Dept. of Ecology and Evolution, University of Chicago, Chicago, IL, USA
Current address: Université Paris-Saclay Evry, Evry, France

ABSTRACT (141 words):

Gene duplication is an important source of evolutionary innovation, but the adaptive division-of-labor between duplicates can be opposed by ongoing gene conversion between them. Here we document a tandem duplication of Na^+,K^+-ATPase subunit α1 (ATP1A1) shared by frogs in the genus *Leptodactylus*, a group of species that feeds on toxic toads. One ATP1A1 paralog evolved resistance to toad toxins while the other copy retained ancestral susceptibility. We show that the two *Leptodactylus* paralogs are distinguished by 12 amino acid substitutions that were maintained by strong selection that counteracted the homogenizing effect of gene conversion. Protein-engineering experiments show that two major-effect substitutions confer toxin resistance, whereas the 10 additional substitutions mitigate deleterious pleiotropic effects on enzyme function. Our results highlight how trans-specific, neofunctionalized gene duplicates can provide unique insights into interactions between adaptive substitutions and the genetic backgrounds on which they arise.

One Sentence Summary: Selection counteracts gene conversion to maintain an adaptive division-of-labor between tandemly duplicated genes.

Main Text:

The repeated evolution of toxin resistance in animals is one of the clearest examples of natural selection at the molecular level and represents a useful paradigm to examine constraints on the evolution of novel protein functions (*1*). Neotropical Grass Frogs of the genus *Leptodactylus* (Leptodactylidae) are widely distributed throughout lowland South America and are known to feed on chemically-defended toads – a predatory tendency that is rare among frogs (*2–6*). A major component of the chemical defense secretions of toads is a class of cardiac glycosides (CGs) called "bufadienolides" (*7*) that inhibit the α-subunit of Na^+,K^+-ATPases (ATP1A). Na^+,K^+-ATPases are transmembrane proteins that are vital to numerous physiological processes in animals including neural signal transduction, muscle contraction, and cell homeostasis (*8, 9*). CGs bind to the extracellular surface of ATP1A and block the flux of ions (*10*), making them potent poisons to most animals. However, some vertebrates have independently evolved the ability to prey on chemically-defended toads, partly via amino acid substitutions to the CG-binding domain of ATP1A1 that confer molecular resistance to CGs (*11–14*).

Most vertebrates share several paralogous copies of ATP1A that have different tissue-specific expression profiles (*15*). For example, ATP1A1 is the most ubiquitously expressed paralog and ATP1A3 has enriched expression in nervous tissue and heart muscle (*16, 17*; Fig. S1). Previous studies on the molecular convergence of CG-resistance in reptiles have focused primarily on the αM1–2 extracellular loop of ATP1A3 (*12–14, 18*), whereas studies of birds, mammals, and amphibians have focused on the same region of ATP1A1 (*11, 18*). A survey of αM1–2 in toads and frogs (*11*) revealed a possible duplication of ATP1A1 in the toad-eating frog, *Leptodactylus latrans* (reported as *L. ocellatus*), where the resistant (R) copy includes substitutions known to

confer resistance to CGs while the sensitive (S) copy appears to have retained the ancestral

susceptibility to CGs. Duplication and neofunctionalization of ATP1A are a recurrent adaptive

mechanism in CG-resistant insects (*19–22*) but appear to be rare among CG-resistant vertebrates.

Further, the fate of duplicate genes and the probability that they will neofunctionalize is

predicted to depend on the strength of selection for duplicate diversification relative to the rate of

non-allelic gene conversion (NAGC)—nonreciprocal recombination between duplicates that

homogenizes sequence variation between them, thereby impeding divergence (*23–25*; Fig. 2A).

The ATP1A1 gene family in *Leptodactylus* therefore provide a unique opportunity to explore

this premise further in vertebrates.

We systematically surveyed the full-length coding sequences for all ATP1A paralogs in

Leptodactylus and other anurans using a combination of RNA-seq-based gene discovery (*19*)

(Tables S1 and S2), targeted sequencing of the full length ATP1A1 gene (Methods), and a *de*

novo genome assembly for *Leptodactylus fuscus* (Table S3). As predicted by previous findings

(*11*), we confirmed that ATP1A1 is duplicated in *L. latrans*. By generating a *de novo* genome

assembly for *L. fuscus* (Table S3), we find that the two duplicate copies are arranged in tandem

(Fig. S2), and are therefore potentially subject to non-allelic gene conversion. Our results also

indicate that duplication of ATP1A1 most likely occurred in the common ancestor of all

surveyed *Leptodactylus* species (Fig. 1A, Table S2). Two other ancient paralogs common to

vertebrates, ATP1A2 and ATP1A3, appear to be present as single-copy genes and lack any

known CG-resistant substitutions in *Leptodactylus* (Fig. S3).

The αM1–2 transmembrane domains of the ATP1A1 paralog in *L. latrans* are distinguished by

four amino acid substitutions (*11*) (Fig 1C, Fig. S4). Two of these substitutions, Q111R and

N122D, were first identified in rat ATP1A1 and have been shown to interact synergistically to

confer CG-resistance to the CG-sensitive sheep ATP1A1 protein *in vitro* (*26*, *27*). By comparing

ATP1A1 sequences in five distantly related *Leptodactylus* species (*28*), we find that all five

harbor both a putatively resistant paralog (R) that includes the Q111R and N122D substitutions

and a putatively sensitive ATP1A1 paralog (S) that lacks these substitutions. In addition to

Q111R and N122D, 10 other amino acid substitutions distinguish the R and S paralogs in most

of the five sampled species (Fig. 1C). We refer to these twelve substitutions as "R/S

distinguishing" substitutions henceforth.

104
To elucidate the history of gene duplication relative to speciation events, we estimated

phylogenies from alignments of ATP1A1 protein-coding nucleotide sequences. Phylogenies

estimated from synonymous and inferred protein sequences support strikingly different

topologies (Fig 1B and C). If we were to assume that R evolve independently within each

species, the phylogeny estimated from nucleotide sequences (primarily based on variation at

synonymous sites) would imply independent duplications in each of the *Leptodactylus* species,

followed by parallel substitutions at the same 12 R/S distinguishing amino acid positions (Fig.

1B). Instead, we propose that this genealogical pattern reflects a single ancestral duplication - as

estimated from inferred amino acid sequences (Fig. 1C, Table S4) - and a history of recurrent

NAGC between paralogs. Frequent NAGC produces a pattern of "concerted evolution" whereby

tandemly linked paralogs from the same species are more similar to one another than they are to

their orthologous counterparts in other species (Fig. 2A). We propose that the persistence of the

12 amino acid differences between the two paralogs is due to selection counteracting the

homogenizing effects of NAGC (*23, 29*) (Fig. 2B), thereby maintaining an adaptive division-of-

labor between toxin resistance and cellular physiology.

The opposing forces of NAGC and selection are predicted to leave a characteristic genealogical

signature at neutral sites closely linked to the targets of selection (*29*) (Fig. 2B). We thus tested

the relationship between the genealogical signature and distance from nonsynonymous variants

putatively under selection. To do this, we classified all informative variant sites as supporting

either an ancient duplication of ATP1A1 that predated all speciation events versus an alternative

in which within-species paralogs are genealogical neighbors (defining support as a difference in

two log likelihoods for these alternative genealogical models). This analysis reveals that

synonymous (presumed to be neutral) variants congruent with an ancient duplication of R and S

have a median distance of 4 bp from nonsynonymous variants exhibiting the same pattern (Fig.

2C). In contrast, randomly sampled synonymous sites supporting the alternative genealogy have

a median distance of 88 bp from those nonsynonymous variants (bootstrap $p<10^{-5}$). This pattern

at synonymous sites is consistent with selection to maintain key amino acid substitutions

distinguishing neofunctionalized gene duplicates in the face of NAGC.

Similar qualitative patterns have been observed in the Amylase gene family in Drosophila (*30*),

the RHCE/RHD antigen genes of humans (*31*), and the red/green opsin genes of primates (*29*),

but the trans-specific sharing of R/S distinguishing substitutions here allows us to quantify the

intensity of these opposing evolutionary forces. We quantified the strength of selection required

at R/S distinguishing substitutions to counteract NAGC between R and S paralogs. We first

considered population genetics theory for the evolution of a single site in tandem duplicates (*32*)

(Supplementary Materials). This analytic model predicts that if, for example, gene conversion is

an order of magnitude more frequent than point mutation, then the maintenance of diverged,

distinct alleles is only likely under strong purifying selection (namely, a population size scaled

selection coefficient, $2Ns$, larger than one) (Fig. 3A). We next developed an inference method

based on simulations of ATP1A1 evolution to estimate the parameters that, jointly, best explain

divergence patterns throughout the gene, including both high and low levels of paralog

divergence observed near and far from the 12 R/S distinguishing substitutions, respectively

(Supplementary Materials). We estimate the rate of NAGC to be an order of magnitude higher

than the point mutation rate (posterior mode 9 with 80% credible interval 4-54 times the point

mutation rate), and $2Ns$ substantially larger than one (posterior mode 9; 80% credible interval 5-

18; Fig. 3B). These estimates fall within the plausible range predicted by the theoretical single-

site model of Fig. 3A. Thus, our evolutionary inference supports the joint, reciprocal action of

frequent NAGC and strong purifying selection at R/S distinguishing substitutions to generate the

observed divergence patterns in ATP1A1.

The inference that selection maintains the co-occurrence of these 12 R/S distinguishing

substitutions on the R copy implies they are functionally important and collectively contribute to

organismal fitness. The functional importance for Q111R and N122D in the context of CG-

insensitivity has previously been demonstrated by *in vitro* enzyme inhibition assays (*9*).

Additionally, the potential importance of substitutions at sites 112 and 116 has been suggested

by molecular evolution analysis and structural studies, respectively (*11*, *33*). However, the

remaining eight R/S distinguishing substitutions are scattered among regions of the protein that

have not been implicated in CG-resistance. Since our analysis suggests that amino acid

divergence between R and S paralogs is maintained by selection, we performed protein-

engineering experiments to elucidate the functional significance of the 12 R/S distinguishing

substitutions. We synthesized and recombinantly expressed eight mutant Na^+,K^+-ATPase

proteins, each harboring different combinations of R-specific replacements on both S- and R-

type genetic backgrounds of the representative species, *L. latrans* (Fig. 4A, Table S6) and we

then quantified the level of CG-resistance of each genotype using enzyme-inhibition assays

(Table S7, Fig. S5, (*34*)). Individually, Q111R and N122D significantly increased CG-resistance

by 21-fold and 14-fold, respectively (ANOVA $p=2.7e-13$ and $p=2.3e-6$; Fig. 4B, Table S8).

When combined, Q111R and N122D produce a >100-fold increase in CG resistance relative to

the S-copy (Tukey's HSD test, adjusted $p<4e-5$, Fig. 4B, Tables S6-S7). In contrast, the

remaining 10 substitutions had no detectable effect on CG-resistance when added to the S

background ($p=0.22$, Fig. 4B).

Given the absence of detectable effects of R/S distinguishing substitutions other than Q111R and

N122D on CG-resistance, we asked whether these substitutions had effects on other aspects of

ATP1A1 function that relate to organismal fitness. Since ATP hydrolysis and ion co-transport

are strongly coupled functions of Na^+,K^+-ATPase (*35*), we used estimates of the rate of ATP

hydrolysis in the absence of ouabain as a proxy for overall protein activity. Based on this assay,

we found that CG-resistance substitutions Q111R and N122D significantly impair activity,

individually reducing ATPase activity by an average of 40% (*p*=0.024 and *p*=7.7e-4

respectively; Fig. 4B; Tables S8). We also detected a significant interaction between Q111R and

N122D that renders their joint effects somewhat less severe than predicted by the sum of their

individual effects (*i.e.*, a 30% reduction rather than the expected 78% reduction, $p=0.022$).

Additionally, adding the remaining 10 R-specific substitutions on the S background containing

Q111R and N122D restores normal ATPase activity close to wild-type levels. An analysis of

variance reveals a highly significant effect of these 10 R-specific substitutions (p=1.2×10^{-4}, Fig.

4B, Table S8). These results indicate that these 10 R/S distinguishing substitutions play a vital

role in compensating for the negative pleiotropic effects of the resistance-conferring

substitutions, Q111R and N122D. We conclude that the evolution of the R protein from a CG-

sensitive ancestral state involved two epistatically interacting substitutions (Q111R and N122D)

in conjunction with compensatory effects of 10 additional substitutions that mitigate the trade-off

between toxin resistance and native enzyme activity.

Apart from the 12 R/S distinguishing substitutions shared by most *Leptodactylus* species, an

additional nine substitutions and a two-amino acid deletion distinguish the S and R paralogs in

the focal species, *L. latrans*. Our experiments revealed that these 10 species-specific

substitutions do not contribute to S vs. R differences in CG resistance or enzyme function (using

all 10 as one co-variate, ANOVA *p*>0.5).

The functional significance of protein divergence between species is a fundamental question in

biology and it is often difficult to assess in the absence of *a priori* functional information. The R-

and S-type ATP1A1 paralogs of *L. latrans* are distinguished by 22 amino acid substitutions

(including one deletion), only two of which are associated with protein function (in this case CG-

resistance). Our evolutionary analyses allowed us to focus experiments on just 12 of these 22

substitutions that we inferred were likely to be important for function. We found that an adaptive

division-of-labor between the R and S paralogs of ATP1A1 in *Leptodactylus* has been

maintained by strong selection that has counteracted the homogenizing effects of frequent

NAGC over the 35 million-year history of this genus. The possession of functionally distinct R-

and S-type ATP1A1 paralogs ensures that *Leptodactylus* frogs are resistant to the CG toxins

produced by their prey without compromising native ATPase function. Our work reveals the

mechanisms by which selection surmounts intragenic constraints on adaptation and demonstrates

how intramolecular epistasis and pleiotropy can shape adaptation.

References and Notes:

1. E. D. Brodie, Toxins and venoms. *Curr. Biol.* **19**, R931–R935 (2009).

2. K. Chen, A. L. CHEN, Notes on the poisonous secretions of twelve species of toads. *J. Pharmacol. Exp. Ther.* **47**, 281–293 (1933).

3. W. R. Heyer, R. W. McDiarmid, D. L. Weigmann, Tadpoles, predation and pond habitats in the tropics. *Biotropica*, 100–111 (1975).

4. M. R. Crossland, C. Azevedo-Ramos, Effects of Bufo (Anura: Bufonidae) toxins on tadpoles from native and exotic Bufo habitats. *Herpetologica*, 192–199 (1999).

5. C. Azevedo-Ramos, W. E. Magnusson, Tropical tadpole vulnerability to predation: association between laboratory results and prey distribution in an Amazonian savanna. *Copeia*, 58–67 (1999).

6. . D. Guimaraes, R. M. Pinto, R. F. Juliano, Bufo granulosus (NCN). Predation. *Herpetol. Rev.* **35**, 259 (2004).

7. L. Krenn, B. Kopp, Bufadienolides from animal and plant sources. *Phytochemistry*. **48**, 1–29 (1998).

8. J.-D. Horisberger, Recent insights into the structure and mechanism of the sodium pump. *Physiology*. **19**, 377–387 (2004).

9. J. B. Lingrel, The physiological significance of the cardiotonic steroid/ouabain-binding site of the Na, K-ATPase. *Annu. Rev. Physiol.* **72**, 395–412 (2010).

10. M. Laursen, J. L. Gregersen, L. Yatime, P. Nissen, N. U. Fedosova, Structures and characterization of digoxin-and bufalin-bound Na+, K+-ATPase compared with the ouabain-bound complex. *Proc. Natl. Acad. Sci.* **112**, 1755–1760 (2015).

11. D. J. Moore, D. C. Halliday, D. M. Rowell, A. J. Robinson, J. S. Keogh, Positive Darwinian selection results in resistance to cardioactive toxins in true toads (Anura: Bufonidae). *Biol. Lett.* **5**, 513–516 (2009).

12. B. Ujvari, H. Mun, A. D. Conigrave, A. Bray, J. Osterkamp, P. Halling, T. Madsen, Isolation breeds naivety: island living robs Australian varanid lizards of toad-toxin immunity via four-base-pair mutation. *Evol. Int. J. Org. Evol.* **67**, 289–294 (2013).

13. B. Ujvari, N. R. Casewell, K. Sunagar, K. Arbuckle, W. Wüster, N. Lo, D. O'Meally, C. Beckmann, G. F. King, E. Deplazes, Widespread convergence in toxin resistance by predictable molecular evolution. *Proc. Natl. Acad. Sci.* **112**, 11911–11916 (2015).

14. S. Mohammadi, Z. Gompert, J. Gonzalez, H. Takeuchi, A. Mori, A. H. Savitzky, Toxin-resistant isoforms of Na+/K+-ATPase in snakes do not closely track dietary specialization on toads. *Proc. R. Soc. B Biol. Sci.* **283**, 20162111 (2016).

15. J. Orlowski, J. B. Lingrel, Tissue-specific and developmental regulation of rat Na, K-
ATPase catalytic alpha isoform and beta subunit mRNAs. *J. Biol. Chem.* **263**, 10436–10442
(1988).

16. L. Fagerberg, B. M. Hallström, P. Oksvold, C. Kampf, D. Djureinovic, J. Odeberg, M.
Habuka, S. Tahmasebpoor, A. Danielsson, K. Edlund, Analysis of the human tissue-specific
expression by genome-wide integration of transcriptomics and antibody-based proteomics.
Mol. Cell. Proteomics. **13**, 397–406 (2014).

17. S. Mohammadi, A. H. Savitzky, J. Lohr, S. Dobler, Toad toxin-resistant snake (Thamnophis
elegans) expresses high levels of mutant Na+/K+-ATPase mRNA in cardiac muscle. *Gene*.
614, 21–25 (2017).

18. B. M. Marshall, N. R. Casewell, M. Vences, F. Glaw, F. Andreone, A. Rakotoarison, G.
Zancolli, F. Woog, W. Wüster, Widespread vulnerability of Malagasy predators to the
toxins of an introduced toad. *Curr. Biol.* **28**, R654–R655 (2018).

19. Y. Zhen, M. L. Aardema, E. M. Medina, M. Schumer, P. Andolfatto, Parallel molecular
evolution in an herbivore community. *science*. **337**, 1634–1637 (2012).

20. G. Petschenka, V. Wagschal, M. von Tschirnhaus, A. Donath, S. Dobler, Convergently
evolved toxic secondary metabolites in plants drive the parallel molecular evolution of
insect resistance. *Am. Nat.* **190**, S29–S43 (2017).

21. J. N. Lohr, F. Meinzer, S. Dalla, R. Romey-Glüsing, S. Dobler, The function and
evolutionary significance of a triplicated Na, K-ATPase gene in a toxin-specialized insect.
BMC Evol. Biol. **17**, 1–10 (2017).

22. L. Yang, N. Ravikanthachari, R. Mariño-Pérez, R. Deshmukh, M. Wu, A. Rosenstein, K.
Kunte, H. Song, P. Andolfatto, Predictability in the evolution of Orthopteran cardenolide
insensitivity. *Philos. Trans. R. Soc. B*. **374**, 20180246 (2019).

23. J. B. Walsh, Sequence-dependent gene conversion: can duplicated genes diverge fast
enough to escape conversion? *Genetics*. **117**, 543–557 (1987).

24. J.-M. Chen, D. N. Cooper, N. Chuzhanova, C. Férec, G. P. Patrinos, Gene conversion:
mechanisms, evolution and human disease. *Nat. Rev. Genet.* **8**, 762–775 (2007).

25. H. Innan, F. Kondrashov, The evolution of gene duplications: classifying and distinguishing
between models. *Nat. Rev. Genet.* **11**, 97–108 (2010).

26. E. M. Price, J. B. Lingrel, Structure-function relationships in the sodium-potassium
ATPase. alpha. subunit: site-directed mutagenesis of glutamine-111 to arginine and
asparagine-122 to aspartic acid generates a ouabain-resistant enzyme. *Biochemistry*. **27**,
8400–8408 (1988).

27. E. Price, D. Rice, J. Lingrel, Structure-function studies of Na, K-ATPase. Site-directed
mutagenesis of the border residues from the H1-H2 extracellular domain of the alpha
subunit. *J. Biol. Chem.* **265**, 6638–6641 (1990).

28. R. O. de Sá, T. Grant, A. Camargo, W. R. Heyer, M. L. Ponssa, E. Stanley, Systematics of
the neotropical genus Leptodactylus Fitzinger, 1826 (Anura: Leptodactylidae): phylogeny,
the relevance of non-molecular evidence, and species accounts. *South Am. J. Herpetol.* **9**
(2014).

29. K. M. Teshima, H. Innan, Neofunctionalization of duplicated genes under the pressure of
gene conversion. *Genetics*. **178**, 1385–1398 (2008).

30. H. Shibata, T. Yamazaki, Molecular evolution of the duplicated Amy locus in the
Drosophila melanogaster species subgroup: concerted evolution only in the coding region
and an excess of nonsynonymous substitutions in speciation. *Genetics*. **141**, 223–236
(1995).

31. H. Innan, A two-locus gene conversion model with selection and its application to the
human RHCE and RHD genes. *Proc. Natl. Acad. Sci.* **100**, 8793–8798 (2003).

32. J. A. Fawcett, H. Innan, Neutral and non-neutral evolution of duplicated genes with gene
conversion. *Genes*. **2**, 191–209 (2011).

33. H. Ogawa, T. Shinoda, F. Cornelius, C. Toyoshima, Crystal structure of the sodium-
potassium pump (Na+, K+-ATPase) with bound potassium and ouabain. *Proc. Natl. Acad.*
Sci. **106**, 13742–13747 (2009).

34. S. Dalla, M. Baum, S. Dobler, Substitutions in the cardenolide binding site and interaction
of subunits affect kinetics besides cardenolide sensitivity of insect Na, K-ATPase. *Insect*
Biochem. Mol. Biol. **89**, 43–50 (2017).

35. G. G. Hammes, Unifying concept for the coupling between ion pumping and ATP
hydrolysis or synthesis. *Proc. Natl. Acad. Sci.* **79**, 6881–6884 (1982).

36. Y.-J. Feng, D. C. Blackburn, D. Liang, D. M. Hillis, D. B. Wake, D. C. Cannatella, P.
Zhang, Phylogenomics reveals rapid, simultaneous diversification of three major clades of
Gondwanan frogs at the Cretaceous–Paleogene boundary. *Proc. Natl. Acad. Sci.* **114**,
E5864–E5870 (2017).

37. A. Harpak, X. Lan, Z. Gao, J. K. Pritchard, Frequent nonallelic gene conversion on the
human lineage and its effect on the divergence of gene duplicates. *Proc. Natl. Acad. Sci.*
114, 12779–12784 (2017).

38. B. J. Haas, A. Papanicolaou, M. Yassour, M. Grabherr, P. D. Blood, J. Bowden, M. B.
Couger, D. Eccles, B. Li, M. Lieber, De novo transcript sequence reconstruction from
RNA-seq using the Trinity platform for reference generation and analysis. *Nat. Protoc.* **8**,
1494 (2013).

39. D. R. Zerbino, E. Birney, Velvet: algorithms for de novo short read assembly using de
Bruijn graphs. *Genome Res.* **18**, 821–829 (2008).

40. M. H. Schulz, D. R. Zerbino, M. Vingron, E. Birney, Oases: robust de novo RNA-seq
assembly across the dynamic range of expression levels. *Bioinformatics*. **28**, 1086–1092
(2012).

41. G. W. Vurture, F. J. Sedlazeck, M. Nattestad, C. J. Underwood, H. Fang, J. Gurtowski, M.
C. Schatz, GenomeScope: fast reference-free genome profiling from short reads.
Bioinformatics. **33**, 2202–2204 (2017).

42. N. I. Weisenfeld, V. Kumar, P. Shah, D. M. Church, D. B. Jaffe, Direct determination of
diploid genome sequences. *Genome Res.* **27**, 757–767 (2017).

43. M. Seppey, M. Manni, E. M. Zdobnov, in *Gene Prediction* (Springer, 2019), pp. 227–245.

44. S. M. Kiełbasa, R. Wan, K. Sato, P. Horton, M. C. Frith, Adaptive seeds tame genomic
sequence comparison. *Genome Res.* **21**, 487–493 (2011).

45. H. Li, seqtk Toolkit for processing sequences in FASTA/Q formats. *GitHub*. **767**, 69
(2012).

46. S. Koren, B. P. Walenz, K. Berlin, J. R. Miller, N. H. Bergman, A. M. Phillippy, Canu:
scalable and accurate long-read assembly via adaptive k-mer weighting and repeat
separation. *Genome Res.* **27**, 722–736 (2017).

47. H. Li, Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*. **34**, 3094–
3100 (2018).

48. R. Vaser, I. Sović, N. Nagarajan, M. Šikić, Fast and accurate de novo genome assembly
from long uncorrected reads. *Genome Res.* **27**, 737–746 (2017).

49. R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).

50. M. Gouy, S. Guindon, O. Gascuel, SeaView version 4: a multiplatform graphical user
interface for sequence alignment and phylogenetic tree building. *Mol. Biol. Evol.* **27**, 221–
224 (2010).

51. B. Subramanian, S. Gao, M. J. Lercher, S. Hu, W.-H. Chen, Evolview v3: a webserver for
visualization, annotation, and management of phylogenetic trees. *Nucleic Acids Res.* **47**,
W270–W275 (2019).

52. M. Stanke, A. Tzvetkova, B. Morgenstern, AUGUSTUS at EGASP: using EST, protein and
genomic alignments for improved gene prediction in the human genome. *Genome Biol.* **7**,
1–8 (2006).

53. B. Q. Minh, H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. von
Haeseler, R. Lanfear, IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

54. Z. Yang, PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**,
1586–1591 (2007).

55. S. Kumar, G. Stecher, M. Li, C. Knyaz, K. Tamura, MEGA X: molecular evolutionary
genetics analysis across computing platforms. *Mol. Biol. Evol.* **35**, 1547–1549 (2018).

56. M. Kimura, On the probability of fixation of mutant genes in a population. *Genetics*. **47**,
713 (1962).

57. J. H. Gillespie, *Population genetics: a concise guide* (JHU Press, 2004).

58. Y.-B. Sun, Z.-J. Xiong, X.-Y. Xiang, S.-P. Liu, W.-W. Zhou, X.-L. Tu, L. Zhong, L. Wang,
D.-D. Wu, B.-L. Zhang, Whole-genome sequence of the Tibetan frog Nanorana parkeri and
the comparative evolution of tetrapod genomes. *Proc. Natl. Acad. Sci.* **112**, E1257–E1262
(2015).

59. A. J. Crawford, Relative rates of nucleotide substitution in frogs. *J. Mol. Evol.* **57**, 636–641
(2003).

60. S. P. Mansai, H. Innan, The power of the methods for detecting interlocus gene conversion.
Genetics. **184**, 517–527 (2010).

61. D. Feng, L. Tierney, Computing and displaying isosurfaces in R. *J. Stat. Softw.* **28**, 1–24
(2008).

62. B. Venables WNRipley, Modern applied statistics with S. *Stat. Comput. N. Y. Springer*
(2002).

63. V. A. Luckow, S. Lee, G. Barry, P. Olins, Efficient generation of infectious recombinant
baculoviruses by site-specific transposon-mediated insertion of foreign genes into a
baculovirus genome propagated in Escherichia coli. *J. Virol.* **67**, 4566–4579 (1993).

64. G. Petschenka, S. Fandrich, N. Sander, V. Wagschal, M. Boppré, S. Dobler, Stepwise
evolution of resistance to toxic cardenolides via genetic substitutions in the Na+/K+-
ATPase of milkweed butterflies (Lepidoptera: Danaini). *Evolution*. **67**, 2753–2761 (2013).

65. H. H. Taussky, E. Shorr, A microcolorimetric method for the determination of inorganic
phosphorus. *J. Biol. Chem.* **202**, 675–685 (1953).

Acknowledgments: We thank C. Natarajan, K. Rohlfing, V. Wagschal, and P. Kowalski for
assistance in the laboratory. **Funding:** This study was funded by grants to PA from the National
Institutes of Health (R01- GM115523) and to JFS from the National Institutes of Health (R01-
HL087216) and the National Science Foundation (OIA-1736249), to SD from Deutsche
Forschungsgemeinschaft (DFG grant DO527/10-1), and by a fellowship from to AH by The
Simons Foundation's Society of Fellows (#633313). **Author contributions:** PA and AJC
conceived of and oversaw the project; LY, MPRO, SHA, JP, and AJC collected samples and
generated data; LY, AH, and KZ performed evolutionary and population genetics analyses; SM,
JFS, SD, AJC and PA designed functional experiments; SM and PA performed experiments and
statistical analyses; SM, JFS, LY, AH and PA wrote the paper; All authors edited the manuscript;
Competing interests: None. **Data and materials availability:** BioProject PRJNA627222, SRA:
SRR11583961-91, nucleotides: MT396181-92, genes: MT422192-MT422203, genome: TBD.

Supplementary Materials:

Materials and Methods

Figures S1-S6

Tables S1-S8

References 38-65

405 **Fig. 1. Molecular evolution of ATP1A1 in anurans. (**A) A maximum likelihood phylogeny of

406 anuran species and mammalian outgroups derived from (*36*). Species names in purple

407 correspond to chemically defended toads, and blue and red colors correspond to the S and R

408 ATP1A1 paralogs in *Leptodactylus* species, respectively. Only variant sites with roles in the CG-

409 binding or sensitivity are shown (reviewed in (*22*)). The numbering of sites is based on sheep

410 ATP1A1 (*Ovis aries*, Genbank: NC019458.2) and appears at the top of the table (e.g. the first

411 position showed is 108). Dots indicate identity with the consensus sequence of anuran species

412 and letters represent amino acid substitutions relative to the reference. The images on the left are

413 the five surveyed *Leptodactylus* species and a representative prey toad (*Rhinella marina*). (B) A

414 maximum likelihood phylogeny estimated with nucleotide sequences with bootstrap support

415 values at nodes. (C) A maximum likelihood genealogy based on protein sequences with

416 bootstrap support values at nodes. To the right is the pattern of amino acid variation at 12

417 positions that most strongly distinguish the S and R paralogs.

A. NAGC homogenizes paralog divergence

B. NAGC changes local genealogy

C. Inferred genealogy along ATP1A1

Fig. 2. (A) Non-allelic gene conversion (NAGC) homogenizes sequence variation between

paralogous genes, and therefore changes the genealogical signal. (This illustration is adapted

from Harpak et al. (*37*)). **(B)** Frequent NAGC can result in a genealogy in which paralogous

genes in the same species appear to share a more recent common ancestor with one another than

with their orthologous counterparts in other species. In principle, the homogenizing effects of

NAGC can be counteracted by selection that favors the differentiation of paralogous genes. **(C)**

Site-by-site difference in the likelihood of two alternative tree topologies—generalizing the grey

and purple extremes of panel (B) to five *Leptodactylus* species. Shaded regions show a log

likelihood difference greater than 2 in support of the corresponding model. Only parsimony-

informative variants in the ATP1A1 coding sequence are shown. Red bars correspond to the 12

R/S distinguishing nonsynonymous substitutions (shown in red in Fig 1C; Table S4).

Fig. 3. (A) Theoretical probability of maintaining distinct alleles at a single site in the face

of non-allelic gene conversion (NAGC). We used a theoretical model to compute the

probability of maintaining alternative alleles at the same site in a pair of paralogous genes, given

an NAGC rate and strength of selection against allele homogenization at the site. The black dot

shows the approximate mode estimate from panel B, which falls in the range in which

maintenance is likely according to this theoretical model. **(B) Estimates of evolutionary**

parameters. Approximate posterior probabilities were inferred based on simulations of the

evolution of ATP1A1 genes in *Leptodactylus*. The x-axis shows the NAGC rate across the gene,

and the y-axis shows the population selection coefficient for the 12 substitutions that distinguish

the R and S paralogs across species.

Fig. 4. Functional analysis of R-copy-specific substitutions. (A) ATP1A1 gene constructs

designed to produce proteins with various combinations of the 12 variants distinguishing the S

and R paralogs. Black circles indicate an amino acid matching the R copy whereas a white circle

indicates a match with the S copy. A dark grey shaded background indicates that the construct

was produced on the R genetic background and a white background indicates that the construct

was produced on the S background. Light grey columns highlight two variants (Q111R and

N122D) previously known to confer CG-resistance. (B) The joint functional properties of

engineered Na$^+$,K$^+$-ATPases. Mean ± SEM Log IC$_{50}$ (i.e., measurement of CG resistance) of the

six biological replicates is plotted on the x-axis and mean ± SEM ATP hydrolysis rate (i.e.,

measurement of protein activity) for the same proteins is plotted on the y-axis. R and S copies

differ by the 12 amino acid substitutions being studied and by 9 additional amino acid

substitutions and one two amino acid insertion-deletion difference (see Table S6).