1

Title: Concerted evolution reveals co-adapted amino acid substitutions in Na$^+$K$^+$-ATPase of frogs that prey on toxic toads

Authors: Shabnam Mohammadi[1]*, Lu Yang[2]*, Arbel Harpak[3]*, Santiago Herrera-Álvarez[4, ψ], María del Pilar Rodríguez-Ordoñez[4, #], Julie Peng, Karen Zhang[2], Jay F. Storz[1], Susanne Dobler[6], Andrew J. Crawford[4]** and Peter Andolfatto[3]**

Affiliations:

[1] School of Biological Sciences, University of Nebraska, Lincoln, NE, USA

[2] Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, USA

[3] Department of Biological Sciences, Columbia University, New York, NY, USA

[4] Department of Biological Sciences, Universidad de los Andes, Bogotá, 111711, Colombia

[5] Lewis-Sigler Institute, Princeton University, Princeton, NJ, USA

[6] Molecular Evolutionary Biology, Zoological Institute, Universität Hamburg, Hamburg, Germany

*Co-first authorship

**Correspondence to: andrew@dna.ac, pa2543@columbia.edu

ψ Current address: Dept. of Ecology and Evolution, University of Chicago, Chicago, IL, USA

Current address: Université Paris-Saclay Evry, Evry, France

ABSTRACT (141 words):

Gene duplication is an important source of evolutionary innovation, but the adaptive division-of-

labor between duplicates can be opposed by ongoing gene conversion between them. Here we

document a tandem duplication of Na^+,K^+-ATPase subunit α1 (ATP1A1) shared by frogs in the

genus *Leptodactylus*, a group of species that feeds on toxic toads. One ATP1A1 paralog evolved

resistance to toad toxins while the other paralog retained ancestral susceptibility. We show that

the two *Leptodactylus* paralogs are distinguished by 12 amino acid substitutions that were

maintained by strong selection that counteracted the homogenizing effect of gene conversion.

Protein-engineering experiments show that two major-effect substitutions confer toxin resistance,

whereas the 10 additional substitutions mitigate deleterious pleiotropic effects on enzyme

function. Our results highlight how trans-specific, neofunctionalized gene duplicates can provide

unique insights into interactions between adaptive substitutions and the genetic backgrounds on

which they arise.

One Sentence Summary: Selection counteracts gene conversion to maintain an adaptive

division-of-labor between tandemly duplicated genes.

Main Text:

The repeated evolution of toxin resistance in animals is one of the clearest examples of natural

selection at the molecular level and represents a useful paradigm to examine constraints on the

evolution of novel protein functions (*1*). Neotropical Grass Frogs of the genus *Leptodactylus*

(Leptodactylidae) are widely distributed throughout lowland South America and are known to

feed on chemically-defended toads – a predatory tendency that is rare among frogs (*2–6*). A

major component of the chemical defense secretions of toads is a class of cardiac glycosides

(CGs) called "bufadienolides" (*7*) that inhibit the α-subunit of Na^+,K^+-ATPases (ATP1A).

Na^+,K^+-ATPases are transmembrane proteins that are vital to numerous physiological processes

in animals including neural signal transduction, muscle contraction, and cell homeostasis (*8, 9*).

CGs bind to the extracellular surface of ATP1A and block the flux of ions (*10*), making them

potent poisons to most animals. However, some vertebrates have independently evolved the

ability to prey on chemically-defended toads, partly via amino acid substitutions to the CG-

binding domain of ATP1A1 that confer resistance to CGs (*11–14*).

Most vertebrates share several paralogous copies of ATP1A that have different tissue-specific

expression profiles (*15*). For example, ATP1A1 is the most ubiquitously expressed paralog and

ATP1A3 has enriched expression in nervous tissue and heart muscle (*16, 17*; Fig. S1). Previous

studies on the molecular convergence of CG-resistance in reptiles have focused primarily on the

αM1–2 extracellular loop of ATP1A3 (*12–14, 18*), whereas studies of birds, mammals, and

amphibians have focused on the same region of ATP1A1 (*11, 18*). A survey of ATP1A1 αM1–2

in toads and frogs (*11*) revealed a possible duplication of this gene in the toad-eating frog,

Leptodactylus latrans (reported as *L. ocellatus*), where the resistant (R) paralog includes

substitutions known to confer resistance to CGs while the sensitive (S) paralog appears to have

retained the ancestral susceptibility to CGs. Neofunctionalization of ATP1A paralogs has

contributed to the evolution of CG-resistance in numerous insect lineages (*19–22*) but appear to

be rare among CG-resistant vertebrates. Further, the fate of duplicated genes and the probability

that they will neofunctionalize is predicted to depend on the strength of selection for functional

differentiation relative to the rate of non-allelic gene conversion (NAGC), a form of

nonreciprocal genetic exchange that homogenizes sequence variation between duplicated genes,

thereby impeding divergence (*23–25*). The ATP1A1 duplication in *Leptodactylus* provides an

ideal opportunity to explore this process because the functional differentiation between R and S

paralogs has clear adaptive significance with regard to CG-resistance.

We surveyed the full-length coding sequences of all ATP1A paralogs in *Leptodactylus* and other

anurans using RNA-seq-based gene discovery (*19*; Tables S1). Our results confirm that ATP1A1

is duplicated in *L. latrans* (*11*) and indicate that the duplication of ATP1A1 most likely occurred

in the common ancestor of all surveyed *Leptodactylus* species (Fig. 1A, Table S2). Two other

ancient paralogs common to vertebrates, ATP1A2 and ATP1A3, appear to be present as single-

copy genes and lack any known CG-resistant substitutions in *Leptodactylus* (Fig. S2). The αM1–

2 transmembrane domains of the ATP1A1 paralogs in *L. latrans* are distinguished by four amino

acid substitutions (*11*; Fig 1C, Fig. S3). Two of these substitutions, Q111R and N122D, were

first identified in rat ATP1A1 and have been shown to interact synergistically to confer CG-

resistance to sheep ATP1A1 protein *in vitro* (*26*, *27*). Comparison of ATP1A1 sequences in five

distantly related *Leptodactylus* species (*28*) reveals that they each harbor a putatively resistant

paralog (R) that includes the Q111R and N122D substitutions and a putatively sensitive ATP1A1

paralog (S) that lacks these substitutions. In addition to Q111R and N122D, there are 10 other

amino acid substitutions distinguishing the R and S paralogs in most of the five sampled species

(Fig. 1C). Hereafter, we refer to these twelve substitutions as "R/S distinguishing".

To infer when duplication occurred relative to speciation events, we estimated phylogenies from

an alignment of ATP1A1 coding sequence. Phylogenies estimated from nucleotide and inferred

amino-acid sequences support strikingly different topologies (Fig 1B and C). Without further

information, the genealogy based on nucleotide sequences (primarily based on variation at

synonymous sites) would imply independent duplications in each of the *Leptodactylus* species,

followed by parallel substitutions at the same 12 R/S distinguishing amino acid positions (Fig.

1B). Instead, it seems much more likely that this genealogical pattern reflects a single ancestral

duplication — as indicated by genealogy based on amino acid sequences (Fig. 1C, Table S4) —

coupled with on-going NAGC between the R and S paralogs of each species. Frequent NAGC

produces a pattern of "concerted evolution" whereby tandemly linked paralogs from the same

species are more similar to one another than they are to their orthologous counterparts in other

species (*29;* Fig. 2A,B). By generating a *de novo* genome assembly of *L. fuscus* based on single-

molecule sequencing, we established that S and R copies are indeed arranged in tandem and in

the same orientation, and are therefore likely to be subject to NAGC (Table S3, Fig. S4). We

thus propose that the persistence of the 12 amino acid differences between the two paralogs is

due to selection counteracting the homogenizing effects of NAGC (*23, 30*; Fig. 2B), thereby

maintaining an adaptive division-of-labor between the R and S copies.

The opposing forces of NAGC and selection are predicted to leave a characteristic genealogical

signature at neutral sites closely linked to the targets of selection (*30*; Fig. 2B). We therefore

tested the relationship between the genealogical signature and distance from nonsynonymous

variants putatively under selection. To this end, for all informative sites, we evaluated the level

of support for an ancient duplication of ATP1A1 in the common ancestor of all *Leptodactylus*

species (with no concerted evolution) relative to support for an alternative in which ATP1A1

paralogs within species are always more closely related to one another than they are to paralogs

in other species (as expected under concerted evolution). This analysis reveals that synonymous

(presumed to be neutral) variants congruent with an ancient duplication of R and S have a

median distance of 4 bp from nonsynonymous variants exhibiting the same pattern (Fig. 2C). In

contrast, randomly sampled synonymous sites supporting the alternative genealogy (i.e.

concerted evolution) have a median distance of 88 bp from those nonsynonymous variants

(bootstrap $p<10^{-5}$). This pattern at synonymous sites is consistent with a scenario in which

purifying selection maintains functionally important sequence differences between

neofunctionalized gene duplicates in the face of NAGC.

We next quantified the strength of purifying selection required to maintain the amino acid

differentiation between R and S duplicates in the face of NAGC. We first considered population

genetics theory for the evolution of a single site in tandem duplicates (*31*) (Supplementary

Materials). This analytic model predicts that if the rate of NAGC is an order of magnitude higher

than the rate of point mutation, then the maintenance of alternative amino acid states is only

likely under sufficiently strong purifying selection — namely, when the selection coefficient

scaled by population size, 2*Ns*, is larger than one (Fig. 3A). We next developed an inference

method based on simulations of ATP1A1 evolution to estimate the combination of parameters

that best explain divergence patterns throughout the gene, including levels of paralog divergence

observed as a function of distance from the 12 R/S distinguishing substitutions (Supplementary

Materials). We estimate the rate of NAGC to be an order of magnitude higher than the point

mutation rate (posterior mode 9 with 80% credible interval 4-54 times the point mutation rate),

and 2Ns substantially larger than one (posterior mode 9; 80% credible interval 5-18; Fig. 3B).

These estimates fall within the plausible range predicted by the theoretical single-site model (Fig.

3A). These results indicate that the observed pattern of divergence between R and S paralogs

reflects a history of strong purifying selection that maintains fixed differences between them

despite high rates of NAGC.

The inference that selection maintains the co-occurrence of the 12 R/S distinguishing

substitutions implies they are functionally important and collectively contribute to organismal

fitness. The effects of Q111R and N122D on CG-insensitivity have previously been

demonstrated by *in vitro* enzyme inhibition assays (*9*). Additionally, while not related directly to

CG-resistance, the potential importance of substitutions at sites 112 and 116 has been suggested

by molecular evolution analysis and structural studies, respectively (*11*, *32*). However, the

remaining eight R/S distinguishing substitutions are located in structural domains that have not

been implicated in CG-resistance. Since our analysis suggests that amino acid divergence

between R and S paralogs is maintained by selection, we performed protein-engineering

experiments to elucidate the functional significance of the 12 R/S distinguishing substitutions.

We synthesized and recombinantly expressed eight mutant Na$^+$,K$^+$-ATPase proteins, each

harboring different combinations of R-specific replacements on both S- and R-type genetic

backgrounds of a representative species, *L. latrans* (Fig. 4A, Table S6), and we then quantified

the level of CG-resistance of each genotype using enzyme-inhibition assays (Table S7, Fig. S7,

(*33*)). Individually, Q111R and N122D significantly increased CG-resistance by 21-fold and 14-

fold, respectively (ANOVA p=2.7e-13 and p=2.3e-6; Fig. 4B, Table S8). When combined,

Q111R and N122D produce a greater than 100-fold increase in CG resistance relative to the S

paralog (Tukey's HSD test, adjusted p<4e-5, Fig. 4B, Tables S6-S8). In contrast, the remaining

10 substitutions had no detectable net effect on CG-resistance when jointly added to the S

background (p=0.22, Fig. 4B).

Given the absence of detectable effects of R/S distinguishing substitutions other than Q111R and

N122D on CG-resistance, we tested whether these substitutions had effects on other aspects of

ATP1A1 function. Since ATP hydrolysis and ion co-transport are strongly coupled functions of

Na^+,K^+-ATPase (*34*), we used estimates of the rate of ATP hydrolysis in the absence of ouabain

as a proxy for overall protein activity. Based on this assay, we found that CG-resistance

substitutions Q111R and N122D significantly impair activity, individually reducing ATPase

activity by an average of 40% (*p*=0.024 and *p*=7.7e-4 respectively; Fig. 4B; Table S8). We also

detected a significant interaction between Q111R and N122D that renders their joint effects

somewhat less severe than predicted by the sum of their individual effects (*i.e.*, a 30% reduction

rather than the expected 78% reduction, p=0.022). Critically, adding the remaining 10 R-specific

substitutions on the S background containing Q111R and N122D restores normal ATPase

activity close to wild-type levels. An analysis of variance reveals a highly significant effect of

these 10 R-specific substitutions (p=1.2 × 10^{-4}, Fig. 4B, Table S8). Our results thus indicate that

these 10 R/S distinguishing substitutions play a vital role in compensating for the negative

pleiotropic effects of the resistance-conferring substitutions, Q111R and N122D. We conclude

that the evolution of the R protein from a CG-sensitive ancestral state involved two epistatically-

interacting substitutions (Q111R and N122D) in conjunction with compensatory effects of 10

additional substitutions that mitigate the trade-off between toxin resistance and native enzyme

activity.

The adaptive division-of-labor between the R and S paralogs of ATP1A1 in *Leptodactylus* has

been maintained by strong selection that has counteracted the homogenizing effects of frequent

NAGC over the 35 million-year history of this genus. Similar signatures of selection to maintain

sequence differentiation between neofunctionalized duplicates have been observed for the

RHCE/RHD antigen proteins of humans (*35*), "major facilitator family" transporter proteins in

Drosophila (*36*) and red/green opsins of primates (*30*). To our knowledge, only the case of

opsins has been linked directly to functional differentiation, notably two closely-linked amino

substitutions contributing to a red to green shift in absorbance maxima (*37*). Our study highlights

similar signatures of selection not only on the two amino acid substitutions directly linked to

adaptive differentiation for CG-resistance, but also at 10 more amino acid substitutions scattered

throughout the protein that facilitate this neofunctionalization. Thus, by identifying interactions

between adaptive substitutions and the genetic backgrounds that permit these changes, our

combination of evolutionary and functional analyses reveals how mechanisms of adaptation are

shaped by intramolecular epistasis and pleiotropy.

References and Notes:

1. E. D. Brodie, Toxins and venoms. *Curr. Biol.* **19**, R931–R935 (2009).

2. K. Chen, A. L. CHEN, Notes on the poisonous secretions of twelve species of toads. *J. Pharmacol. Exp. Ther.* **47**, 281–293 (1933).

3. W. R. Heyer, R. W. McDiarmid, D. L. Weigmann, Tadpoles, predation and pond habitats in the tropics. *Biotropica*, 100–111 (1975).

4. M. R. Crossland, C. Azevedo-Ramos, Effects of Bufo (Anura: Bufonidae) toxins on tadpoles from native and exotic Bufo habitats. *Herpetologica*, 192–199 (1999).

5. C. Azevedo-Ramos, W. E. Magnusson, Tropical tadpole vulnerability to predation: association between laboratory results and prey distribution in an Amazonian savanna. *Copeia*, 58–67 (1999).

6. . D. Guimaraes, R. M. Pinto, R. F. Juliano, Bufo granulosus (NCN). Predation. *Herpetol. Rev.* **35**, 259 (2004).

7. L. Krenn, B. Kopp, Bufadienolides from animal and plant sources. *Phytochemistry*. **48**, 1–29 (1998).

8. J.-D. Horisberger, Recent insights into the structure and mechanism of the sodium pump. *Physiology*. **19**, 377–387 (2004).

9. J. B. Lingrel, The physiological significance of the cardiotonic steroid/ouabain-binding site of the Na, K-ATPase. *Annu. Rev. Physiol.* **72**, 395–412 (2010).

10. M. Laursen, J. L. Gregersen, L. Yatime, P. Nissen, N. U. Fedosova, Structures and characterization of digoxin-and bufalin-bound Na+, K+-ATPase compared with the ouabain-bound complex. *Proc. Natl. Acad. Sci.* **112**, 1755–1760 (2015).

11. D. J. Moore, D. C. Halliday, D. M. Rowell, A. J. Robinson, J. S. Keogh, Positive Darwinian selection results in resistance to cardioactive toxins in true toads (Anura: Bufonidae). *Biol. Lett.* **5**, 513–516 (2009).

12. B. Ujvari, H. Mun, A. D. Conigrave, A. Bray, J. Osterkamp, P. Halling, T. Madsen, Isolation breeds naivety: island living robs Australian varanid lizards of toad-toxin immunity via four-base-pair mutation. *Evol. Int. J. Org. Evol.* **67**, 289–294 (2013).

13. B. Ujvari, N. R. Casewell, K. Sunagar, K. Arbuckle, W. Wüster, N. Lo, D. O'Meally, C. Beckmann, G. F. King, E. Deplazes, Widespread convergence in toxin resistance by predictable molecular evolution. *Proc. Natl. Acad. Sci.* **112**, 11911–11916 (2015).

14. S. Mohammadi, Z. Gompert, J. Gonzalez, H. Takeuchi, A. Mori, A. H. Savitzky, Toxin-resistant isoforms of Na+/K+-ATPase in snakes do not closely track dietary specialization on toads. *Proc. R. Soc. B Biol. Sci.* **283**, 20162111 (2016).

15.   J. Orlowski, J. B. Lingrel, Tissue-specific and developmental regulation of rat Na, K-
ATPase catalytic alpha isoform and beta subunit mRNAs. *J. Biol. Chem.* **263**, 10436–10442
(1988).

16.   L. Fagerberg, B. M. Hallström, P. Oksvold, C. Kampf, D. Djureinovic, J. Odeberg, M.
Habuka, S. Tahmasebpoor, A. Danielsson, K. Edlund, Analysis of the human tissue-specific
expression by genome-wide integration of transcriptomics and antibody-based proteomics.
Mol. Cell. Proteomics. **13**, 397–406 (2014).

17.   S. Mohammadi, A. H. Savitzky, J. Lohr, S. Dobler, Toad toxin-resistant snake (Thamnophis
elegans) expresses high levels of mutant Na+/K+-ATPase mRNA in cardiac muscle. *Gene.*
614, 21–25 (2017).

18.   B. M. Marshall, N. R. Casewell, M. Vences, F. Glaw, F. Andreone, A. Rakotoarison, G.
Zancolli, F. Woog, W. Wüster, Widespread vulnerability of Malagasy predators to the
toxins of an introduced toad. *Curr. Biol.* **28**, R654–R655 (2018).

19.   Y. Zhen, M. L. Aardema, E. M. Medina, M. Schumer, P. Andolfatto, Parallel molecular
evolution in an herbivore community. *Science.* **337**, 1634–1637 (2012).

20.   G. Petschenka, V. Wagschal, M. von Tschirnhaus, A. Donath, S. Dobler, Convergently
evolved toxic secondary metabolites in plants drive the parallel molecular evolution of
insect resistance. *Am. Nat.* **190**, S29–S43 (2017).

21.   J. N. Lohr, F. Meinzer, S. Dalla, R. Romey-Glüsing, S. Dobler, The function and
evolutionary significance of a triplicated Na, K-ATPase gene in a toxin-specialized insect.
BMC Evol. Biol. **17**, 1–10 (2017).

22.   L. Yang, N. Ravikanthachari, R. Mariño-Pérez, R. Deshmukh, M. Wu, A. Rosenstein, K.
Kunte, H. Song, P. Andolfatto, Predictability in the evolution of Orthopteran cardenolide
insensitivity. *Philos. Trans. R. Soc. B.* **374**, 20180246 (2019).

23.   J. B. Walsh, Sequence-dependent gene conversion: can duplicated genes diverge fast
enough to escape conversion? *Genetics.* **117**, 543–557 (1987).

24.   J.-M. Chen, D. N. Cooper, N. Chuzhanova, C. Férec, G. P. Patrinos, Gene conversion:
mechanisms, evolution and human disease. *Nat. Rev. Genet.* **8**, 762–775 (2007).

25.   H. Innan, F. Kondrashov, The evolution of gene duplications: classifying and distinguishing
between models. *Nat. Rev. Genet.* **11**, 97–108 (2010).

26.   E. M. Price, J. B. Lingrel, Structure-function relationships in the sodium-potassium
ATPase. alpha. subunit: site-directed mutagenesis of glutamine-111 to arginine and
asparagine-122 to aspartic acid generates a ouabain-resistant enzyme. *Biochemistry.* **27**,
8400–8408 (1988).

27.   E. Price, D. Rice, J. Lingrel, Structure-function studies of Na, K-ATPase. Site-directed
277   mutagenesis of the border residues from the H1-H2 extracellular domain of the alpha
278   subunit. *J. Biol. Chem.* **265**, 6638–6641 (1990).

279   28.   R. O. de Sá, T. Grant, A. Camargo, W. R. Heyer, M. L. Ponssa, E. Stanley, Systematics of
280   the neotropical genus *Leptodactylus* Fitzinger, 1826 (Anura: Leptodactylidae): phylogeny,
281   the relevance of non-molecular evidence, and species accounts. *South Am. J. Herpetol.* **9**
282   (2014).

283   29.   K. M. Teshima, H. Innan, The effect of gene conversion on the divergence between
284   duplicated genes. *Genetics*. **166**, 1553–1560 (2004).

285   30.   K. M. Teshima, H. Innan, Neofunctionalization of duplicated genes under the pressure of
286   gene conversion. *Genetics*. **178**, 1385–1398 (2008).

287   31.   J. A. Fawcett, H. Innan, Neutral and non-neutral evolution of duplicated genes with gene
288   conversion. *Genes*. **2**, 191–209 (2011).

289   32.   H. Ogawa, T. Shinoda, F. Cornelius, C. Toyoshima, Crystal structure of the sodium-
290   potassium pump (Na+, K+-ATPase) with bound potassium and ouabain. *Proc. Natl. Acad.*
291   *Sci.* **106**, 13742–13747 (2009).

292   33.   S. Dalla, M. Baum, S. Dobler, Substitutions in the cardenolide binding site and interaction
293   of subunits affect kinetics besides cardenolide sensitivity of insect Na, K-ATPase. *Insect*
294   *Biochem. Mol. Biol.* **89**, 43–50 (2017).

295   34.   G. G. Hammes, Unifying concept for the coupling between ion pumping and ATP
296   hydrolysis or synthesis. *Proc. Natl. Acad. Sci.* **79**, 6881–6884 (1982).

297   35.   H. Innan, A two-locus gene conversion model with selection and its application to the
298   human RHCE and RHD genes. *Proc. Natl. Acad. Sci.* **100**, 8793–8798 (2003).

299   36.   N. Osada, H. Innan, Duplication and gene conversion in the Drosophila melanogaster
300   genome. *PLoS Genet.* **4**, e1000305 (2008).

301   37.   S. Yokoyama, F. B. Radlwimmer, The molecular genetics and evolution of red and green
302   color vision in vertebrates. *Genetics*. **158**, 1697–1710 (2001).

303   38.   Y.-J. Feng, D. C. Blackburn, D. Liang, D. M. Hillis, D. B. Wake, D. C. Cannatella, P.
304   Zhang, Phylogenomics reveals rapid, simultaneous diversification of three major clades of
305   Gondwanan frogs at the Cretaceous–Paleogene boundary. *Proc. Natl. Acad. Sci.* **114**,
306   E5864–E5870 (2017).

307   39.   A. Harpak, X. Lan, Z. Gao, J. K. Pritchard, Frequent nonallelic gene conversion on the
308   human lineage and its effect on the divergence of gene duplicates. *Proc. Natl. Acad. Sci.*
309   **114**, 12779–12784 (2017).

40.   B. J. Haas, A. Papanicolaou, M. Yassour, M. Grabherr, P. D. Blood, J. Bowden, M. B.
Couger, D. Eccles, B. Li, M. Lieber, De novo transcript sequence reconstruction from
RNA-seq using the Trinity platform for reference generation and analysis. *Nat. Protoc.* **8**,
1494 (2013).

41.   D. R. Zerbino, E. Birney, Velvet: algorithms for de novo short read assembly using de
Bruijn graphs. *Genome Res.* **18**, 821–829 (2008).

42.   M. H. Schulz, D. R. Zerbino, M. Vingron, E. Birney, Oases: robust de novo RNA-seq
assembly across the dynamic range of expression levels. *Bioinformatics*. **28**, 1086–1092
(2012).

43.   G. W. Vurture, F. J. Sedlazeck, M. Nattestad, C. J. Underwood, H. Fang, J. Gurtowski, M.
C. Schatz, GenomeScope: fast reference-free genome profiling from short reads.
Bioinformatics. **33**, 2202–2204 (2017).

44.   N. I. Weisenfeld, V. Kumar, P. Shah, D. M. Church, D. B. Jaffe, Direct determination of
diploid genome sequences. *Genome Res.* **27**, 757–767 (2017).

45.   M. Seppey, M. Manni, E. M. Zdobnov, BUSCO: Assessing Genome Assembly and
Annotation Completeness in *Gene Prediction* (Springer, 2019), pp. 227–245.

46.   S. M. Kiełbasa, R. Wan, K. Sato, P. Horton, M. C. Frith, Adaptive seeds tame genomic
sequence comparison. *Genome Res.* **21**, 487–493 (2011).

47.   H. Li, seqtk Toolkit for processing sequences in FASTA/Q formats. *GitHub*. **767**, 69
(2012).

48.   S. Koren, B. P. Walenz, K. Berlin, J. R. Miller, N. H. Bergman, A. M. Phillippy, Canu:
scalable and accurate long-read assembly via adaptive k-mer weighting and repeat
separation. *Genome Res.* **27**, 722–736 (2017).

49.   H. Li, Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*. **34**, 3094–
3100 (2018).

50.   R. Vaser, I. Sović, N. Nagarajan, M. Šikić, Fast and accurate de novo genome assembly
from long uncorrected reads. *Genome Res.* **27**, 737–746 (2017).

51.   R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).

52.   M. Gouy, S. Guindon, O. Gascuel, SeaView version 4: a multiplatform graphical user
interface for sequence alignment and phylogenetic tree building. *Mol. Biol. Evol.* **27**, 221–
224 (2010).

53.   Z. He, H. Zhang, S. Gao, M. J. Lercher, W.-H. Chen, S. Hu, Evolview v2: an online
visualization and management tool for customized and annotated phylogenetic trees.
Nucleic Acids Res. **44**, W236–W241 (2016).

54.   M. Stanke, A. Tzvetkova, B. Morgenstern, AUGUSTUS at EGASP: using EST, protein and
346   genomic alignments for improved gene prediction in the human genome. *Genome Biol.* **7**,
347   1–8 (2006).

348   55.   B. Q. Minh, H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. von
349   Haeseler, R. Lanfear, IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
350   Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

351   56.   Z. Yang, PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**,
352   1586–1591 (2007).

353   57.   S. Kumar, G. Stecher, M. Li, C. Knyaz, K. Tamura, MEGA X: molecular evolutionary
354   genetics analysis across computing platforms. *Mol. Biol. Evol.* **35**, 1547–1549 (2018).

355   58.   M. Kimura, On the probability of fixation of mutant genes in a population. *Genetics*. **47**,
356   713 (1962).

357   59.   J. H. Gillespie, *Population genetics: a concise guide* (JHU Press, 2004).

358   60.   Y.-B. Sun, Z.-J. Xiong, X.-Y. Xiang, S.-P. Liu, W.-W. Zhou, X.-L. Tu, L. Zhong, L. Wang,
359   D.-D. Wu, B.-L. Zhang, Whole-genome sequence of the Tibetan frog Nanorana parkeri and
360   the comparative evolution of tetrapod genomes. *Proc. Natl. Acad. Sci.* **112**, E1257–E1262
361   (2015).

362   61.   A. J. Crawford, Relative rates of nucleotide substitution in frogs. *J. Mol. Evol.* **57**, 636–641
363   (2003).

364   62.   S. P. Mansai, H. Innan, The power of the methods for detecting interlocus gene conversion.
365   *Genetics*. **184**, 517–527 (2010).

366   63.   D. Feng, L. Tierney, Computing and displaying isosurfaces in R. *J. Stat. Softw.* **28**, 1–24
367   (2008).

368   64.   W. N. Venables, B. D. Ripley, Modern applied statistics with S. N. Y. Springer (2002).

369   65.   V. A. Luckow, S. Lee, G. Barry, P. Olins, Efficient generation of infectious recombinant
370   baculoviruses by site-specific transposon-mediated insertion of foreign genes into a
371   baculovirus genome propagated in Escherichia coli. *J. Virol.* **67**, 4566–4579 (1993).

372   66.   G. Petschenka, S. Fandrich, N. Sander, V. Wagschal, M. Boppré, S. Dobler, Stepwise
373   evolution of resistance to toxic cardenolides via genetic substitutions in the Na+/K+-
374   ATPase of milkweed butterflies (Lepidoptera: Danaini). *Evolution*. **67**, 2753–2761 (2013).

375   67.   H. H. Taussky, E. Shorr, A microcolorimetric method for the determination of inorganic
376   phosphorus. *J. Biol. Chem.* **202**, 675–685 (1953).

Acknowledgments: We thank M. Przeworski for helpful comments on the manuscript. We
thank C. Natarajan, K. Rohlfing, V. Wagschal, and P. Kowalski for assistance in the laboratory.
Funding: This study was funded by grants to PA from the National Institutes of Health (R01-
GM115523) and to JFS from the National Institutes of Health (R01- HL087216) and the
National Science Foundation (OIA-1736249), to SD from Deutsche Forschungsgemeinschaft
(DFG grant DO527/10-1), and a fellowship to AH from The Simons Foundation's Society of
Fellows (#633313). **Author contributions:** PA and AJC conceived of and oversaw the project;
LY, MPRO, SHA, JP, and AJC collected samples and generated sequence data; LY, AH, PA,
SHA, and KZ performed evolutionary and population genetics analyses; SM, JFS, SD, AJC and
PA designed functional experiments; SM and PA performed experiments and statistical analyses;
SM, JFS, LY, AH and PA wrote the paper; All authors edited the manuscript; **Competing
interests:** None. **Data and materials availability:** BioProject PRJNA627222, SRA:
SRR11583961-91, nucleotides: MT396181-92, genes: MT422192-MT422203.

395 **Supplementary Materials:**

396 Methods

397 Figures S1-S7

398 Tables S1-S8

399 References 38-67
400

Fig. 1. Molecular evolution of ATP1A1 in anurans. (A) Maximum likelihood phylogeny of

anuran species and mammalian outgroups derived from (*38*). Species names in purple

correspond to chemically defended toads, and blue and red colors correspond to the S and R

ATP1A1 paralogs in *Leptodactylus* species, respectively. Only variable sites with documented

roles in CG-binding or sensitivity are shown (reviewed in (*22*)). The numbering of sites is based

on sheep ATP1A1 (*Ovis aries*, Genbank: NC019458.2) and appears at the top of the table (e.g.

the first position shown is 108). Dots indicate identity with the reference sequence and letters

represent amino acid substitutions relative to the reference. The images on the left depict the five

surveyed *Leptodactylus* species and a representative toad species (*Rhinella marina*) as potential

prey. Maximum likelihood phylogeny estimates based on nucleotide sequences (B) and amino

acid sequences (C) yield distinct topologies. Bootstrap support values are indicated at internal

nodes. To the right is the pattern of amino acid variation at 12 positions that distinguish the S and

R paralogs.

Fig. 2. (A) Non-allelic gene conversion (NAGC) homogenizes sequence variation between

paralogous genes, and therefore changes the genealogical signal (adapted from (*39*)). **(B)** NAGC

can result in a genealogy in which paralogous genes in the same species share a more recent

common ancestor with one another than with their orthologous counterparts in other species

("concerted evolution"). The homogenizing effects of NAGC can be counteracted by selection

that favors the differentiation of paralogous genes. **(C)** Site-wise difference in the $\log_{10}$-

likelihood of two alternative tree topologies—generalizing the grey and purple extremes of panel

B to five *Leptodactylus* species. Shaded regions show a log-likelihood difference greater than 2

in support of the corresponding model. Only parsimony-informative variants in the ATP1A1

coding sequence are shown. Black bars correspond to the 12 R/S distinguishing nonsynonymous

substitutions (shown in red or blue in Fig 1C; Table S4).

Fig. 3. (A) Theoretical probability of maintaining distinct alleles at a single site in the face of non-allelic gene conversion (NAGC). We used a theoretical model to compute the probability of maintaining alternative amino acid states at the same site in a pair of paralogous genes, given an NAGC rate and strength of selection against allele homogenization at the site. The black dot shows the approximate mode estimate from panel B, which falls in the range in which maintenance is likely according to this theoretical model. **(B) Estimates of evolutionary parameters.** Approximate posterior probabilities were inferred based on simulations of the evolution of ATP1A1 genes in *Leptodactylus*. The x-axis shows the NAGC rate across the gene, and the y-axis shows the population selection coefficient for the 12 substitutions that distinguish the R and S paralogs across species.

Fig. 4. Functional analysis of substitutions specific to the R-type ATP1A1 paralog. (A)

ATP1A1 gene constructs with various combinations of the 12 substitutions that distinguish the S

and R paralogs. Black circles indicate an amino acid matching the R paralog whereas a white

circle indicates a match with the S paralog. Dark grey shading denotes the R background and

white denotes the S background. Light grey columns highlight two substitutions (Q111R and

N122D) that are known to confer CG-resistance. (B) Functional properties of engineered

Na^+,K^+-ATPases. The mean $\pm$ SEM $\log_{10}IC_{50}$ (i.e., a measure of CG resistance) is plotted on the

x-axis and the mean $\pm$ SEM ATP hydrolysis rate (i.e., a measure of protein activity) for the same

proteins is plotted on the y-axis. Each estimate is based on six biological replicates.