1   Fitness benefits of providing services to others: Grooming predicts survival in a neotropical

2   primate

5   Kotrina Kajokaite*[a], Andrew Whalen[b], Jeremy Koster[a,c], and Susan Perry[d,e]

7   a. Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary

8   Anthropology, Deutscher Platz 6, 04103 Leipzig, Germany

9   b. The Roslin Institute and Royal (Dick) School of Veterinary Studies, The University of

10   Edinburgh, Easter Bush Campus, Midlothian EH25 9RG, UK

11   c. Department of Anthropology, University of Cincinnati, PO BOX 210380 481 Braunstein Hall,

12   Cincinnati, OH 45221-0380, USA

13   d. Department of Anthropology, University of California, 375 Portola Plaza, Los Angeles, CA

14   90095, USA

15   e. Behavior, Evolution and Culture Program, Department of Anthropology, University of

16   California, 375 Portola Plaza, Los Angeles, CA 90095, USA

* Corresponding author: Kotrina Kajokaite, Department of Human Behavior, Ecology and

Culture, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6 04103

Leipzig, Germany

ABSTRACT

Across multiple species of social mammals, evidence is accumulating that sociality is associated

with fitness. In long-lived species, like primates, lifespan is one of the main fitness components.

A growing number of studies shows that interactions with individuals of both sexes, and both

weak and strong bonds predict survival. We used 18 years of data from the Lomas Barbudal

Monkey project to quantify sociality in 11 capuchin (*Cebus capucinus*) groups and tested

whether females' tendencies to interact with their group members are associated with survival.

We found that females who provide more grooming to other group members survive longer.

The association is stronger for older females. These benefits were independent of females'

rank, presence of their adult daughters and mothers, and their group size. Receiving more

grooming from others showed a similar effect on survival, but confidence in the estimated

effect was unclear. Participation in coalitionary aggression and foraging in close proximity of

others were not predictive of increased survival. The results presented here suggest that

females' general propensities to affiliate with members of their group may result in survival-

related benefits. Females might enjoy these benefits through exchanging grooming for other

currencies, such as coalitionary support or tolerance.

44   Keywords: capuchins, C*ebus*, sociality, social relationships, survival, fitness

INTRODUCTION

A key question for understanding the evolution of animal sociality is: Do more social individuals have higher fitness? A number of studies link components of reproductive success, such as fertility (Fedigan et al., 2008; Schülke et al., 2010; Gilby et al., 2013) and offspring survival (Silk, 2009; Silk et al., 2003; Kalbitzer et al., 2017), to differences in the social behavior of individuals. In long-lived, iteroparous species, lifespan is also an important component of life-time reproductive success that contributes to variation in fitness (Clutton-Brock, 1998) and there is evidence that aspects of individual sociality are associated with lifespan in humans and nonhuman animals (Snyder-Mackler et al., 2020).

It has been hypothesized that social animals have evolved cognition characterized by skills and motivations to monitor their social world and interact with conspecifics in fitness-enhancing ways (Jolly, 1966; Humphrey, 1976; Whiten and Byrne, 1997). Individuals cultivate differentiated social relationships through these interactions, which, in turn, affect their fitness. Discussion of causal pathways by which the social integration of an individual contributes to their fitness is just starting to emerge in the literature (Ostner and Schülke, 2018; Thompson, 2019, Snyder-Mackler et al., 2020). The task is complicated by the fact that there are multiple ways one can measure how social animals connect to others (Cheney et al., 2016; Ellis et al., 2019), and different measures may be associated with different pathways linking social behavior to fitness.

Here we ask how an adult white-faced capuchin (*Cebus capucinus*) female's survival is predicted by various kinds of social interactions. To choose a measurement of sociality that is well suited to address this question, we consider three different types of social interactions:

grooming, foraging near others, and coalitionary aggression. These interactions are

characterized by varying qualities that define the social relationships: affiliative touch, affiliation

through spatial proximity in the vicinity of food resources, and cooperation respectively (Hinde,

1976). We quantified interactions in each domain separately and, where possible, we estimated

a female's role as a provider and as a receiver of the behavior. In order to better understand

the pathways by which females translate their social interactions into survival-related benefits

(e.g., reduced wounding, increased rank, better access to resources, reduced predation risk), it

is important to document whether females are providing or receiving these services (*see*

supplemental material for detailed descriptions of pathways). Although we are testing for

predictors of fitness and are not testing specific pathways, our approach of estimating

individuals' roles separately in each behavioral domain generates hypotheses that can be

tested subsequently regarding causal pathways connecting these interactions to fitness.

Measures of individual sociality (Silk et al., 2013; Ostner and Schülke 2018; Thompson

2019; Ellis et al. 2019), vary according to the emphasis on the interaction frequency and/or

numbers of partners being incorporated into the metric. We chose an individual-level metric

that is best conceptualized as a social integration measure, where interaction frequency rather

than the number of partners is quantified. We aimed to estimate a female's overall

participation in social life without making arbitrary distinctions between strong and weak

partnerships (McFarland et al., 2017), because important interindividual variation in affiliation

tendency, number of partners, and number of strong bonds can be neglected when focusing

only on the strongest affiliative relationships (Ostner and Schülke, 2018; Thompson and Cords,

2018).

Our measure of a female's sociality can be conceptualized as an individual's base rate of

engaging in a behavior relative to that of other individuals in her social group, regardless of the

number of partners, their sex, their age or the presence of specific social bonds in her life. We

assume that participating more in some social interactions translates into more survival related

benefits received, from the group members in general. In species where individuals live in

stable, well-defined groups, like white-faced capuchin monkeys, daily interactions occur with all

types of partners (Perry and Manson, 2009; Perry, 2012). Interactions with both males and

females are associated with female longevity in wild yellow baboons (*Papio cynocephalus*)

whose social groups resemble those of capuchin monkeys (Archie et al., 2014).

We hypothesize that females who are more socially integrated experience fitness

benefits associated with survival. Specifically, we predicted that the more females provide and

receive either grooming or coalitionary support, the better they will survive. We also predicted

that females who are more likely to forage in close proximity of others will survive longer. We

tested these predictions using data spanning 18 years of capuchin monkeys' social interactions

and demography from the Lomas Barbudal Monkey Project dataset.

METHODS

Study subjects and the dataset

We studied members of the wild white-faced capuchin population at the Lomas

Barbudal Biological Reserve and surrounding private lands in Guanacaste, Costa Rica (Perry et

al., 2012). The dataset has longitudinal records including demographic information, pedigree

information, and social interactions on individuals living in 11 capuchin social groups. The data

on capuchin behavior were collected between January 2002 and December 2019. All groups in

this study were observed during at least 7 calendar years (mean = 13.5, SD = 4.4). The primary

subjects of this analysis were 132 adult females (adulthood defined as beginning at 5 years of

age). Sixty-two females died during the study period. The behavioral and demographic data on

each group were collected by experienced observers during visits lasting at least 6 hours/day.

Measuring Sociality

As measures of female sociality, we focused on three behavioral domains: grooming,

coalitionary aggression, and foraging. We treated grooming and coalition formation as directed

behaviors, and we used observations of individuals as both initiators and recipients of the

behavior. We did not have information about which individual had initiated the proximity when

foraging, and therefore foraging in proximity was treated as an undirected behavior. We

treated each of the five interaction types (*grooming giving, grooming receiving, support giving,*

support receiving, foraging in proximity) as a separate measure of sociality.

In calculating the frequency with which adult females engaged in these interactions,

we used behavioral records from all the individuals who resided in the 11 respective groups

during the study period. There was a total of 563 monkeys and 13,770 unique dyads. In addition

to the aforementioned theoretical reasons for considering interactions with partners of both

sexes and all age groups, using all recorded interactions provided more information per adult

female. We estimated the posterior mean and standard deviation of each individual-level

sociality measure and propagated that measurement uncertainty when estimating the effect of

sociality on survival (*see supplemental material*).

Grooming (grooming giving and grooming receiving)

Grooming rates were estimated using data collected during 10-minute focal follows. To

estimate individual grooming rates, we calculated dyadic counts of grooming and dyadic

opportunities for grooming. The opportunity for an ij dyad to engage in grooming, was

calculated as the sum of the focal follows of i and the focal follows of j at times when i and j

were co-resident. A count of 1 was assigned if i groomed j at least once during a focal follow,

otherwise 0 was assigned. The same was done when evaluating if j groomed i.

Joining an ongoing conflict (support giving and support receiving)

The behavior of joining a coalitionary conflict was defined as an individual intervening

on one side during an ongoing aggressive conflict. This definition only indicates the functional

aspect of joining a side; it entails no inferences about internal psychological states such as the

intent to help a specific individual. Since aggressive interactions are salient and harder to miss

than quiet activities like grooming, aggressive interactions were collected both *ad libitum* and

during focal follows. The chronological stream of aggressive behaviors was divided into 5-

minute intervals. In order to identify instances of joining a coalitionary conflict, monkey i is

identified as joining monkey j if i performed an aggressive behavior toward either monkey j's

opponent or victim within the context of the intervals. The measure is dichotomous, and a

single instance was recorded for the occasions when there were multiple observations of

monkey i joining monkey j during the interval. To calculate the opportunities to join a

coalitionary conflict, all individuals who were co-resident during the aggressive conflict were

regarded as having the opportunity to join on either side during the conflict.

Foraging in proximity

Foraging in close proximity was estimated from group scans that occurred in the

context of foraging. In group scans, the identity of the scanned individuals, their activity and

their proximity to other individuals within 10 body lengths (~2 m) was noted. We considered

individuals to be foraging in close proximity if they were scanned within 5 body lengths (~1m) of

each other. For each dyad, we scored whether they were observed foraging within close

proximity in 10-minute intervals. The number of opportunities that the dyad had to forage

within close proximity is a sum of group scans in the foraging context that are 10 minutes apart,

where one of the individuals is a subject of a group scan.

Individual sociality measures

The data for these analyses were collected across eighteen years and the number of

observed social groups and individuals generally increased over time. As a result, the density of

data is uneven across time periods, social groups, and individuals. We incorporated uneven

distributions of the data by aggregating the data annually and using adaptations of the

multilevel Social Relations Model (Snijders and Kenny, 1999; Koster et al. 2020) to estimate

individual annual estimates of grooming, coalitionary support, and foraging in close proximity

(*see supplemental material*). This method provided estimates of individual sociality that reflect

measurement uncertainty, with the uncertainty increasing for infrequently observed

individuals.

Sociality estimates from the Social Relations Model can be conceptualized as female's

base rate of either providing or receiving behavior, after accounting for average rates for the

population and the individual's social group. Conceptually, it describes how likely a female is to

participate in an interaction (either as a recipient or as a provider, if applicable) with another

individual in her social group. The measure is estimated as a random effect in the statistical

model and centered on zero, which conceptually represents an average monkey in the

population. The sociality estimate then is an offset from the population average. In general, our

sociality measures are similar to nodal in- and out-degree centrality in the social networks

tradition (Hasenjager and Dugatkin, 2015).

Modeling survival as a function of the individual sociality measure

To investigate whether sociality is associated with adult female survival, we used

Bayesian Cox proportional hazards models (*see supplemental material*). In separate models,

each of the five individual sociality measures (mean and SD) was modeled as a predictor of

survival probability over one-year periods. In the main text, we report models in which survival

in year t is modeled as a function of sociality observed in the same year. As a supplemental

analysis, we model survival in year t as a function of sociality in the preceding year, $t-1$. The

former approach has a larger sample size whereas the second approach assumes that the

benefits sociality are evident over an intermediate timeframe.

In addition to the respective sociality measures, these models included the following

time-varying (calendar year-specific) covariates: the female's age, her dominance rank (ranges

from 0 to 1, where 1 represents the highest rank; *see supplemental material*), the average

number of individuals in her group (including juveniles and infants) throughout the year, the

proportion of time during that year that her mother was alive, and the number of adult

daughters that she had (*see supplemental material for further details on covariates*).

We used a Bayesian approach to fit Cox proportional hazards models (Singer and

Willett, 2003). All of the covariates, except the number of daughters, were standardized by

subtracting the mean and dividing it by the standard deviation. The number of daughters was

centered by subtracting the mean number of daughters for each respective age.

Models were run using Stan (v.2.19.1) and the *rethinking* package (v. 1.93: McElreath,

2020) in *R* (v. 3.6.2; R Core Team 2019).

RESULTS

For perspective on the individual sociality measures, Figure 1 illustrates how the

estimates of *grooming, coalitionary support,* and *foraging in proximity* by adult females

compare to the estimates for other age-sex classes. Females differ the most from the other

age-sex classes in their distribution of *grooming giving* measures. A majority of the females

(80%) groom more than an average monkey in the population (Figure 1, panel A). The female

distributions for the rest of the behavioral domains are comparatively similar to the other age-

sex distributions, indicating that females behave similarly to adult males and immatures in

those domains.

In Figure 2, we plotted the pairwise correlations between each of the individual sociality

measures (*see supplemental materials section 3.A*) in order to investigate if female profiles are

similar across all five measures, e.g. Is a frequent groomer also a frequent recipient of grooming

and coalitional support? Within the coalitionary aggression domain, estimates of female giving

and receiving coalitionary support were highly correlated (r=0.97). In the grooming domain, the

correlation between giving and receiving estimates was moderate (r=0.57). The correlations

across behavioral domains are smaller than within a domain, between 0.15 and 0.37.

Our primary objective was to test the effects of sociality on survivorship. Table 1

presents the Cox proportional hazards model posterior mean estimates and the 95% Highest

Posterior Density Interval (HPDI), representing the narrowest interval containing the 95%

probability mass. The effect of *grooming giving* exhibited a clear positive effect ($\beta = 0.49$),

suggesting that females who groomed others more frequently exhibited greater survivorship.

The other sociality measures (particularly *grooming receiving)* had posterior estimates

consistent with sociality predicting greater survival but their 95% HPDI intervals were wider and

included zero, suggesting uncertainty about the effects of these sociality estimates.

To facilitate the interpretation of the Cox proportional hazards model coefficients, we

plotted model predictions showing the probability of dying in a given year (Figure 3). Across all

five sociality measures, the less social females have a greater probability of dying than more

social females, but only the slopes for *grooming giving* confidently suggest a beneficial effect on

survivorship (Figure 3, panel A).

The supplemental analysis, in which survival in year *t* is modeled as a function of

sociality in year *t−1*, provides qualitatively similar results.

Figure 1. Posterior mean distributions of annual individual sociality measures for each type of
behavior. The orange density represents adult females, whereas the white density represents
the rest of the population. The population base rate is normalized at approximately zero.
235
236

238   **Figure 2.** Bivariate correlations of individual sociality measures. Each point represents the
239   posterior mean of an annual estimate for an individual female in the sample.

241   **Table 1.** Estimates of fixed effects of each of the Cox proportional hazards models: posterior
242   means and 95% HPDI (see supplemental material for graphical representation of this table).
243

	Sociality measure				
Parameter	Grooming giving	Grooming receiving	Support giving	Support receiving	Foraging in proximity
Intercept	9.20 [8.86,9.56]	9.13 [8.78,9.52]	9.07 [8.74,9.42]	9.06 [8.73,9.41]	9.08 [8.74,9.42]
β sociality	0.49 [0.26,0.70]	0.19 [–0.09,0.45]	0.11 [–0.21,0.45]	0.06 [–0.28,0.40]	0.16 [–0.18,0.47]
β rank	−0.20 [–0.51,0.11]	−0.11 [–0.45,0.24]	−0.10 [–0.45,0.24]	−0.10 [–0.47,0.28]	−0.16 [–0.54,0.23]
β age	−0.57 [–0.88,–0.25]	−0.66 [–0.96,–0.37]	−0.66 [–0.96,–0.37]	−0.68 [–0.99,–0.38]	−0.69 [–0.98,–0.39]
β daughters	−0.08 [–0.45,0.28]	−0.08 [–0.45,0.31]	−0.07 [–0.44,0.31]	−0.06 [–0.43,0.30]	−0.05 [–0.43,0.32]
β mother	0.26 [–0.09,0.63]	0.31 [–0.02,0.66]	0.33 [–0.01,0.67]	0.32 [–0.01,0.65]	0.33 [–0.01,0.66]
β group size	0.19 [–0.10,0.49]	0.18 [–0.10,0.46]	0.19 [–0.10,0.48]	0.18 [–0.1,0.46]	0.18 [–0.09,0.46]

245   **Figure 3.** The predicted annual probabilities of dying at two representative ages (10 and 30) as a
246   function of the sociality levels in five behavioral domains, as calculated from the Cox
247   proportional hazards models. For both ages, we assumed median group size, median rank, and
248   average number of daughters for the age. For 10-year-old females, we assumed a co-resident
249   mother, while predictions for the 30-year-old assume the mother is deceased.
250

DISCUSSION

In female white-faced capuchin monkeys, providing grooming to others is associated

with greater survival rates. By contrast, there was no strong evidence that females who receive

more grooming, participate more often in coalitionary aggression, and forage more often in

close proximity of others also have higher survivorship. Our results in the grooming domain are

broadly consistent with previous findings that variation in social behavior is associated with

survival (Holt-Lunstad et al., 2010, Fagen and Fagen, 2004; Yee et al., 2008; Silk et al., 2010;

Archie et al., 2014; Brent et al., 2017), but depart from previous studies by considering

interactions with all social partners. Multiple studies reported that interacting with various

partners (Archie et al., 2014) and both strong (Silk et al., 2010) and weak bonds (McFarland et

al., 2017; Silk et al., 2018) predict survival. Our results suggest that females' propensity to

affiliate with all of her group members predicts survivorship.

Our study also adds a new dimension by considering the direction of the social

interaction: female capuchins who were providing most grooming were the least likely to die,

but there was no reliable relationship between receiving most grooming and survival. If females

are exchanging grooming for food sharing (de Waal 1989, 1997; Jaeggi & Gurven 2013) or

tolerance (Tiddi et al., 2011) their grooming provider estimate should correlate with survival,

while their grooming receiving estimate might not. Notably, whereas capuchins vary

considerably in the amount of grooming they provide, the variance of the *groom receiving* is

relatively modest, which potentially explains why the effect of this latter variable on survival is

positive but statistically inconclusive. Overall, the respective causal pathways that connect

giving and receiving behaviors to survival are likely to be non-overlapping. Accordingly,

estimating individuals' roles as providers or recipients of directed social behaviors enables

insights into the mechanisms by which sociality contributes to fitness.

The causal pathway that connects the providing of grooming to females' survival is

potentially mediated by multiple variables. The survivorship benefits might accrue to avid

groomers via increased spatial centrality within the group, in which case the females are

spending more time around other individuals, plausibly reducing predation risk (Josephs et al.,

2016). Another alternative is that females might be trading grooming for other benefits. For

example, in return for grooming, females might receive more coalitionary support when in

conflict and either avoid injuries or improve their rank, both of which can result in lowered risk

of death. To test the hypothesis that grooming is exchanged for coalitionary support, future

studies could assess whether female grooming is reciprocated by recipients through a different

currency such as coalitionary support (Kajokaite et al. in prep). Another possibility is that

females trade grooming for tolerance and engage in conflicts less often, which results in better

access to resources and/or less wounding. Although we do not directly test this mediator, or

any other pathway, the result that female's grooming giving estimates are only weakly

correlated with their support giving and receiving in coalitionary conflicts suggest that those

who spend a lot of time grooming are unlikely to receive a lot of coalitionary support in return

for these services.

Our analysis does not rule out possible confounders that could influence both grooming

given and survival, in which case the observed correlation could be caused by other factors

(e.g., a favorable genotype that promotes social behavior and survival) (Ostner and Schülke

2018). Also, although our supplemental models examine the effects of sociality and survival in

successive years, it is possible that physical senescence induces a correlation between social

behavior and survival on longer timescales. Although assessments of health are logistically

challenging among habituated primate groups, it would be beneficial for future studies to

consider variation in physical condition as a proximate mechanism of the patterns reported

here.

CONCLUSION

Across taxa, research on sociality has revealed diverse effects on components of fitness.

This study examines sociality in detail, showing that grooming others is the social behavior that

most prominently distinguishes individuals, and this aspect of sociality concomitantly predicts

the survivorship of adult female capuchins. Therefore, this study adds a neotropical primate

species to the list of mammalian species where a similar association between sociality and

survival has been demonstrated. These results accentuate the need for greater attention to the

mechanistic pathways that connect sociality to fitness (Ostner and Schülke, 2018; Thompson,

2019). As longitudinal data become increasingly common in studies of animal behavior, careful

analyses can better elucidate the evolutionary consequences of variation in sociality among

individuals.

ETHICS

The study was strictly observational, all protocols were approved by UCLA's Animal Care

Committee (protocol 2016-022). All necessary permits were obtained from SINAC and MINAE

(the Costa Rican government bodies responsible for research on wildlife) and renewed every 6

months over the course of the study; the most recent scientific passport number being #117-

2019-ACAT and the most recent permit being Resolución # M-P-SINAC-PNI-ACAT-072-2019. This

research is in compliance with the Animal Behavior Society's Guidelines for the Use of Animals

in Research.

DATA AND CODE ACCESSIBILITY

The data and code used to produce the analyses in this paper are available at:

www.doi.org/10.5281/zenodo.4036211

ACKNOWLEDGEMENTS

The following field assistants contributed significant amounts of data to the Lomas Barbudal

Monkey Project dataset: C. Angyal, A. Autor, B. Barrett, L. Beaudrot, M. Bergstrom, R. Berl, A.

Bjorkman, L. Blankenship, T. Borcuch, J. Broesch, D. Bush, J. Butler, F. Campos, C. Carlson, A.

Cobden, M. Corrales, J. Damm, B. Davis, C. deRango, N. Donati, C. Dillis, G. Dower, R. Dower, A.

Duchesneau, K. Feilen, J. Fenton, S. Fiello, K. Fisher, A. Fuentes J., M. Fuentes A., T. Fuentes, C.

Gault, H. Gilkenson, I. Godoy, I. Gottlieb, J. Griciute, L.M. Guevara R., L. Hack, R. Hammond, S.

Herbert, C. Hirsch, M. Hoffman, A. Hofner, C. Holman, J. Hubbard, S. Hyde, M. Jackson, O.

Jacobson, E. Johnson, L. Johnson, S. Lee, S. Leinwand, T. Lord, M. Kay, E. Kennedy, D. Kerhoas-

Essens, E. Johnson, S. Kessler, B. Krimmel, W. Lammers, S. Lopez, S. MacCarter, J. Mackenzie,

M. Mayer, F. McKibben, W. Meno, A. Mensing, M. Milstein, C. Mitchell, Y. Namba, A. Neyer, C.

O'Connell, J. C. Ordoñez, J., N. Parker, B. Pav, R. Popa, K. Potter, K. Ratliff, K. Reinhardt, N.

Roberts Buceta, E. Rothwell, J. Rottman, H. Ruffler, S. Sanford, M. Saul, E. Seabright, I.

Schamberg, S. Schembari, N. Schleissman, C. Schmitt, S. Schulze, A. Scott, S. Sita, J. Shih, K.

Stewart, W. Tucker, E. Urquhart, K. van Atta, J. Vandermeer, L. van Zuidam, J. Verge, V. Vonau,

A. Walker-Bolton, E. Wikberg, E. Williams, L. Wolf, and D. Wood. We thank H. Gilkenson and W.

Lammers for long-term management of the field site during 2002-2013. We thank the Costa

Rican park service (Sistema Nacional de Áreas de Conservación and Área de Conservación

Tempisque), Hacienda Pelon de la Bajura, Hacienda Brin D'Amor, and the residents of San

Ramon de Bagaces for permission to work on their land. D. Cohen assisted with database

management and queries. This paper has benefited from helpful discussions with H. Clark

Barrett, Donald Cohen, Joseph Manson, Mary Brooke McElreath, Oliver Schülke, and Julia

Ostner and two anonymous reviewers.

REFERENCES

Archie, E. A., Tung, J., Clark, M., Altmann, J., & Alberts, S. C. (2014). Social affiliation matters:

Both same-sex and opposite-sex relationships predict survival in wild female baboons.

Proceedings of the Royal Society B: Biological Sciences, *281*(1793), 20141261.

Brent, L. J. N., Ruiz-Lambides, A., & Platt, M. L. (2017). Family network size and survival across

the lifespan of female macaques. *Proceedings of the Royal Society B: Biological Sciences*,

284(1854), 20170515.

Cheney, D. L., Silk, J. B., & Seyfarth, R. M. (2016). Network connections, dyadic bonds and

fitness in wild female baboons. *Royal Society Open Science*, *3*(7), 160255.

Clutton-Brock, T. H. (1988). *Reproductive success: Studies of individual variation in contrasting*

breeding systems. University of Chicago Press.

De Waal, F. B. (1989). Food sharing and reciprocal obligations among chimpanzees. *Journal of*

Human Evolution, *18*(5), 433-459.

De Waal, F. B. (1997). The chimpanzee's service economy: Food for grooming. *Evolution and*

Human Behavior, *18*(6), 375-386.

Ellis, S., Snyder-Mackler, N., Ruiz-Lambides, A., Platt, M. L., & Brent, L. J. (2019). Deconstructing

sociality: the types of social connections that predict longevity in a group-living

primate. *Proceedings of the Royal Society B*, *286*(1917), 20191991.

Fagen, R., & Fagen, J. (2004). Juvenile survival and benefits of play behaviour in brown bears,

Ursus arctos. *Evolutionary Ecology Research*, *6*(1), 89–102.

Fedigan, L. M., Carnegie, S. D., & Jack, K. M. (2008). Predictors of reproductive success in female

white-faced capuchins (Cebus capucinus). *American Journal of Physical Anthropology: The*

Official Publication of the American Association of Physical Anthropologists, *137*(1), 82–90.

Gilby, I. C., Brent, L. J., Wroblewski, E. E., Rudicell, R. S., Hahn, B. H., Goodall, J., & Pusey, A. E.

(2013). Fitness benefits of coalitionary aggression in male chimpanzees. *Behavioral Ecology*

and Sociobiology, *67*(3), 373–381.

Hasenjager, M. J., & Dugatkin, L. A. (2015). Chapter Three—Social Network Analysis in

Behavioral Ecology. In M. Naguib, H. J. Brockmann, J. C. Mitani, L. W. Simmons, L. Barrett,

S. Healy, & P. J. B. Slater (Eds.), *Advances in the Study of Behavior* (Vol. 47, pp. 39–114).

Academic Press. https://doi.org/10.1016/bs.asb.2015.02.003

Hinde, R. A. (1976). Interactions, relationships and social structure. *Man*, 1–17.

Holt-Lunstad, J., Smith, T. B., & Layton, J. B. (2010). Social relationships and mortality risk: A

meta-analytic review. *PLoS Medicine*, *7*(7), e1000316.

Humphrey, N. K. (1976). The social function of intellect. In Growing points in ethology. edited by

B. Bertram, P. Bateson, and R. Hinde (pp. 303-317). Cambridge University Press.

Jaeggi, A. V., & Gurven, M. (2013). Reciprocity explains food sharing in humans and other

primates independent of kin selection and tolerated scrounging: a phylogenetic meta-

analysis. *Proceedings of the Royal Society B: Biological Sciences*, *280*(1768), 20131615.

Jolly, A. (1966). Lemur social behavior and primate intelligence. Science, 153(3735), 501-506.

Josephs, N., Bonnell, T., Dostie, M., Barrett, L., & Henzi, S. P. (2016). Working the crowd:

sociable vervets benefit by reducing exposure to risk. *Behavioral Ecology*, *27*(4), 988-994.

Kalbitzer, U., Bergstrom, M. L., Carnegie, S. D., Wikberg, E. C., Kawamura, S., Campos, F. A., Jack,

K. M., & Fedigan, L. M. (2017). Female sociality and sexual conflict shape offspring survival

in a Neotropical primate. *Proceedings of the National Academy of Sciences*, *114*(8), 1892–

1897.

Koster, J., Leckie, G., & Aven, B. (2020). Statistical methods and software for the Multilevel

Social Relations Model. *Field Methods*, *32*(4), 339-345.

McElreath, R. (2020). rethinking: Statistical rethinking book package (R package Version 1.93).

https://github.com/rmcelreath/rethinking.

McFarland, R., Murphy, D., Lusseau, D., Henzi, S. P., Parker, J. L., Pollet, T. V., & Barrett, L.

(2017). The 'strength of weak ties' among female baboons: fitness-related benefits of

social bonds. *Animal Behaviour*, *126*, 101-106.

Ostner, J., & Schülke, O. (2018). Linking sociality to fitness in primates: A call for mechanisms. In

Advances in the Study of Behavior (Vol. 50, pp. 127–175). Elsevier.

Perry, S. (2012). The behavior of wild white-faced capuchins: Demography, life history, social

relationships, and communication. In *Advances in the study of behavior* (Vol. 44, pp. 135–

181). Elsevier.

Perry, S., Manson, J. H., & Perry, S. (2009). *Manipulative monkeys*. Harvard University Press.

Schülke, O., Bhagavatula, J., Vigilant, L., & Ostner, J. (2010). Social bonds enhance reproductive

success in male macaques. *Current Biology*, *20*(24), 2207–2210.

Silk, J. B. (2009). Nepotistic cooperation in non-human primate groups. *Philosophical

Transactions of the Royal Society B: Biological Sciences*, *364*(1533), 3243–3254.

Silk, J. B., Alberts, S. C., & Altmann, J. (2003). Social bonds of female baboons enhance infant

survival. *Science*, *302*(5648), 1231–1234.

Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., Wittig, R. M.,

Seyfarth, R. M., & Cheney, D. L. (2010). Strong and consistent social bonds enhance the

longevity of female baboons. *Current Biology*, *20*(15), 1359–1361.

Silk, J., Cheney, D., & Seyfarth, R. (2013). A practical guide to the study of social relationships.

Evolutionary Anthropology: Issues, News, and Reviews, *22*(5), 213–225.

Silk, J. B., Seyfarth, R. M., & Cheney, D. L. (2018). Quality versus quantity: do weak bonds

enhance the fitness of female baboons?. *Animal Behaviour*, *140*, 207-211.

Singer, J. D., & Willett, J. B. (2003). Survival analysis. *Handbook of psychology*, 555-580.

Snijders, T. A., & Kenny, D. A. (1999). The social relations model for family data: A multilevel

approach. *Personal Relationships*, *6*(4), 471–486.

Snyder-Mackler, N., Burger, J. R., Gaydosh, L., Belsky, D. W., Noppert, G. A., Campos, F. A., ... &

Tung, J. (2020). Social determinants of health and survival in humans and other

animals. *Science*, *368*(6493).

Thompson, N. A. (2019). Understanding the links between social ties and fitness over the life

cycle in primates. *Behaviour*, *1*, 1–50.

Thompson, N. A., & Cords, M. (2018). Stronger social bonds do not always predict greater

longevity in a gregarious primate. *Ecology and Evolution*, *8*(3), 1604–1614.

Tiddi, B., Aureli, F., Polizzi di Sorrentino, E., Janson, C. H., & Schino, G. (2011). Grooming for

tolerance? Two mechanisms of exchange in wild tufted capuchin monkeys. *Behavioral*

Ecology, *22*(3), 663–669.

Whiten, A., & Byrne, R. W. (Eds.). (1997). *Machiavellian intelligence II: Extensions and*

evaluations (Vol. 2). Cambridge University Press.

Yee, J. R., Cavigelli, S. A., Delgado, B., & McClintock, M. K. (2008). Reciprocal affiliation among

adolescent rats during a mild group stressor predicts mammary tumors and lifespan.

Psychosomatic Medicine, *70*(9), 1050.