Fitness benefits of providing services to others: Sociality predicts survival in a neotropical

primate

Kotrina Kajokaite*[a], Andrew Whalen[b], Jeremy Koster[a,c], and Susan Perry[d,e]

a. Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary

Anthropology, Deutscher Platz 6, 04103 Leipzig, Germany

b. The Roslin Institute and Royal (Dick) School of Veterinary Studies, The University of

Edinburgh, Easter Bush Campus, Midlothian EH25 9RG, UK

c. Department of Anthropology, University of Cincinnati, PO BOX 210380 481 Braunstein Hall,

Cincinnati, OH 45221-0380, USA

d. Department of Anthropology, University of California, 375 Portola Plaza, Los Angeles, CA

90095, USA

e. Behavior, Evolution and Culture Program, Department of Anthropology, University of

California, 375 Portola Plaza, Los Angeles, CA 90095, USA

* Corresponding author: Kotrina Kajokaite, Department of Human Behavior, Ecology and

Culture, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6 04103

Leipzig, Germany

ABSTRACT

Across multiple species of social mammals, evidence is accumulating that sociality is associated

with fitness. In long-lived species, like primates, lifespan is one of the main fitness components.

Here, we provide the first direct evidence that females who provide more in social services live

longer. We used 18 years of data from the Lomas Barbudal Monkey project to quantify sociality

in 11 capuchin (*Cebus capucinus*) social groups using three interaction types: grooming, support

in coalitionary aggression, and foraging in close proximity. We analyzed each domain separately

to understand how giving and receiving social favors predict survivorship. To estimate female

sociality in each of these domains, we implemented an adaptation of the Social Relations

Model. This approach enabled us to estimate individual rates of giving and receiving

interactions while accounting for the inherent measurement uncertainty in these estimates due

to varying amounts of observations of each individual. Subsequently, we modeled adult

females' survival as a function of their sociality estimates, rank, age, group size, and maternal

kin presence using a Bayesian Cox proportional hazards model. We found that females who

provide more grooming to other group members survive longer, but other social interaction

types had little effect on survivorship.

Keywords: capuchins, *Cebus*, sociality, social relationships, survival, fitness

INTRODUCTION

A key question for understanding the evolution of animal sociality is: Do animals that

are more social enjoy greater fitness benefits? A number of studies link components of

reproductive success, such as fecundity (Fedigan et al., 2008; Schülke et al., 2010; Gilby et al.,

2013) and offspring survival (Silk, 2009; Silk et al., 2003; Kalbitzer et al., 2017), to differences in

the social behavior of individuals.

In long-lived, iteroparous species, lifespan is also an important component of

reproductive success contributing to fitness variation (Clutton-Brock, 1998). In human

populations, being socially connected is associated with a lower probability of dying (Holt-

Lunstad et al., 2010; Shor and Roelfs, 2015; Shor et al., 2013), while having few and poor social

relationships is associated with a higher risk of death (Roelfs et al., 2011). In nonhuman species

there is similar evidence that various aspects of individual sociality are associated with lifespan

(e.g., Silk et al. 2010).

Detailed longitudinal behavioral records are necessary for relating an individual's social

behavior patterns to longevity (Clutton-Brock, 1988). Because these data are logistically difficult

to collect in many habitats, the data are often challenging to analyze given heterogeneous

sampling across time periods, social groups, and individuals. Although research on sociality

often ignores this sampling variability (reviewed in Silk et al. 2013), the measurement

uncertainty can be substantial, complicating inferences.

The primary objective of this paper in an investigation of the effects of sociality on

survivorship among female white-faced capuchins (*Cebus capucinus*). We test this relationship

using data spanning 18 years of capuchin monkeys' social interactions and demography from

the Lomas Barbudal Monkey dataset. To quantify individual sociality, we selected interactions

from three domains to represent the variety of social behaviors that capuchin monkeys engage

in: grooming, coalitionary aggression, and foraging in proximity to others. A secondary goal of

the paper is to develop analytical approaches for the measurement uncertainty stemming from

heterogenous sampling, which we address by adapting the multilevel Social Relations Model

(Snijders and Kenny, 1999; Koster et al. 2020). We generate individuals' sociality estimates that

reflect the measurement uncertainty, which are subsequently used to test the hypothesis that

social behavior predicts longevity.

METHODS

Study subjects and the dataset

We studied members of the wild white-faced capuchin population at the Lomas

Barbudal Biological Reserve and surrounding private lands in Guanacaste, Costa Rica (Perry et

al., 2012). The dataset has longitudinal records including demographic information, pedigree

information, and social interactions on individuals living in 11 capuchin social groups. The data

on capuchin behavior were collected between January 2002 and December 2019. All groups in

this study were observed in at least 7 calendar years (mean = 13.27). The primary subjects of

this analysis were 132 adult females, where adulthood is assumed to begin at the age of 5

years. The behavioral and demographic data on each group were collected by experienced

observers during visits lasting at least 6 hours/day.

Measuring Sociality

As measures of female sociality, we focused on three behavioral domains: grooming,

coalitionary aggression, and foraging. We treated grooming and coalition formation as directed

behaviors, and we used observations of individuals as both initiators and recipients of the

behavior. We did not have information about which individual had initiated the proximity when

foraging, and therefore foraging in proximity was treated as an undirected behavior. In

calculating the frequency with which adult females engaged in these interactions, we used

behavioral records from all the individuals who resided in the 11 respective groups during the

study period. There were a total of 563 monkeys and 13,770 unique dyads. We treated each of

the five interaction types (*grooming giving, grooming receiving, support giving, support*

receiving, forage in proximity) as a separate measure of sociality.

Grooming (groom giving and groom receiving)

Grooming rates were estimated using data collected during 10-minute focal follows. To

estimate individual grooming rates, we calculated dyadic counts of grooming and dyadic

opportunities for grooming. The opportunity for a dyad, A-B, to engage in grooming, was

calculated as the sum of the focal follows of A and the focal follows of B at times when A and B

were co-resident. A count of 1 was assigned if A groomed B at least once during a focal follow,

otherwise 0 was assigned. The same was done when evaluating if B groomed A.

Joining a coalitionary conflict (support giving and support receiving)

The behavior of joining a coalitionary conflict was defined as an individual intervening

on one side during an ongoing aggressive conflict. This definition only indicates the functional

aspect of joining a side; it entails no inferences about internal psychological states such as the

intent to help a specific individual. Since aggressive interactions are salient and harder to miss

than quiet activities like grooming, aggressive interactions were collected both *ad libitum* and

during focal follows. The chronological stream of aggressive behaviors was divided into 5-

minute intervals. In order to identify instances of joining a coalitionary conflict, monkey A is

identified as joining monkey B if A performed an aggressive behavior toward either monkey B's

opponent or victim within the context of the intervals. The measure is dichotomous, and a

single instance was recorded for the occasions when there were multiple observations of

monkey A joining monkey B during the interval. To calculate the opportunities to join a

coalitionary conflict, all individuals who were co-resident during the aggressive conflict were

regarded as having the opportunity to join on either side during the conflict.

Foraging in proximity

Foraging in close proximity was estimated from group scans that occurred in the

context of foraging. In group scans, the identity of the scanned individuals, their activity and

their proximity to other individuals within 10 body lengths (~2 m) was noted. We considered

individuals to be foraging in close proximity if they were scanned within 5 body lengths (~1m) of

each other. For each dyad, we scored whether they were observed foraging within close

proximity in 10-minute intervals. The number of opportunities that the dyad had to forage

within close proximity is a sum of group scans in the foraging context that are 10 minutes apart,

where one of the individuals is a subject of a group scan.

Individual sociality measures

The data for these analyses were collected across eighteen years and the number of

observed social groups and individuals generally increased over time. As a result, the density of

data is uneven across time periods, social groups, and individuals. We incorporated uneven

distributions of the data by aggregating the data annually and using adaptations of the

multilevel Social Relations Model (Snijders and Kenny, 1999; Koster et al. 2020) to estimate

individual annual rates of grooming, coalitionary support, and foraging (see supplemental

material). This method provided estimates of individual sociality that reflect the measurement

uncertainty, with the uncertainty increasing for infrequently observed individuals.

Modeling survival as a function of individual sociality measure

To investigate whether sociality is associated with adult female longevity, we used

Bayesian Cox proportional hazards models. In separate models, each of the five individual

sociality measures was modeled as a predictor of survival probability over one-year periods. In

addition to the respective sociality measures, these models included the following time-varying

(calendar year-specific) covariates: the female's age, her dominance rank (ranges from 0 to 1,

where 1 represents the highest rank), the average number of individuals in her group, the

proportion of time during that year that her mother was alive, and the number of adult

daughters that she had (see supplemental material for further details on covariates).

Modeling approach

We specified the following model for the number of days before death, D_i,. The

probability for the number of days before death comes from the cumulative probability

distribution:

$$\Pr\left(D_i \mid \lambda_i\right) = \lambda_i \exp(-\lambda_i D_i) . \tag{3}$$

For females who did not die during the observation period, the probability of surviving D_i

comes from the complementary cumulative probability distribution:

$$\Pr\left(D_i \mid \lambda_i\right) = \exp\left(-\lambda_i D_i\right).\qquad(4)$$

We model the rate of dying, λ_i, as follows:

$$\lambda_i = 1/\mu_i\,.\qquad(5)$$

where μ_i is the expected number of days till death

$$\log(\mu_i) = \alpha + a[\,id\,] + b_{sociality} * true\ sociality\ estimate[id] + b_{rank} * rank\qquad(6)$$

$$+ b_{age} * age + b_{daughters} * daughters$$

$$+ b_{mother} * mother + b_{grsize} * group\ size\,.$$

where α denotes the intercept or the base rate of number of days survived, $\alpha[id]$ denotes

individual female random effects corresponding to the observation period. The model

coefficients $b_{sociality}$, b_{rank}, b_{age}, $b_{daughters}$, b_{mother}, and b_{grsize} describe the impact of

sociality, rank, age, number of daughters, mother's presence, and group size, respectively. We

adopted a latent variable approach to model the sociality estimates given that the individuals'

sociality estimates are not point estimates, but rather posterior distributions with means and

standard deviations that reflect the measurement uncertainty.

We used the Bayesian approach to fit Cox proportional hazards models (Singer and

Willett 2003). We assumed a Normal (8, 0.5) prior for a base rate of survival, α, which places

most of the prior mass between 0 to 20 years with the mean of 8 years and a long tail allowing

more extreme values. For fixed effects, we assumed Normal (0, 1) priors. For individual-level

random effects, $\alpha[id]$, we use a Normal (0, 1) prior (see supplemental material). All of the

covariates, except the number of daughters, were standardized by subtracting the mean and

dividing it by the standard deviation. The number of daughters was centered by subtracting the

mean number of daughters for each age.

Models were run using Stan (v.2.19.1) and the *rethinking* package (v. 1.93: McElreath,

2020) in *R* (v. 3.6.2; R Core Team 2019).

RESULTS

For perspective on the individual sociality measures, Figure 1 illustrates how the

estimates of *grooming, coalitionary support,* and *foraging in proximity* by adult females

compare to the estimates for other age-sex classes. Females differ the most from the other

age-sex classes in their distribution of *grooming giving* measures. A majority of the females

(80%) groom more than an average monkey in the population (Figure 1, panel A). The female

distributions for the rest of the behavioral domains are comparatively similar to the other age-

sex distributions, indicating that females behave similarly to adult males and immatures in

those domains.

We plotted the pairwise correlations between each of the sociality measures in Figure 2

in order to investigate if female profiles are similar across all five measures, e.g. is a frequent

groomer also a frequent recipient of grooming and coalitional support? Within the coalitionary

aggression domain, estimates of female giving and receiving coalitionary support were highly

correlated (r=0.97). In the grooming domain, the correlation between giving and receiving

estimates was moderate (r=0.57). The correlations across behavioral domains are smaller than

within a domain, between 0.15 and 0.37.

Our primary objective was to test the effects of sociality on survivorship. Table 1

presents the Cox proportional hazards model posterior mean estimates and the 95% Highest

Posterior Density Interval (HPDI), representing the narrowest interval containing the 95%

probability mass. Independent of the effects of covariates, females who groomed others more

survived at higher rates than other females. The other sociality measures had mean estimates

consistent with sociality predicting greater survival, but their HPDI intervals were wide and

included zero, suggesting uncertainty about the effects of these estimates.

To facilitate the interpretation of the Cox proportional hazards model coefficients, we

plotted model predictions showing the probability of dying in a given year (Figure 3). Across all

five sociality measures, the less social females have a greater probability of dying than more

social females, but only the slopes for *groom giving* confidently suggest a beneficial effect on

survivorship (Figure 3, panel A).

Figure 1. Posterior mean distributions of annual individual sociality measures for each type of
behavior. The orange density represents adult females whereas the white density represents
the rest of the population. The population base rate is normalized at approximately zero.
213
214

216 **Figure 2.** Bivariate correlations of individual sociality measures. Each point represents the
217 posterior mean of an annual estimate for an individual female in the sample.

219 **Table 1.** Estimates of fixed effects of each of the Cox proportional hazards models: posterior
220 means and 95% HPDI (see supplemental material for graphical representation of this table).
221

Parameter	Sociality measure				
	Grooming giving	Grooming receiving	Support giving	Support receiving	Foraging in proximity
Intercept	9.20 [8.86,9.56]	9.13 [8.78,9.52]	9.07 [8.74,9.42]	9.06 [8.73,9.41]	9.08 [8.74,9.42]
β sociality	0.49 [0.26,0.70]	0.19 [–0.09,0.45]	0.11 [–0.21,0.45]	0.06 [–0.28,0.40]	0.16 [–0.18,0.47]
β rank	−0.20 [–0.51,0.11]	−0.11 [–0.45,0.24]	−0.10 [–0.45,0.24]	−0.10 [–0.47,0.28]	−0.16 [–0.54,0.23]
β age	−0.57 [–0.88,–0.25]	−0.66 [–0.96,–0.37]	−0.66 [–0.96,–0.37]	−0.68 [–0.99,–0.38]	−0.69 [–0.98,–0.39]
β daughters	−0.08 [–0.45,0.28]	−0.08 [–0.45,0.31]	−0.07 [–0.44,0.31]	−0.06 [–0.43,0.30]	−0.05 [–0.43,0.32]
β mother	0.26 [–0.09,0.63]	0.31 [–0.02,0.66]	0.33 [–0.01,0.67]	0.32 [–0.01,0.65]	0.33 [–0.01,0.66]
β group size	0.19 [–0.1,0.49]	0.18 [–0.10,0.46]	0.19 [–0.10,0.48]	0.18 [–0.1,0.46]	0.18 [–0.09,0.46]

Figure 3. The predicted annual probabilities of dying at two representative ages (10 and 30) as a
function of the sociality levels in five behavioral domains, as calculated from the Cox
proportional hazards models. For both ages, we assumed median group size, median rank, and
average number of daughters for the age. For 10-year-old females, we assumed a co-resident
mother, while predictions for the 30-year-old assume the mother is deceased.

DISCUSSSION

In female white-faced capuchin monkeys, providing grooming to others is associated

with greater survival rates. Females who provide more grooming die at lower rates than

females who engage in this behavior less often. By contrast, there was no strong evidence that

females who receive more grooming, participate more often in coalitionary aggression, and

forage more often in close proximity of others also have higher survivorship. Our results in the

grooming domain are consistent with findings that social integration is associated with

longevity in humans (Holt-Lunstad et al., 2010) and adds a neotropical primate species to the

list of mammalian species where a similar association between sociality and longevity has been

demonstrated (Fagen and Fagen, 2004; Yee et al. 2008; Cameron et al. 2009; Silk et al., 2010;

Archie et al., 2014; Brent, 2017; Thompson and Cords, 2018).

A biologically interesting question deals with the mechanisms that facilitate the

relationship between sociality and longevity. In the human literature, two general models have

been proposed for the influence of social relationships on health (Holt-Lunstad et al. 2010).

First, the stress buffering model assumes that social relationships provide resources that affect

either behavioral or neuroendocrine responses which buffer the influence of stress on health.

Second, the main effects model assumes that participating in social relationships itself

encourages healthy behaviors.

In the primate literature, recently there has been discussion of the specific pathways

that connect social behavior to fitness (Ostner and Schülke 2018; Thompson, 2019). Consistent

with the stress buffering model, providing grooming can have either direct effects on longevity

or influence other behaviors that in turn influence survival.

Directly, females might derive benefits when grooming others, such as social bonding

through the release of hormones (Dunbar 2010). Both providing and receiving grooming

seemingly influence oxytocin and glucocorticoid levels in primates (reviewed in Crockford et al.

2017), but our analysis reveals a stronger effect of providing grooming on survival in

comparison to receiving grooming. The variation in engaging in these behaviors among

individuals helps to explain this pattern: Adult females do not vary greatly in how much

grooming they receive; therefore, the neuroendocrine benefits they receive from it are less

likely to differentiate the stress buffering among individuals. However, females vary

considerably in how much they are likely to groom others, and those who are the most avid

groomers are likely to derive the greatest benefits related to stress reduction.

Indirectly, grooming interactions potentially shape females' participation in coalitionary

behavior via the effects of mediating variables. Notably, the females who groom others the

most are usually not the same females who participate in coalitionary aggression the most

frequently. It is possible that avid groomers receive less aggression and, as a result, fewer of the

detrimental consequences that directly reduce survival.

In this study, confidence in the statistical findings is enhanced by a methodological

approach that incorporates the measurement uncertainty that typifies observational data. Data

from the natural habitats are notoriously difficult to gather. Inevitably, the records for social

interactions are sparse (Farine, 2015) and some individuals are observed more often than

others (Silk et al., 2013). Based on limited observational data, some dyads may appear to have

no social relationship, whereas if behavioral sampling were sufficiently dense, they would be

observed to interact, albeit rarely (Farine, 2015).

To address the uneven sampling effort of individuals and social groups in our dataset,

we adapted the Social Relations Model. Based on multilevel partial pooling (McElreath 2002),

the resulting estimates of sociality reflect the latent uncertainty of observational data. We

generated these estimates from dyadic observations of adult females with all other group

members rather than demographic subsets of peers as in previous studies (e.g. Silk et al., 2010;

Archie et al., 2014; Kalbitzer et al., 2017). The methods used in this study, however, offer

similar benefits for research on subgroups of dyads and more generally provide an alternative

that is suited to the imbalanced data structures of observational research.

CONCLUSION

Across taxa, research on sociality has revealed diverse effects on components of fitness.

This study examines sociality in detail, showing that grooming others is the social behavior that

most prominently distinguishes individuals, and this aspect of sociality concomitantly predicts

the survivorship of adult female capuchins. These results accentuate the need for greater

attention to the mechanistic pathways that connect sociality to fitness (Ostner and Schülke,

2018). As longitudinal data become increasingly common in studies of animal behavior, careful

analyses can better elucidate the evolutionary consequences of variation in sociality among

individuals.

ETHICS

The study was strictly observational, all protocols were approved by UCLA's Animal Care

Committee (protocol 2016-022). All necessary permits were obtained from SINAC and MINAE

(the Costa Rican government bodies responsible for research on wildlife) and renewed every 6

months over the course of the study; the most recent scientific passport number being #117-

2019-ACAT and the most recent permit being Resolución # M-P-SINAC-PNI-ACAT-072-2019. This

research is in compliance with the Animal Behavior Society's Guidelines for the Use of Animals

in Research.

DATA AND CODE ACCESSIBILITY

The data and code used to produce the analyses in this paper are available at:

www.doi.org/10.5281/zenodo.4036211

AUTHOR CONTRIBUTIONS

K.K. and S.E.P. designed the study, wrote the queries to assembled the data set. K.K. collected

some of the data. S.E.P. runs the long-term study, wrote the grants funding data collection, was

the primary data collector and data cleaner, supervised the data collection and cleaning done

by others, and manages the database. K.K. and J.K. developed SRM analyses and K.K. set up and

carried out the implementation. K.K. and A.W. developed and analyzed Bayesian Cox

proportional hazards model. K.K. took the lead in writing the manuscript. All authors provided

critical feedback and helped to write the manuscript.

COMPETING INTERESTS

We declare we have no competing interests.

FUNDING

K.K. was supported by a University of California-Los Angeles (UCLA) Pauley Fellowship, a UCLA

Dissertation Year Fellowship, and by the Max Planck Institute for Evolutionary Anthropology.

This material is based on data collection supported by funding provided to S.E.P. by the

National Science Foundation (under grant numbers 1638428, 0613226, 0848360, and 1232371),

318 the Max Planck Institute for Evolutionary Anthropology, the L.S.B. Leakey Foundation (5 grants),

319 the National Geographic Society (5 grants), the Templeton World Charity Foundation (grant

320 #0208), the Wild Capuchin Foundation, and multiple Committee on Research grants from UCLA.

321 ACKNOWLEDGEMENTS

322 The following field assistants contributed significant amounts of data to the Lomas Barbudal

323 Monkey Project dataset: C. Angyal, A. Autor, B. Barrett, L. Beaudrot, M. Bergstrom, R. Berl, A.

324 Bjorkman, L. Blankenship, T. Borcuch, J. Broesch, D. Bush, J. Butler, F. Campos, C. Carlson, A.

325 Cobden, M. Corrales, J. Damm, B. Davis, C. deRango, N. Donati, C. Dillis, G. Dower, R. Dower, A.

326 Duchesneau, K. Feilen, J. Fenton, S. Fiello, K. Fisher, A. Fuentes J., M. Fuentes A., T. Fuentes, C.

327 Gault, H. Gilkenson, I. Godoy, I. Gottlieb, J. Griciute, L.M. Guevara R., L. Hack, R. Hammond, S.

328 Herbert, C. Hirsch, M. Hoffman, A. Hofner, C. Holman, J. Hubbard, S. Hyde, M. Jackson, O.

329 Jacobson, E. Johnson, L. Johnson, S. Lee, S. Leinwand, T. Lord, M. Kay, E. Kennedy, D. Kerhoas-

330 Essens, E. Johnson, S. Kessler, B. Krimmel, W. Lammers, S. Lopez, S. MacCarter, J. Mackenzie,

331 M. Mayer, F. McKibben, W. Meno, A. Mensing, M. Milstein, C. Mitchell, Y. Namba, A. Neyer, C.

332 O'Connell, J. C. Ordoñez, J., N. Parker, B. Pav, R. Popa, K. Potter, K. Ratliff, K. Reinhardt, N.

333 Roberts Buceta, E. Rothwell, J. Rottman, H. Ruffler, S. Sanford, M. Saul, E. Seabright, I.

334 Schamberg, S. Schembari, N. Schleissman, C. Schmitt, S. Schulze, A. Scott, S. Sita, J. Shih, K.

335 Stewart, W. Tucker, E. Urquhart, K. van Atta, J. Vandermeer, L. van Zuidam, J. Verge, V. Vonau,

336 A. Walker-Bolton, E. Wikberg, E. Williams, L. Wolf, and D. Wood. We thank H. Gilkenson and W.

337 Lammers for long-term management of the field site during 2002-2013. We thank the Costa

338 Rican park service (Sistema Nacional de Áreas de Conservación and Área de Conservación

339 Tempisque), Hacienda Pelon de la Bajura, Hacienda Brin D'Amor, and the residents of San

340 Ramon de Bagaces for permission to work on their land. D. Cohen assisted with database

341 management and queries. This paper has benefited from helpful discussions with H. Clark

342 Barrett, Donald Cohen, Joseph Manson, Mary Brooke McElreath, Oliver Schülke, and Julia

343 Ostner.

REFERENCES

Archie, E. A., Tung, J., Clark, M., Altmann, J., & Alberts, S. C. (2014). Social affiliation matters:

Both same-sex and opposite-sex relationships predict survival in wild female baboons.

Proceedings of the Royal Society B: Biological Sciences, *281*(1793), 20141261.

Brent, L. J. N., Ruiz-Lambides, A., & Platt, M. L. (2017). Family network size and survival across

the lifespan of female macaques. *Proceedings of the Royal Society B: Biological Sciences*,

284(1854), 20170515.

Cameron, EZ, Setsaas, TH, & Linklater, WL (2009). Social bonds between unrelated females

increase reproductive success in feral horses. *Proceedings of the National Academy of*

Sciences , *106* (33), 13850-13853.

Clutton-Brock, T. H. (1988). *Reproductive success: Studies of individual variation in contrasting*

breeding systems. University of Chicago Press.

Crockford, C., Deschner, T., & Wittig, R. M. (2017). The role of oxytocin in social buffering: What

do primate studies add? In *Behavioral Pharmacology of Neuropeptides: Oxytocin* (pp.

155-173). Springer, Cham.

Dunbar, R. I. M. (2010). The social role of touch in humans and primates: Behavioural function

and neurobiological mechanisms. *Neuroscience & Biobehavioral Reviews*, *34*(2), 260–

268. https://doi.org/10.1016/j.neubiorev.2008.07.001

Fagen, R., & Fagen, J. (2004). Juvenile survival and benefits of play behaviour in brown bears,

Ursus arctos. *Evolutionary Ecology Research*, *6*(1), 89–102.

Farine, D. R. (2015). Proximity as a proxy for interactions: Issues of scale in social network

analysis. *Animal Behaviour*, *104*, e1–e5.

Fedigan, L. M., Carnegie, S. D., & Jack, K. M. (2008). Predictors of reproductive success in female

white-faced capuchins (Cebus capucinus). *American Journal of Physical Anthropology,*

137(1), 82–90.

Fragaszy, D. M., Visalberghi, E., & Fedigan, L. M. (2004). *The complete capuchin: The biology of*

the genus Cebus. Cambridge University Press.

Gilby, I. C., Brent, L. J., Wroblewski, E. E., Rudicell, R. S., Hahn, B. H., Goodall, J., & Pusey, A. E.

(2013). Fitness benefits of coalitionary aggression in male chimpanzees. *Behavioral*

Ecology and Sociobiology, *67*(3), 373–381.

Gros-Louis, J., Perry, S., & Manson, J. H. (2003). Violent coalitionary attacks and intraspecific

killing in wild white–faced capuchin monkeys (Cebus capucinus). *Primates*, *44*(4), 341–

346.

Hinde, R. A. (1976). Interactions, relationships and social structure. *Man*, 1–17.

Holt-Lunstad, J., Smith, T. B., & Layton, J. B. (2010). Social relationships and mortality risk: A

meta-analytic review. *PLoS Medicine*, *7*(7), e1000316.

Kalbitzer, U., Bergstrom, M. L., Carnegie, S. D., Wikberg, E. C., Kawamura, S., Campos, F. A., Jack,

K. M., & Fedigan, L. M. (2017). Female sociality and sexual conflict shape offspring

survival in a Neotropical primate. *Proceedings of the National Academy of Sciences*,

114(8), 1892–1897.

Koster, J., Leckie, G., Miller, A., & Hames, R. (2015). Multilevel modeling analysis of dyadic

network data with an application to Ye'kwana food sharing. *American Journal of Physical*

Anthropology, *157*(3), 507–512.

Koster, J., Leckie, G., & Aven, B. (2020). Statistical methods and software for the Multilevel

Social Relations Model. *Field Methods*, *32*(4), 339–345.

Manson, J. H., Navarrete, C. D., Silk, J. B., & Perry, S. (2004). Time-matched grooming in female

primates? New analyses from two species. *Animal Behaviour*, *67*(3), 493–500.

McElreath, R. (2020). *Statistical rethinking: A Bayesian course with examples in R and Stan*. CRC

press.

McElreath, R. (2020). rethinking: Statistical rethinking book package (R package Version 1.93).

https://github.com/rmcelreath/rethinking.

Nuñez, C. M., Adelman, J. S., & Rubenstein, D. I. (2014). Sociality increases juvenile survival after

a catastrophic event in the feral horse (Equus caballus). *Behavioral Ecology*, *26*(1), 138–

147.

Ostner, J., & Schuelke, O. (2018). Linking sociality to fitness in primates: A call for mechanisms.

In *Advances in the Study of Behavior* (Vol. 50, pp. 127–175). Elsevier.

Perry, S. (1996). Female-female social relationships in wild white–faced capuchin monkeys,

Cebus capucinus. *American Journal of Primatology*, *40*(2), 167–182.

Perry, S. (2012). The behavior of wild white-faced capuchins: Demography, life history, social

relationships, and communication. In *Advances in the study of behavior* (Vol. 44, pp.

135–181). Elsevier.

Perry, S., Godoy, I., & Lammers, W. (2012). The Lomas Barbudal Monkey Project: Two decades

of research on Cebus capucinus. In *Long-term field studies of primates* (pp. 141–163).

Springer.

Perry, S., Manson, J. H., & Perry, S. (2009). *Manipulative Monkeys*. Harvard University Press.

Porter, C. A., Page, S. L., Czelusniak, J., Schneider, H., Schneider, M. P. C., Sampaio, I., &

Goodman, M. (1997). Phylogeny and Evolution of Selected Primates as Determined by

Sequences of the ε-Globin Locus and 5' Flanking Regions. *International Journal of*

Primatology, *18*(2), 261–295. https://doi.org/10.1023/A:1026328804319

R Core Team. (2020). R: A language and environment for statistical computing. Vienna,

Austria: R Foundation for Statistical Computing.

435
Roelfs, D. J., Shor, E., Kalish, R., & Yogev, T. (2011). The rising relative risk of mortality for

singles: Meta-analysis and meta-regression. *American Journal of Epidemiology*, *174*(4),

379–389.

Schülke, O., Bhagavatula, J., Vigilant, L., & Ostner, J. (2010). Social bonds enhance reproductive

success in male macaques. *Current Biology*, *20*(24), 2207–2210.

Shor, E., & Roelfs, D. J. (2015). Social contact frequency and all-cause mortality: A meta-analysis

and meta-regression. *Social Science & Medicine*, *128*, 76–86.

Shor, E., Roelfs, D. J., & Yogev, T. (2013). The strength of family ties: A meta-analysis and meta-

regression of self-reported social support and mortality. *Social Networks*, *35*(4), 626–

638.

Silk, J. B. (2009). Nepotistic cooperation in non-human primate groups. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *364*(1533), 3243–3254.

Silk, J. B., Alberts, S. C., & Altmann, J. (2003). Social bonds of female baboons enhance infant

survival. *Science*, *302*(5648), 1231–1234.

Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R., Wittig, R. M.,

Seyfarth, R. M., & Cheney, D. L. (2010). Strong and consistent social bonds enhance the

longevity of female baboons. *Current Biology*, *20*(15), 1359–1361.

Silk, J., Cheney, D., & Seyfarth, R. (2013). A practical guide to the study of social relationships.

Evolutionary Anthropology: Issues, News, and Reviews, *22*(5), 213–225.

Snijders, T. A., & Kenny, D. A. (1999). The social relations model for family data: A multilevel

approach. *Personal Relationships*, *6*(4), 471–486.

Stan Development Team. (2020). RStan: The R interface to Stan (Version 2.19.1).

Sterck, E. H. M., Watts, D. P., & van Schaik, C. P. (1997). The evolution of female social

relationships in nonhuman primates. *Behavioral Ecology and Sociobiology*, *41*(5), 291–

309. https://doi.org/10.1007/s002650050390

Thompson, N. A. (2019). Understanding the links between social ties and fitness over the life

cycle in primates. *Behaviour*, *1*(aop), 1–50.

Thompson, N. A., & Cords, M. (2018). Stronger social bonds do not always predict greater

longevity in a gregarious primate. *Ecology and Evolution*, *8*(3), 1604–1614.

Yee, J. R., Cavigelli, S. A., Delgado, B., & McClintock, M. K. (2008). Reciprocal affiliation among

adolescent rats during a mild group stressor predicts mammary tumors and lifespan.

Psychosomatic Medicine, *70*(9), 1050.

468