Fitness benefits of providing services to others: Sociality predicts survival in a

neotropical primate

Kotrina Kajokaite*[a], Andrew Whalen[b], Jeremy Koster[a,c], and Susan Perry[d,e]

a. Department of Human Behavior, Ecology and Culture, Max Planck Institute for

Evolutionary Anthropology, Leipzig, Germany

b. The Roslin Institute and Royal (Dick) School of Veterinary Studies, The University of

Edinburgh, Edinburgh, UK

c. Department of Anthropology, University of Cincinnati, Cincinnati, OH, USA

d. Department of Anthropology, University of California, Los Angeles, CA, USA

e. Behavior, Evolution and Culture Program, Department of Anthropology, University of

California, Los Angeles, USA

* Corresponding author: Kotrina Kajokaite, Department of Human Behavior, Ecology and

Culture, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6 04103

Leipzig, Germany

ABSTRACT

Across multiple species of social mammals, evidence is accumulating that

sociality is associated with fitness. In long-lived species, like primates, lifespan is one of

the main fitness components. Here, we provide the first direct evidence that females

who provide more in social services live longer. We used 18 years of data from the

Lomas Barbudal Monkey project to quantify sociality in 11 capuchin (*Cebus capucinus*)

social groups using three interaction types: grooming, support in coalitionary

aggression, and foraging in close proximity. We analyzed each domain separately to

understand how giving and receiving social favors impact survivorship. To estimate

female sociality in each of these domains, we built an adaptation of the Social Relations

Model. This approach enabled us to estimate individual rates of giving and receiving

interactions and take into account the uncertainty in these estimates due to varying

amounts of observations of each individual. Subsequently, we modeled adult females'

survival as a function of their sociality estimate, rank, age, group size, and maternal kin

presence using a Bayesian Cox proportional hazards model. We found that females who

provide more grooming to other group members survive longer. There was no strong

evidence that females who receive more grooming, participate more often in

coalitionary aggression, and forage more often in close proximity of others also have

higher survivorship. These results add a neotropical primate species to the growing list

of mammals, including humans, where greater sociality is associated with longevity.

    INTRODUCTION

A key question for understanding the evolution of animal sociality is: Do animals

that are more social enjoy greater fitness benefits? A number of studies link

components of reproductive success, such as fecundity (white-faced capuchins, *Cebus*

capucinus: Fedigan et al., 2008; Assamese macaques, *Macaca assamensis*: Schülke et

al., 2010; chimpanzees, *Pan troglodytes schweinfurthii*: Gilby et al., 2013) and offspring

survival (yellow baboons, *Papio cynocephalus ursinus*: Silk, 2009; savannah baboons,

Papio cynocephalus: Silk et al., 2003; white-faced capuchins, *Cebus capucinus:* Kalbitzer

et al., 2017), to differences in the social behavior of individuals.

In long-lived species, lifespan is also an important component of reproductive

success contributing to fitness variation (Clutton-Brock, 1998). The most abundant

evidence that individual sociality affects lifespan has accumulated in human literature.

Both the presence and the absence of social relationships affects lifespan. That is, being

socially connected is associated with a lower probability of dying (Holt-Lunstad et al.,

2010; Shor and Roelfs, 2015; Shor et al., 2013), while having few and poor social

relationships is associated with a higher risk of death (Roelfs et al., 2011).

In nonhuman species there is similar evidence that individual sociality is

associated with lifespan. Among baboons, females who maintain stable social

relationships with other males and females live longer (chacma baboons, *Papio*

hamadryas ursinus, Silk et al., 2010; Archie et al., 2014). Prime-aged female macaques of

Cayo Santiago who have more living adult female relatives live longer (Brent et al.,

2017). Blue monkey females (*Cercopithecus mitis stuhlmanni)* who have strong and

enduring social relationships consistently with the same partners tend to live longer in

comparison to females whose strong partnerships are inconsistent across years

(Thompson and Cords, 2018). Female rats who cultivate reciprocal affiliations with their

sisters survive longer (Yee et al., 2008), and brown bear cubs who play longer during

summer survive better (*Ursus arctos*) (Fagen and Fagen, 2004). More social juvenile feral

horses (*Equus caballus*) live longer than their less social peers. However, facultatively

social yellow-bellied marmots (*Marmote flaviventer*) show the opposite pattern: more

social marmots live shorter lives than less social marmots (Blumstein et al., 2018).

Neotropical primates are not represented among the studies that report the link

between sociality and longevity. Capuchin monkeys are an excellent neotropical primate

species for the investigation of these associations: Despite the long evolutionary history

that separates them from humans and cercopithecines (Porter et al., 1997), their social

structure is similar to cercopithecine primates (Perry, 2012).

In this paper, we investigate whether more social white-faced capuchin females

live longer than less social females. We analyzed eighteen years of capuchin monkeys'

social interactions taken from the Lomas Barbudal Monkey dataset. To quantify

individual sociality, we selected interactions from three domains to represent the

variety of social behaviors that capuchin monkeys engage in: grooming, coalitionary

aggression, and foraging in proximity to others. To deal with different amounts of

observations for each individual, we used the multilevel Social Relations Model (Snijders

and Kenny, 1999; Koster et al. 2019) to decompose the observations of exchanges of

these interactions into separate individual giving, individual receiving, and dyadic

relationship components. This approach allowed us to estimate the uncertainty in

sociality estimates, which can be substantial for individuals with small numbers of

observations. Then, we used the individual giving and individual receiving posterior

means and standard deviations (SD) from these models as sociality measures to model

adult female longevity using the Bayesian Cox proportional hazards model.

We found that adult female capuchins provide most of the grooming in the

population. However, they display considerable variation in their propensity for

grooming others, and some adult females are among the least frequent groomers in the

population. The variation in *giving grooming* measures predicts longevity: Females who

provide more grooming live longer than the females who provide less grooming.

Measures in other interactions – *receiving of grooming*, *giving* and *receiving of*

coalitionary support, and *foraging in close proximity* – were similar in their distribution

to the rest of the population. In addition, females were more similar to each other in

these interactions, especially in how much grooming they received, i.e. the females who

were the most and the least likely to be groomed were not very different from an

average monkey in the population. The females who engaged more in these interactions

seemingly experienced some benefits associated with longevity, but the model

estimates were uncertain. Therefore, the evidence does not support a strong

association between longevity and the sociality measures we considered (*grooming*

receiving, participating in coalitionary support, and *foraging in close proximity of*

others).

METHODS

Study subjects and the dataset

We studied members of the wild white-faced capuchin population at the Lomas

Barbudal Biological Reserve and surrounding private lands in Guanacaste, Costa Rica

(Perry et al., 2012). The Lomas Barbudal Monkey Project dataset has longitudinal

records on individuals living in 11 capuchin social groups. It contains detailed records

including demographic information, pedigree information, and data regarding social

interactions that are necessary for quantifying individual sociality.

White-faced capuchins live in multimale, multifemale social groups (Perry, 2012).

Females are philopatric – they stay in their natal groups for their entire lives – which

allows researchers to collect data about the social interactions from large portions of

their lives (Fragaszy et al., 2004; Perry et al., 2008). As a result, females' demographic

and behavioral records are more complete than males' records because males transfer

to different groups multiple times in their lives and often leave the study area. This

makes female capuchins more suitable subjects for the investigation of the relationships

between individual sociality and survival.

The records on capuchin behavior were collected between January 2002 and

December 2019. The subjects were 132 adult females, living in eleven different social

groups over 146 group years (average years of observation per group = 13.27; range = 7

– 18 years). The females were considered adults once they reached the age of 5 years.

The behavioral and demographic data on each group were collected by experienced

observers during censuses lasting at least 6 hours/day.

Measuring Sociality

To measure female sociality, we chose three behavioral domains: grooming,

coalitionary aggression, and foraging. We treated grooming and coalition formation as

directed behaviors, and we used observations of individuals as both initiators and

recipients of the behavior. We did not have information about which individual had

initiated the proximity when foraging, and therefore foraging in proximity was treated

as an undirected behavior. In calculating the frequency with which adult females

engaged in these interactions, we used behavioral records from all the individuals who

resided in the 11 groups during the study period. There were a total of 563 monkeys,

residing in 11 social groups, comprising 13,770 dyads. We treated each of the five

interaction types (*grooming giving, grooming receiving, support giving, support*

receiving, forage in proximity) as a separate measure of sociality.

Grooming (groom giving and groom receiving)

Allogrooming, or social grooming, is a common behavior in capuchin monkeys

(Perry, 1996). Grooming serves a hygienic and social bonding function and is thought to

impose costs on groomers of loss of time and increased vigilance (Manson et al., 2004).

These features of grooming behavior make it a good candidate for providing information

regarding an individual's sociality.

Grooming rates were estimated using data collected during 10-minute focal

follows. To estimate individual grooming rates, we calculated dyadic counts of grooming

and dyadic opportunities for grooming. The opportunity for a dyad, A-B, to engage in

grooming, was calculated as the sum of the focal follows of A and the focal follows of B

at times when A and B were residents of the same social group. The dyadic counts of

giving and receiving grooming were assigned as follows: a count of 1 was assigned if A

groomed B at least once during a focal follow, otherwise 0 was assigned. The same was

done when evaluating if B groomed A.

Joining a coalitionary conflict (support giving and support receiving)

Coalitionary aggression is frequent in capuchin monkeys (Perry, 2012).

Intergroup encounters are characterized by high rates of coalitionary lethal aggression

(Gros-Louis et al., 2003), and coalitions play an important role in male competition for

reproductive opportunities (Perry, 2012). Intragroup conflicts are far less dangerous;

they often do not result in physical fighting, and individuals are rarely injured (Perry,

2012). Capuchins have stereotyped signals for recruiting allies in conflicts (Perry, 2012),

and participation in coalitionary aggression is likely based on social relationships

between the partners (Kajokaite et al., 2019).

The behavior of joining a coalitionary conflict was defined as an individual

intervening on one side during an ongoing aggressive conflict. This definition only

indicates the functional aspect of joining a side; it entails no inferences about internal

psychological states such as the intent to help a specific individual. Since aggressive

interactions are salient and harder to miss than quiet activities like grooming or foraging

in proximity, aggressive interactions were collected both *ad libitum* and during focal

follows. The chronological stream of aggressive behaviors was divided into 5 min.

segments. In order to identify instances of joining a coalitionary conflict, monkey A is

identified as joining monkey B if A performed an aggressive behavior toward either

monkey B's opponent or victim within the context of the 5-minute segment. Even if

multiple instances of A joining monkey B were observed during the 5 min segment, we

scored only one instance of A joining B. To calculate the opportunities to join a

coalitionary conflict, all individuals who were co-resident during the aggressive conflict

were counted as having had an opportunity to join on either side during the conflict.

Foraging in proximity

Although foraging is a seemingly asocial behavior, the way individuals

distribute themselves across space can be indicative of their social relationships with

group members. It can be costly to tolerate another individual near you (more

competition for food) and monkeys might only tolerate proximity of individuals with

whom they maintain good social relationships.

Foraging in close proximity was estimated from group scans that occurred in

the context of foraging. In group scans, the identity of the scanned individuals, their

activity and their proximity to other individuals within 10 body lengths (~2 m) was

noted. We considered individuals to be foraging in close proximity if they were scanned

within 5 body lengths (~1m) of each other. For each dyad, we scored whether they were

observed foraging within close proximity in 10 min. segments (i.e. if they were observed

doing this more than once within 10 min., only one instance was counted). The number

of opportunities that the dyad had to forage within close proximity is a sum of group

scans in the foraging context that are 10 min. apart, where one of the individuals is a

subject of a group scan.

The Social Relations Model and Individual sociality measures

Detailed longitudinal behavioral records are necessary for relating an individual's

social behavior patterns to longevity (Clutton-Brock, 1988). Such data is hard to collect

and resources are limited; therefore the more individuals we observe, the less

information we collect on each one of them. The data for these analyses were collected

across eighteen years and the number of social groups and individuals observed

increased over time. As a result, data density is uneven across time periods, social

groups, and individuals.

We incorporated uneven distribution of the data into our analyses as follows.

First, we aggregated the data annually and used adaptations of the multilevel Social

Relations Model (Snijders and Kenny, 1999; Koster et al. 2019) to estimate individual

annual rates of grooming, coalitionary support, and foraging. This provided us with

mean and standard deviation of individual sociality measures (see the supplemental

material). Second, we used these distributions of individual sociality measures as

predictors of longevity and to investigate the effects of social behavior on survivorship.

The Social Relations Model (Snijders and Kenny, 1999) decomposes the variance

in giving and receiving behaviors into separate giving, receiving, and dyadic relationship

components. We fitted a Bayesian multilevel Social Relations Model with the following

parameters: the intercept parameter (base rate), social group-level, individual giving,

individual receiving, dyadic symmetric, and dyadic asymmetric random effects

parameters (see Koster at al. 2015). We used data collected on the entire population

(i.e. adult males, females, and immatures) since the Social Relations Model is a

multilevel model and can take advantage of additional data to estimate the average

rates of interactions in the population (McElreath, 2020). Although we estimated the

random effects for all the individuals in the population, we used the adult female annual

individual posterior mean and SD of *grooming giving, grooming receiving, support*

giving, support receiving, and non-directional *foraging in proximity* rate estimates as our

five measures of annual individual sociality.

226   The posterior estimates of mean and SD of female giving and receiving random

effects capture the extent to which an individual female deviates from the estimated

average rate of engaging in this behavior in the population. The individual sociality

measures are centered on zero, which represents the population mean (an average

monkey). Each behavior domain has its own population average estimate. For example,

the individual estimates of *grooming giving* ranges from –3.14 to 3.27 across the entire

population. For a particular individual, a posterior mean closer to the minimum (–3.14)

indicates that this individual grooms others much less than the average monkey grooms

her partners in the population. The inverse is true for a posterior mean estimate close to

the maximum of the range (3.27). The SD reflects the measurement uncertainty from

heterogenous sampling effort, where individuals who were observed less have posterior

estimates with wider distributions around the posterior mean estimate.

Modeling survival as a function of individual sociality measure

239   To investigate whether sociality is associated with adult female longevity, we

used Bayesian Cox proportional hazards models. In separate models, each of the five

individual sociality measures was modeled as a predictor of survival probability over

one-year periods. These models included the following time-varying (calendar year-

specific) covariates: the female's age, her dominance index, the average number of

individuals in her group, the proportion of time during that year that her mother was

alive, and the number of adult daughters that she had.

Age: Focal female ages were assigned based on demographic records of births

and deaths collected from 1990 – 2019. The ages of females who were born before

1990 were estimated in part by retroactively comparing photos taken then with photos

of known-aged females collected later in the study. In addition, we inferred

reproductive histories via genetic maternity data. We assumed that the age of first birth

for each mother was six years, and each mother had a two-year interbirth interval

(Fedigan and Rose, 1995; Perry, 2012).

Average number of individuals in her group: This is the mean number of adult

females, adult males, and immatures that resided in the female's group during the days

when researchers spent at least six hours of observation with the group, averaged for

the year.

Mother's presence: The proportion of the year that the female's mother was

alive and co-resided with her. The measure varies from 0 to 1, where 1 indicates that

the mother was alive and co-resident with the focal female for the entire year, and 0

indicates that the mother died in some previous year.

Number of daughters: The number of daughters that a focal female had was

very highly correlated with her age (0.72). To control for the effect of age on the

number of daughters, we grouped all of the females that were the same age together

and centered the number of daughters for each age group. For every age group we

subtracted the mean number of daughters for that age group to reduce the confounding

between age and number of daughters. The resulting variable was not correlated with

age (~0). This is equivalent to modeling an interaction between age and number of

daughters, with age treated as a categorical variable.

Annual dominance index: The annual dominance index represents the proportion

of group members that the female dominated, on average, that year. For each

observation day that the female (the focal) resided in a social group, we identified all of

the other co-resident individuals (alters). To assess whether the female was dominant to

an alter on a particular day,

we first identified the dominance interaction immediately preceding and following the

day of interest for each focal-alter dyad. The total number of these interactions across

all focal–alter dyads represents the number of opportunities for the focal to dominate

someone. For each interaction, we identified that the focal individual was dominant if

she was either the animal performing the supplanting, or being cowered at, or being

avoided, or fled from. The daily dominance index, DDI_i, of a focal individual, i, is a sum of

dominance interactions where focal was dominant to their alters, w_{i-a}, divided by the

total number of dominance interactions that the focal had with her alters, s_{i-a}:

$$DDI_i = \frac{\sum w_{i-a}}{\sum s_{i-a}}.$$ (1)

Then, the average annual dominance index, ADI_i, is an average of daily dominance

indices:

$$ADI_i = \frac{1}{n} \sum_{i=1}^{n} DDI_i \,.$$
(2)

In some cases, either one or no dominance interactions were available for a focal-alter

dyad. As a result, the individuals who did not have dominance interactions with the

focal did not contribute to the calculation of the daily dominance index.

We included the covariates, because each of them is likely to influence

females' probability of survival. An adult female's probability of dying in a given year is

expected to increase with age (Archie et al. 2014; Brent et al. 2017). Females who live in

larger groups might experience greater survival rates due to reduced predation risks

(Sterk et al., 1997). Higher-ranking females possibly enjoy reduced mortality in

comparison to lower-ranking females due to their central position in the group

(Kalbitzer et al., 2017), which also reduces risks of predation. The presence of a mother

and/or adult daughters approximate female's close kin network benefits: Females who

have a more extensive kin network might experience reduced mortality (Brent et al.,

2017).

Having many covariates in the model can result in "included variable bias"

(McElreath, 2020) where predictors are not only causally influencing the outcome, but

also influencing each other. This can result in predictor induced statistical selection

within the model and manifest itself through misleading statistical, but not causal,

associations between the variables we are interested in (McElreath, 2020). A female's

rank and age are likely to influence how much the female participates in social

interactions. Females who reside in larger groups are likely to have different interaction

networks in comparison to females in smaller social groups due to a larger number of

social partners available (Archie et al., 2014). To check if the above assumed

relationships justify the inclusion of all of the predictor variables in the model, we drew

a directed acyclical graph (DAG) using the package *daggity* (v.3.0) and analyzed implied

functional relationships. The DAG assumed is provided in the supplemental materials.

Modeling approach

To take into account that some females contributed different numbers of

years to the analysis, we modeled individual differences using a random effect term.

Models were run using Stan (v.2.19.1) and the *rethinking* package (v. 1.93: McElreath

2020) in *R* (v. 3.6.2; R Core Team 2019).

More formally, we specified the following model for the number of days

before death, D_i,. The probability for the number of days before death come from the

cumulative probability distribution:

$$\Pr\left(D_i \mid \lambda_i\right) = \lambda_i \exp(-\lambda_i D_i)\,. \tag{3}$$

For females who did not die during the observation period, the probability of waiting D_i

without dying comes from the complementary cumulative probability distribution:

$$\Pr\left(D_i \mid \lambda_i\right) = \exp\left(-\lambda_i D_i\right)\,. \tag{4}$$

We model the rate of dying, λ_i, as follows

332
$$\lambda_i = 1/\mu_i . \tag{5}$$

where μ_i is the expected number of days till death

$\log(\mu_i) = \alpha + a[\,id\,] + b_{sociality} * true\ sociality\ estimate[id] + b_{rank} * rank_s \tag{6}$

$+ b_{age} * age + b_{daughters} * daughters$

$+ b_{mother} * mother + b_{grsize} * group\ size .$

α denotes the intercept or the base rate of number of days survived, $\alpha[id]$ denotes

individual female random effects corresponding to the observation period. The model

coefficients $b_{sociality}$, b_{rank}, b_{age}, $b_{daughters}$, b_{mother}, b_{grsize} describe the impact of

sociality, rank, age, number of daughters, mother's presence, and group size,

respectively. We took a latent variable approach to model the sociality estimates, since

individual sociality estimates are not point estimates, but rather posterior distributions

with mean and SD reflecting measurement uncertainty about the true rate. Under this

approach, we assumed that each female had some true underlying sociality measure,

true sociality estimate[id], which was normally distributed with mean and SD equal to

the mean and SD of the individual estimate of sociality drawn from the Social Relations

Model results.

We used the Bayesian approach to fit Cox proportional hazard model (Singer

and Willett 2003). We assumed Normal (8, 0.5) prior for a base rate of survival, α, which

places most of the prior mass between 0 to 20 years with the mean of 8 years and a

long tail allowing more extreme values. For fixed effects, we assumed Normal (0, 1)

prior. For individual-level random effects, $\alpha[id]$, we compared three types of models:

one with no random effects and two with individual-level random effect priors assumed

as Normal (0, 0.2) or Normal (0, 1). All three model types resulted in similar estimates;

therefore here we are reporting the results with individual-level random effect prior

Normal (0, 1), while the results of the other two types of models are reported in the

supplemental material (Table S1 and Table S2). All of the covariates, except the number

of daughters, were standardized by subtracting the mean and dividing it by the standard

deviation. As noted previously, the number of daughters was centered by subtracting

the mean number of daughters for each age.

RESULTS

(1) Measures of sociality: How do adult females compare to the rest of the population?

Females differ the most from the other age-sex classes in their distribution of

grooming giving measures. A majority of the females (80%) groom more than an

average monkey in the population, while a majority of the adult males and immatures

(78%) groom less than an average monkey (Figure 1, panel A). The female distributions

for the rest of the behavioral domains are very similar to the other age-sex distributions,

indicating that females behave similarly to adult males and immatures in those domains.

Figure 1 illustrates how the individual sociality measures of *grooming,*

coalitionary support, and *foraging in proximity* by adult females compare to the

estimates for other age-sex classes. The distribution in orange represents adult female

posterior means estimated by the Social Relations Model, while the distribution in white

represents a combination of adult males and immature capuchins. The zero represents

an average individual in the population.

Grooming behavioral domain distributions show both the greatest and the

smallest amount of variation. Females display the greatest range in *grooming giving*

posterior estimates (*grooming giving* variance=2.02), and the narrowest range in

grooming receiving posterior estimates (*grooming receiving* variance=0.12). The

difference in variation suggests that the most avid groomers and the least enthusiastic

groomers differ a lot from an average monkey. However, as recipients of grooming,

both females who receive the most and the least amount of grooming are much more

similar to an average monkey in the population in comparison to any other type of

interaction that we measured.

Figure 1. Posterior mean distributions of annual individual sociality measures for each type of behavior. The orange density represents adult females whereas the white density represents the rest of the population. The population base rate is at approximately zero.

(2) Measure of sociality: Do females behave similarly across the behavioral domains?

Each female has five annual sociality measures. We plotted the pairwise

correlations between each of the sociality measures in Figure 2 in order to investigate if

female profiles are similar across all five measures, i.e. is a frequent groomer also a

frequent recipient of grooming and coalitional support? Within the coalitionary

aggression domain, estimates of female giving and receiving coalitionary support were

highly correlated (r=0.97). Females who intervene on someone's side during a conflict

also tend to receive a lot of support from others. This suggests that there is a small

subset of females that participate in coalitionary aggression more frequently than

others. However, females vary less in their *support receiving* than in their *support giving*

estimates (Figure 1, Panel C–D). In the grooming domain, the correlation between giving

and receiving estimates was moderate (r=0.57). Some of the females who groom others

the most are not among those who receive the most grooming. In addition, Figure 1

panel B shows a narrow distribution of the receiving measurements, which implies that

females overall distribute their grooming evenly among their group members.

The correlation across behavioral domains is smaller than within a domain,

between 0.15 and 0.37: If a female is among those who engage a lot in coalitionary

aggression, she is not necessarily involved a lot in grooming interactions and in foraging

in close proximity. The pairwise correlations show that females have different profiles of

interactions depending on the behavioral domain.

Figure 2. Bi-variate correlations of individual sociality measures. Each point represents
an annual observation of an individual female.

(3) Do more social females survive better?

Table 1 presents the Cox proportional hazards model posterior mean estimates

and the 95% Highest Posterior Density Interval (HPDI), representing the narrowest

interval containing the 95% probability mass. Independent of the effects of covariates

(age, dominance rank, group size, number of adult daughters, co-residence with

mother), females who groomed others more survived at higher rates than other

females. The other sociality measures had mean estimates consistent with sociality

predicting greater survival, but their HPDI intervals were wide and included zero

indicating that the models are uncertain about these estimates.

Table 1. Estimates of fixed effects of each of the Cox proportional hazards models:
posterior means and 95% HPDI.

	Sociality measure				
Parameter	Grooming giving	Grooming receiving	Support giving	Support receiving	Foraging in proximity
Intercept	9.20 [8.86,9.56]	9.13 [8.78,9.52]	9.07 [8.74,9.42]	9.06 [8.73,9.41]	9.08 [8.74,9.42]
β sociality	0.49 [0.26,0.70]	0.19 [−0.09,0.45]	0.11 [−0.21,0.45]	0.06 [−0.28,0.40]	0.16 [−0.18,0.47]
β rank	−0.20 [−0.51,0.11]	−0.11 [−0.45,0.24]	−0.10 [−0.45,0.24]	−0.10 [−0.47,0.28]	−0.16 [−0.54,0.23]
β age	−0.57 [−0.88,−0.25]	−0.66 [−0.96,−0.37]	−0.66 [−0.96,−0.37]	−0.68 [−0.99,−0.38]	−0.69 [−0.98,−0.39]
β daughters	−0.08 [−0.45,0.28]	−0.08 [−0.45,0.31]	−0.07 [−0.44,0.31]	−0.06 [−0.43,0.30]	−0.05 [−0.43,0.32]
β mother	0.26 [−0.09,0.63]	0.31 [−0.02,0.66]	0.33 [−0.01,0.67]	0.32 [−0.01,0.65]	0.33 [−0.01,0.66]
β group size	0.19 [−0.1,0.49]	0.18 [−0.10,0.46]	0.19 [−0.10,0.48]	0.18 [−0.1,0.46]	0.18 [−0.09,0.46]

442   We plotted the posterior means and their standard errors across five Cox

443   proportional hazard models in Figure 3 to better illustrate how model estimates

444   compare across all five models. Apart from sociality, all of the covariates have mean

445   estimates with wide HPDI intervals that include zero. The only exception is age,

446   indicating that female survival declines with age.

Figure 3. Posterior means and standard deviations of 5 Cox proportional hazards models
in which annual survivorship varied as a function of five measures of sociality (*foraging*
in proximity, support receiving, support giving, grooming receiving, and *grooming giving*)
and covariates.

To facilitate the interpretation of the Cox proportional hazards model

coefficients, we plotted model predictions showing the probability of dying in a given

year (Figure 4). We have varied females' sociality measure to compute the predictions.

Figure 4 illustrates the annual probability of dying as a function of a female's sociality

level measure where a yellow line represents a 10-year-old female and a blue line

represents a 30-year-old female (holding all other covariates fixed at the values

described above).

Across all five sociality measures, the less social females have a greater

probability of dying than more social females, but only the slopes for *groom giving*

confidently suggest a beneficial effect on survivorship (Figure 4, panel A). Predictions for

groom receiving likewise suggest a beneficial effect of sociality, but the model

predictions are less certain (Figure 4, panel B). Meanwhile, the slopes for *coalitionary*

*suppo*rt and *foraging in close proximity* are quite shallow, indicating that the survival

varies little as a function of these sociality measures (Figure 4, panels C, D, and E). These

model predictions reflect the uncertainty associated with these estimates (Table 1 and

Figure 3). Overall, there is less certainty in the estimates for older females, partly owing

to their scarcity in the sample.

478　**Figure 4.** Cox proportional hazard model predicted annual probabilities of dying at two
479　representative ages (10 and 30) as a function of the sociality levels in five behavioral
480　domains. For both ages, we assumed median group size, median rank, and average
481　number of daughters for the age. For 10-year-old females, we assumed a co-resident
482　mother, while predictions for the 30-year-old assume the mother is deceased.
483

DISCUSSSION

In female white-faced capuchin monkeys, providing grooming to others is

associated with greater survival rates. Females who provide more grooming die at lower

rates than females who engage in this behavior less often. By contrast, there was no

strong evidence that females who receive more grooming, participate more often in

coalitionary aggression, and forage more often in close proximity of others also have

higher survivorship. Our results in the grooming domain are consistent with findings

that social integration is associated with longevity in humans (Holt-Lunstad et al., 2010)

and adds a neotropical primate species to the list of mammalian species where a similar

association between sociality and longevity has been demonstrated (Silk et al., 2010;

Archie et al., 2014; Brent et al., 2017; Yee et al., 2008; Nuñez et al., 2014; Fagen &

Fagen, 2004).

Despite great interest by primatologists in studying social relationships and their

impact on primate lives, there is no established consensus on how to measure the

properties of social relationships in primates (Silk et al., 2013). The conceptual

foundation for studying primate social relationships dates back to Robert Hinde's

seminal paper (1976), where he argued that analyses of multiple types of interactions

should be included in measuring social relationships. However, the details of how to

combine these interactions into a meaningful measure have been under-theorized.

There are some well-established measures that combine the rates of behaviors from

different domains into a single measure of sociality (reviewed in Silk, 2013), but the

implications of averaging over multiple rates of different behaviors have not been

analyzed.

We do not suggest that combining information about interactions from multiple

domains should not be our goal in principle, but we want to draw attention to the idea

that doing so imposes inferential tradeoffs. For example, we demonstrated that the

distributions of the rates for each interaction type are very different (Figure 1) and, for

example, the females who frequently groom others are not always the same females

who frequently participate in coalitionary aggression (Figure 2). Averaging estimates

across all five behaviors would result in the same point estimates for females who have

very different profiles. For example, it is possible that a female who is an avid groomer

and seldom participates in coalitionary aggression through simple averaging could

appear identical to a female who rarely grooms others, but who often joins coalitionary

aggression. The interpretation of such measures is problematic because sociality indices

that draw on multiple behaviors may conceal important differences between

individuals. We suggest that it is useful to understand the variation of these behaviors

within a population before constructing a single measure.

We selected three behavioral domains, each of which represents a different

type of social engagement in capuchins – grooming, coalitionary conflicts, and foraging –

and estimated the individual sociality measures separately. We argue that doing so: (1)

enabled us to incorporate the uncertainty about our measures, and (2) aided our

interpretation of the results.

Incorporating the uncertainty. One of the challenges in analyzing observational

longitudinal datasets is data resolution as a result of uneven sampling. Data from the

natural habitats are notoriously difficult to gather. Inevitably, the records for social

interactions are sparse (Farine 2015) and some individuals are observed more often

than others for multiple reasons (Silk et al., 2013). Analysts therefore face a high risk of

false negatives when estimating dyadic relationships, i.e. some pairs are inferred to

have no social relationship at all, whereas if behavioral sampling were sufficiently dense,

they would be observed to interact, although rarely (Farine, 2015).

To address the uneven sampling of individuals and social groups in our dataset,

we used a Social Relations Model (Snijders and Kenny, 1999). This approach allowed us

to estimate individual rates of giving and receiving interactions and the uncertainty

associated with those estimates due to the varying number of observations. To increase

data resolution, we did not focus on adult female interactions with only other adult

females (e.g. Silk et al., 2010; Kalbitzer et al., 2017; Archie et al., 2014) or with only adult

males (e.g. Archie et al., 2014). We used to use each female's entire observation history

with her group members to ensure as complete as possible a picture of a female's true

sociality (Farine and Whitehead, 2015). As a multilevel model (McElreath 2020), the

Social Relations Model takes advantage of partial pooling across the entire population

when estimating individual female estimates (McElreath, 2020), which reduces

overfitting on the available data. To reflect the latent uncertainty in observational data,

our sociality measures were modeled as distributions, rather than point estimates,

which we subsequently used to model female survival.

Interpretation of results. Our analysis reveals several insights about grooming,

coalitionary support, and foraging interactions by showing how adult females compare

to each other and to the rest of the population (Figures 1–2). The most informative

behavior is grooming. Adult females are the most avid groomers in the capuchin

population with the majority of the females grooming more than an average monkey.

However, some of the individuals who groom the least are also adult females.

Knowing how an individual female's measures in one behavior correlate with

other behaviors that same year provides additional insights into the heterogenous

effects that the respective sociality measures exhibit on survivorship. As noted,

grooming giving measures are reliably associated with longevity. Yet, because *grooming*

giving is only weakly correlated with *support in coalitionary aggression* and *foraging in*

close proximity, females who are providing the most grooming are usually not the same

females who participate most often in coalitionary aggression or most often foraging in

close proximity of others.

Within the grooming domain, our results make sense in light of the respective

grooming distributions (Figure 1). *Grooming giving* and *grooming receiving* are

moderately correlated ($r = 0.57$, Figure 2), suggesting that many females occupy a

similar position relative to other females within the grooming network when quantifying

their grooming rates with other group members. However, there is substantially more

variation in how much grooming females provide, i.e. the most and the least avid

groomers are much more different from an average monkey in the population than the

females who receive the most and the least grooming. The narrow distribution of groom

receiving estimates suggests that grooming is somewhat equally distributed within the

population and that individuals do not differ from each other a lot in how likely they are

to be groomed by a group member. Unsurprisingly, these relatively small differences are

not reliably associated with survival.

Finally, a biologically interesting question deals with the mechanisms that

facilitate the relationship between sociality and longevity. Recently there has been a call

for understanding exactly how sociality is affecting fitness in primates (Ostner and

Schülke 2018; Thompson, 2019). Individual sociality as a construct is likely to be

composed of the many types of relationships that an animal has: affiliative, agonistic

and others. The lifespans that we observe are a consequence of many decisions that

animals make and there is arguably more than one pathway through which sociality can

affect survival in a particular species (Ostner & Schülke, 2018; Thompson, 2019).

Behaviors such as grooming, coalitionary support and foraging in proximity of others

indicate social relationships that individuals cultivate, but are probably not the sole

indicators of those relationships.

Our study was not designed with the purpose of identifying a mechanism linking

sociality and longevity. It is possible that females derive direct benefits from grooming

others, such as social bonding through the release of hormones (Dunbar 2010).

Indirectly, grooming interactions might contribute to structuring a female's overall social

life. Capuchin monkeys spend a lot of time grooming, and providing grooming to

multiple partners might allow females to develop social relationships which eventually

structure other types of interactions. For example, the distributions of coalitionary

aggression interactions suggest that a small group of individuals participate in

coalitionary aggression because female estimates of giving and receiving support are

very highly correlated (Figure 2). Since the females who groom others the most are

usually not the same females who participate in coalitionary aggression the most, it is

possible that the frequent groomers experience less aggression and, as a result, fewer of

the detrimental consequences associated with it. Alloparenting might be another

domain where frequent groomers receive more help from other group members. To

better understand the mechanisms that contribute to the relationship between sociality

and longevity and to design studies to test them, we need to investigate the interaction

patterns themselves. Understanding the natural structure of different types of

interaction types can help in developing better sociality measures and further our

understanding how sociality contributes to fitness.

DATA AND CODE ACCESSIBILITY

The data and code used to produce the analyses in this paper are available at:

www.doi.org/10.5281/zenodo.3961479

ACKNOWLEDGEMENTS

The following field assistants contributed significant amounts of data to the Lomas

Barbudal Monkey Project dataset: C. Angyal, A. Autor, B. Barrett, L. Beaudrot, M.

Bergstrom, R. Berl, A. Bjorkman, L. Blankenship, T. Borcuch, J. Broesch, D. Bush, J. Butler,

F. Campos, C. Carlson, A. Cobden, M. Corrales, J. Damm, B. Davis, C. deRango, N. Donati,

C. Dillis, G. Dower, R. Dower, A. Duchesneau, K. Feilen, J. Fenton, S. Fiello, K. Fisher, A.

Fuentes J., M. Fuentes A., T. Fuentes, C. Gault, H. Gilkenson, I. Godoy, I. Gottlieb, J.

Griciute, L.M. Guevara R., L. Hack, R. Hammond, S. Herbert, C. Hirsch, M. Hoffman, A.

Hofner, C. Holman, J. Hubbard, S. Hyde, M. Jackson, O. Jacobson, E. Johnson, L. Johnson,

S. Lee, S. Leinwand, T. Lord, M. Kay, E. Kennedy, D. Kerhoas-Essens, E. Johnson, S.

Kessler, B. Krimmel, W. Lammers, S. Lopez, S. MacCarter, J. Mackenzie, M. Mayer, F.

McKibben, W. Meno, A. Mensing, M. Milstein, C. Mitchell, Y. Namba, A. Neyer, C.

O'Connell, J. C. Ordoñez, J., N. Parker, B. Pav, R. Popa, K. Potter, K. Ratliff, K. Reinhardt,

N. Roberts Buceta, E. Rothwell, J. Rottman, H. Ruffler, S. Sanford, M. Saul, E. Seabright, I.

Schamberg, S. Schembari, N. Schleissman, C. Schmitt, S. Schulze, A. Scott, S. Sita, J. Shih,

K. Stewart, W. Tucker, E. Urquhart, K. van Atta, J. Vandermeer, L. van Zuidam, J. Verge,

V. Vonau, A. Walker-Bolton, E. Wikberg, E. Williams, L. Wolf, and D. Wood. We thank H.

Gilkenson and W. Lammers for long-term management of the field site during 2002-

2013. We thank the Costa Rican park service (Sistema Nacional de Áreas de

Conservación and Área de Conservación Tempisque), Hacienda Pelon de la Bajura,

Hacienda Brin D'Amor, and the residents of San Ramon de Bagaces for permission to

work on their land. K.K. was supported by a University of California-Los Angeles (UCLA)

Pauley Fellowship, a UCLA Dissertation Year Fellowship, and by the Max Planck Institute

for Evolutionary Anthropology. This material is based on data collection supported by

funding provided to S.E.P. by the National Science Foundation (under grant numbers

1638428, 0613226, 0848360, and 1232371), the Max Planck Institute for Evolutionary

Anthropology, the L.S.B. Leakey Foundation (5 grants), the National Geographic Society

(5 grants), the Templeton World Charity Foundation (grant #0208), the Wild Capuchin

Foundation, and multiple Committee on Research grants from UCLA. D. Cohen assisted

with database management and queries. This paper has benefited from helpful

discussions with H. C. Barrett, D. Cohen, J. Manson, and M.B. McElreath.

659 REFERENCES

660 Archie, E. A., Tung, J., Clark, M., Altmann, J., & Alberts, S. C. (2014). Social affiliation

661   matters: Both same-sex and opposite-sex relationships predict survival in wild

662   female baboons. *Proceedings of the Royal Society B: Biological Sciences*,

663   *281*(1793), 20141261.

664 Blumstein, D. T., Williams, D. M., Lim, A. N., Kroeger, S., & Martin, J. G. (2018). Strong

665   social relationships are associated with decreased longevity in a facultatively

666   social mammal. *Proceedings of the Royal Society B: Biological Sciences*,

667   *285*(1871), 20171934.

668 Brent, L. J. N., Ruiz-Lambides, A., & Platt, M. L. (2017). Family network size and survival

669   across the lifespan of female macaques. *Proceedings of the Royal Society B:*

670   *Biological Sciences*, *284*(1854), 20170515.

671 Clutton-Brock, T. H. (1988). *Reproductive success: Studies of individual variation in*

672   *contrasting breeding systems*. University of Chicago Press.

673 Dunbar, R. I. M. (2010). The social role of touch in humans and primates: Behavioural

674   function and neurobiological mechanisms. *Neuroscience & Biobehavioral*

675   *Reviews*, *34*(2), 260–268. https://doi.org/10.1016/j.neubiorev.2008.07.001

676 Fagen, R., & Fagen, J. (2004). Juvenile survival and benefits of play behaviour in brown

677   bears, Ursus arctos. *Evolutionary Ecology Research*, *6*(1), 89–102.

678 Farine, D. R. (2015). Proximity as a proxy for interactions: Issues of scale in social

679   network analysis. *Animal Behaviour*, *104*, e1–e5.

Farine, D. R., & Whitehead, H. (2015). Constructing, conducting and interpreting animal

social network analysis. *Journal of Animal Ecology*, *84*(5), 1144–1163.

Fedigan, L. M., Carnegie, S. D., & Jack, K. M. (2008). Predictors of reproductive success in

female white-faced capuchins (Cebus capucinus). *American Journal of Physical*

Anthropology, 137(1), 82–90.

Fedigan, L. M., & Rose, L. M. (1995). Interbirth interval variation in three sympatric

species of neotropical monkey. *American Journal of Primatology*, *37*(1), 9–24.

https://doi.org/10.1002/ajp.1350370103

Fragaszy, D. M., Visalberghi, E., & Fedigan, L. M. (2004). *The complete capuchin: The*

biology of the genus Cebus. Cambridge University Press.

Gilby, I. C., Brent, L. J., Wroblewski, E. E., Rudicell, R. S., Hahn, B. H., Goodall, J., & Pusey,

A. E. (2013). Fitness benefits of coalitionary aggression in male chimpanzees.

Behavioral Ecology and Sociobiology, *67*(3), 373–381.

Gros-Louis, J., Perry, S., & Manson, J. H. (2003). Violent coalitionary attacks and

intraspecific killing in wild white–faced capuchin monkeys (Cebus capucinus).

Primates, *44*(4), 341–346.

Hinde, R. A. (1976). Interactions, relationships and social structure. *Man*, 1–17.

Holt-Lunstad, J., Smith, T. B., & Layton, J. B. (2010). Social relationships and mortality

risk: A meta-analytic review. *PLoS Medicine*, *7*(7), e1000316.

Kajokaite, K., Whalen, A., Panchanathan, K., & Perry, S. (2019). White-faced capuchin

monkeys use both rank and relationship quality to recruit allies. *Animal*

Behaviour, *154*, 161–169.

Kalbitzer, U., Bergstrom, M. L., Carnegie, S. D., Wikberg, E. C., Kawamura, S., Campos, F.

A., Jack, K. M., & Fedigan, L. M. (2017). Female sociality and sexual conflict shape

offspring survival in a Neotropical primate. *Proceedings of the National Academy*

of Sciences, *114*(8), 1892–1897.

Koster, J., Leckie, G., Miller, A., & Hames, R. (2015). Multilevel modeling analysis of

dyadic network data with an application to Ye'kwana food sharing. *American*

Journal of Physical Anthropology, *157*(3), 507–512.

Koster, J., Leckie, G., & Aven, B. (2019). Statistical methods and software for the

Multilevel Social Relations Model. *Field Methods*, 1525822X19889011.

711
Manson, J. H., Navarrete, C. D., Silk, J. B., & Perry, S. (2004). Time-matched grooming in

female primates? New analyses from two species. *Animal Behaviour*, *67*(3), 493–

500.

McElreath, R. (2020). *Statistical rethinking: A Bayesian course with examples in R and*

Stan. CRC press.

McElreath, R. (2020). rethinking: Statistical rethinking book package (R package Version

1.93). https://github.com/rmcelreath/rethinking.

719
Nuñez, C. M., Adelman, J. S., & Rubenstein, D. I. (2014). Sociality increases juvenile

survival after a catastrophic event in the feral horse (Equus caballus). *Behavioral*

Ecology, *26*(1), 138–147.

Ostner, J., & Schuelke, O. (2018). Linking sociality to fitness in primates: A call for

mechanisms. In *Advances in the Study of Behavior* (Vol. 50, pp. 127–175).

Elsevier.

Perry, S. (1996). Female-female social relationships in wild white–faced capuchin

monkeys, *Cebus capucinus*. *American Journal of Primatology*, *40*(2), 167–182.

Perry, S. (2012). The behavior of wild white-faced capuchins: Demography, life history,

social relationships, and communication. In *Advances in the study of behavior*

(Vol. 44, pp. 135–181). Elsevier.

Perry, S., Godoy, I., & Lammers, W. (2012). The Lomas Barbudal Monkey Project: Two

decades of research on Cebus capucinus. In *Long-term field studies of primates*

(pp. 141–163). Springer.

Perry, S., Manson, J. H., & Perry, S. (2009). *Manipulative monkeys*. Harvard University

Press.

Porter, C. A., Page, S. L., Czelusniak, J., Schneider, H., Schneider, M. P. C., Sampaio, I., &

Goodman, M. (1997). Phylogeny and Evolution of Selected Primates as

Determined by Sequences of the ε-Globin Locus and 5' Flanking Regions.

International Journal of Primatology, *18*(2), 261–295.

https://doi.org/10.1023/A:1026328804319

R Core Team. (2020). R: A language and environment for statistical computing. Vienna,

Austria: R Foundation for Statistical Computing.

Roelfs, D. J., Shor, E., Kalish, R., & Yogev, T. (2011). The rising relative risk of mortality for

singles: Meta-analysis and meta-regression. *American Journal of Epidemiology*,

174(4), 379–389.

Schülke, O., Bhagavatula, J., Vigilant, L., & Ostner, J. (2010). Social bonds enhance

reproductive success in male macaques. *Current Biology*, *20*(24), 2207–2210.

Shor, E., & Roelfs, D. J. (2015). Social contact frequency and all-cause mortality: A meta-

analysis and meta-regression. *Social Science & Medicine*, *128*, 76–86.

Shor, E., Roelfs, D. J., & Yogev, T. (2013). The strength of family ties: A meta-analysis and

meta-regression of self-reported social support and mortality. *Social Networks*,

35(4), 626–638.

Silk, J. B. (2009). Nepotistic cooperation in non-human primate groups. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *364*(1533), 3243–3254.

Silk, J. B., Alberts, S. C., & Altmann, J. (2003). Social bonds of female baboons enhance

infant survival. *Science*, *302*(5648), 1231–1234.

Silk, J. B., Beehner, J. C., Bergman, T. J., Crockford, C., Engh, A. L., Moscovice, L. R.,

Wittig, R. M., Seyfarth, R. M., & Cheney, D. L. (2010). Strong and consistent social

bonds enhance the longevity of female baboons. *Current Biology*, *20*(15), 1359–

1361.

Silk, J., Cheney, D., & Seyfarth, R. (2013). A practical guide to the study of social

relationships. *Evolutionary Anthropology: Issues, News, and Reviews*, *22*(5), 213–

225.

Snijders, T. A., & Kenny, D. A. (1999). The social relations model for family data: A

multilevel approach. *Personal Relationships*, *6*(4), 471–486.

Stan Development Team. (2020). RStan: The R interface to Stan (Version 2.19.1).
768
Sterck, E. H. M., Watts, D. P., & van Schaik, C. P. (1997). The evolution of female social

relationships in nonhuman primates. *Behavioral Ecology and Sociobiology*, *41*(5),

291–309. https://doi.org/10.1007/s002650050390

Thompson, N. A. (2019). Understanding the links between social ties and fitness over

the life cycle in primates. *Behaviour*, *1*(aop), 1–50.

Thompson, N. A., & Cords, M. (2018). Stronger social bonds do not always predict

greater longevity in a gregarious primate. *Ecology and Evolution*, *8*(3), 1604–

1614.

Yee, J. R., Cavigelli, S. A., Delgado, B., & McClintock, M. K. (2008). Reciprocal affiliation

among adolescent rats during a mild group stressor predicts mammary tumors

and lifespan. *Psychosomatic Medicine*, *70*(9), 1050.

780