Combined deletion of free fatty-acid receptors 1 and 4 minimally impacts

glucose homeostasis in mice

Marine L. Croze[1], Arthur Guillaume[1], Mélanie Ethier[1], Grace Fergusson[1], Caroline Tremblay[1],

Scott A. Campbell[1], Hasna Maachi[1], Julien Ghislain[1], Vincent Poitout[1,2]

[1] Montreal Diabetes Research Center, CRCHUM, Montréal, QC, Canada; [2] Department of

Medicine, Université de Montréal, Montréal, QC, Canada

Running title: FFAR1 and FFAR4 regulation of glucose homeostasis

Corresponding author and reprint requests:

Vincent Poitout, DVM, PhD

CRCHUM

900 rue St Denis, Montréal, QC, H2X 0A9 - CANADA

Tel: (514) 890-8044

Email: vincent.poitout@umontreal.ca

Keywords: GPR120/FFAR4, GPR40/FFAR1, insulin, islet of Langerhans, glucose homeostasis.

This work was supported by a Discovery Grant from the Natural Sciences and Engineering

Research Council of Canada (RGPIN-2016-03952 to V.P.). M.L.C. was supported by a

fellowship from the Société Francophone du Diabète and a postdoctoral fellowship from the

Montreal Diabetes Research Center.

The authors have no relevant conflict of interest to disclose.

ABSTRACT

The free fatty-acid receptors FFAR1 (GPR40) and FFAR4 (GPR120) are implicated in the regulation of insulin secretion and insulin sensitivity, respectively. Although GPR120 and GPR40 share similar ligands, few studies have addressed possible interactions between these two receptors in the control of glucose homeostasis. Here we generated mice deficient in *gpr120* (Gpr120KO) or *gpr40* (Gpr40KO), alone or in combination (Gpr120/40KO), and metabolically phenotyped male and female mice fed a normal chow or high-fat diet. We assessed insulin secretion in isolated mouse islets exposed to selective GPR120 and GPR40 agonists singly or in combination. Following normal chow feeding, body weight and energy intake were unaffected by deletion of either receptor, although fat mass increased in Gpr120KO females. Fasting blood glucose levels were mildly increased in Gpr120/40KO mice, and in a sex-dependent manner in Gpr120KO and Gpr40KO animals. Oral glucose tolerance was slightly reduced in male Gpr120/40KO mice and in Gpr120KO females, whereas insulin secretion and insulin sensitivity were unaffected. In hyperglycemic clamps, the glucose infusion rate was lower in male Gpr120/40KO mice but insulin and c-peptide levels were unaffected. No changes in glucose tolerance were observed in either single or double KO animals under high-fat feeding. In isolated islets from wild-type mice, the combination of selective GPR120 and GPR40 agonists additively increased insulin secretion. We conclude that while simultaneous activation of GPR120 and GPR40 enhances insulin secretion ex vivo, combined deletion of these two receptors only minimally affects glucose homeostasis in vivo in mice.

INTRODUCTION

Insulin resistance and defective insulin secretion are hallmarks of type 2 diabetes (T2D). A number of G protein-coupled receptors (GPCR) that regulate insulin sensitivity and pancreatic β-cell function are validated targets for the treatment of T2D [1]. Among these, the long-chain fatty acid receptors FFAR1 (GPR40) and FFAR4 (GPR120) have been the subject of increasing interest in recent years as their activation has numerous beneficial effects on glucose and energy homeostasis in preclinical T2D models.

In rodents, *gpr120* is expressed in several tissues implicated in energy homeostasis including the central nervous system, enteroendocrine cells, liver, bone, adipose tissue and macrophages [2]. GPR120 activation in adipocytes and macrophages alleviates obesity-induced insulin resistance [3-5] in part via interaction with peroxisome proliferator-activated receptor γ [6]. GPR120 also promotes adipogenesis [7-9] and brown adipose tissue thermogenesis [10,11], regulates food intake [12] and modulates gut endocrine hormone secretion, including ghrelin [13-15] and the incretin hormones glucagon-like peptide-1 (GLP-1) [16], gastric inhibitory polypeptide (GIP) [17], cholecystokinin [18,19] and somatostatin [20]. GPR120 is also expressed in pancreatic islet cells where its activation mitigates β-cell dysfunction [21] and apoptosis [22] and modulates insulin [23-26], glucagon [27], somatostatin [28] and pancreatic polypeptide [29] secretion. Recently, we demonstrated that GPR120 activation potentiates glucose-stimulated insulin secretion (GSIS) and arginine-induced glucagon secretion in part through inhibition of somatostatin release by δ cells [30]. Despite the evidence for a role of GPR120 in the regulation of islet hormone secretion, *gpr120* deficient mice have reportedly normal β-cell function [3,5].

In rodents, *gpr40* is predominantly expressed in enteroendocrine cells and pancreatic β cells

[31,32]. Activation of GPR40 potentiates GSIS [31,33,34] and increases secretion of the incretin

hormones GLP-1, GIP and cholecystokinin [19,35-38] both ex vivo and in vivo. Recent studies

suggest that in the β cell GPR40 primarily responds to endogenous fatty acids released in

response to glucose stimulation and act in an autocrine manner to potentiate insulin secretion

[39]. Although deletion of GPR40 is largely inconsequential for glucose metabolism under

normal physiological conditions [33,40,41], high-fat diet (HFD) fed *gpr40*-null mice develop

fasting hyperglycemia due to insufficient insulin secretion [42].

Complementarity between *gpr120* and *gpr40* tissue expression and function as well as

overlapping ligands prompted us to investigate their possible interaction in the control of glucose

homeostasis. Redundancy between *gpr120* and *gpr40* has been described in the gut where oral

triglyceride-induced increases in plasma GIP and GLP-1 are dramatically reduced in double

gpr120/40 (Gpr120/40KO) but not *gpr120* (Gpr120KO) or *gpr40* (Gpr40KO) knockout mice

[38]. Furthermore, coactivation of GPR120 and GPR40 in vivo with selective synthetic agonists

or a dual agonist improves glucose control in *db/db* mice compared to activation of each receptor

alone [43]. However, whether GPR120 and GPR40 cooperate to improve glucose control under

physiological and pathophysiological conditions has not been addressed. In this study we

investigated possible functional interactions between these two receptors in the control of

glucose homeostasis with a particular emphasis on β-cell function. We generated single and

double *gpr120* and *gpr40* KO mice and compared metabolic parameters, glucose and insulin

tolerance as well as insulin secretion in vivo in both male and female mice fed a normal chow or

HFD.

MATERIALS AND METHODS

Reagents and solutions

RPMI-1640 and FBS were from Life Technologies Inc. (Burlington, ON, Canada). Penicillin/Streptomycin was from Multicell Wisent Inc (Saint-Jean-Baptiste, QC, Canada). Fatty-acid-free BSA was from Equitech-Bio (Kerrville, TX, USA). Insulin (Humulin-R 100U/mL) was from Eli Lilly (Toronto, ON, Canada). Compound A (Cpd A) was from Cayman Chemical (Ann Arbor, MI, USA) and TAK-875 was from Selleckchem (Houston, TX, USA). All other reagents were from MilliporeSigma unless otherwise specified.

Animals

All procedures involving animals were approved by the Institutional Committee for the Protection of Animals at the Centre Hospitalier de l'Université de Montréal. All mice were housed under controlled temperature on a 12h light/dark cycle with unrestricted access to water and laboratory chow. Mutant *gpr40* (RRID:MGI:3713765) [44] and *gpr120* (RRID:MGI:6477328) [45] colonies on a C57BL/6N (RRID:IMSR_JAX:005304) [46] background were maintained and genotyped as previously described. Male and female wild-type (WT, *gpr40$^{+/+}$;gpr120$^{+/+}$*), Gpr40KO (*gpr40$^{-/-}$;gpr120$^{+/+}$*), Gpr120KO (*gpr40$^{+/+}$;gpr120$^{-/-}$*) and Gpr120/40KO (*gpr40$^{-/-}$;gpr120$^{-/-}$*) experimental animals were generated in crosses between double heterozygotes (*gpr40$^{+/-}$;gpr120$^{+/-}$*). Animals were born at the expected Mendelian ratio. Both males and females were used for in vivo experiments and studied in separate cohorts. Animals were fed ad libitum with normal chow diet (CD; #2018 Teklad Global 18% protein rodent diet; 58% carbohydrate, 24% protein and 18% fat on a caloric basis; Harlan Teklad, Madison, WI) throughout the study or, after 8 weeks of age, switched to a HFD (#D12492; 60%

fat, 20% protein and 20% carbohydrate on a caloric basis; Research Diets Inc., New Brunswick,

NJ). Body weight, food intake and fed blood glucose were monitored weekly from 6-8 weeks of

age.

Metabolic tests

Oral Glucose Tolerance Tests (oGTT) were performed on 4-6 hour fasted mice by measuring tail

blood glucose and plasma insulin levels before (t=0 min) and 15, 30, 45, 60, 90 and 120 min

after oral glucose administration (Dextrose 1g/kg BW). Tail blood glucose was measured using

the hand-held glucometer Accu-Chek (Roche, Indianapolis, IN) and plasma insulin was

measured by ELISA (Alpco Diagnostics; RRID:AB_2792981) [47] at 0, 15 and 30 min.

Intraperitoneal Insulin Tolerance Tests (ipITT). Insulin (0.5 UI/kg, BW) was administered

intraperitoneally on 4-6 hour fasted mice and tail blood glucose was measured before (t=0 min)

and 15, 30, 45, 60, 90 and 120 min after insulin injection. Fasting insulinemia was measured by

ELISA (Alpco Diagnostics; RRID:AB_2792981) [47] on plasma samples from tail vein blood

collected before insulin injection.

Body weight composition was assessed with the EchoMRI Analyzer-700 (Echo Medical System,

Houston, TX).

Hyperglycemic clamps (HGC). One-step HGC were performed in conscious, ad libitum fed

animals as described [41]. Briefly, a 20% dextrose solution (Baxter, Mississauga, ON) was

infused via a jugular catheter. Mice initially received a 90-second bolus (140 mg/kg/minute) and

then the glucose infusion rate (GIR) was adjusted to maintain blood glucose between 14.5 and

17.7 mmol/l in females and between 15.7 and 19.7 mmol/l in males for 80 min. The insulin

sensitivity index (M/I) was calculated as the glucose infusion rate (M) divided by the average

plasma insulin during the last 30 min (i.e. from 50 to 80 min) of the clamp (I). The disposition

index (DI) was calculated by multiplying the M/I by the average plasma C-peptide during the

last 30 min of the clamp. Blood samples were collected from the tail to measure glucose, and

plasma insulin (Alpco Diagnostics; RRID:AB_2792981) [47] and C-peptide (Alpco Diagnostics;

RRID:AB_2801468) [48] were measured by ELISA at the time points indicated in the Figure

legends.

Islet isolation and static incubations for insulin secretion

Islets were isolated from 10-12 week-old male WT mice by collagenase digestion and dextran

density gradient centrifugation as described previously [33] and recovered overnight in RPMI

1640 supplemented with 10% (wt/vol) FBS, 100 U/ml penicillin/streptomycin and 11 mM

glucose. After recovery, triplicate batches of 20 islets each were incubated twice in KRBH (pH

7.4) with 0.1% (w/v) fatty-acid-free BSA and 2.8 mM glucose for 20 min., followed by a 1-hour

static incubation in KRBH in the presence of 2.8 or 16.7 mM glucose. Insulin radioimmunoassay

(MilliporeSigma; RRID:AB_2884035) [49] was used to measured insulin in the supernatant and

intracellular insulin content after acid–alcohol extraction.

Statistical analyses

Data are expressed as mean ± SEM. Significance was tested using ordinary one-way ANOVA or

Brown-Forsythe and Welch ANOVA tests with their correction in cases of variance

heterogeneity, two-way ANOVA with no assumption of sphericity and with post hoc adjustment

for multiple comparisons or three-way ANOVA, as appropriate, using GraphPad Instat

(RRID:SCR_000306) [50]. Tukey or Dunnett's post hoc tests were performed as indicated in

figure legends. P<0.05 was considered significant. Data points considered as outliers were

excluded based on Grubbs' test using the online calculator at:

https://www.graphpad.com/quickcalcs/Grubbs1.cfm.

RESULTS

Lean Gpr120/40KO mice exhibit mild glucose intolerance

In a first cohort of animals, WT, Gpr40KO, Gpr120KO and double Gpr120/40KO male and

female mice were CD-fed for 30 weeks. Metabolic parameters were assessed on a weekly basis

and Echo-MRI, glucose and insulin tolerance tests, and HGC were performed as indicated (Fig.

1A). WT, Gpr40KO, Gpr120KO and double Gpr120/40KO male and female mice showed

similar increases in body weight and energy intake over the course of the study (Fig. 1B-E).

Percentages of fat and lean mass as assessed by Echo-MRI at 27-29 weeks of age were

unaffected by genotype in male mice, whereas the proportion of fat to lean mass was slightly

increased in female Gpr120KO compared to WT mice (Fig. 1F and G). Fasting blood glucose

levels were significantly increased at 12 and 20 weeks of age in male and at all time points in

female Gpr120/40KO compared to WT mice (Fig. 2A and B). A significant increase in fasting

blood glucose compared to WT controls was also detected in male Gpr120KO mice at 20 weeks

of age, in female Gpr40KO mice at 16, 20 and 24 weeks of age, and at all time points in female

Gpr120KO mice. Weekly fed blood glucose levels were significantly increased in male

Gpr40KO vs. WT mice and in female Gpr40KO and Gpr120/40KO mice vs. their WT and

Gpr120KO littermates (Supplementary Fig. 1 in reference [51]).

oGTT performed at 10-12 weeks of age revealed prolonged glucose excursions in male

Gpr120/40KO mice that were significantly different between 60-120 min post oral glucose load

compared to WT controls and at 120 min compared to Gpr40KO mice (Fig. 2C). In females,

both single and double Gpr120/40KO mice exhibited reduced glucose tolerance between 45-90

min post oral glucose administration compared to WT mice, but the difference was significant

only for the Gpr120KO mice (Fig. 2D). oGTT performed at 18-20 weeks of age revealed mild

glucose intolerance in male Gpr120/40KO mice at 90 min and in female Gpr120/40KO and

Gpr120KO mice between 90-120 min compared to WT controls (Fig. 2E and F). Plasma insulin

levels determined during the oGTT at 10-12 and 18-20 weeks of age revealed no significant

differences between genotypes in either male or female mice (Supplementary Fig. 2 in reference

[51]).

ipITT performed in male and female mice at 14-16 week of age revealed no significant

difference in insulin tolerance between genotypes (Supplementary Fig. 2 in reference [51]).

We performed HGC at 28-30 weeks of age to assess β cell function in vivo. Blood glucose levels

were within the target range between 50 and 80 min of the clamp (Fig. 3A and B). Although

plasma insulin (Fig. 3C and D) and C-peptide (Fig. 3E and F) levels were similar in the 4

genotypes, the GIR and DI were significantly lower in male Gpr120/40KO vs. Gpr40KO mice

(Fig. 3G and I). A similar, albeit not statistically significant trend was observed in female mice

(Fig. 3H and J). Given that the reduction in DI is mostly driven by a lower GIR without

significant changes in c-peptide levels, we conclude from these data that the combined deletion

of *gpr120* and *gpr40* does not significantly impair insulin secretion in vivo. The absence of

significant differences between Gpr120/40KO and Gpr120KO mice suggests that deletion of

gpr120 accounts for the majority of the observed phenotype.

High-fat fed Gpr120/40KO mice have normal glucose tolerance

In a second cohort of animals, WT, Gpr40KO, Gpr120KO and Gpr120/40KO male and female

mice were fed either CD or HFD beginning at 8 weeks of age for a total of 12 weeks. Metabolic

parameters were assessed on a weekly basis. Echo-MRI and oral glucose tolerance tests were

performed as indicated (Fig. 4A). Both male and female HFD-fed mice accumulated more

weight than CD-fed animals, as expected, but no differences were detected between genotypes

on either feeding regimen (Fig. 4B and C). Male Gpr120/40KO mice fed a HFD exhibited

reduced energy intake compared to Gpr120KO and WT controls (Fig. 4D). In contrast, females

showed no differences in energy intake between genotype following either CD or HFD (Fig. 4E).

Relative fat mass was increased and lean mass decreased in HFD fed compared to CD-fed male

and female animals but no differences were detected between genotypes (Fig. 4F and G).

Mean and/or weekly fed blood glucose levels throughout the study were increased in CD-fed

male Gpr120/40KO compared to all other genotypes and in HFD-fed Gpr120/40KO and

Gpr120KO compared to WT mice but unaffected by genotype in female mice (Supplementary

Fig. 3 in reference [51]).

oGTT performed at 20 weeks of age revealed reduced glucose tolerance in male Gpr120/40KO

mice fed a CD at 60 and 120 min compared to WT controls (Fig. 5A), similar to our previous

findings (Fig. 2). However, plasma insulin levels determined during the test revealed no

significant differences between genotypes (Fig. 5B). A non-significant trend for reduced glucose

tolerance was observed in Gpr120/40KO females (Fig. 5C), without changes in plasma insulin

levels (Fig. 5D). Following HFD, no significant differences in glucose tolerance or plasma

insulin levels were detected between genotypes in either male (Fig. 5E and F) or female (Fig. 5G

and H) mice. Taken together these data suggest that combined deletion of *gpr120* and *gpr40* does

not aggravate glucose intolerance under HFD in either male or female mice.

GPR120 and GPR40 agonists additively potentiate GSIS in isolated mouse islets

Agonists of GPR120 or GPR40 enhance GSIS in isolated mouse islets [23-26,30,31,33,34]. Here

we asked whether simultaneous activation of GPR120 and GPR40 potentiates GSIS additively or

synergistically compared to the activation of each receptor alone. Islets were isolated from lean

male WT mice and insulin secretion was assessed in response to the selective GPR40 agonist

TAK-875 and GPR120 agonist Cpd A, singly or in combination (Fig. 6). Glucose alone

increased insulin secretion (Fig. 6A and B), as expected. TAK-875 and Cpd A alone weakly but

significantly potentiated this effect and combined exposure to TAK-875 and Cpd A additively

increased GSIS (Fig. 6A and B). No significant differences were detected in total insulin content

(Fig. 6C).

DISCUSSION

The objective of this study was to assess possible interactions between GPR120 and GPR40 in

the control of glucose homeostasis and β cell function under physiological conditions and

metabolic stress. By comparing adult, chow-fed WT and *gpr120* and *gpr40* single and double

KO mice we showed that the combined loss of both genes leads to a slight reduction in glucose

tolerance but no substantial impairment of insulin secretion in vivo. In most cases, the

Gpr120/40KO mice phenotype seemed to be mainly related to the loss of GPR120 and slightly

aggravated by the loss of GPR40, suggesting that the two receptors signaling pathways act

additively rather than synergistically. No significant differences between genotypes were

detected in glucose homeostasis following 12 weeks of high-fat feeding in either sex. We then

showed that selective agonists for GPR120 and GPR40 additively potentiate GSIS in isolated

islets ex vivo.

Gpr120KO - Deletion of *gpr120* had no significant impact on body weight in CD-fed male or

female mice, in agreement with previous studies [3,5]. In contrast, Suckow et al. [27] observed

an increase in body mass in *gpr120*-null mice following a similar diet regimen. In our study, fat

mass was increased in female Gpr120KO mice which may be accounted for by the reduced

metabolic rate, inhibition of adipocyte lipolysis and increased size of adipocytes observed

previously in *gpr120*-null mice [5,43].

Fasting glucose levels were elevated in male and female Gpr120KO mice under CD, but oral

glucose tolerance was unaffected in male and mildly reduced in female Gpr120KO mice, and

there were no differences in circulating insulin levels or insulin tolerance. These findings agree

with previous reports [5,24]. They do, however, contrast with some studies which found glucose

intolerance [3,27] and elevated plasma insulin levels and insulin resistance [3] in *gpr120*-null

male mice. Age-related differences may play a role as insulin tolerance test were performed at

14-16 weeks of age in this study whereas Oh et al., analyzed mice using

hyperinsulinemic/euglycemic clamps at 23-28 weeks of age [3].

Following HFD, body weight, glucose tolerance and insulin levels were not affected in either

male or female Gpr120KO mice, similar to previous findings in male mice [3,4,52]. This is in

contrast to the findings of Ichimura et al. [5] who reported that HFD-fed *gpr120*-null male mice

gain more weight and are glucose intolerant. Discrepancies in *gpr120*-null metabolic phenotypes

may be accounted for by differences in diet composition and genetic background, such as the

mixed C57Bl/6/129 used by Ichimura et al. [5] vs. the pure C57Bl/6N used by Bjursell et al. [52]

and in this study. It should be noted, however, that a role of GPR120 signaling has been

demonstrated in the context of chronic administration of specific agonists such as Cpd A [4,6,26]

and GSK137647 [26] which attenuate the negative effects of HFD on energy and glucose

homeostasis in WT but not *gpr120*-null mice. This suggests that during metabolic stress and/or

chronic feeding with a diet poor in omega-3 fatty acids, levels of endogenous GPR120 ligands

may be insufficient to confer a protective effect.

Gpr40KO - Deletion of *gpr40* alone had no significant impact on energy homeostasis in male or

female mice fed either CD or HFD, as shown previously [33,41,42,53].

Glucose and insulin tolerance and insulin secretion in CD-fed Gpr40KO mice were also largely

unaffected, as previously observed [33]. Contrary to our previous findings [42], we did not detect

significant differences in fasting glucose levels in chow-fed, male Gpr40KO mice vs WT. This

could be accounted for by a greater statistical power in our previous study, in which only 2

groups with a larger sample size were compared, and/or differences in the age of the mice (7

weeks in [42] vs 12-24 weeks in this study). In the present study, we found however that female

Gpr40KO mice had higher fasting glucose, and male and female mice had higher fed glucose,

than WT controls. The significant reduction in insulin levels previously detected during HGC in

gpr40-null mice [41] was not seen in the present study, which may be due to difference in the

age and/or strain of the animals. The nicotinamide nucleotide transhydrogenase mutation and

other polymorphisms in the C57Bl/6J background used in our previous study [41] – vs. the

C57Bl/6N background used here - are known to attenuate insulin secretion [54] and may interact

with the *gpr40* mutation to aggravate the secretory defect.

In a HFD study by Kebede et al. [42] we observed a precocious increase in fasting blood glucose

levels after 3 weeks of diet in male Gpr40KO mice which normalized after 8 weeks, similar to

our findings after 12 weeks of diet.

Gpr120/40KO - Although weight gain was similar, male, but not female, Gpr120/40KO mice

exhibited reduced energy intake compared to Gpr120KO and WT controls. GPR120 and GPR40

are expressed in hypothalamic microglia and neurons, respectively [55]. Interestingly,

intracerebroventricular injection of GW9508, a GPR120 and GPR40 dual agonist, increases food

intake in HFD fed mice [55]. Although the underlying reason for the decreased food intake in

Gpr120/40KO mice is unknown, loss-of-function of these genes in the hypothalamus would be

expected to alter food intake.

In mice fed a CD, we observed an increase in fasting glucose levels in both male and female

Gpr120/40KO and a slight decrease in glucose tolerance in males. These changes were not

observed in any of the groups under HFD. The reduction in glucose tolerance was more

pronounced in double Gpr120/40KO compared to single Gpr120KO or Gpr40KO mice,

indicative of an additive effect. As no differences were detected in insulin tolerance between

genotypes, the mild reduction in glucose tolerance in Gpr120/40KO mice might be due to

deletion of the genes in pancreatic islets and/or reduced incretin hormone secretion. Along these

lines, Ekberg et al. [38] demonstrated that *gpr120* and *gpr40* are highly expressed in GLP-1 and

GIP-secreting cells in the gut, and that the combined loss of the two receptors leads to a major

reduction in GLP-1 and GIP levels following oral triglyceride administration, which is not seen

in mice deficient in each receptor alone. However, we cannot exclude that a mild insulin

resistance in Gpr120KO and Gpr120/40KO compared to WT mice, not detected in the ipITT,

contributes to the increased fasting glucose levels and glucose intolerance. This possibility is

consistent with the lower GIR observed during HGC in 28-30 weeks Gpr120/40KO mice, and is

supported by the finding that Gpr120KO mice can develop insulin resistance [3]. However,

glucagon secretion and glucagon sensitivity are elevated in Gpr120KO mice [27] and Gpr40

agonist treatment inhibits glucose production in Goto Kakizaki rat through changes in

gluconeogenesis [56]. Hence, increased glycogenolysis and gluconeogenesis could also

contribute to higher blood glucose levels in Gpr120/40KO mice.

The reduced DI in HGC was largely due to a reduction in GIR with no changes in insulin or c-

peptide levels in male Gpr120/40KO vs single Gpr40KO. Nevertheless, the reduced GIR is

suggestive of a relative functional β-cell deficit, possibly due to a lack of β-cell mass

compensation to age-related insulin resistance in Gpr120/40KO males. Indeed, *gpr120* deletion

is associated with insulin resistance in mice of a similar age [3]. Although our previous studies

indicate that *gpr40* deletion alone does not impact β-cell mass [41], given the role of GPR120

and GPR40 in the regulation of hormones that control β-cell proliferation, including somatostatin

[20,28,30] and GLP-1 [16,24,35,36,38], it is possible that their deletion impacts β-cell mass. We

did not test for this possibility in our study. The reduced GIR seems to be mainly attributable to

the deletion of *gpr120* and slightly aggravated by the additional loss of *gpr40*. A non-significant

trend was detected in female Gpr120/40KO mice. Phenotypic differences between male and

female mice may be due to a number of sex-related factors known to contribute to sex

differences in metabolism [57].

Alterations in glucose homeostasis in Gpr120/40KO mice following CD but not HFD are notable

for their similarities to Gpr120KO mice, which show the same trends. We surmise that HFD may

diminish GPR120 signaling thereby limiting the effect of gene loss in single and double

Gpr120KO mice compared to Gpr40KO and WT animals. This could be in part related to lower

intake of GPR120 ligands, since HFD are poor in ω-3 polyunsaturated FAs compared to CD.

Isolated islets - The stimulatory effects of the GPR120 agonist Cpd A and the GPR40 agonist

TAK-875 were additive and not synergistic. This is consistent with our in vivo observations in

double KO mice and suggests that the signaling mechanisms of the two receptors in islets are

independent. Recently, using a δ-cell specific Gpr120KO mouse we showed that Cpd A

potentiates GSIS largely via δ cell-specific GPR120 inhibitory signaling that dampens cAMP

generation and somatostatin secretion, alleviating a paracrine inhibitory signal that restricts

insulin secretion [30]. In contrast, GPR40 is predominantly expressed in islet β cells where its

activation directly promotes insulin secretion via Gαq/11 coupling and phospholipase C –

protein kinase D1 signaling [34]. However, our studies also revealed that *gpr120* is expressed in

β cells and that Cpd A increases cAMP generation and calcium fluxes in β cells [30].

Furthermore, in a recent study, ciliary GPR120 activation was shown to increase cAMP levels

and promote insulin secretion in intact islets and insulin-secreting cell lines [58]. Together these

studies suggest that although GPR120 inhibitory signaling in δ cells indirectly increases insulin

secretion, GPR120 stimulatory signals in β cells may also contribute to its modulation of islet

hormone secretion. This model is consistent with our observation that the stimulatory effects of

GPR120 and GPR40 on insulin secretion are additive rather than synergistic.

Conclusion - Our results show that combined deletion of GPR120 and GPR40 minimally alters

glucose homeostasis in mice, with only a mild impairment of glucose tolerance in male animals.

Our data also suggest that the two receptors signaling cooperate in an additive rather than in a

synergistic manner to potentiate glucose-stimulated insulin secretion ex vivo. These findings

contribute to our understanding of the role of these fatty-acid GPCR in the regulation of glucose

homeostasis and islet function.

ACKNOWLEDGEMENTS

We thank Annie Levert (CRCHUM) for valuable technical assistance.

DATA AVAILABILITY

The datasets generated during and/or analyzed during the current study are not publicly available

but are available from the corresponding author on reasonable request.

REFERENCES

1. Ahren B. Islet G protein-coupled receptors as potential targets for treatment of type 2 diabetes.

Nat Rev Drug Discov 2009;8(5):369-85.

2. Moniri NH. Free-fatty acid receptor-4 (GPR120): Cellular and molecular function and its role

in metabolic disorders. *Biochem Pharmacol* 2016;110-111:1-15.

3. Oh DY, Talukdar S, Bae EJ, Imamura T, Morinaga H, Fan W, Li P, Lu WJ, Watkins SM,

Olefsky JM. GPR120 is an omega-3 fatty acid receptor mediating potent anti-inflammatory and

insulin-sensitizing effects. *Cell* 2010;142(5):687-98.

4. Oh da Y, Walenta E, Akiyama TE, Lagakos WS, Lackey D, Pessentheiner AR, Sasik R, Hah

N, Chi TJ, Cox JM, Powels MA, Di Salvo J, Sinz C, Watkins SM, Armando AM, Chung H,

Evans RM, Quehenberger O, McNelis J, Bogner-Strauss JG, Olefsky JM. A Gpr120-selective

agonist improves insulin resistance and chronic inflammation in obese mice. *Nat Med*

2014;20(8):942-7.

5. Ichimura A, Hirasawa A, Poulain-Godefroy O, Bonnefond A, Hara T, Yengo L, Kimura I,

Leloire A, Liu N, Iida K, Choquet H, Besnard P, Lecoeur C, Vivequin S, Ayukawa K, Takeuchi

M, Ozawa K, Tauber M, Maffeis C, Morandi A, Buzzetti R, Elliott P, Pouta A, Jarvelin MR,

Korner A, Kiess W, Pigeyre M, Caiazzo R, Van Hul W, Van Gaal L, Horber F, Balkau B, Levy-

Marchal C, Rouskas K, Kouvatsi A, Hebebrand J, Hinney A, Scherag A, Pattou F, Meyre D,

Koshimizu TA, Wolowczuk I, Tsujimoto G, Froguel P. Dysfunction of lipid sensor GPR120

leads to obesity in both mouse and human. *Nature* 2012;483(7389):350-4.

6. Paschoal VA, Walenta E, Talukdar S, Pessentheiner AR, Osborn O, Hah N, Chi TJ, Tye GL,

Armando AM, Evans RM, Chi NW, Quehenberger O, Olefsky JM, Oh DY. Positive Reinforcing

Mechanisms between GPR120 and PPARγ Modulate Insulin Sensitivity. *Cell Metab*

2020;31(6):1173-88.e5.

7. Gotoh C, Hong YH, Iga T, Hishikawa D, Suzuki Y, Song SH, Choi KC, Adachi T, Hirasawa

A, Tsujimoto G, Sasaki S, Roh SG. The regulation of adipogenesis through GPR120. *Biochem*

Biophys Res Commun 2007;354(2):591-7.

8. Song T, Zhou Y, Peng J, Tao YX, Yang Y, Xu T, Peng J, Ren J, Xiang Q, Wei H. GPR120

promotes adipogenesis through intracellular calcium and extracellular signal-regulated kinase 1/2

signal pathway. *Mol Cell Endocrinol* 2016;434:1-13.

9. Hilgendorf KI, Johnson CT, Mezger A, Rice SL, Norris AM, Demeter J, Greenleaf WJ, Reiter

JF, Kopinke D, Jackson PK. Omega-3 Fatty Acids Activate Ciliary FFAR4 to Control

Adipogenesis. *Cell* 2019;179(6):1289-305.e21.

10. Quesada-Lopez T, Cereijo R, Turatsinze JV, Planavila A, Cairo M, Gavalda-Navarro A,

Peyrou M, Moure R, Iglesias R, Giralt M, Eizirik DL, Villarroya F. The lipid sensor GPR120

promotes brown fat activation and FGF21 release from adipocytes. *Nat Commun* 2016;7:13479.

11. Schilperoort M, van Dam AD, Hoeke G, Shabalina IG, Okolo A, Hanyaloglu AC, Dib LH,

Mol IM, Caengprasath N, Chan Y-W, Damak S, Miller AR, Coskun T, Shimpukade B, Ulven T,

Kooijman S, Rensen PC, Christian M. The GPR120 agonist TUG-891 promotes metabolic health

by stimulating mitochondrial respiration in brown fat. *EMBO Mol Med* 2018;10(3):e8047.

12. Auguste S, Fisette A, Fernandes MF, Hryhorczuk C, Poitout V, Alquier T, Fulton S. Central

Agonism of GPR120 Acutely Inhibits Food Intake and Food Reward and Chronically Suppresses

Anxiety-Like Behavior in Mice. *Int J Neuropsychopharmacol* 2016;19(7):pyw014.

13. Lu X, Zhao X, Feng J, Liou AP, Anthony S, Pechhold S, Sun Y, Lu H, Wank S. Postprandial

inhibition of gastric ghrelin secretion by long-chain fatty acid through GPR120 in isolated gastric

ghrelin cells and mice. *Am J Physiol Gastrointest Liver Physiol* 2012;303(3):G367-76.

14. Engelstoft MS, Park WM, Sakata I, Kristensen LV, Husted AS, Osborne-Lawrence S, Piper

PK, Walker AK, Pedersen MH, Nohr MK, Pan J, Sinz CJ, Carrington PE, Akiyama TE, Jones

RM, Tang C, Ahmed K, Offermanns S, Egerod KL, Zigman JM, Schwartz TW. Seven

transmembrane G protein-coupled receptor repertoire of gastric ghrelin cells. *Mol Metab*

2013;2(4):376-92.

15. Gong Z, Yoshimura M, Aizawa S, Kurotani R, Zigman JM, Sakai T, Sakata I. G protein-

coupled receptor 120 signaling regulates ghrelin secretion in vivo and in vitro. *Am J Physiol*

Endocrinol Metab 2014;306(1):E28-35.

16. Hirasawa A, Tsumaya K, Awaji T, Katsuma S, Adachi T, Yamada M, Sugimoto Y, Miyazaki

S, Tsujimoto G. Free fatty acids regulate gut incretin glucagon-like peptide-1 secretion through

GPR120. *Nat Med* 2005;11(1):90-4.

17. Iwasaki K, Harada N, Sasaki K, Yamane S, Iida K, Suzuki K, Hamasaki A, Nasteska D,

Shibue K, Joo E, Harada T, Hashimoto T, Asakawa Y, Hirasawa A, Inagaki N. Free fatty acid

receptor GPR120 is highly expressed in enteroendocrine K cells of the upper small intestine and

has a critical role in GIP secretion after fat ingestion. *Endocrinology* 2015;156(3):837-46.

18. Tanaka T, Katsuma S, Adachi T, Koshimizu TA, Hirasawa A, Tsujimoto G. Free fatty acids

induce cholecystokinin secretion through GPR120. *Naunyn Schmiedebergs Arch Pharmacol*

2008;377(4-6):523-7.

19. Sankoda A, Harada N, Iwasaki K, Yamane S, Murata Y, Shibue K, Thewjitcharoen Y,

Suzuki K, Harada T, Kanemaru Y, Shimazu-Kuwahara S, Hirasawa A, Inagaki N. Long-Chain

Free Fatty Acid Receptor GPR120 Mediates Oil-Induced GIP Secretion Through CCK in Male

Mice. *Endocrinology* 2017;158(5):1172-80.

20. Egerod KL, Engelstoft MS, Lund ML, Grunddal KV, Zhao M, Barir-Jensen D, Nygaard EB,

Petersen N, Holst JJ, Schwartz TW. Transcriptional and Functional Characterization of the G

Protein-Coupled Receptor Repertoire of Gastric Somatostatin Cells. *Endocrinology*

2015;156(11):3909-23.

21. Wang Y, Xie T, Zhang D, Leung PS. GPR120 protects lipotoxicity-induced pancreatic β-cell

dysfunction through regulation of PDX1 expression and inhibition of islet inflammation. *Clin Sci*

(Lond) 2019;133(1):101-16.

22. Taneera J, Lang S, Sharma A, Fadista J, Zhou Y, Ahlqvist E, Jonsson A, Lyssenko V,

Vikman P, Hansson O, Parikh H, Korsgren O, Soni A, Krus U, Zhang E, Jing XJ, Esguerra JL,

Wollheim CB, Salehi A, Rosengren A, Renstrom E, Groop L. A systems genetics approach

identifies genes and pathways for type 2 diabetes in human islets. *Cell Metab* 2012;16(1):122-34.

23. Moran BM, Abdel-Wahab YH, Flatt PR, McKillop AM. Evaluation of the insulin-releasing

and glucose-lowering effects of GPR120 activation in pancreatic beta-cells. *Diabetes Obes*

Metab 2014;16(11):1128-39.

24. Sundstrom L, Myhre S, Sundqvist M, Ahnmark A, McCoull W, Raubo P, Groombridge SD,

Polla M, Nystrom AC, Kristensson L, Nagard M, Winzell MS. The acute glucose lowering effect

of specific GPR120 activation in mice is mainly driven by glucagon-like peptide 1. *PLoS One*

2017;12(12):e0189060.

25. Zhang D, So WY, Wang Y, Wu SY, Cheng Q, Leung PS. Insulinotropic effects of GPR120

agonists are altered in obese diabetic and obese non-diabetic states. *Clin Sci (Lond)*

2017;131(3):247-60.

26. McCloskey AG, Miskelly MG, Flatt PR, McKillop AM. Pharmacological potential of novel

agonists for FFAR4 on islet and enteroendocrine cell function and glucose homeostasis. *Eur J*

Pharm Sci 2020;142:105104.

27. Suckow AT, Polidori D, Yan W, Chon S, Ma JY, Leonard J, Briscoe CP. Alteration of the

glucagon axis in GPR120 (FFAR4) knockout mice: a role for GPR120 in glucagon secretion. *J*

Biol Chem 2014;289(22):15751-63.

28. Stone VM, Dhayal S, Brocklehurst KJ, Lenaghan C, Sorhede Winzell M, Hammar M, Xu X,

Smith DM, Morgan NG. GPR120 (FFAR4) is preferentially expressed in pancreatic delta cells

and regulates somatostatin secretion from murine islets of Langerhans. *Diabetologia*

2014;57(6):1182-91.

29. Zhao YF, Li XC, Liang XY, Zhao YY, Xie R, Zhang LJ, Zhang XC, Chen C. GPR120

Regulates Pancreatic Polypeptide Secretion From Male Mouse Islets via PLC-Mediated Calcium

Mobilization. *Endocrinology* 2020;161(10).

30. Croze ML, Flisher MF, Guillaume A, Tremblay C, Noguchi GM, Granziera S, Vivot K,

Castillo VC, Campbell SA, Ghislain J, Huising MO, Poitout V. Free-fatty acid receptor 4

inhibitory signaling in delta cells regulates islet hormone secretion in mice. *bioRxiv* 2020:2020.07.17.208637.

31. Itoh Y, Kawamata Y, Harada M, Kobayashi M, Fujii R, Fukusumi S, Ogi K, Hosoya M, Tanaka Y, Uejima H, Tanaka H, Maruyama M, Satoh R, Okubo S, Kizawa H, Komatsu H, Matsumura F, Noguchi Y, Shinohara T, Hinuma S, Fujisawa Y, Fujino M. Free fatty acids regulate insulin secretion from pancreatic beta cells through GPR40. *Nature* 2003;422(6928):173-6.

32. Briscoe CP, Tadayyon M, Andrews JL, Benson WG, Chambers JK, Eilert MM, Ellis C, Elshourbagy NA, Goetz AS, Minnick DT, Murdock PR, Sauls HR, Jr., Shabon U, Spinage LD, Strum JC, Szekeres PG, Tan KB, Way JM, Ignar DM, Wilson S, Muir AI. The orphan G protein-coupled receptor GPR40 is activated by medium and long chain fatty acids. *J Biol Chem* 2003;278(13):11303-11.

33. Latour MG, Alquier T, Oseid E, Tremblay C, Jetton TL, Luo J, Lin DC, Poitout V. GPR40 is necessary but not sufficient for fatty acid stimulation of insulin secretion in vivo. *Diabetes* 2007;56(4):1087-94.

34. Ferdaoussi M, Bergeron V, Zarrouki B, Kolic J, Cantley J, Fielitz J, Olson EN, Prentki M, Biden T, MacDonald PE, Poitout V. G protein-coupled receptor (GPR)40-dependent potentiation of insulin secretion in mouse islets is mediated by protein kinase D1. *Diabetologia* 2012;55(10):2682-92.

35. Edfalk S, Steneberg P, Edlund H. Gpr40 is expressed in enteroendocrine cells and mediates free fatty acid stimulation of incretin secretion. *Diabetes* 2008;57(9):2280-7.

36. Xiong Y, Swaminath G, Cao Q, Yang L, Guo Q, Salomonis H, Lu J, Houze JB, Dransfield PJ, Wang Y, Liu JJ, Wong S, Schwandner R, Steger F, Baribault H, Liu L, Coberly S, Miao L,

Zhang J, Lin DC, Schwarz M. Activation of FFA1 mediates GLP-1 secretion in mice. Evidence

for allosterism at FFA1. *Mol Cell Endocrinol* 2013;369(1-2):119-29.

37. Liou AP, Lu X, Sei Y, Zhao X, Pechhold S, Carrero RJ, Raybould HE, Wank S. The G-

protein-coupled receptor GPR40 directly mediates long-chain fatty acid-induced secretion of

cholecystokinin. *Gastroenterology* 2011;140(3):903-12.

38. Ekberg JH, Hauge M, Kristensen LV, Madsen AN, Engelstoft MS, Husted AS, Sichlau R,

Egerod KL, Timshel P, Kowalski TJ, Gribble FM, Reiman F, Hansen HS, Howard AD, Holst B,

Schwartz TW. GPR119, a Major Enteroendocrine Sensor of Dietary Triglyceride Metabolites

Coacting in Synergy With FFA1 (GPR40). *Endocrinology* 2016;157(12):4561-69.

39. Tunaru S, Bonnavion R, Brandenburger I, Preussner J, Thomas D, Scholich K, Offermanns

S. 20-HETE promotes glucose-stimulated insulin secretion in an autocrine manner through

FFAR1. *Nat Commun* 2018;9(1):177.

40. Tan CP, Feng Y, Zhou Y-P, Eiermann GJ, Petrov A, Zhou C, Lin S, Salituro G, Meinke P,

Mosley R, Akiyama TE, Einstein M, Kumar S, Berger JP, Mills SG, Thornberry NA, Yang L,

Howard AD. Selective small-molecule agonists of G protein-coupled receptor 40 promote

glucose-dependent insulin secretion and reduce blood glucose in mice. *Diabetes*

2008;57(8):2211-19.

41. Alquier T, Peyot ML, Latour MG, Kebede M, Sorensen CM, Gesta S, Ronald Kahn C, Smith

RD, Jetton TL, Metz TO, Prentki M, Poitout V. Deletion of GPR40 impairs glucose-induced

insulin secretion in vivo in mice without affecting intracellular fuel metabolism in islets.

Diabetes 2009;58(11):2607-15.

42. Kebede M, Alquier T, Latour MG, Semache M, Tremblay C, Poitout V. The fatty acid

receptor GPR40 plays a role in insulin secretion in vivo after high-fat feeding. *Diabetes*

2008;57(9):2432-7.

43. Satapati S, Qian Y, Wu MS, Petrov A, Dai G, Wang SP, Zhu Y, Shen X, Muise ES, Chen Y,

Zycband E, Weinglass A, Di Salvo J, Debenham JS, Cox JM, Lan P, Shah V, Previs SF, Erion

M, Kelley DE, Wang L, Howard AD, Shang J. GPR120 suppresses adipose tissue lipolysis and

synergizes with GPR40 in antidiabetic efficacy. *J Lipid Res* 2017;58(8):1561-78.

44. RRID:MGI:3713765. http://www.informatics.jax.org/allele/MGI:3713759.

45. RRID:MGI:6477328.

46. RRID:IMSR_JAX:005304. https://www.jax.org/strain/005304.

47. RRID:AB_2792981. http://antibodyregistry.org/AB_2792981.

48. RRID:AB_2801468. http://antibodyregistry.org/AB_2801468.

49. RRID:AB_2884035. http://antibodyregistry.org/AB_2884035.

50. RRID:SCR_000306.

https://scicrunch.org/resources/Tools/record/nlx_144509-

1/SCR_000306/resolver?q=SCR_002798&l=SCR_002798.

51. Croze ML, Guillaume A, Ethier M, Fergusson G, Tremblay C, Campbell SA, Maachi H,

Ghislain J, Poitout V. Supplementary Data from: Combined deletion of free fatty-acid receptors

1 and 4 minimally impacts glucose homeostasis in mice. *figshare*. 2020. Deposited 11 December

2020. https://doi.org/10.6084/m9.figshare.13365962.v1

52. Bjursell M, Xu X, Admyre T, Bottcher G, Lundin S, Nilsson R, Stone VM, Morgan NG,

Lam YY, Storlien LH, Linden D, Smith DM, Bohlooly YM, Oscarsson J. The beneficial effects

of n-3 polyunsaturated fatty acids on diet induced obesity and impaired glucose control do not

require Gpr120. *PLoS One* 2014;9(12):e114942.

53. Lan H, Hoos LM, Liu L, Tetzloff G, Hu W, Abbondanzo SJ, Vassileva G, Gustafson EL,

Hedrick JA, Davis HR. Lack of FFAR1/GPR40 does not protect mice from high-fat diet-induced

metabolic disease. *Diabetes* 2008;57(11):2999-3006.

54. Fergusson G, Ethier M, Guevremont M, Chretien C, Attane C, Joly E, Fioramonti X, Prentki

M, Poitout V, Alquier T. Defective insulin secretory response to intravenous glucose in C57Bl/6J

compared to C57Bl/6N mice. *Mol Metab* 2014;3(9):848-54.

55. Dragano NRV, Solon C, Ramalho AF, de Moura RF, Razolli DS, Christiansen E, Azevedo C,

Ulven T, Velloso LA. Polyunsaturated fatty acid receptors, GPR40 and GPR120, are expressed

in the hypothalamus and control energy homeostasis and inflammation. *J Neuroinflammation*

2017;14(1):91.

56. Miller C, Pachanski MJ, Kirkland ME, Kosinski DT, Mane J, Bunzel M, Cao J, Souza S,

Thomas-Fowlkes B, Di Salvo J, Weinglass AB, Li X, Myers RW, Knagge K, Carrington PE,

Hagmann WK, Trujillo ME. GPR40 partial agonist MK-2305 lower fasting glucose in the Goto

Kakizaki rat via suppression of endogenous glucose production. *PLoS One*

2017;12(5):e0176182.

57. Mauvais-Jarvis F. Epidemiology of Gender Differences in Diabetes and Obesity. *Adv Exp*

Med Biol 2017;1043:3-8.

58. Wu C-T, Hilgendorf KI, Bevacqua RJ, Hang Y, Demeter J, Kim SK, Jackson PK. Discovery

of ciliary G protein-coupled receptors regulating pancreatic islet insulin and glucagon secretion.

bioRxiv 2020:2020.10.21.349423.

   FIGURE LEGENDS

Figure 1. Energy balance in *gpr40* and *gpr120* single and double knockout (KO) mice under

chow diet (CD). (**A**) Study plan to assess energy and glucose homeostasis in wild-type (WT),

Gpr40KO (40KO), Gpr120KO (120KO) and Gpr120/40KO (120/40KO) male and female mice

fed CD. (**B-G**) body weight (**B, C**), cumulative energy intake (**D, E**) and fat and lean mass as a

percentage of body weight measured by Echo-MRI at 27-29 weeks of age (**F, G**) in male (**B, D,**

F) and female (**C, E, G**) mice. Data are presented as mean +/- SEM (n=6-8 animals per group).

P-values were calculated by regular 2-way ANOVA or mixed model when some follow-up

values were missing with no assumption of sphericity (**B-E**) or 2-way ANOVA with post-hoc

Tukey's multiple comparisons test (**F, G**). *$p < 0.05$.

Figure 2. Glucose homeostasis in *gpr40* and *gpr120* single and double knockout (KO) mice

under chow diet (CD). (**A, B**) Fasted blood glucose at 12, 16, 20 and 24 weeks of age in wild-

type (WT), Gpr40KO (40KO), Gpr120KO (120KO) and Gpr120/40KO (120/40KO) male (**A**)

and female (**B**) mice fed CD. (**C-F**) Blood glucose during an oGTT (glucose 1g/kg BW at 0 min)

at 10-12 (**C, D**) and at 18-20 (**E, F**) weeks of age in male (**C, E**) and female (**D, F**) mice. Data

are presented as mean +/- SEM (n=7-13 animals per group). P-values were calculated by regular

2-way ANOVA or mixed model when some values were missing, with no assumption of

sphericity and with Tukey's multiple comparisons tests. **$p<0.01$ and *$p<0.05$, 120/40KO vs

WT; &, $p<0.05$ 120/40KO and 120KO vs WT; £, $p<0.01$ 120/40KO and 120KO vs WT, $p<0.05$

120/40KO and 120KO vs Gpr40KO; $, at least $p<0.05$ for 120KO and 40KO vs WT and at least

$p<0.005$ for the 120/40KO vs WT; #, $p<0.05$ 120/40KO vs 40KO; Φ, at least $p<0.05$ WT vs

120KO; Σ, at least $p<0.05$ 120/40KO and 120KO vs WT.

Figure 3. Hyperglycemic clamp analysis of *gpr40* and *gpr120* single and double knockout

mice under chow diet. (**A-D**) Blood glucose (**A, B**) and plasma insulin (**C, D**) during the

hyperglycemic clamp (HGC) in male (**A, C**) and female mice (**B, D**). (**E-J**) Plasma c-peptide (**E,**

F) and glucose infusion rate (GIR) (**G, H**) during the steady state (50-80 min) and disposition

Index (DI) (**I, J**) in male (**E, G, I**) and female (**F, H, J**) mice. Data are presented as mean +/-

SEM (n=8-10 animals per group). (**A-D**) P-values were calculated by the mixed model of 2-way

ANOVA with Greisser-Greenhouse correction in case of variance heterogeneity (i.e. no

assumption of sphericity) and with Tukey's post-hoc multiple comparisons tests. No significant

effect of genotype on blood glucose or plasma insulin levels was detected during the steady state

(50-80 min) in both males and females. (**E-J**) P-values were calculated by Brown-Forsythe and

Welch's ANOVA (**G**) or ordinary one-way ANOVA (**E, F, H-J**) with Dunnett's T3 or Tukey's

post-hoc MCTs, respectively. **p<0.01, *p<0.05.

Figure 4. Energy balance in *gpr40* and *gpr120* single and double knockout (KO) mice under

chow (CD) or high-fat diet (HFD). (**A**) Study plan to assess energy and glucose homeostasis in

wild-type (WT), Gpr40KO (40KO), Gpr120KO (120KO) and Gpr120/40KO (120/40KO) male

and female mice fed CD or HFD from 8 to 20 weeks of age. (**B-G**) body weight (**B, C**), total

energy intake at 20 weeks of age (**D, E**) and fat and lean mass as a percentage of body weight

measured by Echo-MRI at 19 weeks of age (**F, G**) in male (**B, D, F**) and female (**C, E, G**) mice.

Data are presented as mean +/- SEM (n=7-10 animals per group). P-values were calculated by

regular 2-way ANOVA or mixed model when some follow-up values were missing with no

assumption of sphericity (**B, C**) or 2-way ANOVA with post-hoc Tukey's multiple comparisons

test (**D-G**). *p < 0.05.

Figure 5. Oral glucose tolerance in *gpr40* and *gpr120* single and double knockout (KO)

mice under chow (CD) or high-fat diet (HFD). (**A-H**) Blood glucose (**A, C, E, G**) and plasma

insulin (**B, D, F, H**) levels during an oGTT (glucose 1g/kg BW at 0 min) at 20 weeks of age in

male (**A, B, E, F**) and female (**C, D, G, H**) wild-type (WT), Gpr40KO (40KO), Gpr120KO

(120KO) and Gpr120/40KO (120/40KO) mice fed CD (**A-D**) or HFD (**E-H**) from 8 to 20 weeks

of age. Data are presented as mean +/- SEM (n=6-8 animals per group). P-values were calculated

by regular 2-way ANOVA or mixed model when some values were missing, with no assumption

of sphericity and with Tukey's multiple comparisons tests. **p<0.01 and *p<0.05, 120/40KO vs

WT.

Figure 6. Potentiation of GSIS in response to Cpd A and TAK-875 in islets ex vivo. Insulin

secretion was assessed in 1-h static incubations in response to 2.8 or 16.7 mM glucose with or

without the GPR120 agonists Cpd A (50 µM) and the GPR40 agonist TAK-875 (5 µM). (**A-C**)

Insulin secretion presented as a percentage of islet insulin content (**A**), insulin secretion (**B**) and

insulin content (**C**). Data represent individual values and are expressed as mean +/- SEM of 6

independent experiments. P-values were calculated by regular two-way ANOVA with Dunnett's

or Tukey's (G-J) post hoc adjustment for multiple comparisons and Welch/Brown-Forsythe

correction when necessary to compensate for SD variances. * p<0.05, ** p<0.005, *** p<0.001,

**** p<0.0001.

Figure 1

Figure 2

Figure 3   **Male**   **Female**

Figure 4

Figure 5

Figure 6