Brief Research Report

The effect of object type on building scene imagery – an MEG study

1   **Anna M. Monk, Gareth R. Barnes and Eleanor A. Maguire**[*]

2   Wellcome Centre for Human Neuroimaging, UCL Queen Square Institute of Neurology, University
3   College London, London, UK
4

* Correspondence:
Eleanor A. Maguire
e.maguire@ucl.ac.uk

Keywords: scenes; objects; space-defining; magnetoencephalography (MEG); hippocampus; theta;
vmPFC; DCM

ABSTRACT

Previous studies have reported that some objects evoke a sense of local three-dimensional space (space-
defining; SD), while others do not (space-ambiguous; SA), despite being imagined or viewed in
isolation devoid of a background context. Moreover, people show a strong preference for SD objects
when given a choice of objects with which to mentally construct scene imagery. When deconstructing
scenes, people retain significantly more SD objects than SA objects. It therefore seems that SD objects
might enjoy a privileged role in scene construction. In the current study, we leveraged the high temporal
resolution of magnetoencephalography (MEG) to compare the neural responses to SD and SA objects
while they were being used to build imagined scene representations, as this has not been examined
before using neuroimaging. On each trial, participants gradually built a scene image from three
successive auditorily-presented object descriptions and an imagined 3D space. We then examined the
neural dynamics associated with the points during scene construction when either SD or SA objects
were being imagined. We found that SD objects elicited theta changes relative to SA objects in two
brain regions, the right ventromedial prefrontal cortex (vmPFC) and right superior temporal gyrus
(STG). Furthermore, using dynamic causal modelling, we observed that the vmPFC drove STG
activity. These findings may indicate that SD objects serve to activate schematic and conceptual
knowledge in vmPFC and STG upon which scene representations are then built.

INTRODUCTION

Our lived experience of the world comprises a series of scenes that are perceived between the interruptions imposed by eye blinks and saccades. Indeed, scene mental imagery has been shown to dominate when people engage in critical cognitive functions such as recalling the past, imagining the future and spatial navigation (Clark et al., 2020; Andrews-Hanna et al., 2010; see also Clark et al., 2019). It is not surprising, therefore, that visual scenes have been deployed extensively as stimuli in cognitive neuroscience.

A scene is defined as a naturalistic three-dimensional spatially-coherent representation of the world typically populated by objects and viewed from an egocentric perspective (Dalton et al., 2018; Maguire and Mullally, 2013). Neuroimaging and neuropsychological studies have identified a number of brain areas that seem to be particularly engaged during the viewing and imagination of scenes including the ventromedial prefrontal cortex (vmPFC; Zeidman et al., 2015a; Bertossi et al., 2016; Barry et al., 2019a), the anterior hippocampus (Hassabis et al., 2007a, 2007b; Summerfield et al., 2010; Zeidman et al., 2015a, 2015b; Dalton et al., 2018; reviewed in Zeidman and Maguire, 2016), the posterior parahippocampal cortex (PHC; Epstein and Kanwisher, 1998; reviewed in Epstein, 2008 and Epstein and Baker, 2019), and the retrosplenial cortex (RSC; Park and Chun, 2009; reviewed in Epstein, 2008; Vann et al., 2009; Epstein and Baker, 2019). How are scene representations built, and what specific roles might these brain regions play? While spatial aspects of scenes have been amply investigated and linked to the hippocampus (Byrne et al., 2007; Morgan et al., 2011; Epstein et al., 2017; Epstein and Baker, 2019), the higher-order properties of objects within scenes have received comparatively less attention (Auger et al., 2012; Troiani et al., 2014; Julian et al., 2017; Epstein and Baker, 2019), and yet they could influence how scene representations are constructed by the brain.

One object attribute that seems to play a role in scene construction was reported by Mullally and Maguire (2011; see Kravitz et al., 2011 for related work). They observed that certain objects, when viewed or imagined in isolation, evoked a strong sense of three-dimensional local space surrounding them (space-defining (SD) objects), while others did not (space-ambiguous (SA) objects), and this was associated with engagement of the PHC during functional MRI (fMRI). In other words, SD objects seem to anchor or impose themselves on their surrounding space (e.g., the traffic light in Figure 1A) and in doing so they define the space around them. By contrast, SA objects do not anchor themselves in their surrounding space in the same way (e.g., a worn car tyre in Figure 1A). Participants in the Mullally and Maguire study (2011) were asked to describe the difference between an SD and an SA object, and a typical response included: "SD items conjure up a sense of space whereas SA items float - they go anywhere." Many participants spontaneously used the word "float" when discussing SA objects, emphasizing the detachment of these objects from an explicit sense of surrounding space. Mullally and Maguire (2011) also found that SD objects tended to be less portable, maintaining a permanent location, although a considerable number of SA objects had reasonably high permanence also (e.g., a large hay bale in Figure 1A), suggesting that permanence, while certainly related to SD, is not the sole basis of the SD effect. Indeed, object permanence has been linked to the RSC rather than the PHC (Auger et al., 2012, 2015; Auger and Maguire, 2013; Troiani et al., 2014). Similarly, Mullally and Maguire (2011) found that object size and contextual associations did not account for the SD effect.

In a subsequent behavioural study, participants showed a strong preference for SD objects when given a choice of objects with which to mentally construct scenes, even when comparatively larger and more permanent SA objects were available (Mullally and Maguire, 2013). Moreover, when deconstructing scenes, participants retained significantly more SD objects than SA objects. It therefore seems that SD objects might enjoy a privileged role in scene construction.

Mullally and Maguire (2011) examined SD and SA objects in isolation. However, given their
apparent influence during scene construction (Mullally and Maguire, 2013), in the current study we
compared neural responses to SD and SA objects while they were being used to build imagined scene
representations. We adapted a paradigm from Dalton et al. (2018) and Monk et al. (2020) where
participants gradually built a scene image from three successive auditorily-presented object
descriptions and an imagined 3D space. In order to capture the neural dynamics associated with the
points during scene construction when either SD or SA objects were being imagined, we leveraged the
high temporal resolution of magnetoencephalography (MEG). In previous MEG studies, changes in
vmPFC and anterior hippocampal theta were noted when participants imagined scenes in response to
scene-evoking cue words (Barry et al., 2019a, 2019b), and when scene imagery was gradually built
(Monk et al., 2020), but the effect, if any, of SD and SA objects on brain responses remains unknown.

Here, we performed a whole brain analysis, and characterized the effective connectivity
between any brain regions that emerged from this analysis. While our main interest was in theta, we
also examined other frequencies. The obvious prediction, given the previous Mullally and Maguire
(2011) fMRI study, was that PHC would be engaged by SD objects. However, because all stimuli were
scenes, and the key manipulation of SD and SA objects within scenes was so subtle, we retained an
open mind about which brain areas might distinguish between the two object types.

MATERIALS AND METHODS

Participants

Twenty-three healthy, right-handed people (13 females; mean age = 25.35 years; standard deviation =
3.69) participated in the experiment. Participants were either monolingual native English speakers, or
bilingual native English speakers (i.e., they had English as their first language, but were also but fluent
in a second language). They were reimbursed £10 per hour for taking part which was paid at study
completion. The study was approved by the University College London Research Ethics Committee
(project ID: 1825/005). All participants gave written informed consent in accordance with the
Declaration of Helsinki.

Stimuli

Each stimulus comprised auditorily-presented SD and SA object descriptions (see examples in Figure
1A) and a 3D space. We used the same auditory object descriptions as Dalton et al. (2018). As part of
this previous study, the objects were rated as SD or SA. Using these stimuli, we found that SD and SA
objects were matched on utterance length ($Z = 1.643$, $p < 0.1$) and number of syllables ($Z = 1.788$, $p <$
0.074). Unsurprisingly, as outlined above, SD objects were rated as more permanent than SA objects
($Z = 5.431$, $p < 0.001$). All objects were rated as highly imageable, obtaining a score of at least 4 on a
scale from 1 (not imageable) to 5 (extremely imageable). Objects in each triplet were not contextually
related to each other. Participants in the current MEG study were unaware of the SD-SA distinction.

Task and Procedure

The task, adapted from Dalton et al. (2018) and Monk et al. (2020), involved participants gradually
constructing simple scenes in their imagination from a combination of auditorily-presented SD and SA
object descriptions and a 3D space. The Cogent2000 toolbox (http://www.vislab.ucl.ac.uk/cogent.php)
run in Matlab was used to present stimuli and record responses in the MEG scanner. Auditory stimuli
were delivered via MEG-compatible earbuds. Each trial started with a visual cue (4 sec), which
displayed the configuration of locations at which objects should be imagined in the upcoming trial

(Figure 1B). Four different cue configurations (Figure 2) were randomized across the five scanning
blocks. We emphasized the importance of following the cue configurations as precisely as possible.
This ensured matched eye movements between the scene and array conditions, consistency across
participants, and that objects were imagined as separate and non-overlapping. Participants then fixated
on the screen center (1 sec). During the scene construction task (~9 sec) (Figure 1B), keeping their
eyes open whilst looking at a blank screen, participants first imagined a 3D grid covering
approximately the bottom two-thirds of the blank screen. Upon hearing each of three auditory
descriptions, one at a time, they imagined the objects in the separate, cue-specified positions on the 3D
grid. They were instructed to move their eyes to where they were imagining each object on the screen,
but also to maintain imagery of previous objects and the grid in their fixed positions. Each construction
stage consisted of a ~2 sec object description and a silent 1 sec gap before the presentation of the next
object. An additional 1 sec at the end of scene construction avoided an abrupt end to the task. By the
end of a trial, participants had created a mental image of a simple scene composed of a 3D grid and
three objects. Vividness of the entire scene was then rated on a scale of 1 (not vivid at all) to 5
(extremely vivid). An inter-trial interval (2 sec) preceded the next trial.

Participants imagined a total of 66 scenes (composed of 99 SD, 99 SA objects). Each object
description was heard only once. The order of presentation of SD and SA objects within triplets was
balanced across scenes with an equal number of SD and SA objects in the first, second and third
construction stages (33 in each). A control task (33 trials) involved participants attending to a backward
series of auditorily-presented numbers, and was designed to provide relief from the effortful
imagination task; it was not subject to analysis. Seven additional catch trials (5 scenes, 2 counting)
were also included and pseudorandomly presented across blocks to ensure that participants sustained
attention - participants pressed a button upon hearing a repeated object description or number within a
triplet.

The visit of each participant took 2.5 hours in total (including training, set up in the MEG scanner
and eye tracker calibration, the experimental task (~50 minutes) and the post-scan memory test).

Eye Tracking

Eye movements were recorded during the MEG scan using an Eyelink 1000 Plus (SR Research) eye
tracking system with a sampling rate of 2000 Hz. The right eye was used for calibration and data
acquisition. For some participants the calibration was insufficiently accurate, leaving 19 data sets for
the eye tracking analyses.

Surprise Post-Scan Memory Test

If we found a neural difference between SD and SA objects, it could be argued that this was due to a
difference in encoding success. For example, perhaps SD objects were more readily encoded into
memory than SA objects, even though there was no explicit memory encoding requirement in this task.
By conducting a surprise memory test for SD and SA objects immediately post-scan, we could examine
this issue. Participants were presented with a randomized order of all previously heard auditory object
descriptions and an additional 33 SD and 33 SA new object description (lures). After hearing each
item, they indicated whether or not they had heard the object description during the scan, and their
confidence in their decision (1 = low, to 5 = high).

Behavioral Data Analysis

In-scanner vividness was compared between SD-majority (where 2 out of the 3 objects were SD) and
SA-majority scenes (where 2 out of the 3 objects were SA) using a paired-samples t-test. Eye tracking
data were analyzed using two-way repeated measures ANOVAs. Memory performance was assessed
using the sensitivity index *d'*, based on Signal Detection Theory (Macmillan and Creelman, 1990). We
also quantified the bias of participants, to control for any tendency participants might have to give one
type of response more than another. This response bias *'c'* was calculated using the formula: c = -
0.5(Hit Rate + False Alarm Rate). Differences in *d'* and *c* as a function of object type (SD, SA) and
construction stage (first, second, third) were each analyzed using a two-way repeated measures
ANOVA. Statistical analyses were performed in SPSS25 using a significance threshold of $p < 0.05$. In
cases where Mauchly's test found sphericity violated, Greenhouse-Geisser adjusted degrees of freedom
were applied.

MEG Data Acquisition and Preprocessing

172   MEG data were recorded using a 275 channel CTF Omega MEG system with a sampling rate of
173   1200 Hz. Head position fiducial coils were attached to the three standard fiducial points (nasion, left
174   and right preauricular) to monitor head position continuously throughout acquisition. Recordings were
175   filtered with a 1 Hz high-pass filter, 48-52 Hz stop-band filter, and 98-102 Hz stop-band filter, to
176   remove slow drifts in signals from the MEG sensors and power line interference.

MEG Data Analysis

178   All MEG analyses were conducted using SPM12 (www.fil.ion.ucl.ac.uk/spm). Source reconstruction
179   was performed using the DAiSS toolbox (https://github.com/SPM/DAiSS) and visualized using
180   MRIcroGL (https://www.mccauslandcenter.sc.edu/mricrogl).

Source Reconstruction

Epochs corresponding to each construction period were defined as 0-3 sec relative to the onset of the
SD or SA object description, and concatenated across scanning blocks. Source reconstruction was
performed using a linearly constrained minimum variance (LCMV) beamformer (Van Veen et al.,
1997). This approach allowed us to estimate power differences between SD and SA objects in source
space within selected frequency bands: theta (4-8 Hz), alpha (9-12 Hz), beta (13-30 Hz), and gamma
(30-100 Hz).

For each participant, a set of filter weights was built based on data from the SD and SA conditions
within each frequency band, and a 0-3 sec time window encapsulating a construction period.
Coregistration to MNI space was performed using a 5 mm volumetric grid and was based on nasion,
left and right preauricular fiducials. The forward model was computed using a single-shell head model
(Nolte, 2003). At the first level, power in a particular frequency band was estimated to create one image
per object type (SD or SA) per participant. Images were spatially smoothed using a 12 mm Gaussian
kernel and entered into a second-level random effects paired t-test to determine power differences
between SD objects and SA objects across the whole brain. An uncorrected threshold of $p < 0.001$ with
a cluster extent threshold of >100 voxels was applied to each contrast, in line with numerous other
MEG studies (e.g., Barry et al., 2019a; Cushing et al., 2018; Hanslmayr et al., 2011; Kaplan et al.,
2012; Kveraga et al., 2011; Lieberman & Cunningham, 2009). This is held to provide a balance
between protecting against false positives whilst enabling the detection of subtler signals. The same
beamforming protocol was followed when objects were re-categorized as permanent and non-
permanent, with the number of permanent and non-permanent objects equalized to 65 in each category.

The contrast of primary interest was the direct comparison between SD and SA objects. As a
secondary issue, we were mindful that it was also of interest to know whether an effect represented an
increase or a decrease in power from baseline. Including baseline correction in the original beamformer
would be the usual way to examine this. However, this is challenging to implement in the case of the
current design because an SD or SA object was not necessarily preceded by a clean baseline. On each
trial, three object descriptions were heard one after the other. While the first object in a triplet was
preceded by a clean 1 sec fixation baseline, objects 2 and 3 had an object imagination stage preceding
them. Objects at different construction stages would therefore be corrected against different types of
baseline (i.e., object 1 vs preceding fixation, object 2 vs preceding object 1, and object 3 vs preceding
object 2). Conducting this analysis could have introduced spurious effects. Consequently, having
established that a difference was apparent between SD and SA objects when they were compared
directly, we then sought to ascertain the direction of power change. By using a separate beamformer
where the 1 sec pre-stimulus fixation period was the only baseline for all objects (whether a SD/SA or
first/second/third object), and hence did not overlap with any stimulus period, we were able to establish
the direction of power change for SD and SA objects in a straightforward way.

Dynamic Causal Modelling (DCM)

Brain areas identified in the whole brain SD versus SA beamformer provided the seed regions for the
subsequent effective connectivity analysis, which was conducted using DCM for cross spectral
densities (Moran et al., 2009). This approach permitted us to compare different biologically plausible
models of how one brain region influences another, as well as mutual entrainment between regions
(Friston, 2009; Kahan and Foltynie, 2013). Random-effects Bayesian model selection (BMS) was
performed to compare the evidence for each specified model that varied according to which
connections were modulated by SD relative to SA objects (Klaas et al., 2009). We determined the
winning model to be the one with the greatest exceedance probability. To assess the consistency of the
model fit, we also calculated the log Bayes factor for each participant.

RESULTS

Behavioral Data

There was no significant difference in the vividness of mental imagery between SD-majority ($M =$
3.91, standard deviation = 0.69) and SA-majority ($M = 3.89$, standard deviation = 0.66) scene trials
($t_{(22)} = 0.464$, $p = 0.647$). Participants correctly identified on average 97.52% (standard deviation =
0.39) of catch trials, indicating that they attended throughout the experiment.

The effect of object type (SD, SA) and construction stage (first, second, third) on eye-
movement fixation count (Fix_{Count}) and fixation duration (Fix_{Dur}) showed that there were no significant
main effects of object type (Fix_{Count}: $F_{(1,18)} = 1.908$, $p = 0.184$; Fix_{Dur}: $F_{(1,18)} = 0.086$, $p = 0.772$) or
construction stage (Fix_{Count}: $F_{(2,36)} = 0.292$, $p = 0.748$; Fix_{Dur}: $F_{(2,36)} = 0.535$, $p = 0.590$), and no object
type×construction stage interaction (Fix_{Count}: $F_{(2,36)} = 0.710$, $p = 0.499$; Fix_{Dur}: $F_{(2,36)} = 1.871$, $p =$
0.169). Heat maps of the spatial patterns of fixations during the task demonstrated a consistent
adherence to cue configuration instructions across participants (Figure 2).

In terms of recognition memory (see Table 1), performance exceeded 80% correct for both SD
and SA objects, and for *d'* and *c* there were no significant effects of object type (*d'*: $F_{(1,22)} = 0.469$, p
= 0.500 ; *c*: $F_{(1,22)} = 0.012$, $p = 0.915$), construction stage (*d'*: $F_{(2,44)} = 2.383$, $p = 0.104$; *c*: $F_{(2,44)} =$
0.120, $p = 0.887$), nor were there any interactions (*d'*: $F_{(2,44)} = 1.431$, $p = 0.250$; *c*: $F_{(2,44)} = 0.035$, $p =$
0.965).

MEG Data

Power Changes

A whole brain beamforming analysis revealed significant theta power attenuation for SD compared to
SA objects in only two regions: the right vmPFC (peak MNI = 12, 60, -8; *t*-value = 3.66; cluster size
= 1960) and right superior temporal gyrus (STG; peak MNI = 66, -6, -12; *t*-value = 3.76; cluster size
= 1197) (Figure 3A).

A subsequent contrast between each object type and the baseline revealed that the theta power
changes were decreases, echoing numerous previous reports of power attenuation during the
construction of scene imagery (e.g., Guderian et al., 2009; Barry et al., 2019a, 2019b) and memory
recall (e.g., Solomon et al., 2019; McCormick et al., 2020).

We did not observe any significant differences in theta power between permanent and non-
permanent objects across the whole brain.

Analysis of alpha, beta and gamma showed no significant power differences across the whole
brain when SD and SA objects were compared.

Effective Connectivity

Having established a response to object type in the vmPFC and STG, we next sought to examine the
effective connectivity between these regions. We tested three simple hypotheses: (1) vmPFC and STG
are mutually entrained, (2) STG drives vmPFC, or (3) vmPFC drives STG. We embodied each
hypothesis as a DCM where models differed in which connection could be modulated by SD relative
to SA objects. BMS identified the winning model to be vmPFC driving STG during SD more so than
SA objects, with an exceedance probability of 91.62% (Figure 3B, left panel). This model was also
the most consistent across participants (Figure 3B, right panel).

DISCUSSION

In this study we focused on an object property, SD-SA, that has been shown to influence how scene
imagery is constructed (Mullally and Maguire, 2013). We found that while these object types were
being imagined during scene construction, SD objects elicited significant theta changes relative to SA
objects in two brain regions, the right vmPFC and right STG. Furthermore, the vmPFC drove STG
theta activity.

SD and SA objects were matched in terms of the vividness of mental imagery, eye movements
and incidental memory encoding. All objects were incorporated into the same three-object scene
structures within which the order of SD or SA object presentation and object locations were carefully
controlled. We also examined object permanence, and found that this property did not engage the
vmPFC or STG. Our findings are therefore unlikely to be explained by these factors.

Most of the brain regions typically associated with scenes did not respond to SD objects during
scene construction. This is likely because scene processing was constant throughout the experiment,
and so there was no variation required in the activity of these areas. It is notable that the PHC, which
was active during fMRI in response to SD objects when they were viewed or imagined in isolation and
devoid of a scene context (Mullally and Maguire, 2011), did not exhibit power changes during scene
construction. It may be that examining objects in isolation afforded a "purer" expression of SD

whereas, once these objects were included in scene building, higher-order areas then came online to
direct their use in constructing scene representations, a possibility that we discuss next.

Considering first the right STG, while this region has been linked to speech processing (e.g.,
Hullett et al., 2016), the close matching of auditory stimuli and the absence of activity changes in other
auditory areas suggests this factor does not account for its responsivity to SD objects. Perhaps more
germane is the location of the STG within the anterior temporal lobe, a key neural substrate of semantic
and conceptual knowledge that supports object recognition (Peelen and Caramazza, 2012; Chiou and
Lambon Ralph, 2016). Patients with semantic dementia, caused by atrophy to the anterior temporal
lobe, lose conceptual but not perceptual knowledge about common objects (Campo et al., 2013; Guo
et al., 2013).

This could mean that SD objects provide conceptual information that is registered by the STG.
Why might this be relevant to scene construction? Prior expectations have a striking top-down
modulatory influence on our perception of the world, enabling us to process complex surroundings in
an efficient manner (Summerfield and Egner, 2009), and resolve ambiguity (Chiou and Lambon Ralph,
2016). Without this knowledge, we are unable to understand how and where an object should be used
(Peelen and Caramazza, 2012). Therefore, objects are an important source of information about the
category of scene being imagined (or viewed), facilitating a rapid, efficient interpretation of the scene
'gist' without the need to process every component of a scene (Oliva and Torralba, 2006; Summerfield
and Egner, 2009; Clarke and Tyler, 2015; Trapp and Bar, 2015). For example, if we see a park bench
this might indicate the scene is from a park. Although in the current study the scenes were deliberately
composed of semantically unrelated objects, this may not have impeded the STG in nevertheless
registering SD objects more so than SA objects because SD objects would normally offer useful
conceptual information to help anchor a scene.

The operation of the right STG might be facilitated by the right vmPFC. Converging evidence
across multiple studies has shown that the part of the vmPFC that was active in response to SD objects
plays a role in the abstraction of key features across multiple episodes (Roy et al., 2012). These
contribute to the formation of schemas, which are internal models of the world representing elements
that likely exist in a prototypical scene, based on previous exposure to such scenes (Tse et al., 2007;
van Kesteren et al., 2013; Gilboa and Marlatte, 2017). For instance, a park typically contains benches,
trees and flowers. SD objects may be particularly useful in building scene schema, and hence the
response to them by the vmPFC.

Patients with damage to the vmPFC exhibit deficits that suggest aberrant schema re-activation
(Ciaramelli et al., 2006; Gilboa et al., 2006; Ghosh et al., 2014), and this has led to the proposal that
vmPFC may activate relevant schema to orchestrate the mental construction of scenes performed
elsewhere – for example, in the hippocampus (McCormick et al., 2018; Ciaramelli et al., 2019; Monk
et al., 2020). Our DCM findings extend this work by showing that the right vmPFC also exerts
influence over the right STG, indicating it may be engaging in top-down modulation of conceptual
object processing by the STG, specifically during the processing of SD objects. Our results may
therefore indicate that SD objects help to define a scene by priming relevant schemas in the vmPFC
which then guide conceptual processing in areas such as the STG.

There is another possible explanation for our findings. In the current study, the scenes were
deliberately composed of semantically unrelated objects, and this could have introduced ambiguity
about a scene's identity. vmPFC and STG engagement may therefore be evidence of additional neural
processing that was required to resolve incongruences inherent to acontextual scenes (Chiou and

Lambon Ralph, 2016; Brandman and Peelen, 2017; Epstein and Baker, 2019), perhaps by drawing
upon existing schemas in the pursuit of an appropriate scene template. Indeed, connectivity between
medial prefrontal and medial temporal cortex has been shown to increase when novel information that
was less congruent with pre-existing schematic representations was processed (van Kesteren et al.,
2010; Chiou and Lambon Ralph, 2016). The acontextual nature of the scenes, and the effortful nature
of the scene construction task, may also have precluded observation of a schema-related memory
advantage for SD objects (see more on schema and memory in McCormick et al., 2018; Gilboa and
Marlatte, 2017; Tse et al., 2007). It should be noted, however, that our study was not designed to
investigate schema, and consequently these possible interpretations remain speculative. Future studies
will be needed to further elucidate the SD-SA difference revealed here, perhaps by comparing
semantically related and unrelated objects during scene construction, and by adapting the current
paradigm to test patients with vmPFC or STG damage. Another notable feature of our findings was the
right hemisphere location of the responses. This may be related to the visual nature of the imagined
scenes. This too could be probed further in future studies by comparing patients with left and right-
sided lesions.

In conclusion, this study revealed the neural dynamics associated with a specific object property
during scene construction, and we suggest that SD objects in particular may serve to activate schematic
and conceptual knowledge in vmPFC and STG upon which scene representations are then built.

DATA AVAILABILITY STATEMENT

348 The raw data supporting the conclusions of this manuscript, and the test materials, will be made
349 available by the authors to any qualified researcher upon request. Requests can be sent to
350 e.maguire@ucl.ac.uk.

ETHICS STATEMENT

352 The study was approved by the University College London Research Ethics Committee (project ID:
353 1825/005). All participants gave written informed consent in accordance with the Declaration of
354 Helsinki.

AUTHOR CONTRIBUTIONS

356 AMM and EAM designed the study. AMM collected and analyzed the data with input from EAM and
357 GRB. AMM and EAM wrote the manuscript. All authors contributed to manuscript revision, read and
358 approved the submitted version.

FUNDING

360 This work was supported by a Wellcome Principal Research Fellowship to EAM (210567/Z/18/Z) and
361 the Centre by a Centre Award from the Wellcome Trust (203147/Z/16/Z).

ACKNOWLEDGEMENTS

Thanks to Marshall Dalton for providing the task stimuli, Peter Zeidman for his DCM advice, and
Daniel Bates, David Bradbury and Eric Featherstone for technical support.

CONFLICTS OF INTEREST

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

REFERENCES

Andrews-Hanna, J. R., Reidler, J. S., Sepulcre, J., Poulin, R., and Buckner, R. L. (2010). Functional-
anatomic fractionation of the brain's default network. *Neuron* 65, 550–562.
doi:10.1016/j.neuron.2010.02.005.

Auger, S. D., and Maguire, E. A. (2013). Assessing the mechanism of response in the retrosplenial
cortex of good and poor navigators. *Cortex* 49, 2904–2913. doi:10.1016/j.cortex.2013.08.002.

Auger, S. D., Mullally, S. L., and Maguire, E. A. (2012). Retrosplenial Cortex Codes for Permanent
Landmarks. *PLoS One* 7, e43620. doi:10.1371/journal.pone.0043620.

Auger, S. D., Zeidman, P., and Maguire, E. A. (2015). A central role for the retrosplenial cortex in de
novo environmental learning. *Elife* 4. doi:10.7554/eLife.09031.

Barry, D. N., Barnes, G. R., Clark, I. A., and Maguire, E. A. (2019a). The neural dynamics of novel
scene imagery. *J. Neurosci.* 39, 4375– 4386. doi:10.1523/JNEUROSCI.2497-18.2019.

Barry, D. N., Tierney, T. M., Holmes, N., Boto, E., Roberts, G., Leggett, J., et al. (2019b). Imaging
the human hippocampus with optically-pumped magnetoencephalography. *Neuroimage* 203.
doi:10.1016/j.neuroimage.2019.116192.

Bertossi, E., Aleo, F., Braghittoni, D., and Ciaramelli, E. (2016). Stuck in the here and now:
Construction of fictitious and future experiences following ventromedial prefrontal damage.
Neuropsychologia 81, 107–116. doi:10.1016/j.neuropsychologia.2015.12.015.

Brandman, T., and Peelen, M. V. (2017). Interaction between scene and object processing revealed
by human fMRI and MEG decoding. *J. Neurosci.* 37, 0582–17.
doi:10.1523/JNEUROSCI.0582-17.2017.

Byrne, P., Becker, S., and Burgess, N. (2007). Remembering the past and imagining the future: A
neural model of spatial memory and imagery. *Psychol. Rev.* 114, 340–375. doi:10.1037/0033-
295X.114.2.340.

Campo, P., Poch, C., Toledano, R., Igoa, J. M., Belinchón, M., García-Morales, I., et al. (2013).
Anterobasal temporal lobe lesions alter recurrent functional connectivity within the ventral
pathway during naming. *J. Neurosci.* 33, 12679–12688. doi:10.1523/JNEUROSCI.0645-
13.2013.

Chiou, R., and Lambon Ralph, M. A. (2016). The anterior temporal cortex is a primary semantic
source of top-down influences on object recognition. *Cortex* 79, 75–86.
doi:10.1016/j.cortex.2016.03.007.

Ciaramelli, E., De Luca, F., Monk, A. M., McCormick, C., and Maguire, E. A. (2019). What "wins"
in VMPFC: Scenes, situations, or schema? *Neurosci. Biobehav. Rev.* 100, 208–210.
doi:10.1016/j.neubiorev.2019.03.001.

Ciaramelli, E., Ghetti, S., Frattarelli, M., and Làdavas, E. (2006). When true memory availability
promotes false memory: Evidence from confabulating patients. *Neuropsychologia* 44, 1866–
1877. doi:10.1016/j.neuropsychologia.2006.02.008.

Clark, I. A., Hotchin, V., Monk, A., Pizzamiglio, G., Liefgreen, A., and Maguire, E. A. (2019).
Identifying the cognitive processes underpinning hippocampal-dependent tasks. *J. Exp.*
Psychol. Gen. 148, 1861–1881. doi:10.1037/xge0000582.

Clark, I. A., Monk, A. M., Maguire, E. A. (2020). Characterising strategy use during the performance
of hippocampal-dependent tasks. *Frontiers in Psychol.* 11, 2119.
https://www.biorxiv.org/content/10.1101/807990v3

Clarke, A., and Tyler, L. K. (2015). Understanding what we see: How we derive meaning from
vision. *Trends Cogn. Sci.* 19, 677–687. doi:10.1016/j.tics.2015.08.008.

Cushing, C. A., Im, H. Y., Adams Jr., R. B., Ward, N., Albohn, D. N., Steiner, T. G., and Kverga, K.
(2018). Neurodynamics and connectivity during facial fear perception: The role of threat
exposure and signal congruity. *Scientific Reports* 8. doi: 10.1038/s41598-018-20509-8.

Dalton, M. A., Zeidman, P., McCormick, C., and Maguire, E. A. (2018). Differentiable processing of
objects, associations, and scenes within the hippocampus. *J. Neurosci.* 38, 8146–8159.
doi:10.1523/JNEUROSCI.0263-18.2018.

Epstein, R. A. (2008). Parahippocampal and retrosplenial contributions to human spatial navigation.
Trends Cogn. Sci. 12, 388–396. doi:10.1016/j.tics.2008.07.004.

Epstein, R. A., and Baker, C. I. (2019). Scene perception in the human brain. *Annu. Rev. Vis. Sci.* 5,
373–397. doi:10.1146/annurev-vision-091718-014809.

Epstein, R. A., Patai, E. Z., Julian, J. B., and Spiers, H. J. (2017). The cognitive map in humans:
Spatial navigation and beyond. *Nat. Neurosci.* 20, 1504–1513. doi:10.1038/nn.4656.

Epstein, R., and Kanwisher, N. (1998). A cortical representation the local visual environment. *Nature*
392, 598–601. doi:10.1038/33402.

Friston, K. (2009). Causal modelling and brain connectivity in functional magnetic resonance
imaging. *PLoS Biol.* 7, e1000033. doi:10.1371/journal.pbio.1000033.

Ghosh, V. E., Moscovitch, M., Colella, B. M., and Gilboa, A. (2014). Schema representation in
patients with ventromedial PFC lesions. *J. Neurosci.* 34, 12057–12070.
doi:10.1523/JNEUROSCI.0740-14.2014.

Gilboa, A., Alain, C., Stuss, D. T., Melo, B., Miller, S., and Moscovitch, M. (2006). Mechanisms of
spontaneous confabulations: A strategic retrieval account. *Brain* 129, 1399–1414.
doi:10.1093/brain/awl093.

Gilboa, A., and Marlatte, H. (2017). Neurobiology of schemas and schema-mediated memory.
Trends Cogn. Sci. 21, 618–631. doi:10.1016/j.tics.2017.04.013.

Guderian, S., Schott, B. H., Richardson-Klavehn, A., and Düzel, E. (2009). Medial temporal theta
state before an event predicts episodic encoding success in humans. *Proc. Natl. Acad. Sci. U.*
S. A. 106, 5365–5370. doi:10.1073/pnas.0900289106.

Guo, C. C., Gorno-Tempini, M. L., Gesierich, B., Henry, M., Trujillo, A., Shany-Ur, T., et al. (2013).
Anterior temporal lobe degeneration produces widespread network-driven dysfunction. *Brain*
136, 2979–2991. doi:10.1093/brain/awt222.

Hanslmayr, S., Volberg, G., Wimber, M., Raabe, M., Greenlee, M. W., and Bäuml, K.-H. T. (2011).
The relationship between brain oscillations and BOLD signal during memory formation: A
combined EEG–fMRI study. *J of Neurosci.* 31, 15674-15680. doi:
10.1523/JNEUROSCI.3140-11.2011

Hassabis, D., Kumaran, D., and Maguire, E. A. (2007a). Using imagination to understand the neural
basis of episodic memory. *J. Neurosci.* 27, 14365–14374. doi:10.1523/JNEUROSCI.4549-
07.2007.

Hassabis, D., Kumaran, D., Vann, S. D., and Maguire, E. A. (2007b). Patients with hippocampal
amnesia cannot imagine new experiences. *Proc. Natl. Acad. Sci. U. S. A.* 104, 1726–1731.
doi:10.1073/pnas.0610561104.

Hullett, P. W., Hamilton, L. S., Mesgarani, N., Schreiner, C. E., and Chang, E. F. (2016). Human
superior temporal gyrus organization of spectrotemporal modulation tuning derived from
speech stimuli. *J. Neurosci.* 36, 2014–2026. doi:10.1523/JNEUROSCI.1779-15.2016.

Julian, J. B., Ryan, J., and Epstein, R. A. (2017). Coding of object size and object category in human
visual cortex. Cereb. *Cortex* 27, 3095–3109. doi:10.1093/cercor/bhw150.

Kahan, J., and Foltynie, T. (2013). Understanding DCM: Ten simple rules for the clinician.
Neuroimage 83, 542–549. doi:10.1016/j.neuroimage.2013.07.008.

Kaplan, R., Barnes, G. R., Duzel, E., and Burgess, N. (2012). Movement-related theta rhythm in
humans: Coordinating self-directed hippocampal learning. *PLoS Biology* 10. doi:
10.1371/journal.pbio.1001267.

Kravitz, D. J., Peng, C. S., and Baker, C. I. (2011). Real-world scene representations in high-level
visual cortex: It's the spaces more than the places. *J. Neurosci.* 31, 7322–7333.
doi:10.1523/JNEUROSCI.4588-10.2011.

Kveraga, K., Ghuman, A. S., Kassam, K. S., Aminoff, E. A., Hamalainen, M. S., Chaumon, M. and
Bar, M. (2011). Early onset of neural synchronization in the contextual associations network.
Proc. Natl. Acad. Sci. U. S. A. 108, 3389–3394. doi: 10.1073/pnas.1013760108.

Lieberman, M. D., and Cunningham, W. A. (2009). Type I and Type II error concerns in fMRI
research: Re-balancing the scale. *Soc. Cogn. Affect. Neur.* 4, 423–428. doi:
10.1093/scan/nsp052.

Macmillan, N. A., and Creelman, C. D. (1990). Response bias: Characteristics of detection theory,
473   threshold theory, and "nonparametric" indexes. *Psychol. Bull.* 107, 401–413.
474   doi:10.1037/0033-2909.107.3.401.

475   Maguire, E. A., and Mullally, S. L. (2013). The hippocampus: A manifesto for change. *J. Exp.*
476   *Psychol. Gen.* 142, 1180–1189. doi:10.1037/a0033650.

477   McCormick, C., Barry, D. N., Jafarian, A., Barnes, G. R., and Maguire, E. A. (2020). vmPFC Drives
478   Hippocampal Processing during Autobiographical Memory Recall Regardless of Remoteness.
479   *Cereb. Cortex* (in press). doi:10.1093/cercor/bhaa172.

480   McCormick, C., Ciaramelli, E., De Luca, F., and Maguire, E. A. (2018). Comparing and contrasting
481   the cognitive effects of hippocampal and ventromedial prefrontal cortex damage: A review of
482   human lesion studies. *Neuroscience* 374, 295–318. doi:10.1016/j.neuroscience.2017.07.066.

483   Monk, A. M., Dalton, M. A., Barnes, G. R., and Maguire, E. A. (2020). The role of hippocampal-
484   vmPFC neural dynamics in building mental representations. *J. Cogn. Neurosci.* (in press)
485   https://www.biorxiv.org/content/10.1101/2020.04.30.069765v2.

486   Moran, R. J., Stephan, K. E., Seidenbecher, T., Pape, H. C., Dolan, R. J., and Friston, K. J. (2009).
487   Dynamic causal models of steady-state responses. *Neuroimage* 44, 796–811.
488   doi:10.1016/j.neuroimage.2008.09.048.

489   Morgan, L. K., MacEvoy, S. P., Aguirre, G. K., and Epstein, R. A. (2011). Distances between real-
490   world locations are represented in the human hippocampus. *J. Neurosci.* 31, 1238–1245.
491   doi:10.1523/JNEUROSCI.4667-10.2011.

492   Mullally, S. L., and Maguire, E. A. (2011). A new role for the parahippocampal cortex in
493   representing space. *J. Neurosci.* 31, 7441–7449. doi:10.1523/JNEUROSCI.0267-11.2011.

494   Mullally, S. L., and Maguire, E. A. (2013). Exploring the role of space-defining objects in
495   constructing and maintaining imagined scenes. *Brain Cogn.* 82, 100–107.
496   doi:10.1016/j.bandc.2013.02.013.

497   Nolte, G. (2003). The magnetic lead field theorem in the quasi-static approximation and its use for
498   magnetoenchephalography forward calculation in realistic volume conductors. *Phys. Med.*
499   *Biol.* 48, 3637–3652. doi:10.1088/0031-9155/48/22/002.

500   Oliva, A., and Torralba, A. (2006). Building the gist of a scene: the role of global image features in
501   recognition. *Prog. Brain Res.* 155, 23–36. doi:10.1016/S0079-6123(06)55002-2.

502   Park, S., and Chun, M. M. (2009). Different roles of the parahippocampal place area (PPA) and
503   retrosplenial cortex (RSC) in panoramic scene perception. *Neuroimage* 47, 1747–1756.
504   doi:10.1016/j.neuroimage.2009.04.058.

505   Peelen, M. V, and Caramazza, A. (2012). Conceptual object representations in human anterior
506   temporal cortex. *J. Neurosci.* 32, 15728–15736. doi:10.1523/JNEUROSCI.1953-12.2012.

Roy, M., Shohamy, D., and Wager, T. D. (2012). Ventromedial prefrontal-subcortical systems and
the generation of affective meaning. *Trends Cogn. Sci.* 16, 147–156.
doi:10.1016/j.tics.2012.01.005.

Solomon, E. A., Stein, J. M., Das, S., Gorniak, R., Sperling, M. R., Worrell, G., et al. (2019).
Dynamic theta networks in the human medial temporal lobe support episodic memory. *Curr.*
Biol. 29, 1100-1111.e4. doi:10.1016/j.cub.2019.02.020.

Stephan, K. E., Penny, W. D., Daunizeau, J., Moran, R. J., and Friston, K. J. (2009). Bayesian model
selection for group studies. *Neuroimage* 46, 1004–1017.
doi:10.1016/j.neuroimage.2009.03.025.

Summerfield, C., and Egner, T. (2009). Expectation (and attention) in visual cognition. *Trends Cogn.*
Sci. 13, 403–409. doi:10.1016/j.tics.2009.06.003.

Summerfield, J. J., Hassabis, D., and Maguire, E. A. (2010). Differential engagement of brain regions
within a "core" network during scene construction. *Neuropsychologia* 48, 1501–1509.
doi:10.1016/j.neuropsychologia.2010.01.022.

Trapp, S., and Bar, M. (2015). Prediction, context, and competition in visual recognition. *Ann. N. Y.*
Acad. Sci. 1339, 190–198. doi:10.1111/nyas.12680.

Troiani, V., Stigliani, A., Smith, M. E., and Epstein, R. A. (2014). Multiple object properties drive
scene-selective regions. Cereb. *Cortex* 24, 883–897. doi:10.1093/cercor/bhs364.

Tse D., Langston, R. F., Kakeyama, M., Bethus, I., Spooner, P. A., Wood, E. R, Witter, M. P., and
Morris, R. G. M. (2007). Schemas and memory consolidation. *Science* 316, 76–82. doi:
10.1126/science.1135935

van Kesteren, M. T. R., Beul, S. F., Takashima, A., Henson, R. N., Ruiter, D. J., and Fernández, G.
(2013). Differential roles for medial prefrontal and medial temporal cortices in schema-
dependent encoding: From congruent to incongruent. *Neuropsychologia* 51, 2352–2359.
doi:10.1016/j.neuropsychologia.2013.05.027.

van Kesteren, M. T. R., Fernandez, G., Norris, D. G., and Hermans, E. J. (2010). Persistent schema-
dependent hippocampal-neocortical connectivity during memory encoding and postencoding
rest in humans. *Proc. Natl. Acad. Sci.* 107, 7550–7555. doi:10.1073/pnas.0914892107.

Van Veen, B. D., Van Drongelen, W., Yuchtman, M., and Suzuki, A. (1997). Localization of brain
electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Trans.*
Biomed. Eng. 44, 867–880. doi:10.1109/10.623056.

Vann, S. D., Aggleton, J. P., and Maguire, E. A. (2009). What does the retrosplenial cortex do? *Nat.*
Rev. Neurosci. 10, 792–802. doi:10.1038/nrn2733.

Zeidman, P., Mullally, S. L., and Maguire, E. A. (2015a). Constructing, perceiving, and maintaining
scenes: Hippocampal activity and connectivity. *Cereb. Cortex* 25, 3836–3855.
doi:10.1093/cercor/bhu266.

Zeidman, P., Lutti, A., and Maguire, E. A. (2015b). Investigating the functions of subregions within
anterior hippocampus. *Cortex* 73, 240–256. doi:10.1016/j.cortex.2015.09.002.

Zeidman, P., and Maguire, E. A. (2016). Anterior hippocampus: The anatomy of perception,
imagination and episodic memory. *Nat. Rev. Neurosci.* 17, 173–182.
doi:10.1038/nrn.2015.24.

TABLE 1 Results of the surprise post-scan object recognition memory test.
550

	SD objects	SA objects
	Mean (Standard Deviation)	**Mean (Standard Deviation)**
% correct	81.063 (7.332)	80.074 (8.198)
Hit rate	0.809 (0.079)	0.801 (0.091)
False alarm rate	0.156 (0.108)	0.156 (0.103)
d'	2.045 (0.581)	2.001 (0.611)
c	0.091 (0.282)	0.088 (0.272)

551 *Percent (%) correct, hit rate, false alarm rate, dprime (d') and response bias (c) discrimination*
552 *parameters for each object category.*

FIGURE 1 Example stimuli and trial structure. **(A)** Examples of SD and SA object descriptions. **(B)** The structure and timings of an example trial. Note that participants never saw visual objects. During the task the participants imagined the simple scenes while looking at a blank screen.

564   **FIGURE 2** Eye movement results. Heat maps of the fixation count during the 9 sec mental construction
565   period following each cue configuration. Each heat map is an aggregate of fixations on the blank screen
566   across all trials for that cue configuration across all participants with eye tracking data (n=19). Red
567   indicates higher fixation density and green lower fixation density.

FIGURE 3 MEG results. **(A)** Source reconstruction of theta (4-8 Hz) power changes during SD relative to SA objects revealed attenuation in the right ventromedial prefrontal cortex (R_vmPFC) and right superior temporal gyrus (R_STG). Images are superimposed on the MNI 152 template and thresholded at uncorrected $p < 0.001$. **(B)** Effective connectivity between R_vmPFC and R_STG was examined using DCM. Three models were compared, with R_vmPFC driving R_STG theta activity during SD compared to SA objects being the model that best explained the data (left panel). Log Bayes factors per participant (right panel) showed positive to strong evidence for this model in most participants. Participants for whom there was no conclusive evidence for either model are represented by black bars. Where log Bayes factors exceeded five, bars are truncated and the exact values are adjacently displayed.