Gut microbiome composition predicts summer core range size in a generalist and specialist ungulate

J.F. Wolf [1] (https://orcid.org/0000-0003-0773-4456), K.D. Kriss[2], K.M. MacAulay[2] (https://orcid.org/0000-0003-1001-6906), K. Munro[1,3], B.R. Patterson[1,4], and A.B.A. Shafer[1,5] (http://orcid.org/0000-0001-7652-225X)

[1] Trent University, Department of Environmental and Life Sciences, Peterborough, Ontario, K9L 0G2, Canada

[2] Ministry of Forests, Lands, and Natural Resource Operations, and Rural Development, Smithers, British Columbia V0J 2N0, Canada

[3] Ontario Federation of Anglers and Hunters, 4601 Guthrie Drive, Peterborough, Ontario, K9J 8L5, Canada

[4] Ontario Ministry of Natural Resources and Forestry, Wildlife Research and Monitoring Section, Trent University, DNA Building, Peterborough, Ontario, K9L 0G2, Canada

[5] Trent University, Forensic Science Program, Peterborough, Ontario, K9L 0G2, Canada

Corresponding author: jessewolf@trentu.ca

Keywords

Mountain goat, White-tailed deer, Genomics, GPS tracking, Habitat use, Core range

Abstract

The gut microbiome of animals varies by age, diet, and habitat, and directly influences individual health.

Similarly, variation in an individual's home range can lead to differences in feeding strategies and fitness.

Ungulates (hooved mammals) exhibit species-specific microbiomes and habitat use patterns: here, we

combined gut microbiome and movement data to assess relationships between space use and the gut

microbiome in a specialist and a generalist ungulate. We captured and GPS radiocollared 24 mountain

goats (*Oreamnos americanus*) and 34 white-tailed deer (*Odocoileus virginianus*). We collected fecal

samples and conducted high-throughput sequencing of the 16S rRNA gene. Using GPS data, we estimated

core (50%) and home range (95%) sizes and calculated proportional use for several important habitat

types. We generated metrics related to gut diversity and key bacterial ratios. We hypothesize that larger

Firmicutes to *Bacteroides* ratios confer body size or fat advantages that allow for larger home ranges, and

that relationships between gut diversity and disproportionate habitat use is stronger in mountain goats due

to their restricted niche relative to white-tailed deer. *Firmicutes* to *Bacteroides* ratios were positively

correlated with core range area in both species. Mountain goats exhibited a negative relationship between

gut diversity and use of two key habitat types (treed areas and escape terrain), whereas no relationships

were detected in white-tailed deer. This is the first study to relate core range size to the gut microbiome in

wild ungulates and is an important proof of concept that advances the information that can be gleaned

from non-invasive sampling.

Introduction

The gastrointestinal tract of animals contains trillions of microbes that influence individual health. Gut bacteria, hereafter termed the gut microbiome, can modify immune responses [1], improve and modulate metabolism [2], and affect behaviour [3,4]. While largely stable over time [5,6], disturbance of the gut microbiome can lead to disease [7] and impacts metabolic versatility, meaning the ability to survive equally well when presented with a wide range of dietary compositions and habitat [8,9]. Gut microbiome diversity has been shown to impact behaviour; for example, gut microbiome manipulation in mice resulted in improved memory as measured using a passive-avoidance test [10]. In another example, the presence of specific gut bacteria species suppressed protein appetite, indicating the ability of the gut microbiome to drive dietary decisions [11]. Differences in gut microbiome composition are correlated with landscape alteration [12,13], while levels of daily activity and foraging also appear to be influenced by the gut microbiome [14,15]. Notably, distinct diet types, such as herbivory and carnivory, are associated with unique microbiome profiles [16]. The mechanistic links are not totally understood, but are thought to follow the microbiota-gut-brain axis where bacteria have the ability to generate neurotransmitters that influence cognition [17].

Mammalian herbivores are typified by specific gut microbial taxa as they rely on these bacteria to extract energy and nutrients from food, synthesize vitamins, and detoxify plant defense compounds [18]. For example, *Firmicutes* and *Bacteroidetes* are involved in energy resorption and carbohydrate metabolism; *Firmicutes* can act as a more effective energy source, leading to more efficient calorie absorption and weight gain, while *Bacteroidetes* are involved with energy production and conversion, and amino acid transport and metabolism [19]. Ungulates, and ruminants in particular, have specialized anatomical and physiological adaptations to accommodate the cellulolytic fermentation of low-nutrition, high-fiber plant materials [20]. A specialized gut microbiome allows ruminants to digest typically indigestible plant

biomass [21] and as a result, exploit novel environments. Mountain goats (*Oreamnos americanus*) are

alpine ruminants that are endemic to the mountainous regions of northwestern North America [22].

Mountain goats use lower elevation, forested, and warmer aspect habitat during the winter and higher

elevation, mountainous terrain in summer [23–26]. They are considered intermediate browsers and eat a

variety of forage, with diets generally dominated by grasses [27,28]. In contrast, white-tailed deer

(*Odocoileus virginianus*) exploit a variety of habitat and food resources and cover a large geographic

range that stretches across most of North America and includes part of Central and South America [29].

White-tailed deer use woody cover habitats year-round, but can also thrive in urban and rural settings

[30,31]; they maintain distinct seasonal ranges in the northern parts of their range and are considered

browsing ruminants and both habitat and dietary generalists [32].

Our study integrated high-throughput sequencing and GPS telemetry to evaluate the relationship between

gut microbiome, home range area, and disproportionate habitat use of two ungulates living in contrasting

environments. This is among the first studies to link space use to the gut microbiome in wild systems and

provides a proof of concept that advances the information that can be generated from non-invasive

sampling. We quantified the relationship between key microbiome diversity metrics on home range size

and relative use of different habitat classes inferred from GPS tracking of individuals. From an

evolutionary perspective, this link between variation in phenotype or behaviour and the gut microbiome

assumes selection operates on both the phenotypes of the constituents (microbiome) and host, with the

collective genetic material known as the hologenome [33]. We hypothesized that an increase in gut

diversity would be linked with an increase in area used, as greater gut diversity would reflect, and possibly

drive larger use of space and a more resource-diverse home range, similar to the findings of [34]. As high

Firmicutes to *Bacteroides* ratios correspond to larger body size and fat stores [19,35], we hypothesized

that larger *Firmicutes* to *Bacteroides* ratios would be correlated with larger home ranges, as individuals

building up fat stores for winter would generally use more space to forage. This relationship may be

impacted by resource distribution, as specialists prefer homogenously distributed resources, while

generalists prefer heterogeneously distributed resources, which can impact space use [36]. Consequently,

we hypothesized that relationships between proportional habitat use and the gut microbiome would be

stronger in specialists as they have a more restricted niche with deviations from this having larger

consequences, whereas generalists can make use of a variety of habitat areas.

Methods

Animal captures, sample collection and DNA extraction

We captured and radio-collared male and female mountain goats on three adjacent mountain complexes

using aerial net-gun capture northeast of Smithers, British Columbia, Canada (Fig. 1). We captured and

radio-collared 34 female white-tailed deer using baited Clover traps southwest of Ottawa, Ontario, Canada

(Fig. 1). For more information on animal captures, see [37,38]. VERTEX Plus and VERTEX Lite Global

Positioning System (GPS) collars (VECTRONIC Aerospace, Germany) were used for mountain goats,

while store-on-board (G2110D, Advanced Telemetry Solutions, Isanti, MN) or GSM-upload (Wildcell

SG, Lotek Wireless Wildlife Monitoring, Newmarket, ON) GPS collars were used for white-tailed deer.

Collars recorded locations every four hours for mountain goats and every five hours for white-tailed deer.

During captures we took fresh fecal pellets from each individual and stored them at -20°C; all captures

took place during winter. Lab surfaces were sterilized with 90% EtOH and 10% bleach solution and a

small portion of a single fecal sample (~1/4 including exterior and interior portions) was digested

overnight at 56°C in 20 ul proteinase K and 180 ul Buffer ATL from the Qiagen DNeasy Blood & Tissue

Kit (Qiagen, Valencia, California, USA). DNA was extracted from the digest with the QIAamp

PowerFecal DNA Kit (Qiagen, Valencia, California, USA).

High-throughput sequencing and bioinformatics

The validated Illumina 16S rRNA Metagenomic Sequencing Library Preparation (#15044223 rev. B)

protocol was followed for library preparation using slight modifications [39]. The V3 and V4 regions of

the 16S ribosomal ribonucleic acid (16S rRNA) hypervariable region were targeted with four variants of

341F and 805R primers designed by [40]. A unique combination of Nextera XT indexes, index 1 (i7), and

index 2 (i5) adapters were assigned to each sample for multiplexing and pooling. Four replicates of each

sample of fecal DNA were amplified in 25 µl PCR using the 341F and 805R primers. The replicated

amplicons for each sample were combined into a single reaction of 100 µl and purified using a QIAquick

PCR Purification Kit (Qiagen, 28104) and quantified on the Qubit Fluorometer. Sample indexes were

annealed to the amplicons using an 8-cycle PCR reaction to produce fragments approximately 630 bp in

length that included ligated adaptors; the target amplicon is approximately 430 bp in length (Illumina 16S

rRNA Metagenomic Sequencing Library Preparation; #15044223 rev. B). Samples were purified with the

QIAquick PCR Purification Kit and the final purified library was validated on a TapeStation (Agilent,

G2991AA) and sequenced in 300 bp pair-end reads on an Illumina MiSeq sequencer at the Genomic

Facility at the University of Guelph (Guelph, Ontario).

The quality of the raw sequences was assessed using FastQC v 0.11.9 (https://github.com/s-

andrews/FastQC) and we determined the low-quality cut-off for forward and reverse reads. Forward and

reverse reads were imported into QIIME2 v 2019.4 [41] for quality control, sequence classification, and

diversity analysis. Merged, forward, and reverse reads were analyzed independently using the quality

control function within QIIME2 and DADA2 to perform denoising and to detect and remove chimeras.

QIIME2 follows the curated DADA2 R library workflow (https://benjjneb.github.io/dada2/) that requires

zero mismatches in overlapping reads for successful merging, since reads are denoised and errors are

removed before merging occurs. The taxonomy, to the species level, of all sample reads were assigned

using the Silva 132 reference taxonomy database (https://docs.qiime2.org/2019.4/data-resources/). We

calculated the relative proportion of *Firmicutes* to *Bacteroidetes* for each of the grouped data. Estimates of

diversity included Shannon's Index, observed richness, measured with Amplicon Sequence Variants

(ASVs), and Pielou's evenness, a measure of diversity that is the ratio of observed diversity to the

maximum possible in a sample having the same number of species [42]; these were screened for

correlation to one-another and read-depth. Pielou's evenness, observed richness (ASVs), and the ratio of

Firmicutes to *Bacteroidetes* were retained for subsequent analyses. We rarefied sample reads to the

sample with the least number of reads. A principle coordinates analysis (PCoA) was also conducted using

Bray Curtis dissimilarity matrices between species to determine if gut microbiome communities were

different between hosts.

GPS filtering, home range, and proportional habitat use analysis

We used different filtering approaches and seasonal delineations for each species due to the differences in

landscapes occupied by mountain goats and white-tailed deer. For the mountain goat data, any N.A. or

mortality signals were filtered out as were any GPS points outside of 600m - 2500m in elevation, as this

reflects the maximum and minimum for the study area. Dilution of precision (DOP) values over 10 were

plotted against elevation and landscape type, to ensure there were no patterns in distribution, and that

filtering would not bias downstream analyses. Movement rates between successive GPS points were also

calculated, and any movement rates beyond 15 km/hr were removed from analyses as these were deemed

spurious. Seasons were defined as follows: Summer - May 1st to October 31st and Winter - December 1st to

April 30th [23,43–45]. November was excluded from seasonal data as [26] noted a large increase in male

mountain goat home ranges due to the rut. In white-tailed deer filtering occurred as above, but with no

elevation restrictions as the topography of Marlborough Forest is effectively flat. White-tailed deer in

Marlborough Forest have distinct summer and winter ranges; as considerable variation in migration dates

were observed, movements to and from Marlborough forest were relative to each individual deer's

movement and as such, there was no hard date range. A migration movement was defined as when a deer

moved between non-overlapping seasonal ranges and then occupied one seasonal range until the following

migration movement [38]. No seasonal GPS data were excluded for white-tailed deer, as changes in

movement patterns and home range size during the rut are minimal in females [46].

We used Brownian Bridge Movement Models (BBMMs) to generate individual home ranges using the

BBMM package in R [47]. A BBMM is a continuous-time stochastic movement model that uses

probabilistic and maximum likelihood approaches where observed locations are measured with error to

model home ranges [48]. A minimum of 275 GPS points was required to generate a BBMM and

individual home ranges were calculated with a maximum lag time between successive locations of two

times the expected fix rate. A location error of 20 m was used as per [49], with a cell size of 25 m^2. We

generated 50% and 95% isopleths representing the core and home ranges respectively. Isopleths were

generated separately for each individual during both summer and winter, as ungulates exhibit sex-specific

habitat use patterns, that vary both seasonally and by age-class [22,26,44,50,51]. We focused on summer

isopleths for both species in downstream analyses, as deer were baited in winter, which has been shown to

bias movement and shift core ranges [52]. We report only the 50% isopleths, hereafter termed core ranges,

to maximize seasonal differences, as they were highly correlated to the 95% isopleths (t_{46}=9.3, r=0.81,

p<0.0001) and results were similar between summer 50% and 95% isopleths (Fig. S1). We generated

BBMM isopleths using the R package rgdal [53] and all analyses were conducted in R v.3.6.1. We

imported shapefiles into ArcGIS Pro 2.5.0 to calculate home and core range areas in km^2 for further

analyses.

Proportional use of habitats was assessed by calculating the number of GPS points in a given habitat type

within an isopleth, relative to the total number of GPS points located in that isopleth. Similar to Johnson's

third-order habitat selection, proportional habitat use in this study refers to how specific habitat types were

used within a core range [54]. Proportion values in the 50% isopleths were highly correlated to the 95%

isopleths (t_{117}=106, r=0.99, p <2.2e-16, Table S1, S2). We selected ecologically relevant features that

showed previous evidence for use in both species. Features used in mountain goat models were treed

habitat (landscape classified as sub-boreal spruce, interior cedar, or Engelmann spruce), Heat Load Index

(HLI; a combination of latitude, aspect, and slope), and escape terrain (landscape where slope is $\geq 40º$;

[55]). For HLI [56], the average value of HLI for all GPS points within an isopleth for a given individual

was calculated. These features have shown evidence for disproportionate usage/selection in previous

research on mountain goats [55]. We used the Biogeoclimactic Ecosystem Classification (BEC) dataset

from GeoBC to determine terrestrial landcover type for mountain goats and the Digital Elevation Model

from GeoBC to calculate escape terrain. In the white-tailed deer models, we used forested habitat (tree

cover >60%), treed swamp (tree cover >25%), and thicket swamp (tree cover <25%, open and shrub

communities), as each of the three habitat features exhibited >20% core landcover composition in [38],

and thus, were available for the majority of individuals to use [57]. The Southern Ontario Land Resource

Information System (SOLRIS) data set version 2.0 [58] was used to determine land cover types for white-

tailed deer.

Generalized linear models

We analyzed the associations between core range size, gut microbiome metrics, and age class for both

species individually using Generalized Linear Models (GLMs) with the Gaussian family distribution and

identity link function. The core range GLMs consisted of core range size as a response variable, a single

microbiome metric (*Firmicutes* to *Bacteroidetes* ratio, number of ASVs, or Pielou's evenness) and age

class (adult or subadult) as fixed explanatory effects. The proportional habitat use GLMs consisted of

proportion of habitat used as a response variable, a single microbiome metric (*Firmicutes* to *Bacteroidetes*

ratio, ASV, or Pielou's evenness) and age class, as fixed explanatory effects. One exception to this was the

HLI GLM, as the response variable was the mean HLI value for GPS points in the isopleth, while the

explanatory variables were the same as described above. Individuals 0 - 2 years of age were considered

subadults for white-tailed deer, while individuals 0 - 3 years of age were considered subadults for

mountain goats [22,59]. Sex was included only in mountain goat models as we only had data from female

deer. Effect size and confidence intervals are reported for each model. We conducted five-fold linear

model cross validation using the Caret package in R [60] to test for overfitting of our models and quantify

the model's predictive ability. We reported the Scatter Index (SI) and Root Mean Square Error (RMSE):

low values in RMSE and SI are indicative of a good model fit and low residual variance.

Results

Bioinformatic filtering and taxonomic analysis

Twenty-three mountain goat and twenty-five white-tailed deer fecal sample sequences passed QC and a

total of ~8.17 million paired-end reads (n_{MG}=5,488,856, n_{WTD}=2,679,668) were generated (SRA accession

number PRJNA638162). FastQC analysis indicated that both forward and reverse reads lost quality > 259

bp in length (Phred score <25), so all reads were trimmed to a length of 259 bp. Following DADA2 strict

quality filtering, ~1.16 million paired-end reads (n_{MG}=709,457, n_{WTD}=457,541) were retained for

taxonomic and diversity analyses. Losing this many reads to quality filtering is typical (see [39]), as

permitted error rates are extremely low in DADA2, resulting in high certainty among retained reads [61].

White-tailed deer had higher averages of both Pielou's evenness (mean 0.95, min 0.92, max 0.96, SD

0.012) and *Firmicutes* to *Bacteroidetes* ratios (mean 8.3, min 1.49, max 21.5, SD 6.22) than mountain

goats (Pielou's evenness mean 0.92, min 0.84, max 0.95, SD 0.028; *Firmicutes* to *Bacteroidetes* ratio

mean 6.90, min 3.51, max 12.10, SD 2.43). Conversely, mountain goats had higher averages of ASVs

(mean 31,393.6, min 6638, max 59,481, SD 14,187.1) relative to white-tailed deer (mean 17,828.4, min

4847, max 39,146, SD 7212.12). Age class and sex averages, in addition to winter data, are shown in

Table S3. A Bray-Curtis PCoA clearly separated mountain goats and white-tailed deer (Fig. 2).

Core range and proportional habitat use

Data filtering resulted in 84,932 GPS points for mountain goats (mean per individual 3,679, range 277 -

6,704, SD 761), and 63,900 GPS points for white-tailed deer (mean per individual 2,556, range 831 -

3,558, SD 906). One individual mountain goat was removed due to a small number of GPS points (n =

16). Mountain goats exhibited slightly larger summer core range size of 0.40 km^2 (min 0.01 km^2, max 0.72

km^2, SD 0.17), compared to 0.36 km^2 (min 0.13 km^2, max 0.81 km^2, SD 0.16) for white-tailed deer. The

mean proportional habitat use values are reported in Table S4.

Generalized linear models

We compared 50% isopleths from summer and winter; summer core ranges were reported here, as winter

core ranges produced qualitatively similar results, albeit with weaker signals (Table S5). In both species, a

greater *Firmicutes* to *Bacteroidetes* ratio was associated with larger core ranges with both models

explaining an equal amount of variance (Nagelkerkes's R^2~0.28; Table 1; Fig. 3).

Pielou's evenness was directly correlated with core range size in mountain goats; conversely, Pielou's

evenness was inversely correlated with core range size in white-tailed deer. Here the mountain goat model

explained a relatively large portion of the variance (R^2=0.47; Table 1; Fig. 3). Observed richness (ASVs)

was associated with larger core ranges in mountain goats (Nagelkerkes's R^2~0.45, Table 1; Fig. 3) and

white-tailed deer (Nagelkerkes's R^2~0.08, Table 1; Fig. 3). Age-class was a significant predictor in all

three of the mountain goat core range GLMs, but neither of the white-tailed deer core range models while

sex was a non-significant predictor in all mountain goat core range models (Table 1).

The use of escape terrain and treed areas was moderately correlated in mountain goats (t_{20}=2.94, p<0.01,

r=0.55), and exhibited significant relationships relative to Pielou's evenness and number of ASVs; effect

size confidence intervals did not overlap zero in models that measured the relationship between use of

escape terrain and treed areas and Pielou's evenness, and ASVs for the latter (Table 2). Specifically, a

larger Pielou's evenness value and number of ASVs was seen in individuals using less treed habitat and

less escape terrain (Fig. 4). In HLI GLMs, confidence intervals overlapped zero and exhibited a slight

decrease in R^2 value relative to other mountain goat habitat use models (Table 2). All GLM parameter

estimates in the white-tailed deer models had confidence intervals overlapping zero. Cross validation of

linear models supported retaining age class and microbiome metrics as predictor variables of core range

size. RMSE values in models with core range size as the response variable ranged from 0.12 to 0.18, and

SI values ranged from 0.30 to 0.51, whereas in proportional habitat use models, RMSE values ranged

from 0.11 to 0.33 and SI values from 0.033 to 1.14 (Table S6); this suggests moderate to high support for

the models.

Discussion

The link between the gut microbiome and host space use has implications for foraging, activity levels, and

ability to use energetically costly habitats. We show that certain aspects of the gut microbiome in both a

generalist ungulate and specialist ungulate are linked to patterns of habitat use and home range size.

Although the patterns are nuanced, there were some commonalities that collectively suggest the gut

microbiota plays a role in determining space use patterns. An increase in *Firmicutes* to *Bacteroidetes*

ratios in both species was correlated with an increase in core range sizes (Fig. 3). Increased *Firmicutes* to

Bacteroidetes ratios are linked to increased Body Mass Index (BMI) and obesity in humans [19]. Both

Firmicutes and *Bacteroidetes* are involved in energy resorption and are important for metabolism [62,63].

In a comparable study, home range size was not correlated with the *Firmicutes* to *Bacteroidetes* ratio in

wild rodents [64], however, seasonal weight changes are more dynamic in small mammals [65,66], and we

suspect this pattern might be more ubiquitous across temperate ungulates given their specialized gut

microbiomes and need to put on fat stores. Ultimately, a higher *Firmicutes* to *Bacteroidetes* ratio may aid

in the accumulation of fat stores to survive the winter.

In large ungulates, increasing levels of body fat is important to survive the winter when forage is limited,

and may be impacted by the relative ratios of key bacterial species. In muskoxen (*Ovibos moschatus*), the

abundance of *Firmicutes* stayed similar across seasons, while *Bacteroidetes* increased in the summer

months, meaning the ratio of *Firmicutes* to *Bacteroidetes* is lower in the summer [67]. The ability to

conserve bacteria necessary for adding fat, namely high levels of *Firmicutes,* while exhibiting changes in

gut composition suggests ungulates can prepare for the winter even though the gut microbiome is shifting

(see [39]). This concept is of importance to white-tailed deer, as diet turnover is pronounced in Ontario

[68]. Shifting from herbaceous vegetation in the spring to woody browse in the winter may similarly result

in increased diversity, while simultaneously conserving bacteria necessary for adding fat.

Specialists and generalists shift habitats seasonally, which can impact diet, and may be influenced by the

gut microbiome; mountain goats move from alpine summer habitat to subalpine winter areas and white-

tailed deer exhibit distinct winter and summer ranges [23,30,50]. There was considerably more variation

in proportion of habitat used in the generalist white-tailed deer (Table S2), likely a function of their ability

to use multiple habitat types and diet sources. Here, proportion habitat use-microbiome models had no

relationship, as all proportion habitat use coefficients in white-tailed deer had confidence intervals

overlapping zero (Table 2). Interestingly, the two habitat variables in mountain goats with clear signals

were escape terrain and treed areas (Fig. 4), which are defining habitat characteristics of this species: the

use of alpine terrain (i.e. no trees) and steep slopes [22,69]. Deviations from their specialized and

preferred habitat types comes at a cost, as ungulates typically exhibit trade-offs with respect to forage

quality and predation risk; avoiding predation can lead to decreased forage quality and abundance [70,71].

In mountain goats and bighorn sheep (*Ovis canadensis*), energy expenditures increased when travelling

uphill or downhill - termed a vertical cost [72]. A lower *Firmicutes* to *Bacteroidetes* ratio could be linked

to the vertical cost associated with spending more time in escape terrain, meaning less-fat or prime-

conditioned individuals spend more time in escape terrain. Conversely, using more forage-available treed

habitat may be correlated with an increase in *Firmicutes* to *Bacteroidetes* ratio as individuals have

increased access to forage, with minimal vertical cost. Collectively, the habitat models had stronger

signals in the specialist species compared to the generalist, and we suggest this is reflective of their more

restricted habitat niche relative to generalists, ultimately supporting our predictions, where microbial

deviations may be able to generate more prominent shifts in behaviour.

Deviations from the usage of important habitat types have larger consequences in specialists as they have

a more restricted niche, and thus, the potential for the gut microbiome to modulate habitat use patterns is

important in specialists to increase their competitive ability. While there is potential for modulation of

proportion of specific habitat usage, the impact of the gut microbiome on the amount of space used may

be more explicit. A lower Pielou's evenness value indicates that a given individual has decreased diversity

relative to another individual with the same number of gut microbial species, while a lower observed

richness value represents a decrease in absolute number of gut microbiome species. The relationship

between Pielou's evenness and core range size differed in direction between species; greater gut evenness

in mountain goats was correlated with a larger core range, while a negative relationship was noted in

white-tailed deer (Fig. 3). Greater gut microbiome richness has been linked to increased gut stability in

humans when faced with a shift in diet [73], as well as with changes in reproductive status in birds [74]. A

positive relationship was noted between gut richness and core size in both species but was much stronger

in mountain goats (Table 1; Fig. 3). Increased gut evenness might promote individuals moving through

and foraging in larger areas, resulting in a positive relationship between evenness and core range size,

while richness may act similarly. This is of added importance in a specialist, as the ability to utilize and

forage in a more diverse array of habitat types might allow individuals to access and subsist in areas that

other conspecifics cannot, thus increasing their competitive ability [75–77]. Higher relative gut

microbiome diversity might allow specialists to forage more like generalists, while lower diversity may

limit specialists to their typical niches. The reverse trend in white-tailed deer gut evenness might reflect

the difference in feeding and habitat-use strategies; white-tailed deer are generalists, and thus by

definition, can effectively use a variety of habitat and food sources. The negative relationship observed

between gut evenness and range size in white-tailed deer might suggest that deer with higher gut

microbiome evenness are able to use more diet sources overall and thus can meet their nutritional

requirements within a smaller area relative to individuals with less diverse gut microbiomes. As

generalists, deer are not as constrained by specific food and habitat types, and thus, their space use may be

less heavily influenced by gut microbiome diversity. It is also possible that winter baiting had an impact

on the gut microbiome in white-tailed deer, which could have led to weaker relationships.

We built our models assuming the microbiome influences habitat and home range patterns; but this need

not be cause and effect, and the relationship is likely reciprocal [78]. For example, diet-microbiome

covariance has been observed in multiple large herbivores, where seasonal diet turnover and seasonal

microbiome turnover are positively correlated [79]. In this example, our model would assume the gut

microbiome impacts diet choice (e.g. [11]) and in turn, the correlation between diet turnover and the

microbiome contributes to habitat use of a given species. Assessing this relationship clearly needs

experimental testing, and we view our study as a proof-of-concept that provides a testable hypothesis.

Still, we demonstrated that quantifying the gut microbiome yields information related to space use; linking

these two highly complex components of biology aids in our understanding of selection on the

hologenome through the interplay between the individual and its microbial genomes and potential traits

under selection (e.g. core range size and proportional habitat use). These findings also demonstrate that

pellet sampling is useful in determining space and habitat use in managed populations, as it is conceivable

with a large enough database and validation, one could predict the distribution and behaviour of animals

on the landscape from non-invasively sampled pellets. Similar analyses of this kind should clarify the

extent to which space use is linked to the gut microbiome in other species. Ultimately, we utilized a

specialist and generalist ungulate to explore the link between the gut microbiome and movement to

generate quantitative findings surrounding the impact of the gut microbiome on space use in wild

populations.

Acknowledgements

Acknowledgements

Special thanks to Jeff Bowman, Erin Koen, Kathleen Lo and Kiana Young for their comments on earlier

drafts of this manuscript, Spencer Anderson for lab work, and Florent Déry for allowing us to use their

digital drawings.

The authors declare that they have no conflict of interest.

Funding

This work was supported by a Natural Sciences and Engineering Research Council Discovery Grant,

Canada Foundation for Innovation - John R. Evans Leaders Fund, and Compute Canada awards to ABAS,

Habitat Conservation Trust Foundation Enhancement and Restoration Grant to KK (Project # 6-252), the

Forest Enhancement Society of British Columbia, the British Columbia Mountain Goat Society, and the

Rocky Mountain Goat Alliance grant to JFW.

References

1. Arnolds KL, Lozupone CA. 2016 Striking a balance with help from our little friends – how the gut microbiota contributes to immune homeostasis. *Yale J. Biol. Med.* **89**, 389–95. (doi: 10.1371/journal.pone.0176573)

2. De Angelis M, Ferrocino I, Calabrese FM, De Filippis F, Cavallo N, Siragusa S, et al. 2020 Diet influences the functions of the human intestinal microbiome. *Sci. Rep.* **10**, 4247. (doi:10.1038/s41598-020-61192-y)

3. Zhang N, Ju Z, Zuo T. 2018 Time for food: The impact of diet on gut microbiota and human health. *Nutrition* **51**–**52**, 80–85. (doi:10.1016/j.nut.2017.12.005)

4. Shreiner A, Kao J, Young V. 2015 The gut microbiome in health and in disease. *Curr. Opin. Gastroenterol.* **31**, 69–75. (doi: 10.1097/MOG.0000000000000139)

5. Faith JJ, Guruge JL, Charbonneau M, Subramanian S, Seedorf H, Goodman AL, et al. 2013 The long-term stability of the human gut microbiota. *Science* **5**, 1237439. (doi: 10.1126/science.1237439)

6. Coyte KZ, Schluter J, Foster K. 2015 The ecology of the microbiome: Networks, competition, and stability. *Science,* **350**, 663–666. (doi: 10.1126/science.aad2602)

7. Williams CL, Caraballo-Rodríguez AM, Allaband C, Zarrinpar A, Knight R, Gauglitz JM. 2019 Wildlife-microbiome interactions and disease: exploring opportunities for disease mitigation across ecological scales. *Drug Discov. Today Dis. Model.* **28**, 105–15. (doi:10.1016/j.ddmod.2019.08.012)

8. Tinker KA, Ottesen EA. 2016 The core gut microbiome of the American cockroach, *Periplaneta americana*, is stable and resilient to dietary shifts. *Appl. Environ. Microbiol.* **82**, 6603–6610. (doi: 10.1128/AEM.01837-16)

9. Esposti MD, Romero EM. 2017 The functional microbiome of arthropods. *PLoS One* **12**, e0176573. (doi: 10.1371/journal.pone.0176573)

10. Mao JH, Kim YM, Zhou YX, Hu D, Zhong C, Chang H, et al. 2020 Genetic and metabolic links between the murine microbiome and memory. *Microbiome* **8**, 1–14. (doi: 10.1186/s40168-020-00817-w)

11. Leitão-Gonçalves R, Carvalho-Santos Z, Francisco AP, Fioreze GT, Anjos M, Baltazar C, et al. 2017 Commensal bacteria and essential amino acids control food choice behavior and reproduction. *PLoS Biol.* **15**, 1–29. (doi: 10.1371/journal.pbio.2000862)

12. Murray, M.H., Lankau EW, Kidd AD, Welch CN, Ellison T, H.C A, et al. 2020 Gut microbiome shifts with urbanization and potentially facilitates a zoonotic pathogen in a wading bird. *PLoS One* **15**, e0220926. (doi:10.1371/journal.pone.0220926)

13. San Juan P, Hendershot J, Daily G, Fukami T. 2019 Land-use change has host-specific influences on avian gut microbiomes. *ISME J.* **14**, 318–321. (doi:10.1038/s41396-019-0535-4)

14. Schretter CE, Vielmetter J, Bartos I, Marka Z, Marka S, Argade S, et al. 2018 A gut microbial factor modulates locomotor behaviour in Drosophila. *Nature* **563**, 402–406. (doi:10.1038/s41586-018-0634-9)

15. Jones JC, Fruciano C, Marchant J, Hildebrand F, Forslund S, Bork P, et al. 2018 The gut microbiome is associated with behavioural task in honey bees. *Insectes Soc.* **65**, 419–29. (doi:10.1007/s00040-018-0624-9)

16. Ley RE, Hamady M, Lozupone C, Turnbaugh PJ, Ramey RR, Bircher JS, et al. 2008 Evolution of mammals and their gut microbes. *Science* **320**, 1647–1651. (doi: 10.1126/science.1155725)

17. Cryan JF, Dinan TG. 2012 Mind-altering microorganisms: The impact of the gut microbiota on brain and behaviour. *Nat. Rev. Neurosci.* **13**, 701–712. (doi:10.1038/nrn3346)

18. Dearing MD, Kohl KD. 2017 Beyond fermentation: Other important services provided to

endothermic herbivores by their gut microbiota. *Integr. Comp. Biol.* **57**, 723–731. (doi:
10.1093/icb/icx020)

19. Koliada A, Syzenko G, Moseiko V, Budovska L, Puchkov K, Perederiy V, et al. 2017 Association
between body mass index and Firmicutes/Bacteroidetes ratio in an adult Ukrainian population.
BMC Microbiol. **17**, 4–9. (doi: 10.1186/s12866-017-1027-1)

20. De Tarso S, Oliviera D, Bastos Alfonso J. 2016 Ruminants as part of the global food system: How
evolutionary adaptations and diversity of the digestive system brought them to the future. *J. Dairy,*
Vet. Anim. Res. **3**, 1–7. (doi: 10.15406/jdvar.2016.03.00094)

21. Kruger Ben Shabat S, Sasson G, Doron-Faigenboim A, Durman T, Yaacoby S, Berg Miller ME, et
al. 2016 Specific microbiome-dependent mechanisms underlie the energy harvest efficiency of
ruminants. *ISME J.* **10**, 2958–2972. (doi: 10.1038/ismej.2016.62)

22. Festa-Bianchet M, Côté SD. 2008. Mountain Goats: Ecology, Behavior, and Conservation of an
Alpine Ungulate. Island Press, Washington, DC.

23. Poole KG, Heard DC. 2003 Seasonal habitat use and movements of mountain goats, *Oreamnos*
americanus, in east-central British Columbia. *Can. Field-Naturalist* **117**, 565–576.

24. Poole KG, Stuart-Smith K, Teske IE. 2009 Wintering strategies by mountain goats in interior
mountains. *Can. J. Zool.* **87**, 273–283. (doi: 10.1139/z09-009)

25. Taylor S, Wall W, Kulis Y. 2014 Habitat selection by mountain goats in south coastal British
Columbia Habitat Selection by Mountain Goats in South Coastal British Columbia. *Bienn. Symp.*
North. Wild Sheep Goat Counc. **15**, 141–158.

26. White KS. 2006 Seasonal and Sex-specific Variation in Terrain Use and Movement Patterns of
Mountain Goats in Southeastern Alaska. *Bienn. Symp. North. Wild Sheep Goat Counc.* **15**, 183–
194.

27. Festa-bianchet M, Côté SD. 2003. Mountain Goat (*Oreamnos americanus*). In Wild Mammals of
North America: Biology, Management, Conservation. John Hopkins Press, Baltimore, MD.

28. Hofmann RR. 1989 Evolutionary steps of ecophysiological adaptation and diversification of
ruminants: a comparative view of their digestive system. *Oecologia* **78**, 443–457.
(doi:10.1007/BF00378733)

29. Hewitt DG. 2011. Biology and management of white-tailed deer. CRC Press, Boca Raton, FL.

30. Grund MD, McAninch JB, Wiggers E. 2002 Seasonal movements and habitat use of female White-
tailed deer associated with an urban park. *J. Wildl. Manage.* **66**, 123–130. (doi: 10.2307/3802878)

31. Van Deelen T, Campa III H, Hamady M, Haulfer J. 1998 Migration and seasonal range dynamics
of deer using adjacent deeryards in northern Michigan. *J. Wildl. Manage.* **62**, 205–213.

32. Berry SL, Shipley LA, Long RA, Loggers C. 2019 Differences in dietary niche and foraging
behavior of sympatric mule and white-tailed deer. *Ecosphere* **10**, e02815. (doi: 10.1002/ecs2.2815)

33. Bordenstein SR, Theis KR. 2015 Host biology in light of the microbiome: ten principles of
holobionts and hologenomes. *PLoS Biol.* **13**, e1002226. (doi: 10.1371/journal.pbio.1002226)

34. Ma L, Xu S, Liu H, Xu T, Hu L, Zhao N, et al. 2019 Yak rumen microbial diversity at different
forage growth stages of an alpine meadow on the Qinghai-Tibet Plateau. *PeerJ* **7,** e7645. (doi:
10.7717/peerj.7645)

35. Ley RE. 2010 Obesity and the human microbiome. *Curr. Opin. Gastroenterol.* **26**, 5–11. (doi:
10.1097/MOG.0b013e328333d751)

36. Marrotte RR, Bowman J, Morin SJ. 2020 Spatial segregation and habitat partitioning of bobcat and
Canada lynx. *Facets* **5,** 503–522. (doi: 10.1139/facets-2019-0019)

37. Wolf JF, Kriss KD, Macaulay KM, Shafer ABA. 2020 Panmictic population genetic structure of
northern British Columbia mountain goats (*Oreamnos americanus*) has implications for harvest
management. *Conserv. Genet.* **21**, 613–23. (doi:10.1007/s10592-020-01274-6)

38. Munro K. 2020. White-tailed fear: The human-created landscape of fear and its effect on white-
tailed deer (*Odocoileus virginianus*) behaviour. [PhD dissertation].

39.   Haworth SE, White KS, Côté SD, Shafer BA. 2019 Space, time and captivity: Quantifying the
factors influencing the fecal microbiome of an alpine ungulate. *FEMS Microbiol. Ecol.* **95**, 1–12.
(doi: 10.1093/femsec/fiz095)
40.   Herlemann DPR, Labrenz M, Ju K, Bertilsson S, Waniek JJ, Andersson AF. 2011 Transitions in
bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *Int. Soc. Microb. Ecol.*
5, 1571–1579. (doi: 10.1038/ismej.2011.41)
41.   Boylen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. 2019
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat.*
Biotechnol. **37**, 852-857. (doi: 10.1186/s13059-018-1450-0)
42.   Pielou EC. 1966 The measurement of diversity in different types of biological collections. *J.*
Theor. Biol. **13**, 131–44.
43.   Cadsand B. 2012 Responses of mountain goats to heliskiing activity: movements and resource
selection. [MSc dissertation].
44.   Mountain Goat Management Team. 2010 Conservation Status Report. BC Conservation Data
Centre.
45.   Richard JH, Côté SD. 2016 Space use analyses suggest avoidance of a ski area by mountain goats.
J. Wildl. Manage. **80**, 387–395. (doi: 10.1002/jwmg.1028)
46.   Hölzenbein S, Schwede G. 1989 Activity and movements of female white-tailed deer during the
rut. *J. Wildl. Manage.* **53**, 219–223. (doi: 10.2307/3801337)
47.   Nielson, R. M., Sawyer, H., & McDonald, T. L. (2015). BBMM: Brownian Bridge Movement
Model. R package version 3.0. [Online]. Retrieved from: https://cran.r-
project.org/web/packages/BBMM
48.   Horne JS, Garton EO, Krone SM, Lewis JS. 2007 Analyzing animal movements using Brownian
bridges. *Ecology* **88**, 2354–2363. (doi: 10.1890/06-0957.1)
49.   Sawyer H, Kauffman MJ, Nielson RM, Horne JS. 2009 Identifying and prioritizing ungulate
migration routes for landscape-level conservation. *Ecol. Appl.* **19**, 2016–2025. (doi: 10.1890/08-
2034.1)
50.   Lesage L, Crête M, Huot J, Dumont A, Ouellet JP. 2000 Seasonal home range size and philopatry
in two northern white-tailed deer populations. *Can. J. Zool.* **78**, 1930–1940. (doi: 10.1139/z00-117)
51.   Webb SL, Hewitt DG, Hellickson MW. 2007 Scale of management for mature male white-tailed
deer as influenced by home range and movements. *J. Wildl. Manage.* **71**, 1507–1512. (doi:
10.2193/2006-300)
52.   Kilpatrick HJ, Stober WA. 2002 Effects of temporary bait sites on movements of suburban white-
tailed deer. *Wildl. Soc. Bull.* **30**, 760–766.
53.   Bivand, R., Keitt, T., Pebesma, E., Sumner, M., Hijmans, R., Rouault, E., … Rundel, C. (2019).
Rgdal: Bindings for the 'geospatial' data abstraction library. R package version 1.4-8. [Online]
Retrieved from: https://cran.r-project.org/web/packages/rgdal/index.html
54.   Johnson DH. 1980 The comparison of usage and availability measurements for evaluating resource
preference. *Ecology* **61**, 65–71.
55.   Shafer ABA, Northrup JM, White KS, Boyce MS, Côté SD, Coltman DW. 2012 Habitat selection
predicts genetic relatedness in an alpine ungulate. *Ecology* **93**, 1317–1329. (doi: 10.1890/11-
0815.1)
56.   Mccune B, Keon D. 2002 Equations for potential annual direct incident radiation and heat Lload. *J.*
Veg. Sci. **13**, 603–606. (doi: 10.1111/j.1654-1103.2002.tb02087.x)
57.   Massé A, Côté SD. 2013 Spatiotemporal variations in resources affect activity and movement.
Can. J. Zool. **91**, 252–263. (doi: 10.1139/cjz-2012-0297)
58.   OMNRF. 2015 Southern Ontario Land Resource Information System (SOLRIS) Version 2.0: Data
Specifications.
59.   Delgiudice GD, Fieberg J, Riggs MR, Powell MC, Pan W. 2006 A long-term age-specific survival

analysis of female white-tailed deer. *J. Wildl. Manage.* **70**, 1556–1568. (doi: 10.2193/0022-
541x(2006)70[1556:alasao]2.0.co;2)

60.   Kuhn, M., Wing, J., Weston, S., Williams, A., Keefer, C., Engelhardt, A., … Hunt, T. (2020).
Caret: Classification and regression training. R package version 6.0-86. [Online]. Retrieved from
https://cran.r-project.org/web/packages/caret/index.html

61.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016 DADA2: High-
resolution sample inference from Illumina amplicon data. *Nat. Methods* **13**, 581–583. (doi:
https://doi.org/10.1038/nmeth.3869)

62.   Krajmalnik-Brown R, Ilhan ZE, Kang DW, DiBaise JK. 2012 Effects of gut microbes on nutrient
absorption and energy regulation. *Nutr. Clin. Pract.* **27**, 201–214. (doi:
10.1177/0884533611436116)

63.   Ottman N, Smidt H, de Vos WM, Belzer C. 2012 The function of our microbiota: who is out there
and what do they do? *Front. Cell. Infect. Microbiol.* **2**, 1–11. (doi: 10.3389/fcimb.2012.00104)

64.   Jameson J, Réale D, Kembel S. 2020 Gut microbiome modulates behaviour and life history in two
wild rodents. *bioRxiv* 1–26. (doi: 10.1101/2020.02.09.940981)

65.   Lynch G. 1973 Seasonal changes in thermogenesis, organ weights, and dody composition in the
white-footed mouse, *Peromyscus leucopus*. *Oecologia* **13**, 363–376.

66.   Merritt J, Zegers D. 1991 Seasonal thermogenesis and body-mass dynamics of Clethrionomys
gapperi. *Can. J. Zool.* **69**, 2771–7.

67.   Bird S, Prewer E, Kutz S, Leclerc LM, Vilaça ST, Kyle CJ. 2019 Geography, seasonality, and
host-associated population structure influence the fecal microbiome of a genetically depauparate
Arctic mammal. *Ecol. Evol.* **9**, 13202–13217. (doi: 10.1002/ece3.5768)

68.   Brown DT, Doucet GJ. 1991 Temporal changes in winter diet selection by white-tailed deer in a
northern deer yard. *J. Wildl. Manage.* **55**, 361–376.

69.   Gross J, MC K, Reed D, Reich R. 2002 GIS-based habitat models for mountain goats. *J. Mammal.*
83, 218–28. (doi: 10.1644/1545-1542(2002)083<0218:gbhmfm>2.0.co;2)

70.   Hebblewhite M, Merrill EH. 2009 Trade-offs between predation risk and forage differ between
migrant strategies in a migratory ungulate. *Ecology* **90**, 3445–3454. (doi: 10.1890/08-2090.1)

71.   Hamel S, Côté SD. 2007 Habitat use patterns in relation to escape terrain: Are alpine ungulate
females trading off better foraging sites for safety? *Can. J. Zool.* **85**, 933–943. (doi: 10.1139/Z07-
080)

72.   Dailey T V., Hobbs NT. 1989 Travel in alpine terrain: energy expenditures for locomotion by
mountain goats and bighorn sheep. *Can. J. Zool.* **67**, 2368–2375. (doi: 10.1139/z89-335)

73.   Tap J, Furet JP, Bensaada M, Philippe C, Roth H, Rabot S, et al. 2015 Gut microbiota richness
promotes its stability upon increased dietary fibre intake in healthy adults. *Environ. Microbiol.* **17**,
4954–4964. (doi: 10.1111/1462-2920.13006)

74.   Devaynes A, Antunes A, Bedford A, Ashton P. 2018 Bacterial species richness at three stages of
the breeding season in *Cyanistes caeruleus* (blue tit). *Acta Oecologica* **92**, 123–30.
(doi:10.1016/j.actao.2018.09.001)

75.   McPeek M. 1996 Trade-offs, food web structure, and the coexistence of habitat specialists and
generalists. *Am. Nat.* **148**, S124–38.

76.   Blake JG, Karr JR. 1987 Breeding birds of isolated woodlots: area and habitat relationships. *Ecol.*
Soc. Am. **68**, 1724–1734. (doi: 10.2307/1939864)

77.   Rodewald AD, Vitz AC. 2005 Edge and area-sensitivity of shrubland birds. *J. Wildl. Manage.* **69**,
681–688.

78.   Cryan JF, O'Riordan KJ, Cowan CSM, Sandhu K V., Bastiaanssen TFS, Boehme M, et al. 2019
The microbiota-gut-brain axis. *Physiol. Rev.* **99**, 1877–2013. (doi: 10.1152/physrev.00018.2018)

79.   Kartzinel TR, Hsing JC, Musili PM, Brown BRP, Pringle RM. 2019 Covariation of diet and gut
microbiome in African megafauna. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 23588–23593. (doi:

10.1073/pnas.1905666116)
565

Data Accessibility
The sequence data generated for the mountain goat and white-tailed deer gut microbiota has been
deposited in the Sequence Read Archive (BioProject ID: PRJNA638162; SRA submission: SUB7567698).
R scripts used to perform analyses and generate figures are available at
https://gitlab.com/WiDGeT_TrentU/graduate_theses/-/tree/master/Wolf_et_al.

Author Contributions
J.F.W. and A.B.A.S. conceived the ideas; K.D.K., K.M.M., K.M., B.R.P., J.F.W., and A.B.A.S. collected
materials and data; J.F.W. analysed the data; and J.F.W. wrote the paper with contributions from all co-
authors.

576 **Figures**

578
579 **Fig. 1** An example of habitat of both mountain goat (*Oreamnos americanus*) and white-tailed deer
580 (*Odocoileus virginianus*) along with inset sample maps. Colors are reflective of relative elevation and
581 topography. Average elevation of mountain goat habitat was 1614.7m, relative to 113.3m for white-tailed
582 deer.

583
584 **Fig. 2** Observed richness (ASVs) for mountain goats (*Oreamnos americanus*, n=22) and white-tailed deer
585 (*Odocoileus virginianus*, n=25). Mean observed richness was significantly different as computed by a
586 two-sample t-test (t=4.11, df=29.64, p<0.001).

587
588 **Fig. 3** Microbiome metrics for mountain goats (*Oreamnos americanus*, n=22) northeast of Smithers,
589 British Columbia, and white-tailed deer (*Odocoileus virginianus*, n=25) southwest of Ottawa, Ontario,
590 relative to core summer range size. Females are represented by circles and only mountain goats have
591 mixed-sex data. Adults (≥3 years of age) are represented in yellow, and subadults in blue. Significant
592 Generalized Linear Models (GLMs) are denoted with a black line.

593
Fig. 4 Microbiome metrics for mountain goats (*Oreamnos americanus*, n=21*)* northeast of Smithers,
British Columbia, relative to habitat selection coefficients of both Heat Load Index (HLI) and treed
habitat. Females are represented by circles and only mountain goats have mixed-sex data. Adults (≥3
years of age) are represented in yellow, and subadults in blue. Significant Generalized Linear Models
(GLMs) are denoted with a black line.

Tables
600
601 Table 1. Generalized linear models for core summer range size of mountain goat (*Oreamnos americanus*,
602 n=22) and white-tailed deer (*Odocoileus virginianus*, n=25). The left column refers to the model where
603 *Firmicutes* to *Bacteroides* ratios was a predictor, the middle column refers to the model where Pielou's
604 evenness was a predictor, and the right column refers to the model where observed richness (ASVs) was
605 the predictor.
606

Mountain goat

	Core range size					
Predictors	*Estimate*	*CI*	*Estimate*	*CI*	*Estimate*	*CI*
(Intercept)	0.31	0.10 - 0.51	-2.3	-4.1 - -0.42	0.19	0.026 - 0.35
F:B Ratio	0.010	-0.019 - 0.038	-	-	-	-
Pielou's evenness	-	-	2.8	0.85 - 4.8	-	-
Observed richness (ASVs)	-	-	-	-	5.7×10^{-6}	1.4×10^{-6} - 1.0×10^{-5}
Age class (Subadult)	0.20	0.018 - 0.39	0.17	0.020 - 0.32	0.16	3.3×10^{-4} - 0.31
Sex (Male)	-0.030	-0.17 - 0.10	-0.0047	-0.12 - 0.11	-0.012	-0.13 - 0.11
R^2 Nagelkerke	0.28		0.47		0.45	

White-tailed deer

	Core range size					
Predictors	*Estimate*	*CI*	*Estimate*	*CI*	*Estimate*	*CI*
(Intercept)	0.22	0.11 - 0.32	4.27	-0.67 -9.21	0.26	0.078 - 0.44
F:B Ratio	0.015	0.0050 - 0.025	-	-	-	-
Pielou's evenness	-	-	-4.16	-9.36 - 1.0	-	-
Observed richness (ASVs)	-	-	-	-	3.7×10^{-6}	-4.8×10^{-6} - 1.2×10^{-5}

Age class (Subadult)	0.033	-0.084 - 0.15	0.061	-0.068 - 0.19	0.069	-0.064 - 0.20
R^2 Nagelkerke	0.28		0.21		0.075	

607
608

Table 2. Generalized linear models for habitat proportion use for mountain goats (*Oreamnos americanus*, n=22*)* and white-tailed deer (*Odocoileus virginianus,* n=25). The left column refers to the model where *Firmicutes* to *Bacteroides* ratios was a predictor, the middle column refers to the model where Pielou's evenness was a predictor, and the right column refers to the model where observed richness (ASVs) was the predictor.

Mountain goat

Escape terrain proportion use

Predictors	Estimate	CI	Estimate	CI	Estimate	CI
(Intercept)	0.55	0.35 - 0.75	2.84	1.15 - 4.53	0.54	0.38 - 0.71
F:B Ratio	-0.01	-0.044 - 0.012	-	-	-	-
Pielou's evenness	-	-	-2.59	-4.43 - -0.76	-	-
Observed richness (ASVs)	-	-	-	-	-2.9×10^{-6}	-7.4×10^{-6} - 1.5×10^{-6}
Age class (Subadult)	0.15	-0.017 - 0.32	0.16	0.027 - 0.30	0.14	-0.013 - 0.31
Sex (Male)	-0.12	-0.24 - 0.00051	-0.14	-0.25- -0.039	-0.12	-0.24 - -0.0038
R^2 Nagelkerke	0.29		0.48		0.31	

Treed proportion use

Predictors	Estimate	CI	Estimate	CI	Estimate	CI
(Intercept)	0.62	0.19 - 1.1	5.41	1.85 - 8.98	0.88	0.55 - 1.2
F:B Ratio	-0.0044	-0.066 - 0.057	-	-	-	-
Pielou's evenness	-	-	-5.23	-9.09 - -1.36	-	-
Observed richness (ASVs)	-	-	-	-	-8.9×10^{-6}	-1.7×10^{-6} - -1.9×10^{-7}
Age class (Subadult)	-0.033	-0.40 - 0.33	0.057	-0.23 - 0.34	0.053	-0.26 - 0.36
Sex (Male)	-0.12	-0.38 - 0.14	-0.17	-0.39 - 0.05	-0.13	-0.37 - 0.10

	Estimate	CI	Estimate	CI	Estimate	CI
R^2 Nagelkerke	0.051		0.34		0.24	

Heat Load Index average

Predictors	Estimate	CI	Estimate	CI	Estimate	CI
(Intercept)	0.65	0.48 - 0.81	-0.15	-1.5 - 1.8	0.64	0.43 - 0.71
F:B Ratio	0.0086	-0.015 - 0.033	-	-	-	-
Pielou's evenness	-	-	0.60	-1.2 - 2.4	-	-
Observed richness (ASVs)	-	-	-	-	2.0×10^{-6}	-1.8×10^{-6} - 6.4×10^{-6}
Age class (Subadult)	-0.067	-0.21 - 0.073	-0.059	-0.19 - 0.075	-0.069	-0.22 - 0.072
Sex (Male)	-0.12	-0.22 - -0.17	-0.11	-0.22 - -0.011	-0.12	-0.22 - -0.017
R^2 Nagelkerke	0.28		0.28		0.30	

White-tailed deer

Forest proportion use

Predictors	Estimate	CI	Estimate	CI	Estimate	CI
(Intercept)	0.29	0.11 - 0.47	2.8	-4.6 - 10.4	0.23	-0.038 - 0.49
F:B Ratio	-0.0038	-0.021 - 0.013	-	-	-	-
Pielou's evenness	-	-	-2.7	-10.6 -5.2	-	-
Observed richness (ASVs)	-	-	-	-	2.1×10^{-6}	-1.0×10^{-5} - 1.45×10^{-5}
Age class (Subadult)	0.023	-0.18 - 0.22	0.0081	-0.19 - 0.20	0.014	-0.18 - 0.21
R^2 Nagelkerke	0.025		0.017		0.018	

Thicket swamp proportion use

Predictors	Estimate	CI	Estimate	CI	Estimate	CI
(Intercept)	0.19	0.082 - 0.30	-0.85	-5.4 - 3.7	0.25	0.091 - 0.40
F:B Ratio	-8.2×10^{-5}	-0.010 - 0.010	-	-	-	-

Predictors	Estimate	CI	Estimate	CI	Estimate	CI
Pielou's evenness	-	-	1.10	-3.7 - 5.9	-	-
Observed richness (ASVs)	-	-	-	-	-3.1×10^{-6}	-1.1×10^{-5} - 4.2×10^{-6}
Age class (Subadult)	-0.10	-0.23 - 0.016	-0.10	-0.22 - 0.016	-0.10	-0.22 - 0.012
R^2 Nagelkerke	0.12		0.13		0.15	

Treed swamp proportion use

Predictors	Estimate	CI	Estimate	CI	Estimate	CI
(Intercept)	0.35	0.17 - 0.53	-3.57	-11 - 3.9	0.40	0.14 - 0.67
F:B Ratio	0.0011	-0.016 - 0.018	-	-	-	-
Pielou's evenness	-	-	4.14	-3.7 - 12	-	-
Observed richness (ASVs)	-	-	-	-	-2.3×10^{-6}	-1.49×10^{-5} - 1.0×10^{-5}
Age class (Subadult)	0.049	-0.15 - 0.25	0.060	-0.13 - 0.26	0.052	-0.15 - 0.25
R^2 Nagelkerke	0.012		0.059		0.006	

615