Kinetochore-independent mechanisms of sister chromosome separation

Hannah Vicars[1], Travis Karg[1], Brandt Warecki[1], Ian Bast[1], William Sullivan[1]*.

[1] Department of Molecular, Cell and Developmental Biology, University of California, Santa Cruz, Santa Cruz, CA, USA

*Email: wtsulliv@ucsc.edu (WS)

Author contributions: HV contributed to all aspects of planning, execution of the experiments, analysis, and manuscript preparation. TK contributed to the planning, execution, and analysis of the RNAi screen and the planning and execution of live imaging of Greatwall hypomorphs. BW contributed to the planning, execution, and analysis of the telomere-labeling experiments. IB contributed to the execution and analysis of the telomere-labeling experiments. WS contributed to all aspects of experimental planning and manuscript preparation.

1
2 **ABSTRACT**
3

Although kinetochores normally play a key role in sister chromatid separation and

segregation, chromosome fragments lacking kinetochores (acentrics) can in some cases

separate and segregate successfully. In *Drosophila* neuroblasts, acentric chromosomes undergo

delayed, but otherwise normal sister separation, revealing the existence of kinetochore-

independent mechanisms driving sister chromosome separation. Bulk cohesin removal from the

acentric is not delayed, suggesting factors other than cohesin are responsible for the delay in

acentric sister separation. In contrast to intact kinetochore-bearing chromosomes, we discovered

that acentrics align parallel as well as perpendicular to the mitotic spindle. In addition, sister

acentrics undergo unconventional patterns of separation. For example, rather than the

simultaneous separation of sisters, acentrics oriented parallel to the spindle often slide past one

another toward opposing poles. To identify the mechanisms driving acentric separation, we

screened 117 RNAi gene knockdowns for synthetic lethality with acentric chromosome fragments.

In addition to well-established DNA repair and checkpoint mutants, this candidate screen

identified synthetic lethality with X-chromosome-derived acentric fragments in knockdowns of

Greatwall (cell cycle kinase), EB1 (microtubule plus-end tracking protein), and Map205

(microtubule-stabilizing protein). Additional image-based screening revealed that reductions in

Topoisomerase II levels disrupted sister acentric separation. Intriguingly, live imaging revealed

that knockdowns of EB1, Map205, and Greatwall preferentially disrupted the sliding mode of sister

acentric separation. Based on our analysis of EB1 localization and knockdown phenotypes, we

propose that in the absence of a kinetochore, microtubule plus-end dynamics provide the force to

resolve DNA catenations required for sister separation.

AUTHOR SUMMARY

Kinetochores, the site on the chromosomes to which microtubules attach driving the

separation and segregation of replicated sister chromosomes, have been viewed as essential for

proper cell division and accurate transmission of chromosomes into daughter cells. However

previous studies demonstrated that sister chromosomes lacking kinetochores (acentrics) often

undergo separation, segregation and transmission. Here we demonstrate that sister acentrics are

held together through DNA intertwining. We show that during anaphase, acentric sister

separation is achieved through Topoisomerase activity, an enzyme that resolves these DNA

linkages, as well as forces generated on the acentrics by the growing ends of highly dynamic

microtubule polymers. We found that acentric sister chromatids display unique patterns of

separation using mechanisms independent of the kinetochore. Additionally, we identified the

specific microtubule-associated proteins required for the successful mitotic transmission of

acentric chromosomes to daughter cells. These studies reveal unsuspected, distinct forces that

likely act on all chromosomes during mitosis independent of kinetochore-microtubule

attachments.

INTRODUCTION

Eukaryotic cells have evolved mechanisms to detect and protect against genomic insults.

These mechanisms include checkpoint pathways that delay cell cycle progression allowing time

for repair as well as apoptotic pathways that eliminate the damaged cells from the dividing

population [1]. Although a great deal is known regarding the function of these corrective pathways

during interphase, much less is known about the mechanisms that protect against genomic

instability after exit from metaphase. Studies demonstrate that DNA damage persisting through

metaphase delays anaphase onset. This delay is mediated both by the DNA damage and spindle

assembly checkpoint pathways [2, 3].

Despite these mechanisms, if the DNA damage remains, the checkpoints are overridden,

and the cell exits metaphase [4]. The persistence of unrepaired double-strand breaks (DSBs) at

metaphase is particularly problematic due to the formation of chromosome fragments, one of

which lacks a telomere and the other lacking a kinetochore and a telomere. The latter type are

known as acentrics and are incapable of forming canonical microtubule-kinetochore attachments

that drive sister chromosome separation and segregation. Consequently, acentrics would be

expected to lag on the metaphase plate and exhibit severe segregation defects. In accord with

this expectation, acentrics often fail to segregate, are excluded from daughter nuclei, and

subsequently form cytoplasmic micronuclei [5-7]. However, a growing number of reports

demonstrate poleward migration of acentric chromosome fragments [5, 8-13]. Proposed

mechanisms of acentric segregation include neo-centromere formation [14] and direct

association of the acentric chromosome with microtubules [15] or a kinetochore-bearing

chromosome [16-18].

Acentrics are efficiently induced in *Drosophila* bearing an I-CreI endonuclease

transgene, which fortuitously recognizes a repetitive sequence within the pericentric rDNA repeats

of the *Drosophila* X chromosome [19-22]. Induction of I-CreI expression results in the formation

of acentrics in over 80% of third instar larval neuroblast cells [12]. Although acentrics lag behind

on the metaphase plate well after the intact chromosomes migrate toward opposite poles, the

acentrics have a remarkable ability to accurately separate, segregate and incorporate into

daughter telophase nuclei [23]. A previous study done in the lab found that acentric segregation

relies on the chromokinesin Klp3A and interpolar microtubules [15]. However, it remains unclear

how sister acentrics are held together on the metaphase plate well after the main chromosome

mass has separated. Additionally, it is unknown how acentric sisters are able to initially separate

from one another instead of segregating together poleward. These behaviors reveal the existence

of kinetochore-independent mechanisms maintaining sister chromosome association on the

metaphase plate and driving their separation during anaphase. Possible explanations include

delayed acentric cohesin removal, delayed resolution of sister DNA catenations, or opposing plus-

end directed microtubule forces acting on each sister. Here we employ a combination of synthetic

lethal screens and live imaging to identify factors required for proper separation of sister acentrics.

Live analysis reveals three distinct modes of acentric separation: unzipping, sliding, and

simultaneous dissociation. This candidate screen revealed that Topoisomerase II, the cell-cycle

regulator Greatwall kinase, the microtubule (MT) plus-end tracking protein EB1, and the MT-

associated protein Map205 provide key roles in sister separation of acentrics. In addition, gene

knockdowns of EB1, Map205, and Greatwall preferentially disrupt the sliding mode of sister

separation. As will be discussed, this analysis demonstrates the existence of kinetochore-

independent mechanisms facilitating sister chromosome separation.

RESULTS

Acentric sister separation, but not cohesin removal, is delayed during the metaphase-to-

anaphase transition

As described previously, acentric chromosome fragments are efficiently generated

through heat-shock induction of an I-CreI transgene that specifically targets and creates DSBs in

the rDNA repeats at the base of the *Drosophila* X chromosome [19]. In accord with previous

studies [12], live analysis of the resulting X chromosome acentrics in the larval neuroblasts

reveals that sister separation of the acentrics occurs on average 148 seconds (± 44, N=19) after

separation of the intact chromosomes (**Figure 1**). We define acentric sister separation as the

point in which the sister acentrics can be clearly distinguished. Acentric sister segregation is

defined as the interval between separation of sisters and their migration to the spindle pole. The

delay in timing of acentric separation is defined as the time elapsed between the initiation of intact

sister chromosome separation and that of acentric sister separation. Sister chromatids are held

together at metaphase by cohesin, a tripartite ring-like protein complex comprised of two structural

maintenance of chromosome proteins (SMC1, SMC3) and a kleisin subunit (Rad21/Scc1) [24,

25]. Initially, cohesin removal occurs only along the chromosome arms through proteolytic

cleavage of the kleisin subunit by separase just prior to anaphase onset [26-28]. Due to the

Sgo/PP2A-dependent protection mechanism, cohesin remains at the centromeric regions [29].

Once cohesin is removed, microtubules drive sister separation [30, 31].

Fig 1: Acentric sister separation is delayed relative to kinetochore-bearing chromosomes
during anaphase. (A) Still frames of a time-lapse movie of a mitotic neuroblast labeled with H2Av-RFP
and not expressing I-CreI. (B) Still frames of a time-lapse movie of a mitotic neuroblast with I-CreI induced
acentrics. Separation of sister acentrics (arrows) is delayed. Consequently, they lag on the spindle equator
but eventually separate, segregate, and are reincorporated into daughter nuclei. Bars, 2 μm. Time in
seconds. (C) Scatterplot showing the delay in acentric sister separation after anaphase onset. Delay
(seconds) was measured from when kinetochore-bearing chromosomes initiated separation to when
acentric sister chromosomes initiated separation. Bars represent mean and standard deviations.
115

To investigate if delays in cohesin release are responsible for the delayed separation of

acentric chromosomes, female neuroblast divisions were live imaged with a cohesin component,

Rad21, tagged with EGFP [32]. Time-lapse images of a control neuroblast expressing Rad21-

EGFP are shown in **Figure 2A** (chromosomes in magenta, cohesin in green). No lagging

chromosomes are observed and Rad21 is cleared off of all chromosomes just prior to anaphase

onset and separation of sister chromatids. **Figure 2B** and **Movie S1** show time- lapse images of

an I-CreI-expressing neuroblast division. Separation and segregation of sister acentrics is delayed

relative to the intact chromosomes. Interestingly, cohesin removal just prior to anaphase onset

from the acentrics and the intact kinetochore-bearing chromosomes occurs simultaneously. This

is more clearly seen in the single channel black and white cohesin images of **Figure 2B** (depicting

yellow outlined regions of **Figure 2B**). Quantification of the Rad21 fluorescent signal supports the

conclusion of a relatively synchronous removal of cohesin on the acentric and intact

chromosomes (**Figure 2C**). The finding that sister acentrics remain paired despite the absence

of cohesin and well after the intact chromosomes have separated, indicate additional forces must

hold sister acentrics together.

Fig 2: Cohesin complexes are cleared off of acentric sisters upon anaphase onset. Chromosomes
labeled with H2Av-RFP (magenta) and cohesin labeled with Rad21-EGFP (green). (A) Stills from a time-
lapse movie of a control mitotic neuroblast. (B) Still images from a time-lapse movie of a mitotic neuroblast
with I-CreI-induced acentrics (Movie S1). Acentrics (arrowhead) lag on the spindle equator. (C) Bar graphs
of a compilation of five videos of I-CreI- expressing neuroblasts showing the relative fluorescence intensities
in arbitrary units (AU) of chromosomes (H2Av-RFP, top) cohesin (Rad21-EGFP, bottom) around acentrics
(cyan outlined region) and the main mass of chromosomes (yellow outlined region) at time points 50, 40,

25, 0 s, prior to anaphase onset, respectively. Bars, 2 µm. Time in seconds. Error bars represent standard
deviations of fluorescence intensities at all points tested.
140

Acentric sister separation occurs via three distinct patterns

To examine dynamics of acentric separation, we imaged live neuroblasts expressing

I-CreI, the histone marker H2Av-RFP, and the telomere marker HOAP-GFP [33]. Marked

telomeres enabled us to determine the orientation of acentrics with respect to one another as they

aligned on the metaphase plate, separated, and segregated poleward [15]. We observed three

distinct patterns of sister acentric separation (**Figure 3**). In the most frequent pattern (49%, N=45),

acentric pairs separate by sliding past one another (**Figure 3A**, Top row: histone-labeled

chromosomes in magenta, HOAP-labeled telomeres in green). This is more clearly observed in

the single histone channel movie (**Figure 3A** Bottom row and **Movie S2**). Also, in contrast to intact

kinetochore-bearing chromosomes which always align perpendicular to the spindle, the paired

sister acentrics align either parallel or perpendicular to the spindle and division axis.

Fig 3: Sister acentrics separate via three distinct patterns. Chromosomes labeled with H2Av-RFP
(magenta) and telomeres labeled with HOAP-GFP (green). Telomeres indicated by yellow arrowheads.
Direction of separation indicated by white arrows. (A) Top row: Still images of a time-lapse movie of a mitotic
neuroblast with I-CreI induced acentrics showing sister acentrics (cyan arrows) lagging behind during
anaphase, paired with telomeres opposing, sliding past one another, and ultimately separating. Bottom row:
Black and white images of sister acentrics sliding past one another (see arrows) (Movie S2). (B) Top row:
Still images of a time-lapse movie of a mitotic neuroblast with I-CreI induced acentrics showing acentrics
(cyan arrows) lagging on the metaphase plate. In a process we term "unzipping", sister acentrics are aligned
with telomeres paired, initiate separation at their broken ends followed by separation of telomeres. Bottom
row: Black and white images of sister acentrics unzipping (see arrows). (C) Top row: Still images of a time-
lapse movie of a mitotic neuroblast with I-CreI induced acentrics showing sister acentrics (cyan arrows)
lagging behind during anaphase. In a process we term "simultaneous dissociation", sister acentrics
simultaneously separate along their entire length. Bottom row: Black and white images of sister acentrics
simultaneously separating along their entire length (see arrows). (D) Frequency of acentric sisters paired
with their telomeres aligned or in opposite orientations. (E) Measurements of the frequency of each mode
of acentric sister separation (x-axis) and the timing of acentric sister separation after intact chromosomes
separate (y-axis). Each dot represents on acentric pair. Black dots represent acentric sisters with telomeres
aligned and purple dots represent acentric sisters with telomeres oriented in opposite directions. Boxes show
interquartile ranges and lines show medians of the measured data. Asterisks represent statistical
significance (***P=0.0008) determined by Mann-Whitney tests. (F) Chart showing the distribution of telomere
orientations within each mode of acentric sister separation. Acentrics that separate by unzipping are always
oriented with their telomeres paired.
174

The second pattern of acentric separation occurs via an "unzipping" mechanism (**Figure**

3B). This pattern occurs at a frequency of 31% (N=45). During separation, sister acentrics often

first separate at the broken end followed by separation at the telomere. This is illustrated in **Figure**

3B: the telomeres remain associated while the broken ends are well separated. Rarely, the paired

acentrics unzip from the telomere end first. In contrast to the acentrics that separate by sliding,

these paired sister acentrics are frequently aligned perpendicular to the spindle and division axis.

In the third pattern of separation, the remaining 20% (N=45), acentric sisters cleanly

separate from one another along their entire length similar to that observed for centric

chromosomes (**Figure 3C**). We termed this centric-like pattern: simultaneous dissociation.

Acentrics that separate via simultaneous dissociation are aligned in multiple orientations on the

metaphase plate (from parallel to perpendicular) with respect to the spindle and division axis. At

separation, dissociating sister acentrics simultaneously separate along their entire lengths.

We next analyzed the orientation of sister acentrics with respect to one another at the time

of separation through labeling telomeres with HOAP-GFP. As expected, the vast majority (87%,

39/45) of sister acentrics were oriented with their telomeres paired and aligned (**Figure 3D**).

However, a small but notable fraction (13%, 6/45), aligned with their telomeres opposed (**Figure**

3D). Acentric pairs that align with telomeres opposed presumably have already lost catenation

and/or cohesin in order to adopt this geometry. 14% and 33% of acentric sisters that separated

by sliding and dissociating, respectively, aligned with their telomeres opposed. Interestingly, sister

acentrics that separated by unzipping were never observed aligned with their telomeres opposed.

This suggests that there is an absolute requirement for sister pairing in acentrics that separate by

unzipping.

To further characterize these three modes of acentric separation, we measured the time

from anaphase onset (as determined by separation of the intact chromosomes) to sister acentric

separation (**Figure 3E**). Sliding acentrics separated much later than acentrics (202 ± 51 seconds)

that separated either by unzipping (131 ± 54 seconds) or dissociating (106 ± 55 seconds) (N=45,

Figure 3E). These differences were statistically significant as determined by two-sided Mann-

Whitney tests (P=0.0008 and P=0.0008, respectively). We did not detect any difference in the

timing of acentric sister separation by unzipping or dissociating (P=0.25 as determined by a two-

sided Mann-Whitney test).

Taken together, these results demonstrate three distinct patterns for acentric separation.

We note that based on the movements and orientation of both sliding and dissociating acentrics,

it is possible that catenations are lost along the entire length of the acentric pair simultaneously

at the moment of separation. In contrast, acentrics that separate by unzipping and appear to have

an absolute requirement for sister pairing may be more tightly associated with one another, and

this tight association may be due to lingering catenations.

Synthetic lethal screen identifies genes required for separation of sister acentrics

To identify the mechanisms required for transmission of acentric sister chromosomes, we

screened candidate RNAi gene knockdowns that resulted in synthetic lethality in the presence of

acentrics. The rationale for this screen is based on previous studies demonstrating that I-CreI

induction of acentric chromosomes during third instar larval stage resulted in only slight reductions

in adult survival because the acentrics are efficiently transmitted to daughter nuclei [12]. However,

reducing or partially disrupting the function of genes required for the normal acentric transmission

results in a dramatic reduction in adult survival upon acentric induction [12, 15]. Thus, we

expected that a subset of the gene knockdowns that resulted in synthetic lethality upon I-CreI

induction would be required for proper pairing and segregation of sister acentrics.

To perform the screen, adult flies bearing a heat-shock inducible I-CreI endonuclease and

Gal4 under the control of a ubiquitously expressed Actin enhancer element (Act5) were crossed

to adults bearing UAS-gene specific RNAi constructs. Lethality of heat-shocked (I-CreI induced)

and non-heat-shocked (I-CreI not induced) F1 progeny bearing both constructs were assayed.

RNAi constructs that significantly increased lethality upon I-CreI induction were of particular

interest. We screened 117 candidate genes for synthetic lethality upon acentric induction

(**Table S1**). These included genes encoding proteins spanning a diversity of mitotic

functions, including microtubule-associated proteins, chromatin remodelers, DNA repair genes,

cell cycle kinases, and cell cycle checkpoints. For each RNAi line, we determined the survival

ratio of the RNAi knockdown with I-CreI induced to the RNAi knockdown alone (RNAi knockdown

+ I-CreI/RNAi knockdown) (**Table 1**). Because these RNAi knockdowns do not cause complete

lethality, we classified hits as having a survival ratio of 65% or less (**Table 1**).

	Gene RNAi	Survival rate with RNAi and I-CreI induced / Only RNAi induced	Rate of larvae-to-adult survival of RNAi with I-CreI induction % ± SD (N)	Rate of larvae-to-adult survival of RNAi without I-CreI induction % ± SD (N)	Gene function
	Wildtype (Control)		80 ± 21 (194)	91 ± 7 (194)	—
Chromatin-associated:	hyx (2)	10%	8 (25)	88 (25)	Transcription factor
	cap-D2	12%	6 (17)	50 (12)	Condensin
	chd1	30%	10 ± 17 (69)	33 ± 6 (99)	ATP-dependent chromatin remodeling
	sin3A	33%	8 (25)	24 (25)	Histone deacetylase, transcription regulator
	chrac-14 (2)	48%	48 (25)	100 (25)	Nucleosome mobilization, chromatin assembly
	chd1 (2)	50%	48 (25)	95 (23)	ATP-dependent chromatin remodeling
	spt4	55%	24 (25)	44 (25)	Transcription elongation factor
	sir2	60%	64 (25)	94 (16)	Histone deacetylase, heterochromatin gene silencing
	hp1e	65%	65 (20)	100 (25)	Heterochromatin protein 1 (chromosome segregation)
Microtubule-associated:	msps	27%	18 ± 17 (56)	67 ± 24 (30)	Mitotic spindle, microtubule nucleation
	asp (2)	46%	44 (25)	96 (25)	Astral microtubule organization, spindle orientation
	EB1	47%	44 ± 25 (86)	94 ± 9 (38)	Microtubule plus-end binding, chromosome segregation
	Klp68D	51%	29 (24)	57 (21)	Kinesin motor protein, microtubule-based transport
	asp	52%	52 (25)	100 (15)	Astral microtubule organization, spindle orientation
	map205	65%	54 ± 18 (141)	83 ± 13 (91)	Binds/stabilizes microtubules
DNA repair/ checkpoint:	spnA	2%	2 ± 4 (57)	82 ± 18 (99)	Rad51 double-strand break repair
	mus309	11%	10 ± 1 (42)	95 (22)	Double-strand break repair
	ku80	29%	28 (25)	96 (25)	Non-homologous end joining double-strand break repair
	tefu	43%	43 (30)	100 (25)	DNA damage checkpoint
	cp190	48%	12 (17)	25 (12)	Chromatin silencing, DNA damage response
Cell cycle kinase:	bubR1	3%	1 ± 2 (168)	31 ± 23 (69)	Mitotic exit and spindle checkpoint kinase
	gwl (2)	8%	2 ± 2 (112)	26 ± 10 (99)	Negative PP2A regulator and mitotic exit kinase
	ald	59%	52 (25)	88 (25)	Altered disjunction, chromosome separation kinase

236 **Table 1: Top hits from synthetic lethality screen**
237 Top hits from the synthetic lethality screen were grouped by gene function. Overall survival rate ratio was
238 determined by the following: (percentage of surviving larvae after RNAi and I-CreI induction) / (percentage
239 of surviving larvae after only RNAi induction). Those RNAi transgenes that resulted significant reduction in
240 survival (a ratio of less than 0.65) were considered for follow-up live analysis.
241

The acentrics are generated through I-CreI induced double-strand breaks. As expected,

RNAi knockdowns of DNA repair and cell cycle checkpoints exhibit synthetic lethality upon I-CreI

induction. Genes involved in DNA repair all exhibit strong synthetic lethality upon I-CreI induction.

DSB repair genes *spnA/Rad51* [34], *mus309* [35] and NHEJ gene *Ku80* [36] exhibited survival

ratios of 2%, 11%, and 29%, respectively (**Table 1**). The DNA damage checkpoint gene *tefu* (ATM

[37]) also results in a pronounced synthetic lethality (survival ratio of 43%). The screen also

yielded a number of microtubule-associated proteins. These included *msps* (microtubule

nucleation [38]), *asp* (astral microtubule organization [39]), *eb1* (plus-end microtubule binding

[40]), *klp68D* (kinesin motor protein [41]), and *map205* (microtubule stabilizer [42]). Given that

microtubule-based transport plays a key role in the poleward transport of the acentric

chromosome fragments [15], synthetic lethal interactions with microtubule-associated proteins

were expected. However, it was unclear whether microtubules and their associated proteins also

play a role in the initial separation of sister acentrics during the metaphase-to-anaphase transition.

The screen also yielded a large class of genes involved in chromatin organization. Synthetic lethal

interactions with chromatin organizing proteins were expected due to the presence of I-CreI

induced DSBs. This included *cap-D2* (condensin [43]), *chd1* (ATP-dependent chromatin

remodeler [44]), *sin3A* and *sir2* (histone deacetylases [45]), *chrac* (nucleosome mobilization [46]),

and *hp1* (heterochromatin protein [47]). Whether these are directly required for double-strand

break repair, separation and/or segregation of sister acentrics remains to be determined. In

accord with previous work, reduced levels of BubR1 kinase (spindle-assembly checkpoint protein)

also exhibited a pronounced synthetic lethality upon I-CreI induction [12]. The screen also yielded

two additional cell cycle kinases: *gwl* (greatwall kinase, an inhibitor of the cell cycle phosphatase

PP2A [48]) and *ald* (altered disjunction, chromosome segregation [49]). We chose to focus on

EB1 and Map205 due to their well-documented association with microtubules [40, 42]. Greatwall

was chosen for follow-up because previous studies demonstrated that this kinase is required for

sister separation of intact chromosomes [50].

Live imaging analysis reveals the microtubule-associated proteins Map205 and EB1 and

Topo II are required for separation, but not segregation of sister acentrics.

We conducted live imaging experiments on neuroblasts to investigate the effect of specific RNAi-mediated gene knockdowns on acentric mitotic transmission. Each line contains an RFP-tagged histone transgene facilitating live confocal analysis [51]. Based on previous studies revealing the role of microtubules in acentric transmission [15], we initially focused on the microtubule-stabilizing protein Map205 and the microtubule plus-end associated protein EB1.

In a wild-type background, the majority of acentrics line up at the outer edge of the metaphase plate, separated from the main mass of intact chromosomes [12, 15]. As described above and in previous publications, during anaphase sister acentrics remain paired on the metaphase plate well after the separation of intact chromosomes (**Figure 4A**) [12, 15]. On average, separation of sister acentrics occurs 148 seconds ± 44 (N=19) after separation of the kinetochore-bearing chromosomes (**Table 2**). 83% (19/23) of acentric sister chromatid pairs separate normally with sister acentrics going to opposite cell poles (**Figure 4C**). In the remaining 17% (4/23), acentric sister chromatids line up in metaphase, lag behind at the metaphase plate, and segregate together to one pole of the cell (**Table 3**). Micronuclei form in telophase in 24% (4/17) of neuroblasts expressing I-CreI micronuclei form in telophase (**Figure 4D**).

Fig 4: EB1 RNAi knockdowns specifically disrupt acentric sister separation. (A) Still frames of a time-lapse movie of a mitotic neuroblast with I-CreI induced acentrics. Paired sister acentrics (white and yellow arrowheads) lag behind at the spindle equator but eventually separate, segregate, and are reincorporated into daughter nuclei. (B) Still frames of a time-lapse movie of a mitotic neuroblast with I-CreI induced acentrics and expressing EB1 RNAi. Separation of paired sister acentrics (white and yellow arrowheads) fails, resulting in sisters segregating to and incorporating into the same daughter nucleus. Bars, 2 µm. Time in seconds. (Video 3) (C) Percentages of neuroblast divisions in which acentric sisters failed to completely separate from one another. (D) Percentages of neuroblast divisions in which acentrics failed to incorporate into daughter nuclei and formed one or more micronuclei.

	Avg ± SD (seconds)	N (acentric pairs)
Control	148 ± 44	19
MAP205 RNAi	97 ± 39 **	12
EB1 RNAi	142 ± 57	16
Topo II RNAi	226 ± 77 *	10
gwl^{1028}/gwl^{716}	138 ± 49	22

298 **Table 2: Sister separation of acentrics is delayed relative to sister separation of intact chromosomes**
299 Average time (in seconds) for sister acentrics to initiate separation after sister kinetochore-bearing
300 chromosomes separated. Time (average ± SD) measured from initiation of intact chromosome separation
301 to initiation of acentric sister chromosome separation. Asterisks indicate statistical significance (**P=0.001,
302 *P=0.003) as determined by two-sided Mann-Whitney tests.
303

	Fails to separate	Dissociate	Slides Apart	Unzip	N (acentric pairs)
Control	17%	13%	48%	22%	23
MAP205 RNAi	46%	9%	9%	36%	22
EB1 RNAi	47%	20%	13%	20%	30
Topo II RNAi	54%	14%	27%	5%	22
gwl^{1028}/gwl^{716}	39%	11%	28%	22%	36

305 **Table 3: Acentric separation occurs through three distinct modes**
306 Acentric sister chromatids either fail to separate or separate from one another by three different modes:
307 laterally sliding past one another, unzipping from one another, or simultaneously and evenly dissociating
308 along their lengths.
309

Live imaging of acentric behavior in neuroblasts expressing Map205 RNAi revealed that

46% (10/22) of acentric sister chromatids do not separate from one another and segregate to one

cell pole together, in comparison to 17% (4/23) in a wildtype background (**Figure 4C, Figure 5B,**

Table 3). Additionally, after partial knockdown of Map205, sister acentrics separated from one

another significantly earlier than acentrics in a wildtype background (P=0.001, Mann-Whitney test,

Table 2). In spite of defects in acentric sister chromatid separation, their average poleward

segregation velocity was normal (**Table 4**). Despite the increase in failed acentric separation,

there was not an increase in micronuclei formation (21% compared to 24% in controls) (**Figure**

4D).

Fig 5: Knockdowns of EB1 and Map205 preferentially disrupt the sliding mode of acentric separation. (A) Model of different acentric orientations prior to separation. Polarity axis indicated by the dotted line. (B) Bar graph showing the percentage of acentric pairs that separate via simultaneous dissociation, sliding, or unzipping and the percentage of those that fail to separate. (C-G) Measurements of the frequency of each mode of acentric sister separation (x-axis) and the timing of acentric sister separation after the separation of intact chromosomes (y-axis). Each dot represents one acentric pair. Blue dots represent acentric pairs that were oriented perpendicular then parallel to the polarity axis prior to separation. Orange dots represent acentric pairs that were oriented parallel to the polarity axis prior to separation. Green dots represent acentric pairs that were oriented perpendicular to the polarity axis prior to separation. Boxes show interquartile ranges and lines show medians of the measured data. Asterisks indicate statistical significance (*P = 0.01, **P = 0.005, ***P = 0.009) when comparing the timing of acentric separation between separation modes. Statistical analysis was done using two-sided Mann-Whitney tests. Non-significant values had a P-value greater than 0.05.

	Avg velocity ± SD (nm/s)	N (individual acentrics)
Control	10 ± 4	26
MAP205 RNAi	12 ± 5	30
EB1 RNAi	10 ± 4	27
Topo II RNAi	12 ± 3	26
gwl^{l028}/gwl^{716}	8 ± 4 *	32

Table 4: Average velocities of acentric segregation
Depicted below are the velocities of individual acentrics as they segregate during anaphase. Velocity is measured in nanometers per second. Segregation velocity is defined as beginning when an acentric is clearly observed as separated from its sister and ends when the acentric reaches the daughter nucleus. Asterisks indicate statistical significance (*P=0.006) as determined by a two-sided Mann-Whitney test.

Live analysis of acentric behavior in neuroblasts expressing EB1 RNAi revealed that 47% (14/30) of acentric sister pairs fail to separate (**Figure 4C, Figure 5E, Table 3**). The failure of separation results in acentric sisters segregating together to a single cell pole (**Figure 4B, Movie S3**). Additionally, the delay in acentric separation and their rate of poleward segregation were normal (**Figure 5E, Table 2, Table 4**). There was also not an increase in micronuclei formation (15% compared to 24% in controls) (**Figure 4D**). This indicates that the Map205 and EB1 knockdowns are specifically disrupting sister separation of acentrics and have no effect on the latter stages of acentric transmission.

As described earlier, cohesin removal on the intact chromosomes and acentrics occurs simultaneously, yet acentric sisters remain paired. This raised the possibility of a role for DNA catenation in maintaining acentric pairing. Previous work demonstrated that DNA catenations

preserve sister chromatid cohesion in intact chromosomes until resolution by Topoisomerase II

(Topo II) at the metaphase-anaphase transition [52]. Topo II decatenates intertwined DNA

allowing sister chromatids to segregate to opposing poles of the cell. To test the role of DNA

catenation in maintaining acentric sister pairing, we knocked down levels of Topo II specifically in

the *Drosophila* neuroblast using the Gal4/UAS RNAi technique described above. Topo II

knockdowns revealed that 54% (12/22) of sister acentrics fail to separate and subsequently

segregate together to one pole (**Figure 6, Table 3, Movie S4**). Due to the lasting catenations

between acentric sisters, the initiation of acentric separation was further delayed in neuroblasts

expressing Topo II RNAi (Mann-Whitney test; P=0.003) (**Table 2**). Although acentric separation

is disrupted, their poleward segregation rate and their rate of micronuclei formation were normal

(**Figure 5F, Table 4, Figure 4D**).

Fig 6: Acentrics appear to be held together by DNA catenations. (A) Still images of a time- lapse
movie of a control neuroblast with I-CreI induced acentrics showing sister acentrics associated in early
anaphase (white arrows), lagging behind and ultimately separating and segregating to opposite poles (cyan
and magenta arrowheads). (B) Still images of a time-lapse movie of a neuroblast with partial knockdown of
Topoisomerase II using RNAi and I-CreI induced acentrics. Acentrics associate and lag behind in early
anaphase (white arrow), then fail to completely separate and segregate unequally (cyan and magenta
arrowheads). Bars, 2 µm. Time in seconds. (Movie S4) (C) Bar graph showing the quantification of acentric
behavior in control and Topo II RNAi neuroblasts with the failure of acentric sister separation in magenta
and the successful, even separation of acentric sisters in gray.
372

The screen also yielded the PP2A inhibitor Greatwall kinase (*gwl*) that controls the timing

and events of mitotic exit including sister chromatid separation [50]. Thus, we were particularly

interested in the dynamics of acentric separation and segregation when the levels of *gwl* were

reduced. This analysis uncovered that with reduced levels of *gwl*, 39% (14/36) of acentric sister

pairs failed to separate from one another (**Figure 4C, Figure 5B, Table 3**). Additionally, the delay

in acentric separation and the rate of micronuclei formation were normal (**Figure 5G, Table 2,**

Figure 4D). However, the average velocity of acentrics while segregating during anaphase was

significantly slower in *gwl* mutant background compared to acentrics in wildtype background

(P=0.006; Mann-Whitney test) (**Table 4**).

EB1, Map205, Greatwall, and Topo II differentially influence the dynamics of

acentric separation.

As described above, in a wild-type background acentrics exhibit three distinct patterns of separation: sliding, unzipping, and simultaneous dissociation. Sliding is the most frequent occurring 49% of the time, with unzipping and simultaneous dissociation occurring at frequencies of 31% and 20%, respectively (**Figure 3E**). To determine the role of the genes identified above in these distinct forms of acentric separation, we monitored the frequency of segregation patterns in RNAi knockdowns of the candidate genes. Of those sister acentrics that successfully separated, knockdowns of Map205, EB1, and Greatwall resulted in a decreased frequency of separation via sliding (17%, 25%, 45%, respectively) compared to the control (58%) (**Figure 5**). Additionally, Topoisomerase II resulted in a decreased the frequency of acentric separation via unzipping (10%) compared to the control (26%) (**Figure 5**).

To determine if microtubule plus-ends preferentially accumulate around acentrics, I-CreI-expressing neuroblasts were live imaged with EGFP-tagged EB1. Fluorescence intensities of both GFP and RFP were measured and corrected for brightness in control neuroblasts (N=5) and I-CreI-expressing neuroblasts (N=5). During mitosis, EB1 localizes along microtubules moving towards the plus-ends (**Figure 7 and Movie S5**). Previous studies have shown that EB1 is essential for generating antipolar forces on chromosomes [53, 54]. That finding together with our finding that EB1 is required for sister acentric separation motivated us to examine EB1 localization on the acentrics (**Figure 7**). We find the concentration of EB1 is not significantly increased on or near the acentrics ($P > 0.05$, two sample t-test; **Figure 7**). Thus, while EB1 and the acentrics co-localize, the pattern of EB1 comets is not altered in the presence of acentrics. While these data are consistent with a role for EB1 in acentric sister separation, there is not an increase in recruitment and accumulation of EB1 at the acentric.

Fig 7: Acentrics travel poleward in late anaphase while associated with EB1. EB1 is in green and chromosomes are in magenta. (A) Still images from a time-lapse movie of a control neuroblast from metaphase (0 s) through telophase (85 s). (B) Still images from a time-lapse movie of a mitotic neuroblast

with I-CreI induced acentrics from metaphase (0 s) through telophase (135 s) (Movie S5). The intact
chromosomes and acentrics initiate sister separation at the 15 s and 40 s timepoints, respectively. Sister
acentrics (white arrowheads) separate and move toward opposite cell poles while associated with EB1. (C)
Line graph from a compilation of five control videos showing the corrected fluorescence intensities in
arbitrary units (AU) of EB1 (green) and chromosomes (magenta). Corrected fluorescence intensities were
calculated within the yellow boxes at the time points 15, 20, 25, 30, 35 seconds after anaphase onset. (D)
Line graph from a compilation of five videos of I-CreI-expressing neuroblasts showing the corrected
fluorescence intensities in arbitrary units (AU) of EB1 (green) and chromosomes (magenta). Corrected
fluorescence intensities were calculated within the yellow boxes at the time points 15, 25, 30, 40, 50
seconds after anaphase. Bars, 2 µm. Time in seconds. Error bars represent SDs of the fluorescent
intensities at all points tested.
421

DISCUSSION

These studies are based on the unexpected finding that sister chromosome fragments

lacking a kinetochore undergo relatively normal separation. We sought to identify the kinetochore

independent forces driving acentric sister chromatid separation. A key finding is that while cohesin

removal occurs simultaneously on intact and acentric chromosomes, sister separation of the latter

is significantly delayed. An explanation for this delay comes from studies of intact chromosomes

demonstrating that once cohesin is removed, sister acentrics remain held together through DNA

catenations [52, 55, 56]. Catenations are concentrated at the centromeric DNA and opposing

kinetochore microtubule interactions likely provide the resolving force [57-61]. Support for this

conclusion comes from studies of chromosome rearrangements in which centromeric

heterochromatin is displaced from the centromere [31]. Sister chromatin separation is specifically

delayed in these regions resulting in localized stretching during anaphase. A likely consequence

of being displaced from the centromere is that the ectopic heterochromatic regions no longer

experience sufficient kinetochore forces required to efficiently resolve sister DNA catenations. In

light of these studies, it is likely that acentric sisters remain associated well after separation of the

intact chromosomes through DNA catenations. This conclusion is supported by our finding that

reductions in the Topoisomerase II levels specifically disrupt acentric sister separation. As

described below, our studies suggest both plus-end and lateral microtubule interactions with the

acentrics provide the alternative force driving sister separation.

Intact chromosomes align perpendicular to the spindle. In contrast, we find acentric

chromosomes align either perpendicular or parallel to the spindle. When aligned parallel to the

spindle, acentrics travel with one tip leading towards the cell pole, possibly due to microtubule

lateral interactions with acentrics. Without a kinetochore, a combination of lateral and plus-end

microtubule interactions likely determines the final orientation of acentrics on the metaphase

plate. In addition to the multiple orientations, acentrics undergo distinct patterns of sister

separation that we have termed sliding, unzipping and simultaneous dissociation. Sliding of

sisters past one another toward opposite poles is the most common mode of acentric separation.

This mode occurs primarily when sister acentrics are oriented parallel to the spindle just prior to

separation suggesting lateral microtubule interactions provide the force driving this mode of

acentric separation (**Figure 3A, Figure 5C**). While sister separation of all acentrics is delayed

relative to intact chromosomes, the delay is much more pronounced for acentrics that undergo

separation by sliding. The delay may in part be due to the additional time it takes to establish the

multiple lateral interactions required to generate sufficient separation force. It is likely

chromokinesins provide the force driving separation, but these have yet to be identified.

In contrast to sliding, sister separation by unzipping occurs primarily when the acentrics

are oriented perpendicular to the spindle just prior to separation suggesting a limited role for lateral

microtubule interactions (**Figure 3B, Figure 5C**). During unzipping, acentrics generally initiate

separation at their broken ends with completion of separation occurring at the telomeres. The

delay of acentric separation by unzipping is much less than the delay observed for separation

by sliding. Previous studies demonstrated that the acentric and its centric partner are connected

via a DNA tether [12]. The preference for initiation of separation at the broken-ends suggests that

a DNA tether connecting the broken end to the kinetochore-bearing fragment may provide the

initial separation force (**Figure 3B**). While this tether is not thought to provide the force driving

acentric segregation [15], because of its association with the broken end of the acentric, it may

drive the initial stage of unzipping. In contrast to separation by sliding, telomeres are always

aligned in sisters that separate by unzipping (**Figure 3E, Figure 3F**). This suggests maintaining

gene-for- gene sister pairing is essential for this form of separation.

The third mode of acentric separation, simultaneous dissociation, is characterized by a

synchronized separation along the entire length of sister acentrics (**Figure 3C**). While sister

separation via simultaneous dissociation favors the perpendicular orientation, this bias is much

less dramatic compared to the unzipping mode (**Figure 5C**). The most distinguishing feature of

this form of separation is the high frequency in which sister acentrics aligned with their telomeres

opposed (33% compared to 14% and 0% for sliding and unzipping separation, respectively)

(**Figure 3F**). This high frequency of unaligned telomeres suggests weak connections between

sisters and that sister pairing is not required.

Synthetic lethal screens have proven an effective means of identifying factors required for

the successful mitotic transmission of acentric chromosome fragments [12, 15]. These screens

have identified proteins required for poleward transmission and final incorporation of acentrics

into the telophase nucleus. Here we focused on identifying genes required for successful

separation of sister acentrics. Of the 117 candidate genes screened, we identified 23 RNAi

knockdowns/mutations that resulted in a significant lethality upon acentric induction (**Table 1**).

Live analysis secondary screening revealed knockdowns of EB1 and Map205, and the cell cycle

kinase Greatwall resulted in dramatic defects in acentric sister separation. Live imaging also

revealed reducing levels of Topoisomerase II severely disrupted sister acentric separation.

Given that previous studies demonstrated the chromokinesin Klp3A is required for acentric

poleward transport [15], it is not surprising that microtubule-associated proteins also play a key

role in separation of acentric sisters. Reduced levels of EB1 and Map205 greatly increase the

frequency of failed sister separation. EB1 is a plus-end tracking protein that associates with a

number of regulatory proteins and is essential for generating anti-polar forces on the

chromosomes [62]. EB1 knockdowns most dramatically reduce the frequency of acentric

separation via sliding. In addition, for those acentric sisters that did separate by sliding in EB1

knockdowns, separation was greatly delayed. As lateral interactions between the microtubules

and chromatin are likely to drive sister separation by sliding, we suspect plus-end directed EB1

forces may play a role in orienting the acentric in order to establish lateral interactions. Support

for this idea comes from the finding of a synthetic lethal interaction between I-CreI induction and

reductions in the levels of Nod, a non-motile chromokinesin that associates with EB1 and is

involved in chromosome segregation [15, 63-65]. Surprisingly, a previous study in our lab found

no effect on acentric segregation in mutations in Nod [15]. Examining movies from this analysis

revealed no significant difference in acentric sister separation in a loss-of-function *nod* mutant

compared to the control (**Table S2**). Map205 is a microtubule-associated protein required for

targeting Polo kinase to spindle microtubules [42]. Thus, it is likely that the effect of the Map205

knockdowns on acentric sister separation is through disruption of Polo localization. Previous

studies demonstrated that Polo localizes to the DNA tether associated with the acentric and Polo

knockdowns disrupted acentric transmission with sister acentrics remaining on the metaphase

plate [12]. As with EB1, reductions in Map205 levels dramatically reduce the frequency of acentric

separation by sliding suggesting a disruption in lateral interactions between the acentrics and the

microtubules. While much evidence indicates that Polo functions at the centromere during sister

separation, our studies of the effects of Map205 and Polo knockdowns suggest that it is also

involved in promoting chromatid/spindle interactions that are independent of the centromere.

Reducing the levels of Topoisomerase II leads to a significant increase in failed separation of

sister acentrics, supporting the conclusion that once cohesin is removed sister acentrics remain

held together through DNA catenation. We suspect that reduced Topoisomerase II levels disrupt

sister separation of the acentrics more profoundly than that of intact chromosomes because the

latter primarily rely on kinetochore forces to resolve DNA catenations. Of the four mutants

analyzed through live analysis, reductions in Topoisomerase levels had the greatest effect of the

unzipping mode of acentric sister separation. This result is interesting given our finding that the

unzipping mode of acentric separation is likely the most reliant on sister pairing and consequently

are likely to be highly catenated.

Taken together, these data support a model in which multiple forces drive the separation

of sister acentrics. Acentrics that separate by sliding and unzipping tend to be oriented parallel

and perpendicular to the spindle, respectively. Unzipping could be initially driven by the DNA

tether, which connects the acentric to its centric partner, to initially separate one end of the

acentrics. Then, a combination of Topoisomerase II activity and microtubule forces could facilitate

the separation of the other end of the acentrics. Because the partial knockdown of Topoisomerase

II leads to a decreased frequency of acentrics unzipping, we hypothesize that the resolution of

DNA catenations by Topoisomerase II underlies the unzipping mode of separation. Due to the

finding that the tether does not appear to provide segregation forces on the acentrics [15], sliding

is likely driven by lateral microtubule interactions and possibly motor proteins. This is further

supported by the orientation of acentrics parallel to the axis of polarity and the association of EB1

and bundling of microtubules around acentrics in early anaphase. It remains unclear if acentrics

are directly interacting with microtubules or if microtubule-associated proteins are mediating the

interaction with acentrics. There is not an obvious mechanism for the simultaneous dissociation

of sister acentrics and is likely driven by a combination of factors (**Figure 8**). EB1 and Map205

may be required to establish microtubule interactions with the acentric, while in the absence of a

kinetochore high levels of Topoisomerase are needed to resolve DNA catenations between

sisters.

Fig 8: Schematic showing distinct modes of acentric separation. At some point during the metaphase-
to-anaphase transition, acentrics that separate by sliding orient parallel to the spindle and slide past one
another. This is likely driven by lateral associations between the microtubules and acentrics. The unzipping
mode occurs when separation initiates at the broken end followed by separation of sister telomeres. It may
be that the initial separation is driven by the DNA tether connecting the centric and acentric fragments.
Separation across the entire length of the acentric is referred to as simultaneous dissociation. It may be
that the DNA tether and acentric-microtubule interactions contribute equally for sister acentrics that
separate by simultaneous dissociation. (Chromosomes in magenta, microtubules in green, cohesin in cyan,
telomeres in orange).
547

It should be noted that although these mutants resulted in a high frequency of failed

acentric sister separation, we did not observe an equivalent increase in micronuclei. In contrast,

mutants that disrupted acentric poleward transport or the final stages of incorporation into

daughter nuclei resulted in an accompanying increase in micronuclei [66-68]. This indicates that

the knockdowns in the genes identified in this study specifically disrupt acentric sister separation

as the subsequent transmission occurs normally to allow incorporation of the acentric into the

daughter nuclei. This interpretation is in accord with a model that acentrics experience different

forces at different stages of their separation, transmission, and incorporation into daughter nuclei

[68].

There are numerous examples of lateral interactions between spindle microtubules and

intact chromosomes indicating that non-kinetochore forces influence anaphase chromosome

kinetics [69-71]. However, it has been difficult to pursue the underlying mechanisms because of

the dominance of kinetochore forces during anaphase. In analogy with studies of spindle

formation in cells lacking centrosomes that led to the discovery of unsuspected, yet conserved,

chromosome-based mechanisms of spindle assembly, examining the mitotic behavior of acentric

chromosomes without a kinetochore will provide insights into the kinetochore independent forces

acting on intact chromosomes [70, 71].

566
567
568

MATERIALS AND METHODS

<u>Fly Stocks</u>

All stocks were raised on standard *Drosophila* media at room temperature (20-22°C) as previously described [72]. For generating acentrics, a transgenic fly line bearing the I-CreI endonuclease under heat-shock 70 promoter were kindly provided by Kent Golic at The University of Utah. For synthetic lethality screen, the ubiquitous Gal4 driver under the control of an actin enhancer (Act5) was used (#25708 from Bloomington). Dominant negative allele of ISWI was kindly provided by John Tamkun at UC Santa Cruz. Greatwall hypomorphs ($gwl^{1080, 716, 180, and 2790}$) were kindly provided by Michael Goldberg at Cornell University. The line with *rad21-EGFP* transgene were kindly provided by Stefan Heidmann at University of Bayreuth.

<u>Synthetic Lethality Screen</u>

Third instar progeny with genotype Act5-Gal4/+; I-CreI, Sb/UAS-RNAi were collected from parental genotypes Act5-Gal4/CyO-GFP; I-CreI, Sb/TM6B and were heat shocked for 1.5 hours at 37°C (unless otherwise indicated). After heat shock, the vials were set-aside at room temperature for 10-15 days until adult flies emerged. Synthetic lethality was calculated as the % of larvae that develop into adulthood [12, 15]. Control progeny with genotype Act5-Gal4/+; Sb/UAS-RNAi from parents with the genotype Act5-Gal4/CyO-GFP; Sb/TM6B and UAS-RNAi were heat-shocked for 1.5 hours at 37°C.

<u>Live analysis of acentric behavior in *Drosophila* third instar neuroblasts</u>

As previously described, acentric chromosome fragments were induced by I-CreI expression (under heat shock 70 promoter) in 3rd instar larvae by a 1-hour 37°C heat shock followed by a 1-hour recovery period at room temperature [12]. The larval brains from third instar larvae were dissected in PBS and then transferred to a slide with 20 µl of PBS. A coverslip was dropped on PBS with brain and the excess PBS was wicked out from edge of coverslip to induce squashing

of brain between slide and coverslip. For live analysis, the edge of coverslip was sealed with

halocarbon and was imaged as described below. Neuroblast divisions in all images were from

female 3rd instar larvae.

Microscopy and image acquisition

Wide-field microscopy

Time-lapse imaging for Figure 4C, 4D, 5C, and 6 were performed using a Leica DM16000B

wide-field inverted microscope equipped with a Hamamatsu electron-multiplying charge coupled

device camera (ORCA 9100-02) with a binning of 1 and a 100x Plan-Apochromat objective with

NA 1.4. Successive time points were filmed at 20 s. RFP (585 nm) and GFP (508 nm)

fluorophores were imaged. Samples were imaged in PBS and at room temperature (20-22°C).

Widefield images were acquired with Leica Application Suite Advanced Fluorescence Software

and 3D deconvolved using AutoQuant X2.2.0 software.

Spinning-disk microscopy

Images in Figures 1, 2, 3, 4, 6, and 7 were acquired with an inverted Nikon Eclipse TE2000-E

spinning disk (CSLI-X1) confocal microscope equipped with a Hamamatsu electron-multiplying

charge coupled device camera (ImageEM X2) with a 100X 1.4 NA oil-immersion objective.

Samples were imaged in PBS and at room temperature (20-22°C). Images were acquired with

MicroManager 1.4 software. Time-lapse fluorescent images of neuroblasts divisions were done

with 120 and 100 ms exposures for GFP and RFP respectively with 0.5 µm Z-steps. Time-lapse

videos with both GFP and RFP were done every 5 to 9 seconds and time-lapse movies with

RFP alone were done every 5 seconds. Figures were assembled in Adobe Illustrator. Selected

stills (both experimental and control) were processed with ImageJ (http://rsb.info.nih.gov/ij/).

631
632
633

634 <u>Measurements</u>
635
In Figure 2C, relative fluorescence intensities of chromosomes (H2Av-RFP) and cohesin

(Rad21-EGFP) were done using the plot profile function in ImageJ of the region outlined around

acentrics and the main mass of chromosomes. In Figures 7C and 7D, relative fluorescence

intensities of acentrics (H2Av-RFP) and EB1 (EB1-EGFP) were done using the plot profile

function in ImageJ of the region outlined around acentrics in I-CreI expressing neuroblasts and

the region outlined around the spindle midzone in control neuroblasts. For statistical analyses,

unpaired two-sided t-tests and two-sided Mann-Whitney-Wilcoxon tests were used. Unpaired

two-sided t-tests were performed in Prism Version 8 (GraphPad Software). Two-sided Mann-

Whitney-Wilcoxon tests were performed in R (R Core Team) and Prism Version 8 (GraphPad

Software).

646
647 **ACKNOWLEDGEMENTS**
648

We would like to thank Dr. Susan Strome, Dr. William Saxton, and Dr. John Tamkun for use of

their equipment and providing reagents. We thank Dr. Benjamin Abrams for his advice and

technical expertise regarding microscopy experiments. We thank Dr. Kent Golic for providing us

with the transgenic fly line bearing the I-CreI endonuclease. We thank Dr. Michael Goldberg for

providing us with Greatwall hypomorph flies. We thank Dr. Stefan Heidmann for providing us

with the transgenic fly line bearing *rad21-EGFP*. We thank H. Vicars' service dog Teddy Bear

for his guidance and support.

Funding for these studies was provided by National Institutes of Health grant

NIH1RO1GM120321 awarded to W. Sullivan.

The authors declare no competing financial interests.

661
662

REFERENCES

1. Ermolaeva MA, Schumacher B. Systemic DNA damage responses: Organismal adaptations to genome instability. Trends Genet. 2014;30(3):95–102. pmid:24439457; PubMed Central PMCID: PMC4248340

2. Mikhailov A, Cole RW, Rieder CL. DNA Damage during Mitosis in Human Cells Delays the Metaphase/Anaphase Transition via the Spindle-Assembly Checkpoint. Current Biology. 2002;12(21):1797–806. pmid:12419179

3. Royou A, Macias H, Sullivan W. The Drosophila Grp/Chk1 DNA Damage Checkpoint Controls Entry into Anaphase. Current Biology. 2005;15(4):334–9. pmid:15723794

4. Coutelier H, Xu Z. Adaptation in replicative senescence: a risky business. Curr Genet. 2019;65(3):711–6. pmid:30637477

5. Kanda T, Wahl GM. The dynamics of acentric chromosomes in cancer cells revealed by GFP-based chromosome labeling strategies. J Cell Biochem Suppl. 2000; Suppl 35:107–14. pmid:11389539

6. LaFountain JR, Oldenbourg R, Cole RW, Rieder CL. Microtubule Flux Mediates Poleward Motion of Acentric Chromosome Fragments during Meiosis in Insect Spermatocytes. Mol Biol Cell. 2001;12(12):4054–65. pmid:11739800; PubMed Central PMCID: PMC60775

7. Fenech M, Kirsch-Volders M, Natarajan AT, Surralles J, Crott JW, Parry J, et al. Molecular mechanisms of micronucleus, nucleoplasmic bridge and nuclear bud formation in mammalian and human cells. Mutagenesis. 2011;26(1):125–32. pmid:21164193

8. Bajer A. Cine-micrographic studies on chromosome movements in β-irradiated cell. Chromosoma. 1958;9:319-331. pmid:13608834

9. Liang H, Wright WH, Cheng S, He W, Berns MW. Micromanipulation of chromosomes in PTK2 cells using laser microsurgery (optical scalpel) in combination with laser-induced optical force (optical tweezers). Experimental cell research. 1993;204(1):110–20. pmid:8416789

10. Khodjakov A, Cole RW, Bajer AS, Rieder CL. The force for poleward chromosome motion in Haemanthus cells acts along the length of the chromosome during metaphase but only at the kinetochore during anaphase. J Cell Biol. 1996;132(6):1093–104. pmid:8601587; PubMed Central PMCID: PMC2120764

11. Kanda T, Sullivan KF, Wahl GM. Histone–GFP fusion protein enables sensitive analysis of chromosome dynamics in living mammalian cells. Current Biology. 1998;8(7):377–85. pmid:9545195

12. Royou A, Gagou ME, Karess R, Sullivan W. BubR1- and Polo-Coated DNA Tethers Facilitate Poleward Segregation of Acentric Chromatids. Cell. 2010;140(2):235–45. pmid:20141837; PubMed Central PMCID: PMC2969851

13. Bretscher HS, Fox DT. Proliferation of Double Strand Break Resistant Polyploid Cells Requires Drosophila FANCD2. Dev Cell. 2016;37(5):444–57. pmid:27270041; PubMed Central PMCID: PMC4901310

14. Platero JS, Ahmad K, Henikoff S. A Distal Heterochromatic Block Displays Centromeric Activity When Detached from a Natural Centromere. Molecular Cell. 1999;4(6):995–1004. pmid:10635324

15. Karg T, Elting MW, Vicars H, Dumont S, Sullivan W. The chromokinesin Klp3a and microtubules facilitate acentric chromosome segregation. J Cell Biol. 2017;216(6):1597–608. pmid:28500183; PubMed Central PMCID: PMC5461011

16. Kanda T, Otter M, Wahl GM. Mitotic segregation of viral and cellular acentric extrachromosomal molecules by chromosome tethering. J Cell Sci. 2001; 114:49-58. pmid:11112689

17. Ishii K, Ogiyama Y, Chikashige Y, Soejima S, Masuda F, Kakuma T, et al. Heterochromatin Integrity Affects Chromosome Reorganization After Centromere Dysfunction. Science. 2008;321(5892):1088–91. pmid:18719285

18. Ohno Y, Ogiyama Y, Kubota Y, Kubo T, Ishii K. Acentric chromosome ends are prone to fusion with functional chromosome ends through a homology-directed rearrangement. Nucleic Acids Res. 2016;44(1):232–44. pmid:26433224; PubMed Central PMCID: PMC4705696

19. Rong YS, Titen SW, Xie HB, Golic MM, Bastiani M, Bandyopadhyay P, et al. Targeted mutagenesis by homologous recombination in D. melanogaster. Genes Dev. 2002;16(12):1568–81. pmid:12080094; PubMed Central PMCID: PMC186348

20. Maggert KA, Golic KG. Highly Efficient Sex Chromosome Interchanges Produced By I-CreI Expression in Drosophila. Genetics. 2005;171(3):1103–14. pmid:16020774; PubMed Central PMCID: PMC1456814

21. Paredes S, Maggert KA. Ribosomal DNA contributes to global chromatin regulation. Proc Natl Acad Sci U S A. 2009;106(42):17829–34. pmid:19822756; PubMed Central PMCID: PMC2764911

22. Golic MM, Golic KG. A simple and rapid method for constructing ring-X chromosomes in Drosophila melanogaster. Chromosoma. 2011;120(2):159–64. pmid:21085980; PubMed Central PMCID: PMC4454366

23. Warecki B, Sullivan W. Mechanisms driving acentric chromosome transmission. Chromosome Res. 2020;Online ahead of print. pmid:32712740

24. Guacci V, Koshland D, Strunnikov A. A Direct Link between Sister Chromatid Cohesion and Chromosome Condensation Revealed through the Analysis of MCD1 in S. cerevisiae. Cell. 1997;91(1):47–57. pmid:9335334; PubMed Central PMCID: PMC2670185

25. Michaelis C, Ciosk R, Nasmyth K. Cohesins: Chromosomal Proteins that Prevent Premature Separation of Sister Chromatids. Cell. 1997;91(1):35–45. pmid:9335333

26. Uhlmann F, Lottspeich F, Nasmyth K. Sister-chromatid separation at anaphase onset is promoted by cleavage of the cohesin subunit Scc1. Nature. 1999;400(6739):37–42. pmid:10403247

27. Uhlmann F, Wernic D, Poupart M-A, Koonin EV, Nasmyth K. Cleavage of Cohesin by the CD Clan Protease Separin Triggers Anaphase in Yeast. Cell. 2000;103(3):375–86. pmid:11081625

28. Oliveira RA, Hamilton RS, Pauli A, Davis I, Nasmyth K. Cohesin cleavage and Cdk inhibition trigger formation of daughter nuclei. Nat Cell Biol. 2010;12(2):185–92. pmid:20081838; PubMed Central PMCID: PMC3284228

29. Gutiérrez-Caballero C, Cebollero LR, Pendás AM. Shugoshins: from protectors of cohesion to versatile adaptors at the centromere. Trends in Genetics. 2012;28(7):351–60. pmid:22542109

30. Asbury CL. Anaphase A: Disassembling Microtubules Move Chromosomes toward Spindle Poles. Biology. 2017;6(1):15. pmid:28218660; PubMed Central PMCID: PMC5372008

31. Oliveira RA, Kotadia S, Tavares A, Mirkovic M, Bowlin K, Eichinger CS, et al. Centromere-Independent Accumulation of Cohesin at Ectopic Heterochromatin Sites Induces Chromosome Stretching during Anaphase. PLoS Biol. 2014;12(10):e1001962. pmid:25290697; PubMed Central PMCID: PMC4188515

32. Urban E, Nagarkar-Jaiswal S, Lehner CF, Heidmann SK. The Cohesin Subunit Rad21 Is Required for Synaptonemal Complex Maintenance, but Not Sister Chromatid Cohesion, during Drosophila Female Meiosis. PLoS Genet. 2014;10(8):e1004540. pmid:25101996; PubMed Central PMCID: PMC4125089

33. Cenci G, Siriaco G, Raffa GD, Kellum R, Gatti M. The Drosophila HOAP protein is required for telomere capping. Nat Cell Biol. 2003;5(1):82–4. pmid:12510197

34. Staeva-Vieira E, Yoo S, Lehmann R. An essential role of DmRad51/SpnA in DNA repair and meiotic checkpoint control. EMBO J. 2003;22(21):5863–74. pmid:14592983; PubMed Central PMCID: PMC275421

35. McVey M, Andersen SL, Broze Y, Sekelsky J. Multiple Functions of Drosophila BLM Helicase in Maintenance of Genome Stability. Genetics. 2007;176(4):1979–92. pmid:17507683; PubMed Central PMCID: PMC1950607

36. Melnikova L, Biessmann H, Georgiev P. The Ku Protein Complex Is Involved in Length Regulation of Drosophila Telomeres. Genetics. 2005;170(1):221–35. pmid:15781709; PubMed Central PMCID: PMC1449706

37. Oikemus SR, McGinnis N, Queiroz-Machado J, Tukachinsky H, Takada S, Sunkel CE, et al. Drosophila atm/telomere fusion is required for telomeric localization of HP1 and telomere position effect. Genes Dev. 2004;18(15):1850–61. pmid:15256487; PubMed Central PMCID: PMC517405

38. Moon W, Hazelrigg T. The Drosophila Microtubule-Associated Protein Mini Spindles Is Required for Cytoplasmic Microtubules in Oogenesis. Current Biology. 2004;14(21):1957–61. pmid:15530399

39. Saunders RDC, Avides M do C, Howard T, Gonzalez C, Glover DM. The Drosophila Gene abnormal spindle Encodes a Novel Microtubule-associated Protein That Associates with the Polar Regions of the Mitotic Spindle. J Cell Biol. 1997;137(4):881–90. pmid:9151690; PubMed Central PMCID: PMC2139842

40. Rogers SL, Rogers GC, Sharp DJ, Vale RD. Drosophila EB1 is important for proper assembly, dynamics, and positioning of the mitotic spindle. J Cell Biol. 2002;158(5):873–84. pmid:12213835; PubMed Central PMCID: PMC2173155

41. Pesavento PA, Stewart RJ, Goldstein LS. Characterization of the KLP68D kinesin-like protein in Drosophila: possible roles in axonal transport. The Journal of Cell Biology. 1994;127(4):1041–8. pmid:7525600; pmid:7525600; PubMed Central PMCID: PMC2200055

42. Archambault V, D'Avino PP, Deery MJ, Lilley KS, Glover DM. Sequestration of Polo kinase to microtubules by phosphopriming-independent binding to Map205 is relieved by phosphorylation at a CDK site in mitosis. Genes Dev. 2008;22(19):2707–20. pmid:18832073; PubMed Central PMCID: PMC2559908

43. Savvidou E, Cobbe N, Steffensen S, Cotterill S, Heck MMS. Drosophila CAP-D2 is required for condensin complex stability and resolution of sister chromatids. Journal of Cell Science. 2005;118(11):2529–43. pmid:15923665

44. Stokes DG, Tartof KD, Perry RP. CHD1 is concentrated in interbands and puffed regions of Drosophila polytene chromosomes. Proc Natl Acad Sci USA. 1996;93(14):7137–42. pmid:8692958; PubMed Central PMCID: PMC38949

45. Feller C, Forné I, Imhof A, Becker PB. Global and Specific Responses of the Histone Acetylome to Systematic Perturbation. Molecular Cell. 2015;57(3):559–71. pmid:25578876

46. Corona DFV, Eberharter A, Budde A, Deuring R, Ferrari S, Varga-Weisz P, et al. Two histone fold proteins, CHRAC-14 and CHRAC-16, are developmentally regulated subunits of chromatin accessibility complex (CHRAC). EMBO J. 2000;19(12):3049–59. pmid:10856248; PubMed Central PMCID: PMC203371

47. Eissenberg JC, Elgin SCR. HP1a: A Structural Chromosomal Protein Regulating Transcription. Trends Genet. 2014;30(3):103–10. pmid:24555990; PubMed Central PMCID: PMC3991861

48. Yu J, Fleming SL, Williams B, Williams EV, Li Z, Somma P, et al. Greatwall kinase: a nuclear protein required for proper chromosome condensation and mitotic progression in Drosophila. J Cell Biol. 2004;164:487-492. pmid:14970188; PubMed Central PMCID: PMC2171981

49. O'Tousa J. Meiotic chromosome behavior influenced by mutation-altered disjunction in Drosophila melanogaster females. Genetics. 1982;102:503-524. pmid:6816677; PubMed Central PMCID: PMC1201954

50. Voets E, Wolthuis RMF. MASTL is the human orthologue of Greatwall kinase that facilitates mitotic entry, anaphase and cytokinesis. Cell Cycle. 2010;9(17):3591-601. pmid:20818157

51. Schuh M, Lehner CF, Heidmann S. Incorporation of Drosophila CID/CENP-A and CENP-C into centromeres during early embryonic anaphase. Curr Biol. 2007;17(3):237-243. pmid:17222555

52. Porter ACG, Farr CJ. Topoisomerase II: untangling its contribution at the centromere. Chromosome Res. 2004;12(6):569–83. pmid:15289664

53. Tirnauer JS, Bierer BE. EB1 proteins regulate microtubule dynamics, cell polarity, and chromosome stability. J Cell Biol. 2000;149(4):761-6. pmid:10811817; PubMed Central PMCID: PMC2174556

54. Vitre B, Coquelle FM, Heichette C, Garnier C, Chrétien D, Arnal I. EB1 regulates microtubule dynamics and tubulin sheet closure in vitro. Nat Cell Biol. 2008;10(4):415-21. pmid:18364701

55. Toyoda Y, Yanagida M. Coordinated Requirements of Human Topo II and Cohesin for Metaphase Centromere Alignment under Mad2-dependent Spindle Checkpoint Surveillance. Mol Biol Cell. 2006;17(5):2287–302. pmid:16510521; PubMed Central PMCID: PMC1446084

56. Murray AW, Szostak JW. Chromosome Segregation in Mitosis and Meiosis. Annual Review of Cell Biology. 1985;1(1):289–315. pmid:3916318

57. Farcas A-M, Uluocak P, Helmhart W, Nasmyth K. Cohesin's Concatenation of Sister DNAs Maintains Their Intertwining. Mol Cell. 2011;44(1–3):97–107. pmid:21981921; PubMed Central PMCID: PMC3240746

58. Wang LH-C, Mayer B, Stemmann O, Nigg EA. Centromere DNA decatenation depends on cohesin removal and is required for mammalian cell division. Journal of Cell Science. 2010;123(5):806–13. pmid:20144989

59. Kenney RD, Heald R. Essential roles for cohesin in kinetochore and spindle function in Xenopus egg extracts. Journal of Cell Science. 2006;119(24):5057–66. pmid:17158911

60. Baxter J, Sen N, Martinez VL, De Carandini MEM, Schvartzman JB, Diffley JFX, et al. Positive Supercoiling of Mitotic DNA Drives Decatenation by Topoisomerase II in Eukaryotes. Science. 2011;331(6022):1328–32. pmid:21393545

61. Piskadlo E, Oliveira RA. A Topology-Centric View on Mitotic Chromosome Architecture. Int J Mol Sci. 2017;18(12):2751. pmid:29258269; PubMed Central PMCID: PMC5751350

62. Nehlig A, Molina A, Rodrigues-Ferreira S, Honoré S, Nahmias C. Regulation of end-binding protein EB1 in the control of microtubule dynamics. Cell Mol Life Sci. 2017;74(13):2381–93. pmid:28204846

63. Theurkauf WE, Hawley RS. Meiotic spindle assembly in Drosophila females: Behavior of nonexchange chromosomes and the effects of mutations in the nod kinesin-like protein. J Cell Biol. 1992;116:1167-1180. pmid:1740471; PubMed Central PMCID:PMC2289365

64. Ye AA, Verma V, Maresca TJ. NOD is a plus end–directed motor that binds EB1 via a new microtubule tip localization sequence. J Cell Biol. 2018;217(9):3007–17. pmid:29899040; PubMed Central PMCID: PMC6122986

65. Afshar K, Barton NR, Hawley RS, Goldstein LSB. DNA binding and meiotic chromosomal localization of the Drosophila nod kinesin-like protein. Cell. 1995;81:129-138. pmid:7720068

66. Karg T, Warecki B, Sullivan W. Aurora B–mediated localized delays in nuclear envelope formation facilitate inclusion of late-segregating chromosome fragments. Mol Biol Cell. 2015;26(12):2227–41. pmid:25877868; PubMed Central PMCID: PMC4462941

67. Warecki B, Sullivan W. Micronuclei formation is prevented by Aurora B-mediated exclusion of HP1a from late-segregating chromatin in Drosophila. Genetics. 2018;210(1):171-187. pmid:29986897; PubMed Central PMCID: PMC6116970

68. Warecki B, Ling X, Bast I, Sullivan W. ESCRT-III–mediated membrane fusion drives chromosome fragments through nuclear envelope channels. Journal of Cell Biology. 2020;219(3):e201905091. pmid:32032426; PubMed Central PMCID: PMC7054997

69. Laband K, Le Borgne R, Edwards F, Stefanutti M, Canman JC, Verbavatz JM, et al. Chromosome segregation occurs by microtubule pushing in oocytes. Nature Comm. 2017;8(1):1499. pmid:29133801; PubMed Central PMCID: PMC5684144

70. Heald R, Tournebize R, Blank T, Sandaltzopoulos R, Becker P, Hyman A, et al. Self-organization of microtubules into bipolar spindles around artificial chromosomes in Xenopus egg extracts. Nature. 1996;382(6590):420-5. pmid:8684481

71. Fuge H. Non-kinetochore transport phenomena, microtubule-chromosome associations, and force transmission in nuclear division. Protoplasma. 1990;158:1-9.

72. Royou A, McCusker D, Kellogg DR, Sullivan W. Grapes (Chk1) prevents nuclear CDK1 activation by delaying cyclin B nuclear accumulation. J Cell Biol. 2008;183(1):63-75. pmid:18824564; PubMed Central PMCID: PMC2557043

A — Control (No I-CreI): 0 s, 135, 162, 189, 234, 279, 441; H2Av-RFP

B — I-CreI: 0 s, 15, 65, 85, 125, 145, 205; H2Av-RFP

C — Acentric separation delay; Time (seconds); I-CreI Alone; N = 19

	Metaphase	Anaphase
Sliding		
Unzipping		
Dissociating		

Cohesin   Telomere