1 **Article Title**
2 An acylsugar-deficient *Nicotiana benthamiana* strain for aphid and whitefly research
3
4 **Running title**
5 Acylsugars protect *Nicotiana benthamiana*
6
7

8 **Author names**
9 Honglin Feng[a], Lucia Acosta-Gamboa[b], Lars H. Kruse[c,f], Jake D. Tracy[d], Seung Ho Chung[a],
10 Alba Ruth Nava Fereira[e], Sara Shakir[a,g], Hongxing Xu[a,h], Garry Sunter[e], Michael A. Gore[b], Clare
11 L. Casteel[d], Gaurav D. Moghe[c], Georg Jander[a*]
12
13
Author Affiliations
[a]Boyce Thompson Institute, Ithaca NY, USA
[b]Plant Breeding and Genetics Section, School of Integrative Plant Science, Cornell University,
Ithaca NY, 14853, USA
[c]Plant Biology Section, School of Integrative Plant Science, Cornell University, Ithaca NY,
14853, USA
[d]Plant-Microbe Biology and Plant Pathology Section, School of Integrative Plant Science,
Cornell University, Ithaca NY, 14853, USA
[e]Department of Biology, University of Texas San Antonio, San Antonio TX, 78249, USA
[f]Present address: Michael Smith Laboratories, University of British Columbia, Vancouver, BC,
V6T 1Z4, Canada
[g]Present address: Gembloux Agro-Bio Tech Institute, the University of Liege, Gembloux,
Belgium
[h]Present address: College of Life Science, the Shaanxi Normal University, Xi'an, China
28
29
30
*Correspondence:
Georg Jander
Boyce Thompson Institute
Ithaca, NY 14853
USA
Phone: 607-254-1365
Email: gj32@cornell.edu

Abstract

Nicotiana benthamiana is used extensively as a platform for transient gene expression and as a model system for studying plant-virus interactions. However, many tobacco-feeding generalist herbivores, including *Myzus persicae* (green peach aphid), *Bemisia tabaci* (whitefly), *Macrosiphum euphorbiae* (potato aphid), *Heliothis virescens* (tobacco budworm), *Trichoplusia ni* (cabbage looper), and *Helicoverpa zea* (corn earworm), grow poorly on *N. benthamiana*, limiting its utility for research on plant-insect interactions. Using CRISPR/Cas9, we generated knockout mutations in two *N. benthamiana* acylsugar acyltransferases, *ASAT1* and *ASAT2*, which contribute to the biosynthesis of insect-deterrent acylsucroses. Whereas *asat1* mutations reduced the abundance of two predominant acylsucroses, *asat2* mutations caused almost complete depletion of foliar acylsucroses. The tested hemipteran and lepidopteran species survived, gained weight, and/or reproduced significantly better on *asat2* mutant plants than on wildtype *N. benthamiana*. Furthermore, both *asat1* and *asat2* mutations reduced the water content and increased the temperature of leaves, indicating that foliar acylsucroses can protect against desiccation. Two experiments demonstrated the utility of the *N. benthamiana asat2* mutant line for insect bioassays. Transmission of turnip mosaic virus by *M. persicae* was significantly improved by an *asat2* mutation. Tobacco rattle virus constructs were used for virus-induced gene silencing of *acetylcholinesterase*, *squalene synthase*, *toll-like receptor 7*, and *tubulin-specific chaperon D* genes in *B. tabaci*, an experiment that would have been difficult with wild-type *N. benthamiana* due to high insect mortality. Additionally, the absence of acylsugars in *asat2* mutant lines will simplify transient expression assays for the functional analysis of acylsugar biosynthesis genes from other Solanaceae.

Introduction

Nicotiana benthamiana, a wild tobacco species that is native to Australia, is commonly used by plant molecular biologists as model system for laboratory research. Susceptibility to a wide variety of plant viruses has made *N. benthamiana* a popular model for fundamental studies of plant-virus interactions (Goodin et al., 2008). Scientists have developed *N. benthamiana* as a transgene expression powerhouse by engineering viral vectors to express heterologous genes, including fluorescent reporter genes to visualize cell structures (Bally et al., 2018). Antibodies, biofuel compounds, and other protein and metabolite products have been produced in *N. benthamiana* (Arntzen, 2015; Powell, 2015). Virus-induced gene silencing (VIGS), which is employed to study gene function in a variety of plant species, was originally developed in *N. benthamiana* (Velásquez et al., 2009; Hayward et al., 2011). Recently, high-efficiency CRISPR/Cas9 gene editing of germline cells using virus-encoded guide RNA (gRNA) was demonstrated for the first time in *N. benthamiana* (Ellison et al., 2020). Although *N. benthamiana* is hyper-susceptible to many plant viruses, it is not a good host for three virus-transmitting Hemiptera, *Myzus persicae* (green peach aphid; Thurston, 1961; Hagimori et al., 1993), *Bemisia tabaci* (whitefly; Simon et al., 2003), and *Macrosiphum euphorbiae* (potato aphid) that otherwise grow well on cultivated tobacco (*Nicotiana tabacum*).

The poor growth of generalist insect herbivores on *N. benthamiana* may be attributed in part to glandular trichomes. These epidermal secretory structures on the leaf surface of ~30% of vascular plants (Weinhold and Baldwin, 2011; Glas et al., 2012) have been found to play a crucial defensive role in several ways: as a physical obstacle for insect movement on the plant surface (Cardoso, 2008), entrapment (Simmons et al., 2004), synthesis of volatiles and other defensive metabolites (Laue et al., 2000; Schilmiller et al., 2010; Glas et al., 2012), and production of proteins that repel herbivores (phylloplane proteins, e.g. T-phylloplanin; Shepherd and Wagner, 2007). In addition to their defensive functions, glandular trichomes also protect plants from abiotic stresses such as transpiration water loss and UV irradiation (Karabourniotis et al., 1995).

There are two main types of glandular trichomes on *N. benthamiana* leaves, large swollen-stalk trichomes and small trichomes that are capped by a secretory head with one, two, or four cells (Slocombe et al., 2008). The large trichomes have been shown to secrete phylloplane proteins in *N. tabacum*. The small trichomes, which are the most abundant on

tobacco leaf surfaces, secrete exudates, including acylsugars (Wagner et al., 2004; Slocombe et

al., 2008). Detached trichomes, a mixture of the large and small trichomes, from *N. benthamiana*

are able to synthesize acylsugars (Kroumova and Wagner, 2003), and the secretory head cells

alone are able to synthesize acylsugars in *N. tabacum* (Kandra and Wagner, 1988).

Acylsugars, generally sucrose or glucose esterified with aliphatic acids of different chain

lengths (Figure 1), are abundant insect-deterrent metabolites produced by Solanaceae glandular

trichomes (Arrendale et al., 1990; Slocombe et al., 2008; Moghe et al., 2017). Specific

acylsugars are associated with aphid-resistant *Nicotiana* species, while not being detected in

more susceptible species in this genus (Hagimori et al., 1993). Furthermore, relative to cultivated

tomatoes (*Solanum lycopersicum*), acylsugars provide wild tomatoes (*Solanum pennellii*) greater

resistance against *M. persicae* (Rodriguez et al., 1993). Diacylsucrose protects crops against

tobacco aphids (*Myzus persicae nicotianae*), *B. tabaci*, and two-spotted spider mites

(*Tetranychus urticae*) (Chortyk et al., 1996; Alba et al., 2009). The synthetic sucrose octanoate

(an analog of *Nicotiana gossei* sugar esters) is effective in the field against Asian citrus psyllids

(*Diaphorina citri*), citrus leafminer (*Phyllocnistis citrella*), and a mite complex (including Texas

citrus mite, red spider mite, and rust mite) (McKenzie and Puterka, 2004; McKenzie et al.,

2005). Interestingly, acylsucroses in *Nicotiana attenuata* are metabolized to volatile fatty acids

by neonate *Manduca sexta* (tobacco hornworm) larvae, thereby tagging these larvae and

attracting predatory ants, *Pogonomyrmex rugosus* (Weinhold and Baldwin, 2011).

Acylsugars and leaf surface lipids more generally may contribute to plant drought

tolerance. Transcriptomic studies of drought-tolerant *S. pennellii* populations showed that lipid

metabolism genes are among those that are most responsive to drought stress (Gong et al., 2010;

Egea et al., 2018). Additionally, the abundance of acylsugars on the leaf surface in a native *S.*

pennellii population was associated with drought tolerance (Fobes et al., 1985). Acylsugars also

are reported to provide protection against drought stress conditions in *Solanum chilense* (O'

Connell et al., 2007). Similarly, abundant accumulation of acylsugars with C7-C8 acyl groups in

the desert tobacco (*Nicotiana obtusifolia*) was suggested to provide this species with high

drought tolerance for its desert environment (Kroumova et al., 2016). Although the mechanism is

not completely understood, it has been proposed that the polar lipids reduce the surface tension

of adsorbed dew water, thereby allowing the leaves absorb more condensed water on the surface

(Fobes et al., 1985).

More recently, enzymes involved in the biosynthesis of acylsugars have been identified.

Four acylsugar acyltransferases (ASATs), *Sl*ASAT1, *Sl*ASAT2, *Sl*ASAT3, and *Sl*ASAT4, have

been biochemically characterized in cultivated tomato (Fan et al., 2016). *Sl*ASAT1 catalyzes the

first step of sucrose acylation, using sucrose and acyl-CoA to generate monoacylsucroses via

pyranose R_4 acylation (Fan et al., 2016). *Sl*ASAT2 uses the product of *Sl*ASAT1 (*e.g.*

S1:5(iC5^{R4})) and acyl-CoA donor substrates (*e.g.* iC4-CoA, aiC5-CoA, nC10-CoA, and nC12-

CoA) to generate diacylsucroses (Fan et al., 2016). Further, *Sl*ASAT3 uses the diacylsucroses

generated by *Sl*ASAT2 to make triacylsucroses by acylating the diacylsucrose five-membered

(furanose) ring (Fan et al., 2016). Then, *Sl*ASAT4 makes tetraacylsucroses by acetylating

triacylsucroes using C2-CoA (Schilmiller et al., 2012). Notably, *Sl*ASAT4 (formerly *Sl*ASAT2)

is specifically expressed in the trichomes, where acylsugar acetylation occurs (Schilmiller et al.,

2012). The expression and activity of ASATs varies among different plant species (including the

order of the ASAT reactions in the pathway), which likely contributes to the observed trichome

chemical diversity (Kim et al., 2012). Although ASATs have been most intensively studied in

tomato, *ASAT* genes also have been annotated in the available *Nicotiana* genomes (Gaquerel et

al., 2013; Van et al., 2017; Egan et al., 2019). However, *ASAT* genes in *N. benthamiana* have not

been annotated and characterized previously, and their functions in protection against insect

pests and desiccation remained unknown.

The goal of this study was to confirm the role of acylsugars in *N. benthamiana* resistance

to insect feeding, as well as to create an insect-susceptible *ASAT* mutant line to facilitate use of

N. benthamiana for laboratory research on plant-insect interactions. We identified two *ASAT*

genes in *N. benthamiana*, *NbASAT1 and NbASAT2*. Using CRISPR/Cas9 to create mutant lines,

we showed that knockout of both *NbASAT1* and *NbASAT2* reduced acylsugar content. *NbASAT2*

mutations allowed increased survival, growth, and/or reproduction of six tested insect species.

Additionally, decreased water content and elevated leaf temperature in *NbASAT1* and *NbASAT2*

mutants indicated that *N. benthamiana* acylsugars contribute to protection against desiccation.

Results

Identification of **ASAT1** *and* **ASAT2** *in N. benthamiana*

Using reciprocal comparisons to confirmed Solanaceae *ASAT* genes (Moghe et al., 2017), we

identified three highly homologous sequences in the *N. benthamiana* genome:

Niben101Scf02239Ctg025, Niben101Scf22800Ctg001, and Niben101Scf14179Ctg028

(Bombarely et al., 2012) (gene identifiers are from annotations at solgenomics.net). Whereas

Niben101Scf02239Ctg025 and Niben101Scf22800Ctg001 were annotated as a full-length coding

sequences with strong coverage in available RNAseq datasets, Niben101Scf14179Ctg028 was

annotated as a pseudogene because it appears to be a fragment of the predicted cDNA

Niben101Scf141790g02010.1, but with no coverage in available RNAseq datasets. The

Niben101Scf02239Ctg025 and Niben101Scf22800Ctg001 sequences were confirmed in a more

recently assembled *N. benthamiana* genome (Schiavinato et al., 2019). In this assembly, the

pseudogene Niben101Scf14179Ctg028 was annotated as part of Niben101Scf02239Ctg025, and

there were no additional annotated ASAT candidates.

To infer ASAT evolution and function, we constructed a protein phylogenetic tree of

previously annotated Solanaceae ASATs (Figures 2, S1 and S2; Tables S1 and S2). In this tree,

Niben101Scf02239Ctg025 formed a monophyletic group with other ASATs, including the

biochemically characterized *Ss*ASAT1, *Pa*ASAT1 and *Hn*ASAT1. Therefore, we named

Niben101Scf02239Ctg025 as *N. benthamiana* ASAT1 (*Nb*ASAT1). Niben101Scf22800Ctg001

formed a monophyletic group with other ASATs including the biochemically characterized

*Na*ASAT2, *Hn*ASAT2, and *Pa*ASAT2. Therefore, we named Niben101Scf22800Ctg001 as *N.*

benthamiana ASAT2 (*Nb*ASAT2). Notably, the ASAT2 monophyletic group also included the

biochemically characterized *Sp*ASAT1, *Sl*ASAT1, and *Sn*ASAT1 (Figure 2).

Generation of* ASAT *mutants

Using CRISPR/Cas9 coupled with tissue culture, we obtained two independent homozygous

mutants for both *NbASAT1* and *NbASAT2*. *asat1-1* has a five-nucleotide deletion at the gRNA3

cutting site and a single nucleotide insertion at the gRNA2 cutting site, leading to a frameshift

between gRNA3 and gRNA2. *asat1-2* has a 318-nucleotide deletion between the gRNA3 and

gRNA2 cutting sites (Figure 3A). *asat2-1* has a single-nucleotide deletion at the gRNA3 cutting

site and single-nucleotide insertion at the gRNA2 cutting site, leading to a translation frame shift

between the two sites. *asat2-2* has a 115-nucleotide deletion at the gRNA3 cutting site and a

single-nucleotide insertion at the gRNA2 cutting site (Figure 3B). Homozygous mutants were

confirmed by DNA sequencing in the T2 generation, and this generation was used for all

experiments.

ASAT2 *knockout depletes acylsugar biosynthesis*

We quantified the acylsugar content in the *ASAT* mutants by LC/MS, comparing to that of wildtype *N. benthamiana* plants. In the LC/MS profile of *N. benthamiana* leaf surface washes, we characterized twelve mass features as acylsucroses based on their characteristic peaks and neutral losses (Figure S3A). In negative electron spray ionization mode, the characteristic peak features included the mass of 341.11 for sucrose, 509.22 for sucrose + C2+C8, 467.21 for sucrose + C8, 495.21 for sucrose + C7, and 383.12 for sucrose + C2; the neutral loss peaks included mass for 126.10 for C8 (acyl-chain with 8 carbons), 129.09 for C7 + H_2O, and 59.01 for C2 + H_2O (Figure S3B). Those twelve *m/z* ratios included 383.12, 467. 21, 509.22, 555.23, 593.32, 621.31, M625.31, 635.32, 639.32, 667.32, 671.30, 681.34 (Figure S3A). Based on their MS/MS peak features, retention times and relative abundances, we predicted that the identified mass features are mainly derived from two acylsucroses as formate or chloride adducts, pathway intermediates, and/or resulted from in-source fragmentation. We named the two acylsucroses S3:17(2,7,8) and S3:18(2,8,8) (in the nomenclature, "S" refers to the sucrose backbone, "3:18" indicates three acyl chains with total eighteen carbons, and the length of each acyl chain is shown in parentheses) (Figure 1; Figure S3A).

In wildtype plants, S3:18(2,8,8) is the dominant acylsucrose, whereas S3:17(2,7,8) has relatively low abundance (Figure 4). S2:16(8,8) and S2:15(7,8), which may be biosynthetic pathway intermediates for S3:18(2,8,8) and S3:17(2,7,8), respectively, are present at lower levels (Figure 4). Compared to wildtype *N. benthamiana*, both *asat2-1* and *asat2-2* were almost completely depleted in both S3:17(2,7,8) and S3:18(2,8,8), as well as in the two predicted biosynthetic intermediates S2:15(7,8) and S2:16(8,8) (Figure 4). For *asat1-1* and *asat1-2*, the detected acylsucroses were less abundant, and significantly reduced in *asat1-1* (Figure 4). Although acylsugar content was significantly reduced in the *ASAT* mutants, the structure and abundance of trichomes on the leaf surface were not visibly changed (Figure S4).

*Insect performance is improved on **ASAT2** mutant lines*

We used *asat1* and *asat2* mutant lines to test the role of acylsugars in protecting *N. benthamiana* against insect pests. After placing synchronized first-instar aphids on mutant and wildtype *N. benthamiana* leaves, we monitored survival and growth over time. Significant improvements in

aphid survivorship were observed as early as at 2 days post-feeding ($p < 0.001$) on the *asat2-1*
and *asat2-2* mutants and increased until the end of the 5-day monitoring period ($p < 0.001$)
(Figure 5A and Table S3). After 5 days of feeding, surviving aphids on both *asat1* and *asat2*
plants were larger than those on wildtype plants (Figures 5B). When we measured progeny
production by five adult aphids over a period of seven days, an average of more than 200
nymphs were produced on the *asat2* mutants, significantly more than the number of nymphs
produced on either wildtype or *asat1* mutants ($p < 0.05$, Figure 5C). In aphid choice assays with
detached leaves, they preferentially chose *asat2-1* mutant leaves relative to wildtype and *asat1-1*
leaves ($p < 0.05$, Figures 5D-F). A preference for *asat2-1* and *asat2-2* leaves was consistently
observed in choice assays involving any pairwise combination with wildtype, *asat1-1*, and *asat1-*
2 leaves ($p < 0.001$, *Chi-square* test, Figures 5D-F and S5). No *M. persicae* preference was
observed when comparing wildtype *N. benthamiana* and *asat1* mutants ($p > 0.05$, Figures 5D-F
and S5).
When aphid colonies were allowed to grow long-term on *asat2-1* mutant and wildtype *N.*
benthamiana in the same growth chamber, there were many more aphids on the mutant plants
(Figure S6A,B), likely resulting from a combination of host plant choice and increase growth on
the *asat2-1* mutant. It is noteworthy that, on the *asat2-1* mutant plants, aphids were feeding on
the more nutritious younger leaves, which tend to be better-defended in plants. By contrast, on
wildtype *N. benthamiana* aphids were only were able to feed on older, senescing leaves and were
primarily on the abaxial surface. Consistent with the increased aphid presence, growth of the
asat2-1 mutant plants was visibly reduced relative to wildtype *N. benthamiana* (Figure S6B).
Given the almost identical phenotypes of *asat2-1* and *asat2-2* mutants, subsequent insect assays
were conducted with T2 progeny of the *asat2-1* line, which also were confirmed by PCR to
contain the Cas9 transgene.
As we observed with *M. persicae*, the *asat2-1* mutation improved the *M. euphorbiae*
performance on *N. benthamiana* (Figure 6A-C). Significantly increased *M. euphorbiae* survival
was observed after 24 hours on *asat2-1* compared to wildtype ($p < 0.001$, Figure 6A).
Additionally, significantly more nymphs were produced by adult *M. euphorbiae* in the course of
24 hours on *asat2-1* than on wildtype ($p < 0.05$, Figure 6B). In choice assays, potato aphids
preferentially chose the *asat2-1* leaves over wildtype leaves ($p < 0.001$, Figure 6C). Whereas we

were not able to establish an *M. euphorbiae* colony on wildtype *N. benthamiana*, the aphids

readily formed colonies on the *asat2-1* mutant plants (Figure 6D).

Survival of *B. tabaci* was greatly increased on the *asat2-1* mutant line relative to wildtype

(Figure 6E). Moreover, whiteflies laid significantly fewer eggs on wildtype *N. benthamiana* than

on *asat2-1* mutant plants over three days (Figure 6F). Dead adult whiteflies were observed on

wildtype plants (Figure S7A), and it was not possible to establish a reproducing colony. By

contrast, after 23 days of feeding on *N. benthamiana*, whiteflies of different life stages were

observed on *asat2-1* mutant plants (Figure S7B-D). In choice assays with mutant and wildtype

plants in the same cage, *B. tabaci* preferentially settled on *asat2-1* plants (72%) over wildtype

(26%) in a 24-hour experiment (Figure 6G). Notably, all whiteflies on *asat2-1* were alive after

24 hours, whereas about half of the whiteflies found on the wildtype plants were dead at the

same time point (Figure 6F).

To determine whether depletion of acylsugars in *ASAT2* mutants improves the

performance of lepidopteran herbivores on *N. benthamiana*, we conducted experiments with

Helicoverpa zea (corn earworm), *Heliothis virescens* (tobacco budworm) and *Trichoplusia ni*

(cabbage looper). When neonates were placed on the leaves of wildtype or *asat2-1* mutant *N.*

benthamiana, no *H. zea* caterpillars were recovered (Figure 7A). Survivorship of H. *virescens*

and *T. ni* larvae on *N. benthamiana* was low, and the mass of the surviving larvae after ten days

was not significantly increased on the mutant line relative to wildtype (Figure 7B,C). Due to the

low survival of neonates, we repeated the caterpillar bioassay using five-day-old larvae that had

been reared on artificial diet. Almost all *H. zea* and *H. virescens* larvae survived for seven days

on wildtype and *asat2-1* mutant plants, and survival of *T. ni* caterpillars was higher on *asat2-1*

than on wildtype plants (Figure 7D-F). The relative growth rates of surviving *H. zea*, *H.*

virescens, and *T. ni* larvae were higher on the *asat2-1* mutant by 35%, 47%, and 99%,

respectively, than on wildtype plants (Figure 7D-F).

Virus transmission by aphids is increased on* asat2-1 *mutant plants

To demonstrate the use of *asat2-1* mutants for aphid experiments, we measured *M. persicae*

transmission of a GFP-expressing turnip mosaic virus (TuMV-GFP; Lellis et al., 2002; Casteel et

al., 2015) using all combinations of wildtype and *asat2-1* plants as virus donors and recipients,

respectively (Figure 8). The earliest GFP signals in the recipient plants were visible under UV

light on day 4 after TuMV-GFP transmission. Successful transmission and virus replication was

monitored until day 9, when no plants developed additional visible TuMV-GFP infections

relative to the previous day (Figure 8). The *asat2-1* → *asat2-1* transmission group showed the

highest final TuMV-GFP infection rate (73%), significantly higher than that observed in the

other three groups ($p < 0.05$, ANOVA). Relative to the wildtype → wildtype TuMV-GFP

transmission, transmission was not significantly increased in the *asat2-1* → wildtype and

wildtype → *asat2-1* groups. The significantly higher virus transmission rate by aphids in the

asat2-1 → *asat2-1* group is consistent with the feeding preference of *M. persicae* for *asat2-1*

mutant plants (Figure 5).

The* asat2-1 *mutant is suitable for plant-mediated VIGS in whiteflies

To demonstrate the utility of the *asat2-1* mutant for whitefly experiments, we performed plant-

mediated virus-induced gene silencing (VIGS) using tobacco rattle virus (TRV; Hayward et al.,

2011). We selected two previously validated *B. tabaci* RNAi targets, *AchE* (*acetylcholinesterase*;

Malik et al., 2016) and *TLR7* (*toll-like receptor 7*; Chen et al., 2015), as well as two predicted

horizontally transferred genes (Chen et al., 2016), *SQS* (*squalene synthase*) and *TSCD* (*tubulin-

specific chaperone D*), that had not been previously investigated in whiteflies. When whiteflies

were place on *N. benthamiana* infected with TRV VIGS constructs, expression of all four target

genes was significantly reduced after one day of feeding (Figure 9A). After seven days of

feeding, the expression of *AchE* and *SQS* was significantly reduced relative to whiteflies on

empty vector control plants, but expression of *LTR7* and *TCSD* was not (Figure 9B). After seven

days, survivorship of the whiteflies on VIGS plants was significantly reduced relative to control

plants with empty vector or GFP control TRV infections (Figure 9C). Consistent with the low

survival observed in previous experiments (Figure 6E), whitefly survival on TRV-infected

wildtype *N. benthamiana* was very low (Figure 9D), never exceeding 10%. None of the VIGS

constructs decreased survival to a lower level than on the empty vector and TRV-GFP control

plants.

Water loss is greater in* asat2 *mutant plants that in wildtype

While conducting aphid choice assays with detached leaves (Figures 5D-F and 6C), we noticed

that the mutant leaves dried out faster than wildtype leaves. This effect was quantified using

detached-leaf assays, in which *asat2* leaves lost significantly more water over 24 hours than

leaves from either wildtype or *asat1* leaves (Figures 10A and S8A). When plants are subjected to

drought, high/low temperature, salinity, herbivores, or other stresses, they absorb and reflect

specific wavelengths of light, which have been used to determine vegetation indices. For

example, the water band index (WBI) (Penuelas et al., 1993) has been used to monitor changes

in the plant canopy/leaf water content. Using hyperspectral imaging, we determined that the leaf

water content of intact plants, as measured by the WBI, was significantly lower in *asat2* mutants

than in wildtype (Figures 10B and S8B). Although the *asat1* mutants did not lose water faster

than wildtype in detached leaf assays (Figure 10A), the leaf water content in *asat1* mutants was

significantly lower than wildtype (Figures 10B and S10B). Measurement of leaf temperature by

thermal imaging showed that, consistent with the reduced leaf water content, the leaf temperature

of the acylsugar mutants was significantly higher than that of wildtype plants (Figures 10C and

S10C).

Discussion

We were able to identify only two *ASAT* genes, *NbASAT1* and *NbASAT2*, as well as a fragmented

pseudogene, in the *N. benthamiana* genome (Table S2). By contrast, in other *Nicotiana* species,

there are larger numbers of predicted *ASATs*, *e.g.* one *ASAT1*, one *ASAT2*, and 20 *ASAT3*-like

genes in *N. attenuata* (Gaquerel et al., 2013; Van et al., 2017), 35 *ASAT3*-like genes in *N.*

tabacum, and 19 *ASAT3*-like genes in *N. tomentosiformis* (Egan et al., 2019). Given the

relatively small number of predicted *ASAT* genes in *N. benthamiana*, other enzymes may also be

involved in acylsugar biosynthesis. In the Solanaceae, two common pathways are known for

aliphatic acid elongation via acetate: the fatty acid synthase (FAS) and the alpha-ketoacid

elongation (αKAE) pathways (Kroumova and Wagner, 2003). In the FAS pathway, two carbons

from an acetyl-acyl carrier are retained per elongation cycle, whereas in the αKAE pathway, one

carbon is retained per elongation cycle (Kroumova and Wagner, 2003). Knockdown of *E1-β*

branched-chain α-keto acid dehydrogenase (*BCKD*) significantly reduces acylsugars in *N.*

benthamiana (Slocombe et al., 2008). Additionally, *Isopropylmalate Synthase 3* (*IPMS3*) in

cultivated and wild tomatoes (Ning et al., 2015) and *Acyl-Sucrose Fructo-Furanosidase 1*

(*ASFF1*) in wild tomato (Leong et al., 2019) are involved in determining acylsugar composition.

Further studies will be needed to characterize other genes involved in the *N. benthamiana*

acylsugar biosynthesis pathway.

Acylsugars can be categorized as sucrose or glucose esters based on the sugar cores,

which are decorated with varying numbers or lengths of acyl chains (Kim et al., 2012). Whereas

some wild tomatoes produce a mixture of acylsucroses and acylglucoses, we observed only

acylsucroses (Figures 1 and 4), consistent with previous identification of these compounds in *N.*

benthamiana (Matsuzaki et al., 1989; Matsuzaki et al., 1992; Hagimori et al., 1993; Slocombe et

al., 2008). Nevertheless, it has been reported that *N. benthamiana* produces acylglucoses,

although in lower abundance than acylsucroses (Hagimori et al., 1993), and one glucose ester

structure has been proposed (Matsuzaki et al., 1992). Our failure to detect acylglucoses may be

explained by the use of different isolates of *N. benthamiana*, growth conditions, growth stage of

plants, and/or the detection methods. Whereas we used ~1-month-old plants and LC/MS for our

assays, Matsuzaki et al. (1992) used ~3-month-old plants and GC/MS to detect acylglucoses in

N. benthamiana.

Abundance of the characterized acylsugars was reduced to a greater extent in *N.*

benthamiana asat2 than in *asat1* mutants (Figure 4). The smaller reduction of acylsugars in

asat1 mutants suggests that either *Nb*ASAT2 functions upstream of *Nb*ASAT1 in the acylsugar

biosynthesis pathway, but partially complements *Nb*ASAT1 activity; or *Nb*ASAT1 and

*Nb*ASAT2 have similar functions in the biochemical pathway, but *NbASAT2* had higher

abundance or enzymatic activity. It is not known whether *N. benthamiana* ASATs are

monomeric or multimeric, but BAHD acyltransferases generally are monomeric enzymes

(D'Auria, 2006), suggesting that heterodimers between ASAT1 and ASAT2 are unlikely to affect

the observed phenotypes.

In the ASAT phylogenetic tree (Figure 2), *Nb*ASAT2 is closely related to some

biochemically characterized ASAT1 proteins in other Solanaceae species, including the

*Sp*ASAT1, *Sl*ASAT1, and *Sn*ASAT1. Those ASAT1s have some substrate overlap with the

ASAT2s found in the corresponding species, indicating that ASAT2 has moved toward utilizing

the ASAT1 substrate in these species over time (Moghe et al., 2017). The final activity shift that

has become fixed in the *Solanum* genus, most likely occurred after the divergence of the

Solanum and *Capsicum* clades (Moghe et al., 2017). However, if our hypothesis of partial

complementation of *Nb*ASAT1 by *Nb*ASAT2 is correct, it may flag a transition stage or suggest

independent *Nicotiana*-specific evolution of the ASAT1 and ASAT2 functions. Based on

previous knowledge of BAHD activities (Moghe et al., 2017), we postulate that S2:15 (7,8) and

S2:16 (8,8) are produced by *Nb*ASAT1 and *Nb*ASAT2, whereas the acetylation is carried out by

another unrelated BAHD enzyme – not unlike the distantly related *Sl*ASAT4 and *Salpiglossis*

sinuata ASAT5 enzymes (Schilmiller et al., 2015; Moghe et al., 2017). Further characterization

will be required to identify specific acyltransferase enzyme activities in *N. benthamiana*.

The observed role of acylsugars in protecting against desiccation (Figures 10 and S8) is

consistent with previous reports from other Solanaceae (Fobes et al., 1985; O' Connell et al.,

2007; Kroumova et al., 2016) and is likely an adaptation to the seasonally arid native habitat of

N. benthamiana in northwestern Australia (Goodin et al., 2008; The Australasian Virtual

Herbarium, https://avh.ala.org.au). Relative to *asat1-1* and *asat1-2* mutants, the lower acylsugar

content of *asat2-1* and *asat2-2* mutants (Figure 4), resulted in more rapid water loss in detached

leaves (Figures 10A and S8A). However, despite the only partial decrease in the acylsugar

content of *asat1* mutants, the decreases in water content and increases in leaf temperature of

intact plants were similar to those of *asat2* mutants (Figures 10B,C and S8B,C).

Acylsugars with C_{7-12} chains have been shown to be the most toxic sugar esters for small

phloem-feeding Hemiptera such as aphids, Asian citrus psyllids, and whiteflies (Chortyk et al.,

1996; McKenzie and Puterka, 2004; Song et al., 2006). Synthetic acylsucroses with di-heptanoic

acid (C7), di-octanoic acid (C8), and di-nonanoic acid (C9) acyl groups showed the highest

mortality in bioassays with *M. persicae* and *B. tabaci* (Chortyk et al., 1996; McKenzie and

Puterka, 2004; Song et al., 2006). *Nicotiana gossei*, a tobacco species that produces mainly C7-

C8 acyl group acylsugars, has a high level of insect resistance relative to close relatives with

acylsugar profiles that are not dominated by those with C7-C12 acyl groups (Thurston, 1961;

Kroumova and Wagner, 2003). In *N. benthamiana*, the two most abundant acylsugars that we

found contain C7 and predominantly C8 acyl groups, which is consistent with previous findings

of mainly 5- and 6-methyl heptanoate (C8) in *N. benthamiana* (Kroumova and Wagner, 2003;

Slocombe et al., 2008) and *N. alata* (Moghe et al., 2017). The almost complete depletion of

acylsugars in our *asat2* mutants improved both hemipteran and lepidopteran performance,

suggesting that the identified C8-chain acyl group acylsugars are providing insect resistance for

N. benthamiana.

We cannot rule out the possibility secondary effects that might also influence insect

performance on acylsugar-depleted *N. benthamiana*. Specialized metabolites in other plants, for

instance glucosinolates in *Arabidopsis thaliana* (Clay et al., 2009) and benzoxazinoids in *Zea*

mays (Meihls et al., 2013), regulate callose deposition as a secondary defense response. It is not

known whether acylsugars contribute to the regulation of other defense responses in *N.*

benthamiana. The observation of numerous dead whiteflies on wildtype *N. benthamiana* plants

in choice assays (Figure 6G), despite the option of moving to presumably more desirable *asat2-1*

mutant plants in the same cage, suggests that the acylsugars stickiness also plays a role in plant

defense by immobilizing the insects. Both altered leaf turgor and leaf temperature (Figure

10B,C) could affect insect feeding behavior and growth rate, though the specific effects on the

six tested insect species cannot be determined without further research.

Nicotiana benthamiana has been used for virus transmission assays, RNA interference

and VIGS of *M. persicae* genes, and transient expression of genes from other species to

determine their effects on aphid growth and reproduction (Ramsey et al., 2007; Bos et al., 2010;

Pitino and Hogenhout, 2013; Casteel et al., 2014; Elzinga et al., 2014; Rodriguez et al., 2014;

Krenz et al., 2015; Tzin et al., 2015; Mulot et al., 2016; Mathers et al., 2017; Del Toro et al,

2018; Cui et al., 2019; Worrall et al., 2019). However, the poor growth of many *M. persicae*

isolates on *N. benthamiana* (Thurston, 1961; Hagimori et al., 1993; Figure S6) hampers

experiments of this kind Similarly, due to the very poor survival of *B. tabaci* on wildtype plants

(Simon et al., 2003; Figure 6E), *N. benthamiana* has not been useful as a host plant for whitefly

research.

To demonstrate the utility of *asat2* mutants as a tool for aphid research, we performed

TuMV-GFP transmission experiments using *M. persicae*. Our results show that *asat2-1* plants

significantly increased virus transmission (Figure 8). Interestingly, efficient virus transmission

only occurred when both the virus donor and the recipient plants had the *asat2-1* genotype. This

suggests that aphid behavior on *asat2-1* plants, perhaps faster or more continuous probing,

promotes both uptake and delivery of TuMV. The higher virus transmission on *asat2-1* mutant

N. benthamiana will facilitate use of this model system for future research on plant-aphid-virus

interactions and factors that promote virus transmission.

The low survival of whiteflies on wildtype *N. benthamiana* (Figures 6E and 9D) makes it

difficult to assess the negative effects of gene expression silencing by TRV VIGS. By contrast,

we were able to demonstrate reduced survival with all four tested whitefly genes with TRV

VIGS using *asat2-1* mutant plants (Figure 9C). *TLR7* and *TSCD* gene expression was reduced on

day 1 but not on day 7 in the VIGS experiment (Figure 10A,B), but nevertheless there was a

negative effect on whitefly survival over 7 days (Figure 10C). Two possible explanations for this

finding are: (i) There was a survivor bias in that we could only measure gene expression levels in

surviving aphids, and perhaps all aphids with efficient expression silencing of *TLR7* and *TSCD*

were dead after 7 days. (ii) There may be gene expression compensation at the whole-insect level

over time, but not in specific tissues that affect insect survival. Quantitative PCR of fractionated

whiteflies would be necessary to determine the time course of *TLR7* and *TSCD* expression

silencing and whether VIGS primarily affects gene expression in specific tissue types.

In addition to confirming the importance of two *B. tabaci* genes, *AchE* and *TLR7*, that

have been targeted by RNAi (Malik et al., 2016; Chen et al., 2015), we chose two previously

uncharacterized horizontally transferred genes *SQS* and *TSCD,* as VIGS targets. *Bemisia tabaci*

MEAM1 has at least 142 horizontally transferred genes from bacteria and fungi (Chen et al.,

2016). Given that horizontal transfer of functionally expressed microbial genes into insect

germlines is rare on an evolutionary timescale, there is likely a selective advantage to having

these genes expressed in whiteflies. This was confirmed by the observation that VIGS of both

SQS and *TSCD* reduced whitefly survival relative to control plants (Figure 9C,D). Transient

expression knockdown of *SQS*, *TSCD*, and other horizontally transferred genes will enable future

research to study the functions of these genes in whitefly metabolism. Due to their importance

for whitefly survival, as well their absence in beneficial insects such as ladybugs and lacewings,

horizontally transferred genes also are attractive targets for controlling whiteflies by RNA

interference. The established *N. benthamiana* VIGS system, which allows rapid cloning of

targets by Gateway recombination (Liu et al., 2002), will allow rapid screening of other

horizontally transferred genes to identify ones that would be most suitable for whitefly control on

crop plants by RNA interference.

Although *H. zea*, *H. virescens*, and *T. ni* larvae grow well on cultivated tobacco, neonate

larvae had a low survival rate on both wildtype *asat2-1 N. benthamiana* (Figure 7A-C). There

was a higher survival rate with five-day-old larvae of the three tested species, which all grew

significantly better on *asat2-1* mutants than on wildtype *N. benthamiana* (Figure 7D-F). Thus, *N.*

benthamiana acylsugars likely provide at least some protection against lepidopteran pests.

However, the high mortality of neonate larvae on *asat2-1* plants suggests that either residual

acylsugars or as yet unknown resistance mechanisms in *N. benthamiana* can provide protection.

Additional mutations that decrease insect resistance, perhaps regulatory genes such as *COI1* or

genes affecting the production of other specialized metabolites, will be necessary to facilitate *N.*

benthamiana experiments with *H. zea*, *H. virescens*, *T. ni*, and other commonly studied

lepidopteran species.

The *Cas9* transgene, which is still present in the *asat2-1* mutant line that we used for

most of our experiments, may facilitate further mutagenesis. In a recently described method

(Ellison et al., 2020) *Cas9*-transgenic *N. benthamiana* plants were infected with TRV carrying

gRNAs linked to phloem movement sequences. Seeds harvested from these plants had a high

frequency of homozygous knockout mutations in the CRISPR/Cas9-targeted genes. The ease of

generating germline mutations using this approach will make it possible to test the function of

other predicted insect defense genes by knocking out their expression in the *N. benthamiana*

asat2-1 mutant background using a TRV-expressed gRNA.

Our knockout of acylsugar biosynthesis is an important first step toward improving the

already excellent *N. benthamiana* model system (Goodin et al., 2008; Bally et al., 2018), making

it more suitable for studying plant interactions with *M. persicae*, *B. tabaci*, and other

agriculturally relevant insect pests. Such experiments can include transient expression assays to

test the function of insect elicitors and insect-defensive genes from other plant species in *N.*

benthamiana, as well as VIGS to down-regulate insect gene expression in a targeted manner.

Furthermore, the almost complete absence of acylsugars in the *asat2* mutant lines, in

combination with the facile *Agrobacterium-* and virus-mediated transient gene expression

systems available for *N. benthamiana*, will make these mutants a suitable platform for the

functional analysis of ASATs from other Solanaceae.

Materials and Methods

Insect and plant cultures

A red strain of *M. persicae* (Ramsey et al., 2007; Ramsey et al., 2014), originally collected from

N. tabacum by Stewart Gray (Robert W. Holley Center for Agriculture & Health, Ithaca, NY),

was maintained on *N. tabacum* plants in a growth room at 23°C with a 16:8 h light:dark

photoperiod. Insect bioassays were conducted in the same growth room. Colonies of *B. tabaci*

MEAM1 were provided by Danielle Preston and Angela Douglas (Cornell University) and Jane

Polston (University of Florida). *Macrosiphum euphorbiae* was obtained from Isgouhi Kaloshian

(UC Riverside) and was maintained on tomato cv. Moneymaker. Eggs of *H. zea*, *H. virescens*,

and *T. ni* were purchased from Benzon Research (www.benzonresearch.com). *Nicotiana*

benthamiana wild type and mutant plants for aphid experiments, caterpillar experiments, and

whitefly VIGS assays were maintained at 23°C and a 16:8 h light:dark photoperiod in a

Conviron (Winnipeg, Canada) growth chamber and, for seed production, in a greenhouse at

27/24°C (day/night) with ambient light conditions. *Nicotiana benthamiana* wild type and mutant

plants for whitefly choice and no-choice assays were maintained at 26°C and a 16:8 h light:dark

photoperiod in a growth room and, for seed production, in a growth chamber (Percival Scientific,

Perry, IA) at 27/24°C (day/night) with a 16:8 h light:dark photoperiod.

Identification of ASAT1 and ASAT2 orthologs in N. benthamiana

To identify ASAT1 and ASAT2 orthologs in *N. benthamiana*, protein sequences of *Salpiglossis*

sinuate and *Solanum lycopersicum* ASAT1 and ASAT2 (Moghe et al., 2017) were compared to

predicted proteins encoded by the *N. benthamiana* genome. Sequences with >67% identity were

selected as potential ASAT1 and ASAT2 candidates and nucleotide sequences were obtained

from the Solanaceae Genomics Network (www.solgenomics.net). The candidate ASAT

sequences also were confirmed by comparing them to the most recent published *N. benthamiana*

genome assembly (Schiavinato et al., 2019). To confirm the nucleotide sequences of *N.*

benthamiana ASAT1 and *ASAT2*, genes were amplified with ASAT1F/ASAT1R and

ASAT2F/ASAT2R primers (Table S4) using genomic DNA as the template. Amplified

fragments were cloned in pDONOR™207 (ThermoFisher Scientific, US) and were sequenced in

their entirety using Sanger sequencing. This confirmatory sequencing showed no differences

relative to the published *N. benthamiana* genome.

Phylogenetic analysis of N. benthamiana *ASATs*

A protein phylogenetic tree of previously annotated Solanaceae ASATs (Figures 2, S1, S2;

Tables S1, S2) was constructed using maximum likelihood method. Briefly, the ASATs protein

sequences were aligned in program ClustalW (Thompson et al., 1994). Then the alignment was

improved by removing the spurious sequences and poorly aligned regions (gap threshold at 0.25)

using the program TrimAL v 1.4.rev22 (Capella-Gutierrez et al., 2009). Finally, an unrooted

maximum likelihood tree was generated using the improved alignment with a bootstrap of 1000

in RAxML v8.2.12 (Stamatakis, 2014). The tree was visualized and presented using FigTree

v1.4.4 (http://tree.bio.ed.ac.uk).

sgRNA design and plasmid cloning

Single-guide RNAs (sgRNA) targeting *ASAT1* and *ASAT2* were designed based on the coding

regions using online software, CRISPR-P v2.0 (Liu et al., 2017) and CRISPRdirect

(https://crispr.dbcls.jp/), based on two parameters, cleavage efficiency and potential off-targets.

Additionally, only sgRNAs with >40% GC content were selected. Three Cas9/gRNA constructs

each were constructed for *ASAT1* and *ASAT2* following a previously developed CRISPR/Cas9

system (Jacobs et al., 2015). Four segments of DNA were prepared with 20 bp overlaps on their

ends: ssDNA gRNA oligo, linearized p201N:Cas9 plasmid (Addgene 59175-59178), the

Medicago truncatula (Mt) U6 promoter (377 bp), and a scaffold DNA (106 bp). For ssDNA

gRNA, oligonucleotides targeting either the sense or antisense sequence of target genes were

designed as: sense oligo TCAAGCGAACCAGTAGGCTT--GN$_{19}$--

GTTTTAGAGCTAGAAATAGC, and antisense oligo GCTATTTCTAGCTCTAAAAC--

N$_{19}$C—AAGCCTACTGGTTCGCTTGA. The gRNA oligonucleotide sequences are shown in

Figure 3 and Table S4.

Oligonucleotide sequences were synthesized by Integrated DNA Technologies

(www.idtdna.com). One μl of each 100 μM oligo was added to 500 μl 1x NEB buffer 2 (New

England Biolabs, www.neb.com). The p201N:Cas9 plasmid was linearized by digestion with

*Spe*1 (www.neb.com) in 1x buffer 4 at 37°C for 2 h, followed by column purification and a

second digestion with *Swa*l in 1 x buffer 3.1 at 25°C for 2 h. Complete plasmid digestion was

confirmed on a 0.8% agarose gel. The MtU6 promoter and Scaffold DNAs were PCR-amplified

from the pUC gRNA Shuttle plasmid (Jacobs et al., 2015) using the primers

*Swa*l_MtU6F/MtU6R and ScaffoldF/Spe_ScaffoldR, respectively (Table S4). The PCR reactions

were performed with a high-fidelity polymerase (2x Kapa master mix; www.sigmaaldrich.com)

using the program: 95°C for 3 min followed by 31 cycles of 98°C for 20 sec, 60°C for 30 sec,

72°C for 30 sec, and a final extension of 72°C for 5 min. Finally, cloning was done using the

NEBuilder® HiFi DNA Assembly Cloning Kit. For each reaction, the four pieces of DNA were

mixed in a 20 µl reaction with the NEBbuilder assembly mix with a final concentration of 0.011

pmol (~100 ng) of p201N:Cas9 plasmid, 0.2 pmol of MtU6 amplicon (~ 50 ng), scaffold

amplicon (~ 12 ng) and ssDNA gRNA oligo (60-mer, 1 µl). The reactions were placed in thermal

cycler at 50°C for 1 h.

Two µl of the cloning reaction were transformed into 50 µl of the One Shot™ Top10

chemically competent cells (Invitrogen, www.thermofisher.com) and plated on LB (Bertani,

1951) agar medium with 50 µl/ml kanamycin for selection of transformants. Colonies with the

correct inserts were screened using the Ubi3p218R and IScelR primers (Table S4). Plasmids

carrying the designed gRNA constructs were then transformed into *Agrobacterium tumefaciens*

strain GV3101 for generating transgenic plants. All constructs were confirmed by Sanger

sequencing.

To avoid off-target effects, gRNAs were further checked by comparison against the

reference *N. benthamiana* genome v1.0.1 (www.solgenomics.net). Only two sites in the *N.*

benthamiana genome were found to have non-target matches >17 nt (both with 1 internal

mismatch), and with the NGG PAM sequence on the correct strand. These two sites were

checked by PCR amplification and Sanger sequencing and showed no unexpected editing in our

asat1-1, *asat1-2*, *asat2-1*, or *asat2-2* mutant plants. Primers used for off-target Sanger

sequencing are listed in Table S4.

Confirmation of gRNA functionality by transient expression

For both *ASAT1* and *ASAT2*, we *Agrobacterium*-infiltrated *N. benthamiana* plants at the four-leaf

stage. For *ASAT1*, we used *Agrobacterium* carrying the Cas9/gRNA constructs at OD of 1 and 3,

whereas for *ASAT2*, we used an OD 1.5. Each leaf was saturated with *Agrobacterium* solution.

After infiltration, the plants were cultured in a growth chamber for 2 days and then the infiltrated

whole leaves were collected for genomic DNA extraction and tested by PCR amplification for

detection of insertion/deletion polymorphisms in the target region (Figure S9). During the

method optimization, a positive control construct targeting the *N. benthamiana Drm3* gene

(gRNA: GCCACTATCTGGCCGGGGAC, provided by the Greg Martin lab, Boyce Thompson

Institute) was infiltrated in parallel.

Stable mutagenesis of ASATs using tissue culture

Stable *ASAT* mutant *N. benthamiana* plants were created in the Boyce Thompson Institute plant

transformation facility using CRISPR/Cas9 with gRNAs that had been confirmed to be

functional as described above and a previously described protocol (Van Eck et al., 2019), with

minor modifications. To prepare plants for transformation, we disinfected *N. benthamiana* seeds

with 1.5 ml 1.25% sodium hypochlorite (1:5 dilution of 5.25% sodium hypochlorite Clorox

bleach), with 100 µl Tween-20 added, for 20 min on a shaker platform. After sodium

hypochlorite treatment, seeds were rinsed three times with sterile water. Subsequently, we placed

seeds on Tobacco Seed Germination Medium (composition per liter: 1.08 g Murashige and

Skoog salts (Murashige and Skoog, 1962), 30 g sucrose, and 8 g agar, pH = 5.7±0.1) and

incubated the seeds under yellow filtered light with a 16-h photoperiod at 27°C. After ~2 weeks,

the seedlings were transferred to the Rooting Medium (composition per liter: 2.15 g Murashige

and Skoog salts, 30 g sucrose, 1 ml B5 vitamin/amino acid stock, and 8 g agar, pH = 5.7±0.1).

After ~6 weeks, the plants were ready for *Agrobacterium* infection. First, *Agrobacterium*

carrying the gRNA plasmids was amplified in YEP medium (composition per liter: 10 g yeast

extract, 10 g peptone, and 5 g NaCl with antibiotics, 50 µg/µl gentamycin, 50 µg/µl kanamycin,

and 50 µg/µl rifampicin), and *Agrobacterium* culture then was pelleted and combined to a certain

OD (OD 1.0 for *ASAT1* gRNAs, and OD 1.0 or 1.5 for *ASAT2* gRNAs) with filtered cell

suspension buffer (10 mM MES, 10 mM $MgCl_2$ and 200 µM acetosyringone). Then, fully

expanded but immature *N. benthamiana* leaves were dissected into 5 mm segments by removing

the leaf margins. The leaf segments were incubated in the *Agrobacterium* culture for 30 min with

shaking. The incubated leaf segments were transferred to co-cultivation medium plates

(composition per liter: 4.3 g Murashige and Skoog salts, 30 g sucrose, 1 ml B5 vitamin/amino

acid stock, and 5 g agar, pH = 5.7±0.1) and kept in the dark at room temperature. Leaf segments

treated with YEP medium were used as a negative control. After three days, the leaf segments

were transferred from co-cultivation medium to shoot bud initiation medium with selections

(composition per liter: 4.3 g Murashige and Skoog salts, 30 g sucrose, 1 ml B5 vitamin/amino

acid stock, 0.1 mg NAA, 1.0 mg BA, and 6 g agar, pH=5.7±0.1; with 200 µg/ml kanamycin and

250 µg/ml Timentin). Medium with Timentin but without kanamycin was used as a positive

control. Leaf segments on selection medium were incubated under a 16:8 h light-dark

photoperiod at 24-25°C under yellow filtered light. The leaf segments were sub-cultured every 2-

weeks onto the same medium until shoot buds were visible. Finally, to generate roots, shoot buds

were transferred to rooting medium (composition per liter: 2.15 g Murashige and Skoog salts, 30

g sucrose, 1 ml B5 vitamin/amino acid stock, 8 g agar with 200 μg/ml kanamycin and 250 μg/ml

Timentin, pH=5.7±0.1).

Confirmation of homozygous mutant plants in the T2 generation

Rooted *N. benthamiana* plants from tissue culture were transferred to soil (T0 generation).

CRISPR/Cas9-induced mutations were identified by PCR amplification of genomic regions of

the gRNA target sites in *ASAT1* and *ASAT2* (Figure 3), followed by Sanger sequencing. Lines

with mutations were used to generate T1 plants, which were subjected to PCR amplification and

sequencing to confirm homozygous mutations. T2 seeds from confirmed homozygous mutant

asat1-1, *asat1-2*, *asat2-1*, and *asat2-2* T1 plants were used for all experiments. Homozygous

mutations were confirmed in randomly selected T2 plants by PCR amplification and Sanger

sequencing. The presence of Cas9 in transgenic plants in the T0, T1, and T2 generations was

confirmed by PCR amplification and agarose gel electrophoresis. Primers that were used to

identify mutations and confirm the presence of Cas9 are listed in Table S4.

Acylsugar measurements by LC/MS

Liquid chromatography/mass spectrometry (LC/MS) was used to confirm the effect of the *ASAT*

mutations by measuring acylsugar content in leaf extracts from wildtype and *ASAT* mutant

plants. New leaflets were rinsed in acylsugar extraction solution (3:3:2

acetonitrile:isopropanol:water, 0.1% formic acid, and 1 μM Telmisartan as internal standard) and

gently agitated for 2 min. Then, the extraction solutions were transferred to LC/MS glass vials,

and the leaves were air dried for leaf weight measurements.

Chromatography of leaf surface washes was performed on a ThermoScientific Ultimate

3000 HPLC with a glass vial autosampler and coupled with a Thermo Scientific Q Exactive™

Hybrid Quadrupole-Orbitrap™ Mass Spectrometer (Mass Spectrometry Facility at Boyce

Thompson Institute). Acylsugar extracts were separated on an Ascentis Express C18 HPLC

column (10 cm × 2.1 mm × 2.7 μm) (Sigma☐Aldrich, St. Louis, MO) with a flow rate of 0.3

ml/min, using a gradient flow of 0.1% formic acid (Solvent A) and 100% acetonitrile (Solvent

B). We used a 7-min LC method for metabolite profiling, which involved a linear gradient from

95:5 A:B to 0:98 A:B. Full☐scan mass spectra were collected (mass range: m/z 50–1000) in both

positive and negative electron spray ionization (ESI) modes. Mass spectral parameters were set

as follows: capillary spray voltage 2.00 kV for negative ion-mode and 3.00 kV for positive ion-

mode, source temperature: 100°C, desolvation temperature 350°C, desolvation nitrogen gas flow

rate: 600 liters/h, cone voltage 35 V. Acylsugars were identified and annotated using Thermo

Xcalibur Qual Browser (Thermo Fisher) and MS-DIAL v4.20 based on the MS/MS peak features

and neutral losses. The acylsugar abundances were estimated using peak areas at the respective

m/z channel under negative ESI mode. Acylsugar quantification was first normalized to the

internal control Telmisartan to account for technical variation between samples, and then

normalized to the leaf dry weight to allow comparisons between samples.

Insect choice and no-choice bioassays

To measure *M. persicae* and *M. euphorbiae* growth, we caged aphids on individual leaves of

mutant and wildtype 4 to 5-week-old *N. benthamiana* (Figure S10A, B). Twenty adult *M.*

persicae from *N. tabacum* (naïve to *N. benthamiana*) were placed in each cage and allowed to

generate nymphs for ~12 hrs. Twenty-five nymphs were left in each cage and were monitored for

5 d to assess nymph survival. At the end of the *M. persicae* survival monitoring period, five *M.*

persicae were left in each cage and reproduction was monitored for one week. Finally, the

remaining *M. persicae* were collected to measure aphid size by taking a picture and assessing the

area of each aphid using ImageJ (Schneider et al., 2012). Ten adult *M. euphorbiae* from a colony

on tomato cv. Moneymaker were placed in each individual cage on *N. benthamiana* leaves.

Surviving aphids and progeny were counted after 24 hours.

Myzus persicae and *M. euphorbiae* choice assays were performed with detached leaves

from 4 to 5-week-old *N. benthamiana*. Two similarly-sized leaves from individual *ASAT* mutant

and wildtype plants were cut and placed in 15-cm Petri dishes, with their petioles inserted in

moistened cotton swabs (pairwise comparison is shown in Figure S10C). Ten naïve adult aphids,

which had not previously encountered *N. benthamiana*, were released at the midpoint between

pairs of leaves (wildtype, *asat1*, or *asat2*), and the Petri dishes were placed under 16:8 h

light:dark photoperiod. The aphids on each leaf were counted at 24 h after their release in the

Petri dishes.

To measure whitefly survival and fecundity on wildtype and *asat2-1 N. benthamiana*

plants, cages were set up with *N. benthamiana* plants at the seven-leaf stage (approximately 3

weeks old). Each cage contained three plants, either wildtype or *asat2-1*. Ninety adult whiteflies

reared on *Brassica oleracea* (variety Earliana; www.burpee.com, catalog number 62729A) were

introduced into each cage (60 x 60 x 60 cm) with *N. benthamiana* (30 whiteflies/plant) and were

allowed to feed for three days at 26°C with a 16:8 h light:dark photoperiod. The numbers of

whiteflies surviving on each host plant were counted, after which the remaining insects were

killed with insecticidal soap. The following day, the number of whitefly eggs on each plant was

counted. This experiment was conducted twice with similar results.

For whitefly choice assays, wildtype and *asat2-1* plants at the seven-leaf stage were

placed together in the same cage. Approximately 150 whiteflies from cabbage plants were

moved into each cage. After 24 h at 26°C with a 16:8 h light:dark photoperiod, live and dead

whiteflies were counted on the plants and elsewhere in the cage. This experiment was repeated

three times.

Eggs of *H. zea, H. virescens* and *T. ni* were hatched on artificial diet (Southland Products,

Lake Village, Arkansas). Neonate larvae were confined onto individual *N. benthamiana* leaves,

one larva per plant, using 10 x 15 cm organza mesh bags (www.amazon.com, item

B073J4RS9C). After ten days, the surviving larvae were counted and weighed. In a separate

experiment, *H. zea, H. virescens*, and *T. ni* were reared on artificial diet (beet armyworm diet,

www.southlandproducts.net) for five days. Individual five-day-old caterpillars were weighed and

then confined on 4 to 4.5-week-old *N benthamiana* plants using 30 cm x 60 cm micro-perforated

bread bags (www.amazon.com). After seven days, the surviving larvae were weighed again.

Relative growth rate was calculated as: $\ln(((\text{day-12 mass})/(\text{mean day-5 mass}))/7)$.

Leaf water loss and temperature assays

To measure the leaf water loss, two leaves from each of eight plants were detached. The fresh

weight of each leaf was determined on a Sartorius Ultra Micro Balance. All leaves were placed

at 23°C and a 16:8 h light:dark photoperiod. Each leaf was weighted again after 24 h and the

percentage of water loss was calculated as [(fresh weight - final weight)/fresh weight]*100%.

Thermal images were acquired in the growth chamber environment using a thermal

camera (A655sc, FLIR Systems Inc., Boston, MA, USA) with a spectral range of 7.5–14.0 mm

and a resolution of 640 x 480 pixels. The camera was placed approximately 1 m away from each

plant and a white background was used when the plant images were acquired. One region of

interest, corresponding to the perimeter of each leaf, was specified per leaf for each of 20 leaves

per genotype. Using the FLIR ResearchIR Max software (version 4.40.9.30) thermal images files

were exported as CSV files. Images were segmented from the background using Gaussian

mixture models in MATLAB to determine the temperature of each leaf. After segmentation, the

temperature was averaged across the segmented leaf.

Hyperspectral images were acquired in a dark room using a hyperspectral imager

(SOC710, Series 70-V, Surface Optics Corporation, San Diego, CA, USA) that covered a

spectral range from 400 to 1000 nm for 128 wavebands. Image acquisition and recording were

performed using a Dell DELL XPS 15 9570 laptop computer that controls the camera. The

camera was fixed using a stand with the lens facing the plants and capturing top view images

approximately 1 m away. A Spectralon tile (Labsphere, Inc, North Sutton NH, USA) was placed

next to the plant trays, covering one corner of the image to facilitate subsequent image

processing and calibration. The nominal reflectance value for the Spectralon tile was 99% and it

had a 30.5 cm x 30.5 cm reflective area. Lighting consisted of two halogen lamps placed at ~ 45°

angles on either side of the camera to create an even light distribution. All image analysis was

performed in HSIviewer, a MATLAB package (Stone et al., 2020). White reflectance calibration

was performed using the Spectralon tile. One region of interest ROI was specified for each of 20

leaves per *N. benthamiana* genotype. This ROI corresponded to the perimeter of each leaf. From

each hyperspectral cube image, the vegetation pixels (green portion of the plant) were extracted

using the Normalized Difference Vegetation Index (NDVI). Mean reflectance (R) was calculated

per band per 10 leaves in order to obtained the WBI results

To calculate NDVI and WBI we used the following formulas, where *R* corresponds to the

reflectance at a specific wavelength (nm): WBI = (R970/R900) (Penuelas et al., 1993) and NVDI

= (R750 - R705)/(R750 + R705) (Gitelson and Merzlyak, 1994).

Virus transmission assays

For virus transmission assays, young adult *M. persicae* were starved for 3 h and then allowed to

feed for 30 min on 6-week-old wildtype or *asat2-1 N. benthamiana* plants infected with TuMV-

GFP (Lellis et al., 2002; Casteel et al., 2015). Aphids were transferred in groups of ten to

individual 4-week-old wildtype or *asat2-1* recipient plants. The aphids were allowed to feed on

recipient plants for 24 h and were then removed. Donor and recipient plants were kept in the

Conviron growth chamber at 23°C and a 16:8 h light:dark photoperiod. The development of GFP

fluorescence was monitored every 24 h using a 395 nm UV flashlight (www.walmart.com) in the

dark, until no new GFP signals developed from one day to the next. In each experiment, the virus

donor plant group consisted of eight TuMV-GFP infected plants (2 weeks after TuMV-GFP

infection by rub-inoculation) of each genotype, with aphids randomly spread on the plants. Each

recipient group consisted of 20 plants of each genotype, with the aphids confined on individual

plants. The experiment was repeated three times with similar results, and the results were

combined for statistical analysis of differences between groups using one-way ANOVA with a

block effect followed by a Tukey HSD post hoc test.

Plant-mediated virus induced gene silencing in whitefly with WT and asat2-1

For each target gene, a 200-550 bp RNA sequence was designed using the E-RNA*i* webserver

(https://www.dkfz.de/signaling/e-rnai3/). The designed fragments were checked against the *B.*

tabaci MEAM1 (Chen et al., 2016) and *N. benthamiana* (Bombarely et al., 2012) reference

genomes using local blast (Mount, 2007) to avoid potential off-target sequences with full

matches $\geq$ 19 nt. The identified gene fragments were then PCR-amplified from whitefly cDNA

and cloned into a binary tobacco rattle virus VIGS vector pTRV2 (Liu et al., 2002) using the

Invitrogen Gateway recombination cloning technology (Invitrogen, USA). As a positive control,

we used a pTRV2-PDS construct with an RNA fragment targeting *N. benthamiana* phytoene

desaturase (Velásquez et al., 2009). As negative controls, we used pTRV2-EV (empty vector)

and pTRV2-GFP, with an RNA fragment targeting *Aequorea victoria GFP*, a sequence that is

not present in either *N. benthamiana* or *B. tabaci*. The PCR primers used to generate the VIGS

vectors are in Supplemental Table S4.

To infiltrate plants, plasmids (pTRV1 and pTRV2 containing constructs targeting the

genes of interest) were transformed into *Agrobacterium tumefaciens* GV3101. Equal amounts of

Agrobacterium with TRV1 and TRV2 were resuspended in infiltration medium (10 mM MES,

10 mM $MgCl_2$, 200 μM acetosyringone) to an OD 0.3 and incubated at room temperature for 3 h.

The *Agrobacterium* mixture was infiltrated using a 1-ml needleless syringe to saturate three

leaves of ~4-week-old *N. benthamiana*. Inoculated plants were kept in a Conviron growth

chamber for 2~3 weeks until control plants infiltrated with pTRV2-PDS showed photobleaching

symptoms. Successful growth of TRV targeting the genes of interest in whiteflies was confirmed

by PCR and the confirmed plants were then used for insect bioassays.

For whitefly bioassays, three infiltrated plants were used for each gene of interest, and

three cages were attached to each plant. Ten young adult whiteflies (emerged within the past

week) were placed in each cage. After seven days, the survival rates of whiteflies were assessed

and compared across different treatments. The surviving whiteflies were collected for qPCR

analyses of target gene expression using quantitative PCR using the PowerUp^{TM} SYBR^{TM} green

Master Mix (AppliedBiosystem, Thermo Fisher Scientific) on a QuantStudio^{TM} 6 Flex Real-time

PCR system (AppliedBiosystem, Thermo Fisher Scientific), and results were analyzed using the

$2^{-\Delta\Delta CT}$ method (Schmittgen and Livak, 2008). The whitefly VIGS bioassays, comparing *asat2-1*

and wildtype *N. benthamiana*, were repeated three times with similar results. Two replicates of a

separate experiment was set up on *asat2-1* plants to collect whiteflies after one day for gene

expression analyses.

Statistical analysis

All statistical comparisons were conducted using SPSS v25, R and MATLAB R2019a

(MathWorks, Inc., Natick, MA, USA). ANOVA followed by a Dunnett's post hoc test was used

to determine differences in leaf water loss, leaf temperature, and WBI across genotypes in each

data set. ANOVA followed by a Duncan post hoc test was used for aphid bioassay and LC/MS

results. A Chi-square test was used to test for differences in pairwise aphid choice assays,

whitefly, and lepidopteran assays.

Acknowledgements

We want to thank Patricia Keen and Joyce Van Eck for their help with *N. benthamiana* stable

transformation and access to the tissue culture facility, Danielle Preston, Angela Douglas, and

Jane Polston for providing whitefly colonies, Isgouhi Kaloshian for the potato aphid colony,

Ning Zhang and Greg Martin for sharing the p201N-Cas9 construct and the *Drm3* sgRNA

control construct, and William Stone and Thomas Lawton for providing custom image

processing software. This research was supported by Cornell startup funds to G.D.M., Deutsche

Forschungsgemeinschaft award #411255989 to L.H.K., and United States Department of

Agriculture Biotechnology Risk Assessment Grant 2017-33522-27006, US National Science

Foundation award IOS-1645256, and Defense Advanced Research Projects Agency (DARPA)

agreement HR0011-17-2-0053 to G.J, and US National Science Foundation award #1723926 to

Conflict of interest

The authors declare that there is no conflict of interest.

Author Contributions

G.J. and H.F. conceived the original research plans; H.F., S.S., L.A., H.X., L.K., J.D.T, S.H.C.,

and A.N.F. performed the experiments; H.F., L.A., L.K., and G.D.M. analyzed the data; C.L.C.,

M.A.G., G.D.M., G.S., and G.J. supervised the experiments; H.F. and G.J. wrote the article with

contributions from all of the authors; G.J. agrees to serve as the contact author responsible for

communication and distribution of samples.

References

Alba JM, Montserrat M, Fernandez-Munoz R (2009) Resistance to the two-spotted spider

mite (*Tetranychus urticae*) by acylsucroses of wild tomato (*Solanum pimpinellifolium*)

trichomes studied in a recombinant inbred line population. Exp Appl Acarol 47: 35-47

Arntzen C (2015) Plant-made pharmaceuticals: from 'Edible Vaccines' to Ebola therapeutics.

Plant Biotechnology Journal 13: 1013-1016

Arrendale RF, Severson RF, Sisson VA, Costello CE, Leary JA, Himmelsbach DS,

Vanhalbeek H (1990) Characterization of the sucrose ester fraction from *Nicotiana*

glutinosa. Journal of Agricultural and Food Chemistry 38: 75-85

Bally J, Jung H, Mortimer C, Naim F, Philips JG, Hellens R, Bombarely A, Goodin MM,

Waterhouse PM (2018) The rise and rise of *Nicotiana benthamiana*: a plant for all

reasons. Annual Review of Phytopathology, Vol 56 56: 405-426

Bertani G (1951) Studies on lysogenesis. I. The mode of phage liberation by lysogenic

Escherichia coli. J Bacteriol 62: 293-300

Bombarely A, Rosli HG, Vrebalov J, Moffett P, Mueller LA, Martin GB (2012) A draft
genome sequence of *Nicotiana benthamiana* to enhance molecular plant-microbe biology
research. Mol Plant-Micr Int 25: 1523-1530

Bos JIB, Prince D, Pitino M, Maffei ME, Win J, Hogenhout SA (2010) A functional
genomics approach identifies candidate effectors from the aphid species *Myzus persicae*
(green peach aphid). Plos Genetics 6: e1001216.

Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T (2009) trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25: 1972-1973

Cardoso MZ (2008) Herbivore handling of a plant's trichome: The case of *Heliconius*
charithonia (L.) (Lepidoptera : Nymphalidae) and *Passiflora Lobata* (Killip) Hutch.
(Passifloraceae). Neotropical Entomology 37: 247-252

Casteel CL, De Alwis M, Bak A, Dong H, Whitham SA, Jander G (2015) Disruption of
ethylene responses by turnip mosaic virus mediates suppression of plant defense against
the green peach aphid vector. Plant Physiol 169: 209-218

Casteel CL, Yang CL, Nanduri AC, De Jong HN, Whitham SA, Jander G (2014) The NIa-
Pro protein of *Turnip mosaic virus* improves growth and reproduction of the aphid vector,
Myzus persicae (green peach aphid). Plant Journal 77: 653-663

Chen W, Hasegawa DK, Kaur N, Kliot A, Pinheiro PV, Luan J, Stensmyr MC, Zheng Y,
Liu W, Sun H, Xu Y, Luo Y, Kruse A, Yang X, Kontsedalov S, Lebedev G, Fisher
TW, Nelson DR, Hunter WB, Brown JK, Jander G, Cilia M, Douglas AE, Ghanim
M, Simmons AM, Wintermantel WM, Ling KS, Fei Z (2016) The draft genome of
whitefly *Bemisia tabaci* MEAM1, a global crop pest, provides novel insights into virus
transmission, host adaptation, and insecticide resistance. BMC Biol 14: 110

Chen X, Li L, Hu Q, Zhang B, Wu W, Jin F, Jiang J (2015) Expression of dsRNA in
recombinant *Isaria fumosorosea* strain targets the TLR7 gene in *Bemisia tabaci*. BMC
Biotechnol 15: 64

Chortyk OT, Pomonis JG, Johnson AW (1996) Syntheses and characterizations of insecticidal
sucrose esters. J Ag Food Chem 44: 1551-1557

Clay N, Adio A, Denoux C, Jander G, Ausubel F (2009) Glucosinolate metabolites required
for an Arabidopsis innate immune response. Science 323: 95-101

Cui N, Lu H, Wang TZ, Zhang WH, Kang L, Cui F (2019) Armet, an aphid effector protein,

induces pathogen resistance in plants by promoting the accumulation of salicylic acid.

Philosophical Transactions of the Royal Society B-Biological Sciences 374

D'Auria JC (2006) Acyltransferases in plants: a good time to be BAHD. Curr Opin Plant Biol 9:

331-340

Del Toro FJ, Mencia E, Aguilar E, Tenllado F, Canto T (2018) HCPro-mediated transmission

by aphids of purified virions does not require its silencing suppression function and

correlates with its ability to coat cell microtubules in loss-of-function mutant studies.

Virology 525: 10-18

Egan AN, Moore S, Stellari GM, Kang BC, Jahn MM (2019) Tandem gene duplication and

recombination at the AT3 locus in the *Solanaceae*, a gene essential for capsaicinoid

biosynthesis in *Capsicum*. Plos One 14: e0210510

Egea I, Albaladejo I, Meco V, Morales B, Sevilla A, Bolarin MC, Flores FB (2018) The

drought-tolerant *Solanum pennellii* regulates leaf water loss and induces genes involved

in amino acid and ethylene/jasmonate metabolism under dehydration. Sci Rep 8: 2791

Ellison EE, Nagalakshmi U, Gamo ME, Huang PJ, Dinesh-Kumar S, Voytas DF (2020)

Multiplexed heritable gene editing using RNA viruses and mobile single guide RNAs.

Nat Plants 6: 620-624

Elzinga DA, De Vos M, Jander G (2014) Suppression of plant defenses by a *Myzus persicae*

(green peach aphid) salivary effector protein. Mol Plant Microbe Interact 27: 747-756

Fan P, Miller AM, Schilmiller AL, Liu X, Ofner I, Jones AD, Zamir D, Last RL (2016) In

vitro reconstruction and analysis of evolutionary variation of the tomato acylsucrose

metabolic network. Proc Natl Acad Sci U S A 113: E239-248

Fobes JF, Mudd JB, Marsden MP (1985) Epicuticular lipid accumulation on the leaves of

Lycopersicon pennellii (Corr.) D'Arcy and *Lycopersicon esculentum* Mill. Plant Physiol

77: 567-570

Gaquerel E, Kotkar H, Onkokesung N, Galis I, Baldwin IT (2013) Silencing an N-

acyltransferase-like involved in lignin biosynthesis in *Nicotiana attenuata* dramatically

alters herbivory-induced phenolamide metabolism. Plos One 8: e62336

Gitelson A, Merzlyak MN (1994) Spectral reflectance changes associated with autumn
senescence of *Aesculus hippocastanum* L and *Acer platanoides* L leaves - spectral
features and relation to chlorophyll estimation. J Plant Physiol 143: 286-292

Glas JJ, Schimmel BCJ, Alba JM, Escobar-Bravo R, Schuurink RC, Kant MR (2012) Plant
glandular trichomes as targets for breeding or engineering of resistance to herbivores.
International Journal of Molecular Sciences 13: 17077-17103

Gong P, Zhang J, Li H, Yang C, Zhang C, Zhang X, Khurram Z, Zhang Y, Wang T, Fei Z,
Ye Z (2010) Transcriptional profiles of drought-responsive genes in modulating
transcription signal transduction, and biochemical pathways in tomato. J Exp Bot 61:
3563-3575

Goodin MM, Zaitlin D, Naidu RA, Lommel SA (2008) *Nicotiana benthamiana*: its history and
future as a model for plant-pathogen interactions. Mol Plant Microbe Interact 21: 1015-
1026

Hagimori M, Matsui M, Matsuzaki T, Shinozaki Y, Shinoda T, Harada H (1993) Production
of somatic hybrids between *Nicotiana benthamiana* and *Nicotiana tabacum* and their
resistance to aphids. Plant Science 91: 213-222

Hayward A, Padmanabhan M, Dinesh-Kumar SP (2011) Virus-induced gene silencing in
Nicotiana benthamiana and other plant species. Plant Reverse Genetics: Methods and
Protocols 678: 55-63

Jacobs TB, LaFayette PR, Schmitz RJ, Parrott WA (2015) Targeted genome modifications in
soybean with CRISPR/Cas9. Bmc Biotechnology 15: 16

Kandra L, Wagner GJ (1988) Studies of the site and mode of biosynthesis of tobacco trichome
exudate components. Archives of Biochemistry and Biophysics 265: 425-432

Karabourniotis G, Kotsabassidis D, Manetas Y (1995) Trichome density and its protective
potential against ultraviolet-B radiation damage during leaf development. Canadian
Journal of Botany-Revue Canadienne De Botanique 73: 376-383

Kim J, Kang K, Gonzales-Vigil E, Shi F, Jones AD, Barry CS, Last RL (2012) Striking
natural diversity in glandular trichome acylsugar composition is shaped by variation at
the acyltransferase2 locus in the wild tomato *Solanum habrochaites*. Plant Physiol 160:
1854-1870

Krenz B, Bronikowski A, Lu XY, Ziebell H, Thompson JR, Perry KL (2015) Visual
monitoring of *Cucumber mosaic virus* infection in *Nicotiana benthamiana* following
transmission by the aphid vector *Myzus persicae*. Journal of General Virology 96: 2904-
2912

Kroumova AB, Wagner GJ (2003) Different elongation pathways in the biosynthesis of acyl
groups of trichome exudate sugar esters from various solanaceous plants. Planta 216:
1013-1021

Kroumova ABM, Zaitlin D, Wagner GJ (2016) Natural variability in acyl moieties of sugar
esters produced by certain tobacco and other *Solanaceae* species. Phytochemistry 130:
218-227

Laue G, Preston CA, Baldwin IT (2000) Fast track to the trichome: induction of N-acyl
nornicotines precedes nicotine induction in *Nicotiana repanda*. Planta 210: 510-514

Lellis AD, Kasschau KD, Whitham SA, Carrington JC (2002) Loss-of-susceptibility mutants
of *Arabidopsis thaliana* reveal an essential role for eIF(iso)4E during potyvirus infection.
Curr Biol 12: 1046-1051

Leong BJ, Lybrand DB, Lou YR, Fan PX, Schilmiller AL, Last RL (2019) Evolution of
metabolic novelty: A trichome-expressed invertase creates specialized metabolic
diversity in wild tomato. Science Advances 5: eaaw3754

Liu H, Ding YD, Zhou YQ, Jin WQ, Xie KB, Chen LL (2017) CRISPR-P 2.0: an improved
CRISPR-Cas9 tool for genome editing in plants. Molecular Plant 10: 530-532

Liu Y, Schiff M, Dinesh-Kumar SP (2002) Virus-induced gene silencing in tomato. Plant J 31:
777-786

Malik HJ, Raza A, Amin I, Scheffler JA, Scheffler BE, Brown JK, Mansoor S (2016) RNAi-
mediated mortality of the whitefly through transgenic expression of double-stranded
RNA homologous to acetylcholinesterase and ecdysone receptor in tobacco plants. Sci
Rep 6: 38469

Mathers TC, Chen Y, Kaithakottil G, Legeai F, Mugford ST, Baa-Puyoulet P, Bretaudeau
A, Clavijo B, Colella S, Collin O, Dalmay T, Derrien T, Feng H, Gabaldon T,
Jordan A, Julca I, Kettles GJ, Kowitwanich K, Lavenier D, Lenzi P, Lopez-
Gomollon S, Loska D, Mapleson D, Maumus F, Moxon S, Price DR, Sugio A, van
Munster M, Uzest M, Waite D, Jander G, Tagu D, Wilson ACC, van Oosterhout C,

Swarbreck D, Hogenhout SA (2017) Rapid transcriptional plasticity of duplicated gene
clusters enables a clonally reproducing aphid to colonise diverse plant species. Genome
Biology 18: 27

Matsuzaki T, Shinozaki Y, Hagimori M, Tobita T, Shigematsu H, Koiwai A (1992) Novel
glycerolipids and glycolipids from the surface-lipids of *Nicotiana benthamiana*.
Bioscience Biotechnology and Biochemistry 56: 1565-1569

Matsuzaki T, Shinozaki Y, Suhara S, Ninomiya M, Shigematsu H, Koiwai A (1989)
Isolation of glycolipids from the surface-lipids of *Nicotiana bigelovii* and their
distribution in *Nicotiana* species. Agricultural and Biological Chemistry 53: 3079-3082

McKenzie CL, Puterka GJ (2004) Effect of sucrose octanoate on survival of nymphal and adult
Diaphorina citri (Homoptera : Psyllidae). J Econ Entomol 97: 970-975

McKenzie CL, Weathersbee AA, 3rd, Puterka GJ (2005) Toxicity of sucrose octanoate to
egg, nymphal, and adult *Bemisia tabaci* (Hemiptera: Aleyrodidae) using a novel plant-
based bioassay. J Econ Entomol 98: 1242-1247

Meihls LN, Handrick V, Glauser G, Barbier H, Kaur H, Haribal MM, Lipka AE,
Gershenzon J, Buckler ES, Erb M, Köllner TG, Jander G (2013) Natural variation in
maize aphid resistance is associated with 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-
one glucoside methyltransferase activity. Plant Cell 25: 2341-2355

Moghe GD, Leong BJ, Hurney SM, Jones AD, Last RL (2017) Evolutionary routes to
biochemical innovation revealed by integrative analysis of a plant-defense related
specialized metabolic pathway. Elife 6: e28468

Mount DW (2007) Using the basic local alignment search tool (BLAST) Cold Spring Harbor
Protocols doi:10.1101/pdb.top17

Mulot M, Boissinot S, Monsion B, Rastegar M, Clavijo G, Halter D, Bochet N, Erdinger M,
Brault V (2016) Comparative analysis of RNAi-based methods to down-regulate
expression of two genes expressed at different levels in *Myzus persicae*. Viruses 8: 316

Murashige T, Skoog FA (1962) A revised medium for rapid growth and bioassays with tobacco
tissue cultures. Physiol Plant 15: 473-497

Ning J, Moghe GD, Leong B, Kim J, Ofner I, Wang Z, Adams C, Jones AD, Zamir D, Last
RL (2015) A feedback-insensitive isopropylmalate synthase affects acylsugar
composition in cultivated and wild tomato. Plant Physiol 169: 1821-1835

O' Connell MA, Medina AL, Sanchez Pena Pand Trevino MB (2007) Molecular genetics of
drought resistance response in tomato and related species. In Razdan, M. K., & Mattoo,
A. K. (eds). Genetic Improvement of Solanaceous Crops 2: Enfield USA 261-283

Penuelas J, Filella I, Biel C, Serrano L, Save R (1993) The reflectance at the 950-970 nm
region as an indicator of plant water status. International Journal of Remote Sensing 14:
1887-1905

Pitino M, Hogenhout SA (2013) Aphid protein effectors promote aphid colonization in a plant
species-specific manner. Molecular Plant-Microbe Interactions 26: 130-139

Powell JD (2015) From pandemic preparedness to biofuel production: tobacco finds its
biotechnology niche in North America. Agriculture-Basel 5: 901-917

Ramsey JS, Elzinga DA, Sarkar P, Xin YR, Ghanim M, Jander G (2014) Adaptation to
nicotine feeding in *Myzus persicae*. J Chem Ecol 40: 869-877

Ramsey JS, Wilson AC, De Vos M, Sun Q, Tamborindeguy C, Winfield A, Malloch G,
Smith DM, Fenton B, Gray SM, Jander G (2007) Genomic resources for *Myzus*
persicae: EST sequencing, SNP identification, and microarray design. BMC Genomics 8:
423

Rodriguez AE, Tingey WM, Mutschler MA (1993) Acylsugars of *Lycopersicon pennellii* deter
settling and feeding of the green peach aphid (Homoptera, Aphididae). Journal of
Economic Entomology 86: 34-39

Rodriguez PA, Stam R, Warbroek T, Bos JIB (2014) Mp10 and Mp42 from the aphid species
Myzus persicae trigger plant defenses in *Nicotiana benthamiana* through different
activities. Molecular Plant-Microbe Interactions 27: 30-39

Schiavinato M, Strasser R, Mach L, Dohm JC, Himmelbauer H (2019) Genome and
transcriptome characterization of the glycoengineered *Nicotiana benthamiana* line
DeltaXT/FT. BMC Genomics 20: 594

Schilmiller A, Shi F, Kim J, Charbonneau AL, Holmes D, Jones AD, Last RL (2010) Mass
spectrometry screening reveals widespread diversity in trichome specialized metabolites
of tomato chromosomal substitution lines. Plant Journal 62: 391-403

Schilmiller AL, Charbonneau AL, Last RL (2012) Identification of a BAHD acetyltransferase
that produces protective acyl sugars in tomato trichomes. Proc Natl Acad Sci U S A 109:
16377-16382

Schilmiller AL, Moghe GD, Fan P, Ghosh B, Ning J, Jones AD, Last RL (2015) Functionally
divergent alleles and duplicated Loci encoding an acyltransferase contribute to acylsugar
metabolite diversity in *Solanum* trichomes. Plant Cell 27: 1002-1017

Schmittgen TD, Livak KJ (2008) Analyzing real-time PCR data by the comparative C(T)
method. Nat Protoc 3: 1101-1108

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image
analysis. Nature Methods 9: 671-675

Shepherd RW, Wagner GJ (2007) Phylloplane proteins: emerging defenses at the aerial
frontline? Trends in Plant Science 12: 51-56

Simmons AT, Gurr GM, McGrath D, Martin PM, Nicol HI (2004) Entrapment of
Helicoverpa armigera (Hubner) (Lepidoptera:Noctuidae) on glandular trichomes of
Lycopersicon species. Australian Journal of Entomology 43: 196-200

Simon B, Cenis JL, Demichelis S, Rapisarda C, Caciagli P, Bosco D (2003) Survey of
Bemisia tabaci (Hemiptera: Aleyrodidae) biotypes in Italy with the description of a new
biotype (T) from *Euphorbia characias*. Bull Entomol Res 93: 259-264

Slocombe SP, Schauvinhold I, McQuinn RP, Besser K, Welsby NA, Harper A, Aziz N, Li
Y, Larson TR, Giovannoni J, Dixon RA, Broun P (2008) Transcriptomic and reverse
genetic analyses of branched-chain fatty acid and acyl sugar production in *Solanum*
pennellii and *Nicotiana benthamiana*. Plant Physiol 148: 1830-1846

Song Z, Li S, Chen X, Liu L, Song Z (2006) Synthesis of sucrose esters. Forestry Studies in
China 8: 26-29

Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics 30: 1312-1313

Stone WD, Pellicore MJ, Hagstrom S, Lawton TJ (2020) HSIviewer: Pushbutton
hyperspectral image analysis for rapid plant phenotyping. in review

Thompson JD, Higgins DG, Gibson TJ (1994) Clustal-W - Improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific
gap penalties and weight matrix choice. Nucleic Acids Research 22: 4673-4680

Thurston R (1961) Resistance in *Nicotiana* to the green peach aphids and some other tobacco
insect pest. Journal of Economic Entomology 54: 946-949

Tzin V, Yang X, Jing X, Zhang K, Jander G, Douglas AE (2015) RNA interference against
1046   gut osmoregulatory genes in phloem-feeding insects. J Insect Physiol 79: 105-112

1047   **Van Eck J, Keen P, Tjahjadi M** (2019) *Agrobacterium tumefaciens*-mediated transformation of
1048   tomato. Methods Mol Biol 1864: 225-234

1049   **Van T, Weinhold A, Ullah C, Dressel S, Schoettner M, Gase K, Gaquerel E, Xu SQ,**
1050   **Baldwin IT** (2017) O-acyl sugars protect a wild tobacco from both native fungal
1051   pathogens and a specialist herbivore. Plant Physiology 174: 370-386

1052   **Velásquez AC, Chakravarthy S, Martin GB** (2009) Virus-induced gene silencing (VIGS) in
1053   *Nicotiana benthamiana* and tomato. J Vis Exp doi: 10.3791/1292

1054   **Wagner GJ, Wang E, Shepherd RW** (2004) New approaches for studying and exploiting an
1055   old protuberance, the plant trichome. Annals of Botany 93: 3-11

1056   **Weinhold A, Baldwin IT** (2011) Trichome-derived O-acyl sugars are a first meal for caterpillars
1057   that tags them for predation. Proceedings of the National Academy of Sciences of the
1058   United States of America 108: 7855-7859

1059   **Worrall EA, Bravo-Cazar A, Nilon AT, Fletcher SJ, Robinson KE, Carr JP, Mitter N**
1060   (2019) Exogenous application of RNAii-inducing double-stranded rna inhibits aphid-
1061   mediated transmission of a plant virus. Front Plant Sci 10: 265

1062

Figure 1. Predominant acylsugars in *Nicotiana benthamiana*

Acylsucroses S3:17(2,7,8), S3:18(2,8,8), S2:16(8,8) and S2:15(7,8) are present in *N. benthamiana*. In the acylsugar structure names, S refers to sucrose, followed by the number of acyl chains, the total length of acyl chains, and the length of each individual chain in parentheses. Although the presence of C2, C7, and C8 chain lengths is confirmed, the specific positions of the acyl chains on the sucrose molecule are hypothesized based on previous observations of acylsugars in *Nicotiana alata* (Moghe et al, 2017), the predicted evolution of the acylsugar biosynthetic pathway, and enzyme promiscuities in the Solanaceae family.

Figure 2. ASAT phylogenetic tree
The evolutionary history of ASATs in the Solanaceae (Tables S1 and S2) was inferred using the Maximum
Likelihood method in RAxML. Presented is a subtree of a larger tree that includes all annotated ASATs
(Figure S1). The branch labels indicate the percentage of trees in which the associated taxa clustered
together (bootstrap of 1000). Only values greater than 50 are presented. The two predicted *N. benthamiana*
ASATs are highlighted in orange and ASATs that were previously chemically characterized are highlighted
in blue.

Figure 3. *N. benthamiana ASAT* **mutations produced with CRISPR/Cas9**

Three gRNAs (sequences shown in red) were designed to edit either *ASAT1* or *ASAT2*. Whereas gRNA3 for ASAT1 is on the sense strand, the other gRNAs are on the antisense strand. For both *ASAT1* and *ASAT2*, we obtained two independent mutations resulting from the corresponding gRNA2 and gRNA3. Single-base mutations and deletions are shown in green. (**A**) *asat1-1* has a five-nucleotide deletion at gRNA3 and a single-nucleotide insertion at gRNA2. *asat1-2* has a 318-nucleotide deletion between the gRNA3 and gRNA2 cutting sites. (**B**) *asat2-1* has a single-nucleotide deletion at gRNA3 and single-nucleotide insertion at gRNA2 leading to a translation frame shift between the two mutations. *asat2-2* has a 115-nucleotide deletion at gRNA3 and a single-nucleotide insertion leading to a translation frame shift at gRNA2.

Figure 4. The abundances of two *Nicotiana benthamiana* acylsugars (S2:17(2,7,8) and S2:18(2,8,8)), and two likely pathway intermediates/fragmentations (S2:15(7,8) and S2:16(8,8))

Acyl sugar LC/MS peak areas were normalized relative to the peak area of Telmisartan, which was added as an internal control, and then to the leaf dry weight (per gram). Error bars represent standard errors from measurements of three plants of each genotype. Significant differences for each acylsugar between different genotypes were tested using one-way ANOVA followed by a Duncan's post hoc test ($p < 0.05$). Differences between groups are denoted with letters.

Figure 5. *Myzus persicae* bioassays with wildtype (WT), *asat1,* and *asat2 Nicotiana benthamiana*
Data shown in all panels were combined from two independent experiments, which are shown in color orange and cyan in panels B-F. (**A**) Survival over 5 days of nymphs placed onto mutant and wildtype *N. benthamiana. S*ignificant differences are indicated for the 5-day time point: ns: not significant, *** *p* < 0.001, mean +/- s.e. of n = 15-16. Significant differences were tested using one-way ANOVA with a fixed factor of genotypes and a block effect of experiment followed by a Bonferroni post hoc test for multiple comparisons. Full statistical data for all time points are in Table S3. (**B**) Aphid growth, as measured by body size after 5 days feeding on *N. benthamiana*. (**C**) Aphid reproduction, as measured by the number of nymphs that were produced by five aphids in one week. Significant differences between different groups (*p* < 0.05) were determined using ANOVA with a fixed factor of genotypes and a block effect of experiment followed by a Duncan's post hoc test and are indicated by lowercase letters above each group in panels B and C. (**D-F**) Aphid choice among detached leaves of each plant genotype, significant differences between genotypes were tested using Chi-square test, ns: not significant, *** *p* < 0.001. no choice: aphids were elsewhere in the Petri dish and not on a leaf. The box plots show the median, interquartile range, maximum and minimum after removal of outliers, and the individual data points.

Figure 6. Potato aphid and whitefly bioassays on *asat2-1* and wildtype (WT) *N. benthamiana.*
(**A-D**) potato aphid (*M. euphorbiae*) bioassays. (**A**) aphid survival in 24 hours (n=15). (**B**) aphid reproduction in 24 hours (n=15). (**C**) aphid choices between detached leaves of each plant genotype (n=15). (**D**) An established *M. euphorbiae* colony on an *N. benthamiana asat2-1* leaf. (**E-G**) whitefly bioassays. (**E**) whitefly survival in 3 days (n=6). (**F**) whitefly reproduction measured as number of eggs produced per adults in 3 days (n=3). (**G**) whitefly choices between plants of each genotype (n=3 for 4 experiments). Significantly differences were tested using independent t-test for aphid and whitefly survival and reproduction data, and Chi-square test was used for aphid and whitefly choice assays. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. no choice: insects were not on a leaf at the end of the experiment. The box plots show the median, interquartile range, maximum and minimum after removal of outliers, and the individual data points.

Figure 7. Caterpillar bioassays on wildtype (WT) and *asat2-1* mutant *Nicotiana benthamiana.*
(**A, B, C**) Larval mass of surviving *Helicoverpa zea*, *Heliothis virescens*, and *Trichoplusia ni* after 10 days after being placed on plants as neonates. (**D, E, F**) Relative growth rate of surviving *H. zea*, *H. virescens*, and *T. ni* on wildtype and *nata2-1* plants. Insects were raised for five days on artificial diet, prior to 7 days of feeding on *N. benthamiana*. Percent survival (number of surviving insects/number of total insects) is shown below each figure. Mean +/- s.e. ,***$p < 0.001$, ns: no significant difference ($P > 0.05$, *t*-test).

Figure 8. Aphid transmission of GFP-expressing turnip mosaic virus (TuMV-GFP) on wildtype (WT) and *asat2-1* plants. Aphids were transferred from *N. benthamiana* plants infected with TuMV-GFP to uninfected plants and the percentage of GFP-expressing recipient plants was recorded over time. Data are mean +/- s.e. of N = 60, with data from three separate experiments combined. Statistical analyses were conducted using one-way ANOVA followed by a Tukey HSD post hoc test. Significant differences were observed from day 5 post treatment and thereafter. Lowercase letters denoted the differences between each group at day 9 post treatment ($p < 0.05$). WT: wildtype.

Figure 9. Virus induced gene silencing (VIGS) of whitefly genes using tobacco rattle virus. Whiteflies were placed on plants infected with TRV expressing VIGS constructs and gene expression was measured after (**A**) one and (**B**) seven days. In each experiment, the empty virus vector (EV) and GFP-carrying virus were used as controls. Gene expression was normalized to 1 for EV control plants. Whitefly survival was assessed on TRV-infected plants after (**C**) seven days of feeding on *asat2-1* plants, and (**D**) seven days of feeding on wildtype plants. GFP/AchE/SQS/TLR7/TSCD: VIGS plants with TRV expressing RNA constructs targeting GFP, AchE, SQS, TLR7, and TSCD, respectively; AchE: acetylcholinesterase, SQS: squalene synthase, TLR7: toll-like receptor 7, TSCD: tubulin-specific chaperon D. The survival and the 7-day qPCR experiments were done for three times. The qPCR experiments for 1-day were done twice. Significant differences were determined using one-way ANOVA with a fixed factor of treatments and a block effect of experiment, followed by a Dunnett's post hoc test for comparing VIGS constructs to the empty vector (EV) control. Mean +/- s.e. of n = 3 (A), n = 9 (B), and n = 27 (C,D). *$p < 0.05$; **$p < 0.01$, ***$p < 0.001$.

Figure 10. Water loss and leaf temperature of wildtype (WT), *asat1***, and** *asat2 Nicotiana benthamiana.* (**A**) Percent of water loss from detached leaves in 24 hours, mean +/- s.e. of n = 15. (**B**) Leaf water content measured by the water band index from hyperspectral imaging , mean +/- s.e. of n = 20. (**C**) Leaf temperatures from leaves of different plant genotypes, mean +/- s.e. of n = 20. ns = not significant, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, Dunnett's test relative to wildtype control.

Figure S1. Phylogenetic analyses of annotated ASATs from Solanaceae species

The ASAT proteins used in constructing the tree are listed in Tables S1 and S2. The evolutionary history of ASATs from Solanaceae species was inferred by using the Maximum likelihood method in RAxML. The branch labels indicate the percentage of trees in which the associated taxa clustered together (bootstrap of 1000).

Figure S2 Protein sequence alignment of *Nb*ASAT1 and *Nb*ASAT2 with other functionally characterized ASATs in the Solanaceae species. The alignment was generated using Clustal Omega and visualized using Mview. Colors are coded based on the level of sequence identity.

A

Parent ion *m/z*	MS2 fragment/neutral loss peaks *m/z*	rt*	Annotation
681.34	635.33, 509.22, 491.21, 383.12, 179.06, 143.11, 126.10, 59.01, 18.01	1.76	S3:18(2,8,8) Formate Adduct
671.30	681.34, 671.30, 635.33, 509.22, 143.11, 126.10	1.76	S3:18(2,8,8) Chloride Adduct
635.32	509.22, 491.21, 467.21, 383.12, 365.11, 179.06, 143.11, 126.10, 18.01	1.76	S3:18(2,8,8)
639.32	593.32, 467.21, 449.20, 323.10, 143.11, 126.10	1.70	S2:16(8,8) Intermediate Formate Adduct
593.32	467.21, 323.10, 143.11, 126.10	1.70	S2:16(8,8) Intermediate
667.32	621.31, 509.22, 495.03, 383.12, 179.06, 143.11, 129.09, 126.10, 112.09	1.70	S3:17(2,7,8) Formate Adduct
621.31	495.03, 383.12, 365.11, 143.11, 129.09, 126.10, 59.01, 18.01	1.70	S3:17(2,7,8)
625.31	579.30, 467.21, 453.20, 323.10, 143.11, 129.09, 126.10	1.64	S2:15(7,8) Intermediate Formate Adduct
555.23	509.22, 383.12, 143.11, 126.10, 59.01	1.76	S2:10(2,8) In-Source Fragment Formate Adduct
509.22	449.20, 383.12, 143.11, 126.10, 59.01	1.76	S2:10(2,8) In-Source Fragment
467.21	341.11, 323.10, 144.11, 143.11	1.70	S1:8(8) In-Source Fragment
383.12	383.12, 341.11, 59.01	1.76	S1:2(2) In-Source Fragment

* rt: retention time of the predominant peak for each mass.

B

Figure S3. MS/MS chromatography of *N. benthamiana* acylsugars

(**A**) *N. benthamiana* **a**cylsugar LC/MS peak identification and annotation (**B**) MS[2] chromatography of the two predominant acyl sugars, S2:17(2,7,8) and S2:18(2,8,8). The names of the acylsugars are highlighted in blue and the chromatography is aligned on the x axis based on *m/z*. Highlighted in orange boxes are some of the abundant characteristic mass features, with their compositions in parenthesis.

Figure S4. *ASAT* **mutations do not affect trichome morphology and abundance**
The white arrow indicates the small trichomes and yellow arrow indicates the large swollen-stalk trichomes. Trichome pictures were taken from both the front and back sides of the leaf. Examples of leaves are put side-by-side from different genotypes to facilitate comparisons. Scale bars = 200 μm.

Figure S5. Pairwise aphid choice assay with wildtype and *ASAT* mutant leaves
Each experiment included detached leaves from 5-8 plants of each genotype and was repeated twice
with similar results. Chi-square test was used for difference for all choice assays between plant
genotypes. **$p < 0.01$, ***$p < 0.001$, ns: not significant. no choice: aphids were elsewhere in the Petri
dish and not on leaves at the end of the experiment. The box plots show the median, interquartile range,
maximum and minimum after removal of outliers, and the individual data points.

Figure S6. Aphid bioassays on wildtype and *asat2-1* mutant *N. benthamiana*
(**A**) without *M. persicae* feeding. (**B**) with *M. persicae* feeding for one month. One wildtype (right) and one *asat2-1* (left) plant were placed in the same tray in an insect cage. For the aphid challenge, ten adult *M. persicae* (from an aphid colony on *N. tabacum*) were released on each of the wildtype and the *asat2-1* plants. The pictures were taken about one month after the aphid release. In the case of control plants without aphids (**A**), wildtype and *asat2-1* look similar. (**B**) After one month of aphid feeding, the asat2-1 and wildtype plants look very different, with the wildtype plant looking vigorous and the *asat2-1* line being smaller and wilted. On wildtype *N. benthamiana*, *M. persicae* reproduce slowly and sometimes not at all. If they do survive, they tend to feed on the undersides of senescing leaves. By contrast, *M. persicae* are able to establish a dense colony on the *asat2-1* mutant and feed from the youngest leaves, which tend to be the most nutritious but also the best-defended. Plants in three replications of this experiment looked similar.

Figure S7. Whitefly bioassays on wildtype and *asat2-1* mutant *N. benthamiana*
(**A**) Most whiteflies died within three days after release on wildtype *N. benthamiana*. Arrow indicates dead whiteflies. (**B-D**) Whiteflies survived and completed their life cycle on *asat2-1* mutant *N. benthamiana*. At 23 days after release, whiteflies of different life stages were present on *asat2-1* mutant plants. Arrows indicate eggs (**B**), eggshells and nymphs (**C**), and adults (**D**) on *asat2-1* mutant plants.

Figure S8. Independent repeat of water loss and leaf temperature shown in Figure 8.
(**A**) Percent of water loss from detached leaves after 24 hours, mean +/- s.e. of n =16. (**B**) Leaf water content measured by the water band index from hyperspectral imaging, mean +/- s.e. of n = 10. (**C**) Leaf temperatures from leaves of different plant genotypes, mean +/- s.e. of n=10 for wildtype and n=5 for mutants. ns, not significant, *p < 0.05, **p < 0.01, ***p < 0.001, Dunnett's test relative to wildtype control. Error bars = standard error. NS = not significant.

Figure S9. *ASAT* **mutagenesis by transiently expressing Cas9 and gRNAs in** *N. benthamiana* **leaves via**
Agrobacterium **infiltration**

Three gRNAs each were used for *ASAT1* and *ASAT2* mutagenesis to test efficacy. (**A**) PCR of plants that
were infiltrated with combinations of *ASAT1* gRNA constructs. (**B**) DNA sequence confirmation of a
representative sample (*i.e.* g23OD3) that displayed a shorter band in panel A. (**C**) PCR of plants that were
infiltrated with combinations of *ASAT2* gRNA constructs. (**D**) Restriction enzyme digestion site loss in PCR
of g23OD1.5 (from panel C), where, prior to the PCR reaction, genomic DNA was digested with *Kpn*I to
remove wild type *ASAT2*. The *Kpn*I restriction site is only present in the expected deletion region. (**E**)
Sequence confirmation of sample g23+*Kpn*I from digestion site loss PCR in panel D. Arrow heads indicate
the short versions of target genes after a deletion by the two working gRNA constructs. Arrows indicate the
positions expected to be cut in gene sequences. The label for each lane in the gel: M, 100 bp DNA marker
ladder; WT, samples from wild type plants; ddH2O, no genomic DNA PCR control.

release point

Figure S10. Experimental setups for *Myzus persicae* bioassays

(**A**) *Myzus persicae* and (**B**) *Macrosiphum euphorbiae* no-choice assays. Aphids were caged on individual *Nicotiana benthamiana* leaves. Survival, reproduction, and size were assessed at time points described in the methods section. (**C**) Aphid choice assays. Two *N. benthamiana* leaves were placed in a 15 cm diameter Petri dish with their petioles inserted in moistened cotton swabs, ten aphids were released into the Petri dish at the indicated position, the dish was covered and placed under a 16:8 light:dark photoperiod, and, after 24 hours, the aphids residing on each leaf were counted.