Temperature-dependent effects of house fly proto-Y chromosomes on gene expression

Kiran Adhikari[1], Jae Hak Son[1,4], Anna H. Rensink[2], Jaweria Jaweria[1], Daniel Bopp[3], Leo W.
Beukeboom[2], Richard P. Meisel[1]*

1. Department of Biology and Biochemistry, University of Houston, Houston, TX USA
77204-5001
2. Groningen Institute for Evolutionary Life Sciences, University of Groningen, 9700 CC
Groningen, The Netherlands.
3. Institute of Molecular Life Sciences, University of Zurich, Switzerland CH-8057
4. Current affiliation: Department of Epidemiology of Microbial Diseases, Yale University
School of Public Health, New Haven, CT 06520

* Corresponding author:

Richard P. Meisel
Department of Biology and Biochemistry
University of Houston
3455 Cullen Blvd
Houston, TX
77204-5001

rpmeisel@uh.edu

Abstract

Sex determination, the developmental process by which sexually dimorphic phenotypes are

established, evolves fast. Species with polygenic sex determination, in which master regulatory

genes are found on multiple different proto-sex chromosomes, are informative models to study

the evolution of sex determination. House flies are such a model system, with male determining

loci possible on all six chromosomes and a female-determiner on one of the chromosomes as

well. The two most common male-determining proto-Y chromosomes form latitudinal clines on

multiple continents, suggesting that temperature variation is an important selection pressure

responsible for maintaining polygenic sex determination in this species. To identify candidate

genes that may be under selection, we used RNA-seq to test for temperature-dependent effects

of the proto-Y chromosomes on gene expression in adult house flies. We find no evidence for

ecologically meaningful temperature-dependent expression of sex determining genes between

male genotypes, but we were likely not sampling an appropriate developmental time-point to

identify such effect. In contrast, we identified many other genes whose expression depends on

the interaction between proto-Y chromosome genotype and temperature, including genes that

encode proteins involved in reproduction, metabolism, lifespan, stress response, and immunity.

Notably, genes with genotype-by-temperature interactions on expression are not enriched on

the proto-sex chromosomes. Moreover, there is no evidence that temperature-dependent

expression is driven by chromosome-wide expression divergence between the proto-Y and

proto-X alleles. Therefore, if temperature-dependent gene expression is responsible for

differences in phenotypes and fitness of proto-Y genotypes across house fly populations, these

effects are driven by a small number of temperature-dependent alleles on the proto-Y

chromosomes that may in turn affect the expression of genes on other chromosomes.

Introduction

Sex determination establishes sexually dimorphic developmental pathways, either based on genetic differences between males and females or environmental cues (Beukeboom & Perrin 2014). In species with genotypic sex determination, a single master regulatory locus (e.g., *SRY* on the human Y chromosome) is often enough to initiate whether the embryo develops into a male or female (Sinclair *et al.* 1990; Goodfellow & Lovell-Badge 1993). However, in polygenic sex determination systems, multiple master sex determining loci segregate independently, often on different chromosomes (Moore & Roberts 2013). Most population genetics models predict that polygenic sex determination will be an evolutionary intermediate between different monogenic sex determination systems, and the factors responsible for maintaining polygenic sex determination as a stable polymorphism are poorly understood (Rice 1986; van Doorn 2014). Models that do allow for the stable maintenance of polygenic sex determination require opposing (sexually antagonistic) fitness effects of sex chromosomes in males and females (Orzack *et al.* 1980; van Doorn & Kirkpatrick 2007, 2010). Understanding how other selection pressures can maintain polygenic sex determination would provide valuable insight into the factors that drive the evolution of sex determination.

House fly (*Musca domestica*) is a well suited model for studying polygenic sex determination because multiple male and female determining loci segregate on different chromosomes in natural populations (Hamm *et al.* 2015). Male sex in house fly is initiated by the gene *Musca domestica male determiner, Mdmd* (Sharma *et al.* 2017). *Mdmd* arose via the recent duplication of the ubiquitous splicing factor *nucampholin* (*Md-ncm*) after the divergence of house fly from its close relative *Stomoxys calcitrans*. *Mdmd* promotes male development by causing the house fly ortholog of *transformer* (*Md-tra*) to be spliced into non-functional isoforms with premature stop codons (Hediger *et al.* 2010). The lack of functional Md-Tra protein leads to male-specific splicing of *doublesex* (*Md-dsx*) and *fruitless* (*Md-fru*), the two known downstream targets of *Md-tra* (Hediger *et al.* 2004; Meier *et al.* 2013). In the absence of *Mdmd*, *Md-tra* is spliced into a functional transcript that is translated into a protein that promotes female specific splicing of *Md-dsx* and inhibits splicing of the male isoform of *Md-fru*.

Mdmd can be found on multiple different chromosomes in house fly (Sharma *et al.* 2017), and it is most commonly found on the third (III^M) and Y (Y^M) chromosomes (Hamm *et al.* 2015). While Y^M is conventionally referred to as the Y chromosome, both III^M and Y^M are young proto-Y chromosomes that are minimally differentiated from their homologous proto-X

chromosomes (Meisel *et al.* 2017; Son & Meisel 2021). The proto-Y chromosomes are clinally

distributed—with III^M most common at southern latitudes and Y^M most common at northern

latitudes—across multiple continents (Hiroyoshi 1964; Mcdonald *et al.* 1975; Denholm *et al.*

1986; Hamm *et al.* 2005). The frequencies of III^M and Y^M in natural populations have remained

stable for decades (Kozielska *et al.* 2008; Meisel *et al.* 2016). This clinal distribution of III^M and

Y^M, along with their stable frequencies across populations, suggests that natural selection

maintains the polymorphism.

A female determining allele of *Md-tra* (*Md-tra^D*) is also found in some house fly

populations (McDonald *et al.* 1978; Hediger *et al.* 2010). *Md-tra^D* can initiate female

development in embryos with at least three *Mdmd* chromosomes (Schmidt *et al.* 1997; Hediger

et al. 1998). In some populations, both Y^M and III^M can be found, with some males carrying one

copy of two different proto-Y chromosomes or homozygous for a proto-Y (e.g., Hamm & Scott

2008, 2009). *Md-tra^D* is most common in populations with a high frequency of these "multi-Y"

males, which results in a sex ratio with an equal number of males and females (Meisel *et al.*

2016).

The natural distribution of III^M and Y^M hints at a possible genotype-by-temperature (G×T)

interaction that could explain the stable maintenance of Y^M-III^M clines. Temperature is not the

only selection pressure that could vary along the clines, but seasonality in temperature is the

best predictor of the frequencies of the proto-Y chromosomes across populations (Feldmeyer *et*

al. 2008). There are at least two non exclusive ways in which temperature-dependent selection

pressures could maintain the III^M-Y^M polymorphism. First, alleles on the III^M and Y^M

chromosomes (other than the *Mdmd* locus) could have temperature-dependent fitness effects.

In this scenario, the III^M-Y^M clines would be maintained in a similar way to how temperature

variation maintains opposing clines of heat and cold tolerance in *Drosophila melanogaster*

between tropical and temperate regions (Hoffmann *et al.* 2002). Second, it is possible that the

Mdmd copies on the III^M and Y^M chromosomes differ in their temperature dependent activities,

such that *Mdmd* on the III^M chromosome increases male fitness at warm temperatures and

Mdmd on the Y^M chromosome increases fitness at colder temperatures. This is analogous to

how, in some fish and reptile species, temperature can drive sex determination and override the

outcomes of genotypic sex determining systems (Shine *et al.* 2002; Quinn *et al.* 2007; Radder *et*

al. 2008; Holleley *et al.* 2015).

We investigated if temperature-dependent phenotypic effects of III^M and Y^M could be

driven by differential gene expression in males across temperatures. We selected gene

expression as a phenotypic read-out of G×T interactions because temperature-dependent

differences in gene expression are well documented in clinally distributed genetic variation

(Levine *et al.* 2011; Zhao *et al.* 2015). Specifically, we evaluated how G×T interactions affect

gene expression in male house flies carrying either a III^M or Y^M chromosome. We used RNA-seq

to study gene expression in two nearly isogenic lines of house flies, differing only by their

proto-Y chromosome, reared at two developmental temperatures. This allowed us to assess the

effects of the entire III^M and Y^M chromosomes. We also used quantitative reverse transcription

PCR (qRT-PCR) to investigate the temperature-dependent expression of *Mdmd*.

Materials & Methods

qRT-PCR samples and analysis

We used qRT-PCR to measure the expression of *Mdmd* and its paralog *Md-ncm* in two

Y^M strains and two III^M strains. The strains were grouped into two pairs, with one Y^M strain and

one III^M strain per pair. In the first pair, we used the Y^M strain IsoCS and the III^M strain CSkab

(both from North America). IsoCS and CSkab share a common genetic background of the

Cornell susceptible (CS) strain (Scott *et al.* 1996). IsoCS was previously created by crossing a

Y^M chromosome from Maine onto the CS background (Hamm *et al.* 2009). We created CSkab

by backcrossing the III^M chromosome from the KS8S3 strain collected in Florida (Kaufman *et al.*

2010) onto the CS background, using an approach described previously (Son *et al.* 2019). In the

second pair, we used two European strains: the Y^M strain GK-1 from Gerkesklooster

(Netherlands) and the III^M strain SPA3 from near Girona (Spain). GK-1 and SPA3 were

maintained in the lab, each as inbred populations, for approximately 40 and 50 generations,

respectively.

We raised all strains at 18°C and 27°C for two generations with 12:12-h light:dark

photoperiods. Adult males and females for each GxT combination were housed in cages with *ad*

libitum containers of 1:1 combinations of sugar and non-fat dry milk and *ad libitum* containers of

water. Females were provided with a standard medium of wheat bran, calf manna, wood chips,

yeast, and water in which to lay eggs for 12-24 hrs (Hamm *et al.* 2009). The resulting larvae

were maintained in the same media within 32 oz containers. Adult females did not lay a

sufficient number of eggs at 18°C, so the adults from the 18°C colonies were transferred to

22°C for egg laying for 1-2 days. The eggs collected at 22°C were then moved back to 18°C for

larval development, pupation, and emergence as adults. We maintained the colonies at these

temperatures for two generations. Collecting flies after two generations ensured at least one full egg-to-adult generation at the appropriate temperature.

For qRT-PCR experiments involving the North American IsoCS and CSkab strains, abdomen samples were dissected from 5 day old adult males after being anesthetized with CO_2. For qRT-PCR assessments on the European GK and SPA3 strains, full body samples were collected from 5 day old adult males after being anesthetized with CO_2. Tissue samples from 5-7 males were pooled in each of three biological replicates for each genotype (Y^M and III^M) by temperature (18°C and 27°C) combination. The collected tissues were homogenized in TRIzol reagent (Life Technologies) using a motorized grinder in a 1.5 ml microcentrifuge tube. For the North American strains, the Direct-zol RNA MiniPrep kit (Zymo Research) was used to extract RNA from the homogenized samples. The isolated RNA was reverse transcribed into cDNA with MLV RT (Promega), following the manufacturer's protocol. For the European strains, the RNA phase following centrifugation with TRIzol reagent was separated using chloroform and precipitated by using isopropanol and ethanol. The isolated RNA was reverse transcribed into cDNA using RevertAid H minus 1st strand kit (Fermentas #K1632) according to the manufacturer's protocol.

We conducted qRT-PCR of cDNA from the male flies. We used qRT-PCR primers (Supplementary Table 1) to uniquely amplify *Mdmd* and *Md-ncm* without amplifying the other paralog (Sharma *et al.* 2017). Primers were additionally used to amplify cDNA from a transcript (LOC101888902) that is not differentially expressed between Y^M and III^M males as an internal control for cDNA content in each biological replicate (Meisel *et al.* 2015). The IsoCS and CSkab samples were assayed on a StepOnePlus machine using PowerUp SYBR Green Master Mix (Applied Biosystems). The GK and SPA3 samples were assayed on a Applied Biosystems qPCR cycler 7300 machine using Quanta perfecta SYBR Green Fastmix (Quanta bio). We measured the abundance of PCR products from each primer pair in three technical replicates of three biological replicates for each G×T combination. With the same primer pairs, we also measured the expression of serial dilutions (1/1, 1/5, 1/25, 1/125, and 1/625) of cDNA from independent biological collections of house flies. Samples were interspersed across 96-well microtiter plates to minimize batch effects.

We constructed standard curves for each primer pair by calculating the linear relationship between CT values and $\log_{10}$(concentration) from the serial dilutions using the lm() function in the R statistical programming package (R Core Team 2019). We then used the equations of the standard curves to calculate the concentration of transcripts (i.e., cDNA) from

Mdmd and *Md-ncm* in each technical replicate. We next determined a normalized expression

level of each technical replicate by dividing the concentration of the technical replicate by the

mean concentration of the control transcript (LOC101888902) across the three technical

replicates from the same biological replicate.

We used an analysis of variance (ANOVA) approach to test for the effect of genotype

(Y^M vs III^M), developmental temperature (18°C vs 27°C), and the interaction of genotype and

temperature on the expression of each transcript. To those ends, we used the lmer() function in

the lme4 package (Bates *et al.* 2015) in R to model the effect of genotype (*G*), temperature (*T*),

and the interaction term as fixed effect factors, as well as biological replicate (*r*) as a random

effect, on expression level (*E*):

181                    $$E \sim G + T + G{\times}T + r.$$

We then compared the fit of that full model to a model without the interaction term (*E ~ G + T +*

r) using the anova() function in R. If the full model fits significantly better, that is evidence that

there is a significant G×T interaction on the expression of the transcript.

RNA-seq samples

We used RNA-seq to measure gene expression in the Y^M strain IsoCS and a III^M strain

known as CSrab. IsoCS (described above) and CSrab have different proto-Y chromosomes on

the shared CS genetic background (Scott *et al.* 1996). We created CSrab by backcrossing the

III^M chromosome of a spinosad-resistant strain, rspin (Shono & Scott 2003), onto the CS

background, using the same approach as we used to create CSkab, described elsewhere (Son

et al. 2019). These strains are normally raised at 25°C, but were raised at different temperatures

(18°C or 29°C) in our experiment in order to determine the effect of genotype and temperature

on gene expression.

Colonies of both strains were reared at 18°C and 29°C for two generations with at least

one full egg-to-adult generation, as described above. We therefore had four combinations of

genotype (Y^M and III^M) and temperature (18°C and 29°C). We controlled for the adult density

using 35 adult males and 35 adult females for each G×T combination. We also controlled for

larval density with 100 larvae per 32 oz container. Third generation males obtained from second

generation females were collected and reared separately from the females at their respective

developmental temperatures for 1–8 days before RNA extraction.

For the RNA-seq experiments, head and testis samples from 1–8 day old males were

dissected in 1% PBS solution after being anesthetized with CO_2. We dissected testes from

15–20 house flies per each of three replicates of each G×T combination. Similarly, 5 heads were

dissected for each of three biological replicates for each G×T combination. The collected tissues

were homogenized in TRIzol reagent (Life Technologies) using a motorized grinder in a 1.5 mL

microcentrifuge tube. The Direct-zol RNA MiniPrep kit (Zymo Research) was used to extract

RNA from the homogenized samples. RNA-seq library preparation was carried out using the

TruSeq Stranded mRNA Kit (Illumina). Qualities of these libraries were assessed using a 2100

Bioanalyzer (Agilent Technologies, Inc.). Libraries were then sequenced with 75 bp single-end

reads on high output runs of an Illumina NextSeq 500 at the University of Houston Seq-N-Edit

Core. All testis samples (i.e., all replicates of each G×T combination) were sequenced together

in a single run, and all head samples were sequenced together on a separate run. All RNA-seq

data are available in the NCBI Gene Expression Omnibus under accession GSE136188

(BioProject PRJNA561541, SRA accession SRP219410).

RNA-seq data analysis

RNA-seq reads were aligned to the annotated house fly reference genome

Musca_domestica-2.0.2 (Scott *et al.* 2014) using HISAT2 (Kim *et al.* 2015) with the default

settings of a maximum mismatch penalty of 6 and minimum penalty of 2, and a soft-clip penalty

of maximum 2 and minimum 1 (Supplementary Tables 2 and 3). We next used SAMtools (Li *et*

al. 2009) to sort the aligned reads. The sorted reads were assigned to annotated genes (*M.*

domestica Annotation Release 102) using htseq-count in HTSeq (Anders *et al.* 2015). We only

included uniquely mapped reads, and we excluded reads with ambiguous mapping and reads

with a mapping quality of less than 10.

We analyzed the exon-level expression of the sex determining genes *Md-tra*

(LOC101888218) and *Md-dsx* (LOC101895413) for each G×T combination. To do so, we first

determined the read coverage across *Md-tra* and *Md-dsx* transcripts using the 'mpileup' function

in SAMtools (Li et al., 2009). We then calculated normalized read depth (D_{ijk}) at each site i

within each gene in library j for each G×T combination k by dividing the number of reads

mapped to a site (r_{ijk}) into the total number of reads mapped in that library (R_{jk}), and we

multiplied that value by one million:

231
$$D_{ijk} = \left(\frac{r_{ijk}}{R_{jk}} \right) 10^6$$
.

For each site within each gene, we then calculated the average D_{ijk} across all three libraries for

each G×T combination ($\overline{D}_{ik}$).

We also used the DESeq2 package in R (Love *et al.* 2014) to analyze differential

expression of all annotated genes between all G×T combinations. To do so, we used a linear

model that included genotype (Y^M or III^M), developmental temperature, and their interaction term

to predict gene expression levels:

E ~ G + T + G×T.

Genes for which the interaction term has a false discovery rate (FDR) corrected *P*-value

(Benjamini & Hochberg 1995) of less than 0.05 were considered to be differentially expressed

as a result of the G×T interaction. The same FDR corrected cutoff was used to test for genes

that are differentially expressed according to genotype or temperature, by testing for the effect

of *G* or *T* using results analyzed with the full model. For principal component analysis (PCA),

hierarchical clustering, and non-metric multidimensional scaling (NMDS), we analyzed

regularized log transformed count data generated by the rlog() function in DESeq2. NMDS was

carried out using metaMDS() function from the vegan package in R with the autotransform =

FALSE option (Oksanen *et al.* 2019).

We performed a gene ontology (GO) analysis to test for enrichment of functional classes

amongst differentially expressed genes. To assign GO terms to house fly genes, we first used

BLASTX to search house fly transcripts against a database of all *D. melanogaster* proteins

(Gish & States 1993). We took this approach because GO assignments are missing for most

house fly genes. The top hit for each house fly gene obtained from BLASTX was used to assign

a FlyBase ID to each house fly transcript. These *D. melanogaster* homologs were then used in

DAVID 6.8 (Huang *et al.* 2009a; b) to identify GO terms that are significantly enriched amongst

differentially expressed genes (FDR corrected *P* < 0.05).

Allele-specific expression analysis

We tested for differential expression of third chromosome genes between the allele on

the III^M chromosome and the allele on the standard (non-*Mdmd*) third chromosome in III^M males.

To do so, we followed the Genome Analysis Toolkit (GATK) best practices workflow for single

nucleotide polymorphism (SNP) and insertion/deletion (indel) calling to identify sequence

variants in our RNA-seq data (McKenna *et al.* 2010; Meisel *et al.* 2017). We first used STAR

(Dobin *et al.* 2013) to align reads from the 12 testis libraries and 12 head libraries to the house

fly reference genome (Musca_domestica-2.0.2). We then used the splice junction information

from the first alignment to create a new index that was used to perform a second alignment.

Using *de novo* transcripts identified with STAR serves to reduce read-mapping biases

associated with an incomplete transcript annotation. After adding read group information to the

SAM file thus generated, we marked duplicates. We next used SplitNCigarReads to reassign

mapping qualities to 60 with the ReassignOneMappingQuality read filter for alignments with a

mapping quality of 255. We used RealignerTargetCreator to identify and IndelRealigner to

realign the indels. We used BaseRecalibrator and variant calls from a previous RNA-seq

analysis (Meisel *et al.* 2017) to recalibrate the realigned reads. The realigned reads were then

used for variant calling with HaplotypeCaller with emission and calling thresholds of 20. We

filtered the variants obtained using VariantFiltration with a cluster window size of 35 bp, cluster

size of 3 SNPs, FS > 30, and QD < 2. This filtering was applied because there may be

preferential mapping of reads containing SNPs found in the reference genome relative to reads

with alternative SNPs (Stevenson *et al.* 2013; Zimmer *et al.* 2016). By excluding SNPs found in

clusters of at least 3 in a 35 bp window from our analysis, we can greatly reduce read-mapping

biases from our estimates of allele-specific expression (Son & Meisel 2021).

We then used all the generated gvcf files to carry out joint genotyping using

GenotypeGVCFs. We performed separate joint genotyping for testis and head libraries. The

variants from Joint Genotyping were then filtered using VariantFiltration with FS > 30 and QD <

2. We used the vcfR package in R (Knaus & Grünwald 2017) to extract information from vcf files

obtained from joint genotyping. For downstream analysis, we only kept SNPs (i.e., variants

where the reference and alternate allele are 1 bp), and excluded small indels.

To test for allele-specific expression, we first assigned sequence variants to the III^M and

standard third (III) chromosomes. This was only done for sites that were heterozygous in III^M

males and homozygous in Y^M males (all other variable sites on the third chromosome were

discarded) because these are the only alleles we can assign to either the III^M or III chromosome.

This is because Y^M males are homozygous for the III chromosome (X/Y^M; III/III), and III^M males

are heterozygous (X/X; III^M/III). For every variable site, we assigned the allele shared by both

III^M and Y^M males to the III chromosome, and the allele unique to III^M males to the III^M

chromosome. We calculated the sum of read depth for each allele across all three sequencing

libraries (i.e., replicates) of each G×T combination. For each gene, we calculated the average

normalized read depth across all variable sites within the gene separately for the III^M and III

alleles at each temperature. To compare the expression of the III^M and III alleles, we calculated

the difference in sequencing coverage between III^M and III alleles at each site for each

temperature separately. We calculated the average difference in expression of III^M and III alleles

in each gene at each temperature k, d_k, as follows:

299
$$d_k = \frac{1}{n} \sum_{i=1}^{n} \left(\frac{r_{i1k}}{R_k} - \frac{r_{i2k}}{R_k} \right) 10^6$$
,

300 where r_{i1k} is the expression of the III^M allele at site i (out of n total polymorphic sites) and

301 temperature k (either 18°C or 29°C), r_{i2k} is the expression of the III allele at site i and

302 temperature k, and R_k is the total number of mapped reads in III^M males at temperature k. We

303 then calculated standard error of d_k across all sites for each gene at each temperature.

Results

Genotype and temperature affect genome-wide gene expression profiles

We used RNA-seq to test for the effects of genotype and developmental temperature on

gene expression in heads and testes of Y^M and III^M house flies raised at 18°C and 29°C. The

purpose of raising the strains at two different temperatures is to expose G×T effects of Y^M and

III^M alleles sampled from natural populations (i.e., genotype-dependent plasticity across

temperatures), not to evolve adaptations to each temperature. We first used PCA, NMDS, and

hierarchical clustering to assess the similarities of the overall gene expression profiles of each

of three replicates of each G×T combination in head and testis separately.

The PCA of the head RNA-seq data (using all 16,540 expressed genes) provides some

evidence for an effect of genotype on gene expression. The first principal component (PC1) of

head gene expression explains 34% of the variance in expression, and the second (PC2)

explains 23% of the variation (Figure 1A). However, there is no clear grouping by genotype or

developmental temperature, which can be best explained by an age-effect in our samples. One

biological replicate of III^M heads at each temperature came from older males (4–8 days old, as

opposed to the other samples which were 1–3 days old). The two older samples had head

expression profiles that clustered separately from the remaining samples in our PCA (Figure

1A). Excluding the two older samples, we found a clear grouping by genotype along PC2, which

explains 28% of the variance in head gene expression (Figure 1B). Because of the effect of age

on head gene expression, we describe results both including and excluding the two older

samples in the remainder of the analyses we present.

Figure 1. Effect of genotype and temperature on genome-wide gene expression in house flies.
Graphs show the first two principal components (PC) explaining gene expression levels in all male heads
(A), heads of young males only (B) and testes (C) samples. Each data point represents a biological
replicate, with PC coordinates determined using regularized log transformed read counts.

In testis, PC1 explains 36% of the variance in expression, and it separates III^M males at
18°C from Y^M males at 29°C (Figure 1C). III^M is found at southern, warmer temperatures,
whereas Y^M is found at northern, colder temperatures. PC1 for testis expression therefore
separates the two genotypes at the temperatures that are opposite from their geographic
distribution (i.e., Y^M occurs at relatively low temperature and III^M at high temperature). PC2
explains 20% of the variation in testis expression and separates III^M at 29°C from Y^M at 18°C
(Figure 1C). Therefore, PC2 separates the two genotypes at temperatures that are consistent
with their geographic distribution. We did not observe a meaningful effect of age on gene
expression in testis (Figure 1C), and we thus did not repeat the analysis excluding the older
testis samples.

We performed the following analyses to evaluate the robustness of our PCA results.
First, we carried out PCA by considering only the 500 most variable genes in head and testis
and observed the same patterns as those described above (Supplementary Figure S1). We
additionally carried out PCA for genes on each chromosome, and the results for each
chromosome were consistent with those across all chromosomes (Supplementary Figures S2,

S3, and S4). Notably, there is very strong differentiation of III^M and Y^M males when we consider

the testis expression of X chromosome and third chromosome genes (Supplementary Figure

S4). This can be explained by the fact that the two genotypes only differ in these chromosomes,

and share the same genetic background for the remaining chromosomes. We also carried out

NMDS and hierarchical clustering of the RNA-seq data. We observed a grouping by genotype in

the NMDS for head samples, and grouping by genotype and temperature in the testis samples

(Supplementary Figure S5). In the hierarchical clustering, we did not observe grouping by

genotype or temperature in head samples while including or excluding the older samples

(Supplementary Figure S6). For testis gene expression, we found some evidence for clustering

first by genotype and then by temperature (Supplemental Figure S6), similar to the PCA.

However, the concordance between clusters and G×T combinations is not perfect.

Genotype and temperature affect the expression of individual genes

To further test for genotype- and temperature-dependent gene expression, we next

identified differentially expressed genes in two types of pairwise comparisons: i) between

genotypes at one developmental temperature (either at 18°C or 29°C), and ii) within a genotype

across the two developmental temperatures. Comparing between genotypes, we found 900

genes that are differentially expressed between Y^M and III^M heads at 18°C, and there were 1378

genes differentially expressed between Y^M and III^M heads at 29°C (Supplementary Table 4,

Supplementary Figure S7). Excluding the two older samples, we found 786 genes that are

differentially expressed between Y^M and III^M heads at 18°C, and 1748 genes differentially

expressed between Y^M and III^M heads at 29°C (Supplementary Table 5, Supplementary Figure

S7). The increase in differentially expressed genes at 29°C when the older samples are

excluded can be explained by reduced variation within the III^M male samples, which should

increase our power to detect differences between III^M and Y^M males. The number of differentially

expressed genes is higher in testis than head: 2413 genes at 18°C and 2199 genes at 29°C are

significantly differentially expressed between Y^M and III^M testes (Supplementary Table 6,

Supplementary Figure S7). This is consistent with previous work that identified more genes

differentially expressed between Y^M and III^M males in testis than head (Meisel *et al.* 2015).

Figure 2. Genes that are differentially expressed between house fly genotypes are significantly enriched on the third chromosome. A) The proportion of house fly genes on each chromosome that are differentially expressed (DE) between Y^M and III^M males is plotted for heads (top) and testes (bottom) of flies raised at 18°C (left) or 29°C (right). B) The proportion of house fly genes on each chromosome that are DE between temperatures are plotted for heads (top row) and testes (bottom row) for III^M (left) and Y^M (right) males separately. Each bar represents the proportion of DE genes on a chromosome (# DE genes / # genes on the chromosome), and dashed lines show the the proportion of DE genes across the genome (# DE genes / # genes assigned to any chromosome). Asterisks indicate P values obtained from Fisher's exact test comparing the number of DE genes on a chromosome, the number of non-DE genes on a chromosome, and the number of DE and non-DE genes across all other chromosomes, after Bonferroni correction (*$P < 0.05$, **$P < 0.005$, ***$P < 0.0005$, ****$P < 0.00005$, *****$P < 0.000005$).

In both head and testis, there is an excess of genes on the third chromosome that are significantly differentially expressed between genotypes at both 18°C and 29°C (Figure 2A), regardless of whether the older samples are excluded (Supplementary Figure S8). This is consistent with different third chromosome genotypes between strains, and it is suggestive of a *cis* effect on gene expression levels (Meisel *et al.* 2015; Son *et al.* 2019). The excess differential expression of chromosome III genes is also consistent with the signal that chromosome III gene expression provides to differentiating III^M and Y^M males (Supplementary Figure S4). The observed proportion of differentially expressed X-linked genes also appears to deviate from the expectation based on the genome-wide average (Figure 2A), but it is not significant because of

low power caused by the small number (<100) of genes on the house fly X chromosome (Meisel

& Scott 2018).

When comparing between temperatures within each genotype, we found 739 genes

significantly differentially expressed between heads of Y^M flies raised at the two different

temperatures (Supplementary Table 4, Supplementary Figure S7). Similarly, 744 genes are

differentially expressed between the heads of III^M flies raised at different temperatures

(Supplementary Table 4, Supplementary Figure S7). When we excluded older samples, we

found 828 genes differentially expressed between heads of III^M males raised at different

temperatures and 1280 genes differentially expressed between the heads of Y^M males

(Supplementary Table 5, Supplementary Figure S7). Once again, the increase in differentially

expressed genes when the older samples are excluded can be explained by greater power to

detect differential expression when the outlier III^M males are removed. This also increases

power to detect differences within Y^M males because we analyze the data with a statistical

model that includes all genotypes, temperatures, and their interactions. In testis, there are 2402

genes in Y^M flies and 1649 genes in III^M flies that are differentially expressed between 18°C and

29°C (Supplementary Table 6, Supplementary Figure S7).

There is no significant chromosomal enrichment of genes that are differentially

expressed between temperatures in either head or testis when we include all samples (Figure

2B), consistent with these comparisons being between flies with the same genotype. However,

we found a modest enrichment of third chromosome genes that are significantly differentially

expressed between temperatures in young Y^M heads, i.e., excluding the two older samples

(Supplementary Figure S8). This is surprising because all Y^M males should have the same third

chromosome genotype, and we do not have an explanation for this pattern. As above, the small

number of genes on the house fly X chromosome greatly reduces our power to detect significant

differences between observed and expected proportions (Meisel & Scott 2018).

G×T interactions affect the expression of a small subset of genes

We next identified individual genes that are differentially expressed between Y^M and III^M

males depending on temperature by testing for significant interactions between genotype and

temperature on gene expression levels. We found 50 genes in head and 247 genes in testis

whose expression significantly differs in response to the G×T interaction when we include all

samples (Supplementary Tables 4 and 6, Supplementary Figure S7). We found 108 genes

differentially expressed in heads in response to the GxT interaction when the two older samples

were excluded (Supplementary table 5, Supplementary Figure S7). Of the genes for which the G×T interaction significantly affects expression in head, there are 26 genes that are shared by the analysis of all heads and when the two older samples are excluded (Supplementary Figure S9). We did not find an enrichment of genes with significant G×T interactions on any chromosome in all male heads, younger male heads, or testes (Supplementary Figure S10).

There are 10 genes that are affected by G×T interactions in both head and testis (Supplementary Figure S9). We would expect <1 gene to be affected by G×T interactions in both head and testis if the G×T effects are independent across tissues. The ten genes we observed are significantly greater than this expectation (z = 10.12, P < 2.2e-16, in a test of proportions), suggesting G×T effects on expression are not independent across tissues. We found 9 genes that are affected by G×T interactions in both testis and young male heads (Supplementary Figure S9), which is significantly greater than the expectation of <2 genes (z = 5.42, P = 5.88e-08, in a test of proportions). We also found an excess of genes that are differentially expressed in both head and testis in all pairwise comparisons between genotypes and temperatures (Supplementary Figure S9). A similar non-independence of expression differences across tissues was previously observed between Y^M and III^M males (Meisel *et al.* 2015).

We characterized the functional annotations of genes that are differentially expressed as a result of G×T interactions. We did not find any GO terms associated with genes significantly differentially expressed as a result of GxT interactions in either testis or head, regardless of whether we include all head samples or exclude the two older samples. However, individual genes are suggestive of biological functions that could be affected by GxT interactions on expression. In head, the genes that were differentially expressed because of G×T interactions include an apolipoprotein-D gene (*LOC101893129*). This gene is homologous to *D. melanogaster NLaz*, which is involved in stress response (Hull-Thompson *et al.* 2009), and it is upregulated in III^M males at 29°C (Figure 3A). Two genes encoding immune effectors (*LOC105261620,* which encodes a Defensin; and *LOC101895951*, which encodes a Lysozyme and is homologous to *D. melanogaster LysP*) were also upregulated in III^M at 29°C (Figure 3A). Three DNA repair genes (*LOC101889156*, homologous to *D. melanogaster Gen*, encoding XPG-like endonuclease; *LOC101899772*, homologous to *maternal haploid*, *mh*, which encodes a protease; and *LOC101899952*, homologous to *Stromalin*, *SA*) are upregulated in Y^M at 18°C (Figure 3A). Lastly, an odorant binding protein-coding gene (*LOC105261913*, homologous to *D. melanogaster Obp56h*) was upregulated in Y^M males at 29°C (Figure 3A).

Figure 3. G×T interactions affect gene expression in house fly head. Graphs show normalized read counts (obtained using DESeq2) for genes significantly differentially expressed because of G×T interactions in A) all male head samples or B) young male heads only. Genes are identified based on their house fly gene ID followed by their *Drosophila melanogaster* homologs in parenthesis. Error bars represent standard errors of the mean.

We also identified genes whose expression depends on the G×T interaction when we exclude the two older head samples. *LOC101895951* (*LysP*), *LOC105261913* (*Obp56h*), *LOC101889156* (*Gen*), *LOC101899772* (*mh*), and *LOC101899952* (*SA*) were also significantly differentially expressed in younger heads in the same direction as when we analyze all male head samples (Supplemental Figure 15). A similar pattern was observed for *Nlaz* expression when we only include young heads, although the G×T effect is not significant (Supplementary Figure 15). Other genes only have significant G×T effects in young male heads, including three genes related to muscle performance (*LOC101893720*, homologous to *D. melanogaster bent, bt*; *LOC101895658*, homologous to *Unc-89*; and *LOC101901052*, homologous to *Myofilin, Mf*), which are all upregulated in Y^M at 18°C (Figure 3B). One gene involved in endoplasmic reticulum (ER) stress response (*LOC101901283*, homologous to *Calx*) is upregulated in III^M males at 29°C (Figure 3B).

Figure 4. G×T interactions affect gene expression in house fly testis. Graphs show normalized read counts (obtained using DESeq2) for genes significantly differentially expressed in testis because of G×T interactions. Genes are identified based on their house fly gene ID followed by their *Drosophila melanogaster* homologs in parenthesis. Error bars represent standard errors of the mean.

In testis, genes with significant G×T effects on expression include those coding for proteins related to reproductive functions: the protamine *ProtB* homolog *LOC101887804*; the *asunder* (*asun*) homolog *LOC101899763*; the *sarah* (*sra*) homolog *LOC101894442*, and the *Farnesyl pyrophosphate synthase* (*Fpps*) homolog *LOC101896699* (Figure 4). Differential expression of reproduction related genes is expected because testis is the largest male reproductive organ in house fly. Other notable genes that are differentially expressed in testis because of G×T interactions include three metabolic genes (*LOC109613297*, which encodes a hexokinase and is homologous to *D. melanogster Hex-t2*; *LOC101901027,* which encodes fructose-1,6-bisphosphatase and is homologous to *D. melanogaster fbp*; and *LOC101901154*, which encodes an aldehyde oxidase, homologous to *AOX3*), all of which are upregulated in Y^M males at 18°C (Figure 4). We also identify one adult lifespan related gene (*LOC101897626*, the homolog of *D. melanogaster pointed*, *pnt*) that is downregulated in III^M males at 29°C, and another lifespan related gene (*LOC101897352*, which encodes cystathionine β-synthase, *Cbs*) that is upregulated in Y^M males at 18°C (Figure 4). Lastly, two immunity-related genes are differentially expressed in testis. One of the immune genes (*LOC101887442*, which encodes a

493 Gram-negative bacteria-binding protein and is homologous to *GNPB3*) is upregulated in Y^M

494 males at 18°C, and the other (*LOC101895929*, which is homologous to *D. melanogaster*

495 *Phenoloxidase 1*, *PPO1*) is upregulated in III^M males at 18°C (Figure 4).

G×T interactions affecting expression of genes in the sex determination pathway

We did not find evidence that the sex determining gene *Md-tra* is differentially expressed

according to a G×T interaction in either all male heads (Supplementary Table 4), young male

heads (Supplementary Table 5), or testes (Supplementary Table 6). This suggests that

temperature-dependent misregulation of the sex determination pathway is not responsible for

fitness differences of Y^M and III^M males across the cline. In comparison, we found evidence of

effects of G×T interactions on the expression of most *Md-tra* exons in both head (including or

excluding older samples) and testis (Supplementary Figures S11 and S12). However, if a G×T

interaction affecting the mis-splicing of *Md-tra* were responsible for the latitudinal distribution of

Y^M and III^M, we would expect more female-determining isoforms produced (i.e., misexpressed)

in Y^M males raised at a high temperature, or a higher expression level of female-determining

isoforms in III^M males raised at a low temperature. In contrast to that expectation, the G×T

interactions are not in the directions consistent with misexpression at discordant temperatures

(Supplementary Figure S11 and S12). An analysis of *Md-tra* splicing with qPCR was not

possible because we could not design primers that specifically amplified isoforms for

quantitative assessment.

We further tested if G×T interactions affect the expression and splicing of two direct

downstream targets of *Md-tra* in the sex determination pathway, *Md-dsx* and *Md-fru*. Our

RNA-seq data show that there is no effect of G×T interactions in the expression of *Md-dsx* or

Md-fru in all male heads (Supplementary Table 4), young male heads (Supplementary Table 5),

or testes (Supplementary Table 6). We also found no evidence of G×T interactions affecting the

expression of individual *Md-dsx* exons (Supplementary Figure S13). We did not test for G×T

effects on the expression of *Md-fru* exons because exons that differentiate the male and female

isoforms have not been annotated in the reference genome (Meier *et al.* 2013; Scott *et al.*

2014).

We also used qRT-PCR to examine the expression of the house fly male-determining

gene, *Mdmd*, in two III^M strains and two Y^M strains raised at 18°C and 27°C (Supplementary

Figure S14). One Y^M strain and one III^M strain originated from North America, and the other Y^M

strain and III^M strain came from Europe. If temperature-dependent differential expression of

Mdmd were responsible for the clinal distribution of Y^M and III^M (with higher expression

conferring a fitness advantage), we would expect higher *Mdmd* expression in Y^M (III^M) males at

lower (higher) temperatures. There is a significant G×T interaction affecting the expression of

Mdmd in the European Y^M and III^M strains, with higher *Mdmd* expression in III^M males at lower

temperatures (Supplementary Figure S14). This is the opposite pattern from what would be

expected if the hypothesized G×T effects on *Mdmd* expression were responsible for maintaining

the cline. A similar trend is observed in the North American strains, although the interaction term

is not significant. We observe these similar patterns in both population samples even though

they were assayed with two different types of tissue (abdomen in the North American strains,

and whole fly in the European strains), demonstrating that these results are robust to the tissues

we sampled. We also did not find a significant G×T interaction affecting expression of *Md-ncm*

(the ancestral paralog of *Mdmd*), which is not part of the sex determination pathway

(Supplementary Figure S14). Therefore, there is no evidence that *Mdmd* expression is

increased at the hypothesized favored temperatures for Y^M and III^M males.

G×T interactions on gene expression are not driven by cis-regulatory divergence

We next tested if divergence of *cis*-regulatory sequences between the III^M and standard

third chromosome is responsible for temperature-dependent expression differences between III^M

and Y^M males. III^M males are heterozygous (III^M/III) whereas Y^M males are homozygous (III/III)

for a standard third chromosome. If *cis*-regulatory alleles on the third chromosome are

responsible for differential expression of third chromosome genes between III^M and Y^M males,

the III^M and III alleles of those genes should also be differentially expressed in III^M males. For

example, if a gene is more highly expressed in III^M males than Y^M males, the III^M allele of the

gene should be more highly expressed than the III allele in III^M males. The opposite would be

true if Y^M males have higher expression than III^M males. We used this logic to test if G×T

interactions on gene expression are the result of *cis*-regulatory divergence of third chromosome

genes between the III^M and III chromosomes. To do so, we asked if genes on the third

chromosome that are significantly differentially expressed in head or testis because of G×T

interactions have concordant differences in expression between the III^M and III allele in III^M

males.

Figure 5: G×T interactions affect allele-specific expression in house fly testes. Differences in sequencing coverage in III^M house fly males between III^M and III alleles at either 18°C (circles) or 29°C (squares) are shown for 12 house fly genes where there is a G×T effect on testis expression between Y^M and III^M males. Each circle or square represents the difference in normalized mapped reads in testis between the III^M and III alleles at a single variable site (SNP) within a gene. Circles show expression differences between alleles at 18°C, and squares show expression differences between alleles at 29°C. The small horizontal lines indicate the mean difference in coverage between alleles across all sites in each gene at each temperature. Error bars represent the standard error across all variable sites within a gene at each temperature. The gene names with asterisks along the x-axis are differentially expressed between the III^M and III alleles in the same direction as the differential expression in III^M males between 18°C and 29°C (Table 1).

To test for differences in allelic expression, we first identified 12 genes on the third chromosome with a significant G×T interaction affecting testis expression, at least one heterozygous SNP in III^M males, and homozygous at those SNP sites in Y^M males (Figure 5). We required the variants to be heterozygous in III^M males and homozygous in Y^M males because we are interested in expression differences between the III^M and III allele in III^M males. We assumed that the allele in common between III^M and Y^M males is found on the standard third chromosome, and the allele unique to III^M males is on the III^M chromosome. This assumption is reasonable because the Y^M and III^M flies that we used for our RNA-seq share the same genetic background, and therefore should have the same standard third chromosome. We quantified

the expression of the two alleles (III^M and III) based on allele-specific RNA-seq read coverage.

We asked if the difference in expression of III^M alleles in each gene is consistent with the

difference in overall expression of these genes between 18°C and 29°C within III^M males. For

example, if III^M males have higher expression at 29°C, we expect the difference between the III^M

and III alleles to be greater at 29°C than 18°C.

We first compared the expression of III^M and III alleles in testis. Of the 12 genes with

significant G×T effects and the requisite SNPs to test for allele-specific expression, seven have

a significant effect of temperature on testis gene expression within III^M males (Figure 5; Table 1).

Of those seven genes, five have a pattern of allelic expression consistent with the differential

expression between 18°C and 29°C within III^M males: *LOC101892094* (homologous to

D. melanogaster Pdfr, which is responsible for regulating circadian behaviors), *LOC101891589*

(homologous to *D. melanogaster CG42450*, which is predicted to be involved in G

protein-coupled receptor signaling), *LOC101893483* (encoding a GATA zinc finger

domain-containing protein), *LOC101891938* (homologous to *D. melanogaster mmd*, which is

predicted to encode a membrane protein involved in ectodomain proteolysis), and

LOC101897352 (the cystathionine β-synthase gene associated with lifespan, mentioned

earlier). The two genes with allelic expression that is inconsistent with temperature-dependent

expression in III^M males are *LOC101882943* (homologous to *D. melanogaster Nepl5*) and

LOC101900427 (homologous to *D. melanogaster fne*). The remaining five genes do not differ in

testis expression between III^M males raised at 18°C and 29°C (Table 1).

Table 1: Temperature-dependent allele-specific expression in testis.

	Chromosome III	Rest of genome
Genes with significant temperature effect on III^M and III^M-III in right direction	5	7
Genes with significant temperature effect on III^M and incorrect direction of III^M-III	2	6
Genes with heterozygous sites in III^M males, but without a significant temperature effect on expression in III^M males	5	17

To determine a null expectation for the proportion of genes with allelic expression

consistent with the differential expression between 18°C and 29°C, we tested for concordance

between allele-specific expression and temperature-dependent expression differences for

genes on other chromosomes. We do not expect concordance for genes on other chromosomes

because the inbred Y^M and III^M males used in our RNA-seq experiment share a common genetic

background. We identified 30 genes on other chromosomes with heterozygous sites whose

testis expression depends on the G×T interaction (Table 1). Of those 30 genes, 13 are

differentially expressed between III^M males raised at 18°C and 29°C. We find that 7 out of the 13

genes in the rest of the genome have allele-specific expression that is consistent with the 18°C

vs 29°C expression differences (Table 1). There is not a significant excess of genes on the third

chromosome whose temperature-dependent expression is consistent with changes in

allele-specific expression relative to the rest of the genome (P = 0.64 in Fisher's exact test).

This suggests that the G×T effects on the expression of genes on the third chromosome is not

the result of an excess of *cis*-regulatory differences between the III^M and standard third

chromosomes.

When we analyzed only the younger male head samples, we found 7 genes on the third

chromosome with a significant G×T interaction that also had at least one SNP in III^M males.

Among them, one gene (*LOC101890343*, homologous to *D. melanogaster mahe*, encoding an

ATP-dependent RNA helicase) had a significant effect of temperature on gene expression within

III^M males. The allele specific expression of this gene is consistent with the temperature effect in

III^M males, but there are no genes on other chromosomes with the requisite SNPs in our head

RNA-seq data to test for a significant excess relative to a null expectation. When analyzing all

head samples, we found a single gene on the third chromosome with a significant GxT

interaction that also had a SNP in III^M males. However, we did not find a significant effect of

temperature on expression of this gene within III^M males.

We are limited in the analysis we can perform on allele-specific expression of genes on

the X vs Y^M chromosomes because of small sample sizes. There are only 40 genes assigned to

the house fly X or Y^M chromosome (Meisel & Scott 2018), none of which have a significant G×T

interaction affecting expression in testis (Supplementary Table 3). Only one X or Y^M

chromosome gene has a significant G×T interaction affecting expression in heads when we

analyze all samples (Supplementary Table 2), and it does not have any heterozygous sites.

Similarly, none of the three genes on the X chromosome with a significant G×T interaction
affecting expression in young male heads has any heterozygous sites.

Discussion

We tested for G×T interactions that affect gene expression in Y^M and III^M house fly
males. These G×T effects could lead to differences in temperature-dependent phenotypes
between house fly genotypes, thereby maintaining polygenic sex determination across
latitudinal clines based on temperature-dependent fitness effects of the proto-Y chromosomes.
We used RNA-seq to compare gene expression in heads and testes of two nearly isogenic
strains that differ only in their proto-Y chromosomes (Y^M or III^M) that we raised at two different
temperatures (18°C and 29°C). This 2×2 full factorial design allowed us to compare
genome-wide expression between four G×T combinations, which we combined with targeted
expression measurements of the male-determining gene (*Mdmd*) using qRT-PCR. We found
that G×T interactions lead to differential gene expression in both head and testis, but the
expression of genes involved in the sex determination pathway is not meaningfully affected by
those G×T interactions. We therefore hypothesize that alleles present on either the III^M
chromosome or the Y^M chromosomes, other than *Mdmd*, may be targets of selection.

No evidence that G×T interactions affect the sex determination pathway in a way that
explains the maintenance of polygenic sex determination

Our results suggest that G×T interactions affecting the sex determination pathway do not
explain the maintenance of polygenic sex determination in house fly. Evolutionary transitions
between heritable and temperature-dependent sex determination systems are possible if sex
determination pathways are temperature sensitive (Shine *et al.* 2002; Quinn *et al.* 2007; Radder
et al. 2008; Holleley *et al.* 2015). Sex determination in flies operates by alternative splicing of
multiple genes in the pathway (Salz 2011; Bopp *et al.* 2014). Temperature dependent alternative
splicing has been reported in *Arabidopsis* (Streitner *et al.* 2013; Steffen & Staiger 2017),
Neurospora (Colot *et al.* 2005), *Drosophila* (Jakšić & Schlötterer 2016; Martin Anduaga *et al.*
2019), and mammals (Preußner *et al.* 2017). It is therefore possible that temperature-sensitive
expression or splicing of sex determination factors can establish a clinal distribution of sex
determination genes, such as what is observed in house fly (Schenkel 2021). We did not find
evidence for G×T interactions affecting the expression of the male-determining *Mdmd* gene or
splicing of *Md-tra* in a way that is consistent with the clinal distribution of Y^M and III^M. In addition,

the expression of *Md-dsx* and *Md-fru*, the immediate downstream targets of *Md-tra*, do not

depend on G×T interactions.

It is possible that temperature affects the expression or splicing of sex determination

pathway genes earlier in development than we measured. For example, *Mdmd* expression level

might be more critical during early embryogenesis when *Md-tra* needs to be locked into a male

or female mode of splicing (Sharma *et al.* 2017). Hediger et al. (2010) have shown that the

Md-tra auto-regulatory loop can be effectively shut down in embryos by RNA interference, and

male development proceeds normally without the need of *Mdmd* expression. Similarly, when

Mdmd was removed from *Mdmd*-/+ cells at embryonic stages, the resulting clones developed as

males despite their female genotype (Hilfiker-Kleiner *et al.* 1993). Thus the adult *Mdmd* and

Md-tra expression we observed might not reflect the critical early expression levels. Additional

work is required to further examine temperature-dependent effects on the expression or splicing

of *Mdmd* or *Md-tra* across male genotypes in embryos, larvae, or pupae, rather than in adults.

Even though we did not observe differential expression of *Mdmd* that is consistent with

our hypothesis for the clinal distribution of Y^M and III^M males, we believe that the increased

expression of *Mdmd* in III^M males that we observe at the lower temperature is intriguing. It is

possible that *Mdmd* expression is optimal at an intermediate level between high and low

extremes—lower expression of *Mdmd* might be insufficient for *Md-tra* splicing, whereas higher

expression of *Mdmd* might be toxic because of its proposed role in antagonizing functions of the

generic splicing factor *Md-ncm* (Sharma *et al.* 2017). The increased expression of *Mdmd* in III^M

males at a lower temperature might thus explain the absence of III^M males in northern latitudes.

Moreover, Hediger et al. (1998) found male determining regions on both arms of the Y^M

chromosome that act additively. However, it is not yet resolved whether *Mdmd* is the male

determining factor on both of these arms or only one arm (Sharma *et al.* 2017). Additional work

is required to determine if there is an additional male determining gene other than *Mdmd* on the

Y^M chromosome that may have temperature dependent activity.

Temperature-dependent gene expression is not the result of large-scale cis-regulatory

changes on the III^M chromosome

Previous RNA-seq experiments (Meisel *et al.* 2015; Son *et al.* 2019), as well as the

results presented here (Figure 2A), provide consistent evidence that the third chromosome is

enriched for genes that are differentially expressed between Y^M and III^M males. This is expected

as the comparisons are between flies that differ in their third chromosome genotypes, and it

suggests there are *cis*-regulatory effects on the expression of genes on the third chromosome.

Consistent with this hypothesis, we observe a more pronounced clustering by genotype in our

PCA when we consider only chromosome III genes (Supplementary Figure S4). In contrast, we

find that genes that are differentially expressed because of temperature are not enriched on the

third chromosome in III^M males (Figure 2B). This is not because of lack of power to detect the

enrichment as we see a modest enrichment of differentially expressed third chromosome genes

in young Y^M male heads (Supplementary Figure 8). We also found that G×T interactions

affecting the expression of genes in male heads or testes are not enriched on the third

chromosome either (Supplementary Figure S9).

The lack of an enrichment of genes with temperature-dependent expression in III^M males

on the third chromosome suggests that temperature-dependent effects of the III^M chromosome

are not mediated by large-scale *cis*-regulatory changes across the III^M chromosome. Consistent

with this interpretation, there is not an enrichment of third chromosome genes with

temperature-dependent expression differences between the III^M and III alleles (Figure 5, Table

1). Moreover, an independent analysis of other RNA-seq data also found that there is not an

excess of expression differences between III^M and III alleles in a different house fly strain (Son &

Meisel 2021). We cannot perform a similar statistical analysis of Y^M genes because of the small

number of genes on that chromosome.

Temperature-dependent gene expression and the maintenance of polygenic sex

determination in house fly

We hypothesized that targets of selection responsible for the maintenance of polygenic

sex determination in house fly could be differentially expressed across proto-Y chromosome

genotypes and developmental temperatures. Despite our conclusion that a large number of

cis-regulatory variants on the III^M chromosome cannot explain the effect of the III^M chromosome

on temperature-dependent phenotypes, we still found evidence for temperature-dependent

effects of the III^M and Y^M chromosomes that could explain their divergent phenotypic effects.

First, there is some clustering by G×T combinations in the transcriptome-wide testis gene

expression profiles (Figure 1C). Second, we identify substantial temperature-dependent gene

expression (Figure 2B) and many genes whose expression depend on G×T interactions

(Figures 3 and 4). These temperature-dependent effects on expression could be responsible for

phenotypic differences between Y^M and III^M males, which could in turn provide a substrate upon

which selection acts to maintain the Y^M-III^M clines. If wide-spread *cis*-regulatory differences

across proto-sex chromosomes are not responsible for these G×T effects (as we hypothesize

above), then it is possible that a small number of loci on the proto-Y chromosomes act as

temperature-dependent *trans* regulators of gene expression across the entire genome.

Reproductive traits are a promising target of selection that could depend on G×T

interactions. There are more genes differentially expressed in testis because of G×T interactions

than in head, consistent with previous work that identified more differentially expressed genes in

testis than head between Y^M and III^M males (Meisel *et al.* 2015). Genes associated with

reproductive functions (*LOC101887804*, *LOC101899763*, *LOC101894442*, and

LOC101896699) were amongst the genes whose testis expression depend on G×T effects

(Figure 4). It is therefore possible that selection along the Y^M-III^M cline acts on reproductive

traits, which is consistent with the idea that the strength of sexual selection can vary across

populations (Arnqvist 1992; Payne & Krakauer 1997; Blanckenhorn *et al.* 2006; Connallon 2015;

Allen *et al.* 2017). These reproductive traits, or other variants under selection, could have

sexually antagonistic fitness effects (i.e., opposing fitness effects in males and females) which

may be temperature-sensitive. Sexual antagonism is one of the few selection pressures capable

of maintaining polygenic sex determination (Rice 1986; van Doorn & Kirkpatrick 2007).

Population genetic modeling also predicts that sexually antagonistic effects of Y^M and III^M can

maintain polygenic sex determination within house fly populations (Meisel *et al.* 2016; Meisel

2021), possibly in conjunction with epistatic interactions between either Y^M or III^M and autosomal

loci not linked to either *Mdmd* locus (Schenkel 2021). It is worth pursuing if sexual antagonism

can maintain polygenic sex determination by acting on temperature-dependent gene expression

differences between Y^M and III^M males.

Energy metabolism is a potential phenotype upon which selection acts to affect

reproductive functions. We previously found divergence between III^M and standard third

chromosome sequences surrounding genes encoding mitochondrial proteins (Son & Meisel

2021). Here, we report G×T interactions affecting the testis expression of three genes with

metabolic functions (*LOC101901027*, *LOC101901154*, and *LOC109613297*). All three genes

are upregulated in Y^M males at 18°C, and, to a lesser extent, upregulated in III^M males at 29°C

(Figure 4). None of the *D. melanogaster* homologs of these genes are differentially expressed

between flies raised at high (21.5°C) or low (6°C) temperatures (MacMillan *et al.* 2016), nor are

they differentially expressed between *D. melanogaster* that are evolved in hot or cold laboratory

environments (Hsu *et al.* 2020). However, one of the metabolic genes (*LOC101901154*),

encoding an aldehyde oxidase, has a *D. melanogaster* homolog (*AOX4*) that is expressed

higher at 21°C than 29°C (Zhao *et al.* 2015), consistent with the higher expression of the house

fly gene in Y^M males at lower temperatures. We are cautious to interpret further because there

are four tandemly arrayed *AOX* genes in the *D. melanogaster* genome and at least 3

corresponding genes in house fly; it is therefore not possible to assign orthology across this

family.

One of the other metabolic genes (*LOC101901027*) has a homolog (*fbp*) that is

expressed higher in *D. melanogaster* raised at 29°C than those raised at 21°C, regardless of

whether the flies come from Maine (USA) or Panama (Zhao *et al.* 2015). This is consistent with

the higher expression of this gene at 29°C in III^M testes, but opposite from the lower expression

at 29°C in Y^M testes (Figure 4). It is possible that the Y^M chromosome confers a fitness

advantage via increased production of fructose-1,6-bisphosphatase in testes at lower

temperatures. Consistent with this hypothesis, fructose-1,6-bisphosphatase is necessary for

cold-stress in mice (Park *et al.* 2020) and associated with cold hardiness in plants and insects

(Storey & Storey 2012; Cai *et al.* 2018). There is also evidence that *D. melanogaster fbp* is

differentially *trans*-regulated across genotypes and temperatures (Chen *et al.* 2015). In house

fly, this gene is not found on either the Y^M or III^M chromosome, which would require it to be

differentially regulated in *trans*, consistent with what is observed in *D. melanogaster*. Together

with the other differentially expressed metabolic genes, our results suggest that energy

metabolism related to spermatogenesis or sperm function may be a target of selection driving

the evolution of the III^M and Y^M chromosomes.

Muscle performance might also be under differential selection across the Y^M-III^M cline.

We identified three muscle component related genes (*LOC101893720, LOC101895658*, and

LOC101901052) upregulated in Y^M male heads at 18°C (Figure 3B). One of these genes

(*LOC101893720*) is homologous to *D. melanogaster bt*. Knockdown of *bt* decreases sarcomere

length and reduces climbing ability in *D. melanogaster* (Perkins & Tanentzapf 2014). Another

muscle-related gene (*LOC101895658*) is homologous to *D. melanogaster Unc-89*, which

encodes an obscurin protein. Reduced expression of *Unc-89* using P-element insertion results

in flightless adults in *D. melanogaster* (Katzemich *et al.* 2012). Upregulation of these genes in

Y^M males at lower temperatures might improve muscle performance.

We also find evidence that selection may have acted in response to thermal stress

across environments along the Y^M-III^M cline. A gene (*LOC101893129*) homologous to

D. melanogaster Nlaz, which encodes an extracellular lipid binding protein (similar to

apolipoprotein D and Retinol Binding Protein 4), is upregulated in heads of III^M males at the high

temperature (Figure 3A). *Nlaz* is regulated by the JNK signalling pathway to confer stress and
starvation tolerance, and it reduces oxidative stress by maintaining metabolic homeostasis
(Hull-Thompson *et al.* 2009). *NLaz* mutants in *D. melanogaster* have reduced stress resistance
and shorter lifespans, while over-expressing *NLaz* increases stress tolerance and extends
lifespan. *Nlaz* is also upregulated at extreme low temperature in *D. melanogaster* (Chen *et al.*
2015; MacMillan *et al.* 2016). Upregulation of this gene may therefore help III[M] males tolerate
thermal stress at high temperatures. Our results demonstrate the utility of simultaneously
studying the effects of both genotypic and temperature variation to determine how thermal
stress affects gene expression (Rivera *et al.* 2021).

There is also evidence that improved response to thermal stress may act to increase
lifespan in Y[M] and III[M] males at temperatures concordant with their clinal distribution. For
example, *LOC101897352* encodes cystathionine β-synthase and is homologous to
D. melanogaster Cbs. In *D. melanogaster*, *Cbs* is involved in ER stress response (Chow *et al.*
2013) and is a positive regulator of lifespan (Kabil *et al.* 2011). *LOC101897352* is upregulated in
Y[M] male testes at 18°C (Figure 4), consistent with longer lifespan for Y[M] males at lower
temperatures. *LOC101897352* (the *Cbs* homolog) is also one of the genes with a consistent
direction of allele-specific expression and expression difference between III[M] males at 18°C and
29°C (Figure 5), providing evidence that a *cis*-regulatory allele on the III[M] chromosome drives
temperature-dependent expression of a gene with a potential phenotypic effect. Future work
should aim to identify *cis*-regulatory regions underlying the temperature-dependent expression
differences between the III[M] and III alleles in *LOC101897352* (the *Cbs* homolog) and other such
genes on the third chromosome (Figure 5). Searching for such regulatory sequences in house
fly is currently impeded by the lack of a chromosome-scale genome assembly and
comprehensive gene annotations (Scott *et al.* 2014; Meisel & Scott 2018).

Two other genes are differentially expressed in a way that is suggestive of
temperature-dependent lifespan differences between Y[M] and III[M] males. *LOC101895929* is
homologous to *D. melanogaster pnt*. Knockdown of *pnt* extends lifespan in *D. melanogaster*
(Dobson *et al.* 2019). Interestingly, we see downregulation of this gene in III[M] male testes 29°C
(Figure 4), consistent with longer lifespan for III[M] males at a higher temperature. Lastly,
LOC101901283 is homologous to *D. melanogaster Calx*, which is associated with response to
ER stress (Chow *et al.* 2013). *Calx* mutation reduces *D. melanogaster* lifespan (Mok *et al.*
2020). *LOC101901283* is upregulated in III[M] male heads at 29°C (Figure 3B), suggesting a
longer lifespan for III[M] males at a higher temperature. All three lifespan-related genes

(*LOC101897352*, *LOC101895929*, and *LOC101901283*) therefore have expression profiles

consistent with longer lifespan of Y^M males at lower temperatures or III^M males at higher

temperatures, suggesting that temperature-dependent senescence might be a phenotype under

differential selection between III^M and Y^M males. It remains to be tested if these male genotypes

have different lifespans across temperatures.

Conclusion

The clinal distribution of the house fly proto-Y chromosomes in natural populations hints

at a possible G×T interaction involved in maintaining polygenic sex determination (Hiroyoshi

1964; Mcdonald *et al.* 1975; Denholm *et al.* 1986; Hamm *et al.* 2005; Feldmeyer *et al.* 2008;

Kozielska *et al.* 2008). We did not find evidence that temperature-dependent expression or

splicing of genes in the sex determination pathway explain the maintenance of polygenic sex

determination in house fly. However, such effects may act earlier in development than we

assayed. In contrast, G×T interactions affect gene expression in both somatic and reproductive

tissues across the entire genome. Our results therefore suggest that alleles on the proto-Y

chromosomes other than the male-determining *Mdmd* gene are targets of selection responsible

for maintaining the proto-Y chromosome clines in house fly.

There is no enrichment of G×T effects on the expression of genes on the proto-Y

chromosomes, suggesting that temperature-dependent expression differences between Y^M and

III^M males (and thereby phenotypic and fitness effects of the proto-Y chromosomes) are not

driven by a large-number of *cis*-regulatory changes on the III^M chromosome. Instead, if

temperature-dependent gene expression is responsible for temperature-dependent phenotypic

effects of the III^M and Y^M proto-Y chromosomes, those effects are the result of a small number of

alleles on the III^M (and possibly Y^M) chromosome. One such sex-linked gene, encoding

cystathionine β-synthase, is differentially expressed across genotypes and temperatures in a

way that is consistent with divergence in *cis*-regulatory sequences between the III^M and III

chromosomes. However, most of the differentially expressed genes between Y^M and III^M males

are not sex-linked and therefore likely the result of *trans* G×T effects on gene expression across

the entire genome. This is consistent with our previous work that identified very few differentially

expressed genes as a result of differences in proto-Y chromosome genotypes (Son *et al.* 2019).

Autosomal genes whose expression depends on proto-Y genotype and temperature include

those encoding metabolic proteins in testis, proteins involved in stress response in head, or with

effects on aging. This suggests temperature-dependent sperm function or thermal stress

tolerance may be targets of selection maintaining the Y^M-III^M cline via *trans*-effects of the proto-Y

chromosomes on expression of genes affecting these phenotypes across the genome.

Acknowledgements

855   This work was supported by the National Science Foundation (OISE-1444220 and

856 DEB-1845686 to RPM), Mindlin Foundation (MF16-US04 to JJ), and the University of Houston

857 (start up funds to RPM and Provost's Undergraduate Research Scholarship to JJ). We thank

858 Dalia Aldin and Ashley Lee for assistance establishing the Y^M and III^M strains with a common

859 genetic background.

Data Accessibility

861   RNA-seq data were collected for this manuscript and have been deposited in the NCBI

862 Gene Expression Omnibus under accession GSE136188 (BioProject PRJNA561541, SRA

863 accession SRP219410).

References cited

Allen SL, Bonduriansky R, Sgro CM, Chenoweth SF (2017) Sex-biased transcriptome

divergence along a latitudinal gradient. *Mol. Ecol.*, **26**, 1256–1272.

Anders S, Pyl PT, Huber W (2015) HTSeq--a Python framework to work with high-throughput

sequencing data. *Bioinformatics* , **31**, 166–169.

Arnqvist G (1992) Spatial variation in selective regimes: sexual selection in the water strider,

Gerris odontogaster. *Evolution*, **46**, 914–929.

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models using lme4.

J. Stat. Softw., **67**, 1–48.

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful

approach to multiple testing. *J. R. Stat. Soc. Series B Stat. Methodol.*, **57**, 289–300.

Beukeboom LW, Perrin N (2014) *The Evolution of Sex Determination*. Oxford University Press.

Blanckenhorn WU, Stillwell RC, Young KA, Fox CW, Ashton KG (2006) When Rensch meets

Bergmann: does sexual size dimorphism change systematically with latitude? *Evolution*, **60**,

2004–2011.

Bopp D, Saccone G, Beye M (2014) Sex determination in insects: variations on a common

theme. *Sex Dev.*, **8**, 20–28.

Cai B, Li Q, Liu F, Bi H, Ai X (2018) Decreasing fructose-1,6-bisphosphate aldolase activity

reduces plant growth and tolerance to chilling stress in tomato seedlings. *Physiol. Plant.*,

163, 247–258.

Chen J, Nolte V, Schlötterer C (2015) Temperature stress mediates decanalization and

dominance of gene expression in *Drosophila melanogaster*. *PLoS Genet.*, **11**, e1004883.

Chow CY, Wolfner MF, Clark AG (2013) Using natural variation in *Drosophila* to discover

previously unknown endoplasmic reticulum stress genes. *Proc. Natl. Acad. Sci. U. S. A.*,

110, 9013–9018.

Colot HV, Loros JJ, Dunlap JC (2005) Temperature-modulated alternative splicing and promoter

use in the circadian clock gene frequency. *Mol. Biol. Cell*, **16**, 5563–5571.

Connallon T (2015) The geography of sex-specific selection, local adaptation, and sexual

dimorphism. *Evolution*, **69**, 2333–2344.

Denholm I, Franco MG, Rubini PG, Vecchi M (1986) Geographical variation in house-fly (*Musca*

domestica L.) sex determinants within the British Isles. *Genet. Res.*, **47**, 19–27.

Dobin A, Davis CA, Schlesinger F *et al.* (2013) STAR: ultrafast universal RNA-seq aligner.

Bioinformatics , **29**, 15–21.

Dobson AJ, Boulton-McDonald R, Houchou L *et al.* (2019) Longevity is determined by ETS

transcription factors in multiple tissues and diverse species. *PLoS Genet.*, **15**, e1008212.

van Doorn GS (2014) Patterns and mechanisms of evolutionary transitions between genetic

sex-determining systems. *Cold Spring Harb. Perspect. Biol.*, a017681.

van Doorn GS, Kirkpatrick M (2007) Turnover of sex chromosomes induced by sexual conflict.

Nature, **449**, 909–912.

van Doorn GS, Kirkpatrick M (2010) Transitions between male and female heterogamety

caused by sex-antagonistic selection. *Genetics*, **186**, 629–645.

Feldmeyer B, Kozielska M, Kuijper B *et al.* (2008) Climatic variation and the geographical

distribution of sex-determining mechanisms in the housefly. *Evol. Ecol. Res.*, **10**, 797–809.

Gish W, States DJ (1993) Identification of protein coding regions by database similarity search.

Nat. Genet., **3**, 266–272.

Goodfellow PN, Lovell-Badge R (1993) *SRY* and sex determination in mammals. *Annu. Rev.*

Genet., **27**, 71–92.

Hamm RL, Gao J-R, Lin GG-H, Scott JG (2009) Selective advantage for III^{M} males over Y^{M}

males in cage competition, mating competition, and pupal emergence in *Musca domestica*

L. (Diptera: Muscidae). *Environ. Entomol.*, **38**, 499–504.

Hamm RL, Meisel RP, Scott JG (2015) The evolving puzzle of autosomal versus Y-linked male

determination in *Musca domestica*. *G3*, **5**, 371–384.

Hamm RL, Scott JG (2008) Changes in the frequency of YM versus IIIM in the housefly, Musca

domestica L., under field and laboratory conditions. *Genet. Res.*, **90**, 493–498.

Hamm RL, Scott JG (2009) A high frequency of male determining factors in male Musca

domestica (Diptera: Muscidae) from Ipswich, Australia. *J. Med. Entomol.*, **46**, 169–172.

Hamm RL, Shono T, Scott JG (2005) A cline in frequency of autosomal males is not associated

with insecticide resistance in house fly (Diptera: Muscidae). *J. Econ. Entomol.*, **98**,

171–176.

Hediger M, Burghardt G, Siegenthaler C *et al.* (2004) Sex determination in *Drosophila*

melanogaster and *Musca domestica* converges at the level of the terminal regulator

doublesex. *Dev. Genes Evol.*, **214**, 29–42.

Hediger M, Henggeler C, Meier N *et al.* (2010) Molecular characterization of the key switch *F*

provides a basis for understanding the rapid divergence of the sex-determining pathway in

the housefly. *Genetics*, **184**, 155–170.

Hediger M, Minet AD, Niessen M *et al.* (1998) The male-determining activity on the Y

chromosome of the housefly (*Musca domestica* L.) consists of separable elements.

Genetics, **150**, 651–661.

Hilfiker-Kleiner D, Dübendorfer A, Hilfiker A, Nöthiger R (1993) Developmental analysis of two

sex-determining genes, M and F, in the housefly, *Musca domestica*. *Genetics*, **134**,

1187–1194.

Hiroyoshi T (1964) Sex-limited inheritance and abnormal sex ratio in strains of the housefly.

Genetics, **50**, 373.

Hoffmann AA, Anderson A, Hallas R (2002) Opposing clines for high and low temperature resistance in *Drosophila melanogaster*. *Ecology letters*, **5**, 614–618.

Holleley CE, O'Meally D, Sarre SD *et al.* (2015) Sex reversal triggers the rapid transition from genetic to temperature-dependent sex. *Nature*, **523**, 79–82.

Hsu S-K, Belmouaden C, Nolte V, Schlötterer C (2020) Parallel gene expression evolution in natural and laboratory evolved populations. *Mol. Ecol.*, **30**, 884–894.

Huang DW, Sherman BT, Lempicki RA (2009a) Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat. Protoc.*, **4**, 44–57.

Huang DW, Sherman BT, Lempicki RA (2009b) Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.*, **37**, 1–13.

Hull-Thompson J, Muffat J, Sanchez D *et al.* (2009) Control of metabolic homeostasis by stress signaling is mediated by the lipocalin NLaz. *PLoS Genet.*, **5**, e1000460.

Jakšić AM, Schlötterer C (2016) The interplay of temperature and genotype on patterns of alternative splicing in *Drosophila melanogaster*. *Genetics*, **204**, 315–325.

Kabil H, Kabil O, Banerjee R, Harshman LG, Pletcher SD (2011) Increased transsulfuration mediates longevity and dietary restriction in *Drosophila*. *Proc. Natl. Acad. Sci. U. S. A.*, **108**, 16831–16836.

Katzemich A, Kreisköther N, Alexandrovich A *et al.* (2012) The function of the M-line protein obscurin in controlling the symmetry of the sarcomere in the flight muscle of *Drosophila*. *J. Cell Sci.*, **125**, 3367–3379.

Kaufman PE, Nunez SC, Geden CJ, Scharf ME (2010) Selection for resistance to imidacloprid in the house fly (Diptera: Muscidae). *J. Econ. Entomol.*, **103**, 1937–1942.

Kim D, Langmead B, Salzberg SL (2015) HISAT: a fast spliced aligner with low memory

requirements. *Nat. Methods*, **12**, 357–360.

Knaus BJ, Grünwald NJ (2017) vcfr: a package to manipulate and visualize variant call format

data in R. *Mol. Ecol. Resour.*, **17**, 44–53.

Kozielska M, Feldmeyer B, Pen I, Weissing FJ, Beukeboom LW (2008) Are autosomal

sex-determining factors of the housefly (*Musca domestica*) spreading north? *Genet. Res.*,

90, 157–165.

Levine MT, Eckert ML, Begun DJ (2011) Whole-genome expression plasticity across tropical

and temperate Drosophila melanogaster populations from Eastern Australia. *Molecular*

biology and evolution, **28**, 249–256.

Li H, Handsaker B, Wysoker A *et al.* (2009) The Sequence Alignment/Map format and

SAMtools. *Bioinformatics* , **25**, 2078–2079.

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.*, **15**, 550.

MacMillan HA, Knee JM, Dennis AB *et al.* (2016) Cold acclimation wholly reorganizes the

Drosophila melanogaster transcriptome and metabolome. *Sci. Rep.*, **6**, 28999.

Martin Anduaga A, Evantal N, Patop IL *et al.* (2019) Thermosensitive alternative splicing senses

and mediates temperature adaptation in *Drosophila*. *eLife*, **8**, e44642.

McDonald I c., Evenson P, Nickel CA, Johnson OA (1978) House fly genetics: isolation of a

female determining factor on chromosome 4. *Ann. Entomol. Soc. Am.*, **71**, 692–694.

Mcdonald IC, Overland DE, Leopold RA *et al.* (1975) Genetics of house flies: variability studies

with North Dakota, Texas, and Florida populations. *J. Hered.*, **66**, 137–140.

McKenna A, Hanna M, Banks E *et al.* (2010) The Genome Analysis Toolkit: a MapReduce

framework for analyzing next-generation DNA sequencing data. *Genome Res.*, **20**,

1297–1303.

Meier N, Käppeli SC, Hediger Niessen M *et al.* (2013) Genetic control of courtship behavior in

the housefly: evidence for a conserved bifurcation of the sex-determining pathway. *PLoS*

One, **8**, e62476.

Meisel RP (2021) The maintenance of polygenic sex determination depends on the dominance

of fitness effects which are predictive of the role of sexual antagonism. *G3* , jkab149.

Meisel RP, Davey T, Son JH *et al.* (2016) Is multifactorial sex determination in the house fly,

Musca domestica (L.), stable over time? *J. Hered.*, **107**, 615–625.

Meisel RP, Gonzales CA, Luu H (2017) The house fly Y Chromosome is young and minimally

differentiated from its ancient X Chromosome partner. *Genome Res.*, **27**, 1417–1426.

Meisel RP, Scott JG (2018) Using genomic data to study insecticide resistance in the house fly,

Musca domestica. *Pestic. Biochem. Physiol.*, **151**, 76–81.

Meisel RP, Scott JG, Clark AG (2015) Transcriptome differences between alternative sex

determining genotypes in the house fly, *Musca domestica*. *Genome Biol. Evol.*, **7**,

2051–2061.

Mok J-W, Chung H, Choi K-W (2020) Calx, a sodium/calcium exchanger, may affect lifespan in

Drosophila melanogaster. *MicroPubl Biol*, **2020**.

Moore EC, Roberts RB (2013) Polygenic sex determination. *Curr. Biol.*, **23**, R510–2.

Oksanen J, Guillaume Blanchet F, Friendly M *et al.* (2019) vegan: Community Ecology Package.

Orzack SH, Sohn JJ, Kallman KD, Levin SA, Johnston R (1980) Maintenance of the three sex

chromosome polymorphism in the platyfish *Xiphophorus maculatus*. *Evolution; international*

journal of organic evolution, **34**, 663–672.

Park H-J, Jang HR, Park S-Y *et al.* (2020) The essential role of fructose-1,6-bisphosphatase 2

enzyme in thermal homeostasis upon cold stress. *Exp. Mol. Med.*, **52**, 485–496.

Payne RJH, Krakauer DC (1997) Sexual selection, space, and speciation. *Evolution*, **51**, 1–9.

Perkins AD, Tanentzapf G (2014) An ongoing role for structural sarcomeric components in maintaining *Drosophila melanogaster* muscle function and structure. *PLoS One*, **9**, e99362.

Preußner M, Goldammer G, Neumann A *et al.* (2017) Body temperature cycles control rhythmic alternative splicing in mammals. *Mol. Cell*, **67**, 433–446.e4.

Quinn AE, Georges A, Sarre SD *et al.* (2007) Temperature sex reversal implies sex gene dosage in a reptile. *Science*, **316**, 411.

Radder RS, Quinn AE, Georges A, Sarre SD, Shine R (2008) Genetic evidence for co-occurrence of chromosomal and thermal sex-determining systems in a lizard. *Biol. Lett.*, **4**, 176–178.

R Core Team (2019) R: A Language and Environment for Statistical Computing.

Rice WR (1986) On the instability of polygenic sex determination: the effect of sex-specific selection. *Evolution*, **40**, 633–639.

Rivera HE, Aichelman HE, Fifer JE *et al.* (2021) A framework for understanding gene expression plasticity and its influence on stress tolerance. *Molecular ecology*, **30**, 1381–1397.

Salz HK (2011) Sex determination in insects: a binary decision based on alternative splicing. *Curr. Opin. Genet. Dev.*, **21**, 395–400.

Schenkel MA (2021) Evolutionary genetics and dynamics of transitions in sex determination systems. University of Groningen.

Schmidt R, Hediger M, Roth S, Nöthiger R, Dübendorfer A (1997) The Y-chromosomal and autosomal male-determining M factors of *Musca domestica* are equivalent. *Genetics*, **147**, 271–280.

Scott JG, Sridhar P, Liu N (1996) Adult specific expression and induction of cytochrome P450lpr in house flies. *Arch. Insect Biochem. Physiol.*, **31**, 313–323.

Scott JG, Warren WC, Beukeboom LW *et al.* (2014) Genome of the house fly, *Musca domestica*

    L., a global vector of diseases with adaptations to a septic environment. *Genome Biol.*, **15**,

    466.

Sharma A, Heinze SD, Wu Y *et al.* (2017) Male sex in houseflies is determined by *Mdmd*, a

    paralog of the generic splice factor gene *CWC22*. *Science*, **356**, 642–645.

Shine R, Elphick MJ, Donnellan S (2002) Co-occurrence of multiple, supposedly incompatible

    modes of sex determination in a lizard population. *Ecol. Lett.*, **5**, 486–489.

Shono T, Scott JG (2003) Spinosad resistance in the housefly, *Musca domestica*, is due to a

    recessive factor on autosome 1. *Pestic. Biochem. Physiol.*, **75**, 1–7.

Sinclair AH, Berta P, Palmer MS *et al.* (1990) A gene from the human sex-determining region

    encodes a protein with homology to a conserved DNA-binding motif. *Nature*, **346**, 240–244.

Son JH, Kohlbrenner T, Heinze S *et al.* (2019) Minimal effects of proto-Y chromosomes on

    house fly gene expression in spite of evidence that selection maintains stable polygenic sex

    determination. *Genetics*, **213**, 313–327.

Son JH, Meisel RP (2021) Gene-level, but not chromosome-wide, divergence between a very

    young house fly proto-Y chromosome and its homologous proto-X chromosome. *Mol. Biol.*

    Evol., **38**, 606–618.

Steffen A, Staiger D (2017) Chromatin marks and ambient temperature-dependent flowering

    strike up a novel liaison. *Genome Biol.*, **18**, 119.

Stephens M (2016) False discovery rates: a new deal. *Biostatistics* , **18**, 275–294.

Stevenson KR, Coolon J, Wittkopp PJ (2013) Sources of bias in measures of allele-specific

    expression derived from RNA-seq data aligned to a single reference genome. *BMC*

    Genomics, **14**, 536.

Storey KB, Storey JM (2012) Insect cold hardiness: metabolic, gene, and protein adaptation.

Can. J. Zool., **90**, 456–475.

Streitner C, Simpson CG, Shaw P *et al.* (2013) Small changes in ambient temperature affect alternative splicing in *Arabidopsis thaliana*. *Plant Signal. Behav.*, **8**, e24638.

Zhao L, Wit J, Svetec N, Begun DJ (2015) Parallel gene expression differences between low and high latitude populations of *Drosophila melanogaster* and *D. simulans*. *PLoS Genet.*, **11**, e1005184.

Zimmer F, Harrison PW, Dessimoz C, Mank JE (2016) Compensation of dosage-sensitive genes on the chicken Z chromosome. *Genome Biol. Evol.*, **8**, 1233–1242.

 Supplementary Figures

 Supplementary Figure S1. PCA of the 500 most variable genes in A) all male heads, B) young
 male heads and C) testes.

1067 **Supplementary Figure S2**. PCA plots of gene expression levels in all male heads on
1068 Chromosome I (A), Chromosome II (B), Chromosome III (C), Chromosome IV (D), Chromosome
1069 V (E), and the X Chromosome (F).

Supplementary Figure S4. PCA plots of gene expression levels in testes on Chromosome I (A), Chromosome II (B), Chromosome III (C), Chromosome IV (D), Chromosome V (E), and the X Chromosome (F).

Supplementary Figure S5. Non-metric multidimensional scaling (NMDS) plots showing gene
expression profiles in all male heads (A), young male heads (B) and testes (C).

Supplementary Figure S6. Heatmaps and dendrograms showing hierarchical clustering in all
male heads (A), young male heads (B), and testes (C). Samples were compared using the
Euclidean distance between regularized log transformed read counts. Color (0-80) refers to the
euclidean distance between samples. Asterisks indicate older samples.

Supplementary Figure S7: MA plots showing differential expression of genes because of: genotype effect at 18°C in all male heads (A), young male heads (B), or testes (C); genotype effect at 29°C in all male heads (D), young male heads (E), or testes (F); temperature effect in III^M males in all male heads (G), young male heads (H), or testes (I); and temperature effect in Y^M males in all male heads (J), young male heads (K), or testes (L). The log fold change values were shrunk using the lfcShrink() function in DESeq2 with the adaptive shrinkage estimator (Stephens 2016). Each dot represents a gene. Red dots represent genes that are significantly differentially expressed (p<0.05).

Supplementary Figure S8. Proportion of genes that are differentially expressed in head on each chromosome when the two older samples were excluded. Comparisons are between genotypes at 18°C (A) and 29°C (B), as well as between temperatures in III^M males (C) and Y^M males (D). Each bar represents the proportion of differentially expressed genes on a chromosome, and dashed lines show the genome-wide average. Asterisks indicate P values obtained from Fisher's exact test comparing the number of differentially expressed genes on a chromosome, the number of non-differentially expressed genes on a chromosome, and the number of differentially and non-differentially expressed genes across all other chromosomes, after Bonferroni correction (*$P < 0.05$, **$P < 0.005$, ***$P < 0.0005$, ****$P < 0.00005$, *****$P < 0.000005$). When we exclude the two older III^M head samples, there is a modest enrichment of third chromosome genes that are significantly differentially expressed between temperatures in Y^M heads (Fisher's Exact Test, Odds ratio = 1.2, 95%CI: 1.01-1.42, $P = 0.03$).

Supplementary Figure S9: Venn diagrams show the number of significantly differentially expressed genes in testes vs all male heads (first column), testes vs young male heads (second column), and all male heads vs young male heads (third column). There is an excess of genes in the overlapping portion in all comparisons (*P*<0.05 using a *z*-test of proportions).

Supplementary Figure S10. No chromosomes are enriched for significantly differentially expressed genes as a result of G×T interactions in all male heads (A), young male heads (B), or testes (C). The dashed line represents the genome-wide proportion of significantly differentially expressed genes. The numbers within each bar represent the number of significantly differentially expressed genes on that chromosome. Not all genes with significant G×T effects on expression are assigned to chromosomes, and only genes assigned to chromosomes are plotted. The X chromosome has <100 genes, which is much less than the other chromosomes which have 1000s of genes. Therefore a single G×T interaction on the X chromosome appears as a large proportion of genes. However, the small total number of X chromosome genes means this large proportion is not a significant excess.

Supplementary Figure S11. (A) Schematic representation of the *Md-tra* locus based on DNA sequencing, cDNA clones, and RNA-Seq data (Hediger *et al.* 2010; Scott *et al.* 2014). Splicing of the female-determining transcript is illustrated by the red diagonal lines connecting exons, and exons that contain protein-coding sequence of the female-determining splice variant are in red. Exons found in the male isoforms are shown in blue. The start and stop codon locations are shown. The expression of *Md-tra* across all exons is shown for RNA-seq data collected from all male heads (B and D) and testis (C and E). Error bars represent standard error (most standard error estimates are smaller than the size of the points, and thus cannot be seen in the graph).

A.

B.

Supplementary Figure S12. Expression of *Md-tra* across all exons is shown for RNA-seq
data collected from young male heads. Error bars represent standard error (most standard
error estimates are smaller than the size of the points, and thus cannot be seen in the graph).
Exons that contain protein-coding sequence of the female-determining splice variant (E2a, E4,
E5,and E6) are in red. Exons found in the male isoforms (E2b and E3) are shown in blue.

Supplementary Figure S13. Read depth coverage of *Md-dsx* in all male heads (A) and testis
(B) in each G×T combination. Exons are shown along the x-axis. Exons in the
male-determining isoform are shown in blue, and exons in the female-determining isoform are
shown in red. Expression of *Md-dsx* is not significantly affected by G×T interactions in either
head or testis (Supplementary Tables 2 and 3). Usage of male-specific exon E_m (ANOVA, *P*
= 0.0004), exon E5 (ANOVA, *P* = 0.013), and the female specific exon E4 (ANOVA, *P* <
2.2e-16) are affected by G×T interactions in testis. However, these G×T interactions are not in
the directions expected if mis-splicing of *Md-dsx* is responsible for maintaining Y^M-III^M clines.
In head, usage of E_m (ANOVA, *P* < 2.2 e-16) and Exon 4 (ANOVA, *P* < 2.2 e-16) is affected
by G×T interactions. Similar to testis, effects of the G×T interactions in head are also not in the
directions expected if mis-splicing of *Md-dsx* is responsible for maintaining Y^M-III^M clines.

Supplementary Figure S14. Expression levels of *Mdmd* (A), and *Md-ncm* (B) in the abdomens
of CSkab (III^M) and IsoCS (Y^M), and expression levels of *Mdmd* (C), and *Md-ncm* (D) in the
whole body of SPA3 (III^M) and GK-1 (Y^M) males raised at 18°C and 27°C. Each data point is a
technical replicate that has been normalized by dividing by the control gene within that replicate,
and points with the same shape are from the same biological replicate. The horizontal line
indicates the median across all replicates.

Supplementary Figure S15. Normalized read counts in young male heads (excluding two older
head samples) for genes that are differentially expressed and have consistent expression
patterns with the analysis when the two older head samples are included (Figure 3A). *Nlaz* is
not significantly differentially expressed when only younger heads are considered (*P* =0.08),
however the pattern is consistent with the analysis of all heads. All other genes are significantly
differentially expressed. Error bars represent standard error of mean.

 Supplementary Tables

 Supplementary Table S1. Primers for qPCR

Gene	Forward Primer (5'-3')	Reverse Primer (5'-3')	Anneal
Mdmd	TGGTGCGCCCTTCTTTAAAC	GTTGACGCGGACAATCAACG	55°C
Md-ncm	TTCCGACTCTGAATCATCTGAC	GCACTCCTCATAATCCAAACTG	55°C
LOC101888902 XM_005187313 (control)	GTTGTGTCGCAAATATGGCTTG	ACCACTCATACGCTGCAAAAG	55°C

1155 Supplementary Table S2: RNA-seq reads from head mapped to the reference genome

Sample	Geno-type	Mapped reads	Total reads	% mapped reads	Uniquely mapped reads
CSrab_18C_head1	III^M	23,386,737	30,872,584	75.75	18,211,108
CSrab_18C_head2	III^M	36,498,365	48,036,045	75.98	28,405,569
CSrab_18C_head3 *	III^M	25,899,682	33,922,170	76.35	20,123,259
CSrab_29C_head1	III^M	29,078,715	35,935,602	80.92	24,626,862
CSrab_29C_head2	III^M	32,542,292	40,966,941	79.44	26,979,708
CSrab_29C_head3 *	III^M	45,976,216	60,550,809	75.93	35,694,337
IsoCS_18C_head1	Y^M	23,338,684	30,429,230	76.70	18,470,903
IsoCS_18C_head2	Y^M	23,593,266	30,809,521	76.58	18,657,918
IsoCS_18C_head3	Y^M	20,170,741	26,500,047	76.12	15,690,817
IsoCS_29C_head1	Y^M	30,777,222	40,045,778	76.86	24,278,617
IsoCS_29C_head2	Y^M	36,530,758	47,069,689	77.61	28,848,124
IsoCS_29C_head3	Y^M	29,711,982	38,668,539	76.84	23,352,024

1156 Asterisks indicate the two older sample

 Supplementary Table S3: RNA-seq reads from testis mapped to the reference genome

Sample	Geno-type	Mapped reads	Total reads	% mapped reads	Uniquely mapped reads
CSrab_18C_testis1	III^M	18,951,438	24,457,263	77.49	15,651,117
CSrab_18C_testis2	III^M	18,154,085	23,758,773	76.41	14,916,846
CSrab_18C_testis3	III^M	24,260,793	32,425,899	74.82	19,961,330
CSrab_29C_testis1	III^M	34,972,612	46,278,674	75.57	28,989,480
CSrab_29C_testis2	III^M	25,449,937	32,944,157	77.25	21,232,767
CSrab_29C_testis3	III^M	38,081,673	51,434,420	74.04	31,713,175
IsoCS_18C_testis1	Y^M	39,350,279	49,827,408	78.97	32,970,057
IsoCS_18C_testis2	Y^M	24,820,579	31,850,279	77.93	20,609,110
IsoCS_18C_testis3	Y^M	31,389,010	39,651,139	79.16	26,217,670
IsoCS_29C_testis1	Y^M	48,781,311	65,167,161	74.86	40,005,841
IsoCS_29C_testis2	Y^M	30,454,185	41,083,650	74.13	24,992,406
IsoCS_29C_testis3	Y^M	38,393,746	48,460,189	79.23	32,299,686

1158 Supplementary Table S4. DESeq2 result for all male heads. Data provided in a separate file.

1159 Supplementary Table S5. DESeq2 result for young male heads. Data provided in a separate file.

1160 Supplementary Table S6. DESeq2 result for testes. Data provided in a separate file.