Dynamic patterns of correlated activity in the prefrontal cortex encode information about social behavior

Nicholas A. Frost[1,3,4,5], Anna Haggart[1,3,4,5], and Vikaas S. Sohal[2,3,4,5]*

[1]Department of Neurology, [2]Department of Psychiatry, [3]Center for Integrative Neuroscience, [4]Weill Institute for Neurosciences, [5]Kavli Institute for Fundamental Neuroscience
University of California, San Francisco

*Correspondence to: vikaas.sohal@ucsf.edu

Short title: Changes in correlated activity in PFC neurons encode social behavior

ABSTRACT

New technologies have made it possible to measure activity from many neurons simultaneously.
Nevertheless, most studies still analyze the activity of simultaneously recorded neurons one-at-a-
time, then group together neurons which increase their activity during similar behaviors into an
'ensemble.' This notion of an ensemble ignores the ability of neurons to act collectively, and
encode and transmit information in ways that are not reflected by their individual activity levels.
We used microendoscopic GCaMP imaging to measure prefrontal activity while mice were
either alone or engaged in social interaction. We developed new approaches, using neural
network classifiers and surrogate (shuffled) datasets, to characterize how neurons synergistically
transmit information about social behavior. Surrogate datasets which preserve behaviorally-
specific patterns of coactivity (correlations) outperform those which preserve behaviorally-
driven changes in activity levels but not correlated activity. This shows that prefrontal neurons
act collectively to transmit information about socialization, because social behavior elicits
increases in correlated activity that are not explained simply by the activity levels of the
underlying neurons. Notably, this ability of correlated activity to enhance the information
transmitted by neuronal ensembles is lost in mice lacking the autism-associated gene Shank3.
These results show that synergy is an important concept for the coding of social behavior which
can be disrupted in disease states, reveal a specific mechanism underlying this synergy (social
behavior increases correlated activity within specific ensembles), and outlines methods for
studying how neurons within an ensemble can work together to encode information.

INTRODUCTION

Activity in different neurons often exhibits precise temporal relationships that are modulated by behavior (1–3). For example, subsets of neurons can exhibit correlated activity in which they become co-active at the same time or within short windows of time during specific behaviors. However, correlated activity can occur simply as a byproduct of the interconnected nature of neuronal networks (4). Thus it is unclear whether correlated activity contributes in a meaningful way to information encoding (5), or whether it is simply a reflection of (and thus redundant with) changes in other variables, e.g., firing rates (6).

Many experimental and theoretical studies have studied how noise correlations, e.g., correlations between the trial-to-trial variability of different neurons, affect information coding (7–9). From a population coding perspective, noise correlations can either impair or enhance decoding, depending on whether correlations are present between neurons that have similar or distinct tuning for a behavioral variable of interest. A different question, separate from that of whether variability (noise) between neurons is correlated on a trial-by-trial basis, is whether neurons exhibit correlations that differ based on a behavioral variable. For example, a group of neurons may exhibit correlated activity (co-activity that occurs more often than expected by chance) only during a specific behavior. Groups of co-active neurons represent an attractive computational unit for information processing because they should optimize temporal summation in downstream targets. Thus, even if a group of neurons do not increase their activity levels during a behavioral state, they could increase their correlated activity, and thereby more effectively activate a common downstream target. In this manner, behaviorally-driven changes in correlations can transmit information independent of changes in the activity levels of the underlying neurons. Whereas noise correlations, as their name suggests, reflect correlations in noise, i.e., variability that is unrelated to a behavioral variable of interest, behaviorally-modulated correlations represent potentially important signals in neural networks.

While multiple studies have shown that behavior can modulate correlations (3,10) the functional significance of this has remained unclear, because changes in correlations might simply reflect (and be redundant with) variation in activity levels (6) rather than contributing additional information. Even with the advent of new technologies for simultaneously recording from large numbers of neurons in behaving animals, many studies still focus on information that is transmitted by changes in the activity levels of neurons, while ignoring contributions from correlated activity or other ways in which neurons can act collectively. In particular, it is now commonplace to identify 'neuronal ensembles' by grouping together neurons which significantly increase or decrease their activity during the same behaviors or in response to the same stimuli. However, this approach ignores information that is transmitted collectively, e.g., it would not be sensitive to the hypothetical group of neurons described above, which increase their correlations during a specific behavior while keeping their activity levels constant. Thus current methods for identifying neuronal ensembles might falsely conclude that a group of neurons do not encode a behavioral variable (when they do encode it collectively), incorrectly estimate the amount of information that is being encoded, and/or miss important mechanisms that contribute to encoding.

Correlations have been shown to contribute additional information for small groups (3-8
neurons) of cortical neurons (11). Recently, a few studies have examined how correlations
contribute to encoding within larger cortical ensembles, which have greater potential for both
synergy and redundancy. One found that the identity of a conditioned stimulus was encoded in
mean activity levels, but not in moment-to-moment patterns of co-activity (12). In hippocampal
region CA1, disrupting correlations impairs the decoding of position, head direction and speed
(13). And during a two-choice discrimination task with probabilistic outcomes, correlations
between neurons in secondary motor region M2 enhance the encoding of the choice(14). Thus,
some studies have found evidence that correlations between neurons can augment the
information those neurons encode about behavioral variables(11,13,14). However, all of these
focused on the potential role of noise correlations in enhancing encoding by activity levels. I.e.,
they all hypothesize that noise correlations help disambiguate trial-by-trial variation in activity
levels that is related to behavioral variables of interest, from variation that is attributable to
'noise', e.g., changes in other variables such as internal states. They did not specifically examine
whether correlations might themselves encode unique information by increasing or decreasing in
relation to specific behavioral variables, creating behaviorally-specific patterns of co-activity.
Thus specific mechanisms through which cortical neurons can collectively encode information
currently remain poorly described. A further challenge is the lack of methods for identifying
neurons that contribute to the collective encoding of behavioral variables, but do so without
significantly increasing or decreasing their activity levels. This makes it difficult to directly
examine these neurons in order to assess whether they contribute to ensemble encoding by
altering their correlations, generating behaviorally-specific patterns of co-activity, etc.
103
The goal of this study is to address these two challenges related to neural encoding in the cortex
by focusing on a subject that has already been well-studied: the encoding of rodent socialization
by neurons in the medial prefrontal cortex (mPFC) (15–18). Many studies have already shown
that prefrontal neurons can significantly increase or decrease their activity levels during periods
of social interaction (2,17,18) and/or in response to social stimuli such as odors (19). However,
these studies have not assessed whether prefrontal neurons might act collectively to encode
additional information (beyond what activity levels encode), examined whether this might occur
via behaviorally-driven changes in correlated activity, or identified ensembles based on neurons'
ability to encode information collectively, rather than through increases or decreases in their
individual activity levels.
114
The goal of this study is to examine a neuronal population that is already known to encode social
behavior, and determine whether this encoding occurs solely via changes in the activity levels of
neurons vs. whether additional information is transmitted collectively, and if so, how this occurs.
These are critical questions, because if we do not understand the mechanisms which normally
encode information, we cannot study how they are disrupted in disease states. Thus by design,
we focused on these foundational questions about the nature of information transmission, rather
than trying to tackle more nuanced issues, e.g., does this encoding differ between subpopulations
of prefrontal neurons, etc.
123
We used microendoscopic GCAMP imaging in freely-moving mice to measure activity in
prefrontal ensembles during periods when mice were either alone or engaged in social behavior.
We used a neural network classifier to quantify how well prefrontal ensembles would transmit

information about behavior to downstream neurons. By examining the operation of this neural
network and using surrogate datasets which preserve activity levels but either preserve or disrupt
patterns of correlated activity, we find that behaviorally-driven changes in correlations enhance
the information transmitted by neuronal ensembles. Notably, this was not the case in a mouse
model of autism (Shank3 knockout mice), even though large numbers of prefrontal neurons are
still recruited by socialization in Shank3 knockout mice, and encode socialization through
changes in their activity levels. This illustrates that the ability of prefrontal neurons to
collectively encode information may be selectively disrupted in pathological states.

RESULTS

Social interaction recruits prefrontal ensembles

We implanted microendoscopes (nVoke; Inscopix) into the medial prefrontal cortex (mPFC) of adult wildtype C57/B6 mice (WT) to image calcium transients using GCaMP6f expressed under control of the human synapsin promotor. We imaged freely moving mice during an assay which sequentially introduced 2 novel juvenile mice to the home cage of the subject mouse, first during an initial (novel) epoch and then again during a subsequent (familiar) epoch. These four epochs of social interaction were interleaved with epochs during which the subject mouse was alone in its home cage ('home cage' epochs). The first 5 minutes of each interaction epoch was scored by a blinded observer, and each wild-type mouse spent approximately 10 minutes interacting with the juvenile mice (393 +/- 25 s during the novel epochs and 235 +/- 18 s during the familiar epochs, p = 0.00017, paired t test, n = 10 WT mice).

We processed data using a modified PCA/ICA approach (20,21) to identify neurons which were active during the imaging session. To minimize the influence of the surrounding neuropil on neuronal signals, we calculated the mean signal within each ROI, then subtracted the mean signal calculated from a circular annulus surrounding each ROI (Figure S1). Casual inspection of calcium traces revealed that some neurons were more active during epochs of social interaction (compared to periods of home cage exploration), whereas others exhibited the opposite pattern (**Figure 1A**). Correspondingly, aligning calcium traces to the onset of social interaction revealed many neurons that either increased or decreased their activity levels at the onset of interaction (**Figure 1B**). Fluorescence traces were converted to binary event rasters (see Methods for details of event detection), in which most neurons were "active" in less than 5% of frames (**Figure 1C**). As a population, imaged neurons were more active during social interaction (**Figure 1C**, n = 663 neurons from 10 mice, percent time active in home cage: 1.8% +/- 0.1, percent time active during social interaction: 2.1 +/- 0.1%, p = 0.00002, paired t-test). There was a bimodal distribution of cells that were significantly more (>90th percentile, social: 152/663 neurons, home cage: 80/663 neurons; p < 0.00001, Chi-Squared Test) or less active (<10th percentile, social: 128/663 neurons, home cage: 119/663 neurons; p = 0.5) during either social interaction or matched periods when mice were alone in their home cage, as compared to circularly shuffled datasets (**Figure 1D**). These correspond to conventionally-defined neuronal ensembles, i.e., groups of neurons that increased or decreased their activity levels during social interaction.

Using a neural network classifier to assess how well ensembles transmit information

Next, we sought to determine how well these prefrontal ensembles would transmit information about social behavior to downstream neurons. Specifically, we measured how well downstream neurons could decode whether a mouse was engaged in social behavior based on input from these prefrontal neurons. Later we will study how this was altered in Shank3 knockout mice. For this we used a simple neural network classifier that received input from the recorded neurons. Our rationale for using this kind of neural network classifier was threefold. First, a simple neural network measures information that is immediately and readily available to downstream neurons. Second, for a neural network with only one hidden layer, it is straightforward to examine the weights to determine how the network performs the classification. This can provide insight into exactly how the neural network is able to decode behavior from the input activity. Third,

180 examining how various parameters of the neural network affect its performance can provide
181 additional clues about how information is represented within the input population.
182
Figure 2A shows the design of the neural network classifier. For clarity, we use the terms
'neurons' specifically to refer to actual prefrontal neurons (which provide input to the neural
network), and 'units' to refer to simulated elements within the network. The network consisted of
a hidden layer containing 1000 units. We chose this number because it is both an order of
magnitude larger than the number of input neurons and an order of magnitude smaller than the
number of frames available for training (the latter helps ensure that there will be enough data to
train the output weights). We simulated a different neural network for each mouse. Each hidden
layer unit received input from a random subset of the prefrontal neurons from one mouse. I.e.,
each frame represents one timepoint and if neuron i is active in a frame then it provided an input
of 1 to all the hidden units to which it is connected; otherwise it provides an input of 0. For each
simulation, there was a fixed connection probability between each input neuron and each hidden
layer unit. We tried different values for this connection probability in order to measure how
classifier performance depends on the number of neurons that provide input to each hidden layer
unit. Each hidden layer unit had an output weight which specifies how strongly that unit excites
or inhibits a single output unit which classifies activity as belonging to periods in which a mouse
was actively engaged in social interaction or alone in its home cage. E.g., output unit activity <
0.5 corresponds to the social condition, while output unit activity > 0.5 corresponds to the home
cage condition. These output weights were adjusted during training (see Methods for details of
the training rule) while the pattern of input connectivity was fixed. This models the situation in
which prefrontal neurons transmit information to a downstream population of neurons (the
hidden layer) that decode behavior via their output weights. We initially trained networks on
50% of the data (frames) and used the held-out data for testing. We trained and tested using
intervals during which the mouse was actively engaged in social interaction (i.e. actively sniffing
/ nose in contact with subject mouse) or matched intervals when the mouse was alone in its home
cage.
208
Because each hidden unit simply sums the activity of a fixed population of mPFC neurons, one
can think of each hidden unit as representing one ensemble of mPFC neurons. The size of these
ensembles is determined by the input connection probability. In this framework, training serves
to determine how well each ensemble (corresponding to an individual hidden unit) is correlated
with behavior. Hidden units corresponding to ensembles which are strongly associated with one
type of behavior will be assigned large positive or negative output weights, whereas those that
are not well correlated with behavior will have output weights near zero.
216
Critically, we did not attempt to classify more nuanced behaviors, besides whether the subject
mouse was exploring another mouse vs. alone in its home cage. This was by design because
doing so would have reduced the number of training frames available for each behavioral label.
A large number of training frames for each behavioral label are necessary to train a neural
network with a large number of hidden units, and a large number of hidden units are necessary in
order to effectively capture possible collective encoding (because, as noted above, each hidden
unit effectively represents one neuronal ensemble). Thus, subdividing periods of social
interaction into more nuanced behavior would have compromised our ability to resolve collective

encoding. Therefore, because the focus of this paper is on trying to study collective encoding, we
intentionally used simple, binary classifications of behavior (e.g., social vs. home cage).
227
Classifier performance is optimal for intermediate connection probabilities
Classifier performance was strongly dependent on the probability that each input neuron was
connected to each hidden unit. For the 8/10 datasets that could be classified above chance,
classifier performance (measured on the 50% of data which was held-out during training) was
near chance levels when the connection probability was < 0.1, but increased to a peak of 69 +/-
3% (**Figure 2B; Table S1**) for a connection probability of 0.3. Accuracy decreased dramatically
when the connection probability increased to 0.5 indicating that connection probabilities ~0.2 -
0.4 are optimal.
236
We also validated classifier performance by training and testing on surrogate datasets that were
generated by 'swap shuffling' our original datasets. We created 'swap shuffled' surrogate
datasets by randomly swapping blocks of activity between neurons (each block of activity = a set
of consecutive frames during which the neuron was active). To understand this, think of the
entire raster as a collection of blocks of activity. Each block occurs at a specific time, has a
specific duration, and is associated with a particular neuron. Swap shuffling is equivalent to just
shuffling the neurons associated with each block of activity (the start time and duration of each
block do not change). For example, if neuron i originally became active at time $t1$ for $n1$ frames
and neuron j was active at time $t2$ for $n2$ frames, then in the surrogate dataset neuron i might
become active at $t2$ (but not at $t1$) while neuron j might become active at $t1$ (but not $t2$). Swap
shuffling preserves the number of neurons active at each point in time (because the timing of
blocks of activity does not change). It also preserves the number of blocks of activity for each
neuron, and this tends to preserve the overall level of activity of each neuron. Activity levels are
not perfectly preserved, because blocks of activity can have different durations. Nevertheless, in
practice, blocks of activity tend to have similar durations and the similarity between the mean
activity level in each neuron before and after swap shuffling of entire datasets was 0.97 +/- 0.01.
As expected, we found that neural network classifiers trained and tested on swap shuffled
datasets performed near chance levels (Figure 2B).
255
Prefrontal neurons that drive classifier performance change their correlations during
social behavior
Next, we examined connections in trained networks to reveal factors which enable them to
successfully classify social vs. home cage behavior (we analyzed networks with a connection
probability = 0.3 since this maximized performance of the population). Each hidden layer unit
has an output weight which measures how strongly it excites or inhibits the output unit that
represents the 'decision' (social vs. home cage). Hidden units with output weights ~0 don't
contribute to this decision. By contrast, hidden units with strong negative or positive weights
promote the social or home cage decision, respectively (**Figure 3A**). Therefore, we hypothesized
that there might be important differences in the pattern of input to hidden units, depending on
whether those hidden units have large positive or negative output weights.
267
We arranged hidden layer units based on their output weights, i.e., the unit with the most
negative weight was unit 1 and the unit with the most positive weight was unit 1000. Then we
defined the 25 hidden layer units with the most negative weights as 'social units' and the 25 with

the most positive weights as 'home cage units' (**Figure 3B**). For comparison we also defined the
25 hidden layer units with the weights closest to zero as 'neutral units.' For each pair of hidden
units, we computed the similarity between their inputs (i.e., the correlation between their input
vectors; **Figure 3C**). We then plotted the average input similarity of each hidden unit to either
the social or home cage units (**Figure 3D**) or the neutral units (**Figure 3E**). Social and home
cage units tended to receive input from the same prefrontal neurons as other hidden layer units
with the same preference, i.e., which also had negative or positive output weights. By contrast
neutral units did not exhibit any such relationship.
279
The preceding suggests that distinct ensembles of prefrontal neurons provide input to either
social or home cage hidden units. We hypothesized that there might be important features of
activity in these ensembles that support the classification of social vs. home cage behavior. For
example, one possibility is that prefrontal neurons which provide input to social units might tend
to increase activity during social behavior, whereas prefrontal neurons which provide input to
home cage units do the opposite. Surprisingly, this was not the case. In fact, both ensembles of
prefrontal neurons significantly increased their activity when mice were engaged in social
interaction (**Figure 3F**; social ensemble: mean activity level 1.4 +/- 0.3% in home cage vs. 1.8
+/- 0.3% during social interaction, $p < 0.05$, sign-rank test; home cage ensemble: mean activity
level 1.5 +/- 0.3% in home cage vs. 1.9 +/- 0.3% during social interaction, $p < 0.001$, sign-rank
test). Reasoning that changes in correlations would alter the frequency with which pairs or
groups of neurons became co-active, we examined pairwise correlations between the activity of
prefrontal neurons within each ensemble. Strikingly, mean correlations within the social
ensemble increased during social interaction (**Figure 3G**; (mean correlation coefficient between
neurons in the social ensemble: 0.009 +/- 0.002 in home cage vs. 0.012 +/- 0.002 during social
interaction, $p < 0.05$). By contrast, there was a non-significant decrease in correlations within the
home cage ensemble (Figure 3G; home cage ensemble mean correlation coefficient 0.011 +/-
0.02 in home cage vs. 0.005 +/- 0.003 during social interaction, p=0.99, sign-rank).
298
299 Thus, the ensemble of prefrontal neurons which provide input to the social units form an
300 assembly that collectively becomes more co-active (correlated) during social behavior. In
301 contrast, the prefrontal neurons which provide input to the home cage units increase their
302 activity, but not their co-activity, during social behavior. This suggests that behaviorally-driven
303 changes in correlations may contribute to the encoding of social behavior.
304
Swap shuffling within a behavioral state preserves information encoded by activity levels
but disrupts encoding by correlated activity
How can we quantitatively assess the contribution of these behaviorally-modulated correlations
to classifier performance? Ideally we would first train a neural network on the original data.
Then we would test this network's ability to classify data which maintained behaviorally-driven
changes in activity levels, but either removed or preserved the specific patterns of correlated
activity found in the original dataset. Indeed, we have already developed methods for shuffling
that achieve these goals. First, to shuffle the data in a manner that maintains behaviorally-driven
changes in activity levels, but disrupts the patterns of correlated activity, we can swap shuffle
activity, but do so within each behavioral condition rather than across the entire testing dataset.
In other words, we first divide up the raster into separate subrasters for each 5 minute behavior
epoch (when the mouse was either engaged in social interaction or alone in its home cage). Then

we performed swap shuffling (as described above) separately on each subraster, before
recombining these swap shuffled subrasters to create the swap shuffled surrogate dataset for
testing. Because swap shuffling tends to preserve activity levels, and because we swap shuffled
activity within a behavioral condition, neurons that increase or decrease activity during periods
of social interaction in the original dataset will also do so in the swap shuffled surrogate dataset.
322
We first confirmed that swap shuffling preserves information encoded by activity levels, while
disrupting the ability of correlated activity to transmit information. For this we created synthetic
datasets, swap shuffled them, and then measured classifier performance (**Figure 4**). Specifically,
we generated different types of synthetic datasets, which encoded two different 'behavioral
states' (state A or state B) via correlations or activity levels. In all cases, we first created a raster
for 'state A' by randomly assigning the activity of 100 neurons so that the total activity in each
frame oscillated around a mean level. To encode 'state B' using correlations, we started with the
original 'state A' raster, but shifted the activity of individual neurons in time, to create between 1
and 5 cell assemblies composed of co-active neurons. Specifically, each assembly contained 8
neurons; whenever one of these neurons was active, we probabilistically shifted the activity of
other neurons in that assembly, so that they would be co-active in the same frame (**Figure 4A**).
Notably, the activity level of every neuron was the same in the original 'state A' raster, and these
newly created 'state B' rasters. State B was distinguished simply by an increase in correlated
activity. (When we shifted the activity of one neuron in an assembly into a frame, we swapped
its activity with that of another neuron that was outside of the assembly; as a result, the total
activity in each frame was unchanged).
339
340 After training, our neural network classifiers were able to distinguish between 'state A' activity
341 and 'state B' activity (**Figure 4B**). This confirms that this type of neural network can classify
342 two different behaviors even when they are encoded entirely by changes in correlated activity,
343 not by any differences in activity levels. Furthermore, when we swap shuffled the 'state B' raster
344 to eliminate these correlations, classifier performance was reduced to chance levels (**Figure 4C**).
345
Next tested the ability of our neural networks to classify 'state A' vs. 'state B' utilizing
differences in activity levels. For this, we again started with a random 'state A' raster. Then, to
generate a 'state B' raster, we shifted a proportion of activity from neurons 1-50 to neurons 51-
100 (**Figure 4D**). Thus, in these synthetic datasets, neurons 1-50 have higher activity in 'state A'
whereas neurons 51-100 have higher activity during 'state B'. We varied the proportion of
activity that was transferred between neurons. Once again, after training, our neural networks
could classify activity patterns corresponding to 'state A' vs. 'state B' (**Figure 4E**). However, in
this case, swap shuffling 'state B' rasters did not disrupt classifier performance (**Figure 4F**). This
confirms that our neural networks classifiers can detect encoding that is based on either activity
levels or correlated activity. Swap shuffling a dataset (within one behavioral state) will remove
information encoded by correlated activity without diminishing encoding via activity levels.
357
SHARC shuffling preserves patterns of correlated activity
Before applying this same approach to our actual datasets, we also wanted to create a method for
shuffling that preserves patterns of correlated activity. For this, we used a method that we
published previously: SHuffling Activity to Preserve Correlations, or SHARC (22). Like swap
shuffling, SHARC also re-assigns blocks of activity between neurons, but rather than doing so

randomly, it instead follows an algorithm that achieves a target correlation matrix (in this case,
the original correlation matrix) (**Figure 5B-C**). The full details of SHARC are presented in the
Methods. Briefly: on each iteration, we randomly select one block of activity to be assigned to a
new neuron. Instead of choosing the new neuron randomly, we first compute the difference
between the target correlation matrix and the correlation matrix of the partially reconstructed
surrogate dataset. Then we assign the block of activity to the neuron which will optimally reduce
this difference. Finally, to maintain the mean activity level of each neuron, there is also an
absolute limit on how many blocks of activity can be re-assigned to each neuron. Notably, when
we applied SHARC shuffling to our previously described synthetic datasets, it visibly preserved
assemblies of coactive neurons, and did not disrupt the ability of neural network classifiers to
distinguish between 'state A' and 'state B' based on either activity levels or correlated activity
(**Figure S2**).
375
Patterns of correlated activity contribute to classifier performance
Having confirmed that we can swap or SHARC shuffle individual activity during individual
epochs to preserve activity levels while either disrupting or preserving correlated activity, we
now used this approach to quantitatively measure the encoding of social behavior by activity
levels vs. correlated activity in our actual datasets. Specifically, we swap or SHARC shuffled
each social or home cage subraster separately. Then we concatenated these shuffled subrasters to
create swap or SHARC shuffled surrogate datasets that preserve behaviorally-specific levels of
activity, while disrupting or preserving patterns of correlated activity, respectively.
384
Both swap and SHARC shuffled surrogate datasets preserved levels of activity observed during
either social interaction or periods when mice were alone in their home cages. Specifically, we
computed the correlation between vectors in which each element represents the activity level of
one neuron during one behavioral condition, and quantified the correlation between each real and
surrogate dataset. For swap shuffled surrogate datasets, the similarity of activity levels
(compared to real data) was 0.89 +/- 0.02 in the home cage and 0.82 +/- 0.04 during social
interaction. For SHARC shuffled surrogate datasets, the similarity of activity levels (compared to
real data) was 0.88 +/- 0.03 in the home cage and 0.86 +/- 0.03 during social interaction (n = 10
mice/datasets). We also computed the similarity of the pattern of correlations between each
surrogate dataset and the corresponding real dataset. In this case, only SHARC shuffled
surrogate datasets preserved patterns of correlations. For swap shuffled surrogate datasets, the
similarity of correlations to the real data was 0.01 +/- 0.01 in the home cage, and 0.03 +/- 0.01
during social interaction. For SHARC shuffled surrogate datasets, the similarity was 0.50 +/-
0.05 in home cage and 0.55 +/- 0.03 during social interaction.
399
We then trained classifiers on each dataset and tested each classifier using either swap or
SHARC shuffled surrogate datasets generated from the same dataset using for training (**Figure
5C**). Classifiers performed better than chance when tested with either surrogate dataset
indicating that changes in activity levels encode behavioral information, making it possible to
distinguish bouts of social interaction from periods when a mouse is alone in its home cage.
However, classifier accuracy was significantly higher for SHARC shuffled surrogates datasets
than for swap shuffled ones (**Figure 5D**, left; accuracy for classifying home cage vs. social = 71
+/- 2% for SHARC vs. 67 +/- 2% for swap shuffled datasets, p = 0.005, paired t test).
408

Patterns of correlated activity also help encode mouse identity

We wondered whether we could use this approach to study how correlations contribute to encoding of additional information besides just whether or not a mouse was engaged in social interaction. In our behavioral design, the subject mouse interacted during one mouse for two epochs and with a second mouse during two different epochs. Therefore, we also trained neural networks to classify whether the subject mouse was interacting with Mouse 1 vs. Mouse 2. Again, a neural network classifier was able to perform this classification above chance levels, and performance was higher when the classifier was tested using SHARC shuffled surrogate datasets than when tested using swap shuffled surrogate datasets (Figure 5D, right: accuracy for classifying Mouse 1 vs. Mouse 2 = 71 +/- 3% for SHARC vs 65 +/- 2% for swap shuffled datasets, p = 0.007, paired t test). This shows that behaviorally-modulated patterns of correlated activity transmit additional information, beyond what is readily decodable from activity levels alone.

Combinations of coactive neurons occur in a behaviorally-specific manner

Interestingly, neural networks perform classification better for connection probabilities ~0.2 – 0.4 than for connection probabilities < 0.1. When the connection probability is low, each hidden unit receives input from individual prefrontal neurons or small groups of neurons. By contrast, when the input probability is higher, hidden units receive input from larger groups of prefrontal neurons. This suggests that training proceeds more efficiently when the network represents information about social vs. home cage behavior using multineuron combinations, instead of activity within individual neurons or small groups. Together with the fact that classifier performance was higher for SHARC shuffled datasets than swap shuffled ones, this indicates that multineuron patterns of coactivity, rather than just levels of activity within neuronal ensembles, transmit information about social behavior. Therefore as a proof-of-concept, we directly examined whether 3-neuron patterns of coactivity occur in a behaviorally-specific manner. We examined 3-neuron combinations because they measure network structure beyond pairwise correlations and are the building blocks of larger combinations. One could in principle analyze larger combinations, but because of the limited numbers of neurons and frames in our datasets, there is not always adequate statistical power to resolve larger combinations, i.e., to detect large numbers of combinations that occur more often in real datasets than expected by chance.

First, we quantified how often each possible 3-neuron combination occurred in real datasets. Then we calculated how often each of these combinations occurred in datasets that had been swap-shuffled (across the entirety of the dataset). For each real dataset we constructed 1000 swap-shuffled datasets, and identified 'enriched combinations,' which occurred more often in real datasets than in 95% of swap shuffled surrogate datasets. Enriched combinations are those which occur more often in real datasets than expected based on the chance overlap of activity between marginally independent neurons. Finally, we quantified how many of these enriched combinations were behaviorally-specific, i.e., occurred exclusively during social or home cage epochs. Combinations could appear to be behaviorally-specific simply because they only occurred at a single timepoint. Therefore we also restricted our analysis to enriched combinations which occurred during multiple distinct bouts of social interaction and/or matched sets of intervals during home cage epochs. Many of these repetitively-occurring enriched combinations were behaviorally-specific: 43.5% occurred during social interaction, 26.5% during home cage epochs, and 30% during both conditions.

The selective occurrence of enriched combinations either during social interaction or when a
mouse is alone in its home cage may reflect changes in single neuron activity (i.e., neurons that
form a social combination are only active during the social condition), and/or changes in
correlations (i.e., neurons are active in both conditions but only *co-active* during social
behavior). To test the hypothesis that changes in correlations underlie the behavioral specificity
of significant combinations, we examined the 3-neuron combinations that were specifically
enriched during either periods of home cage exploration or social interaction (**Figure 6**). We
defined specific enrichment as those combinations which occurred more often in real data than in
95% of swap-shuffled surrogate datasets for one behavioral context, and less in real data than in
50% of swap-shuffled surrogate datasets for the other behavioral context. Based on these criteria,
12,408 3-neuron combinations were specifically enriched during social interaction, and 9,572
were specifically enriched during home cage exploration. There were 55,696 instances in which
a social and nonsocial 3-neuron combination overlapped in 2 out of 3 neurons. In 97.0% of these
cases, the neuron which was part of a social 3-neuron combination (triplet) but left out of the
overlapping home cage triplet was part of a different 3-neuron combination that was enriched
during homecage exploration (**Figure 6**, top right). Conversely, the neuron which was part of a
nonsocial triplet but left out of the overlapping social 3-neuron combination was part of a
different socially-enriched 3-neuron combination in 99.1% of cases (**Figure 6**, bottom right).
Overall, an average of 71 enriched homecage combinations contained the neuron missing from
the social triplet, and 85 enriched social combinations contained the neuron missing from
homecage triplets. Thus, the specificity of a combination of co-active neurons for social vs.
nonsocial behavior does not occur simply because some neurons were only active during one
condition, but rather reflects the dynamic reorganization of patterns formed by neurons which are
active in both conditions, i.e., changes in correlations. This – the behaviorally-specific
occurrence of multineuron patterns of coactivity – represents the substrate through which
correlations can add to the behavioral information transmitted by neuronal ensembles.

Socially-enriched combinations are deficient in Shank3 KO mice
We were curious whether there might be conditions under which these phenomena – the
occurrence of multineuron combinations of coactivity during social behavior, and the ability of
correlated activity to enhance the transmission of information about social behavior – might be
impaired. To explore this, we performed microendoscopic GCaMP imaging in mice lacking the
autism-associated gene Shank3 (23–25). These mice have been extensively studied as models of
Phelan-McDermid syndrome, which often includes autism as a clinical feature. Shank3$^{-/-}$ (KO)
mice are known to have social deficits, and indeed, we found that compared to wild-type (WT)
littermates, they spend significantly less time interacting with novel juvenile mice (**Figure 7A**).

We compared patterns of prefrontal activity in Shank3 KO mice and their WT littermates. As in
WT mice, in Shank3 KO mice, many prefrontal neurons either increase or decrease activity
during social interaction. However, compared to WT mice, the fraction of neurons whose activity
increases during social interaction was significantly higher, whereas the fraction whose activity
decreases was significantly lower (**Figure 7B-C;** 22% of 260 WT neurons vs. 39% of 290 KO
neurons increased activity above the 90th percentile of shuffled data during social interaction, chi
squared = 17.7, p < 0.0001; 25% of WT vs. 15% of KO neurons decreased activity below the
10th percentile of shuffled data during social interaction, chi squared = 8.2, p < 0.0001). Thus,

Shank3 KO mice recruit abnormal neuronal ensembles during social behavior. We hypothesized that this might reflect a network-level disorganization that affects the normal patterning of activity during social behavior.

Indeed, we found that in KO mice a significantly smaller fraction of the 3-neuron combinations observed during social interaction were strongly enriched, i.e., occur more often in actual data than in 99.9% of swap-shuffled surrogate datasets (**Figure 7D**). This suggests that even though more neurons (i.e., larger ensembles), were recruited during social behavior in KO mice, these may have been less well-organized, such that the occurrence of socially-enriched patterns of activity is obscured by 'noise,' i.e., patterns formed by the chance overlap of activity between neurons that fire in a largely independent fashion. Notably, this deficiency was specific for social interaction. The fraction of 3-neuron combinations that were strongly enriched during home cage exploration (in comparison to swap-shuffled surrogate datasets) was similar in WT and KO mice (Figure 7D).

The temporal organization of multineuron activity does not enhance the transmission of information about social behavior in Shank3 KO mice

The preceding shows that even though social behavior robustly recruits neuronal ensembles in Shank3 KO mice, the organization of these ensembles into multineuron combinations is disorganized. This suggests that the ability of patterns of correlated activity to encode information about social behavior may be impaired in these mice. To test this, we examined whether correlated activity contributes to the transmission of information about social behavior in Shank3 KO mice. As before, we generated swap and SHARC shuffled surrogate datasets, then tested the ability of classifiers trained on the original datasets (from Shank3 KO mice) to classify activity associated with behavior during social interaction vs. in home cage. We then compared the degree to which classifier accuracy was improved when testing SHARC-shuffled surrogate datasets compared to testing swap-shuffled ones. In datasets from Shank3 KO mice, we observed a significantly diminished improvement in classifier accuracy for SHARC vs. swap shuffled datasets, indicating a diminished role for correlated activity in encoding (relative improvement for classifying home cage vs. social: 0.44 +/- 0.10 in WT vs 0.24 +/- 0.01 in KO; relative improvement for classifying whether the subject was interacting with Mouse 1 vs. Mouse 2: 0.41 +/- 0.15 in WT vs. 0.14 +/- 0.06 in KO; 2-way ANOVA using classification type and genotype as factors: significant for genotype, $p < 0.05$, no effect of classification type, $p = 0.58$, or interaction, $p = 0.76$). Thus, in Shank3 KO mice, the multineuron patterns of coactivity which normally occur during social behavior are disturbed, and as a result patterns of correlated activity contribute less information about social behavior.

DISCUSSION

During complex behaviors, the brain can use many strategies to represent information about the external environment and internal state of the organism. The term 'ensemble' is often used to refer to a group of neurons whose activity is similarly modulated (either increased or decreased) during specific behaviors (1,26–30). It is generally accepted that ensembles transmit behavioral information via changes in the activity levels of their constituent neurons. On the other hand, many studies have also shown that correlations between neurons can change during specific behaviors (3,10) or behavioral states (31–33). Importantly, correlations reflect changes in coactivity which exceed those expected to occur simply because of changes in the activity levels of the individual neurons (6). I.e., when an ensemble becomes more active, correlations between the neurons in that ensemble could go up, down, or remain unchanged. By optimizing synaptic interactions such as temporal summation, changes in correlated activity could potentially act to enhance the behavioral information transmitted by changes in ensemble activity, or transmit additional information.

The role of correlations in information transmission has been studied extensively in the isolated retina (34). In the cortex, noise correlations – correlations in trial-to-trial variability across neurons that are thought to be unrelated to the variable of interest – have also been extensively studied(11,13,14). Noise correlations can enhance information transmission, presumably by helping to disambiguate changes in activity levels due to the variable of interest from those attributable to other causes (noise). However, the ability of correlations to transmit information themselves, by changing as a function of the variable being encoded, has been less well studied. In particular, it is not clear whether changes in correlations encode additional information, beyond what is transmitted by changes in activity levels, and if so whether this reflects the behaviorally-specific occurrence of patterns of correlated activity / coactivity.

Here, we addressed this question using microendoscopic GCaMP imaging to measure activity from many (~40-100) prefrontal neurons during social behavior in mice. We used a simple neural network, in which prefrontal neurons provide input, there is one hidden layer, and a single output unit classifies social vs. nonsocial behavior, to quantify how well prefrontal ensembles would transmit information about social behavior to downstream neurons. We used multiple new approaches to disentangle the respective contributions of changes in activity levels vs. correlated activity to the encoding of social behavior. First, whereas previous studies have compared information transmission by real datasets vs. by shuffled ones(14), we extended a method we previously published, (22), to non-randomly shuffle datasets in order to preserve both behaviorally-modulated correlations and ensemble activity. This enabled us to compare the amount of information about social behavior transmitted by either SHARC-shuffled surrogate datasets or randomly-shuffled surrogates which preserved ensemble activity but not correlations. In this way, we found that correlated activity enhances the amount of information that prefrontal ensembles transmit about social behavior. Notably, we used neural networks to classify periods when a subject mouse was alone in its home cage vs. interacting with another mouse, and separate classifiers to determine whether the subject was interacting with test mouse 1 vs. test mouse 2. We found that correlated activity normally enhances the encoding of both types of information.

Second, when we examined connections within neural network classifiers, we found that
prefrontal neurons which serve to detect social behavior increase their correlations during social
behavior (whereas neurons which detect nonsocial behavior do not). Third, we found that
combinations of coactive neurons which occur more often than expected based on the activity
levels of the constituent neurons, manifest in a behaviorally-specific manner. Positive
correlations measure neuronal coactivity that occurs more often than expected based on the
chance overlap of activity between neurons. Thus, in accordance with our finding that social
behavior modulates correlations, we found that multineuron patterns of coactivity which occur
more often than expected by chance are behaviorally-specific. We also directly examined these
behaviorally-specific and statistically-enriched combinations of coactive neurons. We found that
they tend to be composed of neurons which are active in both conditions but only coactive in
one, rather that neurons which are only active in one condition. Thus, we used multiple new
approaches (comparing classifier accuracy for SHARC vs. swap shuffled datasets, measuring
changes in correlations for neurons in ensembles derived from neural network classifiers, linking
statistically enriched patterns of coactivity to specific behaviors) to show exactly how dynamic
correlations encode information about social behavior.
600
Interestingly, statistically-enriched patterns of coactivity were specifically deficient during social
behavior in mice lacking the autism-associated gene Shank3. Accordingly, in Shank3 KO mice,
surrogate datasets which preserve behaviorally-modulated correlations failed to transmit more
information about social behavior compared to randomly shuffled datasets which only preserved
ensemble activity. This shows that the ability of correlated activity to enhance the transmission
of information about social behavior is not automatic, and can in fact be disrupted in pathological
states.
608
609   Similar to other recent studies (12,13), we have studied activity using binary activity rasters
610   derived from GCaMP imaging. However, an important caveat is that any method of quantifying
611   neural activity has limitations, such that there could be additional ways that neurons encode
612   information which are not well resolved using this approach.
613
What is the meaningful size of ensembles in the cortex?
Complex behavior is possible because the brain reliably encodes features pertaining to the
external environment as well as the internal state of the organism. These features may be
encoded by neuronal ensembles (1,26–30). What size of ensemble reliably encodes an aspect of
behavior? We explored this question, by asking what input connection probability would
optimize the ability of a downstream network to classify behavior based on input from prefrontal
ensembles. Note: input connections in neural network classifiers do not necessarily correspond to
actual connections in the brain – rather they provide information about the size and nature of
neuronal ensembles across which information should be combined to most efficiently decode
behavior. Peak classifier performance occurred for connection probabilities ~0.2 - 0.3.
Performance was markedly lower when the connection probability was 0.5. This is surprising
because a connection probability of 0.5 would maximize the entropy associated with each input
connection. Thus, from the standpoint of encoding social behavior, combining activity from 20-
30% of the input neurons may achieve some synergy that becomes degraded when ensembles are
enlarged beyond this size. In the brain, nonrandom network connectivity (35,36) may similarly
produce correlated activity / coactivity within ensembles of this size (37,38).

630
Combinatorial codes vs. sequential patterns of activity
Like many recent studies, we measured population-level activity in the mouse neocortex using
genetically encoded calcium indicators. These indicators transduce neuronal spiking on
timescales ~100 msec. Thus correlated activity / 'coactivity' imply that neurons jointly increase
their activity within windows ~100 msec, and do not necessarily imply synchronous spiking on
faster timescales (milliseconds or even tens of miliseconds). At the same time, correlated activity
/ coactivity on these timescales should be differentiated from sequential activity of neurons
observed during the performance of sequential behaviors (i.e. spatial navigation or overtrained
tasks) in which the activity of specific neurons corresponds to moving through a specific location
or performing a specific portion of a complex task. As discussed above, in the neocortex
correlations and coactivity likely reflect recurrent neural network connectivity (37). By contrast,
sequential patterns of neuronal activation can occur simply as a byproduct of the arrangement of
spatial locations along a trajectory, the stereotyped order in which cues are encountered during a
task, etc.
645
Relevance to disease states
Interestingly, in *Shank3* KO mice, which exhibit social deficits, the mPFC successfully recruits
specific neuronal ensembles during social interaction. However these ensembles are enlarged,
their organization into statistically-enriched patterns of coactivity is disrupted, and correlations
between neurons fail to enhance the information that these ensembles transmit. Thus, the
computational units by which information is processed in the mPFC appears to be inefficient,
i.e., social behavioral recruits an abnormally large number of neurons at the expense of the
precise temporal patterning of this activity. This central finding is similar to other findings in
rodent models of autism at both the single neuron and network levels (19,23,39). In particular,
we found an increase in the recruitment of prefrontal neurons during social interaction. This
mirrors a recent study which found hyperdynamic response to whisker stimulation in the same
mice (23), possibly reflecting GABAergic circuit dysfunction and/or homeostatic compensations
(40). (Note: these findings cannot be ascribed simply to the fact that Shank3 KO mice spend less
time engaged in social interaction than their wild-type littermates; reduced interaction time
would tend to reduce statistical power and thereby reduce the number of neurons that change
their activity more than expected by chance.)
662
Increased excitatory activity causing decreased signal-to-noise ratio (SNR) has long been posited
to contribute to the pathophysiology of autism (41). However the exact nature of 'signal' and
'noise', and the specific mechanism through which excessive activity degrades the SNR have
been unclear. Here, we show how enlarged neural ensemble recruitment by specific behavioral
conditions disrupts information transmission by degrading the ratio between statistically
meaningful patterns of coactivity (the signal) and the random overlap of activity between
neurons (noise).
670

MATERIALS AND METHODS

Behavior: C57/B6 mice were obtained from Jackson Laboratories. We utilized adult mice of
either sex housed and bred in the UCSF animal facility. Adult mice were habituated to the room
and observer for 3 days prior to test day. All videos were subsequently scored by a blinded
observer. For imaging experiments, 5 WT and 4 KO littermates were generated through crosses
between Shank3 heterozygous parents and injected with AAV5.Syn.GCaMP6f.WPRE.SV40. We
included an additional 5 WT mice which were injected with
AAV5.Syn.GCaMP6m.WPRE.SV40 (42). Viruses were obtained from Penn Viral Core.
Injections and 500 um GRIN lens (Inscopix) implantations were carried out in 8-12 week old
mice to express GCamp6f in prefrontal cortical neurons under control of the human Synapsin
promotor. Mice were anesthetized with 2% isoflurane and mounted in a stereotactic frame.
Craniotomies were made according to stereotaxic coordinates relative to Bregma. Coordinates
for injection into mPFC were (in mm relative to Bregma): +1.7 anterior–posterior (AP), –0.3
mediolateral (ML) and –2.75 dorsoventral (DV), and GRIN lenses were implanted at the same
AP and ML coordinates, to a depth of 2.25. We subsequently attached baseplates for attaching
the microendoscope, ~4 weeks later depending on GCamp expression. Mice were habituated
for three days with the scope attached, prior to test day. On test day, mice were habituated with
the scope turned on, then imaged in alternating home cage and social epochs. During social
epochs, one of 2 novel sex-matched juvenile mouse was introduced to the test mouse's
homecage, in sequential order so that there were two 'novel' epochs, followed by two 'familiar'
epochs interleaved with 'home cage' epochs during which the juvenile mice were removed and
the test mouse was free to explore its home cage. The first and last home cage epoch were 10
minutes in length; the others were 5 minutes in length. Each social epoch lasted 10 minutes but
only the first 5 minutes were recorded and scored. During each behavioral epoch, observer was
not in the room. Interaction epochs were defined from the moment test mouse first sniffed the
juvenile conspecific or object, until the test mouse turned away. Videos were recorded using
Anymaze, and scored by a blinded observer. For the bulk of analysis we pooled data across 10
WT mice. Shank3 KO mice were compared only to recordings from WT littermates.

Image acquisition and segmentation: Images were acquired using an Inscopix nVoke
microendoscope attached to a laptop computer and synced to a separate video acquisition
computer running Anymaze. Frame rate was 20 Hz and the laser power was 0.2 mW. Acquisition
was performed using 2x2 pixel binning, then subsequently downsampled again by 2.

We segmented neuronal signals using a modified PCA/ICA approach (20,21), modified so that
each segment was expressed as a binary ROI consisting of pixels representing a single neuron.
I.e., we used the output from the PCA/ICA to identify a set of contiguous pixels which represent
a neuron, then averaged fluorescence signals across those pixels. To deconvolve neuronal signals
from background neuropil signals, we also subtracted the mean signal from each identified
segment from the mean value in pixels surrounding the edge of the segment (we excluded pixels
that belonged to another ROI). Signals were subsequently lowpass filtered to remove high
frequency noise using the Matlab command: designfilt('lowpassfir', 'PassbandFrequency', 0.5,
'StopbandFrequency', .65, 'PassbandRipple', 1, 'StopbandAttenuation', 25). All signal traces
shown represent normalized versions of the dF/F_0 trace, where F_0 is estimated by the median
value in the surround region. Threshold based event detection was performed on the traces by

detecting increases in dF/F_0 exceeding 3σ over one second, then only keeping those events
which exceeded a 15σ increase over two seconds, and a total area under the curve of 250σ. As
there were occasional downward deflections due to surround subtraction, we instituted a final
parameter requiring that the peak cross an absolute value of $dF/F_0 = 0.0125$. σ is the standard
deviation of dF/F_0, calculated over the least active 50% of the movie. In some cases these
parameters were adjusted slightly to optimize event detection to > 95% sensitivity and
specificity, based on visual inspection, for each movie. After identifying these events in the
GCaMP signal from a cell, the cell was considered "active" during the entire period from the
beginning of an event until the GCaMP signal decreased 30% from the peak of the event, up to a
maximum of 2 seconds. The peak of the event was defined as the local maximum of the entire
event, from the beginning of the event until dF/F_0 returned to the pre-event baseline value.
Calcium traces from segmented neurons were visually inspected and a small number of segments
were removed if they did not appear to represent a single, unduplicated neuron. We restricted
further analysis to those mice with 25 or more active neurons. We then created 2-dimensional
event rasters consisting of detected events for each neuron over the course of the experiment.

Detection of behaviorally modulated neurons: To determine the response of individual neurons
to behavioral context, we averaged the activity of each neuron during frames corresponding to
periods of social interaction, or to a temporally matched set of frames during the preceding home
cage epoch. We then created a 'null distribution' for each neuron that represents the percent of
time active expected in each condition based on chance, by circularly shuffling the data 10,000
times. We then compared the activity of each neuron during either social interaction or home
cage exploration to this null distribution. Neurons were considered positively modulated if they
exceeded the 90th percentile of that observed in circularly-shuffled datasets, and negatively
modulated if the percent of frames that a neuron was active during a given context was below the
10th percentile of observations from circularly-shuffled data.

SHARC: SHARC (SHuffling Activity to Rearrange Correlations) is an iterative method for
generating surrogate datasets. SHARC nonrandomly shuffles blocks of activity within a raster to
generate a new (surrogate) raster in which the pairwise correlations between neurons match a
target correlation matrix (21). Here we apply this previously-published method, with
modifications to also preserve the activity level in each neuron (Figure 4B).

To begin, note that each raster is equivalent to a collection of blocks of activity. Each block of
activity is defined by the time at which it begins, its duration, and the neuron which is active. On
each iteration one block of activity is randomly chosen and assigned to a new neuron as follows.
Suppose block i has been chosen to be re-assigned. First, we find all the blocks of activity that
overlap with block i. Next, we selected the subset of these blocks for which new cell identities
had already been assigned. Call this set X. Let r_j represent the number of timepoints over which
block $j \in X$ overlaps with block i, and let n_j represent the identity of the cell assigned to block $j \in$
X. L_i and L_j are the lengths of blocks i and j, respectively. Then we constructed a vector,

$$\vec{P}_i = \sum_{j \in X} \frac{r_j}{\sqrt{L_i L_j}} \left(\vec{C}_{n_j} - \vec{C'}_{n_j} \right)$$

where $\vec{C}_{n_j}$ represents row j of the target correlation matrix, i.e. the target correlations between

neuron n_j and the other neurons, and $\vec{C'}_{n_j}$ contains the current values of the correlations between

neuron n_j and the other neurons based on the blocks of activity that have already been re-
assigned. This step can be thought of as "guessing" which cell *should* be assigned to a particular
block of activity by first figuring out what other cells are active at the same time, then choosing
cells which are strongly correlated with these known active cells. Note that we assign values of
$\vec{P}_i$ (i.e., construct "guesses" about which cell should be active), using the *difference* between the
current correlation matrix ($\vec{C}'_{n_j}$) and the target correlation matrix ($\vec{C}_{n_j}$), in order to identify cell
pairs for which the current correlation deviates from the target value, and force the new
correlation matrix to progressively approximately the target correlation matrix.
We set elements of $\vec{P}_i$ to zero if the corresponding neuron had already been assigned to a block
of activity that overlaps with block i, i.e. element n_j of $\vec{P}_i$ was set to zero $\forall j \in X$. Finally, we
assigned block i to the neuron corresponding to the maximum value of $\vec{P}_i$. This can be thought of
as choosing the cell that represents the "consensus" based on tallying up all of the "guesses"
about which cells "should" be assigned to the block of activity being considered.
774
When all the elements of $\vec{P}_i$ were zero, e.g. because there no overlapping blocks of activity have
had new cell identities assigned yet, then we chose a cell in order to match the originally
observed level of activity. Specifically, after every iteration, we kept a log of the net number of
blocks of activity that each neuron had donated or received. We used this vector to create a
weighted probability whereby events from neurons which had received a net positive number of
blocks were more likely to be chosen to be reassigned. To further ensure that the total number of
active events for each neuron in the surrogate dataset was similar to the real dataset, if the
difference between the number of blocks gained – lost in the reassignment process exceeded +4
for a particular neuron, then that neurons was no longer eligible to receive additional blocks of
activity; similarly blocks of activity belonging to neurons who had lost greater than 3 blocks of
activity were no longer eligible be reassigned. As the selection of blocks of activity for
reassignment is probabilistic, we first perform a swap shuffle to ensure that all blocks of activity
in the original dataset are swapped at least one time, then perform a total of 5 reassignments for
each block of activity in the source dataset.
789
790 We extended this approach to generate surrogate datasets by shuffling data within shorter time
791 windows (i.e., individual behavioral epochs). Here a discrete set of frames is chosen,
792 corresponding to a subraster of the original raster. By repeating the process described above for
793 each subrasters, then recombining the shuffled subrasters, we generate a complete shuffled
794 dataset.
795
Classifier: We designed and trained a neural network to classify behavior (periods when a mouse
was alone in its home cage vs. engaged in social interaction). This network contained 1000 units
in a hidden layer, each of which received input from specific prefrontal neurons (from the real
dataset). Thus, in each frame the activity of each hidden layer unit was just the summed activity
of the connected prefrontal neurons. Each hidden layer unit had an output weight that
represented the strength of its connection to a single output unit. On each frame the activity of
the output unit was computed as:
803
804
$$y = \frac{1}{1 + e^{-\sum w_i x_i}}$$

805
806   where w_i is the output weight from hidden unit i and x_i is the activity of hidden unit i.

807
When we performed training and testing using the same dataset, we divided the dataset into
alternating blocks of 500 frames for training vs. testing (in other cases we used the real dataset
for training, then tested using a surrogate dataset). We restricted training or testing to frames in
which mice were scored as actively engaged in social interaction (or matched frames during
periods when the mouse was alone in its home cage). We also limited training / testing to frames
with at least 3 active neurons.

814
815   We trained the output weights by performing 500 passes through the training data (each pass
816   visited all of the training frames in a random order). On each training timestep, we calculated y,
817   the activity of the output unit, and then adjusted each output weight based on:

818
819    $$\Delta w_i = \varepsilon y (1 - y)(y - z) x_i$$

820
821   where z is the correct classification of the frame (0 for social behavior, 1 for home cage) and ε,
822   the learning rate, was set to 0.05.

823
Following training, we examined the pattern of input connections and output weights. The
distribution of output weights was roughly gaussian and centered near 0. We identified the
selection of prefrontal neurons most likely to be connected to hidden layer units with large
positive or negative weights. Hidden layer units with large negative or positive output weights
bias classification towards the social or home cage condition, respectively. Therefore, we refer to
the 25 hidden units with the most negative or positive weights as 'social' or 'home cage' units
respectively. We calculated the number of input connections between each prefrontal neuron and
the 25 home cage units or 25 social units. We then defined 'home cage' or 'social' ensembles as
the 20% of prefrontal neurons with the most input connections to home cage or social units,
respectively. As described in the main text, we then analyzed properties of these two ensembles.

834
Quantification of multineuron combinations: Estimating chance overlap between activity of
largely independent neurons requires accounting for two factors. First, neurons with higher
activity are more likely to overlap by chance with other neurons. Second, overall network
activity is dynamic over time, creating a tendency for otherwise independent neurons to be
recruited at similar times. Thus, it is necessary to identify combinations which occur more often
than expected based on 1) the activity levels of the constituent neurons, and 2) the fact that
activity in a network is not constant over time. We can do this by quantifying the occurrence of
combinations in datasets which have been shuffled to preserve 1) the overall level of activity in
each neuron, and 2) the total level of activity in the network at each point in time.

844
3 neuron combinations were quantified by identifying each combination present in frames in
which 2 or more neurons were active. The number of frames each combination was active in
real data was stored in a n-dimensional matrix. Surrogate datasets were then generated from
event rasters by *swapping* the identity of neurons associated with detected events (periods of
activity). As the timing of events themselves is unchanged, and only the identity of the
participating neurons are exchanged, this preserves both the number of events per frame and the

number of events that each neuron participates in. Therefore, the total number of combinations in
each frame and over the course of the experiment (i.e., the sum of occurrences across all
combinations) is also preserved. The total number of combination occurrences in which a given
neuron participates would also tend to be preserved in these swap-shuffled surrogate datasets.
855
We then quantified how often each combination occurred in real vs. swap-shuffled data. By
comparing how often each combination occurred in real data vs. in 1,000 swap-shuffled
surrogate dataset, we were able to quantify how 'enriched' each combination was, compared to
the level of occurrence expected by chance based on the activity levels of its constituent neurons
(and the overall temporal pattern of network activity). We expressed enrichment as a percentile,
calculated relative to swap-shuffled surrogate data, e.g., the 100^{th} percentile indicates that a
particular combination occurred more often in real data than in all 1,000 surrogate datasets.
Further analysis was restricted to 'enriched combinations', i.e., combinations that occurred more
often in real datasets than in 95% of surrogate datasets.
865
Generation of Synthetic datasets: all synthetic datasets consisted of 100 neurons and 6000 frames
with an overall activity A of 5%. Briefly we randomly chose n active neurons in each frame f
where $n = A + sin(f)A$. Nonoverlapping patterns or assemblies each consisting of 8 neurons
were inserted into this oscillating network reciprocally swapping activity from non-assembly
neurons to assembly neurons in each frame in which the first neuron of the assembly/pattern was
active. In frames in which less than 8 neurons were active we randomly chose neurons from the
assembly to be activated. To maintain equivalent levels of activity reciprocal swaps were made
by choosing another frame in which the assembly neuron receiving activity was active and
reassigning activity in that second frame from the assembly neuron to the non-assembly neuron.
To create datasets in which the activity of individual neurons was modulated between 'State A'
and 'State B' we transferred a proportion of active events from half the neurons to partner
neurons so that half the neurons increased their activity and half the neurons activity decreased.
For each pair of neurons the proportion of transferred activity was a random number between 0
and the specified upper bounds of modulation which ranged between 5 and 50%. We trained our
classifier on blocks consisting of 50% of each dataset and testing on the remainder.
881
882 Statistical analysis: Neurons and significant combinations from all animals and groups were
883 pooled and counted as single units. Proportions were compared using chi-squared test. Activity
884 levels were compared using paired *t*-tests (2-sided), unless otherwise noted. Where applicable,
885 error bars denote standard error. Values of the classifier performance (accuracy) were generated
886 by averaging after re-running the training / testing procedure at least 25 times.
887

888 **Acknowledgments:** We thank Josh Berke and Ruchi Malik for helpful comments on the
889 manuscript.

890 **Declaration of interests:** Authors declare no competing interests.

891 **Data and materials availability:** All code used in the analysis will be deposited on GITHUB
892 for any researcher for purposes of reproducing or extending the analysis.

REFERENCES

1. Cai DJ, Aharoni D, Shuman T, Shobe J, Biane J, Song W, et al. A shared neural ensemble links distinct contextual memories encoded close in time. Nature. 2016;

2. Liang B, Zhang L, Barbera G, Fang W, Zhang J, Chen X, et al. Distinct and Dynamic ON and OFF Neural Ensembles in the Prefrontal Cortex Code Social Exploration. Neuron [Internet]. 2018 Nov 7 [cited 2019 Mar 30];100(3):700-714.e9. Available from: https://www.sciencedirect.com/science/article/pii/S0896627318307724?via%3Dihub

3. deCharms RC, Merzenich MM. Primary cortical representation of sounds by the coordination of action-potential timing. Nature [Internet]. 1996 Jun 13 [cited 2019 Aug 25];381(6583):610–3. Available from: http://www.ncbi.nlm.nih.gov/pubmed/8637597

4. Hebb DO. The organization of behavior: A neuropsychological theory. New York: Wiley; 1949.

5. Buzsáki G. Neural Syntax: Cell Assemblies, Synapsembles, and Readers. Neuron [Internet]. 2010 Nov 4 [cited 2019 Mar 30];68(3):362–85. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21040841

6. De La Rocha J, Doiron B, Shea-Brown E, Josić K, Reyes A, Josi K, et al. Correlation between neural spike trains increases with firing rate. Nature. 2007;

7. Ruff DA, Cohen MR. Global cognitive factors modulate correlated response variability between V4 neurons. J Neurosci [Internet]. 2014 Dec 3 [cited 2020 Oct 31];34(49):16408–16. Available from: https://www.jneurosci.org/content/34/49/16408

8. Averbeck BB, Latham PE, Pouget A. Neural correlations, population coding and computation. [cited 2020 Aug 2]; Available from: www.nature.com/reviews/neuro

9. Abbott LF, Dayan P. THE EFFECT OF CORRELATED VARIABILITY ON THE ACCURACY OF A POPULATION CODE.

10. Vaadia E, Haalman I, Abeles M, Bergman H. Dynamics of neuronal interactions in monkey cortex in relation to behavioural events. Nature [Internet]. 1995 Feb 9 [cited 2020 Jul 19];373(6514):515–8. Available from: https://www.nature.com/articles/373515a0

11. Averbeck BB, Lee D. Effects of noise correlations on information encoding and decoding. J Neurophysiol [Internet]. 2006 Jun [cited 2020 Aug 13];95(6):3633–44. Available from: www.jn.org

12. Ahmed MS, Priestley JB, Castro A, Stefanini F, Solis Canales AS, Balough EM, et al. Hippocampal Network Reorganization Underlies the Formation of a Temporal Association Memory. Neuron. 2020 Jul 22;107(2):283-291.e6.

13. Stefanini F, Kushnir L, Jimenez JC, Jennings JH, Woods NI, Stuber GD, et al. A Distributed Neural Code in the Dentate Gyrus and in CA1. Neuron [Internet]. 2020 [cited 2020 Aug 2]; Available from: https://pubmed.ncbi.nlm.nih.gov/32521223/

14. Siniscalchi MJ, Wang H, Kwan AC. Enhanced Population Coding for Rewarded Choices in the Medial Frontal Cortex of the Mouse. Cereb Cortex [Internet]. 2019 Sep 13 [cited 2020 Nov 3];29(10):4090–106. Available from:

https://pubmed.ncbi.nlm.nih.gov/30615132/

15.   Yizhar O, Fenno LE, Prigge M, Schneider F, Davidson TJ, O'Shea DJ, et al. Neocortical
excitation/inhibition balance in information processing and social dysfunction. Nature.
2011;

16.   Selimbeyoglu A, Kim CK, Inoue M, Lee SY, Hong ASO, Kauvar I, et al. Modulation of
prefrontal cortex excitation/inhibition balance rescues social behavior in CNTNAP2-
deficient mice. Sci Transl Med [Internet]. 2017 Aug 2 [cited 2019 Aug
23];9(401):eaah6733. Available from: http://www.ncbi.nlm.nih.gov/pubmed/28768803

17.   Brumback AC, Ellwood IT, Kjaerby C, Iafrati J, Robinson S, Lee AT, et al. Identifying
specific prefrontal neurons that contribute to autism-associated abnormalities in
physiology and social behavior. Mol Psychiatry [Internet]. 2017 Nov 7 [cited 2018 Feb
12]; Available from: http://www.ncbi.nlm.nih.gov/pubmed/29112191

18.   Murugan M, Jang HJ, Park M, Miller EM, Cox J, Taliaferro JP, et al. Combined Social
and Spatial Coding in a Descending Projection from the Prefrontal Cortex. Cell [Internet].
2017 Dec [cited 2019 May 20];171(7):1663-1677.e16. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0092867417313119

19.   Levy DR, Tamir T, Kaufman M, Parabucki A, Weissbrod A, Schneidman E, et al.
Dynamics of social representation in the mouse prefrontal cortex. Nat Neurosci [Internet].
2019 Nov 25 [cited 2020 Jan 13]; Available from:
http://www.ncbi.nlm.nih.gov/pubmed/31768051

20.   Mukamel EA, Nimmerjahn A, Schnitzer MJ. Automated analysis of cellular signals from
large-scale calcium imaging data. Neuron [Internet]. 2009 Sep 24 [cited 2017 Jun
6];63(6):747–60. Available from:
http://linkinghub.elsevier.com/retrieve/pii/S0896627309006199

21.   Luongo FJ, Horn ME, Sohal VS. Putative microcircuit-level substrates for attention are
disrupted in mouse models of autism. Biol Psychiatry. 2016;

22.   Luongo FJ, Zimmerman CA, Horn ME, Sohal VS. Correlations between prefrontal
neurons form a small-world network that optimizes the generation of multineuron
sequences of activity. J Neurophysiol [Internet]. 2016 May 1 [cited 2018 Feb
6];115(5):2359–75. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26888108

23.   Chen Q, Deister CA, Gao X, Guo B, Lynn-Jones T, Chen N, et al. Dysfunction of cortical
GABAergic neurons leads to sensory hyper-reactivity in a Shank3 mouse model of ASD.
Nat Neurosci [Internet]. 2020 Mar 2 [cited 2020 Mar 9];1–13. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/32123378

24.   Peça J, Feliciano C, Ting JT, Wang W, Wells MF, Venkatraman TN, et al. Shank3 mutant
mice display autistic-like behaviours and striatal dysfunction HHS Public Access. Nature
[Internet]. 2011;472(7344):437–42. Available from:
http://www.nature.com/authors/editorial_policies/license.html#terms

25.   Duffney LJ, Zhong P, Wei J, Matas E, Cheng J, Qin L, et al. Autism-like Deficits in
Shank3-Deficient Mice Are Rescued by Targeting Actin Regulators. Cell Rep. 2015;

26.   Corder G, Ahanonu B, Grewe BF, Wang D, Schnitzer MJ, Scherrer G. An amygdalar

neural ensemble that encodes the unpleasantness of pain. Science (80-). 2019 Jan
18;363(6424):276–81.

27. Gründemann J, Bitterman Y, Lu T, Krabbe S, Grewe BF, Schnitzer MJ, et al. Amygdala
ensembles encode behavioral states. Science [Internet]. 2019 Apr 19 [cited 2019 Jul
8];364(6437):eaav8736. Available from: http://www.ncbi.nlm.nih.gov/pubmed/31000636

28. Ghandour K, Ohkawa N, Fung CCA, Asai H, Saitoh Y, Takekawa T, et al. Orchestrated
ensemble activities constitute a hippocampal memory engram. Nat Commun [Internet].
2019 Dec 14 [cited 2019 Jul 8];10(1):2637. Available from:
http://www.nature.com/articles/s41467-019-10683-2

29. Sakurai Y. How do cell assemblies encode information in the brain? Neuroscience and
Biobehavioral Reviews. 1999.

30. Sakurai Y, Osako Y, Tanisumi Y, Ishihara E, Hirokawa J, Manabe H. Multiple
approaches to the investigation of cell assembly in memory research—present and future.
Vol. 12, Frontiers in Systems Neuroscience. Frontiers Media S.A.; 2018.

31. Pinto L, Goard MJ, Estandian D, Xu M, Kwan AC, Lee S-H, et al. Fast modulation of
visual perception by basal forebrain cholinergic neurons. 2013;16(12).

32. Dadarlat MC, Stryker MP. Locomotion enhances neural encoding of visual stimuli in
mouse V1. J Neurosci [Internet]. 2017 Apr 5 [cited 2020 Jul 29];37(14):3764–75.
Available from: https://www.jneurosci.org/content/37/14/3764

33. Abeles M, Bergman H, Gat I, Meilijson I, Seidemann E, Tishby N, et al. Cortical activity
flips among quasi-stationary states. Proc Natl Acad Sci U S A. 1995 Sep 12;92(19):8616–
20.

34. Schneidman E, Berry MJ, Segev R, Bialek W. Weak pairwise correlations imply strongly
correlated network states in a neural population. Nature [Internet]. 2006 Apr 20 [cited
2020 Aug 5];440(7087):1007–12. Available from:
https://pubmed.ncbi.nlm.nih.gov/16625187/

35. Jiang X, Shen S, Cadwell CR, Berens P, Sinz F, Ecker AS, et al. Principles of connectivity
among morphologically defined cell types in adult neocortex. Science (80-). 2015 Nov
27;350(6264).

36. Alex M. Thomson, David C. West, Yun Wang APB. Synaptic Connections and Small
Circuits Involving Excitatory and Inhibitory Neurons in Layers 2–5 of Adult Rat and Cat
Neocortex: Triple Intracellular Recordings and Biocytin Labelling In Vitro. Cereb Cortex
[Internet]. 2002 [cited 2020 Jul 29];12(9):936–53. Available from:
https://academic.oup.com/cercor/article/12/9/936/383201

37. Ko H, Hofer SB, Pichler B, Buchanan KA, Sjöström PJ, Mrsic-Flogel TD. Functional
specificity of local synaptic connections in neocortical networks. Nature [Internet]. 2011
May 10 [cited 2019 Aug 25];473(7345):87–91. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/21478872

38. Litwin-Kumar A, Doiron B. Slow dynamics and high variability in balanced cortical
networks with clustered connections. Nat Neurosci [Internet]. 2012 Nov [cited 2019 Aug
23];15(11):1498–505. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23001062

39.   Hamm JP, Peterka DS, Gogos JA, Yuste R. Altered Cortical Ensembles in Mouse Models
of Schizophrenia. Neuron [Internet]. 2017 Apr 5 [cited 2017 Jun 6];94(1):153-167.e8.
Available from: http://www.ncbi.nlm.nih.gov/pubmed/28384469

40.   Nelson SB, Valakh V. Excitatory/Inhibitory Balance and Circuit Homeostasis in Autism
Spectrum Disorders. Neuron [Internet]. 2015 Aug 19 [cited 2019 May 19];87(4):684–98.
Available from: http://www.ncbi.nlm.nih.gov/pubmed/26291155

41.   Rubenstein JLR, Merzenich MM. Model of autism: increased ratio of excitation/inhibition
in key neural systems. Genes, Brain Behav. 2003;

42.   Chen T-W, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, et al. Ultrasensitive
fluorescent proteins for imaging neuronal activity. Nature [Internet]. 2013 Jul 18 [cited
2019 Apr 17];499(7458):295–300. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/23868258

1028

1029
1030
1031
1032

1033
Figure 1. Social interaction modulates activity levels within prefrontal ensembles.
A. Mice were imaged across 9 consecutive behavioral epochs (each lasting 5 min) during which
they were either alone in their homecage or interacted with one of two novel sex-matched
juvenile mice introduced to the homecage ('M1' or '(M2'). Each novel mouse was
subsequently re-introduced to the home cage during a familiar epoch. GCaMP traces during
show examples of neurons that appear to increase or decrease activity during social epochs
(see arrows at the right of each trace).

B. Mean z-scored GCaMP traces for all neurons recorded from wild-type mice (663 neurons
from 10 mice) aligned to the onset of social interaction during the first bout of interaction
within each social epoch.

C. Cumulative plot showing the distribution of activity levels for individual neurons during
homecage epochs or periods of social interaction (percent time active in homecage: 1.8% +/-
0.1, percent time active during social interaction: 2.1 +/- 0.1%, p = 0.00002, paired t-test; n =
663 neurons from 10 WT mice).

D. Scatter-plot showing the activity of each neuron during each behavioral condition, expressed
as a percentile relative to a null distribution generated by circularly shuffling that neuron's
activity. Activity levels during social interaction or while the mouse was alone in its home
cage are plotted on the x and y axis, respectively. Kernel density plots along the axes indicate
the fraction of neurons whose activity was at a given percentile of the null distribution.
Neurons with activity > 90^{th} percentile of shuffled datasets (green dotted line) were
considered to be positively modulated, whereas neurons with activity < 10^{th} percentile (green
dotted line) were considered to be negatively modulated during each behavior (>90^{th}
percentile, social: 152/663 neurons, home cage: 80/663 neurons; p < 0.00001, chi-squared
test; <10^{th} percentile, social: 128/663 neurons, home cage: 119/663 neurons; p = 0.5, chi-
squared test).

1059
Figure 2. Classifying behavior from prefrontal ensembles using a simple neural network.
A. We constructed a neural network consisting of a single hidden layer (containing 1000 units)
which were connected to a single output unit. The thickness of lines between each hidden
layer unit and the output unit reflects the magnitude of the output weight. Positive and
negative weights are indicated by solid and dashed lines, respectively. Each hidden layer unit
received input from a random subset of prefrontal neurons from one real dataset. For clarity,
we have only shown input connections to two hidden layer units (which are differentiated by
their blue and red colors) – output weights from other hidden units are shown in black. The
output weight from each hidden layer neuron was iteratively updated during training. We
trained the classifier to distinguish periods marked as home cage exploration or social
interaction by dividing a dataset into 500-frame blocks, and then using alternating blocks for
training or testing.
B. The classifier performed poorly (near chance) when the input connection probability
(governing the number of prefrontal neurons that provided input to each hidden layer unit)
was <10%. Classification accuracy was above chance in 8/10 mice and increased to a peak of
69 +/- 3% in these mice, before decreasing again for connection probabilities >30%. The

classifier performed near chance levels when we trained and tested using data that had been
randomly swap-shuffled.

1078

Figure 3. Classifier weights reveals an ensemble that increases correlations during social behavior and detects social behavior.

A. Example histogram depicting the distribution of output weights assigned to connections between hidden layer units and the output unit over the course of training.

B. Matrix of input connections for hidden units which detect the social (left) or home cage condition (right). The hidden layer units (x-axis) have been arranged in order of increasing output weights to identify 'social units' (25 most negative output weights) and 'home cage units' (25 largest positive output weights). Prefrontal neurons (y-axis) have been arranged in order of their preference for social interaction vs. home cage, i.e. the difference between their activity levels in the two conditions.

C. Correlation matrix showing the input similarity, i.e., the pairwise correlation between binary vectors representing the input connections to each pair of hidden layer units. Hidden layer units are arranged in order of increasing output weight. Red and blue rectangles indicate correlations with social or home cage units, respectively. A gaussian filter with a standard deviation of 3 was applied to the 1000x1000 matrix to improve visualization.

D. For each hidden layer unit, we plotted its average input similarity to either the 25 social units (red) or the 25 home cage units (blue). Hidden layer units (x-axis) are again arranged by output weight. Social units had similar patterns of input compared to each other but not to home cage units and vice-versa.

E. The average input similarity of each hidden layer unit to 25 hidden layer units with near-zero output weights ('neutral units'; black rectangle in C).

F. We defined social and home-cage (HC) ensembles as the 20% of prefrontal neurons most likely to provide input to the social or home cage units, respectively. The mean activity of both home cage and social ensembles increased during social interaction compared to the home cage condition (social ensemble: mean activity level 1.4 +/- 0.3% in home cage vs. 1.8 +/- 0.3% during social interaction, $p < 0.05$, sign-rank test; home cage ensemble: mean activity level 1.5 +/- 0.30% in home cage vs. 1.9 +/- 0.3% during interaction, $p < 0.001$, sign-rank test).

G. Correlations between neurons in the same ensemble increased during social interaction for the social ensemble but for the home cage ensemble (mean correlation coefficient between neurons in the social ensemble: 0.009 +/- 0.002 in home cage vs. 0.012 +/- 0.002 during social interaction, $p < 0.05$; home cage ensemble mean correlation coefficient 0.011 +/- 0.02 in home cage vs. 0.005+/- 0.003 during social interaction, $p=0.99$, sign-rank).

1115

Figure 4. Quantification of state-dependent patterned activity with a neural network-based classifier.

A. We generated synthetic datasets consisting of 100 neurons. Each dataset consisted of two activity rasters, corresponding to states 'A' and 'B'. Each raster contained 6,000 timepoints (frames). The overall level of activity in each raster oscillated around a mean level of 5% (e.g., 5% of neurons active in a frame). In 'State A' neuronal activity was randomly assigned. Then we created 1-5 cell assemblies in 'State B' by rearranging the activity raster for 'State A'. Specifically whenever the first neuron in an assembly was active, we swapped activity in the rest of the network, so that other neurons in the same assembly would be co-active in the same frame. This was achieved by making reciprocal swaps between neurons. E.g., suppose a neuron outside the assembly was active in the desired frame. Then we swapped its activity with a neuron within the assembly that was active in a different frame. Thus neither the total number of neurons active in a given frame, nor the total activity level of any neuron differed between the 'State A' and 'State B' rasters. As a result, 'State B' was differentiated from 'State A' only by 1, 2, 3, 4, or 5 assemblies comprised of correlated (co-active) neurons.

B. A sparsely connected binary classifier perfomed above chance in distinguishing 'State A'
from 'State B' when 50% of the data was used for training, and the remainder for testing.
Inset shows the activity level of each neuron in 'State A' plotted against its activity level in
'State B.'
C. The classifier, trained on the original 'State A' and 'State B' rasters, performed at chance
levels when tested on swap-shuffled versions of the 'State A' and 'State B' rasters.
D. We again generated synthetic datasets. The 'State A' rasters were as described in panel *A*.
'State B' rasters were generated by transferring a proportion of the activity from the first 50
neurons in each State A raster to neurons 51-100. In this manner, half the neurons increased
their activity in 'State B' compared to 'State A', while the other half decreased their activity
in 'State B' compared to 'State A.' The proportion of activity transferred was determined by
randomly drawing a number with an upper bound ranging between 5 and 50%.
E. We used 50% of the data to train a sparsely connected binary classifier to distinguish 'State
A' from 'State B'. The remaining data was used for testing. Insert shows the activity level of
each neuron in 'State A' plotted against its activity in 'State B.'
F. Contrary to panel *C*, swap-shuffling these datasets (which differ in the activity levels of
individual neurons) does not disrupt classification accuracy.

1148
Figure 5. Correlations transmit additional information that is not efficiently conveyed by changes in activity levels alone.
A. Cartoon illustrating that information about behavior may be encoded through changes in activity levels, correlations between neurons, or both. When behavior modulates activity levels, correlations in two behavioral conditions may differ or be the same, and vice-versa.
B. To disentangle the roles of activity levels and correlations in transmitting information we used two different methods to create shuffled (surrogate) datasets which preserve changes in

activity levels, but either do or do not preserve patterns of correlations. We made random,
reciprocal swaps of activity between neurons to generate surrogate datasets which maintained
network activity in each frame as well as the number of blocks of activity for each neuron.
However, these datasets destroyed the correlation structure. In a second set of surrogate
datasets we used SHARC to iteratively generate surrogates in which the correlation structure
was also maintained.
C. To maintain dynamic changes in activity levels and correlations that are associated with the
two behavioral conditions we swap-shuffled or performed SHARC separately for each
behavioral epoch, then concatenated the 9 resulting surrogate subrasters to create each
surrogate dataset.
D. We trained a classifier (with a connection probability = 0.3) on each real dataset, then tested
that classifier on swap or SHARC-shuffled surrogate datasets generated from that real
dataset. Accuracy was significantly higher for the SHARC-shuffled surrogates, which
maintain the behaviorally-modulated correlations found in the original dataset. Left: accuracy
for surrogate datasets in classifying periods of home cage (HC) vs. social (Soc) behavior = 71
+/- 2 (SHARC) vs. 67 +/- 2% (swap); $p = 0.005$, paired t test. Right: accuracy for surrogate
datasets in classifying interaction with Mouse 1 vs. interaction with Mouse 2: 71 +/- 3%
(SHARC) vs. 65 +/- 2 % (swap); $p = 0.007$, paired t test. n = 10 mice.

1174

Specifically-enriched combination
observed during social behavior

In the home cage, a partially overlapping combination
is specifically enriched (but neuron C is 'left out')

Other home cage combinations can
contain the 'left-out' neuron (neuron C)

1175
Figure 6. Behaviorally-specific patterns of coactivity are formed by neurons that are active
in both conditions, but coactive with different partners in each condition.
Left: We identified combinations of 3 neurons that are specifically enriched during one
behavioral condition (occurring more often during social interaction than in 95% of surrogate
datasets, and occurring less often during home cage exploration than in 50% of surrogate
datasets, or vice-versa). We then identified overlapping combinations occurring during the
opposite behavioral condition in which a single neuron was 'left out.' In other words, we
identified combinations from the two conditions that overlapped in exactly two neurons.
Right, top: Histogram showing the number of distinct 3 neuron home cage combinations that
contain the neuron which participates in a social combination but is 'left out' during home cage
behavior. **Right, bottom:** Histogram showing the number of distinct 3 neuron social
combinations that contain the neuron which participates in a home cage combination but is 'left

out' during social interaction. In the vast majority of cases, neurons that are 'left out' in one

condition are still active during that condition and participate in other combinations.

1190

Figure 7. Shank3 KO mice have disorganized ensembles for which correlations fail to enhance the transmission of information about social behavior.

A. The mean time that *Shank3* KO mice or wild-type (WT) littermates spend interacting with a novel juvenile mouse of the same sex during a 5 min assay. Data has been pooled from 8 unimplanted WT mice as well as the 5 implanted WT mice used for microendoscopic imaging, and 5 unimplanted KO mice in addition to the 4 implanted mice used for imaging. For implanted mice we used the average of interaction time for the 2 novel mice. Pooled data showed decreased interaction in KO mice (173 +/- 12 s vs. 120 +/- 18 s for WT and KO respectively, $p < 0.05$, t-test). The un-implanted cohort alone shows a similar significant decrease in interaction time for KO mice (165 +/- 15 s vs 110 +/- 16 s for WT and KO respectively, $p < 0.05$, t-test). In the implanted cohort there was a similar trend toward decreased interaction for KO mice (186 +/- 20 s vs 133 +/- 37 s, for WT and KO respectively, $p = 0.21$, t-test).

B. (Similar to Fig. 1D). Scatter-plot showing the activity of each neuron during each behavioral condition, expressed as a percentile relative to a null distribution generated by circularly shuffling that neuron's activity. Activity levels during social interaction or while the mouse was alone in its home cage are plotted on the x and y axis, respectively. Kernel density plots along the axes indicate the fraction of neurons whose activity was at a given percentile of the null distribution. Neurons with activity > 90th percentile of shuffled datasets (green dotted line) were considered to be positively modulated, whereas neurons with activity < 10th percentile (green dotted line) were considered to be negatively modulated during each behavior**.** Data is plotted for Shank3 KO mice (red) and WT littermates (blue) (mean percentile of activity during social interaction, WT: 50 +/- 2 percentile; KO: 64 +/- 2 percentile; $p < 0.0001$ by 2-sample *t*-test; mean percentile of activity during home cage, WT: 47 +/- 2 percentile; KO: 51 +/- 2 percentile; $p = 0.1$, *t*-test).

C. Bar graph showing the fraction of neurons whose activity was positively or negatively modulated (>90th percentile or <10th percentile) during social interaction. The proportion of neurons which increased activity during social interaction was significantly greater in KO mice (22% in WT vs. 39% in KO, chi-squared = 17.7, $p < 0.0001$), whereas the downregulated ensemble was significantly smaller in KO mice (25% in WT vs. 15% in KO, chi-squared test, 8.2, $p < 0.005$). Error bars denote the binomial S.E.M. algebraically derived from total number of neurons and the proportion that were modulated in the specified direction.

D. The proportion of 3 neuron combinations occurring during home cage exploration that are enriched > the 99.9th percentile compared to swap-shuffled datasets was similar across WT (Blue; 7.6%) and KO (Red; 7.8%) mice. By contrast, the proportion of 3 neuron combinations occurring during social interaction that are enriched > 99.9th percentile compared to swap-shuffled datasets was 7.5% in WT compared to only 3.4% in KO mice (total number of home cage combinations: 4,187 in 5 WT mice, and 5,878 in 4 KO mice; total number of social combinations: 5,487 in 5 WT mice, 16,326 in 4 KO mice). The top two plots show histograms of enrichment for the home cage (upper) or social conditions (middle); the lower panel is a bar graph showing the fraction of these combinations that were specifically enriched above the 99.9th percentile (chi-squared = 165, $p < 0.0001$). Error bars denote the S.E.M. algebraically derived from the binomial distribution, the number of 3

neuron combinations in each condition, and the proportion of those combinations that were
enriched.

E. Bar graph showing how much better classifiers trained on real datasets perform whenb tested
on SHARC shuffled datasets, compared to their performance when tested on swap-shuffled
datasets, i.e. $(PerformanceSHARC - PerformanceSwap)/$
$(PerformanceSwap - 0.5)$. Each datapoint represents a classifier from one mouse. Left:
The increase in performance was significantly larger in WT mice (blue) than KO mice (right)
for classification of home cage periods (HC) vs. epochs of social interaction (Social), WT:
0.44 +/- 0.10 vs KO: 0.24 +/- 0.01, p = 0.016, Mann Whitney. Right: The increase in
performance was non-significantly larger in WT mice (blue) than KO mice (right) for
classification of interactions with Mouse 1 vs. those with Mouse 2, WT: 0.41 +/- 0.15 vs.
KO: 0.14 +/- 0.06, p = 0.11, Mann Whitney. Combined analysis of both sets of classifiers
(HC vs. Soc and Mouse 1 vs. 2) shows a significant effect of genotype: p < 0.05 by 2-way
ANOVA.

1257
Figure S1. Spatial Decorrelation of Neuronal Signals.

A. Example image (top) and individual neuron GCaMP traces (bottom) from prefrontal cortex
imaged with implanted endoscope.

B. The average GCaMP signal from a region of interest (ROI), corresponding to one neuron, was
corrected by subtracting the average GCaMP signal from the surrounding pixels, in order to
spatially deconvolve signals from each ROI vs. the surrounding neuropil. Examples traces from a
single neuron are shown.

C. The pairwise correlation matrix between signals from different neurons is shown (calculated
from 550 seconds of activity from a single wildtype mouse), for the original GCaMP signals (top
left), the surround signals (top right), the surround-subtracted signals (bottom left), and the
surround-subtracted signals after lowpass filtering (bottom right).

Encoding by Patterning of Neuronal Activity

Encoding by Modulation of Neuronal Activity

Figure S2. Patterns of correlations and the accuracy of classifications that are based on
correlated activity are both intact after SHARC shuffling

Top: As described in the main text we created synthetic datasets comprising activity of 100
neurons during two states. The activity raster for 'State B' was generated by modifying the 'State
A' raster. Specifically, in State B we 'inserted' between 1 and 5 cell assemblies (groups of
neurons with correlated activity) by reciprocally swapping activity epochs so that both the
overall level activity in each neuron, and total number of neurons active in each frame, were
unchanged between the 'State A' and 'State B' rasters. Left column: the first 200 frames of
activity for 'State A' and 'State B', followed by swap or SHARC-shuffled versions of the
original 'State B' raster. Middle column: correlation matrices for raster. Right column: accuracy
of a neural network trained to classify the original 'State A' and 'State B' rasters, tested using
either the original 'State B' raster or swap or SHARC shuffled versions. Note that SHARC, but
not swap shuffling, maintains both the pattern of correlations as well as the ability of the
classifier to accurately decode 'State A' vs. 'State B'.

Bottom: Similar to Top, but now we created the 'State B' raster from the 'State A' raster by
randomly shifting a fraction of activity from neurons 1-50 to neurons 51-100. Thus, 'State A'
and 'State B' are differentiated by the activity levels of neurons. Note that in this case the
classifier functions independently of correlations and is not affected by swap shuffling the 'State
B' raster.

1289

Mouse	Sex	Neurons	Peak Accuracy
WT			
1	f	84	0.687
2	f	39	0.705
#3	m	25	0.53
4	f	75	0.605
#5	m	37	0.459
6	m	86	0.647
7	m	74	0.848
8	m	73	0.691
9	m	71	0.694
10	f	99	0.633
KO			
1	f	83	0.772
2	m	77	0.832
3	f	45	0.34
4	f	85	0.627

Table S1. Details for each mouse included in this study.

The table shows the genotype, sex, number of imaged neurons, and peak classifier accuracy (performance when half the data was used for training and half for testing). Figure 2 showed how classifier accuracy depends on the input connection probability; because their performance was not >50% for multiple input connection probabilities, WT mice 3 & 5 (marked with #) were not included in this illustrative plot. However, we did not exclude data from these mice in any analyses. WT mice 1-5 were wild-type littermates of Shank3 KO mice. WT mice 6-10 (indicated by italics) were not littermates of Shank3 KO mice and were therefore not included as WT controls for the analyses shown in Figure 6.