Dynamic correlations help prefrontal ensembles transmit information about social behavior

Nicholas A. Frost[1,3,4,5], Anna Haggart[1,3,4,5], and Vikaas S. Sohal[2,3,4,5]*

[1]Department of Neurology, [2]Department of Psychiatry, [3]Center for Integrative Neuroscience, [4]Weill Institute for Neurosciences, [5]Kavli Institute for Fundamental Neuroscience
University of California, San Francisco

*Correspondence to: vikaas.sohal@ucsf.edu

ABSTRACT

How neurons encode behavior is a fundamental question. Neuronal ensembles increase or
decrease activity during specific behaviors. However, it is unclear whether ensembles encode
information solely via changes in activity levels, or whether changes in correlations between
neurons carry additional information. We used microendoscopic GCaMP imaging to measure
prefrontal activity while mice were either alone or engaged in social interaction. Using neural
network classifiers to measure how well prefrontal neurons transmit information about social
behavior to downstream neurons, we find that surrogate datasets which preserve dynamic
correlations outperform those which preserve ensemble activity but not correlations. Notably,
this ability of correlations to enhance the information transmitted by neuronal ensembles is lost
in mice lacking the autism-associated gene Shank3. These results show that dynamically
modulated correlations create patterns of coactive neurons which are behaviorally-specific and
enhance the information transmitted by neuronal ensembles. Furthermore, this process can be
disrupted in pathological states.

INTRODUCTION
32
During behavior, the activity of neurons is organized with precise temporal relationships
(deCharms and Merzenich, 1996; Cai *et al.*, 2016; Liang *et al.*, 2018). For example, during
certain behaviors, subsets of neurons may exhibit correlated activity in which they become active
at the same time or within short windows of time. However, it is unknown whether this sort of
temporal organization is simply a byproduct of the interconnected nature of neuronal networks
(Hebb, 1949), or contributes in a meaningful way to information encoding (Buzsáki, 2010).
Groups of co-active neurons represent an attractive computational unit for information
processing because they should optimize temporal summation in downstream targets. Thus,
increases in correlations might further augment post-synaptic responses when pre-synaptic
activity increases, or enhance post-synaptic responses even when the total level of pre-synaptic
activity remains constant.
44
However, it is unclear whether behaviorally-driven changes in correlations actually encode
additional behavioral information, beyond what is transmitted by changes in neuronal activity
levels. In particular, with the advent of new technologies for simultaneously recording from large
numbers of neurons in behaving animals, many studies have now shown that cortical ensembles
encode behavioral information via increases or decreases in the activity of their constituent
neurons. While correlations have been shown to contribute additional information for small
groups (3-8 neurons) of cortical neurons (Averbeck and Lee, 2006), only a few studies have
examined how correlations contribute to encoding within larger cortical ensembles. One study
found that the identity of a conditioned stimulus was encoded in mean activity levels, but not in
moment-to-moment patterns of co-activity (Ahmed *et al.*, 2020). Another study found that in
hippocampal region CA1, disrupting correlations impairs the decoding of position, head
direction and speed, but did not directly examine whether correlations themselves are
dynamically modulated to encode these behavioral variables (Stefanini *et al.*, 2020). In
particular, while multiple studies have shown that behavior can modulate correlations (Vaadia *et
al.*, 1995; deCharms and Merzenich, 1996) the functional significance of this has remained
unclear, because changes in correlations might simply reflect variation in activity levels (De La
Rocha *et al.*, 2007) rather than contributing additional information.
62
To address these questions, we studied the mouse medial prefrontal cortex during simple social
behaviors. The role of the medial prefrontal cortex in rodent social behavior is well-established
(Yizhar *et al.*, 2011; Brumback *et al.*, 2017; Murugan *et al.*, 2017; Selimbeyoglu *et al.*, 2017).
Many prefrontal neurons are recruited by social interaction (Brumback *et al.*, 2017; Murugan *et
al.*, 2017; Liang *et al.*, 2018) as well as social stimuli such as odors (Levy *et al.*, 2019). These
studies show that the activity levels of neuronal ensembles encode social behavior but have not
examined whether changes in correlations between prefrontal neurons transmit additional
information. Using microendoscopic GCAMP imaging in freely-moving mice, we identified
prefrontal ensembles associated with social behavior. We used a neural network classifier to
quantify how well these would transmit information about social behavior to downstream
neurons. By examining the operation of this neural network and using surrogate datasets which
preserve activity levels but either preserve or disrupt correlations, we find that changes in
correlations enhance the information transmitted by neuronal ensembles. Notably, this was not

the case in a mouse model of autism (Shank3 knockout mice), demonstrating that this form of
information transmission may be disrupted in pathological states.
78
79

80 **RESULTS**
81
Social interaction recruits prefrontal ensembles
We implanted microendoscopes (nVoke; Inscopix) into the medial prefrontal cortex (mPFC) of
adult wildtype C57/B6 mice (WT) to image calcium transients using GCaMP6f expressed under
control of the human synapsin promotor. We imaged freely moving mice during an assay which
sequentially introduced 2 novel juvenile mice to the home cage of the subject mouse, first during
an initial (novel) epoch and then again during a subsequent (familiar) epoch. These four epochs
of social interaction were interleaved with epochs during which the subject mouse was alone in
its home cage ('home cage' epochs). The first 5 minutes of each interaction epoch was scored by
a blinded observer, and each wild-type mouse spent approximately 10 minutes interacting with
the juvenile mice (393 +/- 25 s during the novel epochs and 235 +/- 18 s during the familiar
epochs, p = 0.00017, paired t test, n = 10 WT mice).
93
We processed data using a modified PCA/ICA approach (Mukamel, Nimmerjahn and Schnitzer,
2009; Luongo, Horn and Sohal, 2016) to identify neurons which were active during the imaging
session. To minimize the influence of the surrounding neuropil on neuronal signals, we
calculated the mean signal within each ROI, then subtracted the mean signal calculated from a
circular annulus surrounding each ROI (Supplemental Figure S1). Casual inspection of calcium
traces revealed that some neurons were more active during epochs of social interaction
(compared to periods of home cage exploration), whereas others exhibited the opposite pattern
(**Figure 1A**). Correspondingly, aligning calcium traces to the onset of social interaction revealed
many neurons that either increased or decreased activity at the onset of interaction (**Figure 1B**).
Fluorescence traces were converted to binary event rasters (see Methods for details of event
detection), in which most neurons were "active" in less than 5% of frames (**Figure 1C**). As a
population, imaged neurons were more active during social interaction (**Figure 1C**, n = 663
neurons from 10 mice, percent time active in home cage: 1.8% +/- 0.1, percent time active during
social interaction: 2.1 +/- 0.1%, p = 0.00002, paired t-test). There was a bimodal distribution of
cells that were significantly more (>90th percentile, social: 152/663 neurons, home cage: 80/663
neurons; p < 0.00001, Chi-Squared Test) or less active (<10th percentile, social: 128/663 neurons,
home cage: 119/663 neurons; p = 0.5) during either social interaction or matched periods when
mice were alone in their home cage, as compared to circularly shuffled datasets (**Figure 1D**).
These correspond to neuronal ensembles which are specifically recruited or inhibited during
social interaction, respectively.
114
Using a neural network classifier to assess how well ensembles transmit information
Next, we sought to determine how well these prefrontal ensembles would transmit information
about social behavior to downstream neurons. Since we and others have found that different
prefrontal neurons are recruited during social vs. nonsocial behavior, we measured how well
downstream neurons could decode whether a mouse was engaged in social behavior based on
input from these prefrontal neurons. Later we will study how this was altered in Shank3
knockout mice. For this we used a simple neural network classifier that received input from the
recorded neurons. Our rationale for using this kind of neural network classifier was threefold.
First, a simple neural network measures information that is immediately and readily available to
downstream neurons. Second, for a neural network with only one hidden layer, it is
straightforward to examine the weights to determine how the network performs the

classification. This can provide insight into exactly how the neural network is able to decode
behavior from the input activity. Third, examining how various parameters of the neural network
affect its performance can provide additional clues about how information is represented within
the input population.
130
Figure 2A shows the design of the neural network classifier. For clarity, we use the term
'neurons' specifically to refer to actual prefrontal neurons (which provide input to the neural
network), and 'units' to refer to simulated elements within the network. The network consisted of
a hidden layer containing 1000 units. We chose this number because it is both an order of
magnitude larger than the number of input neurons and an order of magnitude smaller than the
number of frames available for training (the latter helps ensure that there will be enough data to
train the output weights). We simulated a different neural network for each mouse. Each hidden
layer unit received input from a random subset of the prefrontal neurons from one mouse. I.e.,
each frame represents one timepoint and if neuron i is active in a frame then it provided an input
of 1 to all the hidden units to which it is connected; otherwise it provides an input of 0. For each
simulation, there was a fixed connection probability between each input neuron and each hidden
layer unit. We tried different values for this connection probability in order to measure how
classifier performance depends on the number of neurons that provide input to each hidden layer
unit. Each hidden layer unit had an output weight which specifies how strongly that unit excites
or inhibits a single output unit which classifies activity as belonging to periods in which a mouse
was actively engaged in social interaction or alone in its home cage. E.g., output unit activity <
0.5 corresponds to the social condition, while output unit activity > 0.5 corresponds to the home
cage condition. These output weights were adjusted during training (see Methods for details of
the training rule) while the pattern of input connectivity was fixed. This models the situation in
which prefrontal neurons transmit information to a downstream population of neurons (the
hidden layer) that decode behavior via their output weights. We initially trained networks on
50% of the data (frames) and used the held-out data for testing. We trained and tested using
intervals during which the mouse was actively engaged in social interaction or equivalent
intervals when the mouse was alone in its home cage.
155
Classifier performance is optimal for intermediate connection probabilities
Classifier performance was strongly dependent on the probability that each input neuron was
connected to each hidden unit. For the 8/10 datasets that could be classified above chance,
classifier performance (measured on the 50% of data which was held-out during training) was
near chance levels when the connection probability was < 0.1, but increased to a peak of 69 +/-
3% (**Figure 2B; Supplementary Table 1**) for a connection probability of 0.3. Accuracy
decreased dramatically when the connection probability increased to 0.5 indicating that
connection probabilities ~0.2 - 0.4 are optimal.
164
We also validated classifier performance by training and testing on surrogate datasets that were
generated by 'swap shuffling' our original datasets. We created 'swap shuffled' surrogate
datasets by randomly swapping blocks of activity between neurons (each block of activity = a set
of consecutive frames during which the neuron was active). To understand this, think of the
entire raster as a collection of blocks of activity. Each block occurs at a specific time, has a
specific duration, and is associated with a particular neuron. Swap shuffling is equivalent to just
shuffling the neurons associated with each block of activity (the start time and duration of each

block do not change). For example, if neuron *i* originally became active at time *t1* for *n1* frames
and neuron *j* was active at time *t2* for *n2* frames, then in the surrogate dataset neuron *i* might
become active at *t2* (but not at *t1*) while neuron *j* might become active at *t1* (but not *t2*). Swap
shuffling preserves the number of neurons active at each point in time (because the timing of
blocks of activity does not change). It also preserves the number of blocks of activity for each
neuron, and this tends to preserve the overall level of activity of each neuron. Activity levels are
not perfectly preserved, because blocks of activity can have different durations. Nevertheless, in
practice, blocks of activity tend to have similar durations and the similarity between the mean
activity level in each neuron before and after swap shuffling of entire datasets was 0.97 +/- 0.01.
As expected, we found that neural network classifiers trained and tested on swap shuffled
datasets performed near chance levels (Figure 2B).
183
Prefrontal neurons that drive classifier performance exhibit dramatic behaviorally-driven
changes in their correlations
Next, we examined connections in trained networks to reveal factors which enable them to
successfully classify social vs. home cage behavior (we analyzed networks with a connection
probability = 0.3 since this maximized performance of the population). Each hidden layer unit
has an output weight which measures how strongly it excites or inhibits the output unit that
represents the 'decision' (social vs. home cage). Hidden units with output weights ~0 don't
contribute to this decision. By contrast, hidden units with strong negative or positive weights
promote the social or home cage decision, respectively (**Figure 3A**). Therefore, we hypothesized
that there might be important differences in the pattern of input to hidden units, depending on
whether those hidden units have large positive or negative output weights.
195
We arranged hidden layer units based on their output weights, i.e., the unit with the most
negative weight was unit 1 and the unit with the most positive weight was unit 1000. Then we
defined the 25 hidden layer units with the most negative weights as 'social units' and the 25 with
the most positive weights as 'home cage units' (**Figure 3B**). For comparison we also defined the
25 hidden layer units with the weights closest to zero as 'neutral units.' For each pair of hidden
units, we computed the similarity between their inputs (i.e., the correlation between their input
vectors; **Figure 3C**). We then plotted the average input similarity of each hidden unit to either
the social or home cage units (**Figure 3D**) or the neutral units (**Figure 3E**). Social and home
cage units tended to receive input from the same prefrontal neurons as other hidden layer units
with the same preference, i.e., which also had negative or positive output weights. By contrast
neutral units did not exhibit any such relationship.
207
The preceding suggests that distinct ensembles of prefrontal neurons provide input to either
social or home cage hidden units. We hypothesized that there might be important features of
activity in these ensembles that support the classification of social vs. home cage behavior. For
example, one possibility is that prefrontal neurons which provide input to social units might tend
to increase activity during social behavior, whereas prefrontal neurons which provide input to
home cage units do the opposite. Surprisingly, this was not the case. In fact, both ensembles of
prefrontal neurons significantly increased their activity when mice were engaged in social
interaction (**Figure 3F**; social ensemble: mean activity level 1.4 +/- 0.3% in home cage vs. 1.8
+/- 0.3% during social interaction, $p < 0.05$, sign-rank test; home cage ensemble: mean activity
level 1.5 +/- 0.3% in home cage vs. 1.9 +/- 0.3% during social interaction, $p < 0.001$, sign-rank

test). Next, we examined pairwise correlations between the activity of prefrontal neurons within
each ensemble. Strikingly, mean correlations within the social ensemble increased during social
interaction (**Figure 3G**; (mean correlation coefficient between neurons in the social ensemble:
0.009 +/- 0.002 in home cage vs. 0.012 +/- 0.002 during social interaction, $p < 0.05$). By
contrast, there was a non-significant decrease in correlations within the home cage ensemble
(Figure 3G; home cage ensemble mean correlation coefficient 0.011 +/- 0.02 in home cage vs.
0.005 +/- 0.003 during social interaction, p=0.99, sign-rank).
225
226 Thus, the ensemble of prefrontal neurons which provide input to the social units form an
227 assembly that collectively becomes more co-active (correlated) during social behavior. In
228 contrast, the prefrontal neurons which provide input to the home cage units increase their
229 activity, but not their co-activity, during social behavior. This suggests that behaviorally-driven
230 changes in correlations may contribute to the encoding of social behavior.
231

Correlations enhance classifier performance
How can we quantitatively assess the contribution of these correlations, which are behaviorally-
modulated, to classifier performance? Ideally we would first train a neural network on the
original data. Then we would test this network's ability to classify data which maintained
behaviorally-driven changes in activity levels, but either removed or preserved the correlations.
Indeed, we have already developed methods for shuffling that achieve these goals. First, to
shuffle the data in a manner that maintains behaviorally-driven changes in activity levels, but
disrupts correlations, we can swap shuffle activity, but do so within each behavioral condition
rather than across the entire testing dataset. In other words, we first divide up the raster into
separate subrasters for each 5 minute behavior epoch (when the mouse was either engaged in
social interaction or alone in its home cage). Then we performed swap shuffling (as described
above) separately on each subraster, before recombining these swap shuffled subrasters to create
the swap shuffled surrogate dataset for testing. Because swap shuffling tends to preserve activity
levels, and because we swap shuffled activity within a behavioral condition, neurons that
increase or decrease activity during periods of social interaction in the original dataset also tend
to do so in the swap shuffled surrogate dataset.
248
To create surrogate datasets which preserve patterns of correlations as well as behaviorally-
driven changes in activity, we used a method that we published previously: SHuffling Activity to
Preserve Correlations, or SHARC (Luongo *et al.*, 2016). SHARC also re-assigns blocks of
activity between neurons, but rather than doing so randomly, it instead follows an algorithm that
achieves a target correlation matrix (in this case, the original correlation matrix) (**Figure 4B-C**).
The full details of SHARC are presented in the Methods. Briefly: on each iteration, we randomly
select one block of activity to be assigned to a new neuron. Instead of choosing the new neuron
randomly, we first compute the difference between the target correlation matrix and the
correlation matrix of the partially reconstructed surrogate dataset. Then we assign the block of
activity to the neuron which will optimally reduce this difference. Finally, to maintain the mean
activity level of each neuron, there is also an absolute limit on how many blocks of activity can
be re-assigned to each neuron. We SHARC-shuffled each social or home cage subraster
separately, then combined them to create a SHARC-shuffled surrogate dataset that preserves
behaviorally-specific levels of activity and patterns of correlations.
263

We verified that both swap and SHARC shuffled surrogate datasets preserved levels of activity
observed during both social interaction and periods when mice were alone in their home cages.
Specifically, we computed the correlation between vectors in which each element represents the
activity level of one neuron during one behavioral condition, and quantified the correlation
between each real and surrogate dataset. For swap shuffled surrogate datasets, the similarity of
activity levels (compared to real data) was 0.89 +/- 0.02 in the home cage and 0.82 +/- 0.04
during social interaction. For SHARC shuffled surrogate datasets, the similarity of activity levels
(compared to real data) was 0.88 +/- 0.03 in the home cage and 0.86 +/- 0.03 during social
interaction (n = 10 mice/datasets). We also computed the similarity of the pattern of correlations
between each surrogate dataset and the corresponding real dataset. In this case, only SHARC
shuffled surrogate datasets preserved patterns of correlations. For swap shuffled surrogate
datasets, the similarity of correlations to the real data was 0.01 +/- 0.01 in the home cage, and
0.03 +/- 0.01 during social interaction. For SHARC shuffled surrogate datasets, the similarity
was 0.50 +/- 0.05 in home cage and 0.55 +/- 0.03 during social interaction.
278
We then trained classifiers on each dataset and tested each classifier using either swap or
SHARC shuffled surrogate datasets generated from the same dataset using for training (**Figure
4C**). Classifiers performed better than chance when tested with either surrogate dataset
indicating that changes in activity levels encode behavioral information. However, performance
was significantly higher for SHARC shuffled surrogates datasets than for swap shuffled ones
(**Figure 4D**; classifier accuracy for SHARC shuffled surrogate datasets = 68 +/- 4%, classifier
accuracy for swap shuffled surrogate datasets = 61 +/- 4%, p < 0.05, sign-rank test). This
demonstrates that behaviorally-modulated patterns of correlations transmit additional
information, beyond what is readily decodable from activity levels alone.
288
Combinations of coactive neurons occur in a behaviorally-specific manner
Interestingly, neural networks perform classification better for connection probabilities ~0.2 –
0.4 than for connection probabilities < 0.1. When the connection probability is low, each hidden
unit receives input from individual prefrontal neurons or small groups of neurons. By contrast,
when the input probability is higher, hidden units receive input from larger groups of prefrontal
neurons. This suggests that training proceeds more efficiently when the network represents
information about social vs. home cage behavior using multineuron combinations, instead of
activity within individual neurons or small groups. Together with the fact that classifier
performance was higher for SHARC shuffled datasets than swap shuffled ones, this indicates that
multineuron patterns of coactivity, rather than just levels of activity within neuronal ensembles,
transmit information about social behavior. Therefore as a proof-of-concept, we directly
examined whether 3-neuron patterns of coactivity occur in a behaviorally-specific manner. We
examined 3-neuron combinations because they measure network structure beyond pairwise
correlations and are the building blocks of larger combinations. One could in principle analyze
larger combinations, but because of the limited numbers of neurons and frames in our datasets,
there is not always adequate statistical power to resolve larger combinations, i.e., to detect large
numbers of combinations that occur more often in real datasets than expected by chance.
306
First, we quantified how often each possible 3-neuron combination occurred in real datasets.
Then we calculated how often each of these combinations occurred in datasets that had been
swap-shuffled (across the entirety of the dataset). For each real dataset we constructed 1000

swap-shuffled datasets, and identified 'enriched combinations,' which occurred more often in
real datasets than in 95% of swap shuffled surrogate datasets. Enriched combinations are those
which occur more often in real datasets than expected based on the chance overlap of activity
between marginally independent neurons. Finally, we quantified how many of these enriched
combinations were behaviorally-specific, i.e., occurred exclusively during social or home cage
epochs. Combinations could appear to be behaviorally-specific simply because they only
occurred at a single timepoint. Therefore we also restricted our analysis to enriched combinations
which occurred during multiple distinct bouts of social interaction and/or matched sets of
intervals during home cage epochs. Many of these repetitively-occurring enriched combinations
were behaviorally-specific: 43.5% occurred during social interaction, 26.5% during home cage
epochs, and 30% during both conditions.
321
The selective occurrence of enriched combinations either during social interaction or when a
mouse is alone in its home cage may reflect changes in single neuron activity (i.e., neurons that
form a social combination are only active during the social condition), and/or changes in
correlations (i.e., neurons are active in both conditions but only *co-active* during social
behavior). To test the hypothesis that changes in correlations underlie the behavioral specificity
of significant combinations, we examined the 3-neuron combinations that were specifically
enriched during either periods of home cage exploration or social interaction (**Figure 5**). We
defined specific enrichment as those combinations which occurred more often in real data than in
95% of swap-shuffled surrogate datasets for one behavioral context, and less in real data than in
50% of swap-shuffled surrogate datasets for the other behavioral context. Based on these criteria,
12,408 3-neuron combinations were specifically enriched during social interaction, and 9,572
were specifically enriched during home cage exploration. There were 55,696 instances in which
a social and nonsocial 3-neuron combination overlapped in 2 out of 3 neurons. In 97.0% of these
cases, the neuron which was part of a social 3-neuron combination (triplet) but left out of the
overlapping home cage triplet was part of a different 3-neuron combination that was enriched
during homecage exploration (Figure 5, top right). Conversely, the neuron which was part of a
nonsocial triplet but left out of the overlapping social 3-neuron combination was part of a
different socially-enriched 3-neuron combination in 99.1% of cases (Figure 5, bottom right).
Overall, an average of 71 enriched homecage combinations contained the neuron missing from
the social triplet, and 85 enriched social combinations contained the neuron missing from
homecage triplets. Thus, the specificity of a combination of co-active neurons for social vs.
nonsocial behavior does not occur simply because some neurons were only active during one
condition, but rather reflects the dynamic reorganization of patterns formed by neurons which are
active in both conditions, i.e., changes in correlations. This – the behaviorally-specific
occurrence of multineuron patterns of coactivity – represents the substrate through which
correlations can add to the behavioral information transmitted by neuronal ensembles.
348
Socially-enriched combinations are deficient in Shank3 KO mice
We were curious whether there might be conditions under which these phenomena – the
occurrence of multineuron combinations of coactivity during social behavior, and the ability of
correlations to enhance the transmission of information about social behavior – might be
impaired. To explore this, we performed microendoscopic GCaMP imaging in mice lacking the
autism-associated gene Shank3 (Peça *et al.*, 2011; Duffney *et al.*, 2015; Chen *et al.*, 2020). These
mice have been extensively studied as models of Phelan-McDermid syndrome, which often

includes autism as a clinical feature. Shank3$^{-/-}$ (KO) mice are known to have social deficits, and indeed, we found that compared to wild-type (WT) littermates, they spend significantly less time interacting with novel juvenile mice (**Figure 6A**).

We compared patterns of prefrontal activity in Shank3 KO mice and their WT littermates. As in WT mice, in Shank3 KO mice, many prefrontal neurons either increase or decrease activity during social interaction. However, compared to WT mice, the fraction of neurons whose activity increases during social interaction was significantly higher, whereas the fraction whose activity decreases was significantly lower (**Figure 6B-C;** 22% of 260 WT neurons vs. 39% of 290 KO neurons increased activity above the 90th percentile of shuffled data during social interaction, chi squared = 17.7, p < 0.0001; 25% of WT vs. 15% of KO neurons decreased activity below the 10th percentile of shuffled data during social interaction, chi squared = 8.2, p < 0.0001). Thus, Shank3 KO mice recruit abnormal neuronal ensembles during social behavior. We hypothesized that this might reflect a network-level disorganization that affects the normal patterning of activity during social behavior.

Indeed, we found that in KO mice a significantly smaller fraction of the 3-neuron combinations observed during social interaction were strongly enriched, i.e., occur more often in actual data than in 99.9% of swap-shuffled surrogate datasets (**Figure 6D**). This suggests that even though more neurons (i.e., larger ensembles), were recruited during social behavior in KO mice, these may have been less well-organized, such that the occurrence of socially-enriched patterns of activity is obscured by 'noise,' i.e., patterns formed by the chance overlap of activity between neurons that fire in a largely independent fashion. Notably, this deficiency was specific for social interaction. The fraction of 3-neuron combinations that were strongly enriched during home cage exploration (in comparison to swap-shuffled surrogate datasets) was similar in WT and KO mice (Figure 6D).

Correlations do not enhance the transmission of information about social behavior in Shank3 KO mice

The preceding shows that even though social behavior robustly recruits neuronal ensembles in Shank3 KO mice, the organization of these ensembles into multineuron combinations is disorganized. This suggests that the ability of patterns of co-activity to encode information about social behavior may be impaired in these mice. To test this, we directly examined whether correlations contribute to the transmission of information about social behavior in Shank3 KO mice. As before, we generated swap and SHARC shuffled surrogate datasets, then tested the ability of classifiers trained on the original datasets (from Shank3 KO mice) to classify activity associated with behavior during social interaction vs. in home cage. While we still observed above chance classification accuracy using a classifier with a connection probability of 0.3, there was no longer an increase in performance when correlations were preserved in SHARC shuffled surrogate datasets as compared to swap shuffled ones (**Figure 6E**; classifier accuracy: 62 +/ 4% for SHARC vs 63 +/- 2% for swap shuffled surrogate datasets, p = 0.47, sign-rank test). Thus, in Shank3 KO mice, multineuron patterns of coactivity during social behavior are disturbed, and correlations no longer add to the information about social behavior transmitted by prefrontal ensembles.

DISCUSSION

During complex behaviors, the brain can use many strategies to represent information about the external environment and internal state of the organism. The term 'ensemble' is often used to refer to a group of neurons whose activity is similarly modulated (either increased or decreased) during specific behaviors (Sakurai, 1999; Cai *et al.*, 2016; Sakurai *et al.*, 2018; Corder *et al.*, 2019; Ghandour *et al.*, 2019; Gründemann *et al.*, 2019). It is generally accepted that ensembles transmit behavioral information via changes in the activity levels of their constituent neurons. On the other hand, many studies have also shown that correlations between neurons can change during specific behaviors (Vaadia *et al.*, 1995; deCharms and Merzenich, 1996) or behavioral states (Abeles *et al.*, 1995; Pinto *et al.*, 2013; Dadarlat and Stryker, 2017). Importantly, correlations reflect changes in coactivity which exceed those expected to occur simply because of changes in the activity levels of the individual neurons (De La Rocha *et al.*, 2007). I.e., when an ensemble becomes more active, its correlations could go up, down, or remain unchanged. By optimizing synaptic interactions such as temporal summation, changes in correlated activity could potentially enhance the behavioral information transmitted by changes in ensemble activity, or transmit entirely different types of information, e.g., about internal states. Correlations have been studied extensively for the isolated retina responding to visual stimuli (Schneidman *et al.*, 2006). However, how correlations in recurrently connected cortical circuits such as the mPFC encode behavior has been more difficult to discern.

Here, we addressed this question using microendoscopic GCaMP imaging to measure activity from many (~40-100) prefrontal neurons during social behavior in mice. We used multiple approaches to disentangle the respective contributions of activity and correlations to the encoding of behavior. First, we used a simple neural network, in which prefrontal neurons provide input, there is one hidden layer, and a single output unit classifies social vs. nonsocial behavior, to quantify how well prefrontal ensembles would transmit information about social behavior to downstream neurons. Notably, classifier performance was at chance levels when hidden layer units only received input from one or a few prefrontal neurons, but was significantly higher when hidden units combined activity from 20-30% of prefrontal neurons. This suggests the network can be trained to discriminate social vs. nonsocial behavior most efficiently when each output weight corresponds to the activity in a large ensemble (representing ~20-30% of the network), rather than the activity of an individual prefrontal neuron or a small group.

Next, we extended a method we previously published, (Luongo *et al.*, 2016), to non-randomly shuffle datasets in order to preserve both behaviorally-modulated correlations and ensemble activity. This enabled us to compare the amount of information about social behavior transmitted by either SHARC-shuffled surrogate datasets or randomly-shuffled surrogates which preserved ensemble activity but not correlations. In this way, we found that correlations enhance the amount of information that prefrontal ensembles transmit about social behavior. Indeed, when we examined connections within neural network classifiers, we found that prefrontal neurons which serve to detect social behavior increase their correlations during social behavior (whereas neurons which detect nonsocial behavior do not).

Correlations measure neuronal coactivity that occurs more often than expected based on the chance overlap of activity between neurons. Thus, in accordance with our finding that behavior

447   modulates correlations, we found that multineuron patterns of coactivity which occur more often
448   than expected by chance are behaviorally-specific. We then directly examined these
449   behaviorally-specific and statistically-enriched combinations of coactive neurons. We found that
450   they tend to be composed of neurons which are active in both conditions but only coactive in
451   one, rather that neurons which are only active in one condition.
452
Interestingly, these statistically-enriched patterns of coactivity were specifically deficient during
social behavior in mice lacking the autism-associated gene Shank3. Accordingly, in Shank3 KO
mice, surrogate datasets which preserve behaviorally-modulated correlations failed to transmit
more information about social behavior compared to randomly shuffled datasets which only
preserved ensemble activity. This shows that the ability of correlations to enhance the
transmission of information about social behavior is not automatic, and can in fact be disrupted
in pathological states.
460
461   Similar to other recent studies (Ahmed *et al.*, 2020; Stefanini *et al.*, 2020), we have studied
462   activity using binary activity rasters derived from GCaMP imaging. However, an important note
463   is that any method of quantifying neural activity has limitations, such that there could be
464   additional ways that neurons encode information which are not well resolved using this
465   approach.
466
What is the meaningful size of ensembles in the cortex?
Complex behavior is possible because the brain reliably encodes features pertaining to the
external environment as well as the internal state of the organism. These features may be
encoded by the modulation of activity in neuronal ensembles (Sakurai, 1999; Cai *et al.*, 2016;
Sakurai *et al.*, 2018; Corder *et al.*, 2019; Ghandour *et al.*, 2019; Gründemann *et al.*, 2019). How
many neurons are needed to reliably encode an aspect of behavior? This is an important question
because the capacity, robustness against noise, generalization ability, etc., of a network depend
on how many neurons encode specific pieces of information.
475
We explored this question, not by measuring actual connections, but rather by asking what input
connection probability would optimize the ability of a downstream network to classify behavior
based on input from prefrontal ensembles. Note: input connections in neural network classifiers
do not necessarily correspond to actual connections in the brain – rather they provide information
about the size and nature of groups of neurons across which information should be combined to
most efficiently decode behavior. Peak classifier performance occurred for connection
probabilities ~0.2 - 0.3. Performance was markedly lower when the connection probability was
0.5. This is surprising because a connection probability of 0.5 would maximize the entropy of
each connection; correspondingly, the number of distinct input combinations to a hidden unit is
maximized when it receives connections input from half the input neurons. Thus, from the
standpoint of encoding social behavior, combining activity from 20-30% of the input neurons
may achieve some synergy that becomes degraded when the activity of additional neurons is
included. This suggests that whatever mechanism normally generates behaviorally-meaningful
patterns of coactivity in prefrontal neurons, the size of these patterns is limited to about 20-30%
of the network. In the brain, nonrandom network connectivity (Alex M. Thomson, David C.
West, Yun Wang, 2002; Jiang *et al.*, 2015) may similarly produce correlated activity / coactivity

within neurons that represent a specific subset of a larger population (Ko *et al.*, 2011; Litwin-
Kumar and Doiron, 2012).
494
Combinatorial codes vs. sequential patterns of activity
Like many recent studies, we measured population-level activity in the mouse neocortex using
genetically encoded calcium indicators. These indicators transduce neuronal spiking on
timescales ~100 msec. Thus correlated activity / 'coactivity' imply that neurons jointly increase
their activity within windows ~100 msec, and do not necessarily imply synchronous spiking on
faster timescales (milliseconds or even tens of miliseconds). At the same time, correlated activity
/ coactivity on these timescales should be differentiated from sequential activity of neurons
observed during the performance of sequential behaviors (i.e. spatial navigation or overtrained
tasks) in which the activity of specific neurons corresponds to moving through a specific location
or performing a specific portion of a complex task. As discussed above, in the neocortex
correlations and coactivity likely reflect recurrent neural network connectivity (Ko *et al.*, 2011).
By contrast, sequential patterns of neuronal activation can occur simply as a byproduct of the
arrangement of spatial locations along a trajectory, the stereotyped order in which cues are
encountered during a task, etc.
509
Relevance to disease states
Interestingly, in *Shank3* KO mice, which exhibit social deficits, the mPFC successfully recruits
specific neuronal ensembles during social interaction. However these ensembles are enlarged,
their organization into statistically-enriched patterns of coactivity is disrupted, and correlations
between neurons fail to enhance the information that these ensembles transmit. Thus, the
computational units by which information is processed in the mPFC appears to be inefficient,
i.e., social behavioral recruits an abnormally large number of neurons at the expense of the
precise temporal patterning of this activity. This central finding is similar to other findings in
rodent models of autism at both the single neuron and network levels (Hamm *et al.*, 2017; Levy
et al., 2019; Chen *et al.*, 2020). In particular, we found an increase in the recruitment of
prefrontal neurons during social interaction. This mirrors a recent study which found
hyperdynamic response to whisker stimulation in the same mice (Chen *et al.*, 2020), possibly
reflecting GABAergic circuit dysfunction and/or homeostatic compensations (Nelson and
Valakh, 2015). (Note: these findings cannot be ascribed simply to the fact that Shank3 KO mice
spend less time engaged in social interaction than their wild-type littermates; reduced interaction
time would tend to reduce statistical power and thereby reduce the number of neurons that
change their activity more than expected by chance.)
527
Increased excitatory activity causing decreased signal-to-noise ratio (SNR) has long been posited
to contribute to the pathophysiology of autism (Rubenstein and Merzenich, 2003). However the
exact nature of 'signal' and 'noise', and the specific mechanism through which excessive activity
degrades the SNR have been unclear. Here, we show how enlarged neural ensemble recruitment
by specific behavioral conditions disrupts information transmission by degrading the ratio
between statistically meaningful patterns of coactivity (the signal) and the random overlap of
activity between neurons (noise).
535

METHODS

Behavior: C57/B6 mice were obtained from Jackson Laboratories. We utilized adult mice of either sex housed and bred in the UCSF animal facility. Adult mice were habituated to the room and observer for 3 days prior to test day. All videos were subsequently scored by a blinded observer. For imaging experiments, 5 WT and 4 KO littermates were generated through crosses between Shank3 heterozygous parents and injected with AAV5.Syn.GCaMP6f.WPRE.SV40. We included an additional 5 WT mice which were injected with AAV5.Syn.GCaMP6m.WPRE.SV40 (Chen *et al.*, 2013). Viruses were obtained from Penn Viral Core. Injections and 500 um GRIN lens (Inscopix) implantations were carried out in 8-12 week old mice to express GCamp6f in prefrontal cortical neurons under control of the human Synapsin promotor. Mice were anesthetized with 2% isoflurane and mounted in a stereotactic frame. Craniotomies were made according to stereotaxic coordinates relative to Bregma. Coordinates for injection into mPFC were (in mm relative to Bregma): +1.7 anterior–posterior (AP), –0.3 mediolateral (ML) and –2.75 dorsoventral (DV), and GRIN lenses were implanted at the same AP and ML coordinates, to a depth of 2.25. We subsequently attached baseplates for attaching the microendoscope, ~4 weeks later depending on GCamp expression. Mice were habituated for three days with the scope attached, prior to test day. On test day, mice were habituated with the scope turned on, then imaged in alternating home cage and social epochs. During social epochs, one of 2 novel sex-matched juvenile mouse was introduced to the test mouse's homecage, in sequential order so that there were two 'novel' epochs, followed by two 'familiar' epochs interleaved with 'home cage' epochs during which the juvenile mice were removed and the test mouse was free to explore its home cage. The first and last home cage epoch were 10 minutes in length; the others were 5 minutes in length. Each social epoch lasted 10 minutes but only the first 5 minutes were recorded and scored. During each behavioral epoch, observer was not in the room. Interaction epochs were defined from the moment test mouse first sniffed the juvenile conspecific or object, until the test mouse turned away. Videos were recorded using Anymaze, and scored by a blinded observer. For the bulk of analysis we pooled data across 10 WT mice. Shank3 KO mice were compared only to recordings from WT littermates.

Image acquisition and segmentation: Images were acquired using an Inscopix nVoke microendoscope attached to a laptop computer and synced to a separate video acquisition computer running Anymaze. Frame rate was 20 Hz and the laser power was 0.2 mW. Acquisition was performed using 2x2 pixel binning, then subsequently downsampled again by 2.

We segmented neuronal signals using a modified PCA/ICA approach (Mukamel, Nimmerjahn and Schnitzer, 2009; Luongo, Horn and Sohal, 2016), modified so that each segment was expressed as a binary ROI consisting of pixels representing a single neuron. I.e., we used the output from the PCA/ICA to identify a set of contiguous pixels which represent a neuron, then averaged fluorescence signals across those pixels. To deconvolve neuronal signals from background neuropil signals, we also subtracted the mean signal from each identified segment from the mean value in pixels surrounding the edge of the segment (we excluded pixels that belonged to another ROI). Signals were subsequently lowpass filtered to remove high frequency noise using the Matlab command: designfilt('lowpassfir', 'PassbandFrequency', 0.5, 'StopbandFrequency', .65, 'PassbandRipple', 1, 'StopbandAttenuation', 25). All signal traces shown represent normalized versions of the *dF*/*F$_0$* trace, where *F$_0$ is estimated by the median*

value in the surround region. Threshold based event detection was performed on the traces by
detecting increases in dF/F_0 exceeding 3σ over one second, then only keeping those events
which exceeded a 15σ increase over two seconds, and a total area under the curve of 250σ. As
there were occasional downward deflections due to surround subtraction, we instituted a final
parameter requiring that the peak cross an absolute value of $dF/F_0 = 0.0125$. σ is the standard
deviation of dF/F_0, calculated over the least active 50% of the movie. In some cases these
parameters were adjusted slightly to optimize event detection to > 95% sensitivity and
specificity, based on visual inspection, for each movie. After identifying these events in the
GCaMP signal from a cell, the cell was considered "active" during the entire period from the
beginning of an event until the GCaMP signal decreased 30% from the peak of the event, up to a
maximum of 2 seconds. The peak of the event was defined as the local maximum of the entire
event, from the beginning of the event until dF/F_0 returned to the pre-event baseline value.
Calcium traces from segmented neurons were visually inspected and a small number of segments
were removed if they did not appear to represent a single, unduplicated neuron. We restricted
further analysis to those mice with 25 or more active neurons. We then created 2-dimensional
event rasters consisting of detected events for each neuron over the course of the experiment.
598
<u>Detection of behaviorally modulated neurons</u>: To determine the response of individual neurons
to behavioral context, we averaged the activity of each neuron during frames corresponding to
periods of social interaction, or to a temporally matched set of frames during the preceding home
cage epoch. We then created a 'null distribution' for each neuron that represents the percent of
time active expected in each condition based on chance, by circularly shuffling the data 10,000
times. We then compared the activity of each neuron during either social interaction or home
cage exploration to this null distribution. Neurons were considered positively modulated if they
exceeded the 90[th] percentile of that observed in circularly-shuffled datasets, and negatively
modulated if the percent of frames that a neuron was active during a given context was below the
10[th] percentile of observations from circularly-shuffled data.
609
<u>SHARC</u>: SHARC (<u>SH</u>uffling <u>A</u>ctivity to <u>R</u>earrange <u>C</u>orrelations) is an iterative method for
generating surrogate datasets. SHARC nonrandomly shuffles blocks of activity within a raster to
generate a new (surrogate) raster in which the pairwise correlations between neurons match a
target correlation matrix (Luongo, Horn and Sohal, 2016). Here we apply this previously-
published method, with modifications to also preserve the activity level in each neuron (Figure
4B).
616
To begin, note that each raster is equivalent to a collection of blocks of activity. Each block of
activity is defined by the time at which it begins, its duration, and the neuron which is active. On
each iteration one block of activity is randomly chosen and assigned to a new neuron as follows.
Suppose block i has been chosen to be re-assigned. First, we find all the blocks of activity that
overlap with block i. Next, we selected the subset of these blocks for which new cell identities
had already been assigned. Call this set X. Let r_j represent the number of timepoints over which
block $j \in X$ overlaps with block i, and let n_j represent the identity of the cell assigned to block $j \in$
X. L_i and L_j are the lengths of blocks i and j, respectively. Then we constructed a vector,

625
$$\vec{P}_i = \sum_{j \in X} \frac{r_j}{\sqrt{L_i L_j}} \left(\vec{C}_{n_j} - \vec{C'}_{n_j} \right)$$

where $\vec{C}_{n_j}$ represents row j of the target correlation matrix, i.e. the target correlations between

neuron n_j and the other neurons, and $\vec{C'}_{n_j}$ contains the current values of the correlations between

neuron n_j and the other neurons based on the blocks of activity that have already been re-

assigned. This step can be thought of as "guessing" which cell *should* be assigned to a particular

block of activity by first figuring out what other cells are active at the same time, then choosing

cells which are strongly correlated with these known active cells. Note that we assign values of

$\vec{P}_i$ (i.e., construct "guesses" about which cell should be active), using the *difference* between the

current correlation matrix ($\vec{C'}_{n_j}$) and the target correlation matrix ($\vec{C}_{n_j}$), in order to identify cell

pairs for which the current correlation deviates from the target value, and force the new

correlation matrix to progressively approximately the target correlation matrix.

We set elements of $\vec{P}_i$ to zero if the corresponding neuron had already been assigned to a block

of activity that overlaps with block i, i.e. element n_j of $\vec{P}_i$ was set to zero $\forall j \in X$. Finally, we

assigned block i to the neuron corresponding to the maximum value of $\vec{P}_i$. This can be thought of

as choosing the cell that represents the "consensus" based on tallying up all of the "guesses"

about which cells "should" be assigned to the block of activity being considered.

When all the elements of $\vec{P}_i$ were zero, e.g. because there no overlapping blocks of activity have

had new cell identities assigned yet, then we chose a cell in order to match the originally

observed level of activity. Specifically, after every iteration, we kept a log of the net number of

blocks of activity that each neuron had donated or received. We used this vector to create a

weighted probability whereby events from neurons which had received a net positive number of

blocks were more likely to be chosen to be reassigned. To further ensure that the total number of

active events for each neuron in the surrogate dataset was similar to the real dataset, if the

difference between the number of blocks gained – lost in the reassignment process exceeded +10

for a particular neuron, then that neurons was no longer eligible to receive additional blocks of

activity.

We extended this approach to generate surrogate datasets by shuffling data within shorter time

windows (i.e., individual behavioral epochs). Here a discrete set of frames is chosen,

corresponding to a subraster of the original raster. By repeating the process described above for

each subrasters, then recombining the shuffled subrasters, we generate a complete shuffled

dataset.

<u>Classifier</u>: We designed and trained a neural network to classify behavior (periods when a mouse

was alone in its home cage vs. engaged in social interaction). This network contained 1000 units

in a hidden layer, each of which received input from specific prefrontal neurons (from the real

dataset). Thus, in each frame the activity of each hidden layer unit was just the summed activity

of the connected prefrontal neurons. Each hidden layer unit had an output weight that

represented the strength of its connection to a single output unit. On each frame the activity of

the output unit was computed as:

667
$$y = \frac{1}{1 + e^{-\sum w_i x_i}}$$

where w_i is the output weight from hidden unit i and x_i is the activity of hidden unit i.

When we performed training and testing using the same dataset, we divided the dataset into
alternating blocks of 500 frames for training vs. testing (in other cases we used the real dataset
for training, then tested using a surrogate dataset). We restricted training or testing to frames in
which mice were scored as actively engaged in social interaction (or matched frames during
periods when the mouse was alone in its home cage). We also limited training / testing to frames
with at least 3 active neurons.

We trained the output weights by performing 500 passes through the training data (each pass
visited all of the training frames in a random order). On each training timestep, we calculated y,
the activity of the output unit, and then adjusted each output weight based on:

$$\Delta w_i = \varepsilon y(1 - y)(y - z)x_i$$

where z is the correct classification of the frame (0 for social behavior, 1 for home cage) and ε,
the learning rate, was set to 0.05.

Following training, we examined the pattern of input connections and output weights. The
distribution of output weights was roughly gaussian and centered near 0. We identified the
selection of prefrontal neurons most likely to be connected to hidden layer units with large
positive or negative weights. Hidden layer units with large negative or positive output weights
bias classification towards the social or home cage condition, respectively. Therefore, we refer to
the 25 hidden units with the most negative or positive weights as 'social' or 'home cage' units
respectively. We calculated the number of input connections between each prefrontal neuron and
the 25 home cage units or 25 social units. We then defined 'home cage' or 'social' ensembles as
the 20% of prefrontal neurons with the most input connections to home cage or social units,
respectively. As described in the main text, we then analyzed properties of these two ensembles.

Quantification of multineuron combinations: Estimating chance overlap between activity of
largely independent neurons requires accounting for two factors. First, neurons with higher
activity are more likely to overlap by chance with other neurons. Second, overall network
activity is dynamic over time, creating a tendency for otherwise independent neurons to be
recruited at similar times. Thus, it is necessary to identify combinations which occur more often
than expected based on 1) the activity levels of the constituent neurons, and 2) the fact that
activity in a network is not constant over time. We can do this by quantifying the occurrence of
combinations in datasets which have been shuffled to preserve 1) the overall level of activity in
each neuron, and 2) the total level of activity in the network at each point in time.

3 neuron combinations were quantified by identifying each combination present in frames in
which 2 or more neurons were active. The number of frames each combination was active in
real data was stored in a n-dimensional matrix. Surrogate datasets were then generated from
event rasters by *swapping* the identity of neurons associated with detected events (periods of
activity). As the timing of events themselves is unchanged, and only the identity of the
participating neurons are exchanged, this preserves both the number of events per frame and the
number of events that each neuron participates in. Therefore, the total number of combinations in

each frame and over the course of the experiment (i.e., the sum of occurrences across all
combinations) is also preserved. The total number of combination occurrences in which a given
neuron participates would also tend to be preserved in these swap-shuffled surrogate datasets.
718
We then quantified how often each combination occurred in real vs. swap-shuffled data. By
comparing how often each combination occurred in real data vs. in 1,000 swap-shuffled
surrogate dataset, we were able to quantify how 'enriched' each combination was, compared to
the level of occurrence expected by chance based on the activity levels of its constituent neurons
(and the overall temporal pattern of network activity). We expressed enrichment as a percentile,
calculated relative to swap-shuffled surrogate data, e.g., the 100^{th} percentile indicates that a
particular combination occurred more often in real data than in all 1,000 surrogate datasets.
Further analysis was restricted to 'enriched combinations', i.e., combinations that occurred more
often in real datasets than in 95% of surrogate datasets.
728
729   <u>Statistical analysis</u>: Neurons and significant combinations from all animals and groups were
730   pooled and counted as single units. Proportions were compared using chi-squared test. Activity
731   levels were compared using paired *t*-tests (2-sided), unless otherwise noted. Where applicable,
732   error bars denote standard error. Values of the classifier performance (accuracy) were generated
733   by averaging after re-running the training / testing procedure at least 25 times.
734

REFERENCES

Abeles, M. *et al.* (1995) 'Cortical activity flips among quasi-stationary states', *Proceedings of the National Academy of Sciences of the United States of America*. National Academy of Sciences, 92(19), pp. 8616–8620. doi: 10.1073/pnas.92.19.8616.

Ahmed, M. S. *et al.* (2020) 'Hippocampal Network Reorganization Underlies the Formation of a Temporal Association Memory', *Neuron*. Cell Press, 107(2), pp. 283-291.e6. doi: 10.1016/j.neuron.2020.04.013.

Alex M. Thomson, David C. West, Yun Wang, A. P. B. (2002) 'Synaptic Connections and Small Circuits Involving Excitatory and Inhibitory Neurons in Layers 2–5 of Adult Rat and Cat Neocortex: Triple Intracellular Recordings and Biocytin Labelling In Vitro', *Cerebral Cortex*, 12(9), pp. 936–953. Available at: https://academic.oup.com/cercor/article/12/9/936/383201 (Accessed: 29 July 2020).

Averbeck, B. B. and Lee, D. (2006) 'Effects of noise correlations on information encoding and decoding', *Journal of Neurophysiology*. American Physiological Society, 95(6), pp. 3633–3644. doi: 10.1152/jn.00919.2005.

Brumback, A. C. *et al.* (2017) 'Identifying specific prefrontal neurons that contribute to autism-associated abnormalities in physiology and social behavior', *Molecular Psychiatry*. doi: 10.1038/mp.2017.213.

Buzsáki, G. (2010) 'Neural Syntax: Cell Assemblies, Synapsembles, and Readers', *Neuron*, 68(3), pp. 362–385. doi: 10.1016/j.neuron.2010.09.023.

Cai, D. J. *et al.* (2016) 'A shared neural ensemble links distinct contextual memories encoded close in time', *Nature*. doi: 10.1038/nature17955.

Chen, Q. *et al.* (2020) 'Dysfunction of cortical GABAergic neurons leads to sensory hyper-reactivity in a Shank3 mouse model of ASD.', *Nature neuroscience*. Nature Publishing Group, pp. 1–13. doi: 10.1038/s41593-020-0598-6.

Chen, T.-W. *et al.* (2013) 'Ultrasensitive fluorescent proteins for imaging neuronal activity', *Nature*, 499(7458), pp. 295–300. doi: 10.1038/nature12354.

Corder, G. *et al.* (2019) 'An amygdalar neural ensemble that encodes the unpleasantness of pain', *Science*. American Association for the Advancement of Science, 363(6424), pp. 276–281. doi: 10.1126/science.aap8586.

Dadarlat, M. C. and Stryker, M. P. (2017) 'Locomotion enhances neural encoding of visual stimuli in mouse V1', *Journal of Neuroscience*. Society for Neuroscience, 37(14), pp. 3764–3775. doi: 10.1523/JNEUROSCI.2728-16.2017.

deCharms, R. C. and Merzenich, M. M. (1996) 'Primary cortical representation of sounds by the coordination of action-potential timing', *Nature*, 381(6583), pp. 610–613. doi: 10.1038/381610a0.

Duffney, L. J. *et al.* (2015) 'Autism-like Deficits in Shank3-Deficient Mice Are Rescued by Targeting Actin Regulators', *Cell Reports*. doi: 10.1016/j.celrep.2015.04.064.

Ghandour, K. *et al.* (2019) 'Orchestrated ensemble activities constitute a hippocampal memory

engram', *Nature Communications*. Nature Publishing Group, 10(1), p. 2637. doi:
10.1038/s41467-019-10683-2.

Gründemann, J. *et al.* (2019) 'Amygdala ensembles encode behavioral states.', *Science (New*
York, N.Y.). American Association for the Advancement of Science, 364(6437), p. eaav8736.
doi: 10.1126/science.aav8736.

Hamm, J. P. *et al.* (2017) 'Altered Cortical Ensembles in Mouse Models of Schizophrenia',
Neuron, 94(1), pp. 153-167.e8. doi: 10.1016/j.neuron.2017.03.019.

Hebb, D. O. (1949) *The organization of behavior: A neuropsychological theory*. New York:
Wiley.

Jiang, X. *et al.* (2015) 'Principles of connectivity among morphologically defined cell types in
adult neocortex', *Science*. American Association for the Advancement of Science, 350(6264).
doi: 10.1126/science.aac9462.

Ko, H. *et al.* (2011) 'Functional specificity of local synaptic connections in neocortical
networks', *Nature*, 473(7345), pp. 87–91. doi: 10.1038/nature09880.

De La Rocha, J. *et al.* (2007) 'Correlation between neural spike trains increases with firing rate',
Nature. doi: 10.1038/nature06028.

Levy, D. R. *et al.* (2019) 'Dynamics of social representation in the mouse prefrontal cortex.',
Nature neuroscience. doi: 10.1038/s41593-019-0531-z.

Liang, B. *et al.* (2018) 'Distinct and Dynamic ON and OFF Neural Ensembles in the Prefrontal
Cortex Code Social Exploration', *Neuron*. Cell Press, 100(3), pp. 700-714.e9. doi:
10.1016/J.NEURON.2018.08.043.

Litwin-Kumar, A. and Doiron, B. (2012) 'Slow dynamics and high variability in balanced
cortical networks with clustered connections.', *Nature neuroscience*. NIH Public Access, 15(11),
pp. 1498–505. doi: 10.1038/nn.3220.

Luongo, F. J. *et al.* (2016) 'Correlations between prefrontal neurons form a small-world network
that optimizes the generation of multineuron sequences of activity', *Journal of Neurophysiology*,
115(5), pp. 2359–2375. doi: 10.1152/jn.01043.2015.

Luongo, F. J., Horn, M. E. and Sohal, V. S. (2016) 'Putative microcircuit-level substrates for
attention are disrupted in mouse models of autism', *Biological Psychiatry*. doi:
10.1016/j.biopsych.2015.04.014.

Mukamel, E. A., Nimmerjahn, A. and Schnitzer, M. J. (2009) 'Automated analysis of cellular
signals from large-scale calcium imaging data.', *Neuron*, 63(6), pp. 747–60. doi:
10.1016/j.neuron.2009.08.009.

Murugan, M. *et al.* (2017) 'Combined Social and Spatial Coding in a Descending Projection
from the Prefrontal Cortex', *Cell*, 171(7), pp. 1663-1677.e16. doi: 10.1016/j.cell.2017.11.002.

Nelson, S. B. and Valakh, V. (2015) 'Excitatory/Inhibitory Balance and Circuit Homeostasis in
Autism Spectrum Disorders', *Neuron*, 87(4), pp. 684–698. doi: 10.1016/j.neuron.2015.07.033.

Peça, J. *et al.* (2011) 'Shank3 mutant mice display autistic-like behaviours and striatal
dysfunction HHS Public Access', *Nature*, 472(7344), pp. 437–442. doi: 10.1038/nature09965.

Pinto, L. *et al.* (2013) 'Fast modulation of visual perception by basal forebrain cholinergic

neurons', 16(12). doi: 10.1038/nn.3552.

Rubenstein, J. L. R. and Merzenich, M. M. (2003) 'Model of autism: increased ratio of
excitation/inhibition in key neural systems', *Genes, Brain and Behavior*. doi: 10.1034/j.1601-
183X.2003.00037.x.

Sakurai, Y. (1999) 'How do cell assemblies encode information in the brain?', *Neuroscience and
Biobehavioral Reviews*. doi: 10.1016/S0149-7634(99)00017-2.

Sakurai, Y. *et al.* (2018) 'Multiple approaches to the investigation of cell assembly in memory
research—present and future', *Frontiers in Systems Neuroscience*. Frontiers Media S.A. doi:
10.3389/fnsys.2018.00021.

Schneidman, E. *et al.* (2006) 'Weak pairwise correlations imply strongly correlated network
states in a neural population', *Nature*. Nature Publishing Group, 440(7087), pp. 1007–1012. doi:
10.1038/nature04701.

Selimbeyoglu, A. *et al.* (2017) 'Modulation of prefrontal cortex excitation/inhibition balance
rescues social behavior in CNTNAP2-deficient mice.', *Science translational medicine*. American
Association for the Advancement of Science, 9(401), p. eaah6733. doi:
10.1126/scitranslmed.aah6733.

Stefanini, F. *et al.* (2020) 'A Distributed Neural Code in the Dentate Gyrus and in CA1', *Neuron*.
Cell Press. doi: 10.1016/j.neuron.2020.05.022.

Vaadia, E. *et al.* (1995) 'Dynamics of neuronal interactions in monkey cortex in relation to
behavioural events', *Nature*. Springer, 373(6514), pp. 515–518. doi: 10.1038/373515a0.

Yizhar, O. *et al.* (2011) 'Neocortical excitation/inhibition balance in information processing and
social dysfunction', *Nature*. doi: 10.1038/nature10360.

838
839
840

Acknowledgments: We thank Josh Berke and Ruchi Malik for helpful comments on the
manuscript. NAF is funded by K08NS105938 from NINDS and 5R25NS070680-05 from
NINDS.

Author contributions: Experiments were performed by NAF. Analysis was designed by NAF
and VS and performed by NAF; AH assisted with collecting data; the manuscript was written by
NAF and VS.

Declaration of interests: Authors declare no competing interests.

Data and materials availability: All code used in the analysis will be deposited on GITHUB
for any researcher for purposes of reproducing or extending the analysis.

Figure 1. Social interaction modulates activity levels within prefrontal ensembles.
A. Mice were imaged across 9 consecutive behavioral epochs (each lasting 5 min) during which
they were either alone in their homecage or interacted with one of two novel sex-matched
juvenile mice introduced to the homecage ('M1' or '(M2'). Each novel mouse was
subsequently re-introduced to the home cage during a familiar epoch. GCaMP traces during
show examples of neurons that appear to increase or decrease activity during social epochs
(see arrows at the right of each trace).

B. Mean z-scored GCaMP traces for all neurons recorded from wild-type mice (663 neurons
from 10 mice) aligned to the onset of social interaction during the first bout of interaction
within each social epoch.
C. Cumulative plot showing the distribution of activity levels for individual neurons during
homecage epochs or periods of social interaction (percent time active in homecage: 1.8% +/-
0.1, percent time active during social interaction: 2.1 +/- 0.1%, $p = 0.00002$, paired t-test; n =
663 neurons from 10 WT mice).
D. Scatter-plot showing the activity of each neuron during each behavioral condition, expressed
as a percentile relative to a null distribution generated by circularly shuffling that neuron's
activity. Activity levels during social interaction or while the mouse was alone in its home
cage are plotted on the x and y axis, respectively. Kernel density plots along the axes indicate
the fraction of neurons whose activity was at a given percentile of the null distribution.
Neurons with activity > 90^{th} percentile of shuffled datasets (green dotted line) were
considered to be positively modulated, whereas neurons with activity < 10^{th} percentile (green
dotted line) were considered to be negatively modulated during each behavior (>90^{th}
percentile, social: 152/663 neurons, home cage: 80/663 neurons; $p < 0.00001$, chi-squared
test; <10^{th} percentile, social: 128/663 neurons, home cage: 119/663 neurons; $p = 0.5$, chi-
squared test).

Figure 2. Classifying behavior from prefrontal ensembles using a simple neural network.

A. We constructed a neural network consisting of a single hidden layer (containing 1000 units) which were connected to a single output unit. The thickness of lines between each hidden layer unit and the output unit reflects the magnitude of the output weight. Positive and negative weights are indicated by solid and dashed lines, respectively. Each hidden layer unit received input from a random subset of prefrontal neurons from one real dataset. For clarity, we have only shown input connections to two hidden layer units (which are differentiated by their blue and red colors) – output weights from other hidden units are shown in black. The output weight from each hidden layer neuron was iteratively updated during training. We trained the classifier to distinguish periods marked as home cage exploration or social interaction by dividing a dataset into 500-frame blocks, and then using alternating blocks for training or testing.

B. The classifier performed poorly (near chance) when the input connection probability (governing the number of prefrontal neurons that provided input to each hidden layer unit) was <10%. Classification accuracy was above chance in 8/10 mice and increased to a peak of 69 +/- 3% in these mice, before decreasing again for connection probabilities >30%. The

894   classifier performed near chance levels when we trained and tested using data that had been
895   randomly swap-shuffled.

897
898
899 **Figure 3. Classifier weights reveals an ensemble that increases correlations during social**
900 **behavior and detects social behavior.**
901 **A.** Example histogram depicting the distribution of output weights assigned to connections
902   between hidden layer units and the output unit over the course of training.

B. Matrix of input connections for hidden units which detect the social (left) or home cage condition (right). The hidden layer units (x-axis) have been arranged in order of increasing output weights to identify 'social units' (25 most negative output weights) and 'home cage units' (25 largest positive output weights). Prefrontal neurons (y-axis) have been arranged in order of their preference for social interaction vs. home cage, i.e. the difference between their activity levels in the two conditions.

C. Correlation matrix showing the input similarity, i.e., the pairwise correlation between binary vectors representing the input connections to each pair of hidden layer units. Hidden layer units are arranged in order of increasing output weight. Red and blue rectangles indicate correlations with social or home cage units, respectively. A gaussian filter with a standard deviation of 3 was applied to the 1000x1000 matrix to improve visualization.

D. For each hidden layer unit, we plotted its average input similarity to either the 25 social units (red) or the 25 home cage units (blue). Hidden layer units (x-axis) are again arranged by output weight. Social units had similar patterns of input compared to each other but not to home cage units and vice-versa.

E. The average input similarity of each hidden layer unit to 25 hidden layer units with near-zero output weights ('neutral units'; black rectangle in C).

F. We defined social and home-cage (HC) ensembles as the 20% of prefrontal neurons most likely to provide input to the social or home cage units, respectively. The mean activity of both home cage and social ensembles increased during social interaction compared to the home cage condition (social ensemble: mean activity level 1.4 +/- 0.3% in home cage vs. 1.8 +/- 0.3% during social interaction, p < 0.05, sign-rank test; home cage ensemble: mean activity level 1.5 +/- 0.30% in home cage vs. 1.9 +/- 0.3% during interaction, p < 0.001, sign-rank test).

G. Correlations between neurons in the same ensemble increased during social interaction for the social ensemble but for the home cage ensemble (mean correlation coefficient between neurons in the social ensemble: 0.009 +/- 0.002 in home cage vs. 0.012 +/- 0.002 during social interaction, p < 0.05; home cage ensemble mean correlation coefficient 0.011 +/- 0.02 in home cage vs. 0.005+/- 0.003 during social interaction, p=0.99, sign-rank).

932
Figure 4. Correlations transmit additional information that is not efficiently conveyed by
changes in activity levels alone.
A. Cartoon illustrating that information about behavior may be encoded through changes in
activity levels, correlations between neurons, or both. When behavior modulates activity
levels, correlations in two behavioral conditions may differ or be the same, and vice-versa.

B. To disentangle the roles of activity levels and correlations in transmitting information we
used two different methods to create shuffled (surrogate) datasets which preserve changes in
activity levels, but either do or do not preserve patterns of correlations. We made random,
reciprocal swaps of activity between neurons to generate surrogate datasets which maintained
network activity in each frame as well as the number of blocks of activity for each neuron.
However, these datasets destroyed the correlation structure. In a second set of surrogate
datasets we used SHARC to iteratively generate surrogates in which the correlation structure
was also maintained.
C. To maintain dynamic changes in activity levels and correlations that are associated with the
two behavioral conditions we swap-shuffled or performed SHARC separately for each
behavioral epoch, then concatenated the 9 resulting surrogate subrasters to create each
surrogate dataset.
D. We trained a classifier (with a connection probability = 0.3) on each real dataset, then tested
that classifier on swap or SHARC-shuffled surrogate datasets generated from that real
dataset. Accuracy was significantly higher for the SHARC-shuffled surrogates, which
maintain the behaviorally-modulated correlations found in the original dataset (accuracy for
SHARC shuffled surrogate datasets = 68 +/- 4%, classifier accuracy for swap shuffled
surrogate datasets = 61 +/- 4%, $p < 0.05$, sign-rank test, n = 10 mice).

Specifically-enriched combination
observed during social behavior

In the home cage, a partially overlapping combination
is specifically enriched (but neuron C is 'left out')

Other home cage combinations can
contain the 'left-out' neuron (neuron C)

957
Figure 5. Behaviorally-specific patterns of coactivity are formed by neurons that are active
in both conditions, but coactive with different partners in each condition.
Left: We identified combinations of 3 neurons that are specifically enriched during one
behavioral condition (occurring more often during social interaction than in 95% of surrogate
datasets, and occurring less often during home cage exploration than in 50% of surrogate
datasets, or vice-versa). We then identified overlapping combinations occurring during the
opposite behavioral condition in which a single neuron was 'left out.' In other words, we
identified combinations from the two conditions that overlapped in exactly two neurons.
Right, top: Histogram showing the number of distinct 3 neuron home cage combinations that
contain the neuron which participates in a social combination but is 'left out' during home cage
behavior. **Right, bottom:** Histogram showing the number of distinct 3 neuron social
combinations that contain the neuron which participates in a home cage combination but is 'left

970   out' during social interaction. In the vast majority of cases, neurons that are 'left out' in one

971   condition are still active during that condition and participate in other combinations.

973
974

Figure 6. Shank3 KO mice have disorganized ensembles for which correlations fail to enhance the transmission of information about social behavior.

A. The mean time that *Shank3* KO mice or wild-type (WT) littermates spend interacting with a novel juvenile mouse of the same sex during a 5 min assay. Data has been pooled from 8 unimplanted WT mice as well as the 5 implanted WT mice used for microendoscopic imaging, and 5 unimplanted KO mice in addition to the 4 implanted mice used for imaging. For implanted mice we used the average of interaction time for the 2 novel mice. Pooled data showed decreased interaction in KO mice (173 +/- 12 s vs. 120 +/- 18 s for WT and KO respectively, p < 0.05, t-test). The un-implanted cohort alone shows a similar significant decrease in interaction time for KO mice (165 +/- 15 s vs 110 +/- 16 s for WT and KO respectively, p < 0.05, t-test). In the implanted cohort there was a similar trend toward

decreased interaction for KO mice (186 +/- 20 s vs 133 +/- 37 s, for WT and KO
respectively, p = 0.21, t-test).

B. (Similar to Fig. 1D). Scatter-plot showing the activity of each neuron during each behavioral
condition, expressed as a percentile relative to a null distribution generated by circularly
shuffling that neuron's activity. Activity levels during social interaction or while the mouse
was alone in its home cage are plotted on the x and y axis, respectively. Kernel density plots
along the axes indicate the fraction of neurons whose activity was at a given percentile of the
null distribution. Neurons with activity > 90th percentile of shuffled datasets (green dotted
line) were considered to be positively modulated, whereas neurons with activity < 10th
percentile (green dotted line) were considered to be negatively modulated during each
behavior. Data is plotted for Shank3 KO mice (red) and WT littermates (blue) (mean
percentile of activity during social interaction, WT: 50 +/- 2 percentile; KO: 64 +/- 2
percentile; p < 0.0001 by 2-sample *t*-test; mean percentile of activity during home cage, WT:
47 +/- 2 percentile; KO: 51 +/- 2 percentile; p = 0.1, *t*-test).

C. Bar graph showing the fraction of neurons whose activity was positively or negatively
modulated (>90th percentile or <10th percentile) during social interaction. The proportion of
neurons which increased activity during social interaction was significantly greater in KO
mice (22% in WT vs. 39% in KO, chi-squared = 17.7, p < 0.0001), whereas the
downregulated ensemble was significantly smaller in KO mice (25% in WT vs. 15% in KO,
chi-squared test, 8.2, p < 0.005). Error bars denote the binomial S.E.M. algebraically derived
from total number of neurons and the proportion that were modulated in the specified
direction.

D. The proportion of 3 neuron combinations occurring during home cage exploration that are
enriched > the 99.9th percentile compared to swap-shuffled datasets was similar across WT
(Blue; 7.6%) and KO (Red; 7.8%) mice. By contrast, the proportion of 3 neuron
combinations occurring during social interaction that are enriched > 99.9th percentile
compared to swap-shuffled datasets was 7.5% in WT compared to only 3.4% in KO mice
(total number of home cage combinations: 4,187 in 5 WT mice, and 5,878 in 4 KO mice;
total number of social combinations: 5,487 in 5 WT mice, 16,326 in 4 KO mice). The top
two plots show histograms of enrichment for the home cage (upper) or social conditions
(middle); the lower panel is a bar graph showing the fraction of these combinations that were
specifically enriched above the 99.9th percentile (chi-squared = 165, p < 0.0001). Error bars
denote the S.E.M. algebraically derived from the binomial distribution, the number of 3
neuron combinations in each condition, and the proportion of those combinations that were
enriched.

E. Performance of classifier trained on real datasets and tested on surrogate datasets.
Performance was not better (and was non-significantly worse) when correlation structure was
maintained using SHARC (classifier accuracy: 62 +/ 4% for SHARC vs 63 +/- 2% for swap
shuffled surrogate datasets, p = 0.47, sign-rank test).

1031
Supplemental Figure S1. Spatial Decorrelation of Neuronal Signals.

A. Example image (top) and individual neuron GCaMP traces (bottom) from prefrontal cortex imaged with implanted endoscope.

B. The average GCaMP signal from a region of interest (ROI), corresponding to one neuron, was corrected by subtracting the average GCaMP signal from the surrounding pixels, in order to spatially deconvolve signals from each ROI vs. the surrounding neuropil. Examples traces from a single neuron are shown.

C. The pairwise correlation matrix between signals from different neurons is shown (calculated from 550 seconds of activity from a single wildtype mouse), for the original GCaMP signals (top left), the surround signals (top right), the surround-subtracted signals (bottom left), and the surround-subtracted signals after lowpass filtering (bottom right).

1043

Mouse	Sex	Neurons	Peak Accuracy
WT			
1	f	84	0.687
2	f	39	0.705
#3	m	25	0.53
4	f	75	0.605
#5	m	37	0.459
6	*m*	*86*	*0.647*
7	*m*	*74*	*0.848*
8	*m*	*73*	*0.691*
9	*m*	*71*	*0.694*
10	*f*	*99*	*0.633*
KO			
1	f	83	0.772
2	m	77	0.832
3	f	45	0.34
4	f	85	0.627

Supplemental Table S1. Details for each mouse included in this study.

The table shows the genotype, sex, number of imaged neurons, and peak classifier accuracy (performance when half the data was used for training and half for testing). Figure 2 showed how classifier accuracy depends on the input connection probability; because their performance was not >50% for multiple input connection probabilities, WT mice 3 & 5 (marked with #) were not included in this illustrative plot. However, we did not exclude data from these mice in any analyses. WT mice 1-5 were wild-type littermates of Shank3 KO mice. WT mice 6-10 (indicated by italics) were not littermates of Shank3 KO mice and were therefore not included as WT controls for the analyses shown in Figure 6.