Evaluation of Synergy Extrapolation for Predicting Unmeasured Muscle Excitations from Measured Muscle Synergies

Di Ao[1], Mohammad S. Shourijeh[1], Carolynn Patten [2], Benjamin J. Fregly[1*]

[1]Rice Computational Neuromechanics Lab, Department for Mechanical Engineering, Rice University, Houston, TX, USA

[2] Biomechanics, Rehabilitation, and Integrative Neuroscience (BRaIN) Lab, Department of Physical Medicine and Rehabilitation, University of California, Davis, CA, USA

* Correspondence:
Benjamin J. Fregly
fregly@rice.edu

Keywords: muscle synergy, EMG-driven modeling, stroke, principal component analysis, non-negative matrix factorization, muscle excitation, EMG normalization

Abstract

Electromyography (EMG)-driven musculoskeletal modeling relies on high-quality measurements of muscle electrical activity to estimate muscle forces. However, a critical challenge for practical deployment of this approach is missing EMG data from muscles that contribute substantially to joint moments. This situation may arise due to either the inability to measure deep muscles with surface electrodes or the lack of a sufficient number of EMG electrodes. Muscle synergy analysis is a dimensionality-reduction approach to decompose a large number of muscle excitations into a small number of time-varying synergy excitations along with time-invariant synergy weights that define the contribution of each corresponding synergy excitation to a specific muscle excitation. This study evaluates how accurately missing muscle excitations can be predicted using synergy excitations extracted from muscles with available EMGs (henceforth called "synergy extrapolation"). The results were reported on a gait dataset collected from a stroke survivor walking on an instrumented treadmill at self-selected and fastest-comfortable speeds. The evaluation process started with full calibration of a lower-body EMG-driven model using 16-channel EMGs (including surface and indwelling) in each leg. One indwelling EMG (either iliopsoas or adductor longus) was then treated as unmeasured at a time. The synergy weights associated with the unmeasured muscle were predicted through solving a nonlinear optimization problem where the errors between inverse dynamics and EMG-driven joint moments were minimized. We also quantitatively evaluated how synergy analysis algorithms (principal component analysis (PCA) and non-negative matrix factorization (NMF)), EMG normalization methods, and number of synergies affect the accuracy of the predicted unmeasured muscle excitation. Synergy extrapolation performance was most influenced by the choice of synergy analysis algorithm and number of synergies. PCA with 5 or 6 synergies consistently predicted unmeasured muscle excitations most accurately and with greatest robustness to choice of EMG normalization method. Furthermore, the associated joint moment matching accuracy was comparable to that produced by the full EMG-driven calibration. The synergy extrapolation method described in this study may facilitate the assessment of human neuromuscular control and biomechanics in response to surgical or rehabilitation treatment when important EMG signals are missing.

1 Introduction

Prediction of muscle forces can provide valuable insight not only for understanding the control strategies employed by the central nervous system (CNS) (Contessa and Luca, 2013; Del Vecchio et al., 2018), but also for developing effective treatments for neuromusculoskeletal disorders (Shao et al., 2009; Fregly et al., 2012b, 2012a; Allen et al., 2013; Pitto et al., 2019; Sauder et al., 2019) Since direct measurement of muscle force is invasive using current technology, computational techniques has been developed to generate muscle force estimates (Anderson and Pandy, 2001; Lloyd and Besier, 2003; Thelen et al., 2003; Buchanan et al., 2005; Shao et al., 2009) However, since the human musculoskeletal system possesses more muscles than degrees-of-freedom (DOFs) in the skeleton (i.e., the muscle redundancy problem), no unique solution for muscle exists unless muscle activity patterns are defined by measured EMG signals (Lloyd and Besier, 2003; Manal and Buchanan, 2003; Buchanan et al., 2005; Shao et al., 2009; Kumar et al., 2012; Sartori et al., 2012; Meyer et al., 2017), or assumptions are made about how muscles contribute to the joint moments, such as minimization of energetic cost (Anderson and Pandy, 2001; Ackermann and van den Bogert, 2010; S. Shourijeh and McPhee, 2014). EMG-driven musculoskeletal modeling is a computational approach for predicting muscle forces that can bypass the muscle redundancy problem while simultaneously allowing for calibration of unmeasurable musculotendon properties (e.g., optimal muscle fiber length) (Lloyd and Besier, 2003; Amarantini and Martin, 2004; Shao et al., 2009; Sartori et al., 2012; Meyer et al., 2017). In EMG-driven models, processed EMG and muscle-tendon kinematic data are input to a muscle force generation model (typically a Hill-type model) to predict muscle forces and corresponding net muscle moments. Nonlinear optimization is then used to calibrate musculotendon model parameters such that predicted net muscle moments match inverse dynamic joint moments as closely as possible.

In EMG-driven models, the quality of the measured EMG signals affects the reliability of the estimated muscle forces. Surface EMG recording, which is non-invasive and easily applicable, has been the most popular method for measuring muscle electrical activity for biomechanical studies. There are, nevertheless, intrinsic potential challenges with surface EMG data that may limit the accuracy of estimated muscle forces, such as noisy signals from crosstalk between adjacent muscles, movement artifacts, and challenges in attaining the true maximum muscle excitation for EMG normalization (Farina et al., 2002; Racinais et al., 2013; Sartori et al., 2014). Beyond these issues, the inability to acquire EMG data from deep muscles that contribute substantially to joint moments is a practical challenge (Sartori et al., 2014; Zonnino and Sergi, 2019). For instance, it is practically impossible to collect EMG data from deep hip muscles (e.g. iliacus and psoas) using surface electrodes. However, when EMG data from important deep muscles are missing in an EMG-driven model, force estimates for other muscles that have similar roles may be significantly overestimated (Zonnino and Sergi, 2019). Compared to surface electrodes, indwelling electrodes are able to measure the electrical activity of deep muscles with lower levels of crosstalk (Péter et al., 2019). However, the use of indwelling electrodes requires special skills and longer set-up time and may cause discomfort and pain for the subject. Furthermore, in some cases, such as patients who have a cancerous tumor near an important deep muscle, use an indwelling electrode for EMG data collection may be contraindicated for safety reasons. Regardless of the EMG measurement technique, EMG-driven models require EMG data collection from a large number of muscles, which may not be possible due to hardware limitation. Therefore, a computational method is needed that can provide reliably estimate muscle excitations that cannot be measured experimentally for technical or safety reasons.

Previous studies have explored computational methods for predicting unmeasured muscle excitations within the context of EMG-driven modeling. Static optimization (SO) (Crowninshield and Brand, 1981; Anderson and Pandy, 2001; Damsgaard et al., 2006; Heintz and Gutierrez-Farewik, 2007) has

also been embedded into calibration of EMG-driven models, in which while the musculoskeletal model
is being calibrated, missing muscle excitations can be estimated. The objective function for such
technique includes minimizing both activation levels of unmeasured muscles and joint moment
tracking errors (Sartori et al., 2014; Zonnino and Sergi, 2019). Zonnino et al. presented an EMG-driven
forward dynamics estimator that used an SO-based neural model to determine unmeasured muscle
activations, which reduced the estimation error of muscle forces in comparison to the performance of
a conventional estimator that neglected the contribution of unmeasured muscles (Zonnino and Sergi,
2019). Similarly, Satori et al. developed a hybrid EMG-informed model in which experimental EMG
signals were minimally adjusted while missing EMG signals (e.g., from iliacus and psoas) were
predicted via SO. However, none of these studies have shown evidence that the reproduced unmeasured
muscle activations are reliable and in reasonable agreement with experimental measurements.
Furthermore, because time histories are not taken into account in SO, the resulting muscle activations
might contain abrupt unrealistic changes due to the optimization problem being solved at an instant in
time.

Another approach for estimating unmeasured muscle excitations is to use muscle synergy concepts. A
muscle synergy is composed of a time-varying synergy excitation and a corresponding time-invariant
synergy vector containing weights that define how each synergy excitation contributes to the excitation
of each muscle (Tresch et al., 1999; Ting and Chvatal, 2010; Banks et al., 2017; Shourijeh and Fregly,
2020) Muscle synergies have been broadly used in descriptive research to analyze experimental muscle
excitations during a large number of movement tasks (Ivanenko et al., 2005; Torres-Oviedo and Ting,
2007; Bowden et al., 2010; Walter et al., 2014; Kristiansen et al., 2015; Meyer et al., 2016a; Ruiz
Garate et al., 2017; Sauder et al., 2019), but few studies have performed predictive analyses using
muscle synergy information (Ajiboye and Weir, 2009; Bianco et al., 2018). Ajiboye and Weir
demonstrated that subject-specific synergies extracted from muscle activities recorded for a subset of
postures can be used to predict EMG patterns for the remaining postures. Bianco et al. investigated the
theoretical feasibility of using synergy excitations extracted from a group of 8 'included' muscle
excitations treated as measured as building blocks to construct muscle excitations for a group of 8
'excluded' muscle excitations treated as unmeasured. However, the muscle synergy weights associated
with 'remaining' postures in (Ajiboye and Weir, 2009) or 'excluded' muscles in (Bianco et al., 2018)
were not predicted without knowledge of the muscle excitations treated as unmeasured but rather fitted
with knowledge of those excitations using least square algorithms. Several other studies have imposed
synergy structures on muscle excitations or activations through optimization when estimating knee
contact force (Walter et al., 2014), joint stiffness (Shourijeh and Fregly, 2020), or motion (Clark et al.,
2009; Allen and Neptune, 2012; Meyer et al., 2016b; Mehrabi et al., 2019; Falisse et al., 2020).
However, to the best of the authors' knowledge, the reliability with which unmeasured muscle
excitations can be estimated using EMG-driven models with unknown synergy weights has not been
studied previously. Additionally, synergy extrapolation has never been evaluated from a predictive
perspective, and it is therefore unknown how methodological choices for how synergy analysis is
performed affect the outcome of synergy extrapolation.

A synergy-based muscle excitation estimation method involving EMG-driven modeling, termed
"synergy extrapolation," was developed to predict muscle excitations that cannot be measured
experimentally. This study evaluated how reliably synergy extrapolation is able to predict missing
indwelling EMG signals using a well-calibrated EMG-driven lower-body musculoskeletal model.
Since the outcome of muscle synergy analysis is affected by methodological choices, such as EMG
processing (e.g. magnitude normalization), physiological assumptions (e.g. number of synergies), and
numerical analysis approaches (e.g. matrix decomposition algorithm (Tresch et al., 2006; Hug et al.,
2012; Steele et al., 2013; Oliveira et al., 2014; Banks et al., 2017; Shuman et al., 2017; Ebied et al.,

2018; Gallina et al., 2018)), we also evaluated how each of these choices affects synergy extrapolation
results. The evaluated methodological decisions for MSA process included matrix factorization
algorithm (PCA and NMF), EMG normalization approach, and number of synergies. To validate this
approach, we analyzed a gait data set collected from a high-functioning stroke subject performing
treadmill walking at self-selected and fastest-comfortable speeds. Muscle excitations that were
measured using indwelling EMG electrodes, i.e., from iliopsoas and adductor longus, were assumed
unknown and were predicted. By quantitatively evaluating the differences in synergy extrapolation
performance produced by different methodological decisions, this work provides evidence-based
suggestions for which methods are likely to produce the most accurate predictions of missing muscle
excitations.

2   Materials and Methods

2.1   Experimental data

A previously published gait dataset collected from one high-functioning stroke survivor (age 79 years,
LE Fugl-Meyer Motor Assessment 32/34 pts, right-sided hemiparesis, height 1.7 m, mass 80.5 kg) was
used to evaluate the synergy extrapolation process (Meyer et al., 2017). Motion capture (100 Hz, Vicon
Corp., Oxford, UK), ground reaction force (1000 Hz, Bertec Corp., Columbus, OH), and surface EMG
(1000 Hz, Motion Lab Systems, Baton Rouge, LA) data were recorded simultaneously while the
subject walked on a split-belt instrumented treadmill (Bertec Corp., Columbus, OH) at two speeds: 0.5
m/s (self-selected speed) and 0.8 m/s (fastest comfortable speed). Ten cycles of gait data for each
walking speed were analyzed. Motion capture and ground reaction force (GRF) data were low-pass
filtered using a fourth-order zero-phase lag Butterworth filter with a cut-off frequency at $7/t_f$ Hz, where
t_f is the period of the gait cycle being processed (McLean et al., 2005; Meyer et al., 2017). Sixteen
channels of EMG data were collected from each leg using surface and a small number of indwelling
electrodes (i.e. fine-wire), which made available EMG data from important deep muscle groups (e.g.
iliopsoas) (supplementary table 1). After being high-pass filtered at 40 Hz, demeaned, full-wave
rectified, and low-pass filtered at $3.5/t_f$ Hz, each EMG signal was normalized to the maximum value
over all trials. After processing, each trial of EMG data was time-normalized to 101 time frames in
each gait cycle (heel strike to heel strike). See (Meyer et al., 2017) for more details.

2.2   Musculoskeletal model

A generic full-body OpenSim musculoskeletal model (Arnold et al., 2010) was adopted for analyses
in OpenSim v3.3 (Delp et al., 2007; Seth et al., 2018). The model controlled 5 degrees of freedom
(DOFs), including two-DOF hip joints (flexion/extension and adduction/abduction), one DOF knee
joints (flexion/extension), and two-DOF ankle joints (plantarflexion/dorsiflexion and
inversion/eversion). Of the 45 muscles in each leg accounted for in the original model, 35 muscles per
leg were retained. Compartments of large muscles shared common EMG signals (Meyer et al., 2017).
To personalize the model, we performed five steps sequentially on the modified generic OpenSim
model using a combination of OpenSim and custom Matlab analyses: 1) model scaling to match the
subject's anthropometry; 2) kinematic calibration to determine personalized lower body joint positions
and orientations (Reinbolt et al., 2005); 3) inverse kinematics (IK) to calculate joint angle time histories
from the surface marker data using the calibrated kinematic model; 4) surrogate musculoskeletal
geometry creation to fit muscle-tendon lengths and moment arms as polynomial functions of lower
body joint angles and velocities (Menegaldo et al., 2004; Meyer et al., 2017); and 5) inverse dynamics
(ID) to obtain experimental joint moments using experimental GRF data and IK-derived time histories
of joint kinematics as inputs.

2.3 Synergy extrapolation methodology

Our proposed synergy extrapolation method consists of two functional blocks (Fig. 1). First, the
synergy extrapolation block seeks to reconstruct unmeasured muscle excitations with measured
synergy excitations and a set of unmeasured synergy vector weights that is found through an
optimization process. Muscle synergy analysis is performed trial by trial for all measured muscle
excitations e_m to extract a low-dimensional set of time-varying measured synergy excitations W_m and
corresponding sets of time-invariant measured weighting vectors H_m, where each weighting vector
represents the contribution of the corresponding synergy excitation to all measured muscle excitations.
Next, unmeasured muscle excitations e_x are constructed using the measured synergy excitations
W_m along with a set of trial-specific synergy vector weights H_x associated with the unmeasured
muscles and predicted by the optimization. In the present study, several methodological assumptions
and decisions were made prior to MSA, including number of synergies, synergy analysis approach,
and EMG normalization method. The key differences in the implementation of these different
methodologies for synergy extrapolation are described below. Second, the EMG-driven
musculoskeletal model block predicts the net joint moments generated by muscles using a combination
of experimentally measured muscle excitations (e_m) and computationally predicted muscle excitations
(e_x) constructed from measured synergy excitations W_m and predicted synergy vector weights H_x. In
this block, a previously developed and calibrated EMG-driven model, where muscles were treated as
Hill-type models (Hill, 1938; Zajac, 1989) with a rigid tendon, was employed to predict muscle forces
and net joint moments in the lower extremities (Meyer et al., 2017). The joint moment produced by a
muscle spanning a particular joint can be represented as:

$$M = r \cdot F_o^M \cdot \left[a \cdot f_l\big(\tilde{l}^M(t)\big) \cdot f_v\big(\tilde{v}^M(t)\big) + f_p\big(\tilde{l}^M(t)\big) \right] \cos \alpha \qquad (1)$$

where M is the moment generated by the muscle about the joint, r is the moment arm of the muscle
about the same joint, F_o^M is the maximum isometric force of the muscle, a is the muscle activation,
$\tilde{l}^M(t)$ and $\tilde{v}^M(t)$ are the time-varying normalized muscle fiber length and velocity, respectively, and
α is the pennation angle of the muscle. $f_l\big(\tilde{l}^M(t)\big)$ and $f_v\big(\tilde{v}^M(t)\big)$ define the normalized muscle active
force-length and active force-velocity relationships, while $f_p\big(\tilde{l}^M(t)\big)$ defines the normalized muscle
passive force-length relationship (Zajac, 1989)(Meyer et al., 2017). Muscle activation (a) was
calculated from muscle excitation (e) using a model of muscle activation dynamics (He et al., 1991;
Lloyd and Besier, 2003; Meyer et al., 2017). Muscle excitations were derived by multiplying the
processed EMG signals by muscle-specific scale factors between 0.05 and 1 to reflect unknown
maximum excitation levels.

In this study, unmeasured muscle excitations were predicted by following a two-step process. The first
step calibrated an EMG-driven model of each leg using a full set of 16 EMG channels per leg
(henceforth called 'full EMG-driven'), where every muscle included in the EMG-driven model was
associated to an experimentally measured EMG signal. During this step, a sequence of optimizations
was performed to identify the parameter values required by the activation dynamics models, Hill-type
muscle-tendon models, and surrogate musculoskeletal geometric model (described in Meyer et al.,
2017) that reproduced the lower body inverse dynamic joint moments as closely as possible (Eq. 2).
The model parameter values calibrated for each muscle-tendon actuator by the optimization process
included: electromechanical delay, activation time constant, activation nonlinearity constant, EMG
scale factor, optimal muscle fiber length, tendon slack length, and geometric coefficients defining
muscle-tendon lengths, velocities, and moment arms. The second step calibrated the synergy vector
weights, H_x for the unmeasured muscle excitations using the EMG-driven model parameters found in

the first step. During this step, only unmeasured synergy vector weights H_x were identified iteratively
through optimization. Thus, this two-step process estimated missing muscle excitations assuming a
well-calibrated EMG-driven model was already available, including calibrated parameter values for
the muscles with missing excitations. Predicting missing muscle excitations reliably using a well-
calibrated EMG-driven model is an important step toward predicting missing muscle excitations while
simultaneously calibrating the parameter values of the EMG-driven model. The structure of
unmeasured synergy variables was slightly different between the matrix factorization algorithms, and
more details are provided in the section of methodological choices on synergy extrapolation.

The primary cost function for both calibration steps was formulated as:

230
$$J \triangleq \sum_{i=1}^{N} (M_i^{mod} - M_i^{exp})^2 \tag{2}$$

where M_i^{mod} is the model-predicted moment about joint i, M_i^{exp} is the experimental moment about
joint i calculated using inverse dynamics, and N is the total number of joints. Further details on the
specification of initial guesses, variable bounds, overall cost function structure, additional constraints,
and penalty terms for calibrating the EMG-driven model can be found in (Meyer et al., 2017). For the
second step, the missing synergy vector weights, H_x, were unbounded and were initialized by the
optimization using randomly chosen values between 0 and 1. However, the unmeasured muscle
excitations predicted by synergy extrapolation were constrained to be between 0 and 1. All calibrations
were performed using Matlab's built-in "fmincon" optimization algorithm using sequential quadratic
programming.

2.4   Methodological choices for synergy extrapolation

Muscle synergy analysis requires a number of methodological choices that can influence the results of
the analysis (Banks et al., 2017). Methodological choices that have been studied include EMG
processing approaches (e.g. filtering parameters, normalization methods), assumptions about neural
control complexity (e.g. number of synergies, number and choice of muscles, synergy vector variability
across trials), numerical analysis approaches (e.g., matrix decomposition algorithm, selection criteria
for number of synergies), and post-processing of results (Tresch et al., 2006; Hug et al., 2012; Steele
et al., 2013; Oliveira et al., 2014; Shourijeh et al., 2016; Banks et al., 2017; Shuman et al., 2017; Ebied
et al., 2018; Gallina et al., 2018; Mehryar et al., 2020). In the present study, for each subset of measured
muscles, we performed synergy extrapolation using a total of 80 methodological combinations
comprised of 2 algorithms for matrix factorization, 5 methods for EMG normalization, and 8 choices
for number of muscle synergies.

The two matrix factorization algorithms that render the most divergent results for MSA were used to
calculate measured muscle synergies: non-negative matrix factorization (NMF) and principal
component analysis (PCA) (Olree and Vaughan, 1995; Lee and Seung, 1999; d'Avella et al., 2003;
Tresch et al., 2006; Ting and Chvatal, 2010; Banks et al., 2017; Bianco et al., 2018; Ebied et al., 2018).
NMF is a nonlinear decomposition method, the solution of which is found iteratively to minimize the
error between the reconstructed and the original dataset subject to constraints on non-negativity.
Typically, NMF derives components that are similar across multiple decompositions but not
numerically unique due to the non-convexity of its search space (Shourijeh et al., 2016). In contrast,
PCA is a linear analytical operation that seeks to identify the internal structure of the data in a way that
best explains its variance. PCA creates a unique solution of orthogonal principal components composed

of both positive and negative values (Torres-Oviedo and Ting, 2007; Ting and Chvatal, 2010). During
synergy extrapolation with a given number of synergies, muscle excitations in the measured muscle
subset e_m were represented as:

$$e_m = \begin{cases} W_m H_m + \varepsilon_m & (NMF) \\ W_m H_m + \mu_m + \varepsilon_m & (PCA) \end{cases} \tag{3}$$

where e_m denotes an n time points × 16 measured EMG signals matrix that contains measured muscle
excitations in columns, W_m denotes an n time points × p synergies matrix that contains measured
synergy excitations in columns, and H_m denotes a p synergies × 16 measured EMG signals matrix
that contains measured synergy vector weights. In preparation for NMF or PCA, measured EMG
signals from each gait cycle were re-sampled to 101 time frames plus $10/t_f$ time frames before the start
of the cycle to account for a maximum electromechanical delay of 100 ms. For NMF and PCA, ε_m
denotes the part of e_m that cannot be explained by $W_m H_m$, while for PCA, μ_m specifies the average
muscle excitations in e_m. Matlab functions "nnmf" (alternating least squares algorithm with 10
replicates) and "pca" were used to perform NMF and PCA, respectively.

Following MSA by either approach, unmeasured muscle excitations e_x were constructed from the
measured synergy excitations W_m using the relationships below:

$$e_x = \begin{cases} W_m H_x & (NMF) \\ W_m H_x + \mu_x & (PCA) \end{cases} \tag{4}$$

where H_x is a p-synergy × 1 vector representing the unmeasured muscle synergy vector weights.
Unlike NMF, PCA needs an additional design variable μ_x that represents the average value of each
unmeasured muscle excitation. Both H_x and μ_x were calibrated by tracking experimental joint
moments within our EMG-driven modeling framework (Eq. 2).

Because EMG normalization method affects the results of MSA, this study explored five approaches
for normalizing the magnitudes of processed EMG signals in preparation for MSA. EMG normalization
was performed either within individual trials (Per trial) or across all trials (Over all trials). Specifically,
each measured muscle EMG signal was normalized using: 1) maximum value over all trials
(MaxOver), 2) maximum value per trial (MaxPer), 3) unit variance over all trials (VarOver), 4) unit
variance per trial (VarPer), and 5) unit magnitude per trial (MagPer) (Banks et al., 2017). VarOver and
VarPer normalizations involved dividing each EMG signal by its standard deviation over all trials and
in each trial, respectively. MagPer normalization involved dividing each EMG signal by its 2-norm
value for each trial (Banks et al., 2017).

Since the specified number of muscle synergies also affects the outcome of MSA substantially, we
repeated the synergy extrapolation process for three through ten synergies. This range was chosen since
three to six muscle synergies have been shown sufficient to account for over 90% of the variability in
up to thirty muscle excitations during human movement (Ivanenko et al., 2005; Ting and Macpherson,
2005; Cappellini et al., 2006; Bianco et al., 2018). Performance of both synergy extrapolation and joint
moment tracking was taken into account when determining the optimal number for synergies.

2.5   Evaluation of synergy extrapolation performance

To implement the first step of the synergy extrapolation procedure (full EMG-driven calibration), we
randomly selected 10 gait cycles from the two walking speeds (five trials per speed) (henceforth called
"calibration trials") to calibrate the subject-specific EMG-driven musculoskeletal model. In

preparation for synergy extrapolation, we assumed one hip muscle EMG signal at a time (either
iliopsoas or adductor longus) recorded using indwelling electrodes to be unmeasured while the
remaining 15 channels of EMG data were treated as measured. Using the calibrated full EMG-driven
model, we then ran synergy extrapolation on the 10 trials used for calibration as well as on another 10
trials (5 trials per speed) randomly selected from the remaining gait cycles (henceforth called
"evaluation trials").

Several common metrics were employed to score outcomes of muscle synergy analysis, performance
of synergy extrapolation, and accuracy of joint moment estimates with different combinations of
methodological choices. Variance accounted for (VAF) was calculated to compare the ability of
methodological combinations to construct measured muscle excitations (Shourijeh et al., 2016). Root
mean square error (RMSE) and Pearson correlation coefficient r between mean experimental and
model-predicted muscle excitations across all trials were computed to assess quantitatively matching
of magnitude and shape, respectively. In addition, RMSE and r values were calculated for each trial,
and the frequency of number of synergies with which highest r values or lowest RMSE values appeared
was analyzed. The correlation between predicted and experimental unmeasured muscle excitations was
interpreted quantitatively as weak ($r < 0.35$), moderate ($0.35 < r \leq 0.67$), strong ($0.67 < r \leq 0.9$), and very
strong ($r \geq 0.9$) (Taylor, 1990). Mean absolute errors (MAE) between experimental and model-
predicted joint moments were calculated across all gait cycles to evaluate the accuracy of the predicted
joint moments during full EMG-driven calibration (step 1) and synergy extrapolation (step 2).

2.6   Statistical analyses

Multiple statistical analyses were used to assess whether the calculated metrics resulting from different
synergy extrapolation methodological choices were statistically different for each unmeasured muscle-
leg combination. First, to assess whether reconstruction performance of measured muscle excitations
was statistically different between the two matrix factorization algorithms and the five EMG
normalization methods, we performed a two-factor ANOVA with a Tukey-Kramer post-hoc analysis
on VAF values. Second, to compare synergy extrapolation performance for different methodological
choices, we conducted two three-factor (matrix factorization algorithm by EMG normalization method
by number of synergies) ANOVA tests on r and RMSE values between predicted and experimental
unmeasured muscle excitations across all calibration and evaluation trials, respectively. In addition, we
also performed paired t-tests on r and RMSE values to investigate whether matrix factorization
algorithm (i.e., PCA and NMF) had a significant influence on synergy extrapolation performance for
the same number of synergies. Third, a three-factor (matrix factorization algorithm by EMG
normalization method by number of synergies) ANOVA was carried out to compare MAE values
characterizing the accuracy of joint moment tracking from different approaches. All statistical analyses
were performed in Matlab, and significance levels were set at $p \leq 0.05$.

3   Results

3.1   Muscle synergy analysis

The two-way ANOVA for mean VAF values revealed main effects of matrix factorization algorithm
(p < 0.01) and EMG normalization method ($p < 0.01$) on the variance explained by factorization of
measured muscle excitations. For five or fewer synergies, PCA generally had significantly higher VAF
values than did NMF for the same number of synergies (all p values < 0.05, grey shading in Table 1).
Overall, extracted synergy excitations were able to predict the measured muscle excitations with > 90%
VAF using 3 or more synergies in the left leg and 4 or more synergies in the right leg for PCA and 4
or more synergies in both legs for NMF (Table 1). Across all EMG normalization methods, MaxOver

produced significantly higher VAF values than did MaxPer ($p < 0.01$), VarPer ($p = 0.028$), and MagPer
($p < 0.01$), while MaxPer produced the lowest VAF amongst the six EMG normalization methods ($p <$
0.01).

3.2   Synergy extrapolation performance

In the results for both calibration (Fig. 2) and evaluation (Fig. 3) trials, mean predicted unmeasured
muscle excitations using PCA were strongly correlated to the experimental muscle excitations (mean
r always ≥ 0.7), which was not consistently observed for the NMF results. Furthermore, RMSE values
between the average predicted and actual unmeasured muscle excitations across all trials using PCA-
based synergy extrapolation were generally lower than those using NMF-based synergy extrapolation.
In addition, synergy extrapolation performed using either PCA or NMF predicted more accurate
unmeasured muscle excitations with less trial-to-trial variability for the left leg than for the right leg
(Figs. 2 and 3).

The three-factor ANOVA analyses indicated that the number of synergies ($p < 0.01$) and matrix
factorization algorithm ($p < 0.01$) had a significant impact on both r and RMSE values for all
unmeasured muscle-leg combinations, while EMG normalization method did not. For each
unmeasured muscle-leg combination, as the number of synergies increased. PCA produced non-
monotonic changes in r and RMSE values, with r values reaching a maximum and RMSE values
reaching a minimum at 5 or 6 synergies. Unlike PCA, r values for NMF initially rose with an increasing
number of synergies and then remained high with further increases, while RMSE values initially
dropped and then leveled off (Fig. 4). Moreover, for the same number of synergies, PCA generally
exhibited less variance than NMF in mean r and RMSE values across the six EMG normalization
methods (Fig. 4).

PCA offered significantly higher r values and lower RMSE values than NMF when the number of
synergies varied from 3 to 6. The one exception was adductor longus in the left leg, where the only
significant difference between PCA and NMF occurred for 3 synergies ($p < 0.01$). When the number
of synergies increased above 6, NMF provided substantially higher r values for adductor longus in the
left leg (Fig. 4 (A)), whereas PCA had markedly higher r values for adductor longus in the right leg (p
< 0.05, Fig. 4 (B)). Additionally, for the number of synergies above 6, RMSE values using NMF were
significantly smaller than those using PCA for iliopsoas in the right leg (Fig. 4 (A)) and for adductor
longus in the left leg (Fig. 4 (B)). When similarity in both shape and magnitude are taken into account,
based on average r and RMSE values , the best number of synergies for predicting unmeasured
excitations using PCA was 6 for iliopsoas in the left leg ($r = 0.93$; RMSE = 0.043), 5 for iliopsoas in
the right leg ($r = 0.79$; RMSE =0.073), 5 for adductors longus in the left leg ($r = 0.97$; RMSE =0.06),
and 5 for adductors longus in the right leg ($r = 0.93$; RMSE = 0.081). Additionally, for most trials,
PCA required fewer synergies (between 3 and 8) than did NMF (between 4 and 10) to achieve the best
synergy extrapolation performance (see the histogram in Fig. 5). For example, for iliopsoas in the left
leg, with 5 to 8 synergies, 86% of trials could achieve the highest r values with PCA and only 54%
trials could do with NMF. Similarly, with 5 to 8 synergies, the smallest RMSE values could be obtained
for 100% of trials using PCA but only 50% of trials using NMF.

3.3   Joint moment prediction

Results of the three-factor ANOVA revealed that MAE values for joint moment matching were
sensitive to both matrix factorization algorithm and number of synergies ($p < 0.01$) but insensitive to
EMG normalization method ($p = 0.12$) for all muscle-leg-joint combinations. For the hipFE moment,
MAE values for both PCA-based and NMF-based synergy extrapolations decreased as the number of

synergies increased ($p < 0.01$). Nonetheless, PCA produced more accurate joint moment matching, as
indicated by smaller MAE values, than did NMF ($p < 0.01$). In contrast, for the hipAA moment, neither
number of synergies nor matrix factorization algorithm had a significant influence on MAE values.
The one exception was adductor longus in the left leg, where MAE values decreased with an increasing
number of synergies for both algorithms, but PCA had significantly smaller MAE values than NMF (p
< 0.01) (Figs. 6 and 7).

PCA-based and NMF-based synergy extrapolation led to different levels of joint moment tracking
accuracy compared to the full EMG-driven calibration (Figs. 6 and 7). For example, for the hipFE
moment with MaxOver, the surface comprising average MAE values for PCA-based synergy
extrapolation intersected with the flat plane of average MAE for the full EMG-driven calibration at 6
synergies for iliopsoas in the left leg, 6 synergies for adductor longus in the left leg and 10 synergies
for adductor longus in the right leg. In contrast, for the hipFE moment with MaxOver, the surfaces
comprising MAE values for NMF-based synergy extrapolation only dropped below the plane of MAE
values for the full EMG-driven calibration for iliopsoas in the right leg at 6 synergies and for adductor
longus in the left leg at 7 synergies.

When r and RMSE values for muscle excitation matching were plotted as a function of MAE values
for joint moment matching (Fig. 8), the observed trends were different for PCA- versus NMF-based
synergy extrapolation. For PCA-based synergy extrapolation, the trends were parabolic, with a small
region of MAE values corresponding to both largest r values and smallest RMSE values. In contrast,
for NMF-based extrapolation, the observed trends were approximately linear, with the smallest MAE
errors corresponding to the largest r values and smallest RMSE values. For both synergy extrapolation
methods, MAE values for joint moment matching were approximately the same in the region where r
values were the largest, and RMSE values were the smallest.

4   Discussion

This study demonstrated that synergy extrapolation is able to predict unmeasured muscle excitations
with reasonable reliability using a well-calibrated EMG-driven model. However, the reliability of the
predictions was heavily influenced by the matrix factorization algorithm used and number of synergies,
while EMG normalization method had little influence on synergy extrapolation results. Our results
clearly showed that PCA was able to generate estimates of unmeasured muscle excitations that were
more accurate in shape and magnitude, more robust to EMG normalization, and more consistent across
all unmeasured muscle-leg combinations in comparison with NMF The results also highlighted that
for PCA-based synergy extrapolation, a relatively low number of synergies, typically 5 or 6, always
provided the most accurate predictions of unmeasured muscle excitations and joint moments
simultaneously, while NMF was either not able to reproduce unmeasured muscle excitations with
reasonable accuracy or required a large number of synergies, typically above 8, to attain a comparable
outcome to PCA.

The synergy extrapolation method optimizes unmeasured synergy weights by tracking joint moments
as closely as possible. We used the muscle synergy concept for the prediction of missing muscle
excitations for several reasons. First, it is commonly theorized that muscle synergies are generated by
the CNS to regulate highly redundant musculoskeletal systems in an efficient manner (Tresch et al.,
2006; Torres-Oviedo and Ting, 2007). Second, the problem of finding unknown time-varying muscle
excitations is reduced to the problem of finding a small number of synergy weights associated with
unmeasured muscles, which significantly decreases the search space for the optimization. Third, there
are no abrupt changes in predicted muscle excitations as is often observed with SO-based approaches,

since predicted muscle excitations are linear combinations of weighted synergy excitations that are
normally smooth. Fourth, by finding unknown synergy vector weights within the context of an EMG-
driven model framework, the method is practically applicable in contrast to previous work that only
demonstrated theoretical feasibility (Bianco et al., 2018). Sixth, synergy extrapolation results
demonstrated that joint moments were matched as accurately as could be achieved by the full EMG-
driven model (Fig. 6) (Meyer et al., 2017). Given these observations, the proposed synergy
extrapolation method may outperform other EMG-driven muscle force estimators that either ignore
deep muscles or use SO-based methods to estimate EMG signals for inaccessible muscles.

For the same number of synergies, PCA generally predicted unmeasured muscle excitations with
noticeably greater accuracy than did NMF (Fig. 4). PCA and NMF are two of the most popular matrix
factorization algorithms used for performing MSA. Although the MSA literature suggests that NMF
generates synergy components that are highly correlated with those generated by PCA (Ivanenko et
al., 2005; Cappellini et al., 2006; Banks et al., 2017), each of the algorithms decomposes a given dataset
with different assumptions and constraints (Ting and Chvatal, 2010; Gallina et al., 2018). PCA is based
on linear algebra that maintains orthogonality among all principal components during factorization,
while NMF is a nonlinear optimization algorithm with potentially non-unique solutions in the non-
negative space (Fig. 1 in the Supplementary Materials). PCA has been rejected by a few studies due to
the inherent non-negative nature of muscle excitations (Ajiboye and Weir, 2009; Banks et al., 2017).
However, our findings demonstrated several benefits of PCA over NMF for synergy extrapolation
purposes. At least three observations help explain why PCA-based synergy extrapolation generally
worked better than did NMF-based synergy extrapolation. First, for an equal number of synergies,
PCA-derived components accounted for more variance in measured muscle excitations than did those
derived using NMF, for 3 to 5 synergies in particular (Table 1). Our findings agree well with the results
of variance accounted for by different numbers of synergies using when PCA and NMF are applied to
high-density EMG data (Gallina et al., 2018). Second, PCA identifies synergy components that tend to
describe the direction of the largest variance in the measured muscle excitations, with subsequent
components being perpendicular to the previous ones, while NMF finds components that tend to
represent the edge of a convex subspace in which all original EMG measurements lie (Ting and
Chvatal, 2010; Lambert-Shirzad and Van der Loos, 2017). Furthermore, any of the data in the reduced-
dimensionality space could be reconstructed due to negative and positive weights allowed by PCA.
However, only data points representing unmeasured muscle excitations that were located within the
defined subspace could be closely reproduced by NMF due to the constraints on non-negativity of the
solutions (Tresch et al., 2006; Ting and Chvatal, 2010). Third, PCA-based synergy extrapolation had
one more design variable - an offset - for each trial that represents the average value of the unmeasured
muscle excitation, thereby adding one extra degree of freedom to the optimization problem. Therefore,
better performance of PCA-based than NMF-based synergy extrapolation may be on account of the
non-negativity constraints within NMF and the extra design variables for PCA, both of which make
the feasible search space of NMF more restricted than that of PCA.

Synergy extrapolation performance was heavily influenced by the number of synergies used. Previous
studies have reported that 3 to 6 synergies are adequate for reconstructing measured EMG signals with
over 90% variance being accounted for during gait (Torres-Oviedo and Ting, 2007; Clark et al., 2009;
Banks et al., 2017; Bianco et al., 2018) consistent with the observations in this study (Table 1). Not
surprisingly, as the number of synergies increased, reconstruction accuracy of unmeasured muscle
excitations using NMF increased before reaching a plateau, which was similar to reconstruction
behavior for measured muscle excitations (Table 1). However, the change in PCA-based synergy
extrapolation performance with an increasing number of synergies exhibited non-monotonic behavior
(Fig. 4). Since synergy extrapolation iteratively predicted muscle excitations for iliopsoas or adductor

longus through an optimization process, this additional flexibility likely allowed the optimizer to
achieve lower joint moment tracking errors than those achieved by the full EMG-driven optimization.
The same phenomenon can be observed for the hip flexion/extension moment using PCA (Fig.7). There,
the synergy extrapolation performance peaked across different unmeasured muscle-leg combinations
with the same number of synergies (Fig. 4). However, due to the non-negativity constraints built in
NMF, the joint moment tracking errors stayed above or around the levels from the full EMG-driven
optimization, which caused synergy extrapolation performance to level off when the number of
synergies was above a certain number. Even though the variability existed in the number of synergies
that generated the best synergy extrapolation results across trials, muscles, and legs, PCA needed fewer
synergies than did NMF (Fig. 5), which simplifies the implementation of the synergy extrapolation
method.

Though EMG normalization methods had no statistically significant influence on synergy extrapolation
performance using PCA and NMF, the variability of average r and RMSE values across all five EMG
normalization methods was considerably less for PCA than that for NMF using the same number of
synergies (Fig. 4). PCA generates principal components that successively maximize variance (Tresch
et al., 2006), which explains why each PCA-derived synergy excitation representing the corresponding
variance for each primary component exhibited significant magnitude reduction (Supplementary
Materials, Fig. 1). Additionally, the low-dimensional space in PCA defined by orthogonal principal
components rotates as data are scaled differently across different muscles, but the rotated principal
components are still highly descriptive of any points in the low-dimensional orthogonal space (Ting
and Chvatal, 2010). However, NMF decomposition is based on assessing the quality of magnitude
approximation, and scaling of original data differently across muscles would lead to deformation of
the convex subspace that the components surround. With certain scaling schemes or normalization
methods, the points accounting for unmeasured muscle excitations may not be able to stay in the NMF
newly-generated subspace. Therefore, NMF-based synergy extrapolation performance would be
sensitive to EMG normalization method, which is in agreement with the conclusions for NMF-based
decomposition by Tresch et al., 2006. For different EMG normalization methods, the optimizer is
presented with a more consistent searching space for PCA than for NMF, which could explain why
PCA is less sensitive to EMG normalization method than is NMF. Therefore, PCA may eliminate the
need to identify the true maximum muscle excitation for EMG normalization purposes in preparation
for synergy extrapolation.

Another observation in this study was that, generally, for a given number of synergies, the non-paretic
side (left leg) had higher reconstruction quality for both measured and unmeasured muscle excitations
than the paretic side (right leg). For the same number of synergies, VAF values calculated for
reconstructed measured muscle excitations in MSA were slightly higher for the non-paretic leg than
the paretic leg (Table 1), which is not consistent with the findings reported by Clark et al., 2009. The
underlying reason for this discrepancy could be the methodological differences between Clark et al.'s
and our studies. For example, we performed MSA on a different set of muscles, which may have
influenced the overall MSA outcomes (Banks et al., 2017) (Steele et al., 2013). Furthermore, the single
subject post-stroke that participated in this study was high-functioning whereas Clark et al. studied 55
number of subjects; the heterogeneity in this clinical population may have also affected the
conclusions. Our study intriguingly found that the reconstruction of unmeasured muscle excitations
through synergy extrapolation was more accurate for the non-paretic side (left leg) than those for the
paretic leg (right leg) (Fig.4), which is possibly due to less consistent EMG data of the healthy/non-
paretic leg or more variable EMG signals across multiple trials for the paretic leg. For instance, the
peaks appeared in the muscle excitations associated with adductor longus during the swing phase (60%-
100% of gait cycle) were more drastic in the paretic side, and synergy extrapolation was unable to

reproduce those as accurately it could do for the non-paretic side (Figs. 2 and 3). Additionally, to
interpret the model calibration difference between legs, we observed that for the non-paretic side (left
leg), as number of synergies increased, the joint moment tracking error for PCA-based synergy
extrapolation started above and then dropped below the plane for the full EMG-driven calibration (Fig.
7). However, this behavior could not be seen in the paretic side (right leg), and the consequence of this
phenomena was that none of the number of synergies could generate both muscle excitations and joint
moments accurately.

Our study involved several important limitations that suggest areas for future investigation. First, we
calibrated the EMG-driven model with a full set of EMG signals to obtain personalized model
parameters (e.g., EMG scale factors, electromechanical delays, musculotendon parameter values, and
geometric coefficients), and synergy extrapolation was feasible when the calibrated model parameters
were held constant. In the future, a challenge will be to develop an optimization problem formulation
where design variables defining both unmeasured synergy vector weights and EMG-driven model
parameters are found simultaneously, the goal being to reproduce both joint moments and unmeasured
muscle excitations accurately. Second, the extraction of measured muscle synergies was performed
before the optimization process. However, since changing some parameter values (e.g., EMG scale
factor) can change the synergy analysis outcome, MSA would need to be performed iteratively within
optimization. Since PCA leads to unique and fast factorization solutions for synergy excitations while
NMF does not (Tresch et al., 2006), PCA would be the preferred decomposition approach. This
iterative decomposition might itself facilitate improvement in the estimation accuracy of unmeasured
muscle excitations. Third, the synergy extrapolation framework was validated on cases where only one
muscle (e.g. iliopsoas or adductor longus) was assumed to be missing. The approach should be
evaluated further on cases where multiple muscles are assumed to be unmeasured concurrently.
Further, this study used gait data from only one single subject post-stroke with extensive EMG data.
Synergy extrapolation needs to be investigated in diverse subject populations, dynamic movement
conditions, and experimental scenarios with other unmeasured muscles.

5   Conclusion

This study showed that synergy extrapolation is a viable option for estimating an unmeasured muscle
excitation using synergy excitations extracted from measured muscle excitations. The study also
demonstrated that methodological choices (i.e. matrix factorization algorithm and the number of
synergies) made before MSA influence affect the accuracy with which unmeasured muscle excitations
can be predicted. The synergy vector weights associated with unmeasured muscles were optimized by
minimizing errors between the EMG-driven model predicted and experimental joint moments. Our
results highlighted the fact that PCA was able to provide more accurate, reliable, and efficient estimates
of unmeasured muscle excitations than was NMF. In general, PCA required 5 or 6 synergies to achieve
the best prediction of unmeasured muscle excitations, and inclusion of additional synergies reduced
synergy extrapolation performance. Better synergy extrapolation performance for PCA may be the
result of the non-negativity restrictions imposed by NMF. Moreover, PCA was less sensitive to EMG
normalization method than NMF, which may reduce the need to simulate the true maximum muscle
excitations reliably for EMG normalization. Synergy extrapolation is of potential utility to address
difficulties in collecting EMG signals from deep muscles inaccessible by surface electrodes, which is
critical when predicting muscle forces with an EMG-driven musculoskeletal model. Our proposed
synergy extrapolation method may facilitate the assessment of human neuromuscular control and
biomechanics after rehabilitation or surgical treatment when EMG data collection is limited.

6   Acknowledgments

This work was conducted with support from the Cancer Prevention and Research Institute of Texas
(CPRIT) RR170026.

7   Author Contributions Statement

CP and BF designed and performed the experiments; DA analyzed the data, prepared figures, and
drafted the manuscript; MS and BF revised the manuscript; DA, MS, CP, and BF approved the final
version of the manuscript.

8   Reference

Ackermann, M., and van den Bogert, A. J. (2010). Optimality principles for model-based prediction
of human gait. *J. Biomech.* 43, 1055–1060.

Ajiboye, A. B., and Weir, R. F. (2009). Muscle synergies as a predictive framework for the EMG
patterns of new hand postures. *J. Neural Eng.* 6, 36004.

Allen, J. L., Kautz, S. A., and Neptune, R. R. (2013). The influence of merged muscle excitation
modules on post-stroke hemiparetic walking performance. *Clin. Biomech.* 28, 697–704.

Allen, J. L., and Neptune, R. R. (2012). Three-dimensional modular control of human walking. *J.
Biomech.* 45, 2157–2163.

Amarantini, D., and Martin, L. (2004). A method to combine numerical optimization and EMG data
for the estimation of joint moments under dynamic conditions. *J. Biomech.* 37, 1393–1404.
doi:https://doi.org/10.1016/j.jbiomech.2003.12.020.

Anderson, F. C., and Pandy, M. G. (2001). Static and dynamic optimization solutions for gait are
practically equivalent. *J. Biomech.* 34, 153–161.

Arnold, E. M., Ward, S. R., Lieber, R. L., and Delp, S. L. (2010). A Model of the Lower Limb for
Analysis of Human Movement. *Ann. Biomed. Eng.* 38, 269–279. doi:10.1007/s10439-009-9852-
5.

Banks, C. L., Pai, M. M., McGuirk, T. E., Fregly, B. J., and Patten, C. (2017). Methodological
choices in muscle synergy analysis impact differentiation of physiological characteristics
following stroke. *Front. Comput. Neurosci.* 11, 78.

Bianco, N. A., Patten, C., and Fregly, B. J. (2018). Can Measured Synergy Excitations Accurately
Construct Unmeasured Muscle Excitations? *J. Biomech. Eng.* 140, 11011.

Bowden, M. G., Clark, D. J., and Kautz, S. A. (2010). Evaluation of abnormal synergy patterns
poststroke: relationship of the Fugl-Meyer Assessment to hemiparetic locomotion.
Neurorehabil. Neural Repair 24, 328–337.

Buchanan, T. S., Lloyd, D. G., Manal, K., and Besier, T. F. (2005). Estimation of muscle forces and
joint moments using a forward-inverse dynamics model. *Med. Sci. Sports Exerc.* 37, 1911.

Cappellini, G., Ivanenko, Y. P., Poppele, R. E., and Lacquaniti, F. (2006). Motor patterns in human
walking and running. *J. Neurophysiol.* 95, 3426–3437.

Clark, D. J., Ting, L. H., Zajac, F. E., Neptune, R. R., and Kautz, S. A. (2009). Merging of healthy
motor modules predicts reduced locomotor performance and muscle coordination complexity
post-stroke. *J. Neurophysiol.* 103, 844–857.

Contessa, P., and Luca, C. J. De (2013). Neural control of muscle force: indications from a
simulation model. *J. Neurophysiol.* 109, 1548–1570.

Crowninshield, R. D., and Brand, R. A. (1981). A physiologically based criterion of muscle force
prediction in locomotion. *J. Biomech.* 14, 793–801.

d'Avella, A., Saltiel, P., and Bizzi, E. (2003). Combinations of muscle synergies in the construction
of a natural motor behavior. *Nat. Neurosci.* 6, 300–308.

Damsgaard, M., Rasmussen, J., Christensen, S. T., Surma, E., and De Zee, M. (2006). Analysis of
musculoskeletal systems in the AnyBody Modeling System. *Simul. Model. Pract. Theory* 14,
1100–1111.

Del Vecchio, A., Ubeda, A., Sartori, M., Azorin, J. M., Felici, F., and Farina, D. (2018). Central
nervous system modulates the neuromechanical delay in a broad range for the control of muscle
force. *J. Appl. Physiol.* 125, 1404–1410.

Delp, S. L., Anderson, F. C., Arnold, A. S., Loan, P., Habib, A., John, C. T., et al. (2007). OpenSim:
open-source software to create and analyze dynamic simulations of movement. *IEEE Trans.*
Biomed. Eng. 54, 1940–1950.

Ebied, A., Kinney-Lang, E., Spyrou, L., and Escudero, J. (2018). Evaluation of matrix factorisation
approaches for muscle synergy extraction. *Med. Eng. Phys.* 57, 51–60.
doi:https://doi.org/10.1016/j.medengphy.2018.04.003.

Falisse, A., Pitto, L., Kainz, H., Hoang, H., Wesseling, M., Van Rossom, S., et al. (2020). Physics-
based simulations to predict the differential effects of motor control and musculoskeletal deficits
on gait dysfunction in cerebral palsy: a retrospective case study. *Front. Hum. Neurosci.* 14, 40.

Farina, D., Cescon, C., and Merletti, R. (2002). Influence of anatomical, physical, and detection-
system parameters on surface EMG. *Biol. Cybern.* 86, 445–456.

Fregly, B. J., Besier, T. F., Lloyd, D. G., Delp, S. L., Banks, S. A., Pandy, M. G., et al. (2012a).
Grand challenge competition to predict in vivo knee loads. *J. Orthop. Res.* 30, 503–513.

Fregly, B. J., Boninger, M. L., and Reinkensmeyer, D. J. (2012b). Personalized neuromusculoskeletal
modeling to improve treatment of mobility impairments: a perspective from European research
sites. *J. Neuroeng. Rehabil.* 9, 18.

Gallina, A., Garland, S. J., and Wakeling, J. M. (2018). Identification of regional activation by
factorization of high-density surface EMG signals: A comparison of Principal Component
Analysis and Non-negative Matrix factorization. *J. Electromyogr. Kinesiol.* 41, 116–123.
doi:https://doi.org/10.1016/j.jelekin.2018.05.002.

He, J., Levine, W. S., and Loeb, G. E. (1991). Feedback gains for correcting small perturbations to
standing posture. *IEEE Trans. Automat. Contr.* 36, 322–332. doi:10.1109/9.73565.

Heintz, S., and Gutierrez-Farewik, E. M. (2007). Static optimization of muscle forces during gait in
comparison to EMG-to-force processing approach. *Gait Posture* 26, 279–288.

Hill, A. V. (1938). The heat of shortening and the dynamic constants of muscle. *Proc. R. Soc.*
London. Ser. B-Biological Sci. 126, 136–195.

Hug, F., Turpin, N. A., Dorel, S., and Guével, A. (2012). Smoothing of electromyographic signals
can influence the number of extracted muscle synergies. *Clin. Neurophysiol. Off. J. Int. Fed.*
Clin. Neurophysiol. 123, 1895.

Ivanenko, Y. P., Cappellini, G., Dominici, N., Poppele, R. E., and Lacquaniti, F. (2005).
Coordination of locomotion with voluntary movements in humans. *J. Neurosci.* 25, 7238–7253.

Kristiansen, M., Madeleine, P., Hansen, E. A., and Samani, A. (2015). Inter-subject variability of
muscle synergies during bench press in power lifters and untrained individuals. *Scand. J. Med.*
Sci. Sports 25, 89–97.

Kumar, D., Rudolph, K. S., and Manal, K. T. (2012). EMG-driven modeling approach to muscle
force and joint load estimations: Case study in knee osteoarthritis. *J. Orthop. Res.* 30, 377–383.

Lambert-Shirzad, N., and Van der Loos, H. F. M. (2017). On identifying kinematic and muscle
synergies: a comparison of matrix factorization methods using experimental data from the
healthy population. *J. Neurophysiol.* 117, 290–302.

Lee, D. D., and Seung, H. S. (1999). Learning the parts of objects by non-negative matrix
factorization. *Nature* 401, 788–791.

Lloyd, D. G., and Besier, T. F. (2003). An EMG-driven musculoskeletal model to estimate muscle
forces and knee joint moments in vivo. *J. Biomech.* 36, 765–776.
doi:https://doi.org/10.1016/S0021-9290(03)00010-1.

Manal, K., and Buchanan, T. S. (2003). A one-parameter neural activation to muscle activation
model: estimating isometric joint moments from electromyograms. *J. Biomech.* 36, 1197–1202.
doi:https://doi.org/10.1016/S0021-9290(03)00152-0.

McLean, S. G., Huang, X., and van den Bogert, A. J. (2005). Association between lower extremity
posture at contact and peak knee valgus moment during sidestepping: implications for ACL
injury. *Clin. Biomech.* 20, 863–870.

Mehrabi, N., Schwartz, M. H., and Steele, K. M. (2019). Can altered muscle synergies control
unimpaired gait? *J. Biomech.* 90, 84–91.

Mehryar, P., Shourijeh, M. S., Rezaeian, T., Khandan, A. R., Messenger, N., O'Connor, R., et al.
(2020). Differences in muscle synergies between healthy subjects and transfemoral amputees
during normal transient-state walking speed. *Gait Posture* 76, 98–103, doi:
https://doi.org/10.1016/j.gaitpost.2019.10.034

Menegaldo, L. L., de Toledo Fleury, A., and Weber, H. I. (2004). Moment arms and musculotendon
lengths estimation for a three-dimensional lower-limb model. *J. Biomech.* 37, 1447–1453.

Meyer, A. J., Eskinazi, I., Jackson, J. N., Rao, A. V, Patten, C., and Fregly, B. J. (2016a). Muscle
synergies facilitate computational prediction of subject-specific walking motions. *Front. Bioeng.*
Biotechnol. 4, 1–26.

Meyer, A. J., Eskinazi, I., Jackson, J. N., Rao, A. V, Patten, C., and Fregly, B. J. (2016b). Muscle
synergies facilitate computational prediction of subject-specific walking motions. *Front. Bioeng.*
Biotechnol. 4, 77.

Meyer, A. J., Patten, C., and Fregly, B. J. (2017). Lower extremity EMG-driven modeling of walking
with automated adjustment of musculoskeletal geometry. *PLoS One* 12, e0179698.

Oliveira, A. S., Gizzi, L., Farina, D., and Kersting, U. G. (2014). Motor modules of human
locomotion: influence of EMG averaging, concatenation, and number of step cycles. *Front.*
Hum. Neurosci. 8, 335.

Olree, K. S., and Vaughan, C. L. (1995). Fundamental patterns of bilateral muscle activity in human
locomotion. *Biol. Cybern.* 73, 409–414.

Péter, A., Andersson, E., Hegyi, A., Finni, T., Tarassova, O., Cronin, N., et al. (2019). Comparing
surface and fine-wire electromyography activity of lower leg muscles at different walking
speeds. *Front. Physiol.* 10, 1283.

Pitto, L., Kainz, H., Falisse, A., Wesseling, M., Van Rossom, S., Hoang, H., et al. (2019). SimCP: A
Simulation Platform to Predict Gait Performance Following Orthopedic Intervention in Children
With Cerebral Palsy   . *Front. Neurorobotics*   13, 54. Available at:
https://www.frontiersin.org/article/10.3389/fnbot.2019.00054.

Racinais, S., Maffiuletti, N. A., and Girard, O. (2013). M-wave, H-and V-reflex recruitment curves
during maximal voluntary contraction. *J. Clin. Neurophysiol.* 30, 415–421.

Reinbolt, J. A., Schutte, J. F., Fregly, B. J., Koh, B. Il, Haftka, R. T., George, A. D., et al. (2005).
Determination of patient-specific multi-joint kinematic models through two-level optimization.
J. Biomech. 38, 621–626. doi:https://doi.org/10.1016/j.jbiomech.2004.03.031.

Ruiz Garate, V., Parri, A., Yan, T., Munih, M., Molino Lova, R., Vitiello, N., et al. (2017).
Experimental Validation of Motor Primitive-Based Control for Leg Exoskeletons during
Continuous Multi-Locomotion Tasks   . *Front. Neurorobotics*   11, 15. Available at:
https://www.frontiersin.org/article/10.3389/fnbot.2017.00015.

S. Shourijeh, M., and McPhee, J. (2014). Forward Dynamic Optimization of Human Gait
Simulations: A Global Parameterization Approach. *ASME J. Comput. Nonlinear Dyn.* 9, 31018.
doi: https://doi.org/10.1115/1.4026266.

Sartori, M., Farina, D., and Lloyd, D. G. (2014). Hybrid neuromusculoskeletal modeling to best track
joint moments using a balance between muscle excitations derived from electromyograms and
optimization. *J. Biomech.* 47, 3613–3621.

Sartori, M., Reggiani, M., Farina, D., and Lloyd, D. G. (2012). EMG-Driven Forward-Dynamic
Estimation of Muscle Force and Joint Moment about Multiple Degrees of Freedom in the
Human Lower Extremity. *PLoS One* 7, e52618. Available at:

https://doi.org/10.1371/journal.pone.0052618.

Sauder, N. R., Meyer, A. J., Allen, J. L., Ting, L. H., Kesar, T. M., and Fregly, B. J. (2019).
Computational Design of FastFES Treatment to Improve Propulsive Force Symmetry During
Post-stroke Gait: A Feasibility Study . *Front. Neurorobotics* 13, 80. Available at:
https://www.frontiersin.org/article/10.3389/fnbot.2019.00080.

Seth, A., Hicks, J. L., Uchida, T. K., Habib, A., Dembia, C. L., Dunne, J. J., et al. (2018). OpenSim:
Simulating musculoskeletal dynamics and neuromuscular control to study human and animal
movement. *PLoS Comput. Biol.* 14.

Shao, Q., Bassett, D. N., Manal, K., and Buchanan, T. S. (2009). An EMG-driven model to estimate
muscle forces and joint moments in stroke patients. *Comput. Biol. Med.* 39, 1083–1088.
doi:https://doi.org/10.1016/j.compbiomed.2009.09.002.

Shourijeh, M. S., Flaxman, T. E., and Benoit, D. L. (2016). An approach for improving repeatability
and reliability of non-negative matrix factorization for muscle synergy analysis. *J.*
Electromyogr. Kinesiol. 26, 36–43, doi: https://doi.org/10.1016/j.jelekin.2015.12.001

Shourijeh, M. S., and Fregly, B. J. (2020). Muscle Synergies Modify Optimization Estimates of Joint
Stiffness During Walking. *J. Biomech. Eng.* 142, , 011011. doi:
https://doi.org/10.1115/1.4044310.

Shuman, B. R., Schwartz, M. H., and Steele, K. M. (2017). Electromyography Data Processing
Impacts Muscle Synergies during Gait for Unimpaired Children and Children with Cerebral
Palsy . *Front. Comput. Neurosci.* 11, 50. Available at:
https://www.frontiersin.org/article/10.3389/fncom.2017.00050.

Steele, K. M., Tresch, M. C., and Perreault, E. J. (2013). The number and choice of muscles impact
the results of muscle synergy analyses. *Front. Comput. Neurosci.* 7, 105.

Taylor, R. (1990). Interpretation of the correlation coefficient: a basic review. *J. diagnostic Med.*
Sonogr. 6, 35–39.

Thelen, D. G., Anderson, F. C., and Delp, S. L. (2003). Generating dynamic simulations of
movement using computed muscle control. *J. Biomech.* 36, 321–328.

Ting, L. H., and Chvatal, S. A. (2010). Decomposing muscle activity in motor tasks. *Mot. Control*
Theor. Exp. Appl. Oxf. Univ. Press. New York, 102v – 138.

Ting, L. H., and Macpherson, J. M. (2005). A limited set of muscle synergies for force control during
a postural task. *J. Neurophysiol.* 93, 609–613.

Torres-Oviedo, G., and Ting, L. H. (2007). Muscle synergies characterizing human postural
responses. *J. Neurophysiol.* 98, 2144–2156.

Tresch, M. C., Cheung, V. C. K., and d'Avella, A. (2006). Matrix factorization algorithms for the
identification of muscle synergies: evaluation on simulated and experimental data sets. *J.*
Neurophysiol. 95, 2199–2212.

Tresch, M. C., Saltiel, P., and Bizzi, E. (1999). The construction of movement by the spinal cord.
Nat. Neurosci. 2, 162–167.

Walter, J. P., Kinney, A. L., Banks, S. A., D'Lima, D. D., Besier, T. F., Lloyd, D. G., et al. (2014).
Muscle synergies may improve optimization prediction of knee contact forces during walking. *J.*
Biomech. Eng. 136, 21031.

Zajac, F. E. (1989). Muscle and tendon: properties, models, scaling, and application to biomechanics
and motor control. *Crit. Rev. Biomed. Eng.* 17, 359–411.

Zonnino, A., and Sergi, F. (2019). Model-based estimation of individual muscle force based on
measurements of muscle activity in forearm muscles during isometric tasks. *IEEE Trans.*
Biomed. Eng., 1. doi:10.1109/TBME.2019.2909171.

769 **Table 1** mean VAF values across all trials that represent reconstruction quality of measured muscle excitations
770 using 3 to 7 synergies and 6 EMG normalization methods when iliopsoas or adductor longus EMG channel is
771 assumed to be unmeasured (The VAF values represented by PCA/NMF; grey shading represents a significant
772 difference between PCA and NMF with matched EMG normalization method and matched number of synergies,
773 P ≤ 0.05).

Unmeasured muscle	EMG normalization	Left leg (non-paretic)					Right leg (paretic)				
		Number of synergies					Number of synergies				
		3	4	5	6	7	3	4	5	6	7
Iliopsoas	MaxOver	92.7/88.8	97.7/95.4	99.1/98.0	99.7/98.8	99.9/99.3	89.7/82.3	95.2/91.0	98.1/95.4	99.2/97.5	99.7/98.5
	MaxPer	90.8/86.5	97.2/94.7	98.8/97.5	99.5/98.5	99.8/99.1	85.6/77.7	93.1/86.8	97.2/93.1	98.7/96.3	99.5/97.6
	VarOver	90.1/86.4	97.4/94.4	98.9/97.6	99.6/98.5	99.9/99.0	87.8/77.8	94.6/89.1	97.8/94.1	99.1/96.4	99.7/97.6
	VarPer	90.9/87.3	97.2/94.6	98.8/97.5	99.5/98.5	99.8/99.1	85.7/77.9	93.3/86.6	97.2/92.7	98.7/96.2	99.5/97.8
	MagPer	92.5/88.5	97.4/95.0	98.9/97.7	99.6/98.6	99.9/99.2	83.5/75.6	91.9/85.9	96.7/91.5	98.8/95.6	99.5/97.6
Adductor longus	MaxOver	94.0/90.1	97.9/95.8	99.0/97.8	99.7/98.8	99.9/99.3	88.8/81.3	94.3/90.5	97.5/94.8	99.0/97.2	99.7/98.5
	MaxPer	91.9/87.8	97.2/94.7	98.7/97.2	99.5/98.4	99.8/99.0	85.1/77.2	91.7/86.4	96.5/92.6	98.4/95.7	99.5/97.5
	VarOver	93.2/87.6	97.5/94.6	98.8/97.4	99.6/98.4	99.9/99.0	86.1/78.0	93.2/88.6	96.8/93.3	98.6/95.5	99.6/97.5
	VarPer	92.4/88.3	97.3/94.7	98.7/97.2	99.5/98.5	99.8/99.0	84.7/76.5	92.1/86.1	96.5/92.1	98.5/95.8	99.5/97.6
	MagPer	92.0/87.7	96.9/94.3	98.6/97.1	99.5/98.2	99.8/98.8	83.0/74.4	90.8/85.1	96.3/90.4	98.4/95.1	99.5/96.7

Figure captions:

Fig. 1 Flowchart of EMG-driven model-based synergy extrapolation framework. The framework
comprises of two main blocks: synergy extrapolation block and EMG-driven musculoskeletal model
block. In the synergy extrapolation block, measured muscle excitations are decomposed to determine
a set of measured synergy excitations, which are used as building blocks to construct unmeasured
muscle excitations. In the EMG-driven model block, the joint moments are predicted by receiving both
the measured muscle excitations from experimental measurements along with the unmeasured muscle
excitations as the output of the synergy extrapolation block. The unmeasured synergy variables were
calibrated through the optimization process with an objective function of minimizing the difference
between experimental and model-predicted joint moments. For NMF-based synergy extrapolation, the
unmeasured synergy variables include the unmeasured synergy weight vectors, while for PCA-based
synergy extrapolation, the unmeasured synergy variables also include an additional set of design
variables that represent mean values of unmeasured muscle excitations.

Fig. 2 Representative results of reconstructed unmeasured muscle excitations across all calibration walking trials at the same speed using synergy extrapolation framework (black line: average experimental curve; red line: PCA-based synergy extrapolation; green line: NMF-based synergy extrapolation; shade area representing ± 1 standard deviation). The measured synergies were extracted using MaxOver EMG normalization method with 6 synergies. Dataset is reported for the whole gait cycle with 0% being heel-strike and 100% being consecutive heel-strike events for both legs (right leg: paretic, left leg: non-paretic). *r* and RMSE values are computed between average experimental and synergy-extrapolated muscle excitations.

Fig. 3 Representative results of average reconstructed unmeasured muscle excitations across all evaluation walking trials at the same speed using synergy extrapolation framework (black line: average experimental curve; red line: PCA-based synergy extrapolation; green line: NMF-based synergy extrapolation; shade area representing ± 1 standard deviation). The measured synergies, in this case, were extracted using MaxOver EMG normalization method with 6 synergies. Dataset is reported for the whole gait cycle with 0% being heel-strike and 100% being consecutive heel-strike events for both legs (right leg: paretic, left leg: non-paretic). *r* and RMSE values are derived between average experimental and synergy-extrapolated muscle excitations.

Fig. 4 Average *r* and RMSE values for the reconstruction of iliopsoas (A) and adductor longus (B)
muscle excitations across all trials (including both calibration trials and evaluation trials) for both legs
(right leg: paretic, left leg: non-paretic) with 3 to 10 synergies (red markers: PCA-based synergy
extrapolation; green markers: NMF-based synergy extrapolation; 5 EMG normalization methods are
represented by different marker shapes; black bars represent a significant difference between PCA and
NMF when matched for the number of synergies, $p \leqslant 0.05$)

Fig. 5 Distribution of the number of synergies that can provide maximum r values or minimum RMSE values across all calibration and evaluation trials with all the five EMG normalization methods. In each histogram, the horizontal axis reports the number of synergies, and the vertical axis shows the frequency with which the number of synergies can generate the best prediction of iliopsoas or adductor longus muscle excitations in terms of shape (indicated by r values) and magnitude (indicated by RMSE values). Red and green bars represent PCA-based and NMF-based synergy extrapolations, respectively. The right leg is paretic, and the left leg is non-paretic.

Fig. 6 Average joint moment prediction across all calibration trials when performing full EMG-driven model calibration on step 1 (blue dash line) and synergy extrapolation on step 2 with PCA (red line) or NMF (green line), respectively. Black lines stand for the experimental joint moments from inverse dynamics, and the grey shaded areas represent ± 1 standard deviation. Six synergies are used for the generation of these representative results. Data is reported for the whole gait cycle with 0% being heel-strike and 100% being consecutive heel-strike events for both legs (right leg: paretic, left leg: non-paretic).

Fig. 7 3D plots report the average MAE values for hip joint moment prediction across all trials by using 6 EMG normalization methods with an increase of the number of synergies from 3 to 10. The MAE values are averaged across all calibration and evaluation trials at 2 walking speeds for both legs (right leg: paretic, left leg: non-paretic). PCA- and NMF-based synergy extrapolations are indicated in red and green surfaces, respectively. The flat purple planes demonstrate the average MAE values for joint moment prediction with a full set of EMG signals at a specific leg. The circle lines on the bottom walls indicate where 3D surfaces of average MAE values for PCA- and NMF-based synergy extrapolation first intersect with the plane of average MAE for the full EMG-driven calibration, respectively, as the number of synergies increases. The absence of a circle means that there was no intersection corresponding to that synergy.

Fig. 8 The trade-off relationship between the accuracy of joint moment tracking (indicated by MAE values in the horizontal axis) and accuracy of unmeasured muscle excitation reconstruction (indicated by r values or RMSE values in the y-axis) (red markers: PCA-based synergy extrapolation; green markers: NMF-based synergy extrapolation; 5 EMG normalization methods are represented by different marker shapes; the right leg is paretic, and the left leg is non-paretic).