Musculoskeletal Model Personalization Affects Metabolic Cost Estimates for Walking

Marleny Arones[1], Mohammad Shourijeh[1], Carolynn Patten[2], Benjamin J. Fregly[1*]

[1]Rice University, Department of Mechanical Engineering, Houston, TX, USA

[2]University of California, Department of Physical Medicine and Rehabilitation, Davis, CA, USA

* Correspondence:
B.J. Fregly
fregly@rice.edu

Keywords: Metabolic Cost, Model Personalization, Stroke, Gait, EMG-Driven

Abstract

Assessment of metabolic energy cost as a metric for human performance has expanded across various
fields within the scientific, clinical, and engineering communities. As an alternative to measuring
metabolic cost experimentally, musculoskeletal models incorporating metabolic cost models have
been developed. However, to utilize these models for practical applications, the accuracy of their
metabolic cost predictions requires improvement. Previous studies have reported the benefits of using
personalized musculoskeletal models for various applications, yet no study has evaluated how model
personalization affects metabolic cost estimation. This study investigated the effect of
musculoskeletal model personalization on estimates of metabolic cost of transport (CoT) during post-
stroke walking using three commonly used metabolic cost models. We analyzed data previously
collected from two male stroke survivors with right-sided hemiparesis. The three metabolic cost
models were implemented within three musculoskeletal modeling approaches involving different
levels of personalization. The first approach used a scaled generic OpenSim model and found muscle
activations via static optimization (SOGen). The second approach used a personalized EMG-driven
musculoskeletal model with personalized functional axes but found muscle activations via static
optimization (SOCal). The third approach used the same personalized EMG-driven model but
calculated muscle activations directly from EMG data (EMGCal). For each approach, the muscle
activation estimates were used to calculate each subject's cost of transport (CoT) at different gait
speeds using three metabolic cost models (Umberger 2003, Umberger 2010, and Bhargava 2004).
The calculated CoT values were compared with published CoT trends as a function of stance time,
double support time, step positions, walking speed, and severity of motor impairment (i.e., Fugl-
Meyer score). Overall, U10-SOCal, U10-EMGCal, U03-SOCal, and U03-EMGCal were able to
produce slopes between CoT and the different measures of walking asymmetry that were statistically
similar to those found in the literature. Although model personalization seemed to improve CoT
estimates, further tuning of parameters associated with the different metabolic cost models in future
studies may allow for realistic CoT predictions. An improvement in CoT predictions may allow
researchers to predict human performance, surgical, and rehabilitation outcomes reliably using
computational simulations.

1 Introduction

Metabolic energy expenditure has been used to evaluate human performance during daily activities
such as walking (Waters and Mulroy, 1999; Donelan et al., 2002, 2008; Mian et al., 2006; Long III
and Srinivasan, 2013) and athletic activities such as running (Roberts et al., 1998; Chang and Kram,
1999; Teunissen et al., 2007; Franz et al., 2012; Long III and Srinivasan, 2013) and cycling (Davies,
1980; Gnehm et al., 1997; Neptune and Van Den Bogert, 1997; McDaniel et al., 2002; van der
Woude et al., 2008). This quantity has also been adopted as a metric to evaluate the design or
operational settings of assistive devices (Malcolm et al., 2013; Collins et al., 2015; Galle et al.,
2017). Additional applications for which knowledge of metabolic cost is useful include prescription
of training intensities (American College of Sports Medicine, 2000), advancement of geriatric
medicine (Mian et al., 2006; Canavan et al., 2009; Corbett et al., 2017), treatment of clinical gait
disorders (Waters and Mulroy, 1999), and to monitor energy intake and expenditure in obese patients
(Brychta et al., 2010). Though various methods exist to measure metabolic cost (Lam and Ravussin,
2016), technical challenges resulting from cost, feasibility, or medical restrictions of the subject
population limit the availability of metabolic cost measurements.

With advances in computational biomechanics, musculoskeletal and metabolic cost models have
emerged as tools to estimate metabolic energy. These tools have been used in various studies to
predict human movement and response to mechanical interventions. Uchida et al. (Uchida et al.,
2016b) used the OpenSim simulation framework (Seth et al., 2018) to optimize the design of an
assistive device intended to reduce the metabolic cost of running. Dembia et al. (Dembia et al., 2017)
simulated an ideal assistive device to minimize the metabolic cost of several individuals walking with
heavy loads. Fey et al. (Fey et al., 2012) developed an optimization to identify an optimal prosthetic
foot stiffness to minimize metabolic cost for amputee walking. Miller et al. (Miller et al., 2013) used
direct collocation optimal control to predict a gait pattern that reduced metabolic cost while
minimizing peak axial knee joint contact forces. By predicting the factors, designs, or movements
that minimize metabolic energy expenditure, these studies highlight the potential benefits of
combining musculoskeletal and metabolic cost models. However, for these predictions to become
realistic for practical applications, the accuracy of these combined models needs to improve (Miller,
2014), (Koelewijn et al., 2019).

While published studies that estimated metabolic cost during walking have focused on using scaled
generic musculoskeletal models, several studies have reported that personalization of anatomical and
physiological characteristics of a musculoskeletal model can influence prediction of muscle forces,
joint moments, and novel movements, and these factors also play a role in metabolic cost
calculations. Reinbolt et al. (Reinbolt et al., 2007) found that personalization of the functional joint
axes is crucial to obtain reliable inverse dynamic joint moments. Studies like (Lloyd and Besier,
2003; Buchanan et al., 2005; Shao et al., 2009; Sartori et al., 2012; Meyer et al., 2017) reported large
improvements in joint moment matching when muscle force-generating properties were personalized
using an EMG-driven model. Gerus et al. (Gerus et al., 2013) demonstrated that personalizing knee
geometry *and* corresponding muscle-tendon parameter values resulted in improved predictions of
knee contact force compared to predictions generated by a generic model. Since muscle force
estimates depend on joint moments, and the production of muscle forces and activations depends on
muscle parameters, the confounding effect of model personalization may also affect metabolic cost
estimates. However, to the best of the authors' knowledge, no study to date has explored this
possibility.

This study evaluated the influence of musculoskeletal model personalization on metabolic cost
estimates of walking post-stroke. To evaluate the physical realism of different metabolic cost
modeling methods, we compared metabolic cost estimates to trends reported in the literature. For

individuals post-stroke, Finley and Bastian (Finley and Bastian, 2017) reported a decrease in
metabolic cost as walking speed increased and severity of motor impairment decreased. In contrast,
for the same subject population, Finley and colleagues (Finley et al., 2015; Finley and Bastian, 2017)
also reported that metabolic cost increased as differences in step position, stance time, and double
support time increased. The present study evaluated a previously acquired gait data set from which
metabolic cost measurements were not available. Our goal was to determine whether any
combination of musculoskeletal modeling method and metabolic cost modeling method could
reproduce all five physically realistic metabolic cost trends observed in the literature. Since metabolic
energy has been adopted as a tool in various scientific, clinical, and engineering fields to evaluate
human performance, surgical, and rehabilitation outcomes, the results of this study may be useful for
identifying which modeling methods are most likely to predict physically realistic metabolic cost
estimates.

2   Methods

2.1   Experimental Data and Data Processing

Experimental walking data collected from two male stroke survivors – one high functioning and one
low functioning - were used as inputs to the metabolic cost analyses (see Table 1 for subject details).
Data collected from both subjects included marker-based motion capture, ground reaction, and surface
and fine-wire electromyographic (EMG) data (Table 2, 16 channels per leg). The data were collected
at different walking speeds using a split-belt instrumented treadmill with belts tied. Walking speeds
ranged from slower than self-selected to a maximum comfortable speed. For the high functioning
subject, the speed range was from 0.4 to 0.8 m/s in increments of 0.1 m/s, while for the low functioning
subject, the speed range was from 0.35 to 0.65 m/s also in increments of 0.1 m/s. Additional
experimental data were also collected for static and isolated joint movement trials. A static standing
trial was collected for model scaling purposes. Isolated joint motion trials were collected for each hip,
knee, and ankle for purposes of personalizing the model's lower body joint axes. All functional axes
for the joint of interest were exercised during each isolated joint motion trial (Reinbolt et al., 2005,
2008).

The ground reaction and marker motion data were low-pass filtered using a fourth-order zero-phase
lag Butterworth filter with a cut-off frequency of $7/t_f$ Hz (Hug, 2011), where t_f is the period of the gait
cycle being processed (Meyer et al., 2017). On average, this variable cut-off frequency caused data
collected at a normal walking speed to be filtered at approximately 6 Hz.

The EMG data were high-pass filtered (40 Hz), demeaned, rectified, and low-pass filtered ($3.5/t_f$ Hz)
using a fourth-order zero-phase lag Butterworth filter. EMG amplitudes for each muscle were
normalized to the maximum value over all trials and resampled to 101 time points per gait cycle, as
described in (Meyer et al., 2017).

2.2   Musculoskeletal Model

A generic full-body OpenSim musculoskeletal model (Rajagopal et al., 2016; Seth et al., 2018) served
as the starting point for all three metabolic cost analyses. The generic model used for both subjects
started with 40 Hill-type muscle-tendon actuators per leg and 37 degrees of freedom (DOFs), including
3 DOF hip joints, 1 DOF knee joints, 2 DOF ankle joints. For the high functioning subject, six muscles
without related EMG data were eliminated (extensor hallucis longus, flexor hallucis longus, gracilis,
piriformis, sartorius, tensor fascia latae). For the low functioning subject, seven muscles without related
EMG data were eliminated (extensor digitorum longus, flexor digitorum longus, extensor hallucis

longus, flexor hallucis longus, gracilis, piriformis, sartorius). The remaining muscles actuated hip
flexion-extension, hip adduction-abduction, hip internal-external rotation, knee flexion-extension,
ankle flexion-extension, and ankle inversion–eversion on each leg.

2.3   Joint Model Personalization

Personalization of the joint functional axes for the hip, knee, and ankle on each leg was performed by
following a two-step process. First, the geometry of the generic OpenSim model was scaled to match
the dimensions of each subject using OpenSim's Scale tool and the static standing trial. Second, marker
positions and functional axes of the model's lower body's joints were personalized as described in
(Reinbolt et al., 2005, 2008; Meyer et al., 2016; Sauder et al., 2019). The personalization process
involved using nonlinear optimization to adjust the positions and orientations of the model's lower
body joints and marker triads placed on the body segments. The cost function minimized the sum of
squares of errors between the experimental and model marker positions from all isolated joint motion
trials and one walking trial analyzed together. The optimization process was performed using Matlab's
lsqnonlin algorithm, which iteratively ran OpenSim's inverse kinematics analyses to calculate marker
location errors.

2.4   Muscle-tendon Model and Geometry Personalization

Experimental data from ten gait trials collected at each available speed were used to calibrate an EMG-
driven model for each subject. Before performing EMG-driven model calibration, we analyzed marker
data from each gait trial using OpenSim's Inverse Kinematics tool to generate joint angle trajectories.
OpenSim's Inverse Dynamics tool was then used to calculate the joint moments ~~to be~~ produced by
muscle forces. Next, a surrogate model of each subject's musculoskeletal geometry was generated to
allow the EMG-driven model to modify muscle geometry (Meyer et al., 2016, 2017). The surrogate
model was developed by sampling a wide range of joint angle combinations for the lower limbs using
a Latin hypercube design. The muscle-tendon lengths and moment arms for each muscle were then
calculated using OpenSim's Muscle Analysis tool. Linear regression using least squares was used to fit
muscle-tendon lengths as a polynomial function of the corresponding joint angles actuated by each
muscle. Muscle-tendon velocity was then defined as the first derivative of the muscle-tendon length
polynomial with respect to time while muscle moment arms were defined as the negative of the first
derivative of the muscle-tendon length polynomial with respect to corresponding joint angles (An et
al., 1984).

To personalize the model's muscle-tendon force-generating properties, we allowed our EMG-driven
model calibration process to modify three types of parameters: EMG-to-activation parameters
(electromechanical delays, activation dynamics time constants, activation non-linearization shape
factors, and EMG scale factors), Hill-type muscle-tendon model parameters (optimal muscle fiber
lengths, and tendon slack lengths), and surrogate musculoskeletal geometry parameters. With the use
of Matlab's *fmincon* algorithm using sequential quadratic programming, the model parameter values
were adjusted to best match calculated experimental inverse dynamic moments and published passive
joint moments (Silder et al., 2007). Additionally, penalty terms were added to the cost function to
discourage substantial divergence of model parameters from their original values. A detailed
description of our EMG-driven modeling approach can be found in Meyer et al. (Meyer et al., 2017).

2.5   Analysis

We developed three musculoskeletal modeling approaches, each derived from the same generic
OpenSim model and implemented in Matlab, to evaluate the extent to which model personalization

affects estimated metabolic cost. The first approach used a scaled generic OpenSim model where
muscle activations were calculated via static optimization using quadratic programming (SO-Gen). The
second approach used the subject's personalized EMG-driven model but with muscle activations found
via static optimization (SO-Cal). The third approach used the same personalized EMG-driven model
but found muscle activations using the EMG-driven model (EMG-Cal) (Figure 1). The muscle
activations found by each approach were used to calculate each subject's cost of transport (CoT in
J/m/kg) for different gait speeds using metabolic cost models published by Umberger (Umberger et al.,
2003) (U03), Umberger et al. (Umberger, 2010) (U10), and Bhargava et al. (Bhargava et al., 2004)
(B04).

To evaluate the physical realism of each musculoskeletal modeling approach/metabolic cost model
combination, we identified five experimental trends in the literature for how CoT varies with other
quantities for individuals post-stroke. The first three quantities were step position asymmetry, stance
time asymmetry, and double-support time asymmetry that were reported to increase as the cost of
transport increased (Finley and Bastian, 2017). Inversely, the two other quantities were speed and Fugl-
Meyer score that were observed to decrease as the cost of transport increased. Differences or
asymmetries in step positions, stance time, and double-support time were calculated as specified in
(Finley et al., 2015; Finley and Bastian, 2017). The relationship between estimated CoT and the
variables of interest was compared against the linear correlations reported in (Finley and Bastian,
2017). Thus, a t-test analysis, comparing the slopes produced between CoT estimates and variables of
interest against those found in the literature, was performed for each modeling approach and metabolic
cost model. The t-test analysis was performed for all correlations except CoT and Fugl-Meyer score
due to the limitation of only two subjects.

To investigate the effect of either increasing or decreasing muscle specific tension away from its
nominal value of 61 N/cm (Meyer et al., 2017), we iteratively ran static optimization with varying
muscle specific tension values using the personalized model for the high functioning patient. All
iterations of this static optimization were performed at a fixed gait speed of 0.5 m/s (self-selected
speed).

3 Results

When the nine modeling combinations were used to predict the positive or negative correlations of
CoT with other quantities as reported in (Finley and Bastian, 2017), U10-SOCal, U10-EMGCal,
U03SO-Cal, and U03-EMGCal predicted statistically similar slopes for all four variables (excluding
Fugl-Meyer score) ($p \geq 0.05$) (Table 3) (Figures 2-5). In addition, all modeling combinations showed
a decrease in CoT as Fugl-Meyer score increased (Table 4-5).

An inverse relationship was observed between changes in CoT and changes in muscle specific tension
(Figure 6). To compensate for CoT overestimation by both Umberger models (U03 and U10), muscle
specific tension had to be increased to 75-135 N/cm^2 for U03 and 90-135 N/cm^2 for U10. With this
change, the resulting CoT estimates fit within one standard deviation of the average reported by Finley
and Bastian, 2017. Inversely, to compensate for CoT underestimation by the Bhargava model, muscle
specific tension had to be decreased to 37 N/cm^2 for the resulting CoT estimates to lie within the range
of CoT values published by (Miller, 2014; Koelewijn et al., 2019).

4 Discussion

This study evaluated the effect of musculoskeletal model personalization on metabolic cost estimates
for walking obtained using three common metabolic cost models: Umberger 2003 (U03), Umberger *et*

al. 2010 (U10), and Bhargava et al. 2004 (B04). These three models were implemented within three
musculoskeletal modeling approaches: scaled generic musculoskeletal models with muscle activations
found by static optimization (SOGen) (Shourijeh and Fregly, 2020), calibrated EMG-driven
musculoskeletal models with muscle activations found by static optimization (SOCal), and calibrated
EMG-driven musculoskeletal models with muscle activations found from experimental EMG data
(EMGCal). These nine combinations were investigated using previously published walking data
collected from two individuals post-stroke, and correlations between estimated CoT and walking
speed, step position asymmetry, step length asymmetry, stance time asymmetry, double support time
asymmetry, and Fugl-Meyer score were compared with published data (Finley and Bastian, 2017). All
nine modeling combinations predicted the correct positive and negative correlations between CoT and
the five selected quantities as observed in a post-stroke population. However, only the personalized
models (SOCal andEMGCal) paired with Umberger 2003 and Umberger 2010 produced statistically
similar slopes for all four quantities in comparison to the linear models found in the literature. These
findings suggest that a calibrated EMG-driven musculoskeletal model combined with either one of
Umberger's models improved CoT estimates.

In general, only SOGen predicted CoT estimates that were within one standard deviation of the
published experimental average for all three metabolic cost models. EMGCal predicted CoT estimates
that were larger than those obtained from both SOGen and SOCal for all three metabolic cost models.
Additionally, the level of model personalization did not affect the tendency of Bhargava's model to
underestimate CoT nor the tendency of both Umberger's models to overestimate CoT (Figure 2). A
similar pattern was reported in Miller (Miller, 2014), which analyzed the performance of B04 and U10,
among other models. However, in our study, the combination of U03 or U10 used with EMGCal
produced CoT values that were almost twice as large as published CoT averages. Studies such as Ong
(Ong et al., 2019), which used a model based on U03, and Miller (Miller, 2014) which used U10,
reported CoT values within the range of 2 to 8 J/Kg*m, whereas U10-EMG resulted in CoT values
ranging from 7 to 15 J/Kg*m. This magnitude difference was not expected but may be due to
minimization of a metabolic cost term within the optimization cost function used for estimating muscle
activations in both studies. The observation of minimization of metabolic cost may explain why CoT
magnitudes produced by SOGen were closest to those reported in (Finley and Bastian, 2017) for all
three metabolic cost models. Additionally, in contrast to the studies mentioned above, our study also
used calibrated EMG-driven models to calculate CoT. The absence of muscle activation minimization
may help explain why our EMGCal paired with any metabolic model produced CoT estimates that
were larger than those produced by SOGen and SOCal.

Although all modeling combinations produced the positive or negative trends relative to published
data, not all of them resulted in slopes that were statistically similar to those found in the literature.
This finding is similar to a recent study by Koelwijn et al. (Koelewijn et al., 2019), where several
metabolic cost models (including U03 and B04) were shown to predict the correct trends for CoT as a
function of walking slope or gait speed in accordance to trends found in the literature (Margaria, 1968).
Additionally, Koelwijn et al. (Koelewijn et al., 2019) found that B04 tended to underestimate CoT
values in comparison to experimental values, which is in agreement with our findings.

Although future work can be directed toward optimizing parameters within the metabolic cost models,
we tested how changing muscle specific tension affected the magnitude of the predicted CoT values.
As reported in (Russell, Esposito and Miller, 2018), decreasing the muscle specific tension increased
metabolic cost and vice versa. Bhargava's model needed muscle specific tension to be around 37 N/cm^2
rather than 61 N/cm^2 to match experimental averages the best. This new value is closer to values used
within OpenSim (Delp et al., 2007) and studies such as (Miller, 2014; Koelewijn et al., 2019). Future

work to optimize muscle specific tension could improve the magnitude of CoT predictions (Hamner
and Delp, 2013; Uchida et al., 2016a; Falisse et al., 2019).

One of the major limitations of this study was the absence of experimental metabolic cost data for
evaluating our model-based estimates. This limitation makes it difficult to determine the accuracy of
our modeling methods. Furthermore, none of the musculoskeletal models used in our analysis included
muscles to actuate the upper body; therefore, the metabolic cost for all nine cases may be an
underestimation. However, since Finley and Bastian's dataset were collected from subjects who were
allowed to use handrails, their measurements may be underestimates as well. Additionally, we were
limited to two subjects with the extensive EMG datasets required to perform the study, which could
have impacted our statistical results.

In conclusion, this study investigated the effect of musculoskeletal model personalization on estimated
metabolic cost for post-stroke walking as obtained from three commonly used metabolic cost models
and three musculoskeletal model personalization methods. Previously collected walking data from two
stroke survivors were used to analyze correlations between estimated CoT and various variables
commonly associated with gait asymmetry. Although all modeling combinations exhibited the correct
positive and negative correlations observed in (Finley and Bastian, 2017), U10-SOCal, U10-EMGCal,
U03-SOCal, and U03-EMGCal produced statistically similar slopes to those found in the literature for
all gait asymmetry variables (excluding Fugl-Meyer score). While the estimated CoT values were not
close to reference CoT averages (Finley and Bastian, 2017), we observed that additional tuning of
muscle specific tension easily changed the CoT estimates. Therefore, future tuning of muscle specific
tension along with other parameters associated with the metabolic cost models may further improve
CoT estimates. Our results suggest the use of a personalized EMG-driven model to estimate muscle
forces paired with either one of Umberger's energy expenditure models to help the prediction of CoT
estimates during walking for individuals post-stroke. Since metabolic energy has been adopted as a
tool in various scientific, clinical, and engineering fields to evaluate performance, surgical, and
rehabilitation outcomes, the results of this study may help improve the reliability of metabolic cost
predictions generated using musculoskeletal models.

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

6   Author Contributions

CP and BJF performed experiments; MA performed all model personalization tasks, prepared
figures, and drafted manuscript; MA and MS analyzed data; MA, MS, and BF interpreted results of
experiments; MA, MS, CP and BF revised manuscript.

7   Funding

This work was conducted with support from the Cancer Prevention and Research Institute of Texas
(CPRIT) funding RR170026 and NSF Graduate Research Fellowship.

8   References

American College of Sports Medicine (2000). *American College of Sports Medicine Guide to*
Exercise Testing and Prescription. 6th Edition. Lippincott Williams & Wilkins.

An, K. N., Takahashi, K., Harrigan, T. P., and Chao, E. Y. (1984). Determination of Muscle
Orientations and Moment Arms. *J. Biomech. Eng.* 106, 280–282. doi:10.1115/1.3138494.

Bhargava, L. J., M. G. Pandy, and Anderson, F. C. (2004). A phenomenological model for estimating
metabolic energy consumption in muscle contraction. *J. Biomech.* 37, 81–88.

Brychta, R., Wohlers, E., Moon, J., and Chen, K. (2010). Energy expenditure: measurement of
human metabolism. *IEEE Eng. Med. Biol. Mag.* 29, 42–47.

Buchanan, T. S., Lloyd, D. G., Manal, K., and Besier, T. F. (2005). Estimation of muscle forces and
joint moments using a forward-inverse dynamics model. *Med. Sci. Sports Exerc.* 37, 1911–1916.

Canavan, P. K., Cahalin, L. P., Lowe, S., Fitzpatrick, D., Harris, M., and Plummer-D'Amato, P.
(2009). Managing gait disorders in older persons residing in nursing homes: a review of
literature. *J. Am. Med. Dir. Assoc.* 10, 230–237.

Chang, Y.-H., and Kram, R. (1999). Metabolic cost of generating horizontal forces during human
running. *J. Appl. Physiol.* 86, 1657–1662.

Collins, S. H., Wiggin, M. B., and Sawicki, G. S. (2015). Reducing the energy cost of human
walking using an unpowered exoskeleton. *Nature* 522, 212–215.

Corbett, D. B., Wanigatunga, A. A., Valiani, V., Handberg, E. M., Buford, T. W., Brumback, B., et
al. (2017). Metabolic costs of daily activity in older adults (chores xl) study: Design and
methods. *Contemp. Clin. trials Commun.* 6, 1–8.

Davies, C. T. M. (1980). Effect of air resistance on the metabolic cost and performance of cycling.
Eur. J. Appl. Physiol. Occup. Physiol. 45, 245–254.

Delp, S. L., Anderson, F. C., Arnold, A. S., Loan, P., Habib, A., John, C. T., et al. (2007). OpenSim:
open-source software to create and analyze dynamic simulations of movement. *IEEE Trans.*
Biomed. Eng. 54, 1940–1950. doi:10.1109/TBME.2007.901024.

Dembia, C. L., Silder, A., Uchida, T. K., Hicks, J. L., and Delp, S. L. (2017). Simulating ideal
assistive devices to reduce the metabolic cost of walking with heavy loads. *PLoS One* 12, 1–25.
doi:10.1371/journal.pone.0180320.

Donelan, J. M., Kram, R., and Kuo, A. D. (2002). Mechanical work for step-to-step transitions is a
major determinant of the metabolic cost of human walking. *J. Exp. Biol.* 205, 3717–3727.

Donelan, J. M., Li, Q., Naing, V., Hoffer, J. A., Weber, D. J., and Kuo, A. D. (2008). Biomechanical
energy harvesting: generating electricity during walking with minimal user effort. *Science (80-.*
). 319, 807–810.

Falisse, A., Serrancolí, G., Dembia, C. L., Gillis, J., Jonkers, I., and De Groote, F. (2019). Rapid
predictive simulations with complex musculoskeletal models suggest that diverse healthy and
pathological human gaits can emerge from similar control strategies. *J. R. Soc. Interface* 16,
20190402. doi:10.1098/rsif.2019.0402.

Fey, N. P., Klute, G. K., and Neptune, R. R. (2012). Optimization of Prosthetic Foot Stiffness to
Reduce Metabolic Cost and Intact Knee Loading During Below-Knee Amputee Walking: A
Theoretical Study. *J. Biomech. Eng.* 134. doi:10.1115/1.4007824.

Finley, J. M., and Bastian, A. J. (2017). Associations Between Foot Placement Asymmetries and
Metabolic Cost of Transport in Hemiparetic Gait. *Neurorehabil. Neural Repair* 31, 168–177.
doi:10.1177/1545968316675428.

Finley, J. M., Long, A., Bastian, A. J., and Torres-Oviedo, G. (2015). Spatial and temporal control
contribute to step length asymmetry during split-belt adaptation and hemiparetic gait.
Neurorehabil. Neural Repair 29, 786–795.

Franz, J. R., Wierzbinski, C. M., and Kram, R. (2012). Metabolic cost of running barefoot versus
shod: is lighter better? *Med. Sci. Sport. Exerc.* 44, 1519–1525.

Galle, S., Malcolm, P., Collins, S. H., and De Clercq, D. (2017). Reducing the metabolic cost of
walking with an ankle exoskeleton: interaction between actuation timing and power. *J.
Neuroeng. Rehabil.* 14, 35. doi:10.1186/s12984-017-0235-0.

Gerus, P., Sartori, M., Besier, T. F., Fregly, B. J., Delp, S. L., Banks, S. A., et al. (2013). Subject-
specific knee joint geometry improves predictions of medial tibiofemoral contact forces. *J.
Biomech.* 46, 2778–2786. doi:10.1016/j.jbiomech.2013.09.005.

Gnehm, P., Reichenbach, S., Altpeter, E., Widmer, H., and Hoppeler, H. (1997). Influence of
different racing positions on metabolic cost in elite cyclists. *Med. Sci. Sports Exerc.* 29, 818.

Hamner, S. R., and Delp, S. L. (2013). Muscle contributions to fore-aft and vertical body mass center
accelerations over a range of running speeds. *J. Biomech.* 46, 780–787.
doi:10.1016/j.jbiomech.2012.11.024.

Hug, F. (2011). Can muscle coordination be precisely studied by surface electromyography? *J.
Electromyogr. Kinesiol.* 21, 1–12. doi:10.1016/J.JELEKIN.2010.08.009.

Koelewijn, A. D., Heinrich, D., and van den Bogert, A. J. (2019). Metabolic cost calculations of gait
using musculoskeletal energy models, a comparison study. *PLoS One* 14, 1–19.
doi:10.1371/journal.pone.0222037.

Lam, Y. Y., and Ravussin, E. (2016). Analysis of energy metabolism in humans: A review of
methodologies. *Mol. Metab.* 5, 1057–1071.

Lloyd, D. G., and Besier, T. F. (2003). An EMG-driven musculoskeletal model to estimate muscle
forces and knee joint moments in vivo. *J. Biomech.* 36, 765–776.

Long III, L. L., and Srinivasan, M. (2013). Walking, running, and resting under time, distance, and
average speed constraints: optimality of walk–run–rest mixtures. *J. R. Soc. Interface* 10,
20120980.

Malcolm, P., Derave, W., Galle, S., and De Clercq, D. (2013). A simple exoskeleton that assists

plantarflexion can reduce the metabolic cost of human walking. *PLoS One* 8, e56137–e56137.
doi:10.1371/journal.pone.0056137.

Margaria, R. (1968). Positive and negative work performances and their efficiencies in human
locomotion. *Int. Zeitschrift fur Angew. Physiol. Einschl. Arbeitsphysiologie* 25, 339–351.
doi:10.1007/BF00699624.

McDaniel, J., Durstine, J. L., Hand, G. A., and Martin, J. C. (2002). Determinants of metabolic cost
during submaximal cycling. *J. Appl. Physiol.* 93, 823–828.

Meyer, A. J., Eskinazi, I., Jackson, J. N., Rao, A. V., Patten, C., and Fregly, B. J. (2016). Muscle
synergies facilitate computational predict ion of subject-specific walking motions. *Front.*
Bioeng. Biotechnol. 4. doi:10.3389/fbioe.2016.00077.

Meyer, A. J., Patten, C., and Fregly, B. J. (2017). Lower extremity EMG-driven modeling of walking
with automated adjustment of musculoskeletal geometry. *PLoS One* 12, e0179698.

Mian, O. S., Thom, J. M., Ardigò, L. P., Narici, M. V, and Minetti, A. E. (2006). Metabolic cost,
mechanical work, and efficiency during walking in young and older men. *Acta Physiol.* 186,
127–139.

Miller, R. H. (2014). A comparison of muscle energy models for simulating human walking in three
dimensions. *J. Biomech.* 47, 1373–1381. doi:10.1016/j.jbiomech.2014.01.049.

Miller, R. H., Brandon, S. C. E., and Deluzio, K. J. (2013). Predicting sagittal plane biomechanics
that minimize the axial knee joint contact force during walking. *J. Biomech. Eng.* 135.

Neptune, R. R., and Van Den Bogert, A. J. (1997). Standard mechanical energy analyses do not
correlate with muscle work in cycling. *J. Biomech.* 31, 239–245.

Ong, C. F., Geijtenbeek, T., Hicks, J. L., and Delp, S. L. (2019). Predicting gait adaptations due to
ankle plantarflexor muscle weakness and contracture using physics-based musculoskeletal
simulations. *PLoS Comput. Biol.* 15, e1006993. doi:10.1371/journal.pcbi.1006993.

Rajagopal, A., Dembia, C. L., DeMers, M. S., Delp, D. D., Hicks, J. L., and Delp, S. L. (2016). Full-
Body Musculoskeletal Model for Muscle-Driven Simulation of Human Gait. *IEEE Trans.*
Biomed. Eng. 63, 2068–2079. doi:10.1109/TBME.2016.2586891.

Reinbolt, J. A., Haftka, R. T., Chmielewski, T. L., and Fregly, B. J. (2007). Are patient-specific joint
and inertial parameters necessary for accurate inverse dynamics analyses of gait? *IEEE Trans.*
Biomed. Eng. 54, 782–793.

Reinbolt, J. A., Haftka, R. T., Chmielewski, T. L., and Fregly, B. J. (2008). A computational
framework to predict post-treatment outcome for gait-related disorders. *Med. Eng. Phys.* 30,
434–443.

Reinbolt, J. A., Schutte, J. F., Fregly, B. J., Koh, B. Il, Haftka, R. T., George, A. D., et al. (2005).
Determination of patient-specific multi-joint kinematic models through two-level optimization.

J. Biomech. 38, 621–626.

Roberts, T. J., Kram, R., Weyand, P. G., and Taylor, C. R. (1998). Energetics of bipedal running. I.
Metabolic cost of generating force. *J. Exp. Biol.* 201, 2745–2751.

Russell Esposito, E., and Miller, R. H. (2018). Maintenance of muscle strength retains a normal
metabolic cost in simulated walking after transtibial limb loss. *PLoS One* 13, e0191310.
doi:10.1371/journal.pone.0191310.

Sartori, M., Reggiani, M., Farina, D., and Lloyd, D. G. (2012). EMG-Driven Forward-Dynamic
Estimation of Muscle Force and Joint Moment about Multiple Degrees of Freedom in the
Human Lower Extremity. *PLoS One* 7. doi:10.1371/journal.pone.0052618.

Sauder, N. R., Meyer, A. J., Allen, J. L., Ting, L. H., Kesar, T. M., and Fregly, B. J. (2019).
Computational Design of FastFES Treatment to Improve Propulsive Force Symmetry During
Post-stroke Gait: A Feasibility Study   . *Front. Neurorobotics*   13, 80.

Seth, A., Hicks, J. L., Uchida, T. K., Habib, A., Dembia, C. L., Dunne, J. J., et al. (2018). OpenSim:
Simulating musculoskeletal dynamics and neuromuscular control to study human and animal
movement. *PLoS Comput. Biol.* 14, e1006223. doi:10.1371/journal.pcbi.1006223.

Shao, Q., Bassett, D. N., Manal, K., and Buchanan, T. S. (2009). An EMG-driven model to estimate
muscle forces and joint moments in stroke patients. *Comput. Biol. Med.* 39, 1083–1088.
doi:10.1016/j.compbiomed.2009.09.002.

Shourijeh, M. S., and Fregly, B. J. (2020). Muscle Synergies Modify Optimization Estimates of Joint
Stiffness During Walking. *J. Biomech. Eng.* 142, 011011. doi:
https://doi.org/10.1115/1.4044310.

Silder, A., Whittington, B., Heiderscheit, B., and Thelen, D. G. (2007). Identification of passive
elastic joint moment-angle relationships in the lower extremity. *J. Biomech.* 40, 2628–2635.
doi:10.1016/j.jbiomech.2006.12.017.

Teunissen, L. P. J., Grabowski, A., and Kram, R. (2007). Effects of independently altering body
weight and body mass on the metabolic cost of running. *J. Exp. Biol.* 210, 4418–4427.

Uchida, T. K., Hicks, J. L., Dembia, C. L., and Delp, S. L. (2016a). Stretching your energetic budget:
How tendon compliance affects the metabolic cost of running. *PLoS One* 11, 1–19.
doi:10.1371/journal.pone.0150378.

Uchida, T. K., Seth, A., Pouya, S., Dembia, C. L., Hicks, J. L., and Delp, S. L. (2016b). Simulating
Ideal Assistive Devices to Reduce the Metabolic Cost of Running. *PLoS One* 11, e0163417–
e0163417. doi:10.1371/journal.pone.0163417.

Umberger, B. R. (2010). Stance and swing phase costs in human walking. *J. R. Soc. Interface* 7,
1329–1340.

Umberger, B. R., Gerritsen, K. G. M., and Martin, P. E. (2003). A Model of Human Muscle Energy

Expenditure. *Comput. Methods Biomech. Biomed. Engin.* 6, 99–111.

van der Woude, L. H. V, Horstman, A., Faas, P., Mechielsen, S., Bafghi, H. A., and de Koning, J. J.
(2008). Power output and metabolic cost of synchronous and asynchronous submaximal and
peak level hand cycling on a motor driven treadmill in able-bodied male subjects. *Med. Eng.*
Phys. 30, 574–580.

Waters, R. L., and Mulroy, S. (1999). The energy expenditure of normal and pathologic gait. *Gait*
Posture 9, 207–231.

9   Supplementary Material

Additional figures can be found in Supplementary Material

10   Figures

449
450

451   **Table 1.** Clinical characteristics of study participants
452

Gender	Age	Paretic Limb	Lower Limb Fugl-Meyer	Self Selected Speed (m/s)
M	79	R	32	0.5
M	62	R	25	0.35

460   **Table 2.** A list of muscle groups from which surface or fine-wire (*) EMG signal was recorded
461

High Functioning Subject	Low Functioning Subject
Adductor Longus	*Adductor Longus*
Gluteus Maximus	Gluteus Maximus
Gluteus Medius	Gluteus Medius
Iliopsoas	*Iliopsoas*
Semimembranosus	Semimembranosus
Biceps Femoris Long Head	Biceps Femoris Long Head
Rectus Femoris	Tensor Fasciae Latae
Vastus Lateralis	Rectus Femoris
Vastus Medialis	Vastus Lateralis
Gastrocnemius Medialis	Vastus Medialis
Tibialis Anterior	Gastrocnemius Lateralis
Tibialis Posterior	Gastrocnemius Medialis
Peroneus Longus	Tibialis Anterior
Soleus	*Tibialis Posterior*
Extensor Digitorum Longus	Peroneus Longus
Flexor Digitorum Longus	Soleus

463 **Table 3.** Resulting p-values obtained from a t-test analysis between the calculated and experimental
464 CoT estimates. Statistically similar slopes are in bold.
465

	Umberger 2003			Umberger 2010			Bhargava 2004		
	SOGen	**SOCal**	**EMGCal**	**SOGen**	**SOCal**	**EMGCal**	**SOGen**	**SOCal**	**EMGCal**
Speed	0.033	**0.12**	**0.23**	0.033	**0.13**	**0.20**	0.0064	0.0089	0.025
Step Position Contribution	**0.056**	**0.22**	**0.52**	**0.30**	**0.21**	**0.40**	0.0062	0.020	**0.20**
Stance Time Differences	**0.26**	**0.30**	**0.39**	**0.24**	**0.29**	**0.34**	**0.22**	**0.25**	**0.34**
Double Support Time Differences	**0.088**	**0.20**	**0.45**	**0.064**	**0.19**	**0.37**	0.014	0.023	**0.22**

467 **Table 4.** Cost of transport estimates for the three metabolic cost models and all three personalization
468 methods across Fugl-Meyer scores for the high functioning subject.
469

	Umberger 2003			Umberger 2010			Bhargava 2004		
	SOGen	**SOCal**	**EMGCal**	**SOGen**	**SOCal**	**EMGCal**	**SOGen**	**SOCal**	**EMGCal**
Max CoT	4.66	6.24	6.58	6.17	7.10	8.24	2.72	2.45	2.35
Min Cot	7.05	10.98	11.35	8.69	11.96	13.14	3.29	3.67	3.50
Mean CoT	5.56	8.13	8.73	7.10	9.04	10.46	2.89	2.89	2.92

470
471
472
473

474 **Table 5.** Cost of transport estimates for the three metabolic cost models and all three personalization
475 methods across Fugl-Meyer scores for the low functioning subject.
476

	Umberger 2003			Umberger 2010			Bhargava 2004		
	SOGen	SOCal	EMGCal	SOGen	SOCal	EMGCal	SOGen	SOCal	EMGCal
Max CoT	4.68	6.74	7.74	5.91	7.55	9.23	2.37	2.62	3.25
Min Cot	7.65	11.63	12.79	9.21	12.57	14.33	3.76	4.11	4.51
Mean CoT	5.98	8.74	9.82	7.38	9.60	11.30	3.03	3.19	3.77

Figure 1. Flowchart of the three different approaches used to obtain estimates of muscle activations (SOGen: red, SOCal: blue, EMGCal: black).

486
487 **Figure 2.** Cost of transport estimates for the three metabolic cost models and for all three
488 personalization methods (black) across speeds. Experimental data reported in Finley and Bastian
489 (Finley and Bastian, 2017) are shown in grey.

Figure 3. Cost of transport estimates for the three metabolic cost models and for all three
personalization methods (black) across step position contribution to step length differences.
Experimental data reported in Finley and Bastian 2017 are shown in grey.

Figure 4. Cost of transport estimates for the three metabolic cost models and for all three personalization methods across stance time differences. Experimental data reported in Finley and Bastian 2017 are shown in grey.

495
496
497

Figure 5. Cost of transport estimates for the three metabolic cost models and for all three personalization methods across double support time differences. Experimental data reported in Finley and Bastian 2017 are shown in grey.

499
500
501

Figure 6. Increase in cost of transport as the specific muscle strength decreases using the SO-Cal modeling combination for the high functioning patient across three speeds (relative to self-selected speed – 100%).