Rapid evolution promotes fluctuation-dependent species coexistence

Masato Yamamichi[1,2*], Andrew D. Letten[1]

[1]School of Biological Sciences, The University of Queensland, St. Lucia, Brisbane, Queensland

4072, Australia, [2]Department of International Health and Medical Anthropology, Institute of

Tropical Medicine, Nagasaki University, Nagasaki 852-8523, Japan

[*]Email: m.yamamichi@uq.edu.au, Phone: +61-7-3365-1864

Short title: Eco-evolutionary relative nonlinearity

Keywords: Adaptive foraging, Armstrong-McGehee mechanism, breeding suppression,

competition, dormancy, eco-evolutionary dynamics, fitness gradient, gleaner-opportunist trade-

off, phenotypic plasticity, relative nonlinearity

ABSTRACT

Recent studies have demonstrated that rapid contemporary evolution can play a significant role in

regulating population dynamics on ecological timescales. Here we identify a previously

unrecognized mode by which rapid evolution can promote species coexistence via temporal

fluctuations and a trade-off between competitive ability and the speed of adaptive evolution. We

show that this interaction between rapid evolution and temporal fluctuations not only increases

the range of coexistence conditions under a gleaner-opportunist trade-off (i.e., low minimum

resource requirement [R^*] vs. high maximum growth rate), but also yields stable coexistence in

the absence of a classical gleaner-opportunist trade-off. Given the propensity for both oscillatory

dynamics and divergent rates of adaptation (including rapid evolution and phenotypic plasticity)

in the real world, we argue that this expansion of fluctuation-dependent coexistence theory

provides an important overlooked solution to the so-called 'paradox of the plankton'.

Introduction

A wealth of empirical evidence has accumulated over the last few decades indicating that rapid contemporary evolution on ecological timescales (Hendry 2016) can be essential for understanding population dynamics (Yoshida *et al.* 2003; Bell 2017). More recently, evidence has begun to emerge that rapid evolution can also be a significant driver of community dynamics amongst competing species, and in particular play an important role in regulating species coexistence (Lankau 2011; Vasseur *et al.* 2011; Mougi 2013; Hiltunen *et al.* 2017; Wittmann & Fukami 2018; Hart *et al.* 2019; Germain *et al.* 2020; van Velzen 2020; Yamamichi *et al.* 2020).

To date, however, rapid evolution has only been shown to promote species coexistence when there is either a trade-off between traits optimal for intraspecific and interspecific competition (Lankau 2011; Vasseur *et al.* 2011; Mougi 2013; Wittmann & Fukami 2018; Yamamichi *et al.* 2020) or fine-tuning of prey defenses and predator foraging efforts (Kondoh 2003; van Velzen 2020). Here we provide a previously unrecognized pathway for rapid evolution to promote coexistence via temporal fluctuations: differences in the rate of adaptation to fluctuating environments. Specifically, using model simulations, we show that rapidly evolving and non-evolving consumers can stably coexist even when they compete for a single resource.

Understanding how species coexist in nature has been a central topic in community ecology over

the last century (Hutchinson 1961; Wilson 1990). G. E. Hutchinson (1961) famously referred to

the problem as "the paradox of the plankton" on account of the seemingly impossible challenge

of explaining the extraordinary diversity of oceanic plankton (Hutchinson 1961). Independent of

emerging interest into rapid evolution, a growing body of coexistence research has suggested that

temporal fluctuations can promote diversity via two broad classes of mechanisms, the temporal

storage effect and relative nonlinearity (Chesson 1994, 2000; Yuan & Chesson 2015; Barabás *et*

al. 2018; Ellner *et al.* 2019). Although greater attention has been given to the temporal storage

effect (Adler *et al.* 2006; Angert *et al.* 2009; Letten *et al.* 2018; Hallett *et al.* 2019; Zepeda &

Martorell 2019), recent studies have shown that relative nonlinearity can be more important than

previously thought for promoting coexistence (Letten *et al.* 2018; Hallett *et al.* 2019; Zepeda &

Martorell 2019). In this paper, we introduce rapid evolution as a previously neglected driver of

relative nonlinearity.

Relative nonlinearity promotes coexistence when one species benefits from fluctuations in the

intensity of competition (fluctuation-assisted) while the other species dominates in the absence of

fluctuations (fluctuation-impeded). In addition to their responses to fluctuations, for the two

competing species to coexist, they need different impacts on fluctuations. If the fluctuation-

assisted species reduces the amplitude of fluctuations (by "consuming" the variance in the

intensity of competition (Levins 1979)) while the fluctuation-impeded species increases it, each

species produces a favorable environment for the competitor, resulting in negative frequency-

dependence in community dynamics (Hsu *et al.* 1978; Armstrong & McGehee 1980) (Fig. 1a-b).

As the name suggests, two species may coexist via relative nonlinearity when consumer

population growth responses to a limiting resource differ in their degree of nonlinearity (Fig. 1c-

d) (Hsu *et al.* 1978; Armstrong & McGehee 1980; Abrams & Holt 2002; Abrams *et al.* 2003;

Xiao & Fussmann 2013). The differential responses to, and impacts on, fluctuations are most

conspicuous when competing consumers exhibit intermittent cycles (Fig. 1d), where the linear,

competitively inferior species at equilibrium (with larger R^*), increases in abundance when the

system oscillates but simultaneously dampens the consumer-resource cycles induced by the

competitively superior species (with smaller R^*). In contrast, the nonlinear, competitively

superior species generates fluctuations but increases in abundance when the system is stable (Fig.

1d). The species with the lowest R^* value would outcompete its competitor under a stable

equilibrium, but the linear species benefits from fluctuating environments in competition due to

nonlinear averaging of the low R^* species' per capita growth response (Jensen's inequality). The

situation in Fig. 1c is called a gleaner-opportunist trade-off, where species 1 has the lower R^*

whereas species 2 has the higher maximum growth rate (Grover 1990). Because this criterion is

sometimes perceived as being highly restrictive, relative nonlinearity has typically been assumed

to play a minor role in real systems (Adler *et al.* 2006; Angert *et al.* 2009; Letten *et al.* 2018;

Hallett *et al.* 2019; Zepeda & Martorell 2019). By demonstrating how rapid evolution can relax

these restrictions and significantly broaden opportunities for coexistence even without the

classical gleaner-opportunist trade-off, we argue that relative nonlinearity may operate much

more widely in nature than previously thought.

METHODS

The basis for the mechanism presented here is that rapid evolution can enable populations to

adapt to fluctuating environments (response) and to stabilize temporal fluctuations (impact)

(Cortez & Patel 2017). Thus, even when two species with fixed functional responses cannot

coexist, they may have different responses to, and impacts on, temporal fluctuations, facilitating

coexistence. We consider a model of two consumers with saturating consumer functional

responses competing for a single logistically growing resource (Rosenzweig & MacArthur 1963;

Hsu *et al.* 1978; Xiao & Fussmann 2013). In our model, the dynamics of the two consumers, N_1

and N_2, competing for a single resource, R, with quantitative traits, x_1 and x_2 is represented by

$$\frac{dR}{dt} = R\left[r\left(1-\frac{R}{K}\right) - \sum_{j=1}^{2} \frac{a_j(x_j)N_j}{1+h_jR} \right],$$

95   $$\frac{dN_i}{dt} = N_i\left[\frac{c_i a_i(x_i)R}{1+h_iR} - d_i(x_i) \right], \quad (i=1,2), \tag{1}$$

$$\frac{dx_i}{dt} = V_i \frac{\partial}{\partial x_i}\left(\frac{1}{N_i}\frac{dN_i}{dt} \right), \quad (i=1,2),$$

where r is the intrinsic growth rate, K is the carrying capacity, a_i is the parameter that increases

the consumption rate as a whole (including the slope and maximum value), b_i is the parameter

that reduces the maximum consumption rate, c_i is the conversion efficiency, d_i is the consumer

mortality, and V_i is additive genetic variance of the trait x_i ($i = 1, 2$).

We assume a trade-off between parameters for the consumption rate, a_i, and mortality rate, d_i,

such that increasing the consumption rate increase mortality rate in an accelerating manner ($a_i(x_i)$

$= \alpha_i x_i$, $d_i(x_i) = \delta_{0i} + \delta_{1i}x_i + \delta_{2i}x_i^2$). This occurs due to, for example, a foraging-predation risk trade-

off (Verdolin 2006) where predators have Holling type III functional responses. In addition, we

assume the trait changes along the fitness gradient (Abrams 2001; Cortez & Patel 2017). Thus,

trait dynamics can be represented by

$$\frac{dx_i}{dt} = V_i\left(\frac{c_i \alpha_i R}{1 + h_i R} - \delta_{1i} - 2\delta_{2i}x_i \right), \tag{2}$$

and the optimal trait at a certain resource level is obtained by solving $dx_i/dt = 0$, and it is given by

$x_i = (c_i\alpha_i R/(1 + b_i R) - \delta_{1i})/(2\delta_{2i})$. There is stabilizing (instead of disruptive) selection toward the

optimal trait as $\partial(dx_i/dt)/\partial x_i = -2V_i\delta_{2i} < 0$. This indicates that the evolving consumer invests more

in resource uptake when resources are abundant, whereas it tries to reduce its mortality rate when

resources are scarce. Furthermore, by nondimensionalization, we can assume that $r = K = c_i = 1$

without loss of generality (Appendix S1).

As the resource exhibits logistic growth and consumers exploit the resource with a Holling type

II functional response, the system corresponds to the classical Rosenzweig-MacArthur model

when there is a single non-evolving consumer species (Rosenzweig & MacArthur 1963). In the

absence of fluctuations, the consumer with the smaller resource requirement (i.e., smaller R^*)

reduces the resource level to the point ($R_i^* = d_i/(a_i c_i - b_i d_i)$, $i = 1, 2$) where its competitor with a

larger R^* cannot persist and is excluded. On the other hand, in the presence of endogenous limit

cycles (e.g., with larger K values) the two consumers may coexist via relative nonlinearity. Given

Holling type II (concave) functional responses, population growth rates of the two species are

lowered by fluctuations, but the species with lower R^* is more impeded by fluctuations in the

intensity in competition, thereby promoting coexistence (Hsu *et al.* 1978; Xiao & Fussmann

2013).

Results

When we assume that a competitively superior species (with smaller R^* at equilibrium) does not

evolve, whereas an evolving, competitively inferior (larger R^*) species with the optimum trait

either grows faster than the competitively superior species when the resource is abundant (Fig.

2a) or scarce (Fig. 2b), coexistence is possible. Notably, our parameter settings do not satisfy

necessary conditions for coexistence via the classical gleaner-opportunist trade-off identified in

previous studies (Appendix S1, Fig. S1) (Hsu *et al.* 1978; Xiao & Fussmann 2013) for any fixed

trait values. Thus, when the speed of evolution in the species that is competitively inferior at

equilibrium is slow, competitive exclusion ensues (Fig. S2). However, when evolution is faster,

coexistence is possible (Fig. 2). This occurs because relative to the non-evolving species, the

rapidly evolving species benefits from resource fluctuations, but at the same time rapid evolution

dampens the consumer-resource cycles (Appendix S2, Fig. S3) induced by the non-evolving

superior competitor, as can be clearly seen when the system cycles intermittently (Fig. 2f).

Therefore, rapid evolution works as a compensation mechanism for the species that would

otherwise be competitively inferior in a non-fluctuating system.

The underlying mechanism for coexistence (*sensu* Chesson 2000) can be understood by looking

at the very-rapid-evolution limit. When evolution is much faster than ecological processes, the

evolving species is always able to attain the optimal trait value that maximizes per-capita growth

rate for a given resource concentration. The effect of very rapid evolution is to change the

effective functional responses of the evolving species from concave down (gray curves in Fig. 2a-

b) to concave up (especially when the limiting resource is scarce: orange curves in Fig. 2a-b) and

it is this change that allows coexistence by enabling the evolving species to benefit from

fluctuating environments in competition through nonlinear averaging. Although the realized per-

capita growth rates when evolution is not much faster than ecology are different from those based

on the optimal trait values due to the time lag in responses to changing environments (Fig. 2c-d,

Fig. S4), coexistence is still possible because of relative nonlinearity.

It is important to note that rapid evolution not only broadens the parameter conditions that allow

for coexistence via a so-called gleaner-opportunist trade-off (i.e., species 1 has a lower R^*

whereas species 2 has a higher maximum growth rate (Grover 1990): Fig. 2a, c, and e), but it also

yields stable coexistence under conditions previously thought to be incompatible with

coexistence (Fig. 3). More specifically, coexistence is still possible even in the absence of a

classical gleaner-opportunist trade-off, i.e., there is no trade-off between R^* and maximum growth

rate, and the species with the smaller R^* and the higher maximum growth rate (species 1 in Fig.

2b, d, and f) is competitively impeded by fluctuations. Figure 3b shows that coexistence is not

possible when genetic variance is zero, but as the genetic variance of the inferior species

increases, the range of parameter values compatible with coexistence sharply increases. As such,

this relaxation of the perceived restrictions characterizing classical relative nonlinearity can

explain the coexistence of a much greater functional diversity of competing organisms than

would be possible in the absence of rapid evolution.

Discussion

Relative nonlinearity promotes species coexistence via fluctuations when (1) growth rates are

relatively nonlinear functions of the limiting factor in competition (in this case, a shared resource)

and (2) each species, when abundant, drives the amplitude of fluctuations in directions that favor

their competitors (Fig. 1) (Yuan & Chesson 2015). Here we have shown that rapid evolution can

satisfy the above two conditions under conditions previously thought to be incompatible with

coexistence. Even without the classical gleaner-opportunist trade-off (Grover 1990), we show

that coexistence is possible via relative nonlinearity as the evolving consumer species benefits

from resource fluctuations in competition but stabilizes the consumer-resource cycles (Fig. 2-3).

Although our focus here is a system with rapidly evolving consumers competing for a single

resource, a trade-off between competitive ability at equilibrium and the speed of adaptation may

promote coexistence via relative nonlinearity under various other scenarios. For example,

mechanisms of rapid adaptive trait changes include rapid evolution and phenotypic plasticity, and

some organisms plastically suppress foraging activity and reproduction when resources are scarce

(breeding suppression or dormancy), thereby stabilizing population dynamics (Nakazawa *et al.*

2009). This type of adaptive behavioral responses may also promote coexistence via relative

nonlinearity (Appendix S4, Fig. S6). Recently, Tan *et al.* (2020) showed that a competitively

inferior (larger R^*) consumer species with resource-dependent dormancy can coexist with a

superior species in consumer-resource cycles. Although the authors discussed their findings in the

context of game theory, our results indicate that they can also be explained as an outcome of

relative nonlinearity. In addition, phenotypic plasticity in defense traits against predation can

stabilize population dynamics (Yamamichi *et al.* 2019) and may also promote coexistence via

relative nonlinearity. When the limiting factor in competition is predation pressure instead of

resource abundance (i.e., apparent competition), species with higher predation tolerance (P^*) will

persist (Holt 1977). However, coexistence is possible when species with lower predation

tolerance stabilizes predator-prey cycles by inducible defenses (Appendix S5, Fig. S7). This

provides an explanation for previous simulation results demonstrating coexistence of three prey

types (asexually reproducing clones) with a shared predator and a shared resource (Yamamichi *et*

al. 2011). According to classical R^* and P^* theory, only two species (or asexual clones) can

coexist with a single predator and a single resource (Holt *et al.* 1994), but a third prey species

with inducible defense can coexist with two non-plastic (defended and undefended) prey by

relative nonlinearity. The destabilizing non-plastic prey have a competitive advantage when the

dynamics are stable, while the stabilizing plastic prey have an advantage under fluctuations.

While our model assumptions are quite general, further theoretical studies are necessary to

understand the generality and robustness of this novel mode of coexistence in ecosystems

including, for example, the shape of the trade-off between consumption and mortality (Appendix

S3, Fig. S5). Analyzing stochastic models will be particularly helpful to understand how low

genetic variance (and therefore slower rates of evolution) in small populations impedes

coexistence. It will be also valuable to conduct empirical studies that examine the effects of rapid

evolution on species coexistence via relative nonlinearity. In addition to meta-analyses on the

trade-off between foraging and mortality (Verdolin 2006) and the trade-off between competitive

ability at equilibrium and the speed of adaptation, competition experiments will be valuable for

deepening our understandings. Previous studies on the consequences of rapid evolution have

typically used a single clone for competition experiments *after* long-term experimental evolution

(Hart *et al.* 2019). However, it is rapid evolution itself during competition experiments that can

promote coexistence via the mechanism outlined here, and a single clone cannot produce these

kinds of dynamics. Although it is possible to manipulate the amount of genetic variation by

changing the number of clones (Yoshida *et al.* 2003; Becks *et al.* 2010), we note that species with

two clones will not coexist with another species via relative nonlinearity (mathematically, it is

equivalent to three consumers competing for a single resource). In addition, one should note that

rapid evolution does not always stabilize temporal fluctuations (Becks *et al.* 2010; Cortez & Patel

2017), in which case rapid evolution may result in priority effects via positive frequency-

dependence in community dynamics (Ke & Letten 2018). Therefore, using phenotypically plastic

species may be more feasible for showing coexistence driven by rapid adaptive trait dynamics via

relative nonlinearity because phenotypic plasticity tends to strongly stabilize population

dynamics, and the speed of adaptation will not be as slow as genetic evolution in small

populations (Yamamichi *et al.* 2011; Yamamichi *et al.* 2019).

Conclusion

With increasing evidence for rapid contemporary evolution on ecological timescales (Hendry

2016), an interest in developing a broader theory of species coexistence that incorporates rapid

evolution and eco-evolutionary feedbacks has grown (Tachikawa 2008; Kremer & Klausmeier

2013; Wittmann & Fukami 2018; Hart *et al.* 2019; van Velzen 2020; Yamamichi *et al.* 2020). By

allowing for rapid evolution in mechanistic resource competition models, we have identified a

new solution to the paradox of the plankton, deriving from relative nonlinearity of competition.

This mechanism may be important in competition between sexual and asexual species or species

with different mutation rates. Alongside recent empirical work demonstrating relative

nonlinearity across diverse systems (Letten *et al.* 2018; Hallett *et al.* 2019; Zepeda & Martorell

2019), we hope that our findings will help to motivate a reappraisal of the perceived importance

of relative nonlinearity in nature.

ACKNOWLEDGEMENTS

We thank S. P. Ellner, T. Fukami, and three anonymous reviewers for their helpful comments.

MY thanks N. Shinohara for stimulating discussions during the initial stages of this project. MY

was supported by the Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for

Scientific Research (KAKENHI) 18H02509 and 19K16223.

CONFLICT OF INTEREST

The author declares no competing financial interests.

REFERENCES

1.
Abrams, P.A. (2001). Modelling the adaptive dynamics of traits involved in inter- and intraspecific
interactions: An assessment of three methods. *Ecol. Lett.*, 4, 166-175.
2.
Abrams, P.A., Brassil, C.E. & Holt, R.D. (2003). Dynamics and responses to mortality rates of
competing predators undergoing predator–prey cycles. *Theor. Popul. Biol.*, 64, 163-176.
3.
Abrams, P.A. & Holt, R.D. (2002). The impact of consumer–resource cycles on the coexistence of
competing consumers. *Theor. Popul. Biol.*, 62, 281-295.
4.
Adler, P.B., HilleRisLambers, J., Kyriakidis, P.C., Guan, Q. & Levine, J.M. (2006). Climate
variability has a stabilizing effect on the coexistence of prairie grasses. *Proc. Natl. Acad.*
Sci. U.S.A., 103, 12793-12798.

5.

Angert, A.L., Huxman, T.E., Chesson, P. & Venable, D.L. (2009). Functional tradeoffs determine

species coexistence via the storage effect. *Proc. Natl. Acad. Sci. U.S.A.*, 106, 11641-11645.

6.

Armstrong, R.A. & McGehee, R. (1980). Competitive exclusion. *Am. Nat.*, 115, 151-170.

7.

Barabás, G., D'Andrea, R. & Stump, S.M. (2018). Chesson's coexistence theory. *Ecol. Monogr.*, 88,

277-303.

8.

Becks, L., Ellner, S.P., Jones, L.E. & Hairston, N.G., Jr. (2010). Reduction of adaptive genetic

diversity radically alters eco-evolutionary community dynamics. *Ecol. Lett.*, 13, 989-997.

9.

Bell, G. (2017). Evolutionary rescue. *Annu. Rev. Ecol. Evol. Syst.*, 48, 605-627.

10.

Chase, J.M. & Leibold, M.A. (2003). *Ecological Niches: Linking Classical and Contemporary*

Approaches. University of Chicago Press, Chicago, IL.

11.

Chesson, P. (1994). Multispecies competition in variable environments. *Theor. Popul. Biol.*, 45,

227-276.

12.

Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.*, 31,

343-366.

13.

Cortez, M.H. & Patel, S. (2017). The effects of predator evolution and genetic variation on

predator–prey population-level dynamics. *Bull. Math. Biol.*, 79, 1510-1538.

14.

Ellner, S.P., Snyder, R.E., Adler, P.B. & Hooker, G. (2019). An expanded modern coexistence

theory for empirical applications. *Ecol. Lett.*, 22, 3-18.

15.

Germain, R.M., Srivastava, D. & Angert, A.L. (2020). Evolution of an inferior competitor increases

resistance to biological invasion. *Nat. Ecol. Evol.*, 4, 419-425.

16.

Grover, J.P. (1990). Resource competition in a variable environment: phytoplankton growing
according to Monod's model. *Am. Nat.*, 136, 771-789.

17.

Hallett, L.M., Shoemaker, L.G., White, C.T. & Suding, K.N. (2019). Rainfall variability maintains
grass-forb species coexistence. *Ecol. Lett.*, 22, 1658-1667.

18.

Hart, S.P., Turcotte, M.M. & Levine, J.M. (2019). Effects of rapid evolution on species coexistence.
Proc. Natl. Acad. Sci. U.S.A., 116, 2112-2117.

19.

Hendry, A.P. (2016). *Eco-evolutionary Dynamics*. Princeton University Press, Princeton, NJ.

20.

Hiltunen, T., Kaitala, V., Laakso, J. & Becks, L. (2017). Evolutionary contribution to coexistence
of competitors in microbial food webs. *Proc. R. Soc. B*, 284, 20170415.

21.

Holt, R.D. (1977). Predation, apparent competition, and the structure of prey communities. *Theor.*
Popul. Biol., 12, 197-229.

22.

Holt, R.D., Grover, J. & Tilman, D. (1994). Simple rules for interspecific dominance in systems
with exploitative and apparent competition. *Am. Nat.*, 144, 741-771.

23.

Hsu, S.B., Hubbell, S.P. & Waltman, P. (1978). Competing predators. *SIAM Journal on Applied*
Mathematics, 35, 617-625.

24.

Hutchinson, G.E. (1961). The paradox of the plankton. *Am. Nat.*, 95, 137-145.

25.

Ke, P.-J. & Letten, A.D. (2018). Coexistence theory and the frequency-dependence of priority
effects. *Nat. Ecol. Evol.*, 2, 1691-1695.

26.

Kondoh, M. (2003). Foraging adaptation and the relationship between food-web complexity and
stability. *Science*, 299, 1388-1391.

27.

Kremer, C.T. & Klausmeier, C.A. (2013). Coexistence in a variable environment: eco-evolutionary

perspectives. *J. Theor. Biol.*, 339, 14-25.

28.

Lankau, R.A. (2011). Rapid evolutionary change and the coexistence of species. *Annu. Rev. Ecol.*
Evol. Syst., 42, 335-354.

29.

Letten, A.D., Dhami, M.K., Ke, P.-J. & Fukami, T. (2018). Species coexistence through
simultaneous fluctuation-dependent mechanisms. *Proc. Natl. Acad. Sci. U.S.A.*, 115, 6745-
6750.

30.

Levins, R. (1979). Coexistence in a variable environment. *Am. Nat.*, 114, 765-783.

31.

Mougi, A. (2013). Allelopathic adaptation can cause competitive coexistence. *Theor. Ecol.*, 6, 165-
171.

32.

Nakazawa, T., Ohgushi, T. & Yamamura, N. (2009). Resource-dependent reproductive adjustment
and the stability of consumer-resource dynamics. *Popul. Ecol.*, 51, 105-113.

33.

Rosenzweig, M.L. & MacArthur, R.H. (1963). Graphical representation and stability conditions of
predator-prey interactions. *Am. Nat.*, 97, 209-223.

34.

Tachikawa, M. (2008). Fluctuation induces evolutionary branching in a mathematical model of
ecosystems. *PLOS ONE*, 3, e3925.

35.

Tan, Z.X., Koh, J.M., Koonin, E.V. & Cheong, K.H. (2020). Predator dormancy is a stable adaptive
strategy due to Parrondo's paradox. *Advanced Science*, 7, 1901559.

36.

van Velzen, E. (2020). Predator coexistence through emergent fitness equalization. *Ecology*, 101,
e02995.

37.

Vasseur, D.A., Amarasekare, P., Rudolf, V.H.W. & Levine, J.M. (2011). Eco-evolutionary dynamics
enable coexistence via neighbor-dependent selection. *Am. Nat.*, 178, E96-E109.

38.

Verdolin, J.L. (2006). Meta-analysis of foraging and predation risk trade-offs in terrestrial systems.

Behav. Ecol. Sociobiol., 60, 457-464.

39.

Wilson, J.B. (1990). Mechanisms of species coexistence: Twelve explanations for Hutchinson's

'Paradox of the Plankton': Evidence from New Zealand plant communities. *N. Z. J. Ecol.*,

13, 17-42.

40.

Wittmann, M.J. & Fukami, T. (2018). Eco-evolutionary buffering: Rapid evolution facilitates

regional species coexistence despite local priority effects. *Am. Nat.*, 191, E171-E184.

41.

Xiao, X. & Fussmann, G.F. (2013). Armstrong–McGehee mechanism revisited: Competitive

exclusion and coexistence of nonlinear consumers. *J. Theor. Biol.*, 339, 26-35.

42.

Yamamichi, M., Klauschies, T., Miner, B.E. & van Velzen, E. (2019). Modelling inducible defences

in predator–prey interactions: assumptions and dynamical consequences of three distinct

approaches. *Ecol. Lett.*, 22, 390-404.

43.

Yamamichi, M., Kyogoku, D., Iritani, R., Kobayashi, K., Takahashi, Y., Tsurui-Sato, K. *et al.*

(2020). Intraspecific adaptation load: a mechanism for species coexistence. *Trends Ecol.*

Evol., 35, 897-907.

44.

Yamamichi, M., Yoshida, T. & Sasaki, A. (2011). Comparing the effects of rapid evolution and

phenotypic plasticity on predator-prey dynamics. *Am. Nat.*, 178, 287-304.

45.

Yoshida, T., Jones, L.E., Ellner, S.P., Fussmann, G.F. & Hairston, N.G., Jr. (2003). Rapid evolution

drives ecological dynamics in a predator-prey system. *Nature*, 424, 303-306.

46.

Yuan, C. & Chesson, P. (2015). The relative importance of relative nonlinearity and the storage

effect in the lottery model. *Theor. Popul. Biol.*, 105, 39-52.

47.

Zepeda, V. & Martorell, C. (2019). Fluctuation-independent niche differentiation and relative non-

linearity drive coexistence in a species-rich grassland. *Ecology*, 100, e02726.

Figure 1 | Species coexistence by relative nonlinearity. a, Relative nonlinearity promotes

coexistence with negative frequency-dependence (NFD) when one species stabilizes population

cycles but benefits from fluctuating environments in competition (fluctuation-assisted), while the

other species destabilizes dynamics but benefits from stable environments in competition

(fluctuation-impeded). Hence, we need to consider two aspects of the ecological niche: responses

and impacts (the Grinnellian and Eltonian niches, respectively: note that with a feedback cycle,

the ideas of response and effect are complicated since the two terms rely heavily on perspective)

(Chase & Leibold 2003). Priority effects due to positive frequency-dependence (PFD) can arise

when the fluctuation-impeded species stabilizes fluctuations while the fluctuation-assisted species

drive fluctuations (Ke & Letten 2018). **b**, The concept of relative nonlinearity can also be

expressed in terms of zero net growth isoclines (ZNGI) (Chase & Leibold 2003). The fluctuation-

assisted (fluctuation-impeded) species has higher growth rates when the variance in the

competitive factor (e.g., resource) is high (low), but when abundant reduces (increases) the

variance as shown by the impact vector. **c-d**, Coexistence of species with linear and saturating

functional responses (Abrams *et al.* 2003). Species 1 with a saturating functional response (blue

line) has a smaller R^* and causes limit cycles with the resource (grey line), which is

disadvantageous due to nonlinear averaging. Species 2 with a linear functional response (the

dashed orange line) has a larger R^* and stabilizes dynamics. Parameter values are $a_1 = a_2 = 5$, $b_1 =$

10, $b_2 = 0$, $d_1 = 0.33$, and $d_2 = 1.7$.

Figure 2 | Rapid evolution of a competitively inferior (high R^*) species promotes coexistence

by relative nonlinearity. a-b, The per-capita growth rate of species 2 with the optimal trait

(orange dashed line) is larger than that of species 1 (solid blue line) when the resource is

abundant (**a**) or scarce (**b**). When the trait value is fixed (gray lines), species 2 is always

competitively inferior (Fig. S2). **c-d**, The realized per-capita growth rate of species 2 (i.e., the

trajectory over time: orange solid line) is different from that with the optimal trait (orange dashed

line) because of a time lag in adaptive evolution. **e-f**, Rapid evolution of species 2 results in

coexistence with synchronous (**e**) or intermittent (**f**) cycles. Under intermittent cycles, the density

of species 1 (blue solid line) increases when cycle amplitudes are small, whereas the density of

species 2 (orange dashed line) increases when cycle amplitudes are large. The trait value of

species 2 is depicted in black, and resource abundance in gray. Parameter values are $a_1 = 20$, $\alpha_2 =$

84, $b_1 = b_2 = 10$, $d_1 = 0.8$, $\delta_{02} = 1$, $\delta_{12} = 0.2$, $\delta_{22} = 4$, $V_1 = 0$, and $V_2 = 0.1$ in **a**, **c**, and **e**, and $\delta_{02} =$

0.25, $\delta_{22} = 12$, and $V_2 = 0.08$ in **b**, **d**, and **f**.

Figure 3 | Effect of rapid evolution on parameter space yielding stable coexistence. a-b,

Phase diagrams for the genetic variance, V_2, and parameter that reduces the maximum

consumption rate, b_2, of species 2 where the per-capita growth rate of species 2 with the optimal

trait is either larger than that of species 1 when the resource is abundant (**a**) or scarce (**b**). Dashed

grey line indicates the situations where both species have the identical value of b ($b_1 = b_2$), where

species 1 is competitively superior (lower R^*) and therefore excludes species 2 in the absence of

genetic variance (Fig. 2, S2). At the limit of zero genetic variance, the coexistence region in **a**

corresponds to classical relative nonlinearity. With increasing genetic variance, the parameter

space corresponding to coexistence expands. Solid grey line indicates the point at which species 2

becomes the superior competitor and therefore genetic variance (in species 2) no longer promotes

coexistence with smaller b_2. Black dots indicate fixed parameter values in Figure 2. Black dotted

curves indicate Hopf bifurcation in the system with species 2 and resource: the system is locally

stable below the curves (with smaller b_2 and larger V_2).