Rapid evolution promotes fluctuation-dependent species coexistence

Last revision: August 6, 2020

Masato Yamamichi[1,2,3*], Andrew D. Letten[1]

[1]School of Biological Sciences, The University of Queensland, St. Lucia, Brisbane, Queensland

4072, Australia, [2]Department of International Health and Medical Anthropology, Institute of

Tropical Medicine, Nagasaki University, Nagasaki 852-8523, Japan, [3]Department of Ecology and

Evolutionary Biology, Cornell University, Ithaca, NY 14853, U.S.A.

[*]Email: m.yamamichi@uq.edu.au, Phone: +61-7-3365-1864

ABSTRACT

Recent studies have demonstrated that rapid contemporary evolution can play a significant role in

regulating population dynamics on ecological timescales. Here we identify a previously

unrecognized mechanism by which rapid evolution promotes species coexistence via temporal

fluctuations and a trade-off between competitive ability and the speed of adaptive evolution. This

is a special case of relative nonlinearity in coexistence theory, and given the propensity for both

oscillatory dynamics and divergent rates of adaptation in the real world, we propose that this

mechanism may operate widely in nature.

A wealth of empirical evidence has accumulated over the last few decades indicating that rapid

contemporary evolution on ecological timescales[1] can be essential for understanding population

dynamics[2,3]. More recently, evidence has begun to emerge that rapid evolution can also be a

significant driver of community dynamics amongst competing species, and in particular play an

important role in regulating species coexistence[4-12]. Rapid evolution can promote species

coexistence when there is a trade-off between traits optimal for intraspecific and interspecific

competition[4-6,8,12] or fine-tuning of prey defenses and predator foraging efforts[10,13]. Here we

provide a previously unrecognized pathway for rapid evolution to promote coexistence via

temporal fluctuations: differences in the rate of adaptation to fluctuating environments.

Specifically, using model simulations, we show that rapidly evolving and non-evolving

consumers can stably coexist even when they compete for a single resource.

How species coexist in nature has been a central topic in community ecology over the last

century[14,15]. Independent of emerging interest into rapid evolution, a growing body of coexistence

research has suggested that temporal fluctuations can promote diversity via two broad classes of

mechanisms, the temporal storage effect and relative nonlinearity[16-20]. Although greater attention

has been given to the temporal storage effect[21-25], recent studies have shown that relative

nonlinearity can be more important than previously thought for promoting coexistence[23-25]. In

this paper, we also focus on relative nonlinearity, but we introduce rapid evolution as a previously

neglected driver of relative nonlinearity.

Relative nonlinearity promotes coexistence when one species benefits from fluctuations in the

intensity of competition (an opportunist or risk-seeking strategist) while the other species

dominates in the absence of fluctuations (a gleaner or risk-avoiding strategist). In addition to their

responses to fluctuations, for the two competing species to coexist, they need different impacts on

fluctuations. If the opportunist species reduces the amplitude of fluctuations (by "consuming" the

variance in the intensity of competition[26]) while the gleaner species increases it, each species

produces a favorable environment for the competitor, resulting in negative frequency-

dependence[27,28] (Fig. 1a-b).

As the name suggests, two species coexist via relative nonlinearity when consumer functional

responses to a limiting resource differ in their degree of nonlinearity (Fig. 1c-d)[27-31]. The

responses to and impacts on fluctuations are clearly seen in the intermittent cycles (Fig. 1d),

where the linear, competitively inferior species increases in abundance when the system oscillates

but simultaneously dampens the consumer-resource cycles induced by the competitively superior

species. In contrast, the nonlinear, competitively superior species increases when the system is

stable but causes fluctuations (Fig. 1d). Because this criterion is perceived as being highly

restrictive, relative nonlinearity has typically been assumed to play a minor role[21-25], but we

propose that it may be more important than previously realized due to rapid evolution.

The basis for the mechanism presented here is that rapid evolution can enable populations to

adapt to fluctuating environments (response) and to stabilize temporal fluctuations (impact)[32].

Thus, even when two species have the same curvature in their functional responses, they may

have different responses and impacts of temporal fluctuations, facilitating coexistence. We

consider a model of two consumers sharing a single resource with logistic resource growth and

saturating consumer functional responses[27,31,33]. We assume a trade-off between the attack rate

and mortality rate of consumers, and the evolving consumer invests more in resource uptake

when resources are abundant, whereas it invests more in reduced mortality when resources are

scarce. We further assume that a competitively superior species (with the lower resource

requirement) does not evolve, whereas an evolving, competitively inferior species grows faster

than the competitively superior species when the resource is abundant (Fig. 2a) or scarce (Fig.

2b). Our parameter settings do not satisfy necessary conditions for coexistence identified in

previous studies (Appendix S1, Fig. S1)[27,31] for any fixed trait values. Thus, when the speed of

evolution in the competitively inferior species is slow, competitive exclusion ensues (Fig. S2).

However, when evolution is faster, coexistence is possible (Fig. 2). This occurs because the

rapidly evolving species benefits from resource fluctuations, but at the same time rapid evolution

dampens the consumer-resource cycles (Appendix S2, Fig. S3) induced by the non-evolving

superior competitor as shown in the intermittent cycles (Fig. 2f). Therefore, rapid evolution is

working as a compensation mechanism for the competitively inferior species.

The mechanism for coexistence can be understood by looking at the very-rapid-evolution limit.

When evolution is much faster than ecological processes, the evolving species always has the

optimal trait values that maximize per-capita growth rate for a given resource concentration. The

effect of very rapid evolution is to change the effective functional responses of the evolving

species from concave down (gray curves in Fig. 2a-b) to concave up (especially when the

limiting resource is scarce: orange curves in Fig. 2a-b) and it is this change that allows

coexistence. Although the realized per-capita growth rates when evolution is not much faster than

ecology are different from those based on the optimal trait values due to the time lag in responses

to changing environments (Fig. 2c-d), coexistence is still possible because of relative

nonlinearity.

Mechanisms of rapid adaptative trait changes include rapid evolution and phenotypic plasticity.

We showed that both types of rapid adaptation can promote consumer coexistence in exploitative

competition (adaptive foraging and dormancy induction/reproduction suppression: Fig. 2,

Appendix S3-4, Fig. S4-5)[34] as well as prey coexistence in apparent competition (defense

evolution and inducible defense: Appendix S5, Fig. S6)[35]. Recently, Tan et al. showed that a

competitively inferior consumer species with phenotypic plasticity can coexist with a superior

species in synchronous cycles[34]. Although the authors discussed their findings in the context of

game theory, our results indicate that they can also be explained as an outcome of relative

nonlinearity. Furthermore, our work provides an explanation for previous simulation results

demonstrating coexistence of three prey types (asexually reproducing clones) with a shared

predator and a shared resource[35]. According to classical R^* and P^* theory, only two species (or

asexual clones) can coexist with a single predator and a single resource[36], but a third prey species

with inducible defense can coexist with the two non-plastic (defended and undefended) prey by

relative nonlinearity.

It will be valuable to conduct empirical studies that examine the effects of rapid evolution on

species coexistence via relative nonlinearity. Previous studies on the consequences of rapid

evolution have typically used a single clone for competition experiments *after* long-term

experimental evolution[9]. However, it is rapid evolution itself during competition experiments that

can promote coexistence via the mechanism outlined here, and a single clone cannot produce

these kind of dynamics. Although it is difficult to control the presence/absence of rapid evolution,

it may be possible by changing the amount of genetic variation in populations to see dynamical

interactions[2,37]. While our model assumptions are quite general, one should note that rapid

evolution does not always stabilize temporal fluctuations[32,37], in which case rapid evolution may

result in priority effects via positive frequency-dependence[38].

With increasing evidence for rapid contemporary evolution on ecological timescales[1], there has

been an interest in developing a broader theory of species coexistence that incorporates rapid

evolution[8-10,12]. By allowing for rapid evolution in mechanistic resource competition models, we

have identified a new mechanism by which relative nonlinearity of competition promotes

coexistence. This mechanism may be important in competition between sexual and asexual

species or species with different mutation rates. Alongside recent empirical work demonstrating

relative nonlinearity across diverse systems[23-25], we hope that our findings will motivate a

reappraisal of the perceived importance of relative nonlinearity in nature.

METHODS

Consider the dynamics of two consumers, N_1 and N_2, competing for a single resource, R, with

quantitative traits, x_1 and x_2:

$$\frac{dR}{dt} = R\left[r\left(1 - \frac{R}{K}\right) - \sum_{j=1}^{2} \frac{a_j(x_j)N_j}{1 + h_j R} \right],$$

$$\frac{dN_i}{dt} = N_i\left[\frac{c_i a_i(x_i)R}{1 + h_i R} - d_i(x_i) \right], \quad (i = 1,2), \qquad (1)$$

$$\frac{dx_i}{dt} = V_i(x_j)\frac{\partial}{\partial x_i}\left(\frac{1}{N_i}\frac{dN_i}{dt} \right), \quad (i = 1,2),$$

where r is the intrinsic growth rate, K is the carrying capacity, a_i is the attacking rate, h_i is the

handling time, c_i is the conversion efficiency, d_i is the consumer mortality, and V_i is additive

genetic variance of the trait x_i ($i = 1, 2$). We assume that increasing the trait value results in a

larger attacking rate and mortality due to a trade-off ($a_i(x_i) = \alpha_i x_i$, $d_i(x_i) = \delta_{0i} + \delta_{1i}x_i + \delta_{2i}x_i^2$), the

trait changes along the fitness gradient[32,39], and the function $V_i(x_i)$ bounds the trait value, $V_i(x_i) =$

$v_i(x_{i\max} - x_i)(x_i - x_{i\min})$, where the minimum and maximum values of x_i are $x_{i\min}$ and $x_{i\max}$,

respectively.

As the resource exhibits logistic growth and consumers exploit the resource with a Holling type

II functional response, the system is the classical Rosenzweig-MacArthur model when there is a

single consumer species[33]. When the system does not show fluctuations, the consumer with the

smaller resource requirement (i.e., smaller R^*) reduces the resource level to the point ($R_i^* =$

$d_i/(a_ic_i - h_id_i)$, $i = 1, 2$) where its competitor with a larger R^* cannot persist and is excluded. On

the other hand, in the presence of endogenous limit cycles (e.g., with larger K values) the two

consumers may coexist via relative nonlinearity[27,31]. The mechanism with rapid evolution further

broadens the parameter space that allows coexistence mediated by temporal fluctuations.

ACKNOWLEDGEMENTS

We thank S. P. Ellner and T. Fukami for their helpful comments. MY thanks N. Shinohara for

stimulating discussions during the initial stages of this project. MY was supported by the Japan

Society for the Promotion of Science (JSPS) Grant-in-Aid for Scientific Research (KAKENHI)

18H02509 and 19K16223.

CONFLICT OF INTEREST

The author declares no competing financial interests.

REFERENCES

1   Hendry, A. P. *Eco-evolutionary Dynamics*. (Princeton University Press, 2016).
2   Yoshida, T., Jones, L. E., Ellner, S. P., Fussmann, G. F. & Hairston, N. G., Jr. Rapid

evolution drives ecological dynamics in a predator-prey system. *Nature* **424**, 303-306, doi:10.1038/nature01767 (2003).

Bell, G. Evolutionary rescue. *Annu. Rev. Ecol. Evol. Syst.* **48**, 605-627, doi:10.1146/annurev-ecolsys-110316-023011 (2017).

Lankau, R. A. Rapid evolutionary change and the coexistence of species. *Annu. Rev. Ecol. Evol. Syst.* **42**, 335-354, doi:10.1146/annurev-ecolsys-102710-145100 (2011).

Vasseur, D. A., Amarasekare, P., Rudolf, V. H. W. & Levine, J. M. Eco-evolutionary dynamics enable coexistence via neighbor-dependent selection. *Am. Nat.* **178**, E96-E109, doi:10.1086/662161 (2011).

Mougi, A. Allelopathic adaptation can cause competitive coexistence. *Theor. Ecol.* **6**, 165-171, doi:10.1007/s12080-012-0168-y (2013).

Hiltunen, T., Kaitala, V., Laakso, J. & Becks, L. Evolutionary contribution to coexistence of competitors in microbial food webs. *Proc. R. Soc. B* **284**, 20170415, doi:10.1098/rspb.2017.0415 (2017).

Wittmann, M. J. & Fukami, T. Eco-evolutionary buffering: Rapid evolution facilitates regional species coexistence despite local priority effects. *Am. Nat.* **191**, E171-E184, doi:10.1086/697187 (2018).

Hart, S. P., Turcotte, M. M. & Levine, J. M. Effects of rapid evolution on species coexistence. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 2112-2117, doi:10.1073/pnas.1816298116 (2019).

van Velzen, E. Predator coexistence through emergent fitness equalization. *Ecology* **101**, e02995, doi:10.1002/ecy.2995 (2020).

Germain, R. M., Srivastava, D. & Angert, A. L. Evolution of an inferior competitor increases resistance to biological invasion. *Nat. Ecol. Evol.* **4**, 419-425, doi:10.1038/s41559-020-1105-x (2020).

Yamamichi, M. *et al.* Intraspecific adaptation load: a mechanism for species coexistence. *Trends Ecol. Evol.* **35**, doi:10.1016/j.tree.2020.05.011 (in press).

Kondoh, M. Foraging adaptation and the relationship between food-web complexity and stability. *Science* **299**, 1388-1391 (2003).

Hutchinson, G. E. The paradox of the plankton. *Am. Nat.* **95**, 137-145 (1961).

Wilson, J. B. Mechanisms of species coexistence: Twelve explanations for Hutchinson's 'Paradox of the Plankton': Evidence from New Zealand plant communities. *N. Z. J. Ecol.*

13, 17-42 (1990).

16   Chesson, P. Multispecies competition in variable environments. *Theor. Popul. Biol.* **45**, 227-
276, doi:10.1006/tpbi.1994.1013 (1994).

17   Chesson, P. Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.* **31**,
343-366, doi:10.1146/annurev.ecolsys.31.1.343 (2000).

18   Barabás, G., D'Andrea, R. & Stump, S. M. Chesson's coexistence theory. *Ecol. Monogr.* **88**,
277-303, doi:10.1002/ecm.1302 (2018).

19   Ellner, S. P., Snyder, R. E., Adler, P. B. & Hooker, G. An expanded modern coexistence
theory for empirical applications. *Ecol. Lett.* **22**, 3-18, doi:10.1111/ele.13159 (2019).

20   Yuan, C. & Chesson, P. The relative importance of relative nonlinearity and the storage
effect in the lottery model. *Theor. Popul. Biol.* **105**, 39-52, doi:10.1016/j.tpb.2015.08.001
(2015).

21   Adler, P. B., HilleRisLambers, J., Kyriakidis, P. C., Guan, Q. & Levine, J. M. Climate
variability has a stabilizing effect on the coexistence of prairie grasses. *Proc. Natl. Acad.*
Sci. U.S.A. **103**, 12793-12798, doi:10.1073/pnas.0600599103 (2006).

22   Angert, A. L., Huxman, T. E., Chesson, P. & Venable, D. L. Functional tradeoffs determine
species coexistence via the storage effect. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 11641-11645
(2009).

23   Letten, A. D., Dhami, M. K., Ke, P.-J. & Fukami, T. Species coexistence through
simultaneous fluctuation-dependent mechanisms. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 6745-
6750, doi:10.1073/pnas.1801846115 (2018).

24   Zepeda, V. & Martorell, C. Fluctuation-independent niche differentiation and relative non-
linearity drive coexistence in a species-rich grassland. *Ecology* **100**, e02726,
doi:10.1002/ecy.2726 (2019).

25   Hallett, L. M., Shoemaker, L. G., White, C. T. & Suding, K. N. Rainfall variability
maintains grass-forb species coexistence. *Ecol. Lett.* **22**, 1658-1667, doi:10.1111/ele.13341
(2019).

26   Levins, R. Coexistence in a variable environment. *Am. Nat.* **114**, 765-783,
doi:10.1086/283527 (1979).

27   Hsu, S. B., Hubbell, S. P. & Waltman, P. Competing predators. *SIAM Journal on Applied*
Mathematics **35**, 617-625, doi:10.1137/0135051 (1978).

28   Armstrong, R. A. & McGehee, R. Competitive exclusion. *Am. Nat.* **115**, 151-170 (1980).

29 Abrams, P. A. & Holt, R. D. The impact of consumer–resource cycles on the coexistence of
competing consumers. *Theor. Popul. Biol.* **62**, 281-295, doi:10.1006/tpbi.2002.1614 (2002).

30 Abrams, P. A., Brassil, C. E. & Holt, R. D. Dynamics and responses to mortality rates of
competing predators undergoing predator–prey cycles. *Theor. Popul. Biol.* **64**, 163-176,
doi:10.1016/S0040-5809(03)00067-4 (2003).

31 Xiao, X. & Fussmann, G. F. Armstrong–McGehee mechanism revisited: Competitive
exclusion and coexistence of nonlinear consumers. *J. Theor. Biol.* **339**, 26-35,
doi:10.1016/j.jtbi.2013.05.025 (2013).

32 Cortez, M. H. & Patel, S. The effects of predator evolution and genetic variation on
predator–prey population-level dynamics. *Bull. Math. Biol.* **79**, 1510-1538,
doi:10.1007/s11538-017-0297-y (2017).

33 Rosenzweig, M. L. & MacArthur, R. H. Graphical representation and stability conditions
of predator-prey interactions. *Am. Nat.* **97**, 209-223 (1963).

34 Tan, Z. X., Koh, J. M., Koonin, E. V. & Cheong, K. H. Predator dormancy is a stable
adaptive strategy due to Parrondo's paradox. *Advanced Science* **7**, 1901559,
doi:10.1002/advs.201901559 (2020).

35 Yamamichi, M., Yoshida, T. & Sasaki, A. Comparing the effects of rapid evolution and
phenotypic plasticity on predator-prey dynamics. *Am. Nat.* **178**, 287-304,
doi:10.1086/661241 (2011).

36 Holt, R. D., Grover, J. & Tilman, D. Simple rules for interspecific dominance in systems
with exploitative and apparent competition. *Am. Nat.* **144**, 741-771, doi:10.1086/285705
(1994).

37 Becks, L., Ellner, S. P., Jones, L. E. & Hairston, N. G., Jr. Reduction of adaptive genetic
diversity radically alters eco-evolutionary community dynamics. *Ecol. Lett.* **13**, 989-997,
doi:10.1111/j.1461-0248.2010.01490.x (2010).

38 Ke, P.-J. & Letten, A. D. Coexistence theory and the frequency-dependence of priority
effects. *Nat. Ecol. Evol.* **2**, 1691-1695, doi:10.1038/s41559-018-0679-z (2018).

39 Abrams, P. A. Modelling the adaptive dynamics of traits involved in inter- and intraspecific
interactions: An assessment of three methods. *Ecol. Lett.* **4**, 166-175, doi:10.1046/j.1461-
0248.2001.00199.x (2001).

40 Chase, J. M. & Leibold, M. A. *Ecological Niches: Linking Classical and Contemporary*
Approaches. (University of Chicago Press, 2003).

Figure 1 | Species coexistence by relative nonlinearity. a, Relative nonlinearity promotes

coexistence with negative frequency-dependence (NFD) when one species stabilizes population

cycles but benefits from fluctuating environments (risk-seeking), while the other species

destabilizes dynamics but benefits from stable environments (risk-avoiding). Hence, we need to

consider two aspects of the ecological niche: responses and impacts (the Grinnellian and Eltonian

niches, respectively)[40]. Priority effects due to positive frequency-dependence (PFD) can arise

when the risk-avoiding species stabilizes fluctuations while the risk-seeking species drive

fluctuations[38]. **b**, The concept of relative nonlinearity can also be expressed in terms of zero net

growth isoclines (ZNGI)[40]. The risk-seeking (risk-avoiding) species has higher growth rates when

the variance in the competitive factor (e.g., resource) is high (low), but when abundant reduces

(increases) the variance as shown by the impact vector. **c-d**, Coexistence of species with linear

and saturating functional responses[30]. Species 1 with a saturating functional response (blue line)

has a smaller R^*, and causes limit cycles with the resource (grey line), which is disadvantageous

due to nonlinear averaging. Species 2 with a linear functional response (the dashed orange line)

has a larger R^* and stabilizes dynamics. Parameter values are $a_1 = a_2 = 5$, $h_1 = 10$, $h_2 = 0$, $d_1 =$

0.33, and $d_2 = 1.7$.

Figure 2 | Rapid evolution of a competitively inferior species promotes coexistence by

relative nonlinearity. a-b, The per-capita growth rate of species 2 having the optimal trait

(orange dashed line) is larger than that of species 1 (solid blue line) when the resource is

abundant (**a**) or scarce (**b**). When the trait value is fixed (gray lines), species 2 is always

competitively inferior (Fig. S2). **c-d**, The realized per-capita growth rate of species 2 (orange

solid line) is different from that with the optimal trait (orange dashed line) because of a time lag

in adaptive evolution. **e-f**, Rapid evolution of species 2 results in coexistence with synchronous

(**e**) or intermittent (**f**) cycles. Under intermittent cycles, the density of species 1 (blue solid line)

increases when cycle amplitudes are small, whereas the density of species 2 (orange dashed line)

increases under largely amplitude fluctuating. The trait value of species 2 is depicted in black,

and resource abundance in gray. Parameter values are $a_1 = 20$, $\alpha_2 = 84$, $h_1 = h_2 = 10$, $d_1 = 0.8$, δ_{02}

$= 1$, $\delta_{12} = 0.24$, $\delta_{22} = 4$, $\beta_{min} = 0$, $\beta_{max} = 1$, $v_1 = 0$, and $v_2 = 0.5$ in a, c, and e, and $\delta_{02} = 0.3$ and δ_{22}

$= 11$ in b, d, and f.