Genomic Signature of Sexual Reproduction in the Bdelloid Rotifer *Macrotrachella quadricornifera*

Veronika N. Laine[*], Timothy Sackton[†] and Matthew Meselson[‡]

[*]Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland.Department of Animal Ecology, [†]Informatics Group, Faculty of Arts and Sciences, Harvard University, Cambridge, MA 02138; [‡]Department of Molecular and Cellular Biology, Harvard University, Cambridge, MA 02138. email: *msmeselson@gmail.com*

Data Availability

The sequencing reads (Illumina and Nanopore) and assemblies generated in this study are available from NCBI BioProject PRJNA773114.

Difference matrices, phylograms, alignments (tic plots), and raw data are available at *https://github.com/tsackton/rotifer-outcrossing*

.

Running title: Sexual Reproduction in *M. quadricornifera*

Keywords Allele sharing, bdelloid rotifers, life history, *Macrotrachella quadricornifera*,

population structure, Red Queen, sexual reproduction

Corresponding Author**:** Matthew Meselson, Department of Molecular and Cellular

Biology, Harvard University, 16 Divinity Avenue Cambridge, MA 02138. email:

msmeselson@gmail.com

Abstract:

ABSTRACT Bdelloid rotifers, common freshwater invertebrates of ancient origin

and worldwide distribution have long been thought to be entirely asexual, being

the principal exception to the view that in eukaryotes the loss of sex leads to

early extinction. That bdelloids are facultatively sexual is shown by a study of

allele sharing within a group of closely related bdelloids of the species

***Macrotrachella quadricornifera*, making it likely that sexual reproduction is**

essential for long-term success in all eukaryotes.

38   **INTRODUCTION**

Nearly all eukaryotes reproduce sexually, either constitutively or facultatively. That

sexual reproduction may be essential for evolution was first suggested by Weismann

(Weismann 1887) who, noting that, in comparison with asexuals, "in sexual reproduction

twice as many individuals are required to produce any number of descendants"

proposed that the compensating advantage of sexual reproduction is as a "source of

individual variability, furnishing material for the operation of natural selection". More

specific hypotheses for the evolutionary benefit of sexual reproduction put forward since

then include (*i*) bringing favorable mutations in different individuals together in the same

lineage (Fisher 1930; Muller 1932) and reconstituting lines less loaded with detrimental

mutations when, without sexual reproduction, recurring mutation and the continuing

stochastic loss of less-loaded lines might drive the population to extinction (Muller

1964), and, more generally, weakening linkage, allowing selection to operate more

nearly independently on linked loci (Hill and Robertson 1966); (*ii*) producing genotypes

resistant to co-evolving biological antagonists (Hamilton 1980; Lively 2010); and (*iii*)

purging synergistically-acting deleterious mutations from effectively infinite populations

(Kondrashov 1988). An apparent challenge to all hypotheses for the evolutionary

advantage of sexual reproduction is posed by a few groups of eukaryotes that have

been designated "ancient asexuals" (Judson and Normark 1996; Schon *et al.* 2009). Of

these, the most extensively studied are the rotifers of Class Bdelloidia.

59   **BDELLOID ROTIFERS**

First described nearly 350 years ago (van Leewenhoeck 1677, 1702), bdelloid rotifers

are minute freshwater invertebrates commonly found in lakes, ponds and streams and

ephemerally aquatic habitats such as temporary pools and the water films on lichens

and mosses (Figure 1). For review see: (Mark Welch *et al.* 2009). Characterized by their

ciliated head and bilateral ovaries, bdelloids are classified in 4 families, 19 genera and

some 500 morphospecies. The bdelloid radiation began tens of millions of years ago, as

shown by the synonymous site difference between families and by the presence of

bdelloid remains in ancient amber. Although typically only several tenths of a millimeter

in size and containing only *ca.* 1,000 nuclei, mostly in syncytial tissues, bdelloids have

ganglia, muscles, digestive, excretory, reproductive and secretory systems;

photosensitive and tactile sensory organs; and structures for crawling, feeding and

swimming. Bdelloids are degenerate tetraploids, descended from an ancient tetraploid

ancestor (Mark Welch *et al.* 2008; Hur *et al.* 2009; Flot *et al.* 2013; Nowell *et al.* 2018).

The only observed means of bdelloid reproduction is from eggs produced in well-

differentiated ovaries, with no reduction in chromosome number (Hsu 1956a; b). A few

days after deposition a young bdelloid emerges and a few days later commences egg-

laying, producing up to 32 eggs over a period of up to a few weeks during which there is

little death, after which the death rate increases more or less exponentially (Meadow

and Barrows Jr. 1971; Ricci 1983). Depending on species and conditions, the time from

egg deposition to death is about a month (Ricci 1983; Ricci and Fascio 1995). Bdelloids

are eutelic, with no cell division after eclosion except in the germ line.

Bdelloids are extremophiles, being able to survive prolonged desiccation, some

species more than others, as well as starvation and extremes of temperature, and to

resume reproduction upon restoration of favorable conditions, highly unusual abilities

probably apomorphic to the Class (Ricci 1998; Ricci and Caprioli 2005; Ricci and

Perletti 2006; Nowell *et al.* 2018). Bdelloids have a highly effective system of anti-

oxidant protection, as manifested by their extreme resistance to ionizing radiation and to

IR-induced protein carbonylation (Gladyshev and Meselson 2008; Krisko *et al.* 2012),

apparently an adaptation to avoid the oxidative damage caused by desiccation, as known in other systems (França *et al.* 2007; Fredrickson *et al.* 2008).

Although bdelloids have been systematically studied ever since adequate microscopes became available (Ehrenberg 1838; Hudson and Gosse 1886) there is no confirmed observation of males. It has been estimated that contemporary rotifer workers have examined some 500,000 bdelloids from a variety of habitats and from laboratory culture without ever having seen males or hermaphrodites (Birky 2010). The only report to the contrary, in a treatise otherwise devoted to observations of males and sexual periods of the facultatively sexual rotifers of Class Monogononta, is a hesitant account of having twice seen a single male among many bdelloids of species *Rotaria rotatoria* "present in almost incredible numbers" beneath the ice of a frozen lake in Denmark in November 1923 (Wesenberg-Lund 1930). Sampling conducted there in January 2019 found few bdelloids and no males (Martin Sorensen, personal communication) nor was there any ice, the winter of that year having been among the warmest on record.

Despite the failure to document the existence of males, it may be that bdelloids reproduce sexually only rarely and under conditions not adequately investigated—a possibility made less implausible by estimates that outcrossing of *Saccharomyces cerevisiae* in the field may occur as seldom as once in 25-50,000 generations (Ruderfer *et al.* 2006; Magwene *et al.* 2011), owing to the repression of meiosis which, however, can be relieved in the laboratory by growth in specific media.

In the following, we review observations once interpreted as evidence for bdelloid asexuality but now known to have other explanations. We also summarize recent findings strongly suggestive of bdelloid sex. We then present a study of allele sharing in

111   the bdelloid *Macrotrachella quadricornifera* with results in exact accord with

112   expectations for facultative sexual reproduction and not explicable by horizontal gene

113   transfer (HGT) or parasexuality. We also discuss the relation between bdelloid life

114   history and population structure and implications for how bdelloid males and mating

115   might be discovered.

MATERIALS AND METHODS

Sample collection and 10x sequencing

M. quadrifornifera isolates MA, MM and CR are from a group of 29 individuals

morphologically identified as belonging to Family Philodinidae collected from ground

moss in northeast United States in the autumn of 2011 (Signorovitch *et al.* 2015). Six of

the 29, including MA, MM and CR, each collected at a site distant from the others,

belong to the same mitochondrial clade (Lasek-Nesselquist 2012). Cultures of MA, MM

and CR were twice established from single eggs, fed *E. coli* and maintained in 0.24u

Millipore-filtered spring water at 20 ^{o}C in 100X20 mm plastic Petri dishes with

continuous gentle rotation. Washed rotifers were shipped under dry ice in February

2016 to HudsonAlpha (Huntsville AL) for DNA extraction, library preparation, 10x

Illumina sequencing at 56x coverage and provision to us of fastq files.

Nanopore sequencing

Washed flash-frozen rotifers of isolate MA were digested for 17 h at 58 ^{o}C in 100 mM

EDTA, 50 mM tris pH 9.0, 1% sodium sarcosyl, 1 mg/ml freshly dissolved proteinase K.

DNA was isolated with a Qiagen MagAttract HMW DNA Kit and its size distribution

analyzed with an Agilent Technologies 4200 TapeStation. DNA purity was verified by

determining 260/280 and 260/230 ratios with a Nanodrop ND-1000 and its

concentration was determined with a Qubit 3.0 fluorometer. DNA was prepared for

sequencing with an Oxford Nanopore Ligation Sequencing Kit 1D without shearing and

1.1 µg (48µl) of the ligated DNA was loaded into the DNA repair end-end preparation

step. Flow cells were prepared following the protocol from the same kit. DNA libraries

were quantified with the Qubit and 420 ng of DNA was loaded for each sequencing run.

Base calling for Nanopore reads was done with Albacore 2.3.4 and the results

summarized with Nanoplot 1.20.0 (de Coster *et al.* 2018). Reads longer than 10kb were

selected with fastp 0.19.5 (Chen *et al.* 2018) and aligned to the NCBI-nt database with

Blastn (Altschul *et al.* 1990) using an e-value cutoff of 1e-25, removing reads with a best

hit to a non-Animalia sequence, leaving 25,640 reads of length 10-92 kb. Statistics for

the Nanopore reads are given in Table S1.

147   Assembly of 10x reads

Scaffolds assembled from the 10x reads of each of the three isolates were obtained

with Supernova 2.1.1 (Weisenfeld *et al.* 2017) using default parameters. Genome size,

needed as an input parameter for Supernova, was estimated with Jellyfish 2.2.5

(Marçais and Kingsford 2011), based on the distribution of k-mers. Scaffolds were

aligned to the NCBI nt database with Blastn with settings as above and non-Animalia

scaffolds were removed. The mean assembly size was 328 Mb, range 352-371.

Assembly statistics are presented in Table S2. Phased sequences (megabubbles) were

then obtained from the 10x scaffolds with Supernova mkoutput with style=megabubbles.

Alignment of 10x megabubbles to Nanopore reads

In order to select and align groups of homologous megabubbles from 2 or all 3 of the

isolates, we aligned the Nanopore reads longer than 10 kb that did not have a

significant Blast hit to non-Animalia sequences to the megabubbles from the 10x

assemblies with Minimap 2 2.15-r905 (Li 2018). From the resulting alignments longer

than 8kb, those in which megabubbles of at least two of the three isolates aligned to the

same Nanopore read were chosen for analysis with Samtools faidx (Li *et al.* 2009).

These alignments of 10x megabubbles, comprising groups of either 6 or 4 homologous

sequences, were then realigned among themselves with Clustal Omega 1.2.3 (Sievers

et al. 2011), and trimmed with Gblocks 0.91b (Talavera and Castresana 2007) to

remove indels and alignment disruptions caused by repeats of unequal length,

processes that neither add nor rearrange sequences. When two or more alignments

overlapped, only the longest was retained. Lastly, 1 kb was removed from the ends of

each alignment using EMBOSS seqret (Madeira *et al.* 2019) and alignments shorter

than 1 kb were discarded. Pairwise SNP differences between homologs in each

alignment were obtained with snp-dists 0.6.3 (*https://github.com/tseemann/snp-dists*).

The workflow is depicted in Figure S1.

In five alignments MM or CR differed by more than 10% from both homologs of MA

and were rejected. This may occur when, owing to a deletion in MM or CR occurring

since their divergence from MA, there is no MM or CR sequence homologous to the

Nanopore read, leaving only the homeologous sequence to align with it. This left 1,117

non-overlapping genomic regions for analysis, none of which contain homeologous

sequences (Figure S2). That trimming neither adds nor rearranges sequences is

confirmed by manual inspection of 5 randomly chosen regions of MA-MM sharing

(regions 10, 104,142, 179, 262, total length 62,172 bp) in untrimmed MA-MM-CR

alignments revealing only a single departure from perfect alignment, a one base

deletion in a homopolymer run of As.

The sequence accuracy of the 10x assemblies was determined by comparison with

published sequences of the four regions of MA, MA, and CR of total length 19.37 kb

sequenced by Signorovitch *et al.* 2015, revealing a difference of 31 substitutions overall,

or a 99.84% match (Table S3). Comparison of mitochondrial sequences in the 10x

assemblies identified by blast searches against published mitochondrial sequences of

MA, MM and CR (Lasek-Nesselquist 2012) revealed a perfect or near-perfect match for

each isolate.

Test for contamination

It might be asked if, despite precautions taken against it, contamination of MM DNA with

MA DNA or the reverse in the DNA sent to HudsonAlpha for sequencing or occurring

there, has mimicked allele sharing. Although it is most unlikely that contamination could

be so massive and of the particular frequency required to mimic the MA-MM sharing we

observe in half of the 622 MA-MM alignments, a test was conducted to rule out the

possibility. All 10x Illumina reads sequenced from MA, MM, and CR were aligned to

each of the 1,177 alignments. Using each haplotype in turn as a reference, bwa mem

(Li and Durbin 2009) and GATK were used to produce variant calls (McKenna *et al.*

2010), implemented in a Snakemake (Molder *et al.* 2021) pipeline developed by the

Harvard Informatics group

(*https://github.com/harvardinformatics/shortRead_mapping_variantCalling*). The variant

calls were then processed with vcftools (Danecek *et al.* 2011) to generate counts of

reads at each variable nucleotide position supporting each allele. The fraction of

Illumina reads supporting the alternate allele (the nucleotide that differs from the

reference haplotype) shows a characteristic tri-modal pattern for all three isolates, with

peaks at 0 (reference homozygote), 0.5 (heterozygote), and 1 (alternate homozygote),

as shown for MA and MM in Figure S3. Such a pattern is expected for true variable

positions in a diploid. Contamination would instead produce a pattern in which

heterozygous positions are supported by a fraction of reads that departs from ½, and is

determined by the proportion of the total DNA that was from a contaminating source, for

which no evidence is seen in any of the three assemblies.

PREVIOUS STUDIES

Heterozygosity

In sexuals, heterozygosity caused by mutation is limited by haploid drift. The finding of

much greater synonymous difference between gene copies in bdelloids than in

monogononts was therefore initially interpreted as evidence for asexuality (Mark Welch

and Meselson 2000, 2001). Continued investigation, however, showed that bdelloids

are degenerate tetraploids, and that the highly diverged gene copies are homeologs,

not homologs, with many genes present in only one or the other pair of homologs (Mark

Welch *et al.* 2008; Hur *et al.* 2009). Bdelloid silent-site heterozygosity, the difference

between homologs, lies within the range known for sexuals, providing no evidence for

asexuality. Moreover, in asexual *Daphnia pulex* and *Saccharomyces cerevisiae* the

frequency with which a nucleotide site is covered by a tract of homozygosity, as may

result from germline crossing-over at the four-strand stage of mitosis or from certain

processes of DNA damage repair, is much greater than the frequency of nucleotide

substitution (Omilian et al. 2006; Xu et al. 2011; St. Charles and Petes 2013; Flynn et al.

2017). In sexuals, heterozygosity lost by such processes may be regained by

outcrossing. But if bdelloids are ancient asexuals and if loss of heterozygosity were

more frequent than substitution, the absence of outcrossing would be manifested as

unusually *low* heterozygosity, the opposite of what had been thought (Magwene *et al.*

2011; Hartfield *et al.* 2018). A wide range of heterozygosity values could therefore be

consistent with either sexual or asexual reproduction.

In a finite population, computer simulations demonstrate that small populations are

driven to extinction by a Muller's ratchet-like process of element accumulation, but that

large populations can be cured of vertically transmitted TEs, even with excision rates

well below transposition rates.

Paucity of retrotransposons

Sexual reproduction allows vertically transmitted deleterious transposable elements to

proliferate in populations, whereas under some theoretical models, the loss of sex may

eventually free a population of such elements or drive it to extinction (Hickey 1982;

Dolgin and Charlesworth 2006). As a test for asexuality, bdelloids, monogonont rotifers

and sexually-reproducing animals of 23 other animal phyla were examined for genomic

sequences coding for reverse transcriptases of LINE-like retrotransposons. These were

found to be abundant in all the sexually-reproducing taxa but were not detected in

bdelloids (Arkhipova and Meselson 2000, 2005a; b). Nevertheless, although bdelloids

are nearly devoid of LINE-like retrotransposons, later work showed that they are not

entirely absent (Gladyshev *et al.* 2007; Gladyshev and Arkhipova 2010) and that

bdelloids have particularly effective retrotransposon silencing systems (Rodriguez and

Arkhipova 2016). The paucity of LINE-like retrotransposons is therefore non-evidentiary

as regards bdelloid sexuality.

Genome structure

A draft genome sequence of the bdelloid, *Adineta vaga*, with numerous breaks in the

colinearity of homologous regions and individual scaffolds containing genes in direct or

palindromic repeats but no copy elsewhere in the genome was initially taken as

evidence that bdelloids lack homologous chromosome pairs and had therefore evolved

ameiotically (Flot *et al.* 2013). But subsequent genomic sequencing of three other

bdelloid species, including *Adineta ricciae*, a close relative of *A. vaga*, found that the

unusual genomic features that had been interpreted as evidence for ameiotic evolution

are largely absent, suggesting that their apparent presence in *A. vaga* resulted from

mis-assembly (Nowell *et al.* 2018) as later shown to be the case by the demonstration in

A. vaga of homologous chromosome pairs (Simion *et al.* 2020).

Allele sharing

A finding of two individuals closely related with respect to a given genomic region but

more distantly related with respect to its homolog, a form of phylogenetic

noncongruence known as allele sharing, would mean that recently in their ancestry the

region had undergone some form of genetic exchange between individuals. A striking

example of allele sharing among isolates MA, MM and CR was found by (Signorovitch

et al. 2015, 2016) in each of the four genomic regions examined, 2.4 to 9.7 kb in length.

At each region, MA identically shared a homolog with MM while the other homolog of

MA was identical to a homolog of CR in two regions and nearly so in the other two. That

these observations were evidence for sexual reproduction was disputed and attributed

instead to HGT on the basis of observations in *A. vaga* (Debortoli *et al.* 2016; Flot *et al.*

2018). Subsequent analysis, however, showed that what had been interpreted as HGT

could be explained as the result of cross-contamination among isolates (Wilson *et al.*

2018), leaving no agreed evidence for homologous HGT in bdelloids.

Meiosis-associated genes

A survey of the genomes of four bdelloid species, including *A. vaga*, belonging to two

bdelloid families for 11 genes associated with meiosis found all but one, *red1*, to be

present in each species (Nowell *et al.* 2018). But neither was *red1* found in *Drosophila*

melanogaster, a sexual species known to lack it. Although five of these genes had not

been found in the draft assembly of the *A. vaga* genome (Flot *et al.* 2013) their

subsequent detection and conservation in this and the three other bdelloid species

tested suggests that bdelloids engage in meiosis, although it cannot be excluded that

they are retained instead for other functions.

Population Genetics

A recent population genetics survey of 11 individuals of the bdelloid rotifer A. vaga

collected at diverse sites in the wild found a number of features consistent with sexual

reproduction, including genotype proportions in Hardy-Weinberg equilibrium, indicating

some form of genetic exchange between individuals. Recombination either between or

within individuals is indicated by a decay of linkage disequilibrium with increasing

physical distance along scaffolds and by signatures of reciprocal recombination

(Vakhrusheva *et al.* 2020). Phylogenetic analysis of certain genomic segments in a

group of three closely-related individuals revealed incongruence possibly resulting from

sexual reproduction between their ancestors and members of a distinctly different

population. Although the authors caution that neither sexual reproduction nor HGT by

themselves provide a simple explanation for their data, their observations nevertheless

provide strong evidence for recombination and some form of genetic exchange.

307   **RESULTS**

Alignments

As described in METHODS, we obtained alignments of phased sequences from 1,177

non-overlapping genomic regions of *M. quadricornifera* isolates MA, MM and CR. Of

these, 331 alignments are with all three isolates, 291 with MA and MM, 110 with MA

and CR and 445 with MM and CR, having a mean length of 12,490 bp (range 2,051 -

32,937 bp) and altogether covering 14.7 Mb, approximately 4% of the *ca.* 360 Mb

genome (Table 2). Matrices giving pair-wise differences between homologous

sequences, phylograms and plots of the spatial distribution of differences between

homologs ("tic" plots) for four representative regions are given in Table 1 and, for all

1,177 regions, in Supplemental Material Tables 4-6.

Allele sharing

Half of the MA-MM and MA-MM-CR alignments, 315 of 622, comprise a discrete class

in which a homolog of MA is identical to a homolog of MM (Table 2; Fig. 2B). The

frequencies of identical MA-MM sharing in the MA-MM and MA-MM-CR alignments

considered separately are 0.519 and 0.495, respectively or 0.506 overall (S.E. = 0.02).

MA and MM also share identical homologs with CR, but in a much smaller proportion of

the alignments (Fig. 2A, C). CR shares identical homologs with MA in 12 of 441

alignments, and with MM in 18 of 776 alignments, or 2.7% and 2.3% respectively. In

alignments without identical sharing the differences between the homolog of MA most

similar to a homolog of MM form a broad distribution with a mean of 0.96 SNPs per 100

bp (S.E. = 0.61), Fig. 2B. Most or all of the regions identically shared between MA and

MM must be considerably longer than the alignments in which we find them, as shown

in a plot of the frequency of identical MA-MM sharing against alignment length in

consecutive intervals each comprising 76-79 alignments (Fig. 3). The frequency of

identical sharing is not significantly different from 50 percent in even the longest

alignments (18-33 kb).

Genealogy

MA and MM identically share a homolog in a discrete class amounting to half of the 622

MA-MM and MA-MM-CR alignments while their other homologs, within and between MA

and MM, are substantially diverged. MA and MM therefore identically share ¼ of their

genomes as segments at least the length of the alignments, a degree of relationship

expected for grandchild and grandparent, half siblings, aunt or uncle and nephew or

niece from a large panmictic sexually-reproducing population. The near equality of MA-

CR and MM-CR sharing frequencies indicates that CR is equidistant from MA and MM

and therefore that MA and MM are not grandparent and grandchild or aunt/uncle-

nephew/niece but rather half siblings or double first cousins.

Inspection of tic plots for the triple alignments reveals a few with interior regions

of MA-MM identity covering much but not all of the alignment (Supplemental Material

Table 4). In these regions there is substantial divergence from CR, showing that such

identity is not the result of extreme conservation but instead reflects more remote

relationships between MA and MM in addition to their relation as half siblings or double

first cousins. Similarly, alignments in which CR is identical to MA or MM over much but

not all the alignment are likely to reflect remote relationships between CR and MA and

between CR and MM. In general, more distant relations will be manifested as shorter

regions of identity by descent, owing to the crossing-over that occurs at each meiosis.

For individuals related as half-sibs or double first cousins the regions of identity by

descent from their common grandparents, assuming one cross-over per meiosis in each

arm of the 10 chromosomes of *M. quadricornifera*, will average several Mbp in length,

far longer than our longest alignments, consistent with the observation that over the

region examined the frequency of identically shared regions does not fall off with length

(Fig. 3).

Homozygosity

In each of the three isolates there are a few regions that are entirely homozygous (Table 2). No more frequent in the shorter half of the alignments than in the longer half, they must generally be longer than the alignments in which they occur. These regions may be identical by descent or may have arisen by conversion or by germ-line crossing-over at the four-strand stage of mitosis. That these processes may have occurred more frequently in MM than in MA and CR is suggested by its higher frequency of homozygous regions and lower mean heterozygosity (Table 2 and Figure S2). The infrequency of homozygous regions suggests, for each isolate, that its parents cannot have been closely related, as expected for individuals from a large panmictic population. The few alignments in which conversion or mitotic crossing-over may have erased evidence of sharing are not included in the totals given above or in Table 2.

Recombination

The finding that the pattern of sharing was the same in every one of the four regions examined by (Signorovitch *et al.* 2015), with MA sharing one of its homologs with MM and its other homolog with CR, suggested that the regions may not have recombined and therefore that entire haplotypes may have passed through meiosis intact, as in certain plants (Holsinger and Ellstrand 1984), an interpretation made plausible at the time by the earlier but subsequently disproven report that *A. vaga* has genomic features incompatible with meiotic recombination (Flot *et al.* 2016). In contrast, this pattern of sharing occurs in only 4 of our 331 much longer MM-MA-CR alignments. It may therefore be that the pattern encountered in the earlier study resulted from identity

by descent from remote ancestors, in which case such regions would generally be

shorter than the present alignments and therefore not counted as allele sharers. That

entire haplotypes are not conserved and that recombination definitely occurs is seen in

the observation that MA and MM share only a quarter of their genomes and share

considerably less with CR and in the existence of numerous alignments in which

sharing extends over only part of the region.

DISCUSSION

We find that MA and MM identically share homologs in 315 or just half (0.506, S.E. =

0.02) of the 622 MA-MM and MA-MM-CR alignments, therefore sharing ¼ of their

genome in segments at least as long as our alignments and possibly much longer. This

degree of sharing is consistent with sexual reproduction, in which the proportion of the

genome shared by two individuals inherited from their closest common ancestors, their

coefficient of relationship, is a power of ½.

In all of biology there are only three known modes of homologous genetic

exchange between individuals: sexual reproduction, horizontal gene transfer including

its various forms in prokaryotes, and parasexuality. For the observed allele sharing to

have occurred by homologous HGT would require massive horizontal transfer of long

DNA segments between MA and MM, displacing a homolog coincidentally in exactly

half (0.5, S.E. = 0.02) of the 622 alignments containing MA and MM, a degree of

consanguinity consistent with sexual reproduction, yet never displacing both, there

being no alignment in which MA and MM are of the form a/b and a/b.

Nor can the observed allele sharing be explained as the result of parasexuality.

In the parasexual cycle, known only in certain fungi and protozoans, nuclei from two

individuals fuse, forming nuclei of doubled ploidy that during subsequent generations

undergo occasional mis-division, only rarely yielding viable diploids. In bdelloids, this

would require nuclei from different individuals, sequestered in ovaries within the body of

the animal, somehow to undergo fusion, followed by a series of random chromosome

losses to give viable segregants, all having ten chromosomes and with MA-MM identical

sharing in exactly half of the 1,177 genomic regions we examined.

The finding in *A. vaga* of meiosis-related genes (Nowell *et al.* 2018), of homologous

chromosome pairs (Simion *et al.* 2020), and of Hardy-Weinberg equilibrium and decay

of linkage disequilibrium with increasing physical distance and phylogenetic

noncongruence (Vakhrusheva *et al.* 2020), although susceptible of other explanations,

are in entire agreement with our conclusion from the pattern of allele sharing that

bdelloids are facultatively sexual.

Generations since the MA-MM sharing event

The number of clonal generations since the MA-MM sharing event may be estimated

from the frequency of substitution differences between shared homologs, a few of which

differ by one or more substitutions; from the number of generations that would cause

mutational reduction of the frequency of identical sharers to fall significantly below the

observed value of 0.5; from the frequency of homozygosity; and from the mitochondrial

difference between MA and MM.

In addition to the 315 alignments in which MA and MM share identical homologs

there are 6 in which they share homologs that differ by a single substitution. For a mean

alignment length of 15 kb this is a frequency of 1.3×10^{-6} per bp. Substitution rates

measured in accumulation experiments with *Caenorhabditis elegans*, asexual *D. pulex*,

and *D. melanogaster range* from 2.3 to 5.5×10^{-9} per generation (Flynn *et al.* 2017).

Taking a substitution rate of 4×10^{-9} and assuming a Poisson distribution of the few

nucleotide substitution differences between shared homologs, this suggests that the

shared homologs may be separated by 100-200 generations.

If there were as many as 2,000 generations separating the shared homologs of

MA and MM and again assuming a substitution rate of 4×10^{-9} per generation, the

expected number of substitutions in regions 18-33 kb in length, the longest interval in

Figure 3, would be 0.14 – 0.26, reducing the proportion of identical sharers to 0.43 –

0.39, substantially less than the observed value of 0.5, suggesting that the number of

generations between the shared homologs is no more than about 1,000.

As tracts of homozygosity arising in a genetically diverse population are generally

erased by outcrossing, the frequency of substitutions in such tracts will increase with the

number of clonal generations since the last outcross. Assuming that, as in daphnia and

yeast, the likelihood of a site being covered by a tract of homozygosity to be about

4×10^{-5} per generation (Omilian et al. 2006; Xu et al. 2011; St. Charles and Petes 2013;

Flynn et al. 2017) and considering that the total length of MA regions is 13.4 Mbp of

which perfectly homozygous regions comprise some 151 Kbp, or about 1.1%, it appears

that there have been some 340-500 generations from when the sharing event occurred

to MA and 550-800 generations to MM.

A fourth estimate of the number of generations since the MA-MM sharing event

may be obtained by assuming that their mitochondria descend from a common mother

or maternal grandmother. Taking the difference of 5 substitutions or 2.5×10^{-5} between

their 14 kb mitochondria (Lasek-Nesselquist 2012) and a mitochondrial mutation rate of

1.5×10^{-7} (Xu *et al.* 2012; Flynn *et al.* 2017), suggests that the shared homologs

separated some 170 generations ago. These various estimates agree in suggesting that

the shared homologs of MA and MM are separated by no more than about a thousand

clonal generations.

Abundance of close relatives in the sampled population

Isolates MA, MM and CR were collected at widely separated sites as part of a

collection of only 29 individuals. What aspects of bdelloid life history could make finding

relatives as close as MA and MM in so small and widely dispersed a sample of what

must be an enormous population? It must be that the sampled population is largely

made up of relatively few, very large, widely dispersed clones descended from recent

crossing. Such an unusual population structure would result if sexual periods occur only

rarely, during a population bloom, with multiple rounds of mating among two or more

founding types, producing large numbers of closely related individuals. It may be that

males are produced and mating occurs only when particular mating types are present

together, causing one or both to produce haploid eggs and haploid males. At some

stage, from fertilization to zygote development, selfing may be prevented, allowing

mating only between different types, thereby avoiding the production of homozygous

progeny. Such mating, followed by wide dispersion and extensive clonal reproduction

would give rise to very large, widely dispersed clones of the products of recent crossing.

Meanwhile, lines that fail to outcross would suffer loss of heterozygosity caused by

conversion and mitotic crossing-over, and rapid clonal erosion, as seen in asexual

Daphnia pulex (Tucker *et al.* 2013), driving them to extinction unless revived by timely

outcrossing.

On this picture, field observations intended to detect males and mating should be

made during population blooms, as may require specific external stimuli, and in sizeable

bodies of water should it be that different but compatible types must be present in order

to initiate mixis. Further, by analogy with monogononts, the appearance of bdelloid

males may be confined to only a short interval during a population bloom, therefor

requiring frequent sampling for their detection (Wesenberg-Lund 1930).

Bdelloid life history - Eluding the Red Queen

It may be asked if there is a relation between the infrequency of bdelloid outcrossing

and bdelloid life history. A distinctive feature of the latter is the ability of bdelloids to

withstand desiccation and resume reproduction upon rehydration, an ability not present

in most of the fungi and other organisms that infect, parasitize, prey on or compete with

bdelloids (Wilson and Sherman 2010). In habitats that undergo desiccation and

rehydration the population of desiccation-intolerant antagonists will be greatly reduced

at each episode of desiccation while bdelloids will resume reproduction upon

rehydration. Bdelloids gain additional freedom from having to co-evolve with biological

antagonists by their ability to survive prolonged starvation, extremes of temperature and

exposure to toxic conditions lethal to other taxa (Ricci and Perletti 2006; Aguilera *et al.*

2007). Further, owing to their small mass when desiccated, about 10ug, once airborne,

even if associated with a small amount of adhering material, bdelloids may be

transported by wind or vectors over considerable distances (Fontaneto *et al.* 2008),

transferring to an environment where antagonists may be less abundant or less

antagonistic. The combination of anhydrobiosis and resistance to conditions inimical to

other taxa and dispersibility by wind, birds or other vectors therefore affords substantial

protection from what would otherwise be co-evolving biotic antagonists, reducing the

need for frequent recombination by largely eluding the "Red Queen" (Ladle *et al.* 1993;

Wilson and Sherman 2010; Wilson 2011).

Although bdelloids may be substantially freed of the need for sexual reproduction to

keep up with biological antagonists, other benefits of sex apparently maintain it. In

addition to the benefit of recombination in facilitating adaptive selection and limiting the

load of deleterious mutation, a further and more immediate benefit may be the

restoration of lost heterozygosity. An example of this may be seen in the suggestion of

(Tucker *et al.* 2013) that the early extinction of asexual *Daphnia pulex* may result from

the uncovering of preexisting recessive deleterious alleles caused by conversion and

deletion. Whether it is the masking of deleterious recessive alleles and/or some other

benefit of heterozygosity, its restoration by occasional outcrossing may constitute an

important benefit maintaining sex in bdelloid rotifers and more generally.

Literature Cited

Aguilera A., E. Zettler, F. Gomez, L. Amaral-Zettler, N. Rodriguez, *et al.*, 2007

Distribution of Eukaryotes in an extremely acidic, high-metal river. Molecular

Biology 1–41.

Altschul S. F., W. Gish, W. Miller, E. W. Meyers, D. J. Lipman, *et al.*, 1990 Basic

local alignment search tool. Journal of Molecular Biology 215: 403–410.

Arkhipova I. R., and M. Meselson, 2000 Transposable elements in sexual and

ancient asexual taxa. Proceedings of the National Academy of Sciences (USA)

97: 14473–14477. https://doi.org/10.1073/pnas.97.26.14473

Arkhipova I., and M. Meselson, 2005a Deleterious transposable elements and the

extinction of asexuals. BioEssays 27: 76–85.

Arkhipova I. R., and M. Meselson, 2005b Diverse DNA transposons in rotifers of the

class Bdelloidea. Proceedings of the National Academy of Sciences of the

United States of America 102: 11781–11786.

https://doi.org/10.1073/pnas.0505333102

Birky C. W., 2010 Positively negative evidence for asexuality. Journal of Heredity

101: 42–45. https://doi.org/10.1093/jhered/esq014

Chen S., Y. Zhou, Y. Chen, and J. Gu, 2018 Fastp: An ultra-fast all-in-one FASTQ

preprocessor. Bioinformatics 34: i884–i890.

https://doi.org/10.1093/bioinformatics/bty560

Coster W. de, S. D'Hert, D. T. Schultz, M. Cruts, and C. van Broeckhoven, 2018

NanoPack: Visualizing and processing long-read sequencing data.

Bioinformatics 34: 2666–2669. https://doi.org/10.1093/bioinformatics/bty149

Danecek P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, *et al.*, 2011 The variant

call format and VCFtools. Bioinformatics 27: 2156–2158.

https://doi.org/10.1093/bioinformatics/btr330

Debortoli N., X. Li, I. Eyres, D. Fontaneto, B. Hespeels, *et al.*, 2016 Genetic

Exchange among Bdelloid Rotifers Is More Likely Due to Horizontal Gene

Transfer Than to Meiotic Sex. Current Biology 26: 1–10.

https://doi.org/10.1016/j.cub.2016.01.031

Dolgin E. S., and B. Charlesworth, 2006 The fate of transposable elements in

asexual populations. Genetics 174: 817–827.

Ehrenberg C. G., 1838 *Die Infusionsthierchen als vollkommene Organismen.*

EinBlick in das tiefere organische Laben der Natur. L. Voss, Leipzig.

Fisher R. A., 1930 *The Genetical Theory of Natural Selection.* Clarendon Press,

Oxford.

Flot J.-F., B. Hespeels, X. Li, B. Noel, I. Arkhipova, *et al.*, 2013 Genomic evidence

for ameiotic evolution in the bdelloid rotifer Adineta vaga. Nature 500: 453–457.

https://doi.org/10.1038/nature12326

Flot J. F., N. Debortoli, B. Hallet, and K. van Doninck, 2016 Response to

Signorovitch et al. Current Biology 26: R755.

https://doi.org/10.1016/j.cub.2016.06.052

Flot J., N. Debortoli, B. Hallet, J. Narayan, and K. van Doninck, 2018 Reply to

Cross-Contamination Explains "'Inter and Intraspecific Horizontal Genetic

Transfers'"' between Asexual Bdelloid Rotifers (Wilson, Nowell & Barraclough

2018).'" bioRxiv. https://doi.org/10.1101/368209

Flynn J. M., F. J. J. Chain, D. J. Schoen, and M. E. Cristescu, 2017 Spontaneous

mutation accumulation in daphnia pulex in selection-free vs. competitive

environments. Molecular Biology and Evolution 34: 160–173.

https://doi.org/10.1093/molbev/msw234

Fontaneto D., T. G. Barraclough, K. Chen, C. Ricci, and E. A. Herniou, 2008

Molecular evidence for broad-scale distributions in bdelloid rotifers: everything

is not everywhere but most things are very widespread. Molecular ecology 17:

3136–3146. https://doi.org/10.1111/j.1365-294X.2008.03806.x

França M. B., a D. Panek, and E. C. a Eleutherio, 2007 Oxidative stress and its

effects during dehydration. Comparative biochemistry and physiology. Part A,

Molecular & integrative physiology 146: 621–631.

https://doi.org/10.1016/j.cbpa.2006.02.030

Fredrickson J. K., S. W. Li, E. K. Gaidamakova, V. Y. Matrosova, M. Zhai, *et al.*,

2008 Protein oxidation: key to bacterial desiccation resistance? ISME J 2: 393–

403.

Gladyshev E. A., M. Meselson, and I. R. Arkhipova, 2007 A deep-branching clade of

retrovirus-like retrotransposons in bdelloid rotifers. Gene 390: 136–145.

https://doi.org/10.1016/j.gene.2006.09.025

Gladyshev E., and M. Meselson, 2008 Extreme resistance of bdelloid rotifers to

ionizing radiation. Proceedings of the National Academy of Sciences of the

United States of America 105: 5139–5144.

https://doi.org/10.1073/pnas.0800966105

Gladyshev E., and I. Arkhipova, 2010 A subtelomeric non-LTR retrotransposon
        Hebe in the bdelloid rotifer Adineta vaga is subject to inactivation by deletions
        but not 5' truncations. Mobile DNA 1: 12.

Hamilton W. D., 1980 Sex versus non-sex versus parasite. Oikos 35: 282–290.

Hartfield M., S. I. Wright, and A. F. Agrawal, 2018 Coalescence and linkage
        disequilibrium in facultatively sexual diploids. Genetics 210: 683–701.
        https://doi.org/10.1534/genetics.118.301244

Hickey D. A., 1982 Selfish DNA: a sexually-transmitted nuclear parasite. Genetics
        101: 519–531.

Hill W. G., and A. L. Robertson, 1966 The effect of linkage on limits to artificial
        selection. Genetical Research 8: 269–294.

Holsinger K. E., and N. C. Ellstrand, 1984 The Evolution and Ecology of Permanent
        Translocation Heterozygotes. American Naturalist 124: 48–71.

Hsu W. S., 1956a Oogenesis in the Bdelloidea rotifer Philodina roseola Ehrenberg.
        Cellule 57: 283–296.

Hsu W. S., 1956b Oogenesis in Habrotrocha tridens (Milne). Biological Bulletin 111:
        364–374.

Hudson C. T., and P. H. Gosse, 1886 *The Rotifera or Wheel-Animalcules*.
        Longmans, Green, London.

Hur J. H., K. van Doninck, M. L. Mandigo, and M. Meselson, 2009 Degenerate
        tetraploidy was established before bdelloid rotifer families diverged. Molecular
        Biology and Evolution 26: 375–383. https://doi.org/10.1093/molbev/msn260

Judson O. P., and B. B. Normark, 1996 Ancient asexual scandals. Trends in

Ecology and Evolution 11: 41–46.

Kondrashov A. S., 1988 Deleterious mutations and the evolution of sexual

reproduction. Nature 336: 435–440.

Krisko A., M. Leroy, M. Radman, and M. Meselson, 2012 Extreme anti-oxidant

protection against ionizing radiation in bdelloid rotifers. Proceedings of the

National Academy of Sciences of the United States of America 109: 2354–

2357. https://doi.org/10.1073/pnas.1119762109

Ladle R. J., R. A. Johnstone, and O. P. Judson, 1993 Coevolutionary dynamics of

sex in a metapopulation: escaping the Red Queen. Proceedings of the Royal

Society Biological Sciences Series B 253: 155–160.

Lasek-Nesselquist E., 2012 A Mitogenomic Re-Evaluation of the Bdelloid Phylogeny

and Relationships among the Syndermata. PloS one 7: e43554.

https://doi.org/10.1371/journal.pone.0043554

Leewenhoeck A. van, 1677 Observations, Communicated to the Publisher by Mr.

Antony van Leewenhoeck, in a Dutch Letter of the 9th of Octob. 1676. Here

English'd: concerning Little Animals by Him Observed in Rain-Well-Sea- and

Snow Water; as Also in Water Wherein Pepper Had Lain In. Philosophical

Transactions of the Royal Society of London 12: 821–831.

https://doi.org/10.1098/rstl.1677.0003

Leewenhoeck A. van, 1702 Letter to the Royal Society December 25, 1702.

Proceedings of the Royal Society London B 23: 1291–1311.

Li H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-
Wheeler transform. Bioinformatics 25: 1754.

Li H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, *et al.*, 2009 The Sequence
Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352

Li H., 2018 Minimap2: Pairwise alignment for nucleotide sequences. Bioinformatics
34: 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Lively C. M., 2010 A review of red queen models for the persistence of obligate
sexual reproduction. Journal of Heredity 101: 13–20.
https://doi.org/10.1093/jhered/esq010

Madeira F., Y. M. Park, J. Lee, N. Buso, T. Gur, *et al.*, 2019 The EMBL-EBI search
and sequence analysis tools APIs in 2019. Nucleic Acids Research 47: W636–
W641. https://doi.org/10.1093/nar/gkz268

Magwene P. M., Ö. Kayıkçı, J. A. Granek, J. M. Reininga, Z. Scholl, *et al.*, 2011
Outcrossing, mitotic recombination, and life-history trade-offs shape genome
evolution in *Saccharomyces cerevisiae*. Proceedings of the National Academy
of Sciences 108: 1987–1992. https://doi.org/10.1073/pnas.1012544108

Marçais G., and C. Kingsford, 2011 A fast, lock-free approach for efficient parallel
counting of occurrences of k-mers. Bioinformatics 27: 764–770.
https://doi.org/10.1093/bioinformatics/btr011

Mark Welch D. B., and M. Meselson, 2000 Evidence for the evolution of bdelloid
rotifers without sexual reproduction or genetic exchange. Science 288: 1211–
1215.

Mark Welch D. B., and M. S. Meselson, 2001 Rates of nucleotide substitution in

sexual and anciently asexual rotifers. Proceedings of the National Academy of

Sciences (USA) 98: 6720–6724.

Mark Welch D. B., J. L. Mark Welch, and M. Meselson, 2008 Evidence for

degenerate tetraploidy in bdelloid rotifers. Proceedings of the National

Academy of Sciences 105: 5145–5149.

https://doi.org/10.1073/pnas.0800972105

Mark Welch D. B., C. Ricci, and M. Meselson, 2009 Lost Sex, pp. 259–279 in edited

by Schön I., Martens K., Dijk P. Springer Netherlands, Dordrecht.

McKenna A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, *et al.*, 2010 The

Genome Analysis Toolkit: A MapReduce framework for analyzing next-

generation DNA sequencing Data. Genome Research 20: 1297–1303.

Meadow N. D., and C. H. Barrows Jr., 1971 Studies on aging in a bdelloid rotifer .I.

The effect of various culture systems and longevity and fecundity. Journal of

Experimental Zoology 176: 303–314.

Molder F., K. P. Jablonski, B. Letcher, M. B. Hall, C. H. Tomkins-Tinch, *et al.*, 2021

Sustainable data analysis with Snakemake [version 1; peer review: 1 approved,

1 approved with reservations]. F1000Research 10: 1–25.

Muller H., 1932 Some genetic aspects of sex. American Naturalist 66: 118–138.

Muller H. J., 1964 The relation of recombination to mutational advance. Mutation

Research 1: 2–9.

Nowell R. W., P. Almeida, C. G. Wilson, T. P. Smith, D. Fontaneto, *et al.*, 2018

Comparative genomics of bdelloid rotifers: Insights from desiccating and

nondesiccating species, (C. Tyler-Smith, Ed.). PLOS Biology 16: e2004830.

https://doi.org/10.1371/journal.pbio.2004830

Omilian A. R., M. E. A. Cristescu, J. L. Dudycha, and M. Lynch, 2006 Ameiotic

recombination in asexual lineages of Daphnia. PNAS 103: 18638–18643.

https://doi.org/10.1073/pnas.0606435103

Ricci C., 1983 Life histories of some species of Rotifera Bdelloidea. Hydrobiolgia

180: 175–180.

Ricci C., and U. Fascio, 1995 Life-history consequences of resource allocation of

two bdelloid rotifer species. Hydrobiologia 299: 231–239.

https://doi.org/10.1007/BF00767330

Ricci C., 1998 Anhydrobiotic capabilities of bdelloid rotifers. Hydrobiologia 387/388:

321–326.

Ricci C., and M. Caprioli, 2005 Anhydrobiosis in bdelloid species, populations, and

individuals. Integrative and Comparative Biology 45: 759–763.

Ricci C., and F. Perletti, 2006 Starve and survive: stress tolerance and life-history

traits of a bdelloid rotifer. Functional Ecology 20: 340–346.

Rodriguez F., and I. R. Arkhipova, 2016 Multitasking of the pirna silencing

machinery: Targeting transposable elements and foreign genes in the Bdelloid

Rotifer Adineta vaga. Genetics 203: 255–268.

https://doi.org/10.1534/genetics.116.186734

Ruderfer D. M., S. C. Pratt, H. S. Seidel, and L. Kruglyak, 2006 Population genomic

analysis of outcrossing and recombination in yeast. Nature Genetics 38: 1077–

1081. https://doi.org/10.1038/ng1859

Schon I., K. Martens, and P. van Dijk (Eds.), 2009 *Lost Sex : The evolutionary*

biology of parthenogenesis. Springer, Dordrecht.

Sievers F., A. Wilm, D. Dineen, T. J. Gibson, K. Karplus, *et al.*, 2011 Fast, scalable

generation of high-quality protein multiple sequence alignments using Clustal

Omega. Molecular Systems Biology 7. https://doi.org/10.1038/msb.2011.75

Signorovitch A., J. Hur, E. Gladyshev, and M. Meselson, 2015 Allele Sharing and

Evidence for Sexuality in a Mitochondrial Clade of Bdelloid Rotifers. Genetics

200: 581–590. https://doi.org/10.1534/genetics.115.176719

Signorovitch A., J. Hur, E. Gladyshev, and M. Meselson, 2016 Evidence for meiotic

sex in bdelloid rotifers. Current Biology 26: R754–R755.

https://doi.org/10.1016/j.cub.2016.06.051

Simion P., J. Narayan, A. Houtain, A. Derzelle, L. Baudry, *et al.*, 2020 Homologous

chromosomes in asexual rotifer Adineta vaga suggest automixis. 10–12.

https://doi.org/10.1101/2020.06.16.155473

St.Charles J., and T. D. Petes, 2013 High-Resolution Mapping of Spontaneous

Mitotic Recombination Hotspots on the 1.1 Mb Arm of Yeast Chromosome IV.

PLoS Genetics 9: e1003434. https://doi.org/10.1371/journal.pgen.1003434

Talavera G., and J. Castresana, 2007 Improvement of phylogenies after removing

divergent and ambiguously aligned blocks from protein sequence alignments.

Systematic Biology 56: 564–577. https://doi.org/10.1080/10635150701472164

Tucker A. E., M. S. Ackerman, B. D. Eads, S. Xu, and M. Lynch, 2013 Population-

genomic insights into the evolutionary origin and fate of obligately asexual

Daphnia pulex. Proceedings of the National Academy of Sciences of the United

States of America 110: 15740–15745.

https://doi.org/10.1073/pnas.1313388110

Vakhrusheva O. A., E. A. Mnatsakanova, Y. R. Galimov, T. v. Neretina, E. S.

Gerasimov, *et al.*, 2020 Genomic signatures of recombination in a natural

population of the bdelloid rotifer Adineta vaga. Nature Communications 11.

https://doi.org/10.1038/s41467-020-19614-y

Weisenfeld N. I., V. Kumar, P. Shah, D. M. Church, and D. B. Jaffe, 2017 Direct

determination of diploid genome sequences. Genome Research 27: 757–767.

Weismann A., 1887 On the significancation of the polar globules. Nature 36: 607–

609.

Wesenberg-Lund C., 1930 Contributions to the biology of the rotifera, part II: The

periodicity and sexual periods. Kgl. Danske Vidensk. Selsk. Skrifter, Naturv. Og

Math. Afd. 9. II: 1–230.

Wilson C. G., and P. W. Sherman, 2010 Anciently Asexual Bdelloid Rotifers Escape

Lethal Fungal Parasites by Drying Up and Blowing Away. Science 574–577.

Wilson C. G., 2011 Desiccation-tolerance in bdelloid rotifers facilitates

spatiotemporal escape from multiple species of parasitic fungi. Biological

Journal of the Linnean Society 104: 564–574. https://doi.org/10.1111/j.1095-

8312.2011.01737.x

Wilson C. G., R. W. Nowell, and T. G. Barraclough, 2018 Cross-Contamination

Explains "Inter and Intraspecific Horizontal Genetic Transfers" between Asexual

Bdelloid Rotifers. Current Biology 28: 2436-2443.e15.

https://doi.org/10.1016/j.cub.2018.05.070

Xu S., A. R. Omilian, and M. E. Cristescu, 2011 High rate of large-scale hemizygous

deletions in asexually propagating Daphnia: Implications for the evolution of

sex. Molecular Biology and Evolution 28: 335–342.

https://doi.org/10.1093/molbev/msq199

Xu S., S. Schaack, A. Seyfert, E. Choi, M. Lynch, *et al.*, 2012 High mutation rates in

the mitochondrial genomes of Daphnia pulex. Molecular Biology and Evolution

29: 763–769. https://doi.org/10.1093/molbev/msr243

756 **Figure 1.** The bdelloid *M. quadricornifera*. (A) A feeding adult. Water and food particles

757 are swept into the mouth by the motion of a pair of ciliated rings, the corona. The animal

758 has attached itself to the glass slide by an adhesive secreted by its pedal gland,

759 somewhat obscuring the tail. (B) An unhatched egg shown at the same scale. (C) A

760 metaphase nucleus, showing the 10 chromosomes characteristic of the species.

Figure 2. Histograms showing the distribution of divergence between the most similar

homologs in each of the three pairs of isolates. (A) MA-CR, (B) MA-MM, (C) MM-CR.

Alignments with identical or very nearly identical MA-MM sharing form a discrete class

constituting half of the regions (panel B) showing that isolates MA and MM identically

share ¼ of their genomes inherited from their recent ancewstors. Bin size = 0.05

percent difference for the first bar in panel B, otherwise 0.1 percent.

Figure 3. The mean frequency of identical MM-MA sharing in each of 8 bins of

alignments of increasing length, each bin containing 76-79 alignments. Error bars

represent standard errors. It may be seen that the identically shared segments must

generally be longer than the longest regions examined.

Table 1. Representative difference matrices, phylograms and tic plots for four alignments. In the difference matrices, the number of alignments is given above the diagonal and the percent of the corresponding genotype is given below. Vertical lines in the tic plots represent sites of nucleotide difference. Allele numbers (1 or 2) are arbitrary.

Summary Statistics										
		Identical Sharing Number / Proportion			Homozygous Number / Proportion			Average Heterozygosity		
Alignment	Size Range (kb) # of Alignments	MA–MM	MA–CR	MM–CR	MA	MM	CR	MA	MM	CR
MA–MM	2,149-32,937	151			12	8		0.019	0.010	
	291	0.519			0.041	0.028				
MA–CR	2,051–29,791		1		1		0	0.018		0.018
	110		0.009		0.009		0			
MM–CR	4,511–31,052			7		18	12		0.010	0.020
	445			0.016		0.04	0.027			
MA–MM–CR	2,705–29,510	164	11	11	1	4	3	0.018	0.009	0.018
	331	0.495	0.033	0.033	0.003	0.012	0.009			
All alignments	2,051–32,937	315	12	18	14	30	15	0.018	0.010	0.019
	1,177	0.506	0.027	0.023	0.019	0.028	0.017			

784 **Table 2**. Summary statistics for the 1,177 non-overlapping alignments examined.

785 Identical MA-MM sharing occurs in 315 of the 622 alignments in which MA and MM are

786 included. The more frequent homozygosity and lower heterozygosity in alignments of of

787 isolate MM may reflect more frequent occurrence of homozygosing events along its

788 clonal lineage.

Acknowledgements. We thank Nicole El-Ali and Claire Hartmann for Nanopore

sequencing, Jae Hur for the script used in generating tic plots, Janet Montgomery for

overall editing, and Irina Arkhipova, Timothy Barraclough, Brian and Deborah

Charlesworth, Antoine Hout, Paul Simion and Karine van Doninck for critical reading of

the manuscript. This work was supported by Oxford Nanopore, the Harvard Faculty of

Arts and Sciences and by an anonymous donor.

Supplemental Material is available for this paper.

Correspondence and requests for materials should be addressed to Matthew

Meselson, email: msmeselson@gmail.com.

Author Contributions. VL and TS: Genome assembly and data analysis. MM: Rotifer

culturing DNA preparation and data analysis. MM wrote the manuscript, which was

edited and approved by all authors.

Competing Interest Declaration The authors declare no competing interests.