Task-dependent vocal adjustments to optimize biosonar-based information acquisition

Daniel Lewanzik & Holger R. Goerlitz

Acoustic and Functional Ecology, Max Planck Institute for Ornithology, Eberhard-Gwinner-Straße, 82319 Seewiesen, Germany

Authors for correspondence: daniel.lewanzik@gmx.de, hgoerlitz@orn.mpg.de

Running title: Task-dependent vocal adjustments in biosonar

Summary statement: Having the right information for a specific job is crucial. Echolocating bats flexibly and independently adjust different call parameters to match the sensory-motor challenges of four different tasks.

Abstract

Animals need to acquire adequate and sufficient information to guide movements, yet information acquisition and processing is costly. Animals thus face a trade-off between gathering too little and too much information and, accordingly, actively adapt sensory input through motor control. Echolocating animals provide the unique opportunity to study the dynamics of adaptive sensing in naturally behaving animals without interference, since every change in the outgoing echolocation signal directly affects information acquisition and the perception of the dynamic acoustic scene. Here we investigated the flexibility with which bats dynamically adapt information acquisition depending on a task. We recorded the echolocation signals of wild-caught Western barbastelle bats (*Barbastella barbastellus*) while drinking on the wing, flying through an opening, landing on a wall, and capturing prey. We show that the echolocation signal sequences during target approach differed in a task-dependent manner; bats started target approach earlier and increased information update rate more when the task became increasingly difficult, and bats also adjusted dynamics of call duration shortening and peak frequency shifts accordingly. We provide insights into how echolocating animals deal with the constraints they face when sequentially sampling the world through sound by adjusting acoustic information flow in a highly dynamic manner. Our results further highlight the paramount importance of high behavioural flexibility for acquiring information.

Keywords: active sensing, behavioural flexibility, sensory processing, sensory-motor constraints

Introduction

Information is the sensory foundation of adaptive behaviors (Dall, Giraldeau, Olsson, McNamara, & Stephens, 2005). Hence, animals constantly integrate a wide range of information to decide on appropriate actions (Gomes et al 2016; Lewanzik, Sundaramurthy, & Goerlitz, 2019; Prat & Yovel, 2020). Acquiring appropriate information before making decisions is of paramount importance, particularly when decisions have direct fitness consequences, e.g. when failing to evade predators or finding food patches. This universal need for informed decisions is reflected in the evolution of the multitude of senses and their extreme speed and sensitivity (Elemans, Mead, Jakobsen, & Ratcliffe, 2011; Morshedian & Fain, 2015). Yet, animals face the challenge of distinguishing between relevant and irrelevant information (Bates, Simmons, & Zorikov, 2011; Beetz, Hechavarría, & Kössl, 2016), since the central nervous system processing capacity is limited and neural processing is metabolically costly (Laughlin, de Ruyter van Steveninck, & Anderson, 1998; Niven & Laughlin, 2008). Accordingly, selective attention, i.e. the ability to perceptually focus on relevant stimuli while ignoring irrelevant ones, is a crucial feature of cognitive processing (Lavie, 2005) and likely evolved early in evolution (Sridharan, Schwarz, & Knudsen, 2014). In addition, animals apply pre-attentive filters by actively adjusting their primary sensory input, e.g., by controlling the timing (Moss, Chiu & Surlykke 2011; Kurnikova et al 2017; Ladegaard & Madsen 2019) and spatial focus of sensory sampling (Yarbus 1967; Kugler & Wiegrebe, 2017; Geipel et al., 2019; Taub & Yovel, 2020). To study how animals control their primary sensory input, active sensing is an excellent model, as active sensing animals generate and adjust the signal for probing their environment, which we in turn can observe to investigate the animal's information acquisition strategy.

Echolocating bats can orient and forage in complete darkness by sound. They control information quantity by changing call rate, and information quality by adjusting various call features, such as duration, amplitude, frequency, and directionality. Despite the large ecological variety of the more than 1200 bat species that echolocate (Wilson & Mittermeier, 2019), every bat needs to perform specific tasks on a daily basis, such as exiting the roost through an opening, drinking from a water body, capturing prey, and landing on surfaces. Each of those tasks constitutes a very different sensory-motor challenge, which bats might solve by adaptively adjusting their biosonar parameters to optimize the rate and features of acquired information. Yet, an adjustment that is aiding navigation during one task might be disadvantageous in another situation. Increasing call rate, for instance, corresponds to an increase of information update rate, which bats commonly perform when approaching objects (Fenton, Jensen, Kalko, & Tyack, 2014; Schnitzler, Moss, & Denzinger, 2003). Particularly, when approaching flying prey, aerial-hawking bats emit a characteristic call sequence with extremely high call rates, the so-called feeding (or final / terminal) buzz, right before capturing prey (Griffin, 1958; Kalko & Schnitzler, 1989; Lewanzik et al., 2019; Ratcliffe, Elemans, Jakobsen, & Surlykke, 2013). However, high call rates might be energetically expensive especially for bats calling at high amplitudes (Currie, Boonman, Troxell, Yovel, & Voigt, 2020). Also, high call rates can lead to call-echo ambiguity and

thus impair ranging and navigation, when echoes of a call return only after the next call had been

emitted (Beetz, Kössl, & Hechavarría, 2019; Moss, Chiu, & Surlykke, 2011). Thus, bats temporally

arrange their calls in groups, particularly when a task becomes challenging, with longer intervals

between groups allowing also echoes from distant objects to arrive before the next call is emitted

(Wheeler et al., 2016); stable call intervals within a call group might aid object localisation and

tracking (Moss et al., 2011). Similar trade-offs exist also for other call features: long-duration calls are

a means of increasing the energy emitted and accordingly, the volume ensonified, and thus are often

used in open space (Schnitzler & Kalko, 2001). However, long-duration calls increase the probability

that echoes reflected off close-by objects overlap in time with the outgoing call ('forward masking'),

which can impair object detection and evaluation (Schnitzler et al., 2003). The same extension of

detection distance can be achieved by increasing call amplitude, yet further increasing the amplitude of

high-amplitude calls comes at a disproportionately high energetic costs (Currie et al., 2020). Also,

high call amplitudes would result in a large number of 'uninformative' echoes in cluttered

environments, which could distract from relevant echoes and challenge sensory processing (Wahlberg

& Surlykke, 2014). Thus, bats usually reduce call amplitude while closing in on targets (Koblitz et al.,

2011). Simultaneously, bats often increase bandwidth, i.e. the frequency spectrum of their calls when

approaching a target. A broad bandwidth may be helpful to evaluate spectral interference patterns in

target echoes but, on the other hand, broad bandwidth calls usually come with decreased detectability

since the emitted energy is distributed over a large frequency range (Wahlberg & Surlykke, 2014).

To date, biosonar adjustments during object approach and prey capture have been well studied

in a very limited number of species only. Most of these species are Vespertilionid bats that emit their

echolocation calls through the mouth (e.g. Falk, Kasnadi, & Moss, 2015; Hiryu, Hagino, Riquimaroux,

& Watanabe, 2007a; Koblitz et al., 2011; Ratcliffe & Dawson, 2003; Russo, Jones, & Arlettaz, 2007;

Simmons, Brown, Vargas-Irwin, & Simmons, 2020). Some studies focused on the specialised group of

flutter detecting horseshoe bats instead, which emit unique constant frequency calls through the nose

(Tian & Schnitzler, 1997; Yamada et al., 2020). Barbastelle bats differ from the majority of other bat

species in that they emit two different call types alternatingly through mouth and nose (Denzinger,

Siemers, Schaub, & Schnitzler, 2001; Seibert, Koblitz, Denzinger, & Schnitzler, 2015). Also, in

contrast to most other aerial-hawking species, barbastelle bats emit low-amplitude calls that result in

very short detection distances, but that allow them to almost exclusively feed on eared moths

(Goerlitz, Ter Hofstede, Zeale, Jones, & Holderied, 2010; Lewanzik & Goerlitz, 2018). Despite their

low call amplitude, barbastelle bats capture prey in fst flight in (semi-)open habitats; however, they

usually roost and forage inside forest and thus are used to cluttered conditions (Dietz, von Helversen,

& Nill, 2007; Kerth & Melber, 2009; Zeale, Davidson-Watts, & Jones, 2012). Here, we comparatively

investigated the biosonar adjustments of barbastelle bats during four object-approach tasks. We

recorded the biosonar of barbastelle bats under controlled conditions in the flight room while drinking,

flying through an opening, capturing prey and landing, and we compare how they adjust their
information acquisition behaviour to the task.

Materials and methods

Capture and husbandry

In August 2018, we captured three adult male barbastelle bats *Barbastella barbastellus* using mistnets

(16×16 mm^2 mesh size; Ecotone, Gdynia, Poland) at emergence from a roost in eastern Bavaria

(Germany) and transferred them to the husbandry facilities of the Max Planck Institute for Ornithology

in the night of capture. We kept the bats for about three weeks in an aviary ($1.2 \times 2.4 \times 2.0$ m^3, W $\times$ L

$\times$ H) before releasing them at the site of capture. We provided *ad libitum* access to fresh water at all

times. Live moths and/or mealworms (larvae of *Tenebrio molitor*) were provided *ad libitum* during

training and experiment in a flight room. If bats had not reached their initial body mass after foraging

in the flight room, we additionally hand-fed mealworms before releasing bats back into the aviary.

Experimental setup

Experiments were conducted in an echo-attenuated flight room. We divided the flight room into a

'retreat compartment' ($2.7 \times 3.4 \times 3.0$ m^3, L $\times$ W $\times$ H) and a 'foraging compartment' (5.6 x 3.4×3.0

m^3, Fig. S1) by a combination of a wooden wall at the bottom and a thick felt curtain on top. Bats

could fly between compartments through a rectangular opening (size: 1.3 x 0.5 m², W x H) in the

wooden wall, whose centre was at 0.75 m above ground and 1.05 m from the closer side wall. The

foraging compartment contained a large pool (1.4 x 2.3 m²) to drink and a movable stand to offer

tethered mealworms (1.92 m above ground). During half ($17/35$) of the prey capture trials, we ran a

"flutter simulator" about 10 cm next to the mealworm. The flutter simulator, a small plastic propeller

of the size and shape of a single wing of *Noctua pronuba,* one of the barbastelle's main prey insects

(Goerlitz et al., 2010), imitated the wing beat of a flying *Noctua pronuba* (ca. 43 Hz; Waters & Jones,

1994). During the other half of prey capture trials, the propeller engine was running as well but the

wing was detached. Since we did not find any effect of the propeller (see supplementary

information),), we pooled the data for all other analyses. We positioned four microphones

(CM16/CMPA, Avisoft Bioacoustics, Glienicke, Germany) to record the bats' echolocation behaviour

at various positions: two microphones at both short ends of the pool at about 10 cm above ground

pointing towards the centre of the pool; one microphone in a corner of the foraging compartment close

to the site at which bats landed most often during training, pointing towards the bats' common

approach direction; and one microphone at the top of the mealworm stand pointing towards the

tethered mealworm. A fifth microphone (FG-O, Avisoft Bioacoustics) was placed at the side of the

opening at 98 cm above ground, pointing towards the bats approach direction from the foraging

compartment. Calls on CM16 microphones were recorded with custom-written Matlab code
(MathWorks, Natick, MA, USA), SoundMexPro software (HörTech gGmbH, Oldenburg, Germany), a
Scarlett OctoPre preamplifier (Focusrite Plc, High Wycombe, UK), and a soundcard (Fireface 802,
RME, Haimhausen, Germany; 192 kHz sampling rate, 16-bit resolution). Calls on the FG-O
microphone were recorded at 500 kHz sampling rate using UltraSoundGate 416H (Avisoft
Bioacoustics) and Avisoft Recorder software. We further video recorded the bats' drinking, foraging,
and compartment switching behaviour with an infra-red sensitive camera (HDR-CX560, Sony
Corporation, Tokyo, Japan). We placed infrared lights behind the opening inside the retreat
compartment perpendicular to the bats' flight trajectories, such that bats got illuminated in the moment
they passed the opening. Further infrared lights were used to illuminate the pond and the tethered
mealworm. Video recordings were also used to distinguish successful prey captures from unsuccessful
prey capture attempts, both of which ended with a feeding buzz.
158
Experimental procedure
In the first two nights after capture, we let all bats fly together inside the flight room with multiple
tethered live moths and mealworms offered as prey inside the foraging compartment. During the
following days, we let bats fly singly in the flight room in random order. Once bats learned to capture
tethered prey, we progressively reduced the number of prey items until only two fixed positions were
enough to feed the bats. We then started to record the bat when it was drinking or landing while
continuously replacing removed prey at the two fixed positions to feed the bats. After multiple nights
with two fixed prey locations, we started to only offer a single tethered mealworm at a time at one
position (the movable stand), and started to record the bat when attacking this prey item and also when
flying through the wall opening from the foraging compartment into the retreat compartment ('exit').
After bats had captured the mealworm, they usually flew into the retreat compartment. We then closed
the opening, replaced the mealworm and moved its stand to one of six randomly determined positions.
After re-opening the opening between the compartments, bats usually flew immediately into the
foraging compartment. Once a bat stopped foraging and began grooming itself, we captured it using a
hand net, weighed it, hand-fed it more mealworms if necessary, brought it back into the aviary, and
released the next individual into the flight room. Every individual could forage in the flight room
every night.
176
Sound analysis
We analysed the echolocation behaviour of the bats when approaching the pond to drink, the wall to
land, the wall opening to exit the foraging compartment, and the prey to capture. Particularly, we
compared the echolocation behaviour during the approach phase between tasks since - in contrast to
search calls - bats lock their approach calls onto the target (Koblitz et al., 2011; Surlykke, Ghose, &
Moss, 2009), indicating the bats' intention to execute a task and minimizing variation in measured call

parameters due to variation in sonar beam direction. Thus, parameters of approach but not of search phase calls are likely adjusted to a given task.

In total, we analysed the sound recordings (wav-file) of 98 object-approach flights using SASLabPro software (Avisoft Bioacoustics). Since landing and drinking were relatively rare events, we analysed all available recordings of landings (N=15) and chose similar numbers of drinking / drinking attempt recordings (N=18 and 15, respectively; at least 5 per individual) with a good signal-to-noise ratio from across almost all experimental days. Of exit, capture and capture attempt recordings, we only analysed recordings from the last two nights, i.e. 22 and 23 days post capture, when bats were well familiar with the setup and reliably foraged on the meal worm (N=15, 16, and 19 recordings with at least 5 recordings per individual and task,). We first highpass-filtered the recordings at 20 kHz (Hamming window, 1024 taps) and downsampled those recordings that were initially recorded with a higher sampling rate to 250 kHz before setting a fixed threshold (−60 dB FS) to detect bat calls above the noise floor. We then manually removed echoes whose amplitude was above detection threshold. For all calls above threshold (N = 5749), peak frequency (from the spectrogram-derived mean spectrum), call duration (as defined by the call detection threshold), and call interval were calculated automatically. We defined a call's peak frequency to be within the first or second harmonic if it was below or above 55 kHz, respectively (compare Fig. S2). We then manually determined 'call type' (search type I & type II, approach, buzz) of a large subset of calls (3930 calls = 68%) representing all individuals. In barbastelle bats, type I and type II search calls and approach/buzz calls are easily distinguishable by their frequency-time structure (Denzinger et al., 2001; Lewanzik & Goerlitz, 2018; Seibert et al., 2015). We used distributions of inter-call-intervals and peak frequency of these manually determined calls to identify ‚call type' of all remaining calls as follows: We defined calls as search calls when both preceding and following inter-call-interval were ≥36 ms and as approach/buzz calls otherwise. Among search calls we distinguished between type I and type II search calls based on peak frequency (type I: peak frequency either ≤37 kHz or between 55 kHz and 66 kHz; type II otherwise). This might have introduced a few mis-classifications, yet overlap between peak frequency distributions of type I and II search calls is low (Fig. S2) and, importantly, our analyses are not based on search but on approach/buzz calls. Among approach/buzz calls, we classified calls as 'buzz call' if either the preceding or following inter-call-interval was ≤6 ms; otherwise, they were classified as approach calls.

Since bats interspersed approach calls within search call sequences, we defined the onset of the final approach to the object as that approach call after which only approach and buzz but no search calls followed until the 'event' (with the exception of two 'exit trajectories' in which bats used a search call already shortly before exiting through the window; Fig. 1). The onset of the final buzz was defined accordingly, i.e. as that buzz call after which bats only emitted buzz calls. For both successful prey captures and unsuccessful prey capture attempts (distinguished via video recordings), we defined the event time as the time of the last buzz call. We distinguished successful drinking events from

unsuccessful drinking attempts by the splashing sound produced when the bat touched the water, which we used as event time in actual drinking events (Fig. 2b), but which was absent in unsuccessful drinking attempts. In the latter case, we defined the event time as the time of the last approach call, which approximates the behaviour in successful drinking events. For landing events, we used the sound of wall contact as event time, which was not always unambiguously visible in the spectrogram but always audible when listening to the sound file. For exit events, i.e. when flying from the foraging into the retreat compartment, we used the last call above call detection threshold as event time (because bats passed a wooden room divider, their calls were not recorded anymore once they flew through the opening). We termed the duration between event time and first post-event call 'post-event pause'.

Statistical analysis

We used the statistical freeware R version 3.6.2 (R Core Team, 2019) to analyse durations of approach phase and post-event pause as well as temporal dynamics of call interval, duration, and peak frequency, and to compare these parameters between tasks. We analysed only the calls (N = 5002) during the bats' final approach to the object, i.e., approach and buzz calls but not search calls. Since bats grouped their calls, we analysed inter-call-interval separately for shorter within-group intervals and longer between-group intervals. We defined the first intra-group interval of any call group as the interval that was at least 30% shorter than the preceding call interval or that was absolutely smaller than 10 ms. Following call intervals were considered intra-group intervals as long as they differed maximally by 20% or 5 ms compared to the mean intra-group interval of the current call group (Fig. 2 & Fig. S3).

We fit generalised linear mixed-effects models with a Gaussian error distribution to our response variables 'onset of final approach' (rel. to the event time), 'post event pause', 'call duration', and 'inter-call interval' (both within-group and between-group) using the lmer function of the lme4 package (D. Bates, Maechler, Bolker, & Walker, 2015). 'Onset of final object approach' and 'post event pause' were modelled as a function of the fixed effects 'event type' (categorical with six levels: drinking, drinking attempt, exit, landing, prey capture, and unsuccessful prey capture attempt) and – to account for repeated measures – the random effect 'bat ID' (categorical with three levels). 'Call duration' and 'inter-call interval' were modelled as a function of the fixed effects 'time before event' (continuous), 'event type', and the interaction between the two. Further, we included a 3rd order polynomial of 'time before event' as well as its interactions with 'event type' as predictors in the models since 'time before event' had non-linear effects on these response variables. To prevent autocorrelation, we also included 'call duration of previous call' or 'call interval of previous call' (continuous) as predictor in these models. Random effects included were 'trajectory no' (categorical with 98 levels) to account for potential dependency among calls within the same trajectory, and 'bat ID'.

To model changes in the proportion of calls with peak frequency in the 2nd harmonic over

time, we first calculated the number of calls per 0.3 s time bins with peak frequency in the 1st and 2nd

harmonic, respectively (i.e. < or >= 55 kHz; compare Fig. S2). We then fitted a generalised linear

mixed model with binomial error distribution to the response variable 'proportion of calls with peak

frequency in 2nd harmonic' (weighed by the total number of calls per bin) using the glmer function of

the lme4 package; including the fixed effects 'time bin' (modelled as continuous variable) as a 3rd

order polynomial, 'event type', the interaction thereof, and the random effects 'trajectory no' and 'bat

ID'. In contrast to all other models, we here included data of the entire object approach call sequence

(i.e. including also search and approach calls before the beginning of the final approach phase)

because binning the data substantially reduced sample size and would not allow modelling polynomial

effects otherwise.

All continuous predictors except 'time bin' were scaled by their standard deviation and (with

the exception of 'time before event' and 'time bin') centred on their mean. Model assumptions were

verified by plotting residuals versus fitted values and by inspecting QQ□plots of the model residuals

and of the random effects. We assessed residuals for temporal dependency by plotting estimates of the

autocorrelation function using R's acf function.

We used the sim function of the R□package arm (Gelman & Su, 2016) to simulate the values

of the posterior distributions of the model parameters (1,000 simulations) and to obtain posterior mean

parameter estimates. We extracted 95% credible intervals (CIs) around the posterior mean estimates

and considered support for an effect as strong if zero was not included within the 95% CI and as low if

overlap with zero was small, respectively. If 95% CIs were centred on zero, we considered this as

strong support for the absence of an effect. We also used the sim function to obtain mean predictions

of response variables and 95% CIs around these predictions for various combinations of predictors for

plotting purposes. Since we included 'previous call duration' and 'previous call interval' as predictors

in models for call duration and call intervals, respectively, but values of these predictors change over

the course of the approach, we used a 'moving average' approach to calculate predictions of the other

response variables (instead of using the mean value; size of the moving window: 0.3 s and 0.05 s for

call times before/after -0.3 s).

Results

We investigated how flexibly bats adjust their echolocation to control their primary sensory

information when executing various tasks, i.e. when approaching a pond to drink, an opening to fly

through, a wall to land on, and prey to capture. In search flight, i.e. before initiating object approach,

bats usually emitted an alternating pattern of type 1 / type 2 search calls (Fig. 1), which is typical for

barbastelle bats. Peak frequencies of the first harmonic of type I and type II search calls were 33.9 ±

1.6 kHz (mean ± sd; N = 126) and 41.5 ± 2.2 kHz (N = 740), respectively (Fig. S2). At times, search

calls were interspersed with more broadband approach-like calls. Subsequently, bats stopped emitting search calls altogether and switched to approach and buzz calls only (Fig. 1) with first harmonic peak frequencies of 41.3 ± 3.4 kHz (mean ± sd; N = 2656) and 40.2 ± 2.4 kHz (N = 1278), respectively (Fig. S2). In trajectories, in which we recorded search calls, the last search call was mostly (in 72/82 trajectories: 88%) a type II search call. Both the similar peak frequency and the time-frequency structure suggest that approach and buzz calls are more similar to type 2 than type 1 search calls, likely sharing the same mechanism of vocal production and (nasal) emission.

Fig. 1: Dynamic emission of different call types during object approach. The start of the final object approach is highlighted in red for each trajectory. Horizontal error bars indicate mean posterior predictions and credible intervals from model simulations for the start of the final object approach (black) and for the timing of the first post-event call (grey).

We defined the beginning of the approach phase as the time of the first approach call after
which bats did not emit any further search calls. Approach phase duration differed between tasks,
indicating different needs for sensory information. During drinking, landing and prey capture, the bats
started the object approach at about 0.6-0.7 s before the event (Fig. 1, tables S3/S4). In contrast, when
exiting through the opening, the bats started object approach later at about 0.3 s before the event,
resulting in a ca. 50% shorter approach duration. In drinking attempts (in contrast to actual drinking
events), approach duration was longer than in all other tasks (Fig. 1, table S5), likely due to continued
emission of approach calls while the bats continued trying to drink.

Next, we analysed the dynamic changes and task-dependent differences of temporal and
spectral call parameters during the final object approach (i.e., only approach and buzz calls). We will
present results for call intervals, call durations, and call peak frequencies. Neither of these, nor the
duration of approach phase, buzz, or post-event pause differed between capture trials with and without
the flutter simulator (tables S1/S2).

320
321
322

Fig. 2: Waveform (a), spectrogram (b), and call intervals (c) of an exemplary final approach phase call
sequence during drinking. The splash resulting from the bat touching the water surface is visible in the
spectrogram at the event time (x=0). In c), within-group (grey) and between-group (black) call
intervals are highlighted.

Temporal patterning of information acquisition

In all tasks, bats initially decreased their call intervals, thereby increasing the amount and update rate

of sensory information. Bats also grouped their calls into groups, with shorter within-group call

intervals (mean ± sd: 11.2 ± 11.7 ms) and longer between-group call intervals (45.9 ± 35.0 ms; Fig. 2

& Fig. S3). Beyond those common temporal patterns, the dynamics of both within- and between-group

call intervals differed between tasks (Fig. 3, tables S3-S5): bats initially reduced both call intervals

during the approach phase, but then stopped reducing (landing, drinking) or even increased call

intervals towards the end of the approach (exiting, drinking attempts). In contrast, only during prey

capture (attempts), bats kept on reducing call intervals until the event. In addition, the bats started their

approach already with shorter call intervals during prey capture (attempts) than during any other task,

reducing them further to the shortest call intervals right before capturing prey (min: 3.5 ms). Also,

terminal buzzes, i.e. sustained sequences of buzz calls (call interval < 6 ms) were only emitted during

prey capture (attempt) tasks (Fig. 1, Fig. 3b). The onset of the buzz was at 169 ± 76 ms (mean ± sd)

before capture and did not differ between successful and unsuccessful prey capture trials (tables

S3/S4). Note, however, that there was no clear distinction in call interval between approach and buzz

phases, but rather a gradual transition.

As a result of these different dynamics, update rates right before the event were lowest when

approaching the opening to exit the foraging compartment, intermediate when approaching the pool to

drink, high before landing, and highest (up to max. 286 s^{-1}) before capturing prey (Fig. 3). With the

exception of exit and landing trajectories, bats always refrained from emitting echolocation calls for a

short while after any event. This post-event pause was longer after drinking (153 ms) than after any

other event (9-64 ms; Fig. 1, tables S3/S4).

Call duration

With the exception of drinking (attempts), during which call duration initially seemed to increase and

then remained approximately constant, call duration in all other tasks was longest at the beginning of

the approach and subsequently decreased throughout the approach. Maximum call durations varied

between 1.7 ms (landing) and 2.4 ms (capture); minimum durations at the end of the final object

approach were between 1.0 ms (prey captures) and 1.6 ms (drinking / drinking attempts; Fig. 4, tables

S3/S4). As before, call duration dynamics were task-specific beyond those common temporal patterns

(Fig. 4). Bats controlled call duration most dynamically when capturing prey, starting with the longest

(2.4 ms) and ending with the shortest call duration (1.0 ms). Thus, they more than halved their call

durations and, accordingly, the call-echo-overlap zone (Schnitzler & Kalko, 2001; Wilson & Moss,

2004), i.e., the range where returning echoes overlap in time with the outgoing call. Shortening of call

duration started generally slow and was most pronounced just prior to prey captures. In contrast,

landing approaches directly started with shorter call durations (1.7 ms) that also changed only slowly.

Thus, call durations during landing were the shortest of all tasks throughout the approach, except for

the last ~70 ms, when call durations were shorter during capture (attempts). When exiting, shortening

of call duration started the latest of all tasks, and then followed a similar linear trend as during capture.

During drinking (attempts), bats used the longest call duration of all tasks at the end of a task (1.6 ms,

Fig. 4, tables S3-S5).

Frequency changes during object approach

Bats changed the spectral characteristics of their calls during object approach, with some task specific

differences. These spectral changes were not so much achieved by shifting the peak frequency of each

harmonic (Fig. 5a), but by changing energy distribution between harmonics (Fig. 5b). Mean peak

frequencies of the first and second harmonics over the entire final object approach were rather constant

at 39-43 kHz and 71-85 kHz, respectively. At the beginning of the object approach, almost all calls

(91-99%) had peak frequencies within the first harmonic. Only landing trials started with a substantial

proportion of calls (26%) having their peak frequency in the second harmonic (Fig. 5b), which then

further increased earlier and more pronounced than during drinking, exit, and prey captures (Fig. 5b).

Interestingly, bats used a higher proportion of calls with peak frequencies in the second harmonic

during successful than unsuccessful prey capture trials (Fig. 5b, see non-overlapping confidence

intervals; Table S5). Towards the end of the object approach, the proportion of calls with peak

frequency in the second harmonic went back to pre-approach levels in most tasks. Only during

drinking trials, this proportion started to increase later and kept increasing until the very end of the

object approach. In consequence, at the end of the object approach, the proportion of calls with peak

frequency in the second harmonic was substantially higher during drinking trials than during any other

task (Fig. 5b, table S5).

Fig. 3: Dynamics of A) between- and B) within-group call intervals (left y-axis) and resulting

information update rates (right y-axis) during the final object approach in six different tasks. Depicted

are model predictions (solid lines) and 95% credible intervals (coloured bands). Predictions are made

starting from the event-specific median start time of the final (object) approach phase.

Fig. 4: Dynamics of call duration (left y-axis) and call-echo overlap zone (right y-axis) during the final

approach phase prior to six different events. Call-echo overlap zone indicates the bat-object distance

below which returning echoes overlap in time with the outgoing call; it was calculated assuming a

speed of sound of 343 m/s. Depicted are model predictions (solid lines) and 95% credible intervals

(coloured bands). Predictions are made starting from the event-specific median start time of the final

object approach phase.

Fig. 5: Peak frequency dynamics during the final object approach in six different tasks. a) Peak frequency of the first (red) and second (orange) harmonics did not change substantially over the course of the final object approach. Dots: individual calls; coloured lines: mean ± sd per time bin. b) Proportion of calls with most energy in the second harmonic, i.e. at higher frequencies. Depicted are posterior mean predictions (solid lines) and credible intervals (grey areas) from a generalised mixed model with binomial error distribution. We also show the underlying data, i.e. the proportion of calls with peak frequency in the second harmonic per trajectory and time bin (black dots, jittered around bin centre). Further, for each time bin we overlaid the mean (over all trajectories of any given task) proportion of calls with peak frequency in the second harmonic (white dots), which fit well the model predictions.

Discussion

Here, we demonstrate that the aerial-hawking insectivorous barbastelle bat adjusts various aspects of its active sensory sampling strategy in a task-dependent manner. We show that barbastelle bats start their sensory target approach behaviour later and increase call rate less when approaching an opening than when drinking, landing, or capturing prey. On the contrary, bats only emitted buzz sequences with highest information update rates and shortest call durations when attacking prey. Lastly, bats used high information update rates and high frequencies for high spatial accuracy the most when landing, i.e., approaching a solid surface. Thus, we argue that the echolocation adjustments the bats made during target approach reflect the sensory-motor difficulty of any given task. Our results suggest that it is comparably easy for bats to navigate through an opening but challenging to capture small prey, and that bats adjust their echolocation parameters in a task-specific matching its difficulty. Thus, in the following we discuss the challenges bats face in a task-specific manner before widening the scope to other taxa and discussing the lessons learned from comparing across tasks.

Navigating through an opening

Presumably, among the tasks we studied it is easiest for echolocating bats to exit through an opening, particularly since the opening we offered was comparably large (0.5 x 1.3 m, height x width; wingspan of barbastelles is 26-30 cm (Skiba, 2009)), constant and predictable. As long as bats reliably headed towards the opening inside the wall and not the wall *per se*, ranging did not need to be particularly precise. Additionally, the bats' spatial memory (Barchi, Knowles, & Simmons, 2013; Thiele & Winter, 2005) likely reduced the need for high update rates before exiting, particularly since bats had been used to the flight room, the wall and the opening for about three weeks before we started to record exit trajectories. We thus suggest that the type of sensory-motor challenge and the bats' experience and memory rendered the exit task the least challenging one. Accordingly, compared to the other three tasks, bats initiated their approach the latest and used only few approach calls with low update rates when navigating through the opening. Arguably, flight speed during exit is faster than during the other tasks. Together with the long call intervals during exit, this means that the distances flown between subsequent emissions are even larger during exit than in all other tasks, reemphasising that exiting through an opening poses the least sensory-motor challenges of all tasks tested. The larger flight speed and later onset of approach could potentially mean that the bats use distance to object to initiate their approach (in contrast to time to contact that we analysed). However, previous three-dimensional tracking of barbastelle prey capture behaviour under two different conditions showed no such pattern: barbastelles attacking tethered prey in the field and flight room flew at different speeds (7.4 vs. 3.1 m/s) and started their approach at different distances (1.6 vs. 1.2 m) and times to contact (200 vs. 390 ms). This shows that the bats adjust their sensory-motor behaviour flexibly not only to the actual task, but also to various aspects of the environment, such as its spatial layout, its sound reflectivity and the available space for flight manoeuvres.

Interestingly, bats substantially increased between-group intervals and even emitted a few

search calls already before passing the opening, indicating that they already shifted their sonar gaze to

positions behind the opening. Despite this increase in call interval to allow for echoes arriving from

beyond the wall, the bats continued to reduce their call duration until about 20 cm distance (~ 1.2 ms

call duration) to avoid temporal overlap between calls and echoes from the close wall (call-echo

ambiguity; Beetz, Kössl, & Hechavarría, 2019; Moss, Chiu, & Surlykke, 2011). Thus, even in this

rather simple task (without a subsequent mealworm to catch (Surlykke et al., 2009) or a second hole to

navigate through (Falk et al., 2015), the bats used a sensory strategy for simultaneous sampling of

short- and long-range objects.

Drinking

Bats innately recognise water bodies due to their mirror-like echo reflection properties (Greif &

Siemers, 2010). To drink on the wing, bats must get low enough to reach the water with their tongue

while avoiding touching the water surface with wings or body. Simultaneously, they must keep track

of the scene ahead and plan their trajectory. Hence, drinking on the wing likely poses more sensory-

motor challenges than flying through a large opening. Indeed, bats initiated their approach much

earlier and called more frequently when approaching the pond compared to the opening. At the same

time, drinking is likely less challenging than attacking prey since the position of any water surface is

more predictable than that of a small erratically moving prey. Indeed, bats reduced call intervals less

when drinking than when landing or capturing prey, and never emitted buzz calls (<6ms inter-call-

interval), corroborating that barbastelles increase call rate less when drinking as when capturing prey

(Russo *et al.* 2016). Also, when drinking, call intervals need to be long enough to allow echoes from

distant objects ahead to return before the call is emitted, in order to plan the trajectory prospectively.

With a minimum call interval of 10.7 ms during drinking approaches, bats could receive echoes from

about 1.8 m distance without call-echo ambiguity, indicating that bats likely kept track of the edge of

the pool to avoid collisions.

When flying at close distance above water, vertical water echoes arrive with very short delay

(e.g. <0.2 ms at 3 cm above water). Most likely, bats will have difficulties to emit calls that are short

enough to avoid call-echo overlap at all under such close-range conditions, especially given that the

middle-ear reflex prevents echo detection even shortly after call emission (Henson Jr., 1965; Kick &

Simmons, 1984; Suga & Schlegel, 1972). This would explain why barbastelles ended their approach

with longer call durations when approaching the pond compared to all other tasks. Our behavioural

data thus match a spectral interference-based sensory strategy for estimating flight height above water

surfaces (Hoffmann et al., 2015): as the outgoing call and returning echo overlap in time close above

the water surface, the flight height determines position and magnitude of spectral notches generated

through interference between call and echo. Thus, when drinking, bats probably benefit from emitting

comparably long calls that allow for sufficient temporal overlap between call and echo. Our observed

initial increase in call duration, however, is likely a measurement artefact, as calls emitted at large

distance had comparably low received amplitudes and thus relatively short call duration

measurements.

The pause between event and first post-event call was largest after drinking (150 ms),

corroborating long post-drinking pauses (200-350 ms) in other species (Kloepper, Simmons, &

Simmons, 2019). Presumably, bats first need to swallow the water before being able to exhale and

open the mouth for emitting the next call. In contrast, bats might well emit these calls while holding a

prey item with their teeth. This conclusion is supported by shorter post-event pause (~50 ms) after

capture and the observation that barbastelles usually eat their prey in flight, which – depending on

prey size – can take considerable time. Interestingly, the first call(s) after capture and drinking events

where type II calls that presumably are emitted nasally – which could suggest that nasal emission is

easier than mouth emission after chewing and swallowing. However, bats also emitted type II calls

after landing, which could either indicate another reason for preferring nasal over mouth emission

while being stationary, or yet another reason common to all tasks.

Landing

Bats need to execute a complex, almost acrobatic aerial rotation to land head-under-heels (Riskin et

al., 2009). Probably, the major challenge for bats landing on a wall is to execute that manoeuvre at the

right position and right time, such that they are correctly positioned once reaching the wall. Thus,

regular and precise ranging is of paramount importance prior to landing. Accordingly, the information

update rate was higher when approaching the landing site than when approaching the drinking site.

Similarly, both tongue-clicking fruit bats (Danilovich et al., 2015) and laryngeally echolocating bats

(Hiryu, Hagino, Riquimaroux, & Watanabe, 2007b; Melcón, Denzinger, & Schnitzler, 2007; Russo et

al., 2007) increased their call rates just before landing, suggesting that high information update rates

are a general necessity for landing in animals moving fast in 3D space. Likewise, call durations during

landing were the shortest of all tasks for most part of the approach, enabling precise distance

information and little forward-masking early on during the approach (Boonman, Parsons, & Jones,

2003; Schnitzler & Kalko, 2001).

After successful landings, bats continued to call. Theoretically, this might be because we

misidentified the time of landing, but we are confident that this was not the case, even if the wall

contact was not visible in the video or the spectrogram: (i) the sound from hitting the wall was always

audible in the sound file. (ii) Also when the sound was visible in the spectrogram, and the timing thus

less ambiguous, bats continued calling after landing. (iii) At the times defined as wall contact, within-

group call intervals were shortest while between-group intervals were already increasing again (Fig.

3). In addition, also horseshoe bats continued calling after landing (Tian & Schnitzler, 1997). We

assume that bats continue calling after wall contact for multiple reasons: (i) in case landing fails,

continued sensory sampling would allow to evaluate possible 'emergency' flight trajectories. (ii) Bats

often crawled to adjacent and more sheltered sites after landing and therefore sampled the specific
surrounding of the landing spot. (iii) Lastly, bats still need to sense their general environment also
after landing.

Capturing prey

Barbastelles started their approach on average 0.7 s before prey capture, which is just slightly longer
than the 0.4 s we measured previously in the flight room with a stationary prey (Lewanzik & Goerlitz
2018) and similar to drinking and landing. Yet, only when capturing prey did the bats always emit a
buzz sequence with shortest call intervals and durations. Such high sampling rates (up to >200 calls/s)
prior to capture are ubiquitous among echolocators, including all aerial-hawking insectivorous bats
and toothed whales (Fais, Johnson, Wilson, Aguilar Soto, & Madsen, 2016; Jensen, Keller, Tyack, &
Visser, 2020; Madsen & Surlykke, 2013; Ratcliffe et al., 2013; Wisniewska et al., 2016). This suggests
(i) that prey capture is a major sensorimotor challenge for all echolocating animals foraging on prey
moving in three dimensions and (ii) that the terminal buzz is a key functional adaptation for these
animals (Ratcliffe et al., 2013). The maximum call rate measured here, 286 calls/s, is among the
highest measured to date for bats (Ratcliffe et al., 2013; Russo et al., 2016) and might be adaptive for
hunting the barbastelle's typical prey, eared moths (Goerlitz et al., 2010). In contrast to earless prey
that cannot hear and react to attacking bats, eared moths try to escape attacking bats with evasive
flight. The barbastelle's main strategy for hunting eared moths is to emit low-intensity echolocation
calls that allow it to remain undetected until close to the prey (Lewanzik & Goerlitz, 2018). The high
call rates are unlikely to guide capture until contact due to the bats' reaction time (Geberl, Brinkløv,
Wiegrebe, & Surlykke, 2015; Melcón et al., 2007). However, the high call rates might aid in tracking
prey that initiates evasive manoeuvres shortly before capture despite the low call levels, or for
retrospectively evaluating a failed attack (Melcón et al., 2007). Additionally, high call rates allow
perception of wing beat frequency (Goerlitz, Geberl & Wiegrebe 2010, Baier & Wiegebe 2018, Baier,
Stelzer & Wiegrebe 2018), which barbastelles could use to estimate prey size and select more
profitable large moths (Koselj, Schnitzler & Siemers 2011).

During unsuccessful prey capture attempts, and in contrast to successful prey captures, bats
kept the peak frequency of almost all calls in the first harmonic throughout the final approach (Fig.
5b). This difference indicates that the bats either failed to capture prey because they did not control
their sensory information appropriately, or they approached the prey using a suboptimal trajectory and
hence purposely kept most energy at lower frequencies to have a broader sonar beam, i.e. a broader
'field of view', in such trials. If bats approached the prey in a way that put them at risk to collide with
the stand, for instance, a broader beam could have allowed them to track both, prey and stand; yet
focusing only partially on the prey might have been insufficient to successfully capture it.

 General discussion

By comparing the echolocation behaviour of barbastelle bats during object approach across different tasks, we here demonstrate that bats adjusted various call parameters independently from each other in a task- and difficulty-dependent manner – as they do to improve signal detectability in noise (Luo, Goerlitz, Brumm, & Wiegrebe, 2015). Our results imply that bats adjust the acquisition of both information quantity and quality independently to the sensory constraints of each task.

Other echolocating animals, such as toothed whales, face very similar trade-offs. Despite the large differences in body size and physical conditions of their environments, all echolocating mammals resolve these trade-offs in strikingly similar ways (Madsen & Surlykke, 2013). For example, all echolocators need to emit signals frequently enough to avoid obstacles and to track prey, but suffer from echo ambiguity when the next signal is emitted before all echoes have returned. Both bats and toothed whales reduce the time interval between consecutive signals in such a way to maximise information flow while avoiding echo ambiguity when approaching objects (Jensen, Bejder, Wahlberg, & Madsen, 2009; Ladegaard, Jensen, Beedholm, da Silva, & Madsen, 2017; Melcón et al., 2007). Both groups also emit a final buzz for fine-scale tracking prior to capture and reduce output intensity during the approach to reduce the complexity of their auditory scene (Jensen et al., 2020; Ladegaard et al., 2017; Madsen & Surlykke, 2013). Thus, all echolocating animals actively control acoustic information flow and their sensory volume to guide subsequent actions. Here, we confirm that this active control is not connected in a general time- or distant-dependent manner to the approached object, but is finely adjusted in a task-specific way to four different tasks differing in sensory-motor difficulty.

Not only echolocators but every animal including humans needs to constantly acquire information about its environment to make appropriate motor decisions when moving about and interacting with others (Dall et al., 2005). Under some circumstances, social information provided by others might be exploited (Lewanzik et al., 2019; Valone & Templeton, 2002), but social information cannot completely substitute personal information. This study contributes to our understanding of how animals adjust the acquisition of personal information to deal with the constraints they face when sensing the world sequentially through sound. Our results show that they increase information quantity (call rate) only to a level as required by the task but not above that, likely to not minimize the costs required for vocal production (Elemans et al., 2011; Currie et al 2020) and sensory analysis (Laughlin et al., 1998; Niven & Laughlin, 2008) at high update rates. In addition, by flexibly adjusting call features, such as duration and frequency, bats also adapt the quality of the acquired information. Hence, our study highlights the importance of behavioural flexibility to optimize the cost-benefit ratio of sensing and the quantity and quality of sensory information.

Acknowledgements

We thank the authorities for granting capture and experimental permits (55.1-8646.1-1 and 55.2-1-54-2532-18-2015, respectively) and the Max Planck Institute for Ornithology for providing facilities and logistic support. Particularly, we thank Christian Völk and Felix Hartl for valuable logistic support. Further, we are grateful to all members of the Acoustic and Functional Ecology Group for constructive feedback on earlier versions of the manuscript.

Competing interests

We declare that we have no competing interests.

Funding

This research was funded through an Emmy Noether grant to H.R.G. by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, grant # 241711556).

Data availability statement

Data will be made available at the Dryad Digital Repository prior to publication.

Authors' contributions

H.R.G. and D.L. conceived the idea and designed methodology; D.L. collected and analysed the data and led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

References

Barchi, J. R., Knowles, J. M., & Simmons, J. A. (2013). Spatial memory and stereotypy of flight paths

by big brown bats in cluttered surroundings. *Journal of Experimental Biology*, *216*(6), 1053–

1063. doi: 10.1242/jeb.073197

Baier, A. L., & Wiegrebe, L. (2018). Flutter sensitivity in FM bats. Part I: delay modulation. *Journal*

of Comparative Physiology A, 204, 929-939.

Baier, A. L., Stelzer, K.-J., & Wiegrebe, L. (2018). Flutter sensitivity in FM bats. Part II: amplitude

modulation. *Journal of Comparative Physiology A*, 204, 941-951.

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using

lme4. *Journal of Statistical Software*, *67*, 1–48.

Bates, M. E., Simmons, J. A., & Zorikov, T. V. (2011). Bats use echo harmonic structure to

distinguish their targets from background clutter. *Science*, *333*(6042), 627–630. doi:

10.1126/science.1202065

Beetz, M. J., Hechavarría, J. C., & Kössl, M. (2016). Cortical neurons of bats respond best to echoes

from nearest targets when listening to natural biosonar multi-echo streams. *Scientific Reports*,

6(1), 1–12. doi: 10.1038/srep35991

Beetz, M. J., Kössl, M., & Hechavarría, J. C. (2019). Adaptations in the call emission pattern of

frugivorous bats when orienting under challenging conditions. *Journal of Comparative*

Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology, *205*(4), 457–467.

doi: 10.1007/s00359-019-01337-1

Boonman, A. M., Parsons, S., & Jones, G. (2003). The influence of flight speed on the ranging

performance of bats using frequency modulated echolocation pulses. *The Journal of the*

Acoustical Society of America, *113*(1), 617–628. doi: 10.1121/1.1528175

Currie, S. E., Boonman, A., Troxell, S., Yovel, Y., & Voigt, C. C. (2020). Echolocation at high

intensity imposes metabolic costs on flying bats. *Nature Ecology and Evolution*, 1–4. doi:

10.1038/s41559-020-1249-8

Dall, S. R. X., Giraldeau, L. A., Olsson, O., McNamara, J. M., & Stephens, D. W. (2005). Information

and its use by animals in evolutionary ecology. *Trends in Ecology and Evolution*, *20*(4), 187–

193. doi: 10.1016/j.tree.2005.01.010

Danilovich, S., Krishnan, A., Lee, W.-J., Borrisov, I., Eitan, O., Kosa, G., … Yovel, Y. (2015). Bats

regulate biosonar based on the availability of visual information. *Current Biology*, *25*(23),

R1124–R1125. doi: 10.1016/J.CUB.2015.11.003

Denzinger, A., Siemers, B. M., Schaub, A., & Schnitzler, H. U. (2001). Echolocation by the

barbastelle bat, *Barbastella barbastellus*. *Journal of Comparative Physiology A*, *187*(7), 521–

528. doi: 10.1007/s003590100223

Dietz, C., von Helversen, O., & Nill, D. (2007). *Handbuch der Fledermäuse Europas und*

Nordwestafrikas. Stuttgart, Germany: Franckh-Kosmos Verlag.

Elemans, C. P. H., Mead, A. F., Jakobsen, L., & Ratcliffe, J. M. (2011). Superfast muscles set

maximum call rate in echolocating bats. *Science*, *333*(6051), 1885–1888. doi:

10.1126/science.1207309

Fais, A., Johnson, M., Wilson, M., Aguilar Soto, N., & Madsen, P. T. (2016). Sperm whale predator-

prey interactions involve chasing and buzzing, but no acoustic stunning. *Scientific Reports*, *6*(1),

1–13. doi: 10.1038/srep28562

Falk, B., Kasnadi, J., & Moss, C. F. (2015). Tight coordination of aerial flight maneuvers and sonar

call production in insectivorous bats. *The Journal of Experimental Biology*, *218*, 3678–3688. doi:

10.1242/jeb.122283

Fenton, B., Jensen, F. H., Kalko, E. K. V., & Tyack, P. L. (2014). Sonar Signals of Bats and Toothed

Whales. In A. Surlykke, P. E. Nachtigall, R. R. Fay, & A. Popper (Eds.), *Springer handbook of*

auditory research - Biosonar (pp. 11–59). doi: 10.1007/978-1-4614-9146-0_2

Geberl, C., Brinkløv, S., Wiegrebe, L., & Surlykke, A. (2015). Fast sensory-motor reactions in

echolocating bats to sudden changes during the final buzz and prey intercept. *Proceedings of the*

National Academy of Sciences of the United States of America, *112*(13), 4122–4127. doi:

10.1073/pnas.1424457112

Geipel, I., Steckel, J., Tschapka, M., Vanderelst, D., Schnitzler, H. U., Kalko, E. K. V., … Simon, R.

(2019). Bats actively use leaves as specular reflectors to detect acoustically camouflaged prey.

Current Biology, *29*(16), 2731-2736.e3. doi: 10.1016/j.cub.2019.06.076

Goerlitz, H. R., Geberl, C., & Wiegrebe, L. (2010). Sonar detection of jittering real targets in a free-

flying bat. *The Journal of the Acoustical Society of America*, 128(3), 1467-1475.

Goerlitz, H. R., Ter Hofstede, H. M., Zeale, M. R. K., Jones, G., & Holderied, M. W. (2010). An

aerial-hawking bat uses stealth echolocation to counter moth hearing. *Current Biology*, *20*(17),

1568–1572. doi: 10.1016/j.cub.2010.07.046

Gomes, D. G. E., Page, R. A., Geipel, I., Taylor, R. C., Ryan, M. J., & Halfwerk, W. (2016). Bats

perceptually weight prey cues across sensory systems when hunting in noise. *Science*,

353(6305),1277-1280.

Greif, S., & Siemers, B. M. (2010). Innate recognition of water bodies in echolocating bats. *Nature*

Communications, *1*(1), 107. doi: 10.1038/ncomms1110

Griffin, D. R. (1958). *Listening in the dark*. New Haven, Connecticut: Yale University Press.

Henson Jr., O. W. (1965). The activity and function of the middle ear muscles in echolocating bats.

Journal of Physiology, *180*, 871–887. doi: 10.1113/jphysiol.1965.sp007737

Hiryu, S., Hagino, T., Riquimaroux, H., & Watanabe, Y. (2007a). Echo-intensity compensation in

echolocating bats (Pipistrellus abramus) during flight measured by a telemetry microphone. *The*

Journal of the Acoustical Society of America, *121*(3), 1749–1757. doi: 10.1121/1.2431337

Hiryu, S., Hagino, T., Riquimaroux, H., & Watanabe, Y. (2007b). Echo-intensity compensation in

echolocating bats (*Pipistrellus abramus*) during flight measured by a telemetry microphone.

The Journal of the Acoustical Society of America, *121*(3), 1749–1757. doi: 10.1121/1.2431337

Hoffmann, S., Genzel, D., Prosch, S., Baier, L., Weser, S., Wiegrebe, L., & Firzlaff, U. (2015). Biosonar navigation above water I: estimating flight height. *Journal of Neurophysiology*, *113*(4), 1135–1145. doi: 10.1152/jn.00263.2014

Jensen, F. H., Bejder, L., Wahlberg, M., & Madsen, P. T. (2009). Biosonar adjustments to target range of echolocating bottlenose dolphins (*Tursiops sp.*) in the wild. *Journal of Experimental Biology*, *212*(8), 1078–1086. doi: 10.1242/jeb.025619

Jensen, Frants H., Keller, O. A., Tyack, P. L., & Visser, F. (2020). Dynamic biosonar adjustment strategies in deep-diving Risso's dolphins driven partly by prey evasion. *The Journal of Experimental Biology*, *223*(3). doi: 10.1242/jeb.216283

Kalko, E. K. V., & Schnitzler, H.-U. (1989). The echolocation and hunting behavior of Daubenton's bat, *Myotis daubentoni*. *Behavioral Ecology and Sociobiology*, *24*(4), 225–238. doi: 10.1007/BF00295202

Kerth, G., & Melber, M. (2009). Species-specific barrier effects of a motorway on the habitat use of two threatened forest-living bat species. *Biological Conservation*, *142*(2), 270–279. doi: 10.1016/j.biocon.2008.10.022

Kick, S. A., & Simmons, J. A. (1984). Automatic gain control in the bat's sonar receiver and the neuroethology of echolocation. *The Journal of Neuroscience*, *4*(11), 2725–2737. doi: 10.1523/JNEUROSCI.04-11-02725.1984

Kloepper, L. N., Simmons, A. M., & Simmons, J. A. (2019). Echolocation while drinking: Pulse-timing strategies by high- And low-frequency FM bats. *PLoS ONE*, *14*(12), 1–16. doi: 10.1371/journal.pone.0226114

Koblitz, J. C., Stilz, P., Pflästerer, W., Melcón, M. L., & Schnitzler, H.-U. (2011). Source level reduction and sonar beam aiming in landing big brown bats (*Eptesicus fuscus*). *The Journal of the Acoustical Society of America*, *130*(5), 3090–3099. doi: 10.1121/1.3628345

Koselj, K., Schnitzler, H.-U., & Siemers, B. M. (2011). Horseshoe bats make adaptive prey-selection deicsions, informed by echo cues. *Proceedings of the Royal Society of London B*, 278, 3034-3041.

Kugler, K. & Wiegrebe, L. (2017). Echo-acoustic scanning with noseleaf and ears in phyllostomid bats. *The Journal of Experimental Biology*, 220, 2816-2824.

Kurnikova, A., Moore, J. D., Song-Mao, L., Deschênes, M., & Kleinfeld, D. (2017). Coordination of orofacial motor actions into exploratory behavior by rat. *Current Biology* (27), 688-696.

Ladegaard, M., & Madsen, P. T. (2019). Context-dependent biosonar adjustments during active target approaches in echolocating harbour porpoises. *The Journal of Experimental Biology*, 222, jeb.206169.

Ladegaard, M., Jensen, F. H., Beedholm, K., da Silva, V. M. F., Madsen, P. T., Silva, V. M. F. da, & Madsen, P. T. (2017). Amazon river dolphins (Inia geoffrensis) modify biosonar output level and

directivity during prey interception in the wild. *The Journal of Experimental Biology*, *220*, 2654–2665. doi: 10.1242/jeb.159913

Laughlin, S. B., de Ruyter van Steveninck, R. R., & Anderson, J. C. (1998). The metabolic cost of neural information. *Nature Neuroscience*, *1*(1), 36–41. doi: 10.1038/236

Lavie, N. (2005). Distracted and confused?: Selective attention under load. *Trends in Cognitive Sciences*, *9*(2), 75–82. doi: 10.1016/j.tics.2004.12.004

Lewanzik, D., & Goerlitz, H. R. (2018). Continued source level reduction during attack in the low-amplitude bat *Barbastella barbastellus* prevents moth evasive flight. *Functional Ecology*, *32*(5), 1251–1261. doi: 10.1111/1365-2435.13073

Lewanzik, D., Sundaramurthy, A. K., & Goerlitz, H. R. (2019). Insectivorous bats integrate social information about species identity, conspecific activity and prey abundance to estimate cost–benefit ratio of interactions. *Journal of Animal Ecology*, 1365-2656.12989. doi: 10.1111/1365-2656.12989

Luo, J., Goerlitz, H. R., Brumm, H., & Wiegrebe, L. (2015). Linking the sender to the receiver: vocal adjustments by bats to maintain signal detection in noise. *Scientific Reports*, *5*, 18556. doi: 10.1038/srep18556

Madsen, P. T., & Surlykke, A. (2013). Functional convergence in bat and toothed whale biosonars. *Physiology*, *28*, 276–283. doi: 10.1152/physiol.00008.2013

Melcón, M. L., Denzinger, A., & Schnitzler, H.-U. (2007). Aerial hawking and landing: approach behaviour in Natterer's bats, *Myotis nattereri* (Kuhl 1818). *The Journal of Experimental Biology*, *210*, 4457–4464. doi: 10.1242/jeb.007435

Morshedian, A., & Fain, G. L. (2015). Single-photon sensitivity of lamprey rods with cone-like outer segments. *Current Biology*, *25*(4), 484–487. doi: 10.1016/j.cub.2014.12.031

Moss, C. F., Chiu, C., & Surlykke, A. (2011). Adaptive vocal behavior drives perception by echolocation in bats. *Current Opinion in Neurobiology*, *21*, 645–652. doi: 10.1016/J.CONB.2011.05.028

Niven, J. E., & Laughlin, S. B. (2008). Energy limitation as a selective pressure on the evolution of sensory systems. *The Journal of Experimental Biology*, *211*, 1792–1804. doi: 10.1242/jeb.017574

Prat, Y., & Yovel, Y. (2020, February 1). Decision making in foraging bats. *Current Opinion in Neurobiology*, Vol. 60, pp. 169–175. doi: 10.1016/j.conb.2019.12.006

R Core Team. (2019). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.

Ratcliffe, J. M., Elemans, C. P. H., Jakobsen, L., & Surlykke, A. (2013). How the bat got its buzz. *Biology Letters*, *9*(2), 20121031–20121031. doi: 10.1098/rsbl.2012.1031

Ratcliffe, John M., & Dawson, J. W. (2003). Behavioural flexibility: The little brown bat, Myotis lucifugus, and the northern long-eared bat, M. septentrionalis, both glean and hawk prey. *Animal*

Behaviour, *66*(5), 847–856. doi: 10.1006/anbe.2003.2297

Riskin, D. K., Bahlman, J. W., Hubel, T. Y., Ratcliffe, J. M., Kunz, T. H., & Swartz, S. M. (2009). Bats go head-under-heels: The biomechanics of landing on a ceiling. *Journal of Experimental Biology*, *212*(7), 945–953. doi: 10.1242/jeb.026161

Russo, D., Ancillotto, L., Cistrone, L., & Korine, C. (2016). The buzz of drinking on the wing in echolocating bats. *Ethology*, *122*(3), 226–235. doi: 10.1111/eth.12460

Russo, D., Jones, G., & Arlettaz, R. (2007). Echolocation and passive listening by foraging mouse-eared bats *Myotis myotis* and *M. blythii*. *The Journal of Experimental Biology*, *210*, 166–176. doi: 10.1242/jeb.02644

Schnitzler, H.-U., & Kalko, E. K. V. (2001). Echolocation by insect-eating bats. *BioScience*, *51*(7), 557–569. doi: 10.1641/0006-3568(2001)051[0557:EBIEB]2.0.CO;2

Schnitzler, H.-U., Moss, C. F., & Denzinger, A. (2003). From spatial orientation to food acquisition in echolocating bats. *Trends in Ecology & Evolution*, *18*(8), 386–394. doi: 10.1016/S0169-5347(03)00185-X

Seibert, A. M., Koblitz, J. C., Denzinger, A., & Schnitzler, H. U. (2015). Bidirectional echolocation in the bat *Barbastella barbastellus*: different signals of low source level are emitted upward through the nose and downward through the mouth. *PLoS ONE*, *10*(9), 1–17. doi: 10.1371/journal.pone.0135590

Simmons, J. A., Brown, P. E., Vargas-Irwin, C. E., & Simmons, A. M. (2020). Big brown bats are challenged by acoustically-guided flights through a circular tunnel of hoops. *Scientific Reports*, *10*(1), 1–10. doi: 10.1038/s41598-020-57632-4

Skiba, R. (2009). *Europäische Fledermäuse* (2nd editio). Hohenwarsleben: Westarp Wissenschaften.

Sridharan, D., Schwarz, J. S., & Knudsen, E. I. (2014). Selective attention in birds. *Current Biology*, *24*(11). doi: 10.1016/j.cub.2013.12.046

Suga, N., & Schlegel, P. (1972). Neural attenuation of responses to emitted sounds in echolocating bats. *Science*, *177*, 82–84.

Surlykke, A., Ghose, K., & Moss, C. F. (2009). Acoustic scanning of natural scenes by echolocation in the big brown bat, *Eptesicus fuscus*. *Journal of Experimental Biology*, *212*, 1011–1020. doi: 10.1242/jeb.024620

Taub, M., & Yovel, Y. (2020). Segregating signal from noise through movement in echolocating bats. *Scientific Reports*, *10*(1), 382. doi: 10.1038/s41598-019-57346-2

Thiele, J., & Winter, Y. (2005). Hierarchical strategy for relocating food targets in flower bats: Spatial memory versus cue-directed search. *Animal Behaviour*, *69*(2), 315–327. doi: 10.1016/j.anbehav.2004.05.012

Tian, B., & Schnitzler, H.-U. (1997). Echolocation signals of the Greater Horseshoe bat (*Rhinolophus ferrumequinum*) in transfer flight and during landing. *The Journal of the Acoustical Society of America*, *101*(4), 2347–2364. doi: 10.1121/1.418272

Valone, T. J., & Templeton, J. J. (2002). Public information for the assessment of quality: a
widespread social phenomenon. *Philosophical Transactions of the Royal Society B*, *357*(1427),
1549–1557. doi: 10.1098/rstb.2002.1064

Wahlberg, M., & Surlykke, A. (2014). Sound Intensities of Biosonar Signals from Bats and Toothed
Whales. In A. Surlykke, P. E. Nachtigall, R. R. Fay, & A. Popper (Eds.), *Springer handbook of*
auditory research - Biosonar (pp. 107–141). doi: 10.1007/978-1-4614-9146-0_4

Waters, D. A., & Jones, G. (1994). Wingbeat-generated ultrasound in noctuid moths increases the
discharge rate of the bat-detecting A1 cell. *Proceedings of the Royal Society B: Biological*
Sciences, *258*(1351), 41–46. doi: 10.1098/rspb.1994.0139

Wheeler, A. R., Fulton, K. A., Gaudette, J. E., Simmons, R. A., Matsuo, I., & Simmons, J. A. (2016).
Echolocating Big Brown Bats, Eptesicus fuscus, Modulate Pulse Intervals to Overcome Range
Ambiguity in Cluttered Surroundings. *Frontiers in Behavioral Neuroscience*, *10*(Jun), 125. doi:
10.3389/fnbeh.2016.00125

Wilson, D. E., & Mittermeier, R. A. (2019). *Handbook of the mammals of the world Vol. 9: Bats*.
Barcelona, Spain: Lynx Edicions.

Wilson, W. W., & Moss, C. F. (2004). Sensory-motor behavior of free-flying FM bats during target
capture. In J. A. Thomas, C. F. Moss, & M. Vater (Eds.), *Echolocation in bats and dolphins* (pp.
22–27). Chicago, Illinois, USA: Chicago University Press.

Wisniewska, D. M., Johnson, M., Teilmann, J., Rojano-Donate, L., Shearer, J., Sveegaard, S., …
Madsen, P. T. (2016). Ultra-high foraging rates of harbor porpoises make them vulnerable to
anthropogenic disturbance. *Current Biology*, *26*, 1–6. doi: 10.1016/j.cub.2016.03.069

Yarbus, A. L. (1967). Eye movements and vision. New York, Plenum Press.

Yamada, Y., Mibe, Y., Yamamoto, Y., Ito, K., Heim, O., & Hiryu, S. (2020). Modulation of acoustic
navigation behaviour by spatial learning in the echolocating bat Rhinolophus ferrumequinum
nippon. *Scientific Reports*, *10*(1), 10751. doi: 10.1038/s41598-020-67470-z

Zeale, M. R. K., Davidson-Watts, I., & Jones, G. (2012). Home range use and habitat selection by
barbastelle bats (*Barbastella barbastellus*): implications for conservation. *Journal of*
Mammalogy, *93*(4), 1110–1118. doi: 10.1644/11-MAMM-A-366.1