Common synaptic phenotypes arising from diverse mutations in the human NMDA receptor subunit GluN2A

Elmasri, M.[1], Hunter, D.[1], Winchester, G. [1], Bates, E. [1], Aziz, W.[1], Moolenaar Van Der Does, D.[1], Karachaliou, E.[1], Sakimura, K.[2] and Penn, A.C.[1,*]

[1.] Sussex Neuroscience, School of Life Sciences, University of Sussex, Brighton, BN1 9QG, United Kingdom.

[2.] Department of Cellular Neurobiology, Brain Research Institute, Niigata University, Niigata, 951-8585, Japan.

* Corresponding author

Abstract

Dominant mutations in the human gene *GRIN2A*, encoding NMDA receptor (NMDAR) subunit GluN2A, make a significant and growing contribution to the catalogue of published single-gene epilepsies. Understanding the disease mechanism in these epilepsy patients is complicated by the surprising diversity of effects that the mutations have on NMDARs. We have examined the cell-autonomous effect of five GluN2A mutations, 3 loss-of-function and 2 gain-of-function, on evoked NMDAR-mediated synaptic currents (NMDA-EPSCs) in CA1 pyramidal neurons in cultured hippocampal slices. Despite the mutants differing in their functional incorporation at synapses, prolonged NMDA-EPSC current decays (with only marginal changes in charge transfer) were a common effect for both gain- and loss-of-function mutants. Modelling NMDA-EPSCs with mutant properties in a CA1 neuron revealed that the effect of *GRIN2A* mutations can lead to abnormal temporal integration and spine calcium dynamics during trains of concerted synaptic activity. Investigations beyond establishing the molecular defects of GluN2A mutants are much needed to understand their impact on synaptic transmission.

1 *Introduction*

N-methyl-D-aspartate receptors (NMDARs) are conserved synaptic proteins underling the

late component of postsynaptic potentials at excitatory synapses[1]. The receptor plays key

roles in sculpting nervous system development[2–4] and spine morphology[5], shaping temporal

integration of synaptic potentials[6–8], regulating synaptic efficacy[9–11], and the formation of

new declarative memories[12]. Core to NMDARs achieving these roles is their high-calcium

permeability, depolarization-dependent unblock of extracellular magnesium ions from the

channel pore, and the slow time course of NMDAR channel closure (deactivation) following

brief exposure to neurotransmitter glutamate[1,10,13]. These properties are determined to a

large extent by subunit composition and stoichiometry of NMDARs[14]. In adulthood,

conventional NMDARs are heterotetramers composed of obligatory GluN1 subunits and

regulatory GluN2 subunits, while unconventional GluN1/3 heterotetramers are transiently

expressed during early postnatal development[10,15]. The GluN2 subunits contain the ligand-

binding site indispensable for the glutamate-gated operation of NMDARs at excitatory

synapses[16]. Remarkably, four GluN2 subunits, suffixed by A-D, account for much of the

diversity observed for a wide range of functional characteristics of NMDARs[10,17,18].

NMDAR subunit expression varies between neuronal types and during development thus

providing of means of adapting the NMDAR-mediated component of excitatory postsynaptic

potentials (EPSPs) to the physiological context. Principal neurons account for the majority of

neurons in the adult forebrain, where they predominantly express GluN2A and 2B, and to a

much lesser extent GluN2C or D[19–21]. While GluN2B predominates at birth and continues to

make a major contribution to synaptic NMDARs into adulthood, GluN2A is upregulated

during early postnatal development[19,22–25], by neuronal activity[26–28] and sensory

experience[29–32], and accounts for much of the observed developmental speeding of NMDAR-

mediated excitatory postsynaptic currents (NMDA-EPSCs)[22,23]. Mice with germline genetic

knockout of the GluN2A gene (called *grin2a*) are incapable of undergoing the subunit switch

and have impaired spatial learning[33], altered vocalizations and spontaneous epileptiform

discharges[34,35]. Their learning defects are also characterised by raised thresholds for the

induction of contextual fear learning and long term synaptic potentiation[36].

The significance of NMDARs is well-demonstrated by the range of clinical phenotypes of patients with mutations in the genes encoding NMDAR subunits. Since the development of next generation sequencing methods in the mid-1990's there has been a steady rise in number of epilepsy-associated genes[37] with the addition of the human GluN2A receptor gene, *GRIN2A*, to this list only a decade ago[38–40]. Published figures recently showed that >250 patients are known to have *GRIN2A* mutations, either *de novo* or transmitted with dominant inheritance[41,42]. Epilepsy occurs in about 90% of these patients[42,43], with *GRIN2A* mutations representing the best known genetic cause of epilepsy-aphasia spectrum (EAS) disorders, where they account for 9-20% of cases[41,44–48]. The clinical phenotype of EAS varies considerably from the more common and benign childhood (Rolandic) epilepsy with centrotemporal spikes (CECTS), to the rarer and more severe Landau-Kleffner syndrome (LKS) and continuous spike-and-wave during slow wave sleep syndrome (CSWSS)[41,49]. Although rare, the severe epilepsy cases are often also accompanied by progressive cerebral dysfunction (epileptic encephalopathy) and are frequently comorbid with intellectual disability and language disorders[42]. The response to existing treatments in patients with severe EAS can be unpredictable and no anti-epileptic drug (AED) has been demonstrated to be effective specifically in *GRIN2A*-related disorder[41].

Most research on *GRIN2A* mutations has investigated the effects of missense mutations on GluN2A protein function in heterologous expression systems[44,46,50–57]. A common outcome of these studies is that epilepsy-associated *GRIN2A* mutations have strikingly variable functional consequences on NMDARs[50,51,56,58]. Until recently, the impact of GluN2A mutations on the NMDAR-mediated component of synaptic transmission has received relatively little attention[59]. While modelling of synaptic charge transfer has been made for some *GRIN2A* mutations[56], these models do not consider how expressing the mutant GluN2A receptor could feedback on synaptic transmission[7], or to what extent and how other native NMDAR subunits continue to support NMDAR-mediated synaptic transmission[22]. In this study, we have evaluated and compared the phenotypes and mechanisms of synaptic dysfunction in CA1 neurons expressing epilepsy-associated gain- and loss-of function *GRIN2A* mutants.

Results

Contrasting gain- and loss-of-function epilepsy-associated missense mutations in GRIN2A

More than 40% of mutations discovered in *GRIN2A* are missense mutations (Figure 1a), with the majority of them (approx. 60%) being in the sequence encoding the ligand binding domain (LBD) or the ion channel[42]. Missense mutations cause substitutions of amino acids and can vary considerably in their consequence on protein function and expression[44,46,50–53,53–57]. Considerable experimental data has been published characterizing how disease-linked mutations in *GRIN2A* affect the surface expression and biophysical properties of GluN2A in heterologous expression systems, namely in HEK cells and *Xenopus* Oocytes. We compiled data for 20 different mutations from various publications and databases[44,46,50–57] (Figure 1b: blue spheres). The most comprehensive sets of measurements have been made for mutations in the LBD and its linkers to the ion channel, which are the fundamental operating units for glutamate-gated opening of the channel[60,61] (Figure 1b). The measured variables graphed in Figure 1c reflect receptor activation (agonist potencies), channel activity (open probability and current deactivation), surface expression, or a combination of the above (current density). The effects of the mutations were up to 4 orders magnitude different from WT receptors with the effects being generally larger for glutamate potency and current density (Figure 1c). Loss-of-function effects were most frequently observed for current density (79%), surface expression (77%) and open probability (63%), while receptor deactivation time was the only receptor property to be dominated by gain-of-function effects (59%) (Figure 1c).

Although mutation effects can be easily identified for specific receptor properties (Figure 1c), overall classification of mutations can become difficult as we illustrate with some examples. The R518H mutation was identified in a patient with CSWSS and was initially described as a gain-of-function mutation based on the longer open time of the channel in single-channel recordings[46]. Since then, others have characterized the mutation further and reported that the mutant is poorly expressed on the cell surface and gives negligible whole-cell currents[55,56], which raises uncertainty on the overall effect of this mutation. Another complicated example is the case of K669N. This mutant expresses normally and exhibits a 3-

4-fold higher agonist potency and deactivation time constant compared to WT receptors[56].

However, the open probability of the channel is estimated to be 4-5 times lower making the

overall impact of this mutation also unclear.

The examples described above motivated us to adopt factor analysis to assist in interpreting

the data set of GluN2A mutation effects (Figure 1d, e). Principle component analysis

provides a means to reduce multivariate data down to fewer, more manageable variables

(components). The first two components calculated from the 6 variables collectively

accounted for the majority of the variance (71 + 16 = 87 %). After extraction and oblique

rotation of these components, the first and second components (i.e. factors) had the highest

loadings for glutamate potency ($1/EC_{50}$) and current density respectively (Figure 1d).

Consistent with previous observations that GluN2 mutants with low glutamate potency also

tend to exhibit poor surface expression[56] as we found a moderate but significant positive

correlation between the factors 1 and 2 (Figure 1e).

The fragmented bivariate distribution of factor scores for the mutations suggests possible

segregation of the mutants into different classes (Figure 1e). Mutations with similar effects

were classified by k-means clustering. The four clusters identified were: a) gain-of-function

mutants for factor 1 (4/20), referred to as GOF type 1; b) loss-of-function mutants mostly for

the factor 1 (3/20), referred to as LOF type 1; c) severe loss-of-function mutants mostly for

factor 2 (3/20), referred to as LOF type 2; and d) WT-like mutations with overall little effect

in one or both factors (11/20); (Figure 1e). While the clustering of mutations is rather

arbitrary and the wider population of mutations may exist rather as a continuum, the

analysis serves to summarize and broadly classify this set of mutations. Given some of the

assumptions and ambiguity in estimating glutamate potency for R518H and T531M (see

methods), their cluster membership may not be accurate but it is clear that these represent

mutations with a severe LOF phenotype. We cloned representative mutations from the

different clusters that had most extensive functional characterisation that was published at

the time (labelled in Figure 1e), and we went on to examine their functional incorporation at

synapses.

1 *Functional incorporation of GOF but not LOF GluN2A mutants at synapses in CA1*

2 *neurons*

To assess functional incorporation of mutant GluN2A at excitatory synapses we have

adopted a molecular replacement strategy in mouse organotypic hippocampal slices[62]. CA1

neurons in slices from adult mice are reported to express GluN2A and GluN2B subunits and

their combined deletion effectively depletes excitatory synapses of NMDA receptors[22].

Conditional deletion of the native mouse *grin2a* and *b* alleles was achieved by crossing

grin2a^{fl/fl} grin2b^{fl/fl} mouse lines . In these mice, loxP sites are orientated as direct repeats and

inserted into introns flanking exons that encode critical structural components of

receptor[22,63]. This manipulation predisposes the intervening region to deletion by Cre-

recombinase[64]. We prepared slices of hippocampus from double homozygote neonatal mice

(*grin2a^{fl/fl} grin2b^{fl/fl}*) and cultured them *in vitro* for a week before co-transfecting individual

CA1 neurons with cDNA for Cre recombinase (fused in-frame with Green Fluorescent

Protein, Cre-GFP) together with wildtype or mutant human *GRIN2A* cDNA. Slices were

cultured for a further 1.5 weeks before performing recordings of evoked NMDA-mediated

synaptic currents simultaneously from transfected and untransfected neurons at 15-20 DIV

(Figure 2a).

Using the experimental paradigm described above, we were able to compare the amplitudes

of NMDA-EPSCs both within pairs of transfected and untransfected neurons, and the

statistical interaction for the NMDA-EPSC properties between the GluN2A mutants (see

Experimental procedures). Cre-GFP alone reduced the peak NMDA-EPSCs to 12% of the peak

amplitude in neighbouring untransfected neurons (95% *CI* [9, 17]) (Figure 2b: DKO) indicating

that the NMDA-EPSCs were largely mediated by GluN2A and/or GluN2B two subunits. With

the further addition of DPQ-1105 (10 µM), a selective antagonist of GluN2C/D, NMDA-EPSCs

were barely resolvable above baseline noise with the peak amplitudes, being 6% that of

neighbouring untransfected neuons (95% *CI* [4, 9]) (Figure 2b: DKO-DPQ). Since most of the

current was mediated by GluN2A/B, we continued with experiments to rescue NMDA-EPSCs

from the GluN2A/B double knockout (DKO) neurons using mutant human GluN2A variants.

We found strong evidence in favour of an effect of expressing different *GRIN2A* mutants on

NMDA-EPSC amplitudes (Figure S1.1, $F(5,87)=11.4$, $p <.001$, $BF_{10} = 4.14 \times 10^6$). Overall, the

mutants differed significantly from WT in their ability to rescue NMDA-EPSCs (Figure 2c and

S1.1, $F_{(1,87)}=31.2$, $p<.001$) and we could resolve a significant difference between putative

LOF (C436R, T531M, R518H) and GOF (K669N and L812M) mutations (Figure 2c and S1.1,

$F_{(1,87)}=25.9$, $p <.001$). The NMDA-EPSC peak amplitudes for LOF mutants relative to WT

(100%) were 35% for C436R, 38% for T531M and 37% for R518H, 95% *CIs* [25, 49], [25, 57]

and [26, 53], respectively (Figure 2c). In contrast, the K669N and L812M exhibited more

effective functional rescue of the peak NMDA-EPSC, them being 64% for both K669N and

L812M, 95% *CIs* [46, 89] and [46, 91], respectively (Figure 2c). Together, this data shows that

the GOF GluN2A mutants were more effective (than LOF GluN2A mutants) at rescuing NMDA

function at synapses in neurons largely devoid of native GluN2 expression.

The GOF mutants tested were previously reported to have much slower deactivation time

constants when expressed in heterologous expression systems[56,57]. We next examined the

decay time constants of the NMDA-EPSCs from the GOF mutations and compared these to

WT and GluN2A and the more slowly deactivating GluN2B subunit (Figure 2d and S1.2).

Compared to WT (100%), the decay of NMDA-EPSCs in neurons rescued with K669N was

184%, 95% *CI* [146, 231], whereas L812M was 492%, 95% *CI* [386, 625], which is more like

the slower NMDA-EPSC decay observed in neurons rescued with GluN2B WT (Figure 2d). In

summary, these rescue experiments broadly support the mutation classification determined

by analysis of heterologous expression data and presented in Figure 1[56,57].

GRIN2A loss-of-function mutants vary in their ability to deliver NMDA receptors to
synapses

To test whether the poor functional incorporation of LOF mutants reflected an absence of

the mutant NMDARs at synapses, we adopted a simple assay that takes advantage of the

obligatory assembly of GluN1 subunits into NMDARs[26] and the ER-retention of unassembled

GluN1 in neurons[65]. GFP (or its pH-sensitive variant superecliptic pHlourin (SEP)) fused to the

extracellular (or ER-lumenal) amino terminus of GluN1 has previously been used to

investigate trafficking competence of different GluN2 variants to synapses[26,54,66]. Exogenous

expression of SEP-GluN1 alone in cultured hippocampal neurons gave largely dendritic shaft

fluorescence (Figure 3a). However, co-expression with WT GluN2A drove GluN1 into the cell

membrane within dendritic spines, as evidenced by an almost 2-fold enrichment of SEP-

GluN1 at spine heads colocalizing with a postsynaptic density marker, Homer1C-tdTomato.

We rationalized that GluN2 mutations that disrupt the normal enrichment of NMDARs at

synapses would also be incompetent at driving SEP-GluN1 to synapses. Indeed, we found

strong evidence in favour of an effect of expressing different *GRIN2A* mutants on the spine

enrichment of NMDA receptors relative to colocalized Homer1c (Figure S1.3, $F(5,185)=10.3$,

$p < .001$, $BF_{10} = 4.82 \times 10^4$). Significant differences could be resolved for all orthogonal

contrasts indicating that mutations differed within, as well as between, mutation types

(Figures 3 and S1.3. Synaptic levels of NMDARs in cells expressing the C436R GluN2A mutant

were 56% that of WT , 95% *CI* [46, 68], similar to that seen in neurons not co-transfected

with any GluN2A (Figure 3bii). On the other hand, GluN2A T531M and L812M exhibited

intermediate levels of SEP-GluN1 spine fluorescence (75% and 74%, 95% CIs [64, 88] and [60,

92], respectively), whereas K669N efficiently rescued spine SEP-GluN1 levels to 109% of WT,

95% CI [90, 132]. Most striking of all, despite the poor functional rescue we observed earlier

for R518H, expression of this mutant restored NMDA receptor levels in spines to 97% that of

WT, 95% CI [82, 114]. Together, this suggests that R518H loss-of-function arises from

functional defects while C436R (and to some extent T531M) exhibit defective enrichment at

synapses. We note that for the L812M mutant, transfected cell density appeared very low in

cell counts and was rescued by incubating the cells after transfection with NMDA receptor

blocker memantine, which suggests that overexpressing GluN2A L812M could be neurotoxic

(Figure S2). In summary, there is substantial heterogeneity in the capability of GluN2A

mutants to facilitate enrichment of NMDA receptors at synapses.

Both severe gain- and loss-of-function mutation types can prolong the decay of
NMDA-EPSCs

The GluN2A mutants examined above were extremely varied in their abilities to contribute

functional receptors at synapses. To understand how the missense mutations are likely to

affect synaptic transmission with native GluN2B expression intact, we next assessed the

effect of GluN2A mutant rescue in *grin2a* knockout neurons Experiments with WT GluN2A

showed that rescue of the peak amplitude and decay time constant of NMDA-EPSCs was

highly sensitive to the level of GluN2A expression (Figure S3). The concentration of WT

GRIN2A plasmid that effectively rescued peak amplitude and decay time of NMDAR EPSCs in

GluN2A KO (3.8 ng/uL, Figure S3) was used to assess rescue of NMDA-EPSCs for GluN2A

mutants (Figure 4). Despite mutant NMDA-EPSCs being consistently smaller than WT, we

1 could not resolve a significant difference for peak amplitude in the contrast between

2 mutants and WT (Figure 4a, Figure S1.4, F(1,119)=2.47, p =.12), and the evidence was

3 overall strongly in favour of there being no effect among the peak amplitudes across the

4 mutants (Figure S1.4, BF_{10} = 0.084). Relative to WT, estimates of the peak NMDA-EPSCs

5 amplitudes appeared slightly smaller for LOF mutants (80% for C436R, 80% for T531M and

6 74% for R518H, 95% CIs [57, 111], [57, 111] and [53, 103] respectively), than for GOF

7 mutants (87% for K669N and 91% for L812M, 95% CIs [65, 116] and [68, 122], respectively).

8 While examining the NMDA-EPSCs, we noticed that when scaling the NMDA-EPSCs to

9 normalize their amplitudes, the decay kinetics appeared to differ between the genotypes

10 (Figure 4bi). In quantifying and comparing the time constant of the NMDA-EPSCs we found

11 strong support for an effect of GluN2A mutation on NMDA-EPSC decay (Figure S1.5,

12 F(5,119)=12.1, p <.001, BF_{10} = 3.43 x 10^8). Overall, the decay time constants of the mutants

13 were significantly different from WT (Figures 4b and S1.5, F(1,119)=37.1, p <.001) as well as

14 between GOF and LOF mutation types(Figures 4b and S1.5, F(1,119)=28.3, p <.001). Relative

15 to WT (100%), the decay time constants were slower for GOF mutants, it being 138% for

16 K669N and 143% for L812M, 95% CIs [111, 171] and [115, 178], respectively. Surprisingly,

17 not only was the direction of effect the same for LOF mutants, but the effect was also larger,

18 with the decay time constants for LOF mutants relative to WT being 201% for C436R, 213%

19 for T531M and 199% for R518H, 95% CIs [158, 256], [167, 272] and [156, 254], respectively.

20 Since we detected changes in the decay of NMDA-EPSCs, we next assessed whether there

21 would be any effect of *GRIN2A* mutations on NMDA-EPSC charge transfer by taking the

22 integral of the current traces (Figure 4ci). Significant differences could be resolved in

23 orthogonal contrasts comparing all mutants with WT (Figure 4c and Figure S1.6,

24 F(5,119)=4.18, p =.04), and between the GOF and LOF mutation types (Figure 4c and Figure

25 S1.6, F(5,119)=4.26, p =.04). However the effects were relatively small, the evidence was

26 more in favour of there being no effect of mutation on NMDA-EPSC charge transfer (Figure

27 S1.6, BF_{10} = 0.632). Compared to WT (100%), NMDA-EPSC charge transfer was slightly larger

28 for LOF mutations C436R (162%), T531M (177%) and R518H (126%), than it was for GOF

29 mutations K669N (125%) and L812M (114%), 95% CIs [106, 247], [116, 270], [83, 193], [86,

30 182] and [78, 167] respectively.

Overall, the effects LOF and GOF GluN2A mutations on NMDA-EPSC decay kinetics were

large and in the same direction, but because of the slightly smaller mutant NMDA-EPSC

amplitudes, the effect that the slowed kinetics had on NMDA-EPSC charge transfer was

much more subtle.

Marginal effects of LOF mutations on the amplitude of AMPA-EPSCs

NMDARs are key regulators of synaptic strength and genetic deletion of NMDAR subunits

has been shown to upregulate AMPAR-mediated synaptic transmission[22,67]. To assess

whether alterations to levels of synaptic AMPAR receptors could occur as a consequence of

less synaptic NMDA receptors for some GluN2A mutants, we also measured evoked AMPAR-

receptor mediated EPSCs (AMPA-EPSCs). While we resolved a significant difference for

AMPA-EPSC amplitude in the contrast between GOF and LOF mutations (Figure S4 and S1.7,

$F(5,119)=7.23$, $p =.008$) the effects were relatively small and overall the evidence was more

in favour there being no effect of mutation on synaptic AMPA receptors (Figure S1.7, $BF_{10} =$

0.276). Peak amplitudes of LOF mutants relative to WT (100%) were 122% for C436R, 117%

for T531M and 112% for R518H, 95% *CI*s [82, 183], [78, 175] and [75, 167], respectively. In

contrast, the amplitudes of AMPA-EPSCs in neurons with GOF mutations were 86% for

K669N and 85% for L812M, 95% *CI*s [60, 123] and [59, 122], respectively. The evidence here

for an effect of NMDA receptor loss-of-function on AMPA-EPSCs is not as convincing as the

results from NMDA receptor subunit deletion experiments reported by others[22,67].

Intermediate effects of a heterozygote grin2a loss-of-function allele on NMDA-EPSCs

The data thus far raise two important questions: 1) Why do LOF mutations have the same

direction of effect on NMDA-EPSC time course as the GOF mutations? and 2) What effect

can we expect in heterozygotes for LOF alleles, especially considering that patients with

GRIN2A mutations have only one mutant allele? We addressed these questions by

examining the effect of homozygous and heterozygous null alleles on evoked NMDA-EPSCs.

We achieved this genetic manipulation by transfecting Cre-recombinase in pyramidal

neurons in CA1 of organotypic slices prepared from *grin2a*$^{+/+}$, *grin2a*$^{fl/+}$ and *grin2a*$^{fl/fl}$ mouse

pups. We found evidence (albeit it weak) against differing effects of Cre across the

genotypes for NMDA-EPSC charge transfer (Figure S1.8, p = .30, BF_{10} = .277), and even the

apparent upward trend was not statistically significant for increasing doses of mutant allele

in a linear polynomial contrast (Figures 4a and S1.8, $F(1,96)$ = 2.39, p = 0.13). However, a

differing effect of Cre across the genotypes was strongly supported for NMDA-EPSC peak

amplitude, decay time constant and rise-time (Figures S1.9-1.11, BF_{10} = 227, 4.67 x 10^{19} and

2.33 x 10^{13} respectively, p <0.001 for each). Significant differences could be resolved for all

pairwise comparisons between *grin2a* $^{+/+}$, *grin2a* $^{-/+}$ and *grin2a* $^{-/-}$, except for the comparison

of peak amplitudes between *grin2a* $^{+/+}$ and *grin2a* $^{-/+}$ (Figures 5c-d, and S1.9-1.11). Consistent

with the NMDA-EPSCs observed in the rescue experiments of the LOF GluN2A mutants, the

decay time constant was 207% slower for *grin2a* $^{-/-}$ (95% *CI* [183, 233]) relative to WT g*rin2a*

$^{+/+}$, 100%). Together with the negligible NMDA-ESPC in the double knockout (*grin2a* $^{-/-}$ *grin2b* $^{-}$

$^{/-}$, Figure 2b), this suggests that the residual current is mediated by GluN2B, which has slower

deactivation kinetics and could explain the more prolonged NMDA-EPSCs observed also for

LOF GluN2B mutants (Figure 4B). The effect observed for NMDA-EPSC decays in *grin2a* $^{-/+}$

neurons (i.e. heterozygotes) was 120% (95% *CI* [106, 137]), it being intermediate between

wildtype and homozygous cases. Overall, these results illustrate that even a single null allele

of *grin2a* is sufficient to cause defects in synaptic transmission.

Mutation-associated changes in NMDA-EPSCs can shape synaptic excitability and

calcium dynamics

The results raise questions about how the changes in NMDA-EPSCs by the mutations will

ultimately affect excitability. In particular, how could smaller and slower NMDA-EPSCs alter

excitability when the amount of charge transfer is effectively unchanged? Using

computational models of a CA1 pyramidal neuron, complete with a wide range of active

conductances[68]and synaptic AMPA and NMDA receptors[69] (Figure 6ai), we simulated the

effects of slowing the kinetics of NMDA-EPSCs to 150% and 200% compared to WT, which

roughly correspond to the effects observed for GOF and LOF mutants respectively. In each

case the synaptic weight (a.k.a peak amplitude) was also scaled to equalize charge transfer

for single EPSCs (Figure 6aii and iii). We then simulated short trains of synchronous synaptic

activation and compared spine head currents and intracellular calcium levels at activated

synapses between the different NMDA-EPSC profiles.

Trains of 40 randomly positioned synapses, simultaneously activated over a range of

frequencies (20, 40, 80 and 200 Hz) were simulated and the membrane potential (in soma

and spine heads) and inward currents, outward currents and calcium concentrations in spine

heads were stored, averaged across spines and summarised as current scapes[70] (Figure 6b).

The simulation was repeated with 10 different random seeds for each NMDA-EPSC profile

and stimulation protocol. While an individual EPSP burst was subthreshold across the three

groups, WT-like (100%) EPSP trains drove neuronal spiking by the 4th or 5th EPSP during short

high frequency (200 Hz) trains, and the action potentials backpropagated to spine heads

(Figures 6b, ci). The voltage changes in spine heads were associated with robust activation of

NMDARs, R-type Ca^{2+} channels (CaR) and Ca^{2+}-activated K^+ channels (KCa) (Figures 6b, cii)

and large increases in Ca^{2+} reaching a peak of >10 μM, and an integral of 1.3 mmol.ms/L

(Figures 6b, ciii). At the same frequency, neuronal spiking was either lower or absent for the

GOF-like (150%) or LOF-like NMDA-EPSP condition respectively (Figures 6b, ci). These

changes in spiking were also associated with less activation of NMDARs, CaR and KCa, and

less accumulation of Ca^{2+}, in spines for mutant-like NMDA-EPSC profiles (Figures 6b, ciii).

Summation of EPSPs at slower frequencies did not result in any spiking and the differences

between NMDA-EPSC profiles for depolarization and intracellular Ca^{2+} became smaller

(Figures 6ciii).

The full set of simulations above were repeated with 200 synchronous synaptic inputs to

evaluate the effect of NMDA-EPSC profiles on the summation of suprathreshold EPSPs. WT-

like (100%) (but not mutant-like) EPSP trains at 80 Hz and above triggered plateau potentials

that were associated with robust activation of spine NMDARs, CaR and KCa (Figure 6d).

Interestingly, the spiking output was very frequency-dependent, with there being greater

spiking in mutant-like NMDA-EPSCs at 80 Hz, but less spiking at 40 Hz and similar spiking at

the more extreme train frequencies (Figure 6d, ei). Examining the voltage traces (and axonal

Na currents, data not shown) at 80 Hz revealed that spiking was strongly attenuated during

the plateau potential likely due to Na-channel inactivation. Overall spine Ca^{2+} accumulation

reached up to 2-3 times the levels compared to 40 active synaptic inputs, and the mutant-

like NMDA-EPSC profiles still lead to less accumulation of spine Ca^{2+} (Figure 6eiii).

Overall, these results indicate that even in the absence of changes in charge transferred

during synaptic activation, mutant NMDAR kinetics could still effectively modulate synaptic

excitability and spine calcium dynamics, with the effects being very frequency-dependent.

Discussion

Our investigation revealed how mutations from patients with *GRIN2A*-related epilepsy

disorders result in defective synaptic transmission. The robust differences in the ability of

mutant GluN2 subunits to rescue NMDA-EPSCs in GluN2-null ($grin2a^{-/-}\ b^{-/-}$) neurons (Figure

2) was largely consistent with the classification of mutations, as severe GOF and LOF, based

on analysis of data obtained in heterologous expression systems by others (Figure 1).

Remarkably though, replacing only the native GluN2A with the mutant receptor subunit gave

rise to NMDA-EPSCs with very similar defects for LOF and GOF mutations: NMDA-EPSCs that

were slower in time course but similar in charge transfer (Figure 4). In general, mutations

phenocopied the GluN2A knockout (Figure 5), but the mechanism underlying the defect

varied, with some mutants being associated with poor trafficking to synapses (Figures 3 and

7b) while others reached the synapse but were either functionally silent (Figures 2, 3 and 7c)

or contributed directly to the NMDA-EPSCs through altered functional properties of the

mutant receptor (Figure 2 and 7d). In the case of the LOF mutations, the changes in the

NMDA-EPSC time course likely reflect the contribution of natively expressed GluN2B (Figures

5 and 7b-c).

The ability of GluN2A mutants to incorporate into synaptic NMDARs varied considerably

(Figures 2 and 3). The LOF mutation C436R gave similarly low levels of synaptic enrichment

of NMDARs as the control without GluN2A (Figure 3). The C436R mutation disrupts the

formation of a disulfide bridge, which is likely key to the folding and stability of the GluN2A

protein structure (Figure S5). In contrast, the LOF R518H gave similar levels of NMDARs at

synapses as wildtype GluN2A (Figure 3), and the lack of functional incorporation of R518H

(Figure 2c) may be explained by the effect of the mutation on lowering glutamate potency

(Figure 1c-e) by breaking coulombic interactions between GluN2A and the glutamate agonist

(Figure S5). Some mutations, like the T531M mutation, may even show a combination of

these mechanisms (Figures 2-3 and S5). It is important to point out that some mutations may

have effects not represented by this sample, the LOF mutants of which were all from the

LBD. It is conceivable that mutations in other parts of the receptor could give receptors that

traffic to the synapse but whose functional defects lead to distinct synaptic phenotypes.

Nonetheless, our results show that mutations with severe and contrasting LOF and GOF

effects on GluN2A-containing receptors can lead to effects on synaptic function that while

abnormal, are in the same direction.

A current question in the field of *GRIN2A*-related disorders is whether expression or synaptic

delivery of the native GluN2B subunit is specifically upregulated and compensates for

GluN2A loss-of-function[42]. The smaller NMDA-EPSCs in neurons that are homozygous for the

null GluN2A allele indicates that compensation by GluN2B could be incomplete (Figure 5a,

2D). However, in neurons heterozygous for a null allele, the slower, GluN2B-like time course

of NMDA-EPSCs in the absence of changes in amplitude suggest that there could be some,

albeit small, compensation. The mechanism of this compensation may simply reflect a

greater propensity for existing GluN2B-containing NMDARs to accumulate at synapses when

synaptic content of GluN2A is reduced. This means of compensation would still be consistent

with GluN2A deletion not having any effect on either GluN2B total protein or the amplitude

of ifenprodil-sensitive whole-cell currents[42]. It should be mentioned though that since

GluN2A and GluN2B subunits couple to distinct downstream signalling pathways through

their distinct intracellular C-tails[11,71], changes in the GluN2A/B subunit composition at

synapses could be expected to have an impact on long term regulation of synaptic function

(e.g. synaptic plasticity). As such, more complex aspects of synapse phenotype between GOF

and LOF mutants may indeed differ in more than ways we can determine from measuring

basal synaptic transmission. While unlikely in our single-cell genetic manipulations, we also

cannot rule out the possibility that disturbances in NMDA receptor function lead to

downstream changes in the relative expression of GluN1 splice variants, which could also

explain changes in NMDA-EPSC time course[72,73]. Although it is also possible that

manipulations of post-synaptic GluN2 NMDA receptor subunits could change presynaptic

release probability and interfere with interpreting effects of GluN2A mutations on NMDA-

EPSC amplitude, the lack of effect of single-cell KO of NMDA receptor subunits on paired-

pulse facilitation at Schaffer collateral synapses makes this seem unlikely[22].

In general, we were surprised by the relatively modest effects of GluN2A mutations on

synaptic NMDA-EPSCs in our *grin2a* [-/-] rescue experiments (Figure 4) compared to our results

in the GluN2A/B double knockout (Figure 2). Together with the association of these

mutations with seizure disorders (Table S2), this highlights just how sensitive the excitability

of neurons is to the intrinsic properties and abundance of synaptic NMDARs. The actual

impact of *GRIN2A* mutations on NMDA-EPSCs may even be overestimated here since we

performed our rescue experiments in *grin2a*[-/-] (homozygote) neurons, with the risk being

that the effects we observed may not be truly representative of what occurs in the neurons

of patients that are heterozygous for the mutations. This could be of particular concern for

LOF mutations, in which the remaining wildtype GluN2A allele could compensate for the

mutant allele. Our experiment measuring NMDA-EPSCs in neurons that were heterozygous

for a *grin2a* null allele addressed this concern (Figure 5) and suggests that qualitatively, our

inferences throughout could hold in the heterozygote case, although we cannot rule out that

heterozygosity for GOF mutations could lead to unpredictable effects on NMDA-EPSCs.

The overall effects of GluN2A mutations on the properties of the NMDA-EPSCs raised some

interesting questions about how they may modify synaptic excitability. In particular, the

slower time course and smaller amplitudes of the NMDA-EPSCs in *grin2a*[-/-] neurons rescued

with GluN2A mutants resulted in charge transfer through the NMDAR channels being similar

to wildtype. This prompted us to model the effect of the mutations on NMDA-EPSCs and

simulate the depolarizations and spince calcium dynamics during trains of concerted

synaptic activity onto CA1 neurons (Figure 6). These simulations serve to illustrate that the

slower time course of NMDA-EPSCs mediated by GluN2A mutants can affect EPSP

summation even when charge transfer is the same. However, there may be other effects of

the GluN2A that we have not measured or modelled (e.g. presynaptic or extrasynaptic

receptors, synaptic plasticity) that contribute to synaptic excitability and neuronal signalling.

In addition, GluN2A is expressed in other neuronal cell types (including inhibitory neurons[74])

and how the mutation effects in all cells sum together to affect excitatory-inhibitory balance

and network activity is unknown. Together, it is clear that understanding how NMDAR

mutations affect excitability and network stability warrants further investigation and may

ultimately help in designing therapeutic strategies to treating *GRIN2A*-related disorders.

Methods

Mutant classification

The functional consequences of 20 *GRIN2A* mutations were collected from publications and

databases[44,46,50–57]. The functional properties were Glutamate EC_{50} and Glycine EC_{50} (in

oocytes), and deactivation time constant, current density, open probability and relative cell surface levels (in HEK-293T cells). Potency for R518H was approximated from the fold change in potency for R518C documented in the CFERV database[51]. Potency for T531M was approximated by scaling the R518H potency according to the R518H/T531M ratio for single-channel opening frequency corrected for the differences in surface expression[56]. To handle other remaining instances of missing data, single imputation was achieved by linear interpolation following orthogonal regression on the rank-transformed data using principal component analysis (PCA) by the alternating least-squares algorithm in Matlab (Mathworks). The complete data set was then normalised to the wildtype (WT) GluN2A measurement for each property and transformed to the log scale. The data set had a Kaiser-Meyer-Olkin test statistic of >0.5 and was considered suitable for PCA. More than 85% of the variance was explained by the first two principal components, which were then extracted from the loadings matrix and subjected to oblique (Varimax) rotation. The resulting pattern matrix was used to calculate rotated component scores. The number of clusters chosen for K-means clustering of the scores was determined from the elbow in a plot of the sum-of-squares error against the cluster number.

Molecular Biology

Mutagenesis was performed on the cDNA of human *GRIN2A* in the pCI-Neo vector (pCMV GluN2A). GluN2A mutations with respect to the amino acid numbering in NP_000824.1 were: C436R, T531M, R518H, K669N, L812M. Oligonucleotide primers (Integrated DNA Technologies) were designed to incorporate single-base substitutions using the GeneArt® Site-Directed Mutagenesis Kit (Thermofisher Scientific) (Table S3). The presence of each mutation was confirmed by DNA sequencing (Eurofins Genomics) and the whole coding sequence was also examined to ensure that no other mutations were introduced by polymerase chain reaction (PCR) errors. Plasmid DNA was amplified and purified using EZNA endo-free plasmid maxi kit (VWR International).

Animals

C57Bl/6J mice and Sprague Dawley rats were purchased from Charles River Laboratories. Rodents were maintained under conventional housing conditions and a 12-12 light-dark cycle. Heterozygous *grin2a*-flox (*grin2a+/fl) and grin2b*-flox (*grin2b+/fl)* mice were a gift from

Kenji Sakimura (Niigata University, Japan). Generation of these mice has been described previously[22,63]. Homozygous *grin2a*-flox (*grin2a$^{fl/fl}$*) and *grin2b*-flox (*grin2b$^{fl/fl}$*) mice were generated by mating the respective heterozygous mice and the colonies were subsequently maintained as homozygous lines taking care to avoid brother-sister matings. Double-flox (*grin2a$^{fl/fl}$ 2b$^{fl/fl}$*) mice were generated by crossing the *grin2a$^{fl/fl}$* and *grin2b$^{fl/fl}$* mouse lines. Ear biopsies taken for identification were used for genotyping by PCR. Primers used for *grin2a*-flox genotyping were: 5'-GAAATGTGGTAAAATCCAGTTAG-3' (forward), and 5'-TAGGCAGTAAACTTTCCTCATC-3' (reverse). Product sizes were 800 base pairs (bp) for the WT allele and 1100 bp for the *grin2a*-flox allele. Primers used for *grin2b*-flox genotyping were: 5'-GTCAGAGAGTTTAACCTCAG-3' (forward), 5'-ACAAGCACCTTCTTGGTCTC-3' (reverse) and 5'-GCTGCTAAAGCGCATGCTCC-3' (Neo) Product sizes were 940 bp for the WT allele and 200 bp for the *grin2b*-flox allele. Neonatal animals used for experiments were sacrificed by an appropriate Schedule 1 procedure (cervical dislocation) in accordance with the Animals Scientific Procedures Act 1986 amendment regulations 2012. Schedule 1 training of the experimenters was overseen and approved by the local Named Training and Competency Officer (NTCO). Experiments were reviewed *a priori* by the Animal Welfare Ethical Review Body at the University of Sussex and the project was licenced by the Home Office (UK, PPL# P6CF775B9).

Cultures

Organotypic hippocampal slice cultures were prepared from neonatal mice. The hippocampi were dissected from mice at postnatal age 6-8 days in cold sterile filtered dissection medium composed of (in mM): $CaCl_2$ (0.5), KCl (2.5), KH_2PO_4 (0.66), $MgCl_2$ (2), $MgSO_4$ (0.28), NaCl (50), Na_2HPO_4 (0.85), glucose (25), $NaHCO_3$ (2.7), sucrose (175) and HEPES (2) (pH 7.3, 330 mOsm). Transverse hippocampal slices (350 µm) were cut using a McIlwain tissue chopper (Abbotsbury Engineering Ltd., UK), placed on pieces of membrane (FHLC01300, Millipore, UK) placed in Millicell cell culture inserts (PICM03050, Millipore, UK) and cultured in 6-well plates. The culture medium (1 ml per well) was composed of Minimum Essential Medium (MEM) supplemented with 15% heat-inactivated horse serum (26050-088, Gibco), B27 supplement (17504044, Gibco), 25 mM HEPES, 3 mM L-glutamine, 1 mM $CaCl_2$, 1 mM $MgSO_4$, 0.25 mM ascorbic acid and 5 g/L glucose. Slice culture plates were kept in an

incubator at 34°C with 5% CO_2 and 95% humidity. The culture medium was changed twice a

week, with prewarmed slice culture medium supplemented with 10 µg/ml gentamicin.

Dissociated cultures of hippocampal neurons were prepared from neonatal Sprague Dawley

rats. The hippocampus was dissected on ice and the meninges were removed in dissection

medium composed of HBSS (14155048, ThermoFisher Scientific) supplemented with 10 mM

HEPES (H0887, Sigma) and 100 U/ml Penicillin and Streptomycin (Pen-Strep, 15140122,

ThermoFisher Scientific). The hippocampi were then triturated in 2 ml of pre-warmed plating

medium, which was composed of MEM (51200087, ThermoFisher Scientific) supplemented

with 20 mM glucose, 100 U/ml Pen-Strep, 1 mM sodium pyruvate (Sigma S8636), 25 mM

HEPES, N2 supplement (17502048, ThermoFisher Scientific) and 10% heat-inactivated horse

serum (26050088, ThermoFisher Scientific). The homogenised tissue was then diluted with

plating medium to 3.25 ml per hippocampus. 500 µl of this was plated per well into 12-well

plates that contained 16 mm diameter coverslips (10755354, ThermoFisher Scientific) that

had been previously coated with 50 µg/ml poly-D-lysine (P0899, Sigma) and 2 µg/ml laminin

(L2020, Sigma). Neurons were allowed to settle and adhere for 3 hours at 37°C, 5% CO_2,

after which, they were supplemented with 1.5 ml culture medium containing Neurobasal-A

(12349015, ThermoFisher Scientific), B-27 supplement (17504044, ThermoFisher Scientific),

100 U/ml Pen-Strep and GlutaMAX (35050038, ThermoFisher Scientific). Cultures were

returned to 37°C, 5% CO_2 and fed twice a week with culture medium supplemented with

antimitotic agents: 0.1 µM uridine, 0.1 µM fluorodeoxyuridine and 0.1 µM cytosine

arabinoside (all from Sigma).

Transfections

Neurons in organotypic slice cultures were transfected by single-cell electroporation[75]. At 6-

8 days *in vitro* (DIV) CA1 pyramidal neurons in organotypic slices were electroporated with

plasmids expressing pCMV Cre-GFP with or without WT or mutant pCMV *GRIN2A* cDNA

(C436R, T531M, R518H, K669N, or L812M). Plasmid DNA was precipitated upon addition of

0.5 volume of PEG solution (30% PEG 8000 (w/v) in 30 mM $MgCl_2$ to isolate it from residual

lipopolysaccharide endotoxin. The DNA precipitate was pelleted by centrifugation at 10,000

relative centrifugal force (rcf) for 15 minutes, washed with 70% ethanol and the air-dried

pellet was re-dissolved in TE buffer (10 mM Tris-HCl, 1 mM EDTA). Within a week of

transfection, pCMV Cre-GFP was diluted to 0.53 nM (3 ng/μL) in intracellular solution

containing (in mM): CH_3SO_3H (135), KOH (135), NaCl (4), $MgCl_2$ (2), HEPES (10), Na-ATP (2),

Na-GTP (0.3), spermine dihydrate (0.15), EGTA (0.06) and $CaCl_2$ (0.01) (pH 7.25, 285 mOsm).

In molecular replacement experiments, the solution also included 0.62 nM (3.8 ng/μL) of WT

or mutant pCMV *GRIN2A*. DNA-containing intracellular solution was centrifuged at > 10,000

rcf for 15 minutes at 4°C to remove debris before using it to fill patch pipettes (8-10 MΩ),

which were pulled with a Flaming/Brown Micropipette puller from thick-walled borosilicate

glass capillaries (GB150F-8P, Science Products). Slices were transferred to the recording

chamber on an upright microscope (BX51, Olympus) containing room temperature

extracellular solution composed of (in mM): NaCl (140), KCl (3), $MgCl_2$ (1), $CaCl_2$ (2), glucose

(10), Na-pyruvate (1), $NaHCO_3$ (2), HEPES (6), Na-HEPES (4) (pH 7.35, 300 mOsm). The patch

pipette was positioned in the slice under visual guidance using a mechanical manipulator

(PatchStar, Scientifica). CA1 pyramidal cells were approached with positive pressure (20

mbar). Upon dimple formation, the pressure was released to form a putative loose-patch

seal. Immediately, a 12 V stimulus train was applied (100–200 Hz for 0.25–0.5 s; pulse-width

0.25–0.5 ms) from a stimulus isolator (IsoFlex, A.M.P.I.) triggered from ACQ4 software

(v0.9.3)[76] through a USB-X Series Multifunctional DAQ interface (NI USB-6341, National

Instruments). Using the same patch-pipette, ten CA1 pyramidal neuron transfections were

attempted in each slice. Typically, expression (judged from the fluorescent protein marker)

was observed in half the neurons in which transfection was attempted.

Cultures of dissociated hippocampal neurons were transfected at 8 DIV using the calcium

phosphate method[77]. DNA constructs cotransfected (μg/coverslip) were pCaMKII-α

Homer1c-tdTomato (1), empty pcDNA3.1+ (1.5), and pCMV constructs driving expression of

SEP-GluN1 (1.5) (Addgene plasmid #23999) and GluN2A (1.5). Per well of a 12-well plate, a

mixture of plasmid DNA and calcium chloride (50 μl, 2.5 M) was added dropwise to 50 μl 2x

HEPES buffered saline (E1200, Promega) and incubated for 20 minutes at room temperature

in the dark. Coverslips containing neurons were transferred to a fresh 12-well plate

containing 500 μl pre-warmed culture medium supplemented with 1 mM kynurenic and

followed by the addition of DNA/$CaCl_2$-HEPES solution. After a 90-minute incubation at 37°C

with 5% CO_2, the coverslips were transferred to a new 12-well plate containing culture

medium, which had been pre-incubated at 37°C and 10% CO_2. The 12-well plate was then

incubated at 37°C with 5% CO_2 for 20 minutes before being transferred back to the original

12-well plate (containing conditioned medium) and incubated at 37°C with 5% CO_2 until the

day of imaging.

Electrophysiology

The electrophysiology method utilized throughout was current measurements in whole-cell

patch-clamp configuration. Hippocampal slices had a cut made between the CA3 and CA1

regions to reduce recurrent excitation during subsequent recordings of evoked synaptic

transmission. Slices on their membrane were transferred to the chamber of an upright

microscope (SliceScope Pro 2000, Scientifica), visualised using infrared light (780 nm)

through an oblique condenser and 40x 0.8 NA objective (LUMPLFLN40XW, Olympus) and

captured using a mono charge-coupled device (CCD) monochrome camera (SciCam,

Scientifica) controlled using Scight software (Scientifica). The tissue was held down with a

slice anchor and perfused at 2 ml/min with artificial cerebrospinal fluid (aCSF) solution

containing (in mM): NaCl (125), KCl (2.5), $NaHCO_3$ (25), glucose (10), Na-pyruvate (1), $MgCl_2$

(1) and $CaCl_2$ (2). 5 µM 2-chloroadenosine (ab120037, Abcam) was added to prevent

epileptic activity. Addition of other drugs depended on the experiment (see below). The

aCSF was bubbled with carbogen (95% O_2, 5% CO_2) and maintained at 30°C with an in-line

heater (SM-4600, Scientifica) and heated chamber (HCS, ALA Scientific; TC-10, NPI

Electronic). A bipolar tungsten stimulation electrode was positioned (LBM-7, Scientifica) in

Stratum radiatum to stimulate the Schaffer collateral pathway. 2-4 MΩ patch pipettes for

whole-cell patch clamp recordings were filled with intracellular solution composed of (in

mM): CH_3SO_3H (120), CsOH (120), CsCl (20), $MgCl_2$ (2.5), HEPES (10), Na-ATP (4), Na-GTP

(0.4), phosphocreatine disodium salt (5) and EGTA (0.2) (pH 7.25). Osmolarity of intracellular

and extracellular solutions was checked and adjusted on the day of each experiment and

adjusted by addition of water or the major salt so that the aCSF osmolarity was 10-15

mOsmol/kg greater than the intracellular solution. To account for inter-slice variability in our

recordings, two CA1 neurons were patched simultaneously – one transfected neuron

identified by the Cre-GFP fluorescence (pE-300 white, CoolLED; 49018 EGFP long-pass,

Chroma) and one neighbouring untransfected neuron. The intensity and polarity of the

constant voltage stimulus (50 µs duration) was adjusted (IsoFlex, A.M.P.I.) to obtain reliable

NMDAR responses that were < 250 pA, typically 10 V. The configuration of constant voltage

stimulus enabled us to induce rapid changes in the potential difference across the electrodes

with minimal influence of stray capacitance. Stimuli were delivered at an inter-sweep

interval of 10 s. Signals were acquired with a MultiClamp 700B (Molecular Devices), low-pass

filtered (4 kHz, 4-pole Bessel) and digitized (25 kHz) directly to hard disk using a USB-X Series

Multifunctional DAQ interface (NI USB-6363, National Instruments). Hardware was

controlled using Python-based ACQ4 software (v0.9.3)[76]. The series resistance was left

uncompensated during data acquisition (but see later). Pairs of neurons were discarded

during data collection if the difference in series resistance varied by more than ~8 MΩ.

For molecular replacement experiments (Figures 2, 4 and S4), 10 evoked mixed AMPAR- and

NMDA-EPSCs were recorded at a range of voltages (-100 to +20 mV in 20 mV increments).

These holding potentials account for the liquid-liquid junction potential, which we calculated

to be +10 mV for our solutions (PowerPatch Tools Igor Pro Tool collection). The aCSF

included 50 µM picrotoxin, 10 µM gabazine as above, and 2 µM CGP52432 to inhibit $GABA_B$

receptors. The intracellular solution was supplemented with 3 mM QX-314-Cl to attenuate

currents through voltage-activated sodium and calcium channels. For each voltage, 10

evoked NMDA-EPSCs were recorded per neuron at an inter-sweep interval of 10 s.

Transfections and recordings for GluN2A mutant molecular replacements were randomized

and performed double blind. All molecular replacement experiments were carried out in

slice cultures prepared from the left hemisphere since the measurements of NMDA-EPSC

properties tended to show a lower variance compared to those in neurons of slices from the

right hemisphere.

The mutants selected for recordings in this study are known to have no more than 2.5-fold

changes in magnesium potency[51,57]. We thus separated AMPAR- and NMDAR-components of

evoked mixed EPSCs during the analysis using the following protocol: 1) The current traces at

each holding potential were converted to conductance using Ohm's law; 2) The AMPAR-

mediated conductance at -100 mV was subtracted from the conductance trace at +20 mV.

The resulting conductance traces thus reflect AMPA-EPSCs (-100 mV) and NMDA-EPSCs (-40

mV and +20 mV).

For experiments where the NMDAR current was isolated pharmacologically (Figure 5), the

aCSF included 50 µM picrotoxin (ab120315, Abcam) and 10 µM gabazine (SR95531, Abcam)

to block ionotropic gamma-amino-butyric acid receptors (GABA$_A$Rs) and 10 µM NBQX (1044,

Tocris Biosciences) to inhibit α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid

receptors (AMPARs). The membrane potential was raised from -75 mV to +20 mV at a rate

of 0.5 mV per second to provide relief from the magnesium block on NMDARs. For each

neuron, 30 evoked NMDA-EPSCs were recorded.

Since the current-voltage relationship for AMPA-EPSCs$_{-100 mV}$ and NMDA-EPSCs$_{+20 mV}$ relative

to the reversal potential (approximately 0) can be assumed linear, we performed offline

corrections to compensate for series resistance[78]. The stimulus artifact was interpolated and

the series resistance (Rs) and cell membrane capacitance (that were estimated from the test

pulse in each recording) were used to perform offline series resistance compensation. For

consistency, final, uncompensated current in each recording (Rs_{final}) was set to a constant 3

Mohm, where the % of compensation applied to each recording was calculated using the

following equation.

14
$$Rs_{comp}(\%) = 100 \times \max\left(1 - \frac{Rs_{final}}{Rs}, 0\right)$$

Series resistance compensated recording traces were analysed using a custom python

module (https://github.com/acp29/penn) within Stimfit software (v0.13 or 0.15.8)[79] or

custom scripts in Matlab (available at https://github.com/acp29/Elmasri_GRIN2A). Decay of

the EPSC currents was fit by a two-component exponential decay with offset initially using

the Chebyshev algorithm. The weighted exponential time constant was calculated as the

sum of component time constants multiplied by their fractional amplitude. Where the two-

component fits failed, a single exponential was found to fit well.

Imaging

Transfected cultures of dissociated hippocampal neurons were imaged at DIV11. Neurons

were imaged using an inverted microscope (AxioVert.A1, Zeiss), a 40x 0.55 NA objective and

a complementary metal-oxide-semiconductor (CMOS) camera (ProgRes GRYPHAX Arktur 8

MPix, Jenoptik). Images were acquired using GRYPHAX software (v2.2.0, Jenoptik). Coverslips

were placed in ASCF composed of (in mM): NaCl (140), KCl (3), MgCl$_2$ (4), CaCl$_2$ (4), glucose

(10), Na-pyruvate (1), HEPES (6), Na-HEPES (4) (pH 7.35, 300 mOsm). Neurons were imaged

live for the GluN1-SEP signal at (LED λ_{ex} 470 nm) and the Homer1c-tdTomato signal at (LED

λ_{ex} 530 nm). Five images (8-bit) were taken for each coverslip and converted to grayscale in

FIJI software (v1.53)[80,81]. Red and green images were aligned if necessary, with custom

macros using a modified version of the Align_RGB_planes plugin available at

https://github.com/acp29/pimage/tree/master/Penn/align_rgb). The background

fluorescence was subtracted from both images, and puncta in the Homer1C-tdTomato signal

were selected semi-automatically as regions of interest (ROIs). The ROIs were overlaid onto

the GluN1-SEP signal and the mean pixel intensity was measured at these ROIs. Custom

macros used for the measurement of spine fluorescence analysis are available at

https://github.com/acp29/pimage/blob/master/Penn/pimage.ijm. The experimenter was

blind to the experimental groups during image analysis, and the results were reproduced by

multiple lab members.

NEURON simulations

Simulations were conducted in NEURON 7.5 software[82]. Simulations were carried out using a

multicompartmental model of a CA1 pyramidal neuron developed by Graham and

colleagues[68]. The model includes spatially distributed ion channels responsible for fast

sodium (Na) currents, delayed rectifier potassium (K) currents, A-type potassium (K)

currents, h-current, HVA R-type calcium (Ca^{2+}) current, and Ca^{2+}-activated mAHP potassium

(K) currents. In all cases, simulations were conducted with a time step of 25 μs and at a

temperature of 35°C. Simulated synaptic inputs consisted of both AMPAR and NMDAR

conductances described by the sum of two exponentials. AMPAR receptor rise and decay

time constants were set to 0.3 ms and 3 ms respectively and maximal conductance (g_{max})

was set to 500 pS. WT NMDAR rise and decay time constants were set to 3 ms and 80 ms

respectively. The rise and decay time constants of NMDAR conductances were scaled by a

factor of 1.5 or 2.0 to approximate the kinetics of GOF and LOF mutations on NMDA-EPSCs.

The NMDAR gmax was set to 1 nS, 666 pS and 500 pS to emulate WT, GOF and LOF NMDARs

respectively, with constant charge transfer. Activation of 40 or 200 excitatory synapses (as

two-compartment spines) located randomly along the apical dendrites was used to simulate

a barrage of synchronous synaptic events. For each experimental run, barrages of synaptic

events were generated in trains of 5 stimuli at frequencies of 20, 40, 80 or 200Hz. With the

exception of the randomized distribution of activated synapses controlled by setting the

random seed, the models and experiments that were conducted were all deterministic in

nature.

Statistics and reproducibility statement

Data was collected until the number of cell pairs reached between a total of about 15-30 per

mutation (or genotype). The experiment was repeated in 3-6 animals. The choice of which

slices/animals to allocate to which mutation or genotype was random. The experimenter

was blind to the mutation or genotype of the sample. Since it is possible that the occasional

neuron we patched was not representative of the population (e.g. an interneuron, or had

poor co-transfection with *GRIN2A* cDNA), we performed outlier detection prior to data

analysis. Whole-cell properties and measurements of NMDA-EPSCs in matched transfected

and untransfected where expressed as $\log_e$ response ratios and included in multivariate

outlier detection by robust principal component analysis (ROBPCA) using the LIBRA Matlab

library[83,84] in Matlab 9.2 (R2017a, Mathworks). ROBPCA produces principal components

(PCs) that are representative of the trend of the majority of the data with lower influence

from extreme values. The number of PCs used was the number that explained 90% of the

variance. Samples that were extreme orthogonal outliers, bad leverage points and were

beyond the cut-off value of $\sqrt{X^2_{q,0.975}}$ were excluded from all subsequent analysis. Typically,

about 2 recording pairs (range from 0 - 4) were excluded per group of data (e.g. per

mutation). A summary of the sample sizes after multivariate outlier detection is presented in

Table S4.

Statistical analysis was carried out in R (v 4.1.0) and RStudio (v 1.4). Each response datum

from the patch-clamp recordings (in Figures 2, 4, 5 and S4) was collected along with its

assignment to (levels of) the following factors: mutation (WT, C436R, T531M, R518H, K669N,

L812M) or genotype (wildtype, heterozygote, homozygote), transfection (unt. or tr., for

untransfected or transfected respectively), and cell pair, slice and animal (with unique

identifiers for each to make the nested data hierarchy explicit). For the imaging experiments

(in Figure 3), each fluorescence measurement was collected along with its assignment to

(levels of) the following factors: mutation (WT, C436R, T531M, R518H, K669N, L812M),

protein (GFP-GluN1 or Homer1c-tdTomato), region of interest (ROI) and experimental repeat

(with unique identifiers for each to make the nested and crossed data hierarchy explicit).

Factors were appropriately assigned as fixed effects, or (nuisance) random effects. Response

data from each experimental data set was $\log_e$-transformed and then fit by linear mixed

effects regression with univariate linear mixed models (LMMs) using the `lmer` function from

the R package `lme4`. Model fitting was set to optimize the restricted maximum likelihood

(REML) criterion. The `lmer` formula for each random intercept model is reported in Figures

S1.1-S1.11 and follows described convention for mixed models[85]. Where hypothesis testing

was based on *a priori* clustering of the data (as in Figure 1e), orthogonal contrasts (Table S1)

were setup and applied before model fitting. Where hypothesis testing was based on testing

the significance of trends across ordered (categorical) factors, polynomial contrasts were

setup and applied before model fitting.

Model assumptions were evaluated by graphing plots of the residuals. Homoscedasticity was

assessed by examining the scatter of the standardized model residuals against the fitted

values. Normality of the model residuals was assessed by: 1) superimposing a histogram and

kernel density estimate of the standardized model residuals with a standard normal

distribution; and by 2) creating a quantile-quantile (Q-Q) plot of sample quantiles against

normal quantiles together with 95% tail-sensitive confidence bands. Model outliers were

assessed from the above plots of the standardized model residuals and the degree to which

they were influential was assessed using a stem-and-leaf plot of the Cook's distances

calculated using the `hlm_influence` function from the `HLMdiag` package. Cook's distances

were all < 0.15, far below the threshold for being considered influential (1.0). Graphs were

created using functions from the `ggplot2` package.

The LMM results for the fixed effects were summarised as ANOVA tables using Type III Wald

F-tests and with degrees of freedom calculated by the Kenward-Roger method using the

`Anova` function from the `car` package. To summarise the random effects structure, the

variance of each group level was used to calculate its respective intraclass correlation

coefficient (ICC) using the `icc` function from the `performance` package. When *posthoc*

tests were requested, they were pairwise comparisons with type I error rate controlled by

Westfall stepwise procedure and were calculated using the `multcomp` R package. Parameter

estimates and effect sizes are reported in the main text and figures as response ratios

(expressed as %) with 95% confidence intervals calculated using functions from the

`emmeans` package. Effect sizes were standardized by conversion to correlation coefficients

using the `t_to_r` function from the `effectsize` package and reported in the figure

legends. Details of LMM results and diagnostics are provided in the supplement (Figures S1).

All summaries of statistics annotated in graphs have used the following convention: *ns* = not

significant (at α = 0.05), * = $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$. All reported *p*-values are

from two-tailed tests. Code and data relating to the analysis are provided at

https://github.com/acp29/Elmasri_GRIN2A

To evaluate the strength of evidence for the null hypothesis for omnibus ANOVA tests of the

fixed effects, Bayes Factors (BF_{10}) were calculated using the `anovaBF` and

`bayesfactor_inclusion` functions from the `BayesFactor` (v0.9.12-4.3) and

`bayestestR` packages. BF_{10} indicates the likelihood ratio of evidence for ($BF_{10} < 1$), or

against ($BF_{10} > 1$), matched (null) models. Default (Cauchy) priors for fixed and random

effects had the usual scales of 0.5 ('medium') and 1 ('nuisance') respectively.

Confidence intervals and *p*-values for Kendall's tau-b (τ_b) (non-parametric) correlation

coefficients were calculated in Matlab using *iboot*[86], now part of the Octave statistics-

bootstrap package (https://gnu-octave.github.io/packages/statistics-bootstrap). Central

coverage of bias-corrected and accelerated (BCa) intervals was calibrated to reduce small

sample bias using a double bootstrap procedure, with 20,000 and 200 resamples for the first

and second bootstrap respectively.

Acknowledgements

We gratefully acknowledge the following kind gifts: pCMV Cre-GFP from Christophe Mulle

(IINs, Bordeaux, France), pCMV GluN2A from Kasper Hansen, Hongjie Yuan and Stephen

Traynelis (Emory, Atlanta, Georgia, USA) and pCaMKII-αHomer1c-tdTomato from Daniel

Choquet (IINs, Bordeaux, France). The PhD studentship of M.E. was funded on a University of

Sussex Life Sciences School studentship. A.C.P. was funded on MRC Career Development

Award (MR/M020746/1), in which W.A. and M.E. were funded as post-docs.

Figure Legends

Figure 1: Missense GRIN2A mutations can impart severe overall gain or loss of function effects on NMDARs.

a) A pie chart illustrating the proportion of different types of mutations in human *GRIN2A*. Missense mutations make over 40% of all *GRIN2A* mutations but their impact on NMDAR function can be hard to predict. **b)** Surface representation of an NMDAR heterotetramer of GluN1 and GluN2 (pink and yellow respectively). The ligand-binding domain (LBD) of GluN2A is boxed out and magnified to illustrate the glutamate (Glu) binding site, the adjacent gating ring and the location of some functionally characterized disease-associated missense mutations (blue spheres)[44,46,50–57]. NTD = Amino terminal domain. **c)** Raincloud plot showing quantification of the gain (> 0) and loss (< 0) of function of the mutations shown in **b** with respect to different properties measured for mutant GluN1/2A receptors expressed in HEK cells or Oocytes[44,46,50–57]. The black dashed line at 0 corresponds to WT GluN2A. The tick dashed maroon line corresponds to the K669N mutation, which serves to illustrate that some mutations can be simultaneously gain-of-function, loss-of-function, or have no effect across different properties of the receptor. **d)** Stem and leaf plot of the loadings (pattern matrix) of the first two components of a principal component analysis after oblique rotation. The loading for first factor is highest for glutamate potency ($EC_{50,Glu}^{-1}$), while the loading for the second factor is highest for peak current density (I_{peak}). **e)** Scatter plot of the scores of the 20 *GRIN2A* mutations (shown 2B-C) for each of the factors 1 and 2 from D. The graph is annotated with the results of K-means clustering: mutants that are GOF for factor 1 (red, Type 1 GOF); mutants that are mostly LOF for factor 1 (blue, Type 1 LOF); mutants that are LOF mostly for factor 2 (purple, Type 2 LOF); and WT-like mutants (black);. Filled symbols represent mutations used for experiments in this study. The graph is annotated with Kendall's tau-b correlation coefficient accompanied by bootstrap confidence intervals and *p*-values (see methods).

Figure 2: Gain-of-function GluN2A mutants are more effective than loss-of-function GluN2A mutants at rescuing NMDAR-mediated EPSCs in GluN2A/B double knockout neurons

a) Experimental protocol used to test functional incorporation of GluN2A mutants. Plasmids expressing Cre-GFP and human *GRIN2A* cDNA were co-transfected into CA1 pyramidal neurons of organotypic hippocampal slices from $grin2a^{fl/fl} b^{fl/fl}$ mice. With transfection, the action of Cre-GFP at floxed *grin2a* and *grin2b* alleles knocks-out native mouse GluN2A and 2B protein expression concurrently with expressing human GluN2A variants (wildtype or mutant). Untransfected neurons continue to express native GluN2A and B. **b)** NMDA-EPSC$_{+20}$ $_{mV}$ peak amplitudes from $grin2a^{fl/fl} b^{fl/fl}$ (untransfected, -Cre), $grin2a^{-/-} b^{-/-}$ (transfection +Cre, double KO or DKO) neurons, or DKO neurons with aCSF supplemented with selective GluN2C/D antagonist DQP-1105 (10 µM). The residual current in the DKO was effectively blocked by DQP-1105. **c)** NMDA-EPSC$_{+20 mV}$ peak amplitudes in $grin2a^{fl/fl} b^{fl/fl}$ (untransfected) neurons and $grin2a^{-/-} b^{-/-}$ neurons rescued with human GluN2A WT, LOF (C436R, T531M or R518H) or GOF (K669N or L812M) mutants (transfected). **d)** NMDA-EPSC$_{+20 mV}$ decay time constant (tau) in $grin2a^{fl/fl} b^{fl/fl}$ (untransfected) neurons, and $grin2a^{-/-} b^{-/-}$ neurons rescued with human GluN2A WT, GOF (K669N and L812M) mutants or GluN2B (transfected). **b-d i)**. Representative NMDA-EPSCs from transfected neurons (and untransfected neuron in **b**) for each experimental group **ii)** Data points of measurements made in individual neurons. Matched data points, for simultaneously recorded untransfected and transfected neurons, are connected by a line. **iii)** Response ratios (transfected/untransfected) are expressed as a percentage and plotted for each pair of transfected-untransfected neurons. Crossbars in **ii)** and **iii)** show the estimated marginal means with 95% confidence intervals backtransformed from the linear mixed models (Figure S1.1 and 1.2 for c and d respectively). Hypothesis tests in: **ciii)** are orthogonal contrasts based on *a priori* clustering of the mutations in Figure 1e (Table S1); and in **diii)** are *posthoc* pairwise comparisons using Westfall stepwise procedure. Standardized effect sizes (*r*) for comparisons of each mutant with WT for response ratios of: **ciii)** peak amplitude were -.55, -.46, -.50, -.28 and -.26 for mutants C436R, T531M, R518H, K669N and L812M respectively (*N* = 93); or of **diii)** decay tau were .60 and .88 for mutants K669N and L812M respectively (*N* = 52).

Figure 3: Loss-of-function GluN2A mutants vary in their ability to deliver NMDA

receptors to synapses

a) Representative images of cultured neurons overexpressing GluN2A, SEP-GluN1 (green)

and Homer1c-tdTomato (red). The latter was used as a synaptic marker. Arrow heads point

to examples ofsynapses that have high synaptic levels of SEP-GluN1. Circles highlight

examples of synapses that exhibit low-to-moderate levels of SEP-GluN1. **b)** Levels of synaptic

enrichment of NMDA receptors and Homer1c in cultured hippocampal neurons transfected

with or without human GluN2A WT, LOF (C436R, T531M or R518H) or GOF (K669N or

L812M) mutants. Each data point represents mean fluorescence intensity of SEP-GluN1

(green) or Homer1c-tdTomato (red) across all visible putative synaptic regions-of-interest

(ROIs) in a neuron. **b i)** Data points for SEP-GluN1 (green) and Homer1c-tdTomato (red)

fluorescence measured from the same neuron are connected by a line. **b ii)** Response ratios

(SEP-GluN1 / Homer1C-tdTomato) are expressed as a percentage and plotted as a point for

each neuron. Crossbars in **i)** and **ii)** show the estimated marginal means with 95% confidence

intervals backtransformed from the fitted linear mixed models (Figure S1.3). Hypothesis

tests in **bii** are orthogonal contrasts based on *a priori* clustering of the mutations in Figure 1e

(Table S1). Standardized effect sizes (*r*) for comparisons of each mutant with WT were -.39, -

.26, -.03, .06 and -.20 for mutants C436R, T531M, R518H, K669N and L812M respectively (*N*

= 191).

Figure 4: Gain- and loss-of-function LBD mutants are both associated with more

prolonged NMDAR-mediated EPSCs

NMDA-EPSC$_{+20\,mV}$ peak amplitudes (**a**), decay time constants (**b**), or charge transfer (**c**) in

grin2a $^{fl/fl}$ (untransfected) neurons, and *grin2a* $^{-/-}$ neurons rescued with human GluN2A WT,

LOF (C436R, T531M or R518H) or GOF (K669N or L812M) mutants (transfected).

a-c i). Representative NMDA-EPSCs from transfected neurons for each experimental group.

ii) Data points of measurements made in individual neurons. Matched data points, for

simultaneously recorded untransfected and transfected neurons, are connected by a line. **iii)**

Response ratios (transfected/untransfected) are expressed as a percentage and plotted for

each pair of transfected-untransfected neurons. Crossbars in **ii)** and **iii)** show the estimated

marginal means with 95% confidence intervals backtransformed from the linear mixed

models (Figure S1.4-S1.6). Hypothesis tests are orthogonal contrasts based on *a priori*

clustering of the mutations in Figure 1e (and Table S1). Standardized effect sizes (*r*) for comparisons of each mutant with WT for response ratios of: **aiii)** peak amplitude were -.12, -.12, -.16, -.09 and -.06; **biii)** decay tau were .46, .49, .46, .26 and .28; and **ciii)** charge transfer were .20, .24, .10, .11 and .06, for mutants C436R, T531M, R518H, K669N and L812M respectively (*N* = 125).

Figure 5: Dose-dependent effects of grin2a null alleles on NMDA-EPSC kinetics.

NMDA-EPSC$_{+20\text{ mV}}$ charge transfer (**a**), peak amplitude (**b**), decay time constant (**c**), or 20-80% risetime (**d**) in *grin2a$^{+/+}$, grin2a$^{+/fl}$ or grin2a $^{fl/fl}$* (untransfected) neurons and *grin2a$^{+/+}$, grin2a$^{+/-}$ or grin2a $^{-/-}$* neurons (transfected, with Cre-GFP). **a-d i)**. Representative NMDA-EPSCs from transfected neurons for each experimental group. **ii)** Data points of measurements made in individual neurons. Matched data points, for simultaneously recorded untransfected and transfected neurons, are connected by a line. **iii)** Response ratios (transfected/untransfected) are expressed as a percentage and plotted for each pair of transfected-untransfected neurons. Crossbars in **i)** and **ii)** show the estimated marginal means with 95% confidence intervals backtransformed from the fitted linear mixed models (Figure S1.8-S1.11). Hypothesis tests are *posthoc* pairwise comparisons using Westfall stepwise procedure. Standardized effect sizes (*r*) for comparisons of each genotype with WT for response ratios of: **aiii)** charge transfer were .06 and .16; **biii)** peak amplitude were .00 and -.35; **ciii)** decay tau were .28 and .77; and **diii)** risetime were .22 and .68, for heterozygous and homozygous genotypes respectively (*N* = 99).

Figure 6: Spine calcium dynamics, temporal summation and spiking output during EPSP trains are shaped by mutation-associated changes in NMDA-EPSCs

ai) Model of the CA1 pyramidal neuron[68] used for the simulations. **aii)** The kinetics of the NMDAR-component of the synaptic conductance was slowed by 150% and 200% and the amplitude of the NMDAR-component was reduced to equalise charge transfer to mimic the effects of GOF (red) and LOF (blue) mutants. **aiii)** Bar graph illustrating the constant charge transfer during the EPSCs of varying kinetics to mimic GOF and LOF mutants. **b)** Current scapes for 40 activated excitatory synapses stimulated 5 times at 200 Hz. The NMDA component of the synaptic conductance had WT-like (100%), GOF-like (150%) or LOF-like (200%) kinetics with synaptic weight compensated to normalize charge transfer. From top to

bottom, current scapes show the membrane potential at the soma and ensemble mean membrane potential, outward current (amplitude and %), inward current (% and amplitude) and intracellular calcium concentration in spine heads at activated synapses. **c)** Data points overlaying bar charts of mean integral of soma membrane potential (**i**) and spine head membrane potential (**ii**) or intracellular calcium (**iii**) for 40 excitatory synapses stimulated 5 times at 20, 40, 80 and 200 Hz. Lines connecting the points reflect data simulated with the same random seed (n = 10). Numbers labelling bars in **i** represent mean number of action potentials per train. **d)** Current scapes as in **b** but for 200 activated excitatory synapses stimulated 5 times at 80 Hz. **e)** As in c but for 200 activated excitatory synapses.

Figure 7: Model summarizing how different molecular defects can converge on similar NMDAR-mediated EPSCs

a) WT GluN2A and GluN2B mostly contribute to NMDA-EPSCs in CA1 pyramidal neurons. GluN2B-containing NMDARs have more prolonged time course and cation influx as illustrated by the bold purple arrows. **b)** GluN2A mutants with defects culminating in low levels of GluN2A at synapses result in NMDA-EPSCs that are dominated by residual GluN2B-containing NMDARs. **c)** GluN2A mutants that traffic to the synapse but are functionally defective result in NMDA-EPSCs that are dominated by residual GluN2B-containing NMDARs. **d)** GluN2A mutants that traffic to the synapse but have slower deactivation kinetics directly contribute to prolonging NMDA-EPSCs. **b-d)** The effect of mutations may result in a combination of these different mechanisms.

References

1. Zito, K. & Scheuss, V. NMDA Receptor Function and Physiological Modulation. in *Encyclopedia of Neuroscience* (ed. Squire, L. R.) 1157–1164 (Academic Press, 2009). doi:10.1016/B978-008045046-9.01225-0.

2. Forrest, D. *et al.* Targeted disruption of NMDA receptor 1 gene abolishes NMDA response and results in neonatal death. *Neuron* **13**, 325–338 (1994).

3. Hamada, S. *et al.* The glutamate receptor GluN2 subunit regulates synaptic trafficking of AMPA receptors in the neonatal mouse brain. *Eur. J. Neurosci.* **40**, 3136–3146 (2014).

4. Kutsuwada, T. *et al.* Impairment of suckling response, trigeminal neuronal pattern formation, and hippocampal LTD in NMDA receptor epsilon 2 subunit mutant mice. *Neuron* **16**, 333–344 (1996).

5. Lai, K.-O. & Ip, N. Y. Structural plasticity of dendritic spines: The underlying mechanisms and its dysregulation in brain disorders. *Biochim. Biophys. Acta BBA - Mol. Basis Dis.* **1832**, 2257–2263 (2013).

6. Antic, S. D., Zhou, W.-L., Moore, A. R., Short, S. M. & Ikonomu, K. D. The decade of the dendritic NMDA spike. *J. Neurosci. Res.* **88**, 2991–3001 (2010).

7. Hunt, D. L. & Castillo, P. E. Synaptic plasticity of NMDA receptors: mechanisms and functional implications. *Curr. Opin. Neurobiol.* **22**, 496–508 (2012).

8. Major, G., Larkum, M. E. & Schiller, J. Active properties of neocortical pyramidal neuron dendrites. *Annu. Rev. Neurosci.* **36**, 1–24 (2013).

9. Nicoll, R. A. A Brief History of Long-Term Potentiation. *Neuron* **93**, 281–290 (2017).

10. Paoletti, P., Bellone, C. & Zhou, Q. NMDA receptor subunit diversity: impact on receptor properties, synaptic plasticity and disease. *Nat. Rev. Neurosci.* **14**, 383–400 (2013).

11. Shipton, O. A. & Paulsen, O. GluN2A and GluN2B subunit-containing NMDA receptors in hippocampal plasticity. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **369**, 20130163 (2014).

12. Morris, R. G. M. NMDA receptors and memory encoding. *Neuropharmacology* **74**, 32–40 (2013).

13. Higley, M. J. & Sabatini, B. L. Calcium Signaling in Dendritic Spines. *Cold Spring Harb. Perspect. Biol.* **4**, a005686 (2012).

14. Cull-Candy, S., Brickley, S. & Farrant, M. NMDA receptor subunits: diversity, development and disease. *Curr. Opin. Neurobiol.* **11**, 327–335 (2001).

15. Crawley, O., Conde-Dusman, M. J. & Pérez-Otaño, I. GluN3A NMDA receptor subunits: more enigmatic than ever? *J. Physiol.* **n/a**,.

16. Laube, B., Hirai, H., Sturgess, M., Betz, H. & Kuhse, J. Molecular determinants of agonist discrimination by NMDA receptor subunits: analysis of the glutamate binding site on the NR2B subunit. *Neuron* **18**, 493–503 (1997).

17. Vicini, S. *et al.* Functional and Pharmacological Differences Between RecombinantN-Methyl-d-Aspartate Receptors. *J. Neurophysiol.* **79**, 555–566 (1998).

18. Wyllie, D. J. A., Livesey, M. R. & Hardingham, G. E. Influence of GluN2 subunit identity on NMDA receptor function. *Neuropharmacology* **74**, 4–17 (2013).

19. Monyer, H., Burnashev, N., Laurie, D. J., Sakmann, B. & Seeburg, P. H. Developmental and regional expression in the rat brain and functional properties of four NMDA receptors. *Neuron* **12**, 529–540 (1994).

20. Scherzer, C. R. *et al.* Expression of N-Methyl-D-Aspartate receptor subunit mRNAs in the human brain: Hippocampus and cortex. *J. Comp. Neurol.* **390**, 75–90 (1998).

21. Chipman, P. H. *et al.* Astrocyte GluN2C NMDA receptors control basal synaptic strengths of hippocampal CA1 pyramidal neurons in the stratum radiatum. *eLife* **10**, e70818 (2021).

22. Gray, J. A. *et al.* Distinct modes of AMPA receptor suppression at developing synapses by GluN2A and GluN2B: single-cell NMDA receptor subunit deletion in vivo. *Neuron* **71**, 1085–1101 (2011).

23. Hestrin, S. Developmental regulation of NMDA receptor-mediated synaptic currents at a central synapse. *Nature* **357**, 686–689 (1992).

24. Liu, X.-B., Murray, K. D. & Jones, E. G. Switching of NMDA receptor 2A and 2B subunits at thalamic and cortical synapses during early postnatal development. *J. Neurosci. Off. J. Soc. Neurosci.* **24**, 8885–8895 (2004).

25. Sheng, M., Cummings, J., Roldan, L. A., Jan, Y. N. & Jan, L. Y. Changing subunit composition of heteromeric NMDA receptors during development of rat cortex. *Nature* **368**, 144–147 (1994).

26. Barria, A. & Malinow, R. Subunit-specific NMDA receptor trafficking to synapses. *Neuron* **35**, 345–353 (2002).

27. Bellone, C. & Nicoll, R. A. Rapid Bidirectional Switching of Synaptic NMDA Receptors. *Neuron* **55**, 779–785 (2007).

28. Groc, L. *et al.* NMDA receptor surface mobility depends on NR2A-2B subunits. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 18769–18774 (2006).

29. Carmignoto, G. & Vicini, S. Activity-dependent decrease in NMDA receptor responses during development of the visual cortex. *Science* **258**, 1007–1011 (1992).

30. Philpot, B. D., Sekhar, A. K., Shouval, H. Z. & Bear, M. F. Visual experience and deprivation bidirectionally modify the composition and function of NMDA receptors in visual cortex. *Neuron* **29**, 157–169 (2001).

31. Quinlan, E. M., Philpot, B. D., Huganir, R. L. & Bear, M. F. Rapid, experience-dependent expression of synaptic NMDA receptors in visual cortex in vivo. *Nat. Neurosci.* **2**, 352–357 (1999).

32. Roberts, E. B. & Ramoa, A. S. Enhanced NR2A Subunit Expression and Decreased NMDA Receptor Decay Time at the Onset of Ocular Dominance Plasticity in the Ferret. *J. Neurophysiol.* **81**, 2587–2591 (1999).

33. Sakimura, K. *et al.* Reduced hippocampal LTP and spatial learning in mice lacking NMDA receptor epsilon 1 subunit. *Nature* **373**, 151–155 (1995).

34. Salmi, M. *et al.* Impaired vocal communication, sleep-related discharges, and transient alteration of slow-wave sleep in developing mice lacking the GluN2A subunit of N-methyl-d-aspartate receptors. *Epilepsia* **60**, 1424–1437 (2019).

35. Salmi, M. *et al.* Transient microstructural brain anomalies and epileptiform discharges in mice defective for epilepsy and language-related NMDA receptor subunit gene Grin2a. *Epilepsia* **59**, 1919–1930 (2018).

36. Kiyama, Y. *et al.* Increased thresholds for long-term potentiation and contextual learning in mice lacking the NMDA-type glutamate receptor epsilon1 subunit. *J. Neurosci.* **18**, 6704–6712 (1998).

37. Myers, K. A., Johnstone, D. L. & Dyment, D. A. Epilepsy genetics: Current knowledge, applications, and future directions. *Clin. Genet.* **95**, 95–111 (2019).

38. Endele, S. *et al.* Mutations in GRIN2A and GRIN2B encoding regulatory subunits of NMDA receptors cause variable neurodevelopmental phenotypes. *Nat. Genet.* **42**, 1021–1026 (2010).

39. Firth, H. V. *et al.* DECIPHER: Database of Chromosomal Imbalance and Phenotype in Humans Using Ensembl Resources. *Am. J. Hum. Genet.* **84**, 524–533 (2009).

40. Reutlinger, C. *et al.* Deletions in 16p13 including GRIN2A in patients with intellectual disability, various dysmorphic features, and seizure disorders of the rolandic region. *Epilepsia* **51**, 1870–1873 (2010).

41. Myers, K. A. & Scheffer, I. E. GRIN2A-Related Speech Disorders and Epilepsy. in *GeneReviews®* (eds. Adam, M. P. et al.) (University of Washington, Seattle, 2016).

42. Strehlow, V. *et al.* GRIN2A-related disorders: genotype and functional consequence predict phenotype. *Brain* **142**, 80–92 (2019).

43. XiangWei, W., Jiang, Y. & Yuan, H. De novo mutations and rare variants occurring in NMDA receptors. *Curr. Opin. Physiol.* **2**, 27–35 (2018).

44. Carvill, G. L. *et al.* GRIN2A mutations cause epilepsy-aphasia spectrum disorders. *Nat. Genet.* **45**, 1073–1076 (2013).

45. Lemke, J. R. *et al.* Mutations in GRIN2A cause idiopathic focal epilepsy with rolandic spikes. *Nat. Genet.* **45**, 1067–1072 (2013).

46. Lesca, G. *et al.* GRIN2A mutations in acquired epileptic aphasia and related childhood focal epilepsies and encephalopathies with speech and language dysfunction. *Nat. Genet.* **45**, 1061–1066 (2013).

47. Turner, S. J., Morgan, A. T., Perez, E. R. & Scheffer, I. E. New genes for focal epilepsies with speech and language disorders. *Curr. Neurol. Neurosci. Rep.* **15**, 35 (2015).

48. Yang, X. *et al.* GRIN2A mutations in epilepsy-aphasia spectrum disorders. *Brain Dev.* **40**, 205–210 (2018).

49. Gobbi, G., Boni, A. & Filippini, M. The spectrum of idiopathic Rolandic epilepsy syndromes and idiopathic occipital epilepsies: from the benign to the disabling. *Epilepsia* **47 Suppl 2**, 62–66 (2006).

50. Addis, L. *et al.* Epilepsy-associated GRIN2A mutations reduce NMDA receptor trafficking and agonist potency - molecular profiling and functional rescue. *Sci. Rep.* **7**, 66 (2017).

51. CFERV. Center for Functional Evaluation of Rare variants. http://functionalvariants.emory.edu/index.html (2017).

52. Chen, W. *et al.* Functional Evaluation of a De Novo GRIN2A Mutation Identified in a Patient with Profound Global Developmental Delay and Refractory Epilepsy. *Mol. Pharmacol.* **91**, 317–330 (2017).

53. Gao, K. *et al.* A de novo loss-of-function GRIN2A mutation associated with childhood focal epilepsy and acquired epileptic aphasia. *PLoS One* **12**, e0170818 (2017).

54. Ogden, K. K. *et al.* Molecular Mechanism of Disease-Associated Mutations in the Pre-M1 Helix of NMDA Receptors and Potential Rescue Pharmacology. *PLoS Genet.* **13**, e1006536 (2017).

55. Sibarov, D. A. *et al.* Functional Properties of Human NMDA Receptors Associated with Epilepsy-Related Mutations of GluN2A Subunit. *Front. Cell. Neurosci.* **11**, 155 (2017).

56. Swanger, S. A. *et al.* Mechanistic Insight into NMDA Receptor Dysregulation by Rare Variants in the GluN2A and GluN2B Agonist Binding Domains. *Am. J. Hum. Genet.* **99**, 1261–1280 (2016).

57. Yuan, H. *et al.* Functional analysis of a de novo GRIN2A missense mutation associated with early-onset epileptic encephalopathy. *Nat. Commun.* **5**, 3251 (2014).

58. Serraz, B., Grand, T. & Paoletti, P. Altered zinc sensitivity of NMDA receptors harboring clinically-relevant mutations. *Neuropharmacology* **109**, 196–204 (2016).

59. Hansen, K. B. *et al.* Structure, Function, and Pharmacology of Glutamate Receptor Ion Channels. *Pharmacol. Rev.* **73**, 298–487 (2021).

60. Madry, C., Mesic, I., Betz, H. & Laube, B. The N-terminal domains of both NR1 and NR2 subunits determine allosteric Zn2+ inhibition and glycine affinity of N-methyl-D-aspartate receptors. *Mol. Pharmacol.* **72**, 1535–1544 (2007).

61. Puddifoot, C. A., Chen, P. E., Schoepfer, R. & Wyllie, D. J. A. Pharmacological characterization of recombinant NR1/NR2A NMDA receptors with truncated and deleted carboxy termini expressed in Xenopus laevis oocytes. *Br. J. Pharmacol.* **156**, 509–518 (2009).

62. Granger, A. J., Gray, J. A., Lu, W. & Nicoll, R. A. Genetic analysis of neuronal ionotropic glutamate receptor subunits. *J. Physiol.* **589**, 4095–4101 (2011).

63. Akashi, K. *et al.* NMDA receptor GluN2B (GluR epsilon 2/NR2B) subunit is crucial for channel function, postsynaptic macromolecular organization, and actin cytoskeleton at hippocampal CA3 synapses. *J. Neurosci. Off. J. Soc. Neurosci.* **29**, 10869–10882 (2009).

64. Nagy, A. Cre recombinase: The universal reagent for genome tailoring. *genesis* **26**, 99–109 (2000).

65. Horak, M., Petralia, R. S., Kaniakova, M. & Sans, N. ER to synapse trafficking of NMDA receptors. *Front. Cell. Neurosci.* **8**, 394 (2014).

66. Kopec, C. D., Li, B., Wei, W., Boehm, J. & Malinow, R. Glutamate receptor exocytosis and spine enlargement during chemically induced long-term potentiation. *J. Neurosci.* **26**, 2000–2009 (2006).

67. Adesnik, H., Li, G., During, M. J., Pleasure, S. J. & Nicoll, R. A. NMDA receptors inhibit synapse unsilencing during brain development. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 5597–5602 (2008).

68. Graham, B. P., Saudargiene, A. & Cobb, S. Spine head calcium as a measure of summed postsynaptic activity for driving synaptic plasticity. *Neural Comput.* **26**, 2194–2222 (2014).

69. Moradi, K. *et al.* A fast model of voltage-dependent NMDA receptors. *J. Comput. Neurosci.* **34**, 521–531 (2013).

70. Alonso, L. M. & Marder, E. Visualization of currents in neural models with similar behavior and different conductance densities. *eLife* **8**, e42722 (2019).

71. Keith, R. E. *et al.* Direct Intracellular Signaling by the Carboxy terminus of NMDA Receptor GluN2 Subunits Regulates Dendritic Morphology in Hippocampal CA1 Pyramidal Neurons. *Neuroscience* **396**, 138–153 (2019).

72. Regan, M. C. *et al.* STRUCTURAL MECHANISM OF FUNCTIONAL MODULATION BY GENE SPLICING IN NMDA RECEPTORS. *Neuron* **98**, 521-529.e3 (2018).

73. Yi, F., Zachariassen, L. G., Dorsett, K. N. & Hansen, K. B. Properties of Triheteromeric N-Methyl-d-Aspartate Receptors Containing Two Distinct GluN1 Isoforms. *Mol. Pharmacol.* **93**, 453–467 (2018).

74. Booker, S. A., Sumera, A., Kind, P. C. & Wyllie, D. J. A. Contribution of NMDA Receptors to Synaptic Function in Rat Hippocampal Interneurons. *eNeuro* **8**, (2021).

75. Rathenberg, J., Nevian, T. & Witzemann, V. High-efficiency transfection of individual neurons using modified electrophysiology techniques. *J. Neurosci. Methods* **126**, 91–98 (2003).

76. Campagnola, L., Kratz, M. B. & Manis, P. B. ACQ4: an open-source software platform for data acquisition and analysis in neurophysiology research. *Front. Neuroinformatics* **8**, 3 (2014).

77. Jiang, M. & Chen, G. High Ca2+-phosphate transfection efficiency in low-density neuronal cultures. *Nat. Protoc.* **1**, 695–700 (2006).

78. Traynelis, S. F. Software-based correction of single compartment series resistance errors. *J. Neurosci. Methods* **86**, 25–34 (1998).

79. Guzman, S. J., Schlögl, A. & Schmidt-Hieber, C. Stimfit: quantifying electrophysiological data with Python. *Front. Neuroinformatics* **8**, 16 (2014).

80. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).

81. Rueden, C. T. *et al.* ImageJ2: ImageJ for the next generation of scientific image data. *BMC Bioinformatics* **18**, 529 (2017).

82. Carnevale, N. T. & Hines, M. L. *The NEURON Book*. (Cambridge University Press, 2006).

83. Hubert, M., Rousseeuw, P. J. & Branden, K. V. ROBPCA: A New Approach to Robust Principal Component Analysis: Technometrics: Vol 47, No 1. (2005).

84. Verboven, S. & Hubert, M. LIBRA: a MATLAB library for robust analysis. *Chemom. Intell. Lab. Syst.* **75**, 127–136 (2005).

85. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).

86. Penn, A. C. *iboot: Iterated bootstrap for small samples and samples with complex dependence structures [https://github.com/acp29/iboot]*. (Zenodo, 2020). doi:10.5281/zenodo.3992392.

Figure 1

a

Nonsense
Frameshift
Del/Dup
Splice site
Missense
Other

3%
15%
17%
9%
14%
42%

b

NTD
LBD
CHANNEL

GluN1   GluN2

GluN2A
Glu
Gating ring

c

LOF   WT   K669N   GOF

57%   Glu potency   43%
52%   Gly potency   48%
41%   Deactivation   59%
63%   Open probability   37%
79%   Current density   21%
77%   Surface expression   23%

Log10 (Normalized response)

d

Factor 1   Factor 2

Loadings

1) $EC_{50,Glu}^{-1}$
2) $EC_{50,Gly}^{-1}$
3) τ_{deact}
4) I_{peak}
5) P_o
6) Surf. expr.

e

WT   GOF

$\tau_b = 0.49\ [0.11\text{-}0.71]$
$p = 0.015$

WT
K669N
WT
L812M
GOF Type 1

LOF Type 1
R518H

LOF Type 2
T531M   C436R

LOF

Factor 2

Factor 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

a **GluN2A wildtype**

b **GluN2A mutant**
Trafficking defect
(e.g. C436R)

c **GluN2A mutant**
Functional defect
(e.g. R518H)

d **GluN2A mutant**
Prolong channel opening
(e.g. K669N)

Neuron A

Na⁺
Ca²⁺

ER

Neuron B

NMDAR-EPSC

GluN1 / GluN2A   GluN1 / GluN2B   • Glutamate   ★ Mutation