TITLE

PELP1/SRC-3-dependent regulation of metabolic kinases drives therapy resistant ER+ breast cancer

AUTHORS

Thu H. Truong[1], Elizabeth A. Benner[1], Kyla M. Hagen[1], Nuri A. Temiz[1,4], Carlos Perez Kerkvliet[1], Ying Wang[1], Emilio Cortes-Sanchez[6,7,8], Chieh-Hsiang Yang[6,7,8], Thomas Pengo[5], Katrin P. Guillen[6,8], Bryan E. Welm[6,7,8], Sucheta Telang[9], *Carol A. Lange[1,2,3], and *Julie H. Ostrander[1,2]

* co-senior authors

AFFILIATIONS

[1] Masonic Cancer Center, [2] Department of Medicine (Division of Hematology, Oncology, and Transplantation), [3] Department of Pharmacology, [4] Institute for Health Informatics, and [5] University of Minnesota Informatics Institute, University of Minnesota, Minneapolis MN 55455 USA

[6] Department of Oncological Sciences, [7] Department of Surgery, and [8] Huntsman Cancer Institute, University of Utah, Salt Lake City UT 84112,

[9] James Graham Brown Cancer Center, Department of Medicine (Division of Medical Oncology and Hematology), University of Louisville, Louisville KY 40202 USA

RUNNING TITLE

PELP1/SRC-3 regulate metabolic kinases in ER+ breast cancer

KEYWORDS

PELP1, SRC-3, cancer stem cells

FINANCIAL SUPPORT

This work was supported by NIH grants R01 CA236948 (JHO, CAL), F32 CA210340 (THT), T32 HL007741 (THT), and U54 CA224076 (BEW), ACS Institutional Research Grant #124166-IRG-58-001-52-IRG5 (JHO), University of Minnesota Masonic Cancer Center (CAL, JHO), the Tickle Family Land Grant Endowed Chair in Breast Cancer Research (CAL), National Center for Advancing Translational Sciences of the NIH Award UL1TR000114 (JHO), and Department of Defense W81XWH-14-1-0417 (BEW).

CORRESPONDING AUTHORS

Julie H. Ostrander

hans1354@umn.edu (email), 612-625-1996 (phone)

University of Minnesota Masonic Cancer Center

Cancer and Cardiovascular Research Building

Delivery Code 2812

2231 6th St SE, Minneapolis MN 55455, USA

Carol A. Lange

lange047@umn.edu (email), 612-626-0621 (phone)

University of Minnesota Masonic Cancer Center

Cancer and Cardiovascular Research Building

Delivery Code 2812

2231 6th St SE, Minneapolis MN 55455, USA

CONFLICT OF INTEREST

Carol A. Lange is a Scientific Advisory Board Member for Context Therapeutics, Inc. The remaining authors have nothing to disclose.

WORD COUNT: 5,814

FIGURES: 7

TABLES: 0

ABSTRACT

Recurrence of metastatic breast cancer stemming from acquired endocrine and chemotherapy resistance remains a health burden for women with luminal (ER+) breast cancer. Disseminated ER+ tumor cells can remain viable but quiescent for years to decades. Contributing factors to metastatic spread include the maintenance and expansion of breast cancer stem cells (CSCs). Breast CSCs frequently exist as a minority population in therapy resistant tumors. In this study, we show that cytoplasmic complexes composed of steroid receptor (SR) co-activators, PELP1 and SRC-3, modulate breast CSC expansion through upregulation of the HIF-activated metabolic target genes *PFKFB3* and *PFKFB4*. Seahorse metabolic assays demonstrated that cytoplasmic PELP1 influences cellular metabolism by increasing both glycolysis and mitochondrial respiration. PELP1 interacts with PFKFB3 and PFKFB4 proteins, and inhibition of PFKFB3 and PFKFB4 kinase activity blocks PELP1-induced tumorspheres and protein-protein interactions with SRC-3. PFKFB4 knockdown inhibited *in vivo* emergence of circulating tumor cell (CTC) populations in mammary intraductal (MIND) models. Application of PFKFB inhibitors in combination with ER targeted therapies blocked tumorsphere formation in multiple models of advanced breast cancer, including tamoxifen (TamR) and paclitaxel (TaxR) resistant models and ER+ patient-derived organoids (PDxO). Together, our data suggest that PELP1, SRC-3, and PFKFBs cooperate to drive ER+ tumor cell populations that include CSCs and CTCs.

Significance: Identifying non-ER pharmacological targets offers a useful approach to blocking metastatic escape from standard of care ER/estrogen (E2)-targeted strategies to overcome endocrine and chemotherapy resistance.

INTRODUCTION

Metastatic recurrence is an incurable but common complication of ER+ breast cancer. Treatment of metastatic breast cancer typically results in endocrine resistance, and chemotherapy is largely ineffective in advanced disease. Altered signaling pathways drive therapy resistance and offer potential targets for metastatic ER+ breast cancer. PELP1 (proline, glutamic acid, leucine-rich protein 1) and SRC-3 (steroid receptor [SR] co-activator-3) have independently been shown to drive endocrine resistance. PELP1 and SRC-3 are both SR co-activators involved in normal development and cancer (1,2). Increased PELP1 expression is associated with higher tumor grade, tumor proliferation, and decreased breast cancer-specific survival (3,4). PELP1 is primarily nuclear in normal breast tissue; however, altered cytoplasmic PELP1 localization is observed in 40-58% of PELP1+ breast tumors (5). Analysis of breast tumor samples revealed that patients

with high cytoplasmic PELP1 levels were less likely to respond to tamoxifen (tam) (4). Similarly,

SRC-3 mRNA and protein overexpression are correlated with higher tumor grade and decreased

overall and disease-free survival (6). SRC-3 overexpression is also linked to tam resistance in

breast cancer models and human breast tumors (7,8). Both PELP1 and SRC-3 have essential

nuclear functions, but also dynamically shuttle to the cytoplasm where they associate with

signaling molecules and act as scaffolds for growth factor or SR pathways. These SR co-

activators have emerged as promising targets in ER+ breast cancer and as potential mediators

of therapy resistance.

Cancer stem cells (CSCs) are poorly proliferative and frequently exist as a minority sub-population

of cells that drive therapy resistance and metastasis (9). In contrast to non-CSCs, breast CSCs

form colonies in serum-free suspension culture (i.e. tumorspheres), express stem cell markers

(e.g. ALDH+ or CD44^{hi}/CD24^{lo}), exhibit enhanced resistance to chemo and endocrine therapies,

and express markers of epithelial to mesenchymal transition (EMT). The ability to survive and

self-renew following treatment allows CSCs to evade standard chemo and endocrine therapies

aimed at rapidly dividing cancer cells and to drive metastatic tumor growth.

Growing evidence has implicated SR co-activators as mediators of CSC self-renewal. For

example, SRC-3 drives CSC formation and tumor outgrowth in breast cancer models. Treatment

with SI-2, an SRC-3 inhibitor, decreased SRC-3-induced CSC formation in breast cancer cell and

xenograft models (10). Our laboratory reported that cytoplasmic complexes composed of PELP1

and SRC-3 mediate breast CSC expansion (11). Targeting SRC-3 using shRNA or

pharmacological inhibitors (i.e. SI-2) abrogated PELP1/SRC-3 complex formation, PELP1-

induced tumorspheres, and expression of PELP1 target genes that promote cancer cell survival.

These studies imply that inhibiting PELP1 and its binding partners may provide a way to target

the breast CSC population in order to improve patient outcomes.

Herein we sought to identify the molecular mechanisms that contribute to PELP1-driven CSC

survival and self-renewal in ER+ breast cancer. Using endocrine and chemotherapy resistant

breast cancer models, our findings suggest that PELP1/SRC-3 complexes modulate the CSC

compartment through gene programs associated with metabolic adaptation. In contrast to current

therapies that fail to adequately target slow-growing breast CSCs, our studies reveal therapy

combinations that inhibit cooperating signaling cascades, while simultaneously targeting ER. By

targeting CSCs directly, this approach promises to significantly improve the lives of patients with

recurrent ER+ breast cancer.

MATERIALS AND METHODS

Cell Culture. STR authentication was performed by ATCC (October 2018). MCF-7 PELP1 and

J110 cells were cultured as described (11). MCF-7 (12) and T47D TamR (13) cells were cultured

in 100 nM tamoxifen. MCF-7 TaxR (14) cells were cultured in 2 µM Taxol. For 3D (tumorsphere)

conditions, cells were cultured as described (11).

Patient-Derived Organoids (PDxO). PDxOs (HCI-003, -011, -017) were cultured in Advanced

DMEM/F12 (Thermo Fisher) containing 5% FBS, 1X HEPES, 1X GlutaMax, 50 µg/ml Gentamicin

(Genesee), 1 µg/ml hydrocortisone, 10 ng/ml EGF, and supplemented with 10 µM Y-27632

(Selleckchem), 100 ng/ml FGF2 (PeproTech), and 1 mM N-acetyl cysteine (Sigma). PDxOs were

embedded into Matrigel (growth factor reduced; Corning) using the hanging drop method into 6-

well plates and passaged every ~14-18 days.

Synergy Screens. PDxOs were grown as described above in 384-well plates: Prior to assay,

PDxOs were cultured in PDxO media containing charcoal stripped FBS and allowed to incubate

overnight. PDxOs were disbursed with Dispase and re-seeded in PDxO media on 384-well plates

coated with Matrigel. Following 24h after seeding, plates were dosed with SI-2 (0-2 µM), 5MPN

(0-50 µM), and tam (0-10 µM) using a robotic assisted pipette (ViaFlo). On Day 6 post-treatment,

a Cell Titer Glo 3D assay (Promega) was performed per manufacturer's instructions.

Luminescence readings were measured using an Envision Spectrophotometer. Fold changes

were calculated using Day 0 baseline readings. % survival was calculated for each well, and ZIP

synergy model (15) were run to determine synergy scores with the R package synergy finder (16).

Independent drug response was calculated using the four- parameter logistic model (4-PL). 3

biological replicates, each one consisting of 4 technical replicates per dosing, was performed.

Mammary Intraductal (MIND) Model. Intraductal injections of single cells were performed as

described (17,18). Seven-week old female NSG mice (#005557) were purchased from Jackson

Laboratory. Five mice/group were injected with 5×10^4 cells into each nipple of the 4th inguinal

glands with the indicated breast cancer cell line. Mammary glands were harvested 8 weeks after

injection, fixed in 4% PFA, and processed for H&E staining. H&E sections were analyzed using

ImageJ or Q-path to quantitate the total mammary gland area (%) that contained tumor cells.

Statistical Analysis. Data were tested for normal distribution using Shapiro-Wilks normality test

and homogeneity of variances using Bartlett's Test. Once data met these two requirements,

statistical analyses were performed using one-way or two-way ANOVA in conjunction with Tukey

multiple comparison test for means between more than two groups or Student *t* test for means

between two groups, where significance was determined with 95% confidence. For the MIND

study with four groups defined by two factors (cyto PELP1 vs. WT PELP1, and shPFKFB4 vs.

shGFP), a regression model identified a significant interaction due to shPFKFB4 at an alpha level

of 0.1 (p=0.084).

RESULTS

Cytoplasmic PELP1 promotes CSCs and HIF-regulated gene expression

Breast CSCs represent a minority of the total cell population (1-5%) (19), making it difficult to

detect CSC-specific changes in bulk tumor populations. We therefore measured breast CSC

frequency by comparing ALDH activity (**Figure 1A**, **Supplementary Figure 1**) and CD44^{hi}/CD24^{lo}

ratios (**Figure 1B**, **Supplementary Figure 2**) in MCF-7 cells stably expressing LXSN (vector

control), WT PELP1, or cytoplasmic (cyto) PELP1 cultured in either 2D (adherent) or 3D

(tumorsphere) conditions. Relative to 2D, 3D conditions increased breast CSC markers in MCF-

7 cells expressing LXSN, WT PELP1, or cyto PELP1 (**Figure 1A, 1B**). In 2D conditions, cyto

PELP1 expressing cells had no significant changes in ALDH activity when compared to LXSN or

WT PELP1; however, when the same models were cultured in 3D conditions, ALDH activity was

significantly increased in cells expressing cyto PELP1 (12.0% ± 2.9) compared to LXSN (6.6% ±

0.67, p = 0.023) and WT PELP1 (2.6% ± 0.76, p = 0.0015). In 2D conditions, CD44^{hi}/CD24^{lo}

populations were increased in cyto PELP1 expressing cells (13.0% ± 0.49) compared to LXSN

(2.6% ± 0.042, p < 0.0001) or WT PELP1 (1.2% ± 0.19, p < 0.0001), and this trend was enhanced

in 3D conditions (cyto PELP1, 19.4% ± 1.4; LXSN, 9.0% ± 1.1, p = 0.0045; WT PELP1, 2.3% ±

0.18, p = 0.0011). WT PELP1 displayed lower ALDH activity and CD44^{hi}/CD24^{lo} ratios relative to

LXSN controls, suggesting that nuclear PELP1 limits CSC behavior. These results indicate that

both 3D culture and cyto PELP1 expression independently increase CSC expansion in MCF-7

cell models.

We performed RNA-seq on MCF-7 PELP1 models grown as 3D tumorspheres and compared

these data to studies conducted in 2D culture (11) to identify candidate genes and pathways

differentially regulated in cyto PELP1 expressing cells. Comparison of 3D versus 2D conditions

identified 206 upregulated and 114 downregulated genes similarly regulated by >2-fold in all cell

lines (LXSN, WT PELP1, cyto PELP1) (**Figure 1C, Supplementary Figure 3**). Ingenuity Pathway

Analysis (IPA) of these 320 genes revealed activation of estrogen, growth factor, cytokine, and

NF-κB pathways (**Supplementary Table 1**). Significantly activated and inhibited "Diseases and

Functions" are summarized in **Supplementary Table 2**. 3D to 2D comparison in cyto PELP1

expressing cells identified 173 differentially expressed genes (93 upregulated, 80 downregulated)

compared to LXSN or WT PELP1 (**Figure 1C, Supplementary Figure 4**). These 173 genes were

analyzed with IPA to identify cyto PELP1-specific pathways (**Figure 1D**), biological functions, or

disease states (**Figure 1E**), and predicted increased HIF activation, estradiol, ATF4, and

glycolytic-mediated pathways. We created representative heatmaps to illustrate 3D-specific

regulation in upstream regulator analysis associated with HIF and ATF4 pathway activation (>2-

fold; **Figure 1F**) and generated a cyto PELP1 upregulated gene signature (**Supplementary Table

3**). Volcano plots of differentially regulated genes are shown in **Figure 1G**; red dots indicate genes

in the cyto PELP1 signature. We then used the cyto PELP1 upregulated gene signature to query

the METABRIC breast cancer database. Higher expression of this gene signature was associated

with lower overall survival (OS) in the METABRIC cohort (hazard ratio = 1.485, p < 0.0001, **Figure

1H**). We tested this on the ER+ only subtype within the METABRIC cohort and found similar

results (hazard ratio = 1.483, p < 0.0001, **Figure 1I**). A similar query of the TCGA database

revealed no significant differences in OS (**Supplementary Figure 5**). Taken together, these data

identify genes involved in cyto PELP1-mediated pathways that promote CSCs, including those

associated with HIF-activated and glycolytic pathways.

Cytoplasmic PELP1 drives metabolic plasticity

Given the strong activation of HIF and metabolic pathways detected in the RNA-seq analysis, we

used qPCR to test HIF-activated target genes. HIF activates the *PFKFB* family, which are

metabolic bi-functional kinase/phosphatases (20). We found that mRNA levels of *EPAS1* (i.e.

HIF2α), *PFKFB3*, and *PFKFB4* were upregulated in cells expressing cyto PELP1 relative to LXSN

or WT PELP1 in 3D, but not 2D conditions (**Figure 2A**). Additional validation of HIF-activated

metabolic and stem cell genes include *NDRG1* and *SOX9* (**Figure 2A**)**.** Given the central role of

HIF pathways in metabolism (21), we investigated the effect of PELP1 on metabolic pathways

using the Seahorse Cell Energy Phenotype test to measure oxygen consumption rate (OCR) and

extracellular acidification rate (ECAR). At baseline, MCF-7 cells expressing cyto PELP1 exhibited a significant increase in OCR levels compared to LXSN and WT PELP1. Under stressed conditions (i.e. after FCCP and oligomycin), OCR was increased in cyto PELP1 expressing cells compared to LXSN (p = 0.0096). ECAR was significantly different in cyto PELP1 expressing cells compared to LXSN at baseline, but WT and cyto PELP1 displayed an increase in ECAR compared to LXSN controls (p = 0.046 and 0.0045) under stressed conditions (**Figure 2B**). To systematically test effects on key parameters of mitochondrial function, we performed the Seahorse Mito Stress test. Cyto PELP1 expression significantly increased basal respiration, compared to LXSN and WT PELP1 (p < 0.0001 and 0.0001). Furthermore, cyto PELP1 increased ATP-linked respiration, proton leak, maximal respiration, and non-mitochondrial respiration (**Figure 2C**). Cyto PELP1 expressing cells had a 4-fold increase in glucose uptake compared to WT PELP1 and LXSN, as measured by 2-NBDG (**Figure 2D, Supplementary Figure 6**). Collectively, these results indicate cyto PELP1 drives HIF-activated metabolic programs (i.e. *PFKFB3*, *PFKFB4*) in 3D culture, and affects mitochondrial respiration and glycolysis, indicative of metabolic plasticity.

Inhibition of PFKFBs disrupts PELP1/SRC-3 complexes and tumorsphere formation

We hypothesized that HIF-activated targets PFKFB3 and PFKFB4 are required components of the PELP1/SRC-3 complex. Co-immunoprecipitation of PFKFB3 or PFKFB4 demonstrated increased association with PELP1 in cells expressing cyto PELP1 relative to LXSN or WT PELP1 (**Figure 3A, 3B**). Treatment with PFK158 and 5MPN, inhibitors of PFKFB3 and PFKFB4 respectively, reduced the PELP1/SRC-3 interaction (**Figure 3C, 3D**). These inhibitors also blocked PELP1/PFKFB3 and PELP1/PFKFB4 (**Supplementary Figure 7A, 7B**) interactions in cyto PELP1 expressing cells; similar results were observed with another PFKFB3 inhibitor (PFK15; **Supplementary Figure 7C, 7D**).

Next, we tested the effect of PFKFB inhibition on cyto PELP1-induced tumorspheres, an *in vitro* assay to assess breast CSC activity (11). PFKFB4 knockdown (**Supplementary Figure 8**) decreased tumorsphere formation in cyto PELP1 expressing cells by ~50%, but not in LXSN or WT PELP1 (**Figure 3E**, p = 0.0103). Attempts to stably knockdown PFKFB3 were not successful, suggesting that PFKFB3 is crucial for cell viability (22). Inhibitors of PFKFB3 and PFKFB4 reduced cyto PELP1-induced tumorspheres, but had no effect on cells expressing either LXSN or WT PELP1. (**Figure 3F, 3G; Supplementary Figure 7E**). To evaluate PFKFB inhibitors in an alternative PELP1/SRC-3 model, we used a murine tumor cell line (J110) established from the MMTV-SRC-3 mouse (23). Similar to MCF-7 PELP1 models, PFK158 or 5MPN inhibited

tumorsphere formation by ~40% in J110 cells (**Figure 3H**). Western blotting indicated that

PFKFB3 and PFKFB4 protein levels remained unchanged in response to E2, while ER levels

decreased, presumably due to ligand-induced turnover (**Figure 3H,** right). These results indicate

that blocking PFKFB3 or PFKFB4 through knockdown or pharmacological inhibition disrupts

expansion and self-renewal of PELP1-driven CSC populations.

PFKFB4 reduces *in vivo* expansion of CTCs in cyto PELP1 MIND xenografts

To evaluate if PELP1 promotes tumor formation *in vivo*, we injected MCF-7 WT and cyto PELP1

expressing cells (5×10^4) into the inguinal mammary glands of adult female mice (6-8 week old,

4 mice/group) to generate mammary intraductal (MIND) tumors. Both cell lines had 100%

engraftment rates (**Figure 4A, Supplementary Figure 9**). Tumor area (%) calculated from H&E

images of each mammary gland revealed increased tumor volume in cyto PELP1 (25.7% $\pm$ 16.5)

compared to WT PELP1 MIND xenografts (10.9% $\pm$ 9.5, p = 0.046) (**Figure 4B**).

Based on our *in vitro* data showing that inhibition of PFKFB4 (knockdown and 5MPN) decreased

tumorspheres, we queried PFKFB4 mRNA levels on OS in METABRIC datasets. High PFKFB4

mRNA expression is associated with decreased OS in all subtypes and ER+ only patient cohorts

(**Supplementary Figure 10**). Therefore, we tested whether PFKFB4 knockdown would impact

MIND tumor growth or the presence of circulating tumor cells (CTCs); a marker of metastatic

potential and associated CSC behavior (24). 5 mice/group were injected with MCF-7 WT or cyto

PELP1 expressing cells harboring either shGFP control or shPFKFB4. 8 weeks post-injection,

mammary glands were fixed and processed for H&E staining (**Supplementary Figure 11**). As in

Figure 4B, the difference in means between WT PELP1 shGFP (26.8% $\pm$ 10.2) and cyto PELP1

shGFP (41.2% $\pm$ 17.2) tumor area remained significant (p = 0.036, **Figure 4C**). However,

knockdown of PFKFB4 in MCF-7 cells expressing either WT PELP1 or cyto PELP1 failed to

significantly affect primary tumor growth. To assess disseminated tumor cells, blood samples

were collected during euthanization and seeded into soft agar assays to detect CTCs. Mice

injected with WT PELP1 (shGFP or shPFKFB4) expressing cells did not exhibit CTC colony

formation. In sharp contrast, blood samples from mice engrafted with cyto PELP1 cells developed

large viable colonies, indicating the presence of CTCs. Knockdown of shPFKFB4 in MCF-7 cyto

PELP1 expressing cells reduced both colony formation (p < 0.0492) and colony size (p < 0.0016)

(**Figure 4D-4F**). These data demonstrate a requirement for PFKFB4 in cyto PELP1-driven CTC

formation and expansion *in vivo*.

Targeting PELP1/SRC-3 complexes in therapy resistant breast cancer and PDxO models

Paclitaxel (Taxol) is a chemotherapy used to treat late stage breast cancer. Increased PELP1, HIF1α, and HIF-2α expression has been observed in triple negative breast cancer (TNBC) cells in response to Taxol (14). To evaluate whether PELP1 expression affects response to Taxol in ER+ breast cancer, we treated MCF-7 PELP1 cells (LXSN, WT PELP1, cyto PELP1) cultured as tumorspheres with Taxol (0 to 125 nM). We assessed tumorsphere formation and calculated IC50 values for each cell line (**Figure 5A**). IC50 (Taxol) for cyto PELP1 expressing cells was ~2-fold higher than LXSN or WT PELP1. These results suggest that cyto PELP1 expression confers enhanced Taxol resistance compared to LXSN or WT PELP1.

Next, we determined if PELP1/SRC-3 signaling mediates therapy resistance in tamoxifen resistant (TamR) and paclitaxel-resistant (TaxR) cell lines. HIF and cyto PELP1 regulated genes, *EPAS1, PFKFB3, and PFKFB4*, mRNA levels were increased in MCF-7 TaxR (**Figure 5B,** *top*) and TamR cells (**Figure 5B,** *bottom*) relative to MCF-7 parental controls, particularly in 3D conditions. 3D PELP1 target genes, *NDRG1* and *SOX9* were also upregulated in TaxR and TamR cells relative to parental MCF-7 cells (**Supplementary Figure 12**). To determine if similar changes in cellular metabolism occur in MCF-7 TaxR models, we performed Seahorse metabolic assays. The Cell Energy Phenotype test showed TaxR cells exhibit increased OCR and ECAR at baseline and stressed conditions relative to controls (**Figure 5C**), indicating increased mitochondrial respiration and glycolysis. To look at individual effects on OCR, we performed the Mito Stress test in MCF-7 TaxR models. Similar to cyto PELP1 expressing cells, TaxR cells showed significant increases in basal and maximal respiration compared to controls (**Figure 5D**). TaxR cells increased proton leak, spare respiratory capacity, and non-mitochondrial respiration, but not ATP production as observed in MCF-7 cyto PELP1 expressing cells. TaxR cells also displayed ~2-fold increase (p = 0.0006) in glucose uptake compared to controls (**Figure 5E**). Together, these data reveal that TamR and TaxR models phenocopy HIF-associated target gene expression and metabolic plasticity of MCF-7 cyto PELP1 expressing cells, and suggest PELP1 may be a key mediator in therapy resistance.

The PELP1/SRC-3 interaction was similarly increased in MCF-7 TaxR (**Figure 5F**, top) and TamR cells (**Figure 5F**, bottom). Additionally, CD44^{hi}/CD24^{lo} ratios were increased in MCF-7 TaxR cells compared to parental controls (**Supplementary Figure 13**). To test the pharmacological effect of PFKFB3, PFKFB4, and SRC-3 inhibition, MCF-7 TaxR and TamR cells were seeded as tumorspheres and treated with PFK158, 5MPN, and SI-2. Both resistant models exhibited

increased basal tumorsphere formation when compared to parental controls. 5MPN and SI-2
effectively decreased secondary tumorsphere formation by 71% and 75% in TaxR (**Figure 5G**),
and 88% and 92% in TamR models (**Figure 5H**) compared to vehicle controls. PFK158 (PFKFB3
inhibitor) modestly decreased TaxR and TamR tumorspheres by 17% and 27%. These findings
highlight the overlap of key players involved in PELP1-driven CSC biology and suggest that
PFKFB4 and SRC-3 play a more significant role than PFKFB3 within resistant cell models.
345
We hypothesized that tam in combination with PELP1/SRC-3 complex inhibitors (i.e. SI-2 or
5MPN) would be more effective than either inhibitor alone. Combination treatments were
evaluated in several cell lines. In MCF-7 PELP1 models, we tested tam/SI-2, tam/5MPN, and SI-
2/5MPN combinations (**Figure 6A-6C**). Tam/SI-2 and tam/5MPN reduced tumorsphere formation
in cyto PELP1 expressing cells by ~85% (p < 0.0001) and 80% (p < 0.0001) compared to vehicle.
Single agent treatment with tam or SI-2 also reduced tumorspheres, but to a lesser degree than
combinations. PFK158 co-treatment with tam was not more effective than tam alone and was not
further pursued (**Supplementary Figure 14A**). Effective combinations were then tested in J110
cells (**Supplementary Figure 14B-14D**). Tam, SI-2, and 5MPN alone inhibited tumorspheres by
39, 41, and 28%, while co-treatment did not have dramatic effects. The SI-2/5MPN combination
was most effective in J110 cells, and decreased tumorsphere formation by 60%, most likely
because J110 cells are an SRC-3-derived transgenic mouse mammary tumor cell line (25).
358
Because PELP1 confers tamoxifen and Taxol resistance (**Figure 5A**), we also tested the effect
of these agents in resistant cell models. Similar to observations in MCF-7 PELP1 models, tam co-
treatments were more effective when combined with SI-2 or 5MPN in MCF-7 TaxR models
(**Figure 6D, 6E**). The SI-2/5MPN combination was not more effective than individual agents in
TaxR models (**Figure 6F**), suggesting that SRC-3 and PFKFB4 cooperation occurs in tam-
sensitive models. Accordingly, SI-2/5MPN co-treatment in MCF-7 and T47D TamR models
reduced tumorsphere formation by 77% (p < 0.0001) and 75% (p < 0.0001) (**Supplementary
Figure 14E, 14F**).
367
To further explore the therapeutic potential of inhibitor combinations, we utilized pre-clinical
patient-derived organoid models (PDxO;). Synergy screens were used to test combinations
identified from **Figure 6** on proliferation using CellTiter Glo assays in PDxO models (HCI-003, -
011, and -017). Zero Interaction Potency (ZIP) scores are shown in contour maps for tam/SI-2,
tam/5MPN, and SI-2/5MPN treatments (**Figure 7A-7C, Supplementary Figure 15A-15C**), and

indicate the percent a response is higher (>1) or lower (<1) than the expected response for the

dose combination (δ-score). While the δ-score across the range of dose combinations tested were

relatively weak, significant peaks of synergism (δ-score >5) were observed. The most synergistic

area scores are summarized in **Figure 7D**. The tam/SI-2 δ-scores (~1 to 3) were the lowest and

contour maps indicate antagonism. The SI-2/5MPN δ-scores (~5 to 13.5) are lower than the

tam/5MPN scores (~12 to 27) suggesting the tam/5MPN combination is more effective at inhibiting

PDxO proliferation. Next, we evaluated two of the ER+ PDxO models (HCI-003, HCI-017) for

expression of PELP1, SRC-3, PFKFB3, PFKFB4, and ER mRNA and protein (**Supplementary**

Figure 15D, Figure 7E). MCF-7 and T47D cell lines were included as controls. Both HCI-003 and

HCI-017 express all proteins tested. Interestingly, PDxO models have higher levels of PFKFB

proteins compared to MCF-7 and T47D cells. Next, we tested inhibitor combinations on CSC

activity in PDxO models. PDxOs were grown to maturity, pre-treated for 3 days, then dissociated

and seeded into tumorspheres in the presence of inhibitors. Individual treatments (tam, SI-2,

5MPN) reduced tumorsphere formation in both PDxO models by 36 to 62% (**Figure 7F-7H**). The

tam/SI-2 combination was not more effective than individual treatment (**Figure 7F**). In contrast,

tam/5MPN was more effective than tam or 5MPN alone and reduced tumorspheres by ~71% and

~90% in HCI-003 and HCI-017 (**Figure 7G**)**.** SI-2/5MPN co-treatment was more effective than SI-

2 or 5MPN alone and reduced tumorsphere formation by ~71% ($p < 0.0001$) and ~74% ($p <$

0.0001) in HCI-003 and HCI-017 (**Figure 7H**). These results demonstrate that blocking the

PELP1/SRC-3 complex and associated binding partners is an effective approach to targeting CSC

populations in multiple models of advanced breast cancer. Taken together, these studies provide

promising alternative approaches to target non-ER mediators and overcome emergence of

chemotherapy and endocrine resistance.

DISCUSSION

The CSC hypothesis postulates that tumors contain a subset population (i.e. CSCs) that share

properties of normal stem cells including self-renewal, differentiation, and capacity to repopulate

the heterogeneous tumor (9). CSCs are proposed to have heightened resistance to cancer

therapies due to their relative quiescent state (26), enabling this population to evade standard of

care treatments that target proliferating bulk tumor cells. Herein, we sought to define mechanisms

of SR co-activator driven CSC survival and expansion in ER+ breast cancer. We conclude that

SR co-activator complexes enhance CSC activity and therapy resistance by promoting metabolic

plasticity. Inhibiting these complexes and/or associated binding partners in combination with

endocrine therapies may be an effective strategy to block CSC survival and self-renewal, and
breast cancer progression.

Our findings further implicate PELP1/SRC-3 complexes as mediators of CSC activity. We
observed similarities in gene expression, cell metabolism, and sensitivity to inhibitors of PELP1
binding partners in endocrine and chemotherapy resistant ER+ cell lines. Although PELP1
expression contributes to cell survival in response to Taxol in TNBC (14), our studies are the first
to demonstrate enhanced Taxol tolerance in the context of cyto PELP1 in ER+ breast cancer. Our
results in TaxR models highlight the impact of targeting PELP1 binding partners involved in
PELP1-mediated CSC self-renewal (**Figure 5**). Mesenchymal stem cells (27) and ovarian cancer
cells (28) achieve Taxol resistance by shifting to G0 and entering quiescence. PELP1 is a
substrate of CDKs and modulates G1/S cell cycle progression (29). PELP1 may confer Taxol
resistance in part through cell cycle regulation, albeit further studies are needed to define
cytoplasmic PELP1-specific contributions in this context.

Contributing factors to CSC survival include metabolic plasticity, which enables adaptation to
diverse tumor environments. For example, inhibition of glycolysis reduces breast and lung CSCs
(30). Glycolytic reprogramming has been documented in breast cancer cells during EMT, resulting
in acquisition of CSC-like characteristics and tumorigenicity (31). In contrast, breast CSCs utilize
oxidative phosphorylation (OXPHOS) as their primary metabolic program (32). Bulk tumor cells
depend chiefly on glycolysis, whereas tumors enriched for breast CSCs rely mainly on OXPHOS
(33). RNA-seq analysis indicated cytoplasmic PELP1 imparts increased HIF-activated pathways
under normoxic 3D conditions to enrich for CSCs. ChIP assays demonstrated *EPAS1* (i.e. HIF-
2α) recruitment to HRE regions of the PELP1 promoter in TNBC cells (34). Thus, PELP1-induced
HIF pathways may serve as a feed-forward mechanism to drive metabolic genes programs.
PFKFB3 and PFKFB4 are required for glycolytic response to hypoxia via HIF-1α activation (20).
We demonstrated that cyto PELP1 expressing cells increased glycolysis and mitochondrial
respiration. Additional studies are needed to define the bioenergetics driving this plasticity.
PFKFB4-mediated SRC-3 Ser857 phosphorylation has essential functions in lung and breast
cancer metastasis and metabolism (35). Phosphorylation of SRC-3 Ser857 promotes SRC-3
association with transcription factor *ATF4* to mediate non-oxidative pentose phosphate pathway
and purine synthesis**.** This study (35) did not evaluate SRC-3 in the context of CSCs, although
SRC-3 has been linked to CSC activity (10,11). Our IPA studies also identified ATF4 pathway

activation (**Figure 1**); upregulation of *ATF4* could explain the correlation between PFKFB4 and

PELP1/SRC-3-driven CSCs.

PFKFB inhibitors are emerging as promising treatments in endocrine and chemotherapy-resistant

ER+ breast cancer (36). PFKFB3 inhibitor, PFK158, displays broad anti-tumor and

immunomodulatory effects in human and preclinical mouse models (37) and was evaluated in a

Phase I clinical trial with no significant adverse effects (38). The prognostic value of PFKFB4

expression was evaluated in 200 tumor samples from stage I to III breast cancer patients. Similar

to our METABRIC analysis (**Supplementary Figure 10**), elevated PFKFB4 expression was

associated with poor disease-free survival and overall survival in ER+, HER2+, or TNBC patients

(39). PFKFB4 inhibitors (e.g. 5MPN) have not yet moved to clinical trials. Studies have suggested

correlative and mechanistic links between PFKFBs and CSCs. *PFKFB3* was upregulated in a

CD44^{hi}CD24^{lo} gene signature correlated to risk of distant metastasis and poor outcome in breast

cancer patients (40). A cleaved product of CD44 (CD44ICD) promoted breast cancer stemness

via PFKFB4-mediated glycolysis (41). Notably, 5MPN treatment suppressed CD44ICD-induced

tumorigenesis. We have further implicated PFKFBs as drivers of CSC activity by demonstrating

5MPN reduces tumorspheres as a single agent or in combination treatments in multiple ER+

breast cancer models, including treatment resistant cells (TaxR, TamR), murine tumor cells, and

pre-clinical PDxOs. Our data shows that treatment with 5MPN in combination with SI-2 or tam

inhibits PDxO proliferation (**Figure 7**), but importantly also targets the CSC population, Studies in

breast cancer patients indicate that EMT and CSC markers are present in CTC populations, which

have high metastatic potential (42). Our MIND xenografts demonstrate PFKFB4 knockdown does

not have an effect on primary tumor burden, but reduces CTC populations (**Figure 4**). These data

suggest PFKFB4 inhibition is an effective strategy for targeting CSCs and CTCs in ER+ breast

cancer. Future work should involve assessing overlap between PFKFB4-modulated CSC and

CTC populations by evaluating the impact of 5MPN inhibitor combinations *in vivo*.

To evaluate the impact of SR co-activators on CSCs, it will be important that future studies

consider SR-driven contributions. Breast CSCs are reported to be mostly ER negative (43), which

may explain their poor response to anti-estrogens. However, SR+ cells contribute to CSC biology

through SR-dependent (namely PR) paracrine factors (44)**.** For example, breast CSC self-renewal

was stimulated after anti-estrogen treatment of breast cancer cells or PDX models (45,46). These

studies suggest anti-estrogen therapies may initially slow tumor growth, but concurrently evoke

plasticity and CSC activity in non-proliferative tumor cells. Notably, PELP1-containing complexes

include ER and PR (47). PRs but not ER are potent drivers of stem and progenitor cell expansion

in normal and neoplastic breast tissues (48). We have recently defined a requirement for

phosphorylated and inducible PR in CSC biology (49), insulin hypersensitivity, and tam resistance

in ER+ breast cancer (13). CSC outgrowth in therapy resistant ER+/PR-low breast cancer models

is blocked by PR knockdown or antiprogestins (13). These findings suggest PELP1/SRC-3

complexes enable constitutive SR activity in sub-populations that easily bypass endocrine

therapies. Antagonizing estrogen signaling may select for cells that display ligand-independent

ER and PR, resulting in increased proportions of breast CSCs, and subsequently promote

metastasis. Therefore, treatment should include endocrine therapy in combination with targeted

therapies that block mediators of CSC survival and self-renewal as defined herein (i.e. PFKFBs).

CONCLUSION

Our work demonstrates that targeting SR co-activators and associated binding partners involved

in driving CSC survival, self-renewal, and metabolic plasticity may impede breast cancer

progression and has the potential to lead to improved outcomes. Identifying the mechanisms that

mediate recurrent ER+ tumor cell populations (e.g. CSCs, CTCs) will enable specific targeting

within heterogeneous breast tumors to overcome endocrine and chemotherapy resistance.

ACKNOWLEDGEMENTS

We thank Bruce Lindgren for biostatistics support, and the Masonic Cancer Center Biostatistics

and Bioinformatics, Analytical Biochemistry, and Flow Cytometry shared resources. We also

thank Zohar Sachs and Michael Franklin for critical reading of this manuscript.

Figure 1. PELP1-induced gene expression is altered in 3D conditions. (**A**) ALDH activity and (**B**)

CD44^{hi}/CD24^{lo} populations in MCF-7 PELP1 cells. (**C**) Venn diagrams showing unique genes up

or downregulated >2-fold in MCF-7 PELP1 cells (3D vs. 2D). IPA analysis of (**D**) upstream

regulators and (**E**) diseases or functions. (**F**) Representative heat-map showing log2(FPKM)

values of cyto PELP1 gene signature. (**G**) Volcano plots of 3D vs. 2D comparison of MCF-7

PELP1 cells. X-axis is Log2(fold change) and Y-axis represent -Log 10 Benjamini-Hochberg

corrected Q-values. Kaplan-Meier curves for upper and lower 50th percentile of cyto PELP1 gene

signature expression in the METABRIC (**H**) all subtypes and (**I**) ER+ only patient cohorts. Graphed

data represent the mean ± SD (n = 3).   * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure 2. PELP1 cytoplasmic signaling upregulates HIF-activated metabolic pathways. (**A**) mRNA levels of *EPAS1, PFKFB3, PFKFB4, NDRG1,* and *SOX9* in MCF-7 PELP1 cells. (**B**) OCR and ECAR measured in MCF-7 PELP1 cells by Seahorse Cell Energy Phenotype test. (**C**) OCR measured in MCF-7 PELP1 cells by Seahorse Mito Stress test. (**D**) Glucose uptake in cells treated with 2-NBDG (10 μM). 2-NBDG uptake is represented as % cells relative to control. Graphed data represent the mean ± SD (n = 3). * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure 3. PFKFB inhibition blocks PELP1/SRC-3 signaling. Co-immunoprecipitation of (**A**) PELP1

and PFKFB3 or (**B**) PFKFB4 in MCF-7 PELP1 cells. Co-immunoprecipitation of PELP1 and SRC-

3 in MCF-7 PELP1 cells treated with vehicle (DMSO), (**C**) PFK158 (100 nM), or (**D**) 5MPN (5 μM).

Cell lysate controls (*right*). (**E**) Secondary tumorsphere assays in MCF-7 PELP1 shGFP control

or shPFKFB4 knockdown cells. Secondary tumorsphere assays in MCF-7 PELP1 cells treated

with vehicle, (**F**) PFK158 or (**G**) 5MPN. (**H**) Secondary tumorsphere assays in J110 cells treated

with vehicle, PFK158, or 5MPN. Western blot shows PFKFB3 and PFKFB4 protein in J110 cells.

Graphed data represent the mean ± SD (n = 3). * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure 4. PFKFB4 knockdown abrogates cyto PELP1 CTCs in MIND xenograft models. (**A**) Representative H&E stains from MIND glands (WT and cyto PELP1). (**B**) Tumor area (%) calculated from H&E sections from (**A**). (**C**) Tumor area (%) calculated from H&E sections from WT and cyto PELP1 (shGFP, shPFKFB4) MIND glands. (**D**) Representative images of CTCs from blood samples collected from mice injected with WT or cyto PELP1 (shGFP, shPFKFB4) cells. (**E**) Average size of soft agar colonies (CTCs) from (**D**). (**F**) Average number of colonies/well (CTCs). Graphed data represent the mean ± SD (n = 5). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 5. Therapy resistant models phenocopy cyto PELP1 cancer biology. (**A**) Taxol dose
response in MCF-7 PELP1 cells (0-125 nM Taxol). (**B**) mRNA levels of *EPAS1, PFKFB3*, and

PFKFB4 in MCF-7 TaxR (*top*) or TamR (*bottom*) cells cultured in 2D or 3D conditions. (**C**) OCR
and ECAR measured in MCF-7 TaxR cells by Seahorse Cell Energy Phenotype test. (**D**) OCR
measured in MCF-7 TaxR cells by Seahorse Mito Stress test. (**E**) Glucose uptake in cells treated
with 2-NBDG (10 µM). (**F**) Co-immunoprecipitation of PELP1 and SRC-3 in MCF-7 TaxR (*top*) or
TamR (*bottom*) cells. Secondary tumorsphere assays in (**G**) MCF-7 TaxR and (**H**) MCF-7 TamR
cells treated with vehicle (DMSO), PFK158 (100 nM), 5MPN (5 µM), or SI-2 (100 nM). Graphed
data represent the mean ± SD (n = 3). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

641 **Figure 6.** Endocrine therapies exhibit combinatorial effects with PELP1 complex inhibitors.

642 Tumorsphere assays in MCF-7 PELP1 cells treated with: (**A**) tam/SI-2, (**B**) tam/5MPN, or (**C**) SI-

643 2/5MPN. Tumorsphere assays in MCF-7 TaxR cells treated with: (**D**) tam/SI-2, (**E**) tam/5MPN, or

644 (**F**) SI-2/5MPN. Concentrations: tam (100 nM), 5MPN (5 µM), SI-2 (100 nM). Graphed data

645 represent the mean ± SD (n = 3). * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure 7. Co-treatments in preclinical ER+ PDxO models target CSCs. CellTiter Glo assays in HCI-003 and -017 PDxOs co-treated with (**A**) tam/SI-2, (**B**) tam/5MPN, or (**C**) SI-2/5MPN. (**D**) Tables summarizing most synergistic area scores from **7A-C and SFigure 15A-15C**. (**E**) Western blot of PELP1, SRC-3, PFKFB3, PFKFB4, and ER protein levels in HCI-003 and HCI-017. Tumorsphere assays in HCI-003 and HCI-017 PDxOs co-treated with (**F**) tam/SI-2, (**G**) tam/5MPN, or (**H**) SI-2/5MPN. Prior to assay, PDxO models were pre-treated with the indicated compounds for 3 days and subjected to continued treatment during the assay. Concentrations: tam (100 nM), 5MPN (5 µM), SI-2 (100 nM). Graphed data represent the mean ± SD (n = 3). * p < 0.05, ** p < 0.01, *** p < 0.001.

REFERENCES

1.   Vadlamudi RK, Wang RA, Mazumdar A, Kim Y, Shin J, Sahin A, et al. Molecular cloning
and characterization of PELP1, a novel human coregulator of estrogen receptor alpha. J
Biol Chem 2001;276(41):38272-9.
2.   Xu J, Liao L, Ning G, Yoshida-Komiya H, Deng C, O'Malley BW. The steroid receptor
coactivator SRC-3 (p/CIP/RAC3/AIB1/ACTR/TRAM-1) is required for normal growth,
puberty, female reproductive function, and mammary gland development. Proc Natl
Acad Sci U S A 2000;97(12):6379-84.
3.   Habashy HO, Powe DG, Rakha EA, Ball G, Macmillan RD, Green AR, et al. The prognostic
significance of PELP1 expression in invasive breast cancer with emphasis on the ER-
positive luminal-like subtype. Breast Cancer Res Treat 2010;120(3):603-12.
4.   Kumar R, Zhang H, Holm C, Vadlamudi RK, Landberg G, Rayala SK. Extranuclear
coactivator signaling confers insensitivity to tamoxifen. Clin Cancer Res
2009;15(12):4123-30.
5.   Vadlamudi RK, Manavathi B, Balasenthil S, Nair SS, Yang Z, Sahin AA, et al. Functional
implications of altered subcellular localization of PELP1 in breast cancer cells. Cancer Res
2005;65(17):7724-32.
6.   Burandt E, Jens G, Holst F, Janicke F, Muller V, Quaas A, et al. Prognostic relevance of
AIB1 (NCoA3) amplification and overexpression in breast cancer. Breast Cancer Res
Treat 2013;137(3):745-53.
7.   Louie MC, Zou JX, Rabinovich A, Chen HW. ACTR/AIB1 functions as an E2F1 coactivator
to promote breast cancer cell proliferation and antiestrogen resistance. Mol Cell Biol
2004;24(12):5157-71.
8.   Osborne CK, Bardou V, Hopp TA, Chamness GC, Hilsenbeck SG, Fuqua SA, et al. Role of
the estrogen receptor coactivator AIB1 (SRC-3) and HER-2/neu in tamoxifen resistance
in breast cancer. J Natl Cancer Inst 2003;95(5):353-61.
9.   Visvader JE, Lindeman GJ. Cancer stem cells: current status and evolving complexities.
Cell Stem Cell 2012;10(6):717-28.
10.   Rohira AD, Yan F, Wang L, Wang J, Zhou S, Lu A, et al. Targeting SRC Coactivators Blocks
the Tumor-Initiating Capacity of Cancer Stem-like Cells. Cancer Res 2017;77(16):4293-
304.
11.   Truong TH, Hu H, Temiz NA, Hagen KM, Girard BJ, Brady NJ, et al. Cancer Stem Cell
Phenotypes in ER(+) Breast Cancer Models Are Promoted by PELP1/AIB1 Complexes.
Mol Cancer Res 2018;16(4):707-19.
12.   Yang Y, Chan JY, Temiz NA, Yee D. Insulin Receptor Substrate Suppression by the
Tyrphostin NT157 Inhibits Responses to Insulin-Like Growth Factor-I and Insulin in
Breast Cancer Cells. Horm Cancer 2018;9(6):371-82.
13.   Dwyer AR, Truong TH, Perez Kerkvliet C, Paul KV, Kabos P, Sartorius CA, et al. Insulin
Receptor Substrate-1 (IRS-1) Mediates Progesterone Receptor-Driven Stemness and
Endocrine Resistance in Estrogen Receptor+ Breast Cancer Br J Cancer 2020;Manuscript
Accepted.

14.   Regan Anderson TM, Ma S, Perez Kerkvliet C, Peng Y, Helle TM, Krutilina RI, et al. Taxol
Induces Brk-dependent Prosurvival Phenotypes in TNBC Cells through an AhR/GR/HIF-
driven Signaling Axis. Mol Cancer Res 2018;16(11):1761-72.

15.   Yadav B, Wennerberg K, Aittokallio T, Tang J. Searching for Drug Synergy in Complex
Dose-Response Landscapes Using an Interaction Potency Model. Comput Struct
Biotechnol J 2015;13:504-13.

16.   Ianevski A, Giri AK, Aittokallio T. SynergyFinder 2.0: visual analytics of multi-drug
combination synergies. Nucleic Acids Res 2020;48(W1):W488-W93.

17.   Behbod F, Kittrell FS, LaMarca H, Edwards D, Kerbawy S, Heestand JC, et al. An
intraductal human-in-mouse transplantation model mimics the subtypes of ductal
carcinoma in situ. Breast Cancer Res 2009;11(5):R66.

18.   Sflomos G, Dormoy V, Metsalu T, Jeitziner R, Battista L, Scabia V, et al. A Preclinical
Model for ERalpha-Positive Breast Cancer Points to the Epithelial Microenvironment as
Determinant of Luminal Phenotype and Hormone Response. Cancer Cell
2016;29(3):407-22.

19.   Liu Y, Nenutil R, Appleyard MV, Murray K, Boylan M, Thompson AM, et al. Lack of
correlation of stem cell markers in breast cancer stem cells. Br J Cancer
2014;110(8):2063-71.

20.   Chesney J, Clark J, Klarer AC, Imbert-Fernandez Y, Lane AN, Telang S. Fructose-2,6-
bisphosphate synthesis by 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase 4
(PFKFB4) is required for the glycolytic response to hypoxia and tumor growth.
Oncotarget 2014;5(16):6670-86.

21.   Kim JW, Dang CV. Cancer's molecular sweet tooth and the Warburg effect. Cancer Res
2006;66(18):8927-30.

22.   Shi L, Pan H, Liu Z, Xie J, Han W. Roles of PFKFB3 in cancer. Signal Transduct Target Ther
2017;2:17044.

23.   Torres-Arzayus MI, Font de Mora J, Yuan J, Vazquez F, Bronson R, Rue M, et al. High
tumor incidence and activation of the PI3K/AKT pathway in transgenic mice define AIB1
as an oncogene. Cancer Cell 2004;6(3):263-74.

24.   Kilgour E, Rothwell DG, Brady G, Dive C. Liquid Biopsy-Based Biomarkers of Treatment
Response and Resistance. Cancer Cell 2020;37(4):485-95.

25.   Torres-Arzayus MI, Yuan J, DellaGatta JL, Lane H, Kung AL, Brown M. Targeting the AIB1
oncogene through mammalian target of rapamycin inhibition in the mammary gland.
Cancer Res 2006;66(23):11381-8.

26.   Moore N, Lyle S. Quiescent, slow-cycling stem cell populations in cancer: a review of the
evidence and discussion of significance. J Oncol 2011;2011.

27.   Bosco DB, Kenworthy R, Zorio DA, Sang QX. Human mesenchymal stem cells are
resistant to Paclitaxel by adopting a non-proliferative fibroblastic state. PLoS One
2015;10(6):e0128511.

28.   Wang X, Pan L, Mao N, Sun L, Qin X, Yin J. Cell-cycle synchronization reverses Taxol
resistance of human ovarian cancer cell lines. Cancer Cell Int 2013;13(1):77.

29.   Nair BC, Nair SS, Chakravarty D, Challa R, Manavathi B, Yew PR, et al. Cyclin-dependent
kinase-mediated phosphorylation plays a critical role in the oncogenic functions of
PELP1. Cancer Res 2010;70(18):7166-75.

30.   O'Neill S, Porter RK, McNamee N, Martinez VG, O'Driscoll L. 2-Deoxy-D-Glucose inhibits
aggressive triple-negative breast cancer cells by targeting glycolysis and the cancer stem
cell phenotype. Sci Rep 2019;9(1):3788.
31.   Dong C, Yuan T, Wu Y, Wang Y, Fan TW, Miriyala S, et al. Loss of FBP1 by Snail-mediated
repression provides metabolic advantages in basal-like breast cancer. Cancer Cell
2013;23(3):316-31.
32.   Vlashi E, Lagadec C, Vergnes L, Reue K, Frohnen P, Chan M, et al. Metabolic differences
in breast cancer stem cells and differentiated progeny. Breast Cancer Res Treat
2014;146(3):525-34.
33.   Banerjee A, Arvinrad P, Darley M, Laversin SA, Parker R, Rose-Zerilli MJJ, et al. The
effects of restricted glycolysis on stem-cell like characteristics of breast cancer cells.
Oncotarget 2018;9(33):23274-88.
34.   Regan Anderson TM, Ma SH, Raj GV, Cidlowski JA, Helle TM, Knutson TP, et al. Breast
Tumor Kinase (Brk/PTK6) Is Induced by HIF, Glucocorticoid Receptor, and PELP1-
Mediated Stress Signaling in Triple-Negative Breast Cancer. Cancer Res 2016;76(6):1653-
63.
35.   Dasgupta S, Rajapakshe K, Zhu B, Nikolai BC, Yi P, Putluri N, et al. Metabolic enzyme
PFKFB4 activates transcriptional coactivator SRC-3 to drive breast cancer. Nature
2018;556(7700):249-54.
36.   Mondal S, Roy D, Sarkar Bhattacharya S, Jin L, Jung D, Zhang S, et al. Therapeutic
targeting of PFKFB3 with a novel glycolytic inhibitor PFK158 promotes lipophagy and
chemosensitivity in gynecologic cancers. Int J Cancer 2019;144(1):178-89.
37.   Telang S, Yaddanapudi K, Grewal J, Redman R, Fu S, Pohlmann P, et al. Abstract B90:
PFK-158 is a first-in-human inhibitor of PFKFB3 that selectively suppresses glucose
metabolism of cancer cells and inhibits the immunosuppressive Th17 cells and MDSCs in
advanced cancer patients. Cancer Res 2016;76(24 Supplement):B90.
38.   Redman R, Pohlmann P, Kurman M, Tapolsky GH, Chesney J. Abstract CT206: PFK-158,
first-in-man and first-in-class inhibitor of PFKFB3/ glycolysis: A phase I, dose escalation,
multi-center study in patients with advanced solid malignancies. Cancer Res 2015;75(15
Supplement):CT206.
39.   Yao L, Wang L, Cao ZG, Hu X, Shao ZM. High expression of metabolic enzyme PFKFB4 is
associated with poor prognosis of operable breast cancer. Cancer Cell Int 2019;19:165.
40.   Marotta LL, Almendro V, Marusyk A, Shipitsin M, Schemme J, Walker SR, et al. The
JAK2/STAT3 signaling pathway is required for growth of CD44(+)CD24(-) stem cell-like
breast cancer cells in human tumors. J Clin Invest 2011;121(7):2723-35.
41.   Gao R, Li D, Xun J, Zhou W, Li J, Wang J, et al. CD44ICD promotes breast cancer stemness
via PFKFB4-mediated glucose metabolism. Theranostics 2018;8(22):6248-62.
42.   Giordano A, Gao H, Anfossi S, Cohen E, Mego M, Lee BN, et al. Epithelial-mesenchymal
transition and stem cell markers in patients with HER2-positive metastatic breast
cancer. Mol Cancer Ther 2012;11(11):2526-34.
43.   Harrison H, Simoes BM, Rogerson L, Howell SJ, Landberg G, Clarke RB. Oestrogen
increases the activity of oestrogen receptor negative breast cancer stem cells through
paracrine EGFR and Notch signalling. Breast Cancer Res 2013;15(2):R21.

44.   Joshi PA, Jackson HW, Beristain AG, Di Grappa MA, Mote PA, Clarke CL, et al.
808   Progesterone induces adult mammary stem cell expansion. Nature 2010;465(7299):803-
809   7.
810   45.   Simoes BM, O'Brien CS, Eyre R, Silva A, Yu L, Sarmiento-Castro A, et al. Anti-estrogen
811   Resistance in Human Breast Tumors Is Driven by JAG1-NOTCH4-Dependent Cancer Stem
812   Cell Activity. Cell Rep 2015;12(12):1968-77.
813   46.   Shea MP, O'Leary KA, Fakhraldeen SA, Goffin V, Friedl A, Wisinski KB, et al. Antiestrogen
814   Therapy Increases Plasticity and Cancer Stemness of Prolactin-Induced ERalpha(+)
815   Mammary Carcinomas. Cancer Res 2018;78(7):1672-84.
816   47.   Daniel AR, Gaviglio AL, Knutson TP, Ostrander JH, D'Assoro AB, Ravindranathan P, et al.
817   Progesterone receptor-B enhances estrogen responsiveness of breast cancer cells via
818   scaffolding PELP1- and estrogen receptor-containing transcription complexes. Oncogene
819   2015;34(4):506-15.
820   48.   Rajaram RD, Buric D, Caikovski M, Ayyanan A, Rougemont J, Shan J, et al. Progesterone
821   and Wnt4 control mammary stem cells via myoepithelial crosstalk. EMBO J
822   2015;34(5):641-52.
823   49.   Truong TH, Dwyer AR, Diep CH, Hu H, Hagen KM, Lange CA. Phosphorylated
824   Progesterone Receptor Isoforms Mediate Opposing Stem Cell and Proliferative Breast
825   Cancer Cell Fates. Endocrinology 2019;160(2):430-46.
826