Article Type: Original Article

Title: **Egg size variation in the context of polyandry: a case study using long-**
term field data from snowy plovers

Short Running Title: Egg size variation in a polyandrous shorebird

Keywords: breeding phenology, *Charadrius nivosus*, female-female competition, mating
system, reproductive investment, season, senescence

Abstract:

Gamete size variation *between* the sexes is central to the concept of sex roles, however, to
what extent gamete size variation *within* the sexes relates to sex role variation remains
unclear. Comparative and theoretical studies suggest that, when clutch size is invariable,
polyandry is linked to a reduction of egg size, while increased female-female competition for
mates favors early breeding when females cannot monopolize multiple males. To understand
whether and how breeding phenology, egg size, and mating behavior are related at the
individual level, we studied the reproductive histories of 424 snowy plover females observed
in the wild over a 15-year period. Egg size, but not polyandry, were highly repeatable for
individual females. Consistent with theoretical predictions, we found that polyandrous
females were the earliest breeders and that early clutches contained smaller eggs than
clutches initiated later. Neither egg size nor mating behavior showed clear signs of an age-
related deterioration, on the contrary, prior experience acquired either through age or local
recruitment enabled females to nest early. Taken together, these results suggest that gamete
size variation is not linked to mating behavior at the individual level, and, consequently, the
adaptive potential of such variation appears to be limited.

Teaser Text:

Comparative avian analyses have linked female polyandry and sex-role reversal to the
production of smaller eggs: the notion being that smaller eggs allow polyandrous females to
lay multiple sequential clutches quickly. Our research, however, has found that the notion is
more complex at the individual level. We found that females who started breeding early in
the season were more likely to be polyandrous and produce smaller eggs. Experience was
also found to give older and locally-raised females a competitive advantage over younger and
inexperienced females vying for early breeding opportunities. Despite this, we found limited

evidence that egg size reduction gives a competitive advantage in the scramble for mates.

Instead, seasonal variations in egg size can likely be attributed to other factors, such as the

survival of early-hatching chicks, resource limitations during egg production, and individual

differences. We found that senescence had little impact on egg size or mating behavior.

Future research should also examine how seasonality, age, and mating behavior impact the

reproductive fitness of males, who provide more parental care than females in many sex role

reversed systems. By studying these effects in concert, we can gain a deeper understanding of

sex roles in natural populations.

INTRODUCTION

Anisogamy is a fundamental prerequisite for the concept of sex roles (e.g., Darwin 1871;

Bateman 1948; Wade and Shuster 2002; Schärer et al. 2012; Lehtonen et al. 2016). In species

with conventional sex roles, males, the producers of many small gametes, invest into

acquiring matings to sire as many offspring as possible, whereas females, the producers of

few large gametes, aim to improve the survival prospects of their offspring by obtaining

critical resources and providing parental care. However, there are many species where sex

roles do not conform to this conventional paradigm (reviewed in Janicke et al., 2016).

Moreover, there can also be considerable gamete size variation within the sexes – however,

the causes and consequences of this intrasexual variation remains unclear (Bernardo, 1996;

Immler et al., 2011). For example, the largest eggs produced by some females are typically at

least 50% larger than the smallest eggs produced in the same population (Christians, 2002)

and occasionally the largest egg can be more than twice as heavy as the smallest viable egg

(Svagelj & Quintana, 2011). Such among-individual variation in egg size may have important

consequences for individual fitness and could be linked to different reproductive strategies of

females. Egg size is an important life history trait under maternal control that is positively

associated with offspring fitness (Smith & Fretwell, 1974; Williams, 1994; Christians, 2002;

Krist, 2011). Egg size typically reflects hatchling quality: the amount of nutrients provided to

the developing offspring are especially important during early life (Williams, 1994; Krist,

2011). Because resources for maternal investment during oogenesis are typically limited,

among- and within-female variation in egg size may signal a trade-off between offspring

quality and quantity.

In oviparous organisms, egg size represents a basic measure of maternal investment (Kaplan,

1980; Fox, 1994; Williams, 1994, 2012; Starck & Ricklefs, 1998; Moran & Emlet, 2001; Xu

et al., 2019) and is shown to be related to inter- and intra-specific variation in several life

history traits. For example, egg size is associated with developmental mode, with precocial

species typically producing larger eggs than altricial species (Smith & Fretwell, 1974;

Williams, 1994; Christians, 2002; Krist, 2011). Within species a substantial proportion of the

egg size variation can be explained by consistent individual differences between females

which are highly repeatable across breeding attempts (Christians, 2002). In some cases,

intraspecific variation in egg size is related to among-female variation in reproductive

strategies (Slagsvold *et al.*, 1984) and is occasionally linked to discrete demographic classes

such as reproductive morphs (Beck *et al.*, 2022; Giraldo-Deck *et al.*, 2022).

Given the strong association between egg size variation and life-history traits or maternal

investment it is not surprising that there are strong temporal components related to egg size

variation. For example, reproductive output is often age-dependent (Bouwhuis et al., 2009;

Dingemanse et al., 2020; Hammers et al., 2012; Jankowiak et al., 2018; Lemaître et al., 2015;

Reznick et al., 2004; Riecke et al., 2023; Salguero-Gómez et al., 2016; Zhang et al., 2015)

whereby an individual's performance increases over early life to a maximum, followed by

senescent decline. Senescence is a consequence of age-dependent trade-offs between energy

investments in reproduction at the expense of somatic repair (Kirkwood & Rose, 1991).

Decreasing egg and/or clutch size can indicate reproductive senescence that reduces female

reproductive performance with older ages (Reid *et al.*, 2003; Beamonte-Barrientos *et al.*,

2010; Dingemanse *et al.*, 2020; Fay *et al.*, 2021). Conversely, polyandry has been associated

to female age, with older more experienced females being more likely to be polyandrous than

younger females (Oring *et al.*, 1994). Such observations of higher mating rates at older ages

suggest that differences in individual quality or experience are related to female mating

behavior (Whittingham & Dunn, 2010), which could in turn be linked to among- and within-

female variation in egg size.

Another source for temporal variation in egg size is seasonality. Egg size often changes

throughout the reproductive season, especially in birds (Dittmann & Hötker, 2001; Skrade &

Dinsmore, 2013; Weiser *et al.*, 2018; Kubelka *et al.*, 2020; Verhoeven *et al.*, 2020). At the

population level, reproductive performance often declines with season which may coincide

with declining egg sizes (Varpe, 2017; Weiser *et al.*, 2018). Several processes may contribute

to this trend. First, in environments with limited or seasonally dependent resources, breeding

early may provide favorable conditions for raising offspring, thereby reducing the amount of

parental care required and maximizing offspring survival prospects (Varpe, 2017; Weiser *et*
al., 2018). Consistent with this, offspring that are born early in the season often have higher
survival prospects than offspring born late in the season (Festa-Bianchet, 1988; Reznick *et*
al., 2006; Plard *et al.*, 2015; Cruz-López *et al.*, 2017; Varpe, 2017). Second, only high quality
and experienced females may be able to reproduce early in the season because of potential
increased physiological stress for early egg-laying females, and/or the increased likelihood of
brood failure in environments with high stochasticity, such as low food availability,
inclement weather, or frequency-dependent predation risk (Borgmann *et al.*, 2013; Ockendon
et al., 2013). Therefore, less experienced or low-quality females may need to postpone their
reproduction to a later date when conditions are more benign. Third, alternation between
extended breeding and non-breeding periods may contribute to differences in egg and,
consequently, offspring quality: when breeding is restricted to a certain period of the year,
females can use the preceding non-reproductive period to amass resources for investment into
egg or clutch size of their initial breeding attempt (Yohannes *et al.*, 2010). As the breeding
season advances, less time is available for successful breeding attempts and as a consequence
females may acquire less resources for egg production resulting in fewer and or/smaller eggs
in late clutches (Wallander & Andersson, 2003).

An understudied evolutionary process explaining egg size variation within species is sexual
selection acting on females. Although sexual selection is predominantly considered to shape
male-male competition, it is increasingly acknowledged that sexual selection also operates on
females with strong female-female competition for mates (Clutton-Brock, 2007; Kraaijeveld
et al., 2007; Fromonteil *et al.*, 2023). In some species, competition over mating opportunities
is stronger in females than in males, leading to a reversal of conventional sex roles (Oring &
Lank, 1982; Rosenqvist, 1990; Andersson, 2004; Goymann *et al.*, 2015; Fritzsche *et al.*,
2021). Theoretical models have linked intrasexual competition to variation in gamete size,
with more intense competition leading to smaller gametes (Wade & Shuster, 2002; Lehtonen
et al., 2016). Notably, this relationship is not restricted to males and can also be applied to
females in polyandrous species with female-female competition over mates, resulting in a
reduction of egg size. For instance, at the species level, a comparative analysis of birds
showed that a lineage's egg size tends to decrease with the evolution of polyandry (Oring &
Lank, 1982; Rosenqvist, 1990; Andersson, 2004; Goymann *et al.*, 2015; Fritzsche *et al.*,
2021) – supposedly due to the selective advantages that producing smaller eggs has on
minimizing the laying period between matings (Goymann *et al.*, 2015).

Andersson (2004) presented a theoretical quantitative genetics model in which intense female
scramble competition (i.e., whereby females compete to find sexually receptive males to
commence breeding) shapes breeding phenology and egg size variation in females.
According to the model, early breeding provides substantial advantages as it allows these
females enough time to pursue sequential breeding attempts or to replace a failed first attempt
(i.e., "re-nesting"; Morrison *et al.*, 2019). Early breeding should therefore be tightly
correlated with female polyandry and, ultimately enhance the fitness of polyandrous females.
Laying smaller eggs could provide another competitive edge in female-female competition
over males: when clutch sizes are uniform, females that produce small eggs can complete
their clutch faster than those that produce larger eggs and hence gain a competitive edge in
the competition for mates especially when breeding synchrony is high and additional males
are limited. A lower boundary for egg size is provided by reduced offspring viability because
smaller eggs have fewer nutrients (Williams, 1994; Blomqvist *et al.*, 1997; Krist, 2011;
Giraldo-Deck *et al.*, 2022). However, the predictions on breeding phenology, egg size, and
their relationship with female mating behavior provided in Andersson's (2004) theoretical
model remain to be tested empirically at the individual level.

Among oviparous animals, shorebirds (part of the order Charadriiformes) produce some of
the largest eggs in relation to body mass due to the needs of their precocial and
(predominantly) nidifugous young (Lack, 1968; Rahn *et al.*, 1975). Clutch size is highly
uniform within shorebird species and rarely exceeds four eggs (Rahn *et al.*, 1975) meaning
that egg size represents an adequate proxy for maternal investment. As a clade, shorebirds
also exhibit a disproportionately high prevalence of polyandry (Andersson, 2005; Colwell,
2010; Oring, 1986). Here, we investigate breeding phenology, egg size variation, and their
relation with female mating strategy in snowy plovers (*Charadrius nivosus*) using a 15-year
longitudinal mark-recapture dataset. The snowy plover is a long-lived shorebird (longevity
record: 20 years; Colwell et al., 2017; M. A. Colwell *personal communication,* 15 March,
2021) exhibiting flexible sex-specific parental roles with sequential social monogamy and
polygamy occurring in the same population (Warriner *et al.*, 1986; Eberhart-Phillips *et al.*,
2017). In parts of the species' range, the breeding season can extend up to six months
allowing for multiple nesting attempts following successful fledging of young or replacing
failed nests due to depredation or flooding (Eberhart-Phillips, 2019; Plaschke *et al.*, 2019).
On average, females have higher polygamy potential and more partners per season than

males, who in turn, provide more parental care than females, specifically for boods
(Carmona-Isunza *et al.*, 2017; Cruz-López *et al.*, 2017; Eberhart-Phillips *et al.*, 2017). The
reason for the higher polygamy potential in females is a surplus of males in the adult
breeding population (Stenzel *et al.*, 2011; Carmona-Isunza *et al.*, 2017; Eberhart-Phillips *et*
al., 2017). Consequently, sequential polyandry, whereby females regularly desert their broods
after hatching to start a new breeding attempt with another male, is commonly observed in
snowy plovers (Warriner *et al.*, 1986). Female desertion for re-mating declines with season in
polyandrous plovers (Amat *et al.*, 1999; Székely *et al.*, 1999; Cruz-López *et al.*, 2017; Kupán
et al., 2021) suggesting that female-female competition for mates should be most pronounced
early in the season.

To disentangle the relationships between gamete size, breeding phenology, and female
mating behavior, we examined the reproductive histories of 426 snowy plover females. First,
we investigated the correlates of re-nesting and female polyandry. We hypothesized that early
nesting females would have higher probabilities of re-nesting or becoming sequentially
polyandrous: early nesting females can rejoin the mating pool faster and therefore have
higher potential to secure an additional breeding attempt compared to later nesting females.
Secondly, we explored the predictors of clutch initiation date for a given female in a season.
We hypothesized that early nesting females would have a competitive advantage over late
nesters – either due to a direct body size advantage or an indirect advantage gained through
prior experience as a previous breeder or local natal recruit. To evaluate this, we explored
how body size, age, and origin (i.e., immigrant or natal recruit) were related to female nesting
phenology and the likelihood of re-nesting or polyandry. Third, we examined the pattern of
egg size variation in relation to season, age, and female body size. We predicted that early-
laying females would produce the smallest eggs compared to later laying females due to the
costs associated with intense female-female competition at the start of the season, i.e., the
'race' to breed early and maximize re-nesting or polyandry potential (Andersson, 2004). If
reproductive senescence affects egg size variation, we predicted egg size to follow a
quadratic relationship with female age. Alternatively, if female polyandry increases with age,
we predicted a negative linear relationship of egg size with female age.

MATERIALS AND METHODS
Data collection

We studied the reproductive effort and breeding schedules of snowy plovers at Bahía de

Ceuta – an important breeding site located on the coast of Sinaloa, western Mexico (23°54'N,

106°57'W). Details on the study site and population are provided elsewhere (e.g., Cruz-

López et al., 2017; Eberhart-Phillips et al., 2020). In brief, we annually monitored breeding

birds from mid-April until early July and collected mark-recapture data following the

methods described in Székely et al. (2011). We searched for nests using telescopes and

mobile hides to minimize disturbance. Upon finding a nest, we measured the length and

width of each egg to the nearest tenth of a mm to determine their size (Figs. 1a, b). Using

these egg dimensions, we calculated egg volume (Fig. 1c) following (Hoyt, 1979) as:

Eq. 1 $$egg\ volume = K \times length \times width^2,$$

where K is 0.486, a volume-index constant for snowy plovers determined by (Székely *et al.*,

1994) through the use of an egg volumeter (Hanson, 1954). The modal clutch size of snowy

plovers is three (91% of 836 nests in our sample) and is the maximum number of eggs we

have observed in this population (Eberhart-Phillips *et al.*, 2020a). We regularly checked

incomplete nests until the clutch was complete and assigned the age of these nests according

to the date of the last egg laid (Plaschke *et al.*, 2019). If the clutch was complete upon

discovery and had not been incubated longer than 10 days (92.5%; $N = 773$ nests), we

determined its lay date by floating the egg and estimating the stage of embryonic

development. For successful clutches that had been incubated for more than 10 days (5.5%, N

= 46 nests), we back-calculated the laying date based on the hatching date assuming an

incubation period of 25 days (Plaschke *et al.*, 2019). In the rare case that the nest did not

hatch and we discovered it after day 10 of incubation (2%, $N = 17$ nests), we assumed that the

nest was 17 days old upon discovery (i.e., the midpoint between the minimum age of 11 days

and the 25-day incubation period).

We identified previously marked nesting adults based on their unique colour ring

combination. We captured unmarked adults on their nests during incubation using funnel

traps (Hall & Cavitt, 2012) and assigned a unique colour ring combination for subsequent

recognition. In the rare circumstance when we were unable to identify parents before

hatching, we attempted to capture parents while they tended the chicks. As snowy plovers

only show a small degree of sexual dimorphism, we determined the sex of all captured

plovers in the field through a combination of plumage characteristics (Argüelles-Tico *et al.*,

2015), time of capture (i.e. females typically incubate during the day and males during the

night; Vincze *et al.*, 2017), and other behavioral cues (e.g., sex-specific brood care; Kupán et

al., 2021). For a subset of adults (57.5%), we confirmed sex molecularly from DNA extracted

from blood samples by PCR amplification of Z and W specific DNA regions with two sex-

typing markers: P2/P8 and Calex-31 (Griffiths *et al.*, 1998; Küpper *et al.*, 2009b; dos

Remedios *et al.*, 2010). Over our 15-year study we conducted 1168 captures of individuals

that had been molecularly sex-typed: in only 2.4% of these captures did we mis-identify the

sex of an individual using field-based cues.

We visited known active nests every four or five days to determine the status of the nest (e.g.,

active, depredated, etc.) until the 20th day after egg laying and thereafter daily until the eggs

hatched or failed. We weighed the chicks shortly after hatching (879 [84.7%] within 24 hours

after hatching, 159 [15.3%] during the second day after hatching) and marked them with an

alphanumeric metal and a single color ring for subsequent identification in the chance that

these individuals would be recruited into the breeding population as adults in future years.

For the years 2006 to 2016 all longitudinal data collected has been compiled as part of the

CeutaOPEN project – an open-access database for individual-based field studies in

evolutionary ecology and conservation biology (Eberhart-Phillips *et al.*, 2020a). We accessed

these data directly from the open source repository (Eberhart-Phillips *et al.*, 2020b) and

supplemented them with data from four additional field seasons: 2017–2020. Please refer to

our RMarkdown vignette that connects to *CeutaOPEN* and reproduces all analytical methods

and results presented below (Supplementary File 1). All of our field activities were

performed and permitted in accordance with the approved ethical guidelines outlined by

SEMARNAT (Secretariat of Environment and Natural Resources, Mexico).

Statistical Analyses

To evaluate the relationship between female breeding behavior, nesting phenology, and egg

size, our investigation implemented three step-wise analyses that: 1) established the factors

related to a female's potential to have multiple breeding attempts either through polyandry or

re-nesting, 2) identified the key female traits linked to lay date, and 3) dissected the elements

explaining within- and among-female variation in egg volume investment. A central trait of

interest in these three analyses was female age, as we needed to acknowledge age-dependent

variation in the repeated measures of individuals over the 15-year period of data collection.

We first present our statistical methods for estimating the age of females that entered the

population as adults, then we describe our models for investigating the interplay between
female breeding behavior, nesting phenology, and egg size.

Statistical estimation of age for females first encountered as adults—To estimate the ages of
unknown individuals in our marked population, we conducted a capture-mark-recapture
analysis using the 'Bayesian Survival Trajectory Analysis' (BaSTA) package in R (v1.9.4,
Colchero et al., 2012), which uses a Bayesian hierarchical framework to fit parametric
survival functions of the marked population while accounting for imperfect detection (see
Supplementary Text A for details). In short, the capture histories of unknown-age individuals
(90%, $N = 405$) are cast against the survival function of the known-age population (10%, $N =$
45) to produce posterior age estimates. To acknowledge uncertainty in the BaSTA age
estimate of an unknown age individual: we ran bootstrap simulations for each of the
following models 1000 times, with every iteration randomly drawing a birth year estimate for
unknown aged individuals from their posterior distributions provided by BaSTA. For each
simulation, we evaluated the influence of birth year uncertainty by examining the effect size
distribution of the 1000 bootstraps in relation to the 95% confidence interval for effect sizes
of the original model that used the median birth year estimate from BaSTA. Furthermore, to
test the accuracy of the BaSTA age estimates, we implemented a cross-validation simulation
which iteratively estimated the age of each known-age female under an *in silicio* scenario
whereby their age was unknown. The simulation confirmed that the BaSTA age-estimates
were adequately accurate, with 51% of estimated ages being correct and a combined 82% of
estimated ages being within 1-year of the true age (Fig. S1) – a notable result considering that
our mark-recapture sample is from a wild and highly vagile avian species. Moreover, BaSTA
age-estimates that deviated from reality were slightly biased towards an under-estimation
(i.e., estimated to be younger; Fig. S1), hence making our age-specific models more
conservative. Lastly, we considered a time-since marking capture-mark-recapture model and
determined that the recapture probability of our dataset was not biased by individual
heterogeneity in post-marking processes (see Supplementary Text B for details).

A key methodological issue for studying age related processes in wild populations is that
stochastic extrinsic mortality reduces the frequency of individuals in older age classes, hence
making it challenging to disentangle between- vs. within-individual age-dependent variation
– a phenomenon known as "selective disappearance" (van de Pol & Verhulst, 2006a; Nussey
et al., 2008). Investigations using longitudinal data to test for age-related processes are

particularly suitable for this task, as they can control for the confounding effects of selective

disappearance through repeated measures of individuals as they age (van de Pol & Verhulst,

2006a; Nussey *et al.*, 2008; Dingemanse *et al.*, 2020). We controlled for selective appearance

and disappearance of females differing in the trait of interest by fitting 'first observed age'

and 'last observed age' as fixed effects – a method that estimates between-individual age

effects introduced by selective disappearance and appearance (Dingemanse et al., 2020; van

de Pol & Verhulst, 2006a, 2006b). We also modelled within-individual age effects by fitting

linear and quadratic forms of a within-group deviation score for age (henceforth 'age-

deviance'), calculated for individual *i* at age *j* as: age_{ij} – [*first observed age*]$_i$ (van de Pol &

Verhulst, 2006a; Snijders & Bosker, 2011) – essentially describing the number of years since

the bird's arrival to the population. To ensure that intercepts of our age-dependent models

represented the reproductive performance for the earliest age at reproduction (i.e., age 1 in

snowy plovers; Page et al., 2009), we fitted age as '*age* – 1' – otherwise it would represent

reproduction at age 0, which is an empirically meaningless estimate. Moreover, we followed

common statistical approaches to investigate age-related processes in birds (e.g., Bouwhuis et

al., 2009, 2010; Dingemanse et al., 2020; Graham et al., 2019; Herborn et al., 2016;

Schroeder et al., 2012) by fitting a quadratic function of age to model age-specific trends.

Lastly, in all models, tarsus length was included as a fixed effect proxy of female structural

size, and was averaged over an individual's lifetime of repeated measurements, i.e., our *a*

priori expectation was that tarsus length is static throughout adult life and that any variation

in this trait was due to measurement error.

Modelling seasonal variation in polyandry potential ("Polyandry model")—Our sample for

studying seasonal polyandry dynamics included 424 females for which the identity of their

mates had been verified through observation. We defined observed polyandry as a binomial

variable that scored an individual as being monogamous or polyandrous each year based on

our observations of them having one or multiple breeding partners, respectively (see Fig. S2

for an example of the sampling distribution). Overall, we observed 90 cases of polyandry

from 74 females over the 15-year period (annual average incidence of observed polyandry:

13.5%, range: 0–25.6%). To assess the relationship between the likelihood of polyandry and

lay date, age, female structural size, and maternal investment, we fitted a binomial linear

mixed effects model that tested the likelihood of polyandry predicted by the fixed effects of

first lay date, age-deviance (see above), first observed age, tarsus length, and the average

egg-volume (of the first clutch of the season). We included individual ID and year as random

effects.

Modelling seasonal variation in re-nesting potential ("Re-nesting model")—Our sample for

studying seasonal re-nesting dynamics included 177 females for which the fate of their first

nest of the season had been verified as a failure. We defined re-nesting as a binomial variable

that scored an individual as being a re-nester or a single-nester each year based on our

observations of them attempting to re-nest after the loss of their first clutch or not,

respectively. Overall, we observed 62 cases of re-nesting from 53 females over the 15-year

period following a failed attempt (annual average incidence of observed re-nesting: 28.6%,

range: 0–58.3%). To evaluate re-nesting dynamics, we fitted a binomial linear mixed effects

model that tested the likelihood of re-nesting predicted by the fixed effects of first lay date,

age-deviance, first observed age, and tarsus length. We included individual ID and year as

random effects.

*Modelling individual variation in lay dat*e *("Lay date model")*— We modeled the effect of

age on a female's first lay date using a univariate mixed-effect structure that included age-

deviance, age-deviance-squared, first observed age, and last observed age as fixed covariates,

and individual and year as random intercepts. Furthermore, we included average tarsus length

as a fixed covariate in addition to recruitment status as a two-level fixed-effect that described

whether a breeding female hatched locally ("local recruit") or was first encountered as an

adult of unknown origin ("immigrant"). Our sample for studying lay date dynamics used the

same nest-level sample as the polyandry model above, however, as we were interested in how

the recruitment status of an individual influenced breeding phenology, we excluded data from

2006 as this was the first year when we began individual marking of all birds in the

population. This subset resulted in 565 nests from 375 females. We visualized the distribution

of lay dates to confirm normality and to assess the population-level variance in breeding

schedule – an indication of inter-female breeding asynchrony and the intensity of female-

female competition for mates (Andersson, 2004).

Modelling individual variation in egg volume ("Egg volume model")—Our sample for

studying egg volume dynamics included 2393 eggs from 836 nests belonging to 424 females.

56 (13.2%) females had three or more years of repeated measures (Fig. S2), 83 (19.6%) had

two years of repeated measures, and 285 (67.2%) were measured in a single year.

Furthermore, 43 (10.1%) individuals in our sample were marked as hatchlings but later
recruited as breeding adults in subsequent years ("known age"), with the remaining 381
(89.9%) individuals being initially marked as adults ("unknown age"). We modelled within-
individual age effects on egg volume by fitting a univariate mixed-effect model that included
linear and quadratic forms of local tenure, and assessed between-individual age effects by
additionally incorporating first observed age and last observed age as fixed covariates.
Female tarsus length was also included as a covariate to control for female body size, and we
incorporated a quadratic function of lay date to assess seasonal variation in egg volume as
several shorebird studies report seasonal increases (Skrade & Dinsmore, 2013; Kwon *et al.*,
2018) or decreases in egg volume (Dittmann & Hötker, 2001; Skrade & Dinsmore, 2013;
Kwon *et al.*, 2018; Kubelka *et al.*, 2020; Verhoeven *et al.*, 2020). We also included clutch
size in our model to control for this alternate source of variation in maternal investment.
Because clutch size variation in snowy plovers is constrained to 1, 2, or 3 eggs and 1- or 2-
egg nests are rare in our sample (collectively 9% of nests, $N = 75$), we modelled clutch size
as a binary trait (i.e., "3-eggs" vs. "1- or 2-eggs"). To disentangle within- from between-
individual effects of lay date on egg size, we used the same logic as with age above: first lay
dates of all individuals each year represented the between-individual seasonal effect, whereas
the deviation in first lay dates of an individual relative across all successive breeding attempts
for a given season represented the within-individual seasonal effect. We included random
intercepts for nest, individual ID, and year, and assumed a Gaussian error distribution for egg
volume.

Evaluating effect sizes and uncertainty—We used packages "lme4" (Bates *et al.*, 2015),
"rptR" (Stoffel *et al.*, 2017) and "partR2" (Stoffel *et al.*, 2020) in R version 4.3.1 (R Core
Team, 2023) to conduct our statistical modelling and assessed homoscedasticity by visually
examining residuals (see Fig. S2). For each of the four mixed-effect models described above,
we evaluated the uncertainty in our parameter estimates by simulating 1000 parametric
bootstraps via the "partR2::partR2" function (Stoffel *et al.*, 2020). Likewise, we derived nest-
, individual-, and year-level repeatabilities (i.e., intra-class correlations) by simulating 1000
parametric bootstraps of the four mixed-effect models using "rptR::rpt". We report fixed
effects as standardized regression coefficients (i.e., beta weights) and repeatability as the
'adjusted repeatability' – interpreted as the repeatability of a given hierarchical group after
controlling for fixed effects (Nakagawa & Schielzeth, 2010).

RESULTS

Sample summary

The modal clutch size of the 836 clutches was 3 eggs (761 nests, 91%; 2-eggs: 72 nests,

8.6%, 1-egg: 3 nests, 0.4%). Average egg length was 3.09 cm (0.11 cm SD, range: 2.09–3.49

cm, Fig. 1a) and width was 2.24 cm (0.05 cm SD, range: 2.02–2.40 cm, Fig. 1b), which

translated into an average egg volume of 7.58 cm^3 (0.48 cm^3 SD, range: 5.10–8.99 cm^3). The

average egg volume of a clutch strongly predicted the average hatch weight of the subsequent

brood (Figs. 1c, Table S1, see Supplementary Text C for methods). Based on BaSTA's

estimated birth year, 184 of the 381 unknown-age females in our sample were first observed

nesting at age one (48.3%), 120 at age two (31.5%), 71 at age three (18.6%), five at age 4

(1.3%), and one at age 5 (0.3%). Of the 43 locally hatched females in our sample, 29 first

nested at age one (67.4%), six were first observed nesting at age two (14.0%), two at age 3

(4.7%), three at age 4 (7.0%), three at ages 5, 7, and 8, respectively (7.0%). Female tarsus

size was not related to their annual apparent survival (see Supplementary Text D for details).

The average local tenure of all females in the sample was 1.57 years (2.15 SD) with an

average age span of 3.12 years (2.03 SD, median: 3, range: 1–14 years) and an average of

1.56 years of observed ages per female (1.04 SD, median: 1, range: 1–8 age-specific

observations). Females in our sample were typically observed nesting every consecutive year

since their first observation, however, some individuals skipped years or breed elsewhere

(Fig. S2; average yearly interval between nesting attempts = 1.07, 0.27 SD). On average,

females made 1.43 (0.56 SD) nesting attempts per season (median = 1, range 1 to 3). The

number of hatched chicks produced by females in years when they were polyandrous, was

2.19 (1.83–2.55 95% CI) chicks more than in years when they were monogamous (Fig. S5; *N*

= 14 years, 424 females, 836 nests).

Seasonal variation in polyandry and re-nesting potential

A female's probability of being polyandrous in a given season was strongly dependent on the

date she laid her first nest, with early breeders being more likely to be polyandrous than late

breeders (Figs. 2a, Table 1). Likewise, a female's likelihood of re-nesting following a failed

attempt decreased with advancing first lay date (Table 2). We found no effect of female

tarsus length on the likelihood of being polyandrous (Table 1) or re-nesting (Table 2). We

found a potential effect of age at first breeding on the likelihood of re-nesting, with females

that initiated local breeding at an older age being the most likely to re-nest (Table 2). This

effect was not found in the polyandry model (Table 1). We found no evidence that polyandry

or re-nesting potential was related to age since first breeding (i.e., age-deviance; Table 1).

The lay date distribution of polyandrous females was bimodal, with peaks in the first and

second nests occurring 11.1 days before and 29.4 days after the unimodal seasonal peak for

monogamous females (Fig. 2b). Similarly, the lay date distribution of re-nesting females was

also bimodal, with peaks in the first and replacement clutches occurring 9.8 days before and

28.6 days after the unimodal seasonal peak for single nesters. There was no association

between egg size and the likelihood of being polyandrous (Table 1). Females had low

repeatability among years in polyandry (Table 1) and re-nesting (Table 2). When pooled

together to assess multi-clutching behavior (i.e., the probability of simply laying two nests or

one regardless of the outcome of the first nest or the identity of the second male), females had

higher repeatability than when polyandry or re-nesting were assessed separately (Table S3).

Individual variation in lay date

Females showed moderate repeatability in their first lay date among years (Table 3). We

found strong support for the effect of origin on first lay date: females that locally hatched and

later recruited into the breeding population initiated nests 7.80 days earlier (95% CI: [5.09,

10.50]) on average compared to conspecifics whose origin was unknown (Table 3 and Fig.

3b). The next strongest fixed effect was within-individual age: young females laid nests later

compared to older females with lay date advancing by ~2.17 days per year until age six (95%

CI: [1.41, 2.93]; Fig. 3a), after which the uncertainty in the trend became unwieldly in the

oldest age classes of our sample (Fig. 3a). Notably, female size did not affect lay date (Table

3 and S3a), nor did between-individual effects of first or last age at breeding (Table 3).

Individual variation in egg volume

Overall, our model accounted for 74.4% of variation in egg volume, with fixed effects

explaining 6.6% of this variation (Table 4). Females were highly repeatable in their egg

volumes between clutches ($r = 0.47$; Table 4). Furthermore, eggs within the same clutch were

moderately repeatable in volume ($r = 0.16$; Table 4). The strongest fixed effect explaining

egg volume variation was the structural size of the mother (Fig. S3b, Table 4): larger females

laid larger eggs than smaller females (model predicted difference in egg size between largest

and smallest females in the sample: 0.57 cm^3 [0.33, 0.80] 95% CI). The second strongest

effect was the between-individual quadratic season function (Fig. 2c): eggs were smallest at

the start of the season (model prediction: 7.18 cm^3 [6.98, 7.38] 95% CI) and then increased,

to be largest shortly after the middle of the season (model prediction: 7.68 cm^3 [7.60, 7.75]

95% CI). Average egg volume also increased between sequential clutches within individuals

but with smaller magnitude than the population-level trend (predicted increase of 0.20 cm^3

[0.03, 0.37] 95%CI). Senescence in egg volume was not supported (Table 4). Furthermore,

we found no support for selective (dis)appearance of individuals according to egg volume

(Table 4).

To further acknowledge uncertainty in the age-estimates provided by BaSTA, we re-analyzed

the egg volume model using only known-aged females. Notably, this strengthened the effect

sizes of 'mother tarsus length' and the within-individual age effects (i.e., 'linear age' and

'quadratic age'). For example, the 'mother tarsus length' effect (β_{tarsus}) increased to 0.32

[0.11, 0.50 95% CI], the 'linear age' effect (β_{age}) increased to -0.05 [-0.21, 0.12], and the

'quadratic age' effect (β_{age}^{2}) increased to -0.10 [-0.24, 0.05] (Table S2). However, given that

the sample size of the known-aged female data set was greatly reduced, this analysis is

arguably underpowered: the 95% confidence intervals bounding the effect sizes increased

and, in many cases, overlapped zero. Furthermore, the bootstrap simulations incorporating

the individual birth-year posteriors estimated from BaSTA confirmed the results of all age-

dependent models (Fig. S4).

DISCUSSION

Increased competition for mating opportunities in species with conventional sex roles is

typically seen as a consequence of anisogamy where the production of many small gametes

allows polygamous males to maximize their reproductive potential (Darwin, 1871; Bateman,

1948; Wade & Shuster, 2002; Schärer *et al.*, 2012; Lehtonen *et al.*, 2016). Polyandry,

whereby females compete more intensively for mates than males, could lead to the evolution

of smaller eggs in a similar way. Andersson (2004) developed a theoretical quantitative

genetic model of female sexual selection by scramble competition over males in an attempt to

explain reduced egg size in polyandrous species. Andersson's (2004) model predicted that

scramble competition for males should lead to egg size reduction and/or early breeding in

sequentially polyandrous females of species with largely invariable clutch sizes. We tested

the predictions of Andersson's (2004) model empirically at the individual level: examining

the links between breeding phenology, individual mating strategies, and egg size variation in

a population of snowy plovers that contains monogamous and polyandrous females (Cruz-

López *et al.*, 2017; Eberhart-Phillips *et al.*, 2017; Kupán *et al.*, 2021). Consistent with

Andersson's (2004) predictions, we found that female polyandry was associated with early

breeding and that early clutches typically contained smaller eggs than clutches laid later in
the season – both between and within individual females. However, we interpret this pattern
differently than Andersson (2004). In our population, early breeding females whose nesting
attempts were successful invariably deserted the brood shortly after hatching (Cruz-López *et*
al., 2017; Kupán *et al.*, 2021) resulting in collinearity between mating behavior and first lay
date. We considered lay date therefore as a proxy for polyandry potential, with early nesting
females having a higher probability to desert their offspring and become polyandrous (unless
their first nest got predated or failed).

Although the statistical associations between breeding phenology, egg size, and female
mating behavior are intriguing, we note that their impact on the egg size variation observed in
this population is rather limited. First, despite snowy plovers' unusual mating system with
sequential polyandry, monogamy, and occasionally polygyny occurring in the same
population (Warriner *et al.*, 1986; Eberhart-Phillips *et al.*, 2017), the egg size variation we
report is not more pronounced than in other birds: the proportional difference between the
largest and smallest egg in our population was approximately 50%, which is of very similar
magnitude to the intraspecific range observed within other bird populations (Christians,
2002). Other constraints may therefore provide limitations on egg size variation in snowy
plovers. Waders produce relatively large eggs that provide vital reserves for their precocial
young and the moderate egg size variation observed in snowy plovers could indicate that
females are unable to reduce egg size further without jeopardizing chick viability. Second, we
note that structural size of females had a stronger effect on egg size variation than breeding
phenology. This is consistent with other studies (Christians, 2002; Giraldo-Deck *et al.*, 2022)
showing that larger females produce larger eggs than smaller females. Third, we found that,
as demonstrated in other studies (e.g., Christians, 2002), female identity explained the largest
part of the variation in egg size, with repeatability of egg size in females was very high. High
repeatability suggests that egg size may have considerable genetic variation, although it has
been suggested that many loci with small effects seem to encode this life history trait
(Santure *et al.*, 2013).

In contrast to egg size, female mating behavior showed a very low individual repeatability
across years. The observed low individual consistency in polyandry is most likely related to
the highly stochastic nature of nest survival, where the main sources of breeding failure in
our population are flooding, predation, and chick mortality through starvation (Cruz-López *et*

al., 2017). Early breeding females whose nests failed, typically stayed together with their
mate for rapid re-nesting (Halimubieke *et al.*, 2019). These females may have intended to
become polygamous but since their replacement clutch hatched later in the season when
conditions for chicks or remating are less favorable, they switched to a monogamous strategy
that involves longer brood care (Kupán *et al.*, 2021). This was supported by the higher
repeatability multi-clutching behavior (i.e., pooling polyandrous and re-nesting females).
Snowy plover females can adjust their brood care efforts flexibly according to the value and
needs of the young (Kupán *et al.*, 2021). Female care and re-mating efforts are inversely
related to each other and show strong seasonality. Early in the breeding season when survival
prospects for the chicks are favorable, re-mating opportunities are high, and their mate can
successfully rear the chicks alone, females typically desert their broods right after hatching of
the chicks to quickly start a new breeding attempt with another male (Székely *et al.*, 1999;
Cruz-López *et al.*, 2017; Kupán *et al.*, 2021). Females whose chicks hatch later in the season,
typically remain with the brood and provide care jointly with their mate. However, desertion
during brood care is tightly associated with chick mortality (Kupán *et al.*, 2021). When
survival prospects of their offspring are deteriorating and chicks die despite the joint care,
initially caring females desert their broods and presumably remate elsewhere where
environmental conditions are better. Such high plasticity in female care and mating strategies
allows females to flexibly respond to nest and chick survival, and is an important ecological
modulator of breeding behavior in snowy plovers. As expected, in years that females could
be polyandrous, they produced more chicks than in years when they were monogamous,
which we attribute to 1) these females laying early enough to allow time for multiple
breeding attempts, and 2) having good fortune that their first nest was not depredated, so that
they could desert and pursue a sequential mate.

Most studies conducted on temperate or high latitude breeding shorebirds have found a
negative association between time of the season and egg size (Byrkjedal & Kalas, 1985;
Sandercock *et al.*, 1999; Kubelka *et al.*, 2020) although in polyandrous red-necked
phalaropes (*Phalaropus lobatus*) egg size increased throughout the breeding season (Kwon et
al., 2018, n.b., the presented effect size is small). However, many investigations of seasonal
egg size dynamics did not disentangle whether the observed changes were due to within- or
between-individual effects. A study that included 15 arctic shorebirds suggested that
between-individual variation may account for more of the seasonal variation in egg size than
within-individual variation (Weiser *et al.*, 2018). In our population, early season nesters

produced smaller eggs on average than females initiating nests at a later date. These seasonal

changes were driven by within individual changes: females generally increased egg volume

between consecutive nesting attempts (albeit the effect size was small), suggesting that

maternal investment during early breeding attempts could be impaired by physiological

limitations. Mutually non-exclusive constraints may include low food availability

(Steigerwald *et al.*, 2015), and carry-over costs associated with intense competition

(Duckworth, 2006) or migratory status (Crossin *et al.*, 2010). In contrast, delayed breeders

can take advantage of a slower pace to gather more resources and maximize investment prior

to clutch initiation – whilst having only enough time for a single breeding attempt. Although

we did not collect any direct empirical data on temporal changes in food availability during

our study (e.g., with pit-fall traps), the invertebrate prey communities of salina ecosystems

(i.e., like that of our study population) are well documented to have dynamic emergence and

quiescent phenologies (Britton & Johnson, 1987), providing important ecological context and

support of our notion that resource availability for egg-laying female snowy plovers is highly

seasonal. The observed decrease of egg size in clutches produced at the end of season is

consistent with such constraints for breeding females as females are in a rush to complete

their clutches and finish incubation before the onset of catastrophic floods caused by

increased precipitation and tide height at the end of the breeding season (Plaschke *et al.*,

2019).

Despite being long-lived – the oldest documented female in our sample was at least 13 years

old (species' longevity record: 20 years; Colwell et al., 2017; M. A. Colwell *personal*

communication, 15 March, 2021) – and investing substantially in reproduction year-after-

year, we found no evidence of age-dependent trade-offs in egg size or polyandry potential.

Contrary to our predictions derived from senescence theory, older females initiated nests

earlier in the season compared to their younger conspecifics – indicating age-dependent

competitive ability that may initially increase the polygamy potential of females (Oring *et al.*,

1994). Age-dependent variation in lay date followed a non-linear pattern: lay date advanced

with each year of age until a peak at age six, after which the age-lay date trajectory was

unclear because of limited sampling in older age classes. In snowy plovers, occasionally

females re-mate with or retain their last mate in the following season (Halimubieke *et al.*,

2019), presumably to skip the time consuming mate search and start breeding early – a

process that could also explain the age-dependent advance in lay date we observed.

Furthermore, locally recruited females (i.e., hatched locally) bred earlier than immigrant

females indicating a role for prior experience giving competitive advantage for breeders in
our population.

Egg size was not clearly associated with female age. Several studies of oviparous organisms
have observed age-dependent variation in egg size, with some studies finding a positive
relationship (Cooch *et al.*, 1992; Flint & Sedinger, 1992; Robertson *et al.*, 1994; Warner *et*
al., 2016; Verhoeven *et al.*, 2020) and others observing a negative relationship (Reid, 1988;
Potti, 1993; Ito, 1997). Recent longitudinal studies of wild avian systems showed a quadratic
relationship with a prominent increase in egg size in early life, followed by a peak and then a
late-life decline (Bouwhuis *et al.*, 2009; Jankowiak *et al.*, 2018). In captive avian systems, the
early life increase in egg size is also observed, but not the late-life decline likely due to *ad*
libitum food (Woodard & Abplanalp, 1971). In our study, a quadratic-age model of
senescence in egg size was not statistically clear, and did not change when we repeated the
analysis including only the 45 females of known age (10% of the original sample). We
acknowledge that our relatively small sample of known-age individuals presents a limitation
to our study – we cannot exclude a moderate effect of senescence on egg size. Nevertheless,
potential explanations for the lack of a clear senescence trend observed in egg size are that 1)
our open- and free-living population lacks extremely old individuals that would exhibit the
intrinsic "costs paid" to somatic loss, and/or 2) there is little room for further reduction
because of viability constraints for the chicks. In shorebirds, chicks are not fed by the parents
but rather must forage for themselves immediately after hatching and larger chicks that hatch
from larger eggs typically survive better than smaller chicks (Blomqvist *et al.*, 1997; Giraldo-
Deck *et al.*, 2022). The extra nutrients provided by a large egg may be crucial for the chick
survival during the first few days of life when foraging efficiency is reduced as chicks still
learn how to catch prey (Ricklefs, 1968). Consequently, additional reductions of egg size
may not be feasible as the knock-on effects for chick survival are too costly (Williams, 1994;
Starck & Ricklefs, 1998), therefore limiting a females' potential to save resources by
reducing egg volume. Chick-viability constraints may also provide an explanation why the
variation in egg size at the population level is relatively low.

Conclusions
Past species-level studies have linked polyandry and sex-role reversal to reduced female
gamete size (Slotow, 1996; Andersson, 2004), as smaller eggs would permit females to
produce several clutches in rapid succession (Liker *et al.*, 2001). In our individual-level

investigation we found that early nesting females were more likely to be polyandrous and
produced clutches containing smaller eggs than females that started their first breeding
attempt later in the season. Starting to breed early in the season is an important pre-
determinant of female polygamy. Prior experience can provide both older and local females a
competitive advantage over younger and naïve conspecifics scrambling for early breeding
opportunities that are at the heart of the polyandrous mating strategy (Andersson, 2004).
However, we found no evidence supporting an adaptive egg-size adjustment to obtain a
competitive advantage in the scramble competition for mates. Rather we suspect that the
observed seasonal variation in egg size can also be explained by other constraints that include
a survival advantage for early hatching chicks, resource constraints during the egg-producing
phase and, most importantly, between individual differences. Senescence did not clearly have
a major impact on egg size variation nor on the mating behavior of females. Future research
of polyandrous systems should also examine the impact of seasonality, age, and mating
behavior on the reproductive fitness of males, as males provide a larger share of parental care
than females in these systems and therefore should have a more direct impact on the survival
prospects of the young. Examining the effects of season- and age-dependent trade-offs on
individual fitness and variation in mating tactics will enable a deeper understanding about
selective pressures and constraints of diverse and plastic sex roles in natural populations.

DATA ACCESSIBILITY

We provide all computer code and documentation as a html vignette written in Rmarkdown

(click here: File S1) together with all the raw datasets needed to reproduce our modeling and

analyses – these files can be found in this project's Open Science Framework repository:

doi.org/ 10.17605/OSF.IO/UCW6J

LITERATURE CITED

Amat, J.A., Fraga, R.M. & Arroyo, G.M. 1999. Brood desertion and polygamous breeding in
the Kentish plover Charadrius alexandrinus. *Ibis* **141**: 596–607.

Andersson, M. 2005. Evolution of Classical Polyandry: Three Steps to Female Emancipation.
Ethology **111**: 1–23.

Andersson, M. 2004. Social polyandry, parental Investment, sexual selection, and evolution
of reduced female gamete size. *Evolution* **58**: 24–34. [Society for the Study of
Evolution, Wiley].

Argüelles-Tico, A., Küpper, C., Kelsh, R.N., Kosztolányi, A., Székely, T. & van Dijk, R.E.
2015. Geographic variation in breeding system and environment predicts melanin-
based plumage ornamentation of male and female Kentish plovers. *Behavioral
Ecology and Sociobiology* **70**: 49–60.

Bateman, A.J. 1948. Intra-sexual selection in Drosophila. *Heredity* **2**: 349–368. Essex.

Bates, D., Mächler, M., Bolker, B. & Walker, S. 2015. Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software* **67**.

Beamonte-Barrientos, R., Velando, A., Drummond, H. & Torres, R. 2010. Senescence of
Maternal Effects: Aging Influences Egg Quality and Rearing Capacities of a Long-
Lived Bird. *The American Naturalist* **175**: 469–480.

Beck, S.V., Räsänen, K., Kristjánsson, B.K., Skúlason, S., Jónsson, Z.O., Tsinganis, M., *et
al.* 2022. Variation in egg size and offspring phenotype among and within seven
Arctic charr morphs. *Ecology and Evolution* **12**: e9427.

Bernardo, J. 1996. The Particular Maternal Effect of Propagule Size, Especially Egg Size:
Patterns, Models, Quality of Evidence and Interpretations1. *American Zoologist* **36**:
216–236.

Blomqvist, D., Johansson, O.C. & Götmark, F. 1997. Parental quality and egg size affect
chick survival in a precocial bird, the lapwing Vanellus vanellus. *Oecologia* **110**: 18–
24.

Borgmann, K.L., Conway, C.J. & Morrison, M.L. 2013. Breeding phenology of birds:
mechanisms underlying seasonal declines in the risk of nest predation. *PLoS ONE* **8**:
e65909.

Bouwhuis, S., Charmantier, A., Verhulst, S. & Sheldon, B.C. 2010. Trans-generational
effects on ageing in a wild bird population. *Journal of Evolutionary Biology* **23**: 636–
642.

Bouwhuis, S., Sheldon, B.C., Verhulst, S. & Charmantier, A. 2009. Great tits growing old:
selective disappearance and the partitioning of senescence to stages within the
breeding cycle. *Proceedings of the Royal Society B: Biological Sciences* **276**: 2769–
2777.

Britton, R.H. & Johnson, A.R. 1987. An ecological account of a Mediterranean salina: The
Salin de Giraud, Camargue (S. France). *Biological Conservation* **42**: 185–230.

Byrkjedal, I. & Kalas, J.A. 1985. Seasonal variation in egg size in golden plover Pluvialis
apricaria and dotterel Charadrius morinellus populations. *Ornis Scandinavica* **16**:
108–112. WileyNordic Society Oikos.

Carmona-Isunza, M.C., Ancona, S., Székely, T., Ramallo-González, A.P., Cruz-López, M.,
Serrano-Meneses, M.A., *et al.* 2017. Adult sex ratio and operational sex ratio exhibit
different temporal dynamics in the wild. *Behavioral Ecology* **28**: 523–532.

Christians, J.K. 2002. Avian egg size: variation within species and inflexibility within
individuals. *Biological reviews* **77**: 1–26. Cambridge University Press.

Clutton-Brock, T. 2007. Sexual Selection in Males and Females. *Science* **318**: 1882–1885.
American Association for the Advancement of Science.

Colchero, F., Jones, O.R. & Rebke, M. 2012. BaSTA: an R package for Bayesian estimation
of age-specific survival from incomplete mark–recapture/recovery data with
covariates. *Methods in Ecology and Evolution* **3**: 466–470.

Colwell, M.A. 2010. *Shorebird Ecology, Conservation, and Management*. University of
California Press.

Colwell, M.A., Feucht, E.J., McAllister, S.E. & Transou, A.N. 2017. Lessons learned from
the oldest Snowy Plover. *Wader Study* **124**: 157–159.

Cooch, E.G., Lank, D.B., Rockwell, R.F. & Cooke, F. 1992. Is There a Positive Relationship
between Body Size and Fecundity in Lesser Snow Geese? *The Auk* **109**: 667–673.

Crossin, G.T., Trathan, P.N., Phillips, R.A., Dawson, A., Le Bouard, F. & Williams, T.D.
2010. A carryover effect of migration underlies individual variation in reproductive
readiness and extreme egg size dimorphism in macaroni penguins. *The American*
Naturalist **176**: 357–366.

Cruz-López, M., Eberhart-Phillips, L.J., Fernández, G., Beamonte-Barrientos, R., Székely,
T., Serrano-Meneses, M.A., *et al.* 2017. The plight of a plover: Viability of an
important snowy plover population with flexible brood care in Mexico. *Biological*
Conservation **209**: 440–448.

Darwin, C. 1871. *The descent of man and selection in relation to sex*. Murray, London.

Dingemanse, N.J., Moiron, M., Araya-Ajoy, Y.G., Mouchet, A. & Abbey-Lee, R.N. 2020.
Individual variation in age-dependent reproduction: Fast explorers live fast but
senesce young? *Journal of Animal Ecology* **89**: 601–613.

Dittmann, T. & Hötker, H. 2001. Intraspecific Variation in the Egg Size of the Pied Avocet.
Waterbirds: The International Journal of Waterbird Biology **24**: 83–88.

dos Remedios, N., Lee, P.L., Székely, T., Dawson, D.A. & Küpper, C. 2010. Molecular sex-
typing in shorebirds: a review of an essential method for research in evolution,
ecology and conservation. *Wader Study Group Bulletin* **117**: 109–118.

Duckworth, R.A. 2006. Aggressive behaviour affects selection on morphology by influencing
settlement patterns in a passerine bird. *Proceedings of the Royal Society B: Biological*
Sciences **273**: 1789–1795.

Eberhart-Phillips, L.J. 2019. Plover Breeding Systems. In: *The Population Ecology and*
Conservation of Charadrius Plovers (M. A. Colwell & S. M. Haig, eds), pp. 65–88.
New York.

Eberhart-Phillips, L.J., Cruz-López, M., Lozano-Angulo, L., del Ángel, S.G., Rojas-Abreu,
W., Bucio-Pacheco, M., *et al.* 2020a. CeutaOPEN, individual-based field observations
of breeding snowy plovers Charadrius nivosus. *Sci Data* **7**: 149.

Eberhart-Phillips, L.J., Cruz-López, M., Lozano-Angulo, L., Gómez del Ángel, S., Rojas-
Abreu, W., Bucio-Pacheco, M., *et al.* 2020b. CeutaOPEN v1.5. Open Science
Framework.

Eberhart-Phillips, L.J., Küpper, C., Miller, T.E.X., Cruz-López, M., Maher, K.H., Dos
Remedios, N., *et al.* 2017. Sex-specific early survival drives adult sex ratio bias in
snowy plovers and impacts mating system and population growth. *Proceedings of the*
National Academy of Sciences of the United States of America **114**: e5474–e5481.

Fay, R., Ravussin, P.-A., Arrigo, D., von Rönn, J.A.C. & Schaub, M. 2021. Age-specific
reproduction in female pied flycatchers: evidence for asynchronous aging. *Oecologia*
196: 723–734.

Festa-Bianchet, M. 1988. Birthdate and survival in bighorn lambs (Ovis canadensis). *Journal*
of Zoology **214**: 653–661.

Flint, P.L. & Sedinger, J.S. 1992. Reproductive Implications of Egg-Size Variation in the
Black Brant. *The Auk* **109**: 896–903.

Fox, C.W. 1994. The Influence of Egg Size on Offspring Performance in the Seed Beetle,
Callosobruchus maculatus. *Oikos* **71**: 321–325.

Fritzsche, K., Henshaw, J.M., Johnson, B.D. & Jones, A.G. 2021. The 150th anniversary of
The Descent of Man: Darwin and the impact of sex-role reversal on sexual selection
research. *Biological Journal of the Linnean Society* **134**: 525–540.

Fromonteil, S., Marie-Orleach, L., Winkler, L. & Janicke, T. 2023. Sexual selection in
females and the evolution of polyandry. *PLOS Biology* **21**: e3001916. Public Library
of Science.

Giraldo-Deck, L.M., Loveland, J.L., Goymann, W., Tschirren, B., Burke, T., Kempenaers,
B., *et al.* 2022. Intralocus conflicts associated with a supergene. *Nat Commun* **13**:
1384. Nature Publishing Group.

Goymann, W., Makomba, M., Urasa, F. & Schwabl, I. 2015. Social monogamy vs.
polyandry: ecological factors associated with sex roles in two closely related birds
within the same habitat. *Journal of Evolutionary Biology* **28**: 1335–1353.

Graham, J.L., Bauer, C.M., Heidinger, B.J., Ketterson, E.D. & Greives, T.J. 2019. Early-
breeding females experience greater telomere loss. *Molecular Ecology* **28**: 114–126.

Griffiths, R., DOUBLE, M.C., ORR, K. & DAWSON, R.J.G. 1998. A DNA test to sex most
birds. *Molecular Ecology* **7**: 1071−1075.

Halimubieke, N., Valdebenito, J.O., Harding, P., Cruz-López, M., Serrano-Meneses, M.A.,
James, R., *et al.* 2019. Mate fidelity in a polygamous shorebird, the snowy plover
(Charadrius nivosus). *Ecology and Evolution* **9**: 10734−10745.

Hall, L.K. & Cavitt, J.F. 2012. Comparative Study of Trapping Methods for Ground-Nesting
Shorebirds. *Waterbirds* **35**: 342−346.

Hammers, M., Richardson, D.S., Burke, T. & Komdeur, J. 2012. Age-dependent terminal
declines in reproductive output in a wild bird. *PLoS ONE* **7**: e40413.

Hanson, H.C. 1954. Apparatus for the Study of Incubated Bird Eggs. *The Journal of Wildlife*
Management **18**: 191−198.

Herborn, K.A., Daunt, F., Heidinger, B.J., Wilding, H.M.V.G., Burthe, S.J., Newell, M.A., *et*
al. 2016. Age, oxidative stress exposure and fitness in a long-lived seabird.
Functional Ecology **30**: 913−921.

Hoyt, D.F. 1979. Practical Methods of Estimating Volume and Fresh Weight of Bird Eggs.
The Auk **96**: 73−77.

Immler, S., Pitnick, S., Parker, G.A., Durrant, K.L., Lüpold, S., Calhim, S., *et al.* 2011.
Resolving variation in the reproductive tradeoff between sperm size and number.
Proceedings of the National Academy of Sciences **108**: 5325−5330. National Acad
Sciences.

Ito, K. 1997. Egg-size and -number variations related to maternal size and age, and the
relationship between egg size and larval characteristics in an annual marine
gastropod, Haloa japonica (Opisthobranchia; Cephalaspidea). *Marine Ecology*
Progress Series **152**: 187−195.

Janicke, T., Häderer, I.K., Lajeunesse, M.J. & Anthes, N. 2016. Darwinian sex roles
confirmed across the animal kingdom. *Science Advances* **2**: e1500983. American
Association for the Advancement of Science.

Jankowiak, \Lukasz, Zyskowski, D. & Wysocki, D. 2018. Age-specific reproduction and
disposable soma in an urban population of Common Blackbirds Turdus merula. *Ibis*
160: 130−144.

Kaplan, R.H. 1980. The Implications of Ovum Size Variability for Offspring Fitness and
Clutch Size within Several Populations of Salamanders (Ambystoma). *Evolution* **34**:
51−64.

Kirkwood, T.B.L. & Rose, M.R. 1991. Evolution of senescence: late survival sacrificed for
reproduction. *Philosophical Transactions of the Royal Society B-Biological Sciences*
332: 15−24.

Kraaijeveld, K., Kraaijeveld-Smit, F.J.L. & Komdeur, J. 2007. The evolution of mutual
ornamentation. *Animal Behaviour* **74**: 657−677.

Krist, M. 2011. Egg size and offspring quality: a meta-analysis in birds. *Biological Reviews*
86: 692–716.

Kubelka, V., Sládeček, M., Zámečník, V., Vozabulová, E. & Šálek, M. 2020. Seasonality
Predicts Egg Size Better Than Nesting Habitat in a Precocial Shorebird. *Ardea* **107**:
239.

Kupán, K., Székely, T., Cruz-López, M., Seymour, K. & Küpper, C. 2021. Offspring
desertion with care? Chick mortality and plastic female desertion in Snowy Plovers.
Behavioral Ecology.

Küpper, C., Augustin, J., Kosztolányi, A., Burke, T., Flguerola, J. & Székely, T. 2009a.
Kentish Versus Snowy Plover: Phenotypic and Genetic Analyses of Charadrius
alexandrinus Reveal Divergence of Eurasian and American Subspecies. *The Auk* **126**:
839–852.

Küpper, C., Augustin, J., Kosztolányi, A., Burke, T., Flguerola, J. & Székely, T. 2009b.
Kentish Versus Snowy Plover: Phenotypic and Genetic Analyses of Charadrius
alexandrinus Reveal Divergence of Eurasian and American Subspecies. *The Auk* **126**:
839–852.

Kwon, E., English, W.B., Weiser, E.L., Franks, S.E., Hodkinson, D.J., Lank, D.B., *et al.*
2018. Delayed egg-laying and shortened incubation duration of Arctic-breeding
shorebirds coincide with climate cooling. *Ecology and Evolution* **8**: 1339–1351.

Lack, D. 1968. *Ecological adaptations for breeding in birds*. Methuen Ltd, London.

Lehtonen, J., Parker, G.A. & Schärer, L. 2016. Why anisogamy drives ancestral sex roles.
Evolution **70**: 1129–1135.

Lemaître, J.-F., Berger, V., Bonenfant, C., Douhard, M., Gamelon, M., Plard, F., *et al.* 2015.
Early-late life trade-offs and the evolution of ageing in the wild. *Proceedings of the
Royal Society B: Biological Sciences* **282**: 20150209.

Liker, A., Reynolds, J.D. & Székely, T. 2001. The evolution of egg size in socially
polyandrous shorebirds. *Oikos* **95**: 3–14.

Moran, A.L. & Emlet, R.B. 2001. Offspring size and performance in variable environments:
field studies on a marine snail. *Ecology* **82**: 1597–1612.

Morrison, C.A., Alves, J.A., Gunnarsson, T.G., Þórisson, B. & Gill, J.A. 2019. Why do
earlier-arriving migratory birds have better breeding success? *Ecology and Evolution*
9: 8856–8864.

Nakagawa, S. & Schielzeth, H. 2010. Repeatability for Gaussian and non-Gaussian data: a
practical guide for biologists. *Biological Reviews* **85**: 935–956.

Nosály, G. & Székely, T. 1993. Clutch and egg-size variation in the Kentish Plover
(\emphCharadrius alexandrinus) during the breeding season. *Aquila* **100**: 161–179.

Nussey, D.H., Coulson, T., Bianchet, M.F. & Gaillard, J.M. 2008. Measuring senescence in
wild animal populations: towards a longitudinal approach. *Functional Ecology* **22**:
393–406.

Ockendon, N., Leech, D. & Pearce-Higgins, J.W. 2013. Climatic effects on breeding grounds
are more important drivers of breeding phenology in migrant birds than carry-over
effects from wintering grounds. *Biology Letters* **9**: 20130669.

Oring, L.W. 1986. Avian Polyandry. In: *Current Ornithology, vol 3* (R. F. Johnston, ed), pp.
309–351.

Oring, L.W. & Lank, D.B. 1982. Sexual selection, arrival times, philopatry and site fidelity in
the polyandrous spotted sandpiper. *Behavioral Ecology and Sociobiology* **10**: 185–
191.

Oring, L.W., Reed, M.J. & Alberico, A.J. 1994. Mate acquisition tactics in polyandrous
spotted sandpipers (Actitis macularia): the role of age and experience. *Behavioral*
Ecology **5**: 9–16.

Page, G.W., Stenzel, L.E., Warriner, J.S., Warriner, J.C. & Paton, P.W. 2009. Snowy Plover
(Charadrius nivosus).

Plard, F., Gaillard, J.-M., Coulson, T., Hewison, A.J.M., Douhard, M., Klein, F., *et al.* 2015.
The influence of birth date via body mass on individual fitness in a long-lived
mammal. *Ecology* **96**: 1516–1528.

Plaschke, S., Bulla, M., Cruz-López, M., Gómez del Ángel, S. & Küpper, C. 2019. Nest
initiation and flooding in response to season and semi-lunar spring tides in a ground-
nesting shorebird. *Frontiers in Zoology* **16**: 15–11.

Potti, J. 1993. Environmental, ontogenetic, and genetic variation in egg size of Pied
Flycatchers. *Canadian Journal of Zoology-revue Canadienne De Zoologie - CAN J*
ZOOL **71**: 1534–1542.

R Core Team. 2023. *R: A Language and Environment for Statistical Computing*. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Rahn, H., Paganelli, V.C. & Ar, A. 1975. Relation of Avian Egg Weight to Body Weight.
The Auk **92**: 750–765.

Reid, J.M., Bignal, E.M., Bignal, S., McCracken, D.I. & Monaghan, P. 2003. Age-specific
reproductive performance in red-billed choughs Pyrrhocorax pyrrhocorax: patterns
and processes in a natural population. *Journal of Animal Ecology* **72**: 765–776.

Reid, V.W. 1988. Age-Specific Patterns of Reproduction in the Glaucous-Winged Gull:
Increased Effort with Age. *Ecology* **69**: 1454–1465.

Reznick, D., Schultz, E., Morey, S. & Roff, D. 2006. On the virtue of being the first born: the
influence of date of birth on fitness in the mosquitofish, Gambusia affinis. *Oikos* **114**:
135–147.

Reznick, D.N., Bryant, M.J., Roff, D., Ghalambor, C.K. & Ghalambor, D.E. 2004. Effect of
extrinsic mortality on the evolution of senescence in guppies. *Nature* **431**: 1095–
1099. Nature Publishing Group.

Ricklefs, R.E. 1968. Patterns of growth in birds. *Ibis* **110**: 419–451.

Riecke, T.V., Hegelbach, J. & Schaub, M. n.d. Reproductive senescence and mating tactic
interact and conflict to drive reproductive success in a passerine. *Journal of Animal
Ecology* **n/a**.

Robertson, G.J., Cooch, E.G., Lank, D.B., Rockwell, R.F. & Cooke, F. 1994. Female Age
and Egg Size in the Lesser Snow Goose. *Journal of Avian Biology* **25**: 149.

Rosenqvist, G. 1990. Male mate choice and female-female competition for mates in the
pipefish Nerophis ophidion. *Animal Behaviour* **39**: 1110–1115.

Salguero-Gómez, R., Jones, O.R., Jongejans, E., Blomberg, S.P., Hodgson, D.J., Mbeau-
Ache, C., *et al.* 2016. Fast-slow continuum and reproductive strategies structure plant
life-history variation worldwide. *Proceedings of the National Academy of Sciences of
the United States of America* **113**: 230–235.

Sandercock, B.K., Lank, D.B. & Cooke, F. 1999. Seasonal Declines in the Fecundity of
Arctic-Breeding Sandpipers: Different Tactics in Two Species with an Invariant
Clutch Size. *Journal of Avian Biology* **30**: 460. JSTOR.

Santure, A.W., De Cauwer, I., Robinson, M.R., Poissant, J., Sheldon, B.C. & Slate, J. 2013.
Genomic dissection of variation in clutch size and egg mass in a wild great tit (Parus
major) population. *Molecular Ecology* **22**: 3949–3962.

Schärer, L., Rowe, L. & Arnqvist, G. 2012. Anisogamy, chance and the evolution of sex
roles. *Trends in Ecology & Evolution* **27**: 260–264.

Schroeder, J., Burke, T., Mannarelli, M.E., Dawson, D.A. & Nakagawa, S. 2012. Maternal
effects and heritability of annual productivity. *Journal of Evolutionary Biology* **25**:
149–156.

Skrade, P. & Dinsmore, S.J. 2013. Egg-size investment in a bird with uniparental incubation
by both sexes. *The Condor* **115**: 508–514.

Slagsvold, T., Sandvik, J., Rofstad, G., Lorentsen, Ö. & Husby, M. 1984. On the Adaptive
Value of Intraclutch Egg-Size Variation in Birds. *The Auk* **101**: 685–697.

Slotow, R. 1996. Black coucal Centropus grillii egg volume predicts their polyandrous
mating system. *Journal of Avian Biology* 171–173.

Smith, C.C. & Fretwell, S.D. 1974. The Optimal Balance between Size and Number of
Offspring. *The American Naturalist* **108**: 499–506. The University of Chicago Press.

Snijders, T. & Bosker, R.J. 2011. \emphMultilevel analysis: An introduction to basic and
advanced multilevel modeling*, 2nd ed. Sage.

Starck, J.M. & Ricklefs, R.E. 1998. *Avian growth and development. Evolution within the*
altricial-precocial spectrum. New York.

Steigerwald, E.C., Igual, J.-M., Payo-Payo, A. & Tavecchia, G. 2015. Effects of decreased
anthropogenic food availability on an opportunistic gull: evidence for a size-mediated
response in breeding females. *Ibis* **157**: 439–448.

Stenzel, L.E., Page, G.W., Warriner, J.C., Warriner, J.S., Neuman, K.K., George, D.E., *et al.*
2011. Male-skewed adult sex ratio, survival, mating opportunity and annual
productivity in the snowy plover \emphCharadrius alexandrinus. *Ibis* **153**: 312–322.

Stoffel, M.A., Nakagawa, S. & Schielzeth, H. 2020. partR2: Partitioning R2 in generalized
linear mixed models. *bioRxiv* 2020.07.26.221168.

Stoffel, M.A., Nakagawa, S. & Schielzeth, H. 2017. rptR: repeatability estimation and
variance decomposition by generalized linear mixed-effects models. *Methods in*
Ecology and Evolution **8**: 1639–1644.

Svagelj, W.S. & Quintana, F. 2011. Egg-size variation in the Imperial Cormorant: on the
importance of individual effects. *The condor* **113**: 528–537. Oxford University Press.

Székely, T., Argüelles-Tico, A., Kosztolányi, A. & Küpper, C. 2011. Practical guide for
investigating breeding ecology of Kentish plover. 1–16.

Székely, T., Cuthill, I.C. & Kis, J. 1999. Brood desertion in Kentish plover sex differences in
remating opportunities. *Behavioral Ecology* **10**: 185–190.

Székely, T., Kosztolányi, A. & Küpper, C. 2008. *Practical guide for investigating breeding*
ecology of Kentish plover Charadrius alexandrinus. University of Bath, Bath, UK.

Székely, T., Kozma, J. & Piti, A. 1994. The Volume of Snowy Plover Eggs (El Volumen de
los Huevos de Charadrius alexandrinus). *Journal of Field Ornithology* **65**: 60–64.

van de Pol, M. & Verhulst, S. 2006a. Age-Dependent Traits: A New Statistical Model to
Separate Within- and Between-Individual Effects. *The American Naturalist* **167**: 766–
773.

van de Pol, M. & Verhulst, S. 2006b. Age☐Dependent Traits: A New Statistical Model to
Separate Within☐ and Between☐Individual Effects. *The American Naturalist* **167**:
766–773. The University of Chicago Press.

Varpe, Ø. 2017. Life History Adaptations to Seasonality. *Integrative and Comparative*
Biology **57**: 943–960.

Verhoeven, M.A., Loonstra, A.H.J., McBride, A.D., Tinbergen, J.M., Kentie, R.,
Hooijmeijer, J.C.E.W., *et al.* 2020. Variation in Egg Size of Black-Tailed Godwits.
Ardea **107**: 291.

Vincze, O., Kosztolányi, A., Barta, Z., Küpper, C., Alrashidi, M., Amat, J.A., *et al.* 2017.
Parental cooperation in a changing climate: fluctuating environments predict shifts in
care division. *Global Ecology and Biogeography* **26**: 347–358.

Wade, M.J. & Shuster, S.M. 2002. The evolution of parental care in the context of sexual
selection: a critical reassessment of parental investment theory. *The American*
Naturalist **160**: 285–292. The University of Chicago Press.

Wallander, J. & Andersson, M. 2003. Reproductive tactics of the ringed plover Charadrius
hiaticula. *Journal of Avian Biology* **34**: 259–266.

Warner, D.A., Miller, D.A.W., Bronikowski, A.M. & Janzen, F.J. 2016. Decades of field data
reveal that turtles senesce in the wild. *Proceedings of the National Academy of*
Sciences of the United States of America **113**: 6502–6507.

Warriner, J.S., Warriner, J.C., Page, G.W. & Stenzel, L.E. 1986. Mating System and
Reproductive Success of a Small Population of Polygamous Snowy Plovers. *The*
Wilson Bulletin **98**: 15–37. Wilson Ornithological Society.

Weiser, E.L., Brown, S.C., Lanctot, R.B., Gates, H.R., Abraham, K.F., Bentzen, R.L., *et al.*
2018. Life-history tradeoffs revealed by seasonal declines in reproductive traits of
Arctic-breeding shorebirds. *Journal of Avian Biology* **49**: jav-01531.

Whittingham, L.A. & Dunn, P.O. 2010. Fitness benefits of polyandry for experienced
females. *Molecular Ecology* **19**: 2328–2335.

Williams, T.D. 1994. Intraespecific Variation in Egg Size and Egg Composition in Birds:
Effects on Offspring Fitness. *Biological Reviews of the Cambridge Philosophical*
Society **69**: 35–59.

Williams, T.D. 2012. *Physiological adaptations for breeding in birds*. Princenton.

Woodard, A.E. & Abplanalp, H. 1971. Longevity and Reproduction in Japanese Quail
Maintained Under Stimulatory Lighting. *Poultry Science* **50**: 688–692.

Xu, F., Yang, W. & Li, Y. 2019. Enlarged Egg Size Increases Offspring Fitness of a Frog
Species on the Zhoushan Archipelago of China. *Scientific Reports* **9**: 11653.

Yohannes, E., Valcu, M., Lee, R.W. & Kempenaers, B. 2010. Resource use for reproduction
depends on spring arrival time and wintering area in an arctic breeding shorebird.
Journal of Avian Biology **41**: 580–590.

Zhang, H., Vedder, O., Becker, P.H. & Bouwhuis, S. 2015. Age-dependent trait variation: the
relative contribution of within-individual change, selective appearance and
disappearance in a long-lived seabird. *Journal of Animal Ecology* **84**: 797–807.

 TABLES
 Table 1. Predictors of a female's polyandry probability in a given year: the lay date and
963   average egg volume of their first nest of the year, their age-deviance (i.e., years since first
964   local breeding attempt), their age at first breeding, and their structural body size. Fixed effect
965   sizes shown are standardized estimates on the logit scale.

Polyandry probability	Mean estimate	95% confidence interval
Fixed effects β (logit standardized)		
First nest lay date	-2.12	[-2.87, -1.68]
Avg. egg volume of first nest	0.00	[-0.29, 0.29]
Age-deviance	0.00	[-0.28, 0.26]
First breeding age	0.22	[-0.05, 0.5]
Female tarsus length	0.19	[-0.09, 0.5]
Partitioned R^2		
Total Marginal R^2	0.34	[0.24, 0.46]
Total Conditional R^2	0.36	[0.25, 0.54]
First nest lay date	0.33	[0.22, 0.45]
Avg. egg volume of first nest	0.00	[0, 0.2]
Age-deviance	0.00	[0, 0.2]
First breeding age	0.01	[0, 0.2]
Female tarsus length	0.00	[0, 0.2]
Random effects σ^2		
Individual	0.07	[0, 1.78]
Year	0.16	[0, 0.57]
Adjusted repeatability r		
Individual	0.01	[0, 0.17]
Year	0.02	[0, 0.06]
Sample sizes n		
Years	14	
Individuals	424	
Observations (i.e., Nests)	661	

966 **Table 2.** Predictors of a female's re-nesting probability in a given year: the lay date of their
967 first nest of the year, their age-deviance (i.e., years since first local breeding attempt), their
968 age at first breeding, and their structural body size. Fixed effect sizes shown are standardized
969 estimates on the logit scale.

Re-nesting probability	Mean estimate	95% confidence interval
Fixed effects β (logit standardized)		
First nest lay date	-1.79	[-5.79, -1.31]
Age-deviance	0.20	[-0.19, 0.73]
First breeding age	0.41	[0.06, 1.7]
Female tarsus length	-0.09	[-0.47, 0.29]
Partitioned R^2		
Total Marginal R^2	0.42	[0.29, 0.61]
Total Conditional R^2	0.44	[0.3, 0.96]
First nest lay date	0.40	[0.27, 0.6]
Age-deviance	0.01	[0, 0.43]
First breeding age	0.03	[0, 0.44]
Female tarsus length	0.00	[0, 0.43]
Random effects σ^2		
Individual	0.39	[0, 33.96]
Year	0.00	[0, 1.67]
Adjusted repeatability r		
Individual	0.03	[0, 14.04]
Year	0.00	[0, 0.21]
Sample sizes n		
Years	13	
Individuals	177	
Observations (i.e., Nests)	209	

Table 3. Sources of lay date variation. Note: within the *Partitioned R^2* sub-table the term
971   'Senescence' describes the collective variation explained by the linear and quadratic within-
972   individual age-deviance effects in the top sub-table, the term 'Selective appearance' and
973   'Selective disappearance' describe the variation explained by the between individual first-
974   and last-breeding age fixed effects of the top sub-table, respectively.

Lay date of first nest	*Mean estimate*	*95% confidence interval*
Fixed effects β (standardized)		
Mother tarsus length	0.02	[-0.07, 0.1]
Local recruit	-7.80	[-13.17, -3.05]
Within ind. age-deviance (linear)	-0.16	[-0.26, -0.05]
Within ind. age-deviance (quadratic)	0.09	[0.01, 0.17]
Between ind. first breeding age	0.06	[-0.04, 0.16]
Between ind. last breeding age	-0.04	[-0.18, 0.1]
Partitioned R^2		
Total Marginal R^2	0.06	[0.04, 0.12]
Total Conditional R^2	0.23	[0.13, 0.38]
Mother tarsus length	0.00	[0, 0.06]
Local recruit	0.02	[0, 0.08]
Senescence	0.02	[0, 0.07]
Selective appearance	0.01	[0, 0.06]
Selective disappearance	0.00	[0, 0.06]
Random effects σ^2		
Individual	60.69	[18.94, 117.67]
Year	2.00	[0, 12.89]
Residual	283.38	[229.95, 326.82]
Adjusted repeatability r		
Individual	0.19	[0.06, 0.32]
Year	0.01	[0, 0.03]
Residual	0.80	[0.67, 0.93]
Sample sizes n		
Years	13	
Individuals	376	
Observations (i.e., Nests)	568	

Table 4. (*displayed on next page*) Sources of egg size variation. Note: within the *Partitioned R^2* sub-table the term 'Senescence' describes the collective variation explained by the linear and quadratic within-individual age-deviance effects in the top sub-table, the term 'Selective appearance' and 'Selective disappearance' describe the variation explained by the between individual first- and last-breeding age fixed effects of the top sub-table, respectively; the term 'Within ind. seasonality' describes the variation explained by the within individual lay-date deviance effect in the top sub-table, and the term 'Between ind. seasonality' describes the collective variation explained by the linear and quadratic lay date effects in the top sub-table.

 Table 4. (*caption on previous page*)

Egg volume	Mean estimate	95% confidence interval
Fixed effects β (standardized)		
Mother tarsus length	0.22	[0.14, 0.28]
Clutch size (< 3 eggs)	-0.03	[-0.11, 0.04]
Within ind. age-deviance (linear)	-0.01	[-0.08, 0.05]
Within ind. age-deviance (quadratic)	-0.04	[-0.09, 0]
Between ind. first breeding age	0.04	[-0.04, 0.13]
Between ind. last breeding age	0.03	[-0.1, 0.15]
Within ind. lay-date-deviance	0.09	[0.05, 0.13]
Between ind. linear lay date	0.16	[0.1, 0.21]
Between ind. quadratic lay date	-0.06	[-0.12, -0.01]
Partitioned R^2		
Total Marginal R^2	0.07	[0.05, 0.12]
Total Conditional R^2	0.66	[0.63, 0.7]
Mother tarsus length	0.04	[0.02, 0.09]
Clutch size (< 3 eggs)	0.00	[0, 0.05]
Senescence	0.00	[0, 0.05]
Selective appearance	0.00	[0, 0.05]
Selective disappearance	0.00	[0, 0.05]
Within ind. seasonality	0.01	[0, 0.05]
Between ind. seasonality	0.01	[0, 0.06]
Random effects σ^2		
Nest / Individual	0.03	[0.03, 0.04]
Individual	0.10	[0.08, 0.12]
Year	0.01	[0, 0.01]
Residual	0.08	[0.07, 0.08]
Adjusted repeatability r		
Nest / Individual	0.16	[0.12, 0.2]
Individual	0.45	[0.39, 0.5]
Year	0.03	[0.01, 0.06]
Residual	0.37	[0.33, 0.41]
Sample sizes n		
Years	14	
Individuals	424	
Nests	836	
Observations (i.e., Eggs)	2393	

Figure 1. Egg size variation for snowy plovers breeding in Bahía de Ceuta, Mexico between
2006 and 2020. Egg volume was estimated using an allometric equation combining length
(A) and width (B) measurements collected in the field (see *Methods*). C) Average egg volume
of a clutch was highly correlated with the clutch's average chick weight at hatch (β [95%
CIs]: 0.628 [0.552–0.704]; $R^2_{marginal}$ = 0.370 [0.310–0.436]). Ribbon shows the 95% CI of the
model prediction, vertical and horizontal error bars visualize within clutch variation in egg
volume and chick hatch weight (1 SD); see Supplementary Text D for modelling methods.

Figure 2. Phenology of mating behaviors, egg laying, and egg size in 424 female snowy plovers breeding at Bahía de Ceuta. A) Relationship between polyandry potential and lay date of a female's first nest of the season. Each datum is the lay date of an individual's first nest and their observed local mating behavior of each year. B) Lay date distributions of all nests for females that were polyandrous (yellow; 90 nests from 74 females) or monogamous (green; 571 nests from 402 females). C) Seasonal variation in egg volume – trend shows the between-individual polynomial function of the model prediction. Each datum is an egg's volume (cm^3) and lay date. Late date is standardized for each year across all panels.

Figure 3. Age- and origin-dependent breeding phenology of female snowy plovers. A) Within-individual variation in age-specific nest initiation date – as females gained more experience in the local population, they started nesting earlier, however this trend reversed at older ages (albeit with high uncertainty). Each datum represents an individual's 'age-deviance' (i.e., a within-group centered measure of the number of years since the individual's first observed local breeding attempt, see *Methods* for more details) and the lay date of its first nest each year. B) Origin-specific variation in nest initiation date – females that hatched locally and recruited into the breeding population (orange) tended to nest earlier than birds originating from elsewhere (purple). Inner-most distributions show the model estimates and 95% CI, outer-most box plots show the inter-quartile ranges of the raw data (point-cloud).

 SUPPLEMENTARY TABLES

1012   **Table S1.** Relationship between the average egg volume of a clutch and the average chick
1013   weight at hatching. See Supplementary Text D for methods.

Chick hatch weight	Mean estimate	95% confidence interval
Fixed effects β (logit standardized)		
Average egg volume of clutch	0.60	[0.55, 0.66]
Partitioned R^2		
Total Marginal R^2	0.37	[0.3, 0.43]
Total Conditional R^2	0.47	[0.38, 0.56]
Random effects σ^2		
Mother identity	0.01	[0, 0.02]
Year	0.01	[0, 0.02]
Residual	0.09	[0.08, 0.11]
Adjusted repeatability r		
Mother identity	0.05	[0, 0.12]
Year	0.05	[0.01, 0.11]
Residual	0.90	[0.81, 0.98]
Sample sizes n		
Year	14	
Individuals	426	
Observations (i.e., Nests)	841	

Table S2 *(shown on next page).* Sources of egg volume variation when using only known-
aged females. Note: within the *Partitioned R²* sub-table the term 'Senescence' describes the
collective variation explained by the linear and quadratic within-individual age effects in the
top sub-table, the term 'Selective appearance' and 'Selective disappearance' describe the
variation explained by the between individual first- and last-breeding age fixed effects of the
top sub-table, respectively; the term 'Within ind. seasonality' describes the variation
explained by the within individual lay date effect in the top sub-table, and the term 'Between
ind. seasonality' describes the collective variation explained by the linear and quadratic lay
date effects in the top sub-table.

 Table S2 *(caption on previous page).*

Egg volume (known-age subset)	Mean estimate	95% confidence interval
Fixed effects β (logit standardized)		
Mother tarsus length	0.32	[0.11, 0.5]
Clutch size (< 3 eggs)	-0.02	[-0.22, 0.19]
Within ind. age-deviance (linear)	-0.05	[-0.21, 0.12]
Within ind. age-deviance (quadratic)	-0.10	[-0.24, 0.05]
Between ind. first breeding age	0.03	[-0.31, 0.35]
Between ind. last breeding age	-0.01	[-0.38, 0.37]
Within ind. lay-date-deviance	0.06	[-0.09, 0.2]
Between ind. linear lay date	0.14	[-0.02, 0.31]
Between ind. quadratic lay date	-0.16	[-0.32, 0.01]
Partitioned R^2		
Total Marginal R^2	0.12	[0.06, 0.32]
Total Conditional R^2	0.66	[0.58, 0.77]
Mother tarsus length	0.08	[0.02, 0.28]
Clutch size (< 3 eggs)	0.00	[0, 0.22]
Senescence	0.01	[0, 0.23]
Selective appearance	0.00	[0, 0.22]
Selective disappearance	0.00	[0, 0.21]
Within ind. seasonality	0.04	[0, 0.25]
Between ind. seasonality	0.00	[0, 0.22]
Random effects σ^2		
Nest / Individual	0.05	[0.03, 0.08]
Individual	0.06	[0.02, 0.12]
Year	0.00	[0, 0.02]
Residual	0.07	[0.06, 0.09]
Adjusted repeatability r		
Nest / Individual	0.28	[0.13, 0.43]
Individual	0.33	[0.12, 0.51]
Year	0.01	[0, 0.08]
Residual	0.38	[0.28, 0.51]
Sample sizes n		
Years	14	
Individuals	43	
Nests	112	
Observations (i.e., Eggs)	320	

1024 **Table S3.** Relationship between the lay date of an individual's first nest and their likelihood
1025 of re-nesting or being polyandrous (i.e., "multi-clutching") in a given year. Fixed effect size
1026 of 'first nest lay date' is the standardized estimate on the logit scale.

Multi-clutching probability	Mean estimate	95% confidence interval
Fixed effects β (logit standardized)		
First nest lay date	-1.83	[-2.3, -1.52]
Age since first breeding	-0.04	[-0.26, 0.19]
First breeding age	0.11	[-0.12, 0.33]
Female tarsus length	-0.03	[-0.25, 0.19]
Partitioned R^2		
Total Marginal R^2	0.37	[0.28, 0.47]
Total Conditional R^2	0.40	[0.3, 0.51]
First nest lay date	0.37	[0.28, 0.46]
Age since first breeding	0.00	[0, 0.17]
First breeding age	0.00	[0, 0.17]
Female tarsus length	0.00	[0, 0.17]
Random effects σ^2		
Individual	0.35	[0, 0.86]
Year	0.40	[0, 0.69]
Adjusted repeatability r		
Individual	0.02	[0, 0.12]
Year	0.03	[0, 0.08]
Sample sizes n		
Years	14	
Individuals	426	
Observations (i.e., Nests)	665	

SUPPLEMENTARY FIGURES

Figure S1. Results of simulation to cross-validate the birth year estimates of known-aged
females produced by BaSTA. Panel A) shows a correlation between the estimated BaSTA
birth year and the true birth year (size of points refers to the frequency of females), with the
dashed diagonal illustrating a slope of 1 (i.e., a perfect estimation). Panel B) shows the
distribution of the birth year error for all known-aged females based on the cross-validation
simulation.

Figure S2. (caption on next page).

Figure S2. Mating strategy and clutch number of female snowy plovers according to age. Each row shows an individual female in the population for which we have at least three years of observations (note that our analysis also includes females with one or two years of observation, but given space constraints only individuals with a minimum of three years are plotted in this graph). Panel A) shows known-aged females which were born locally, whereas B) shows females that were initially captured as adults and are therefore of unknown age. Points illustrate the age at which we collected observations of egg volume, with the size of the point corresponding to the number of clutches measured at a given age, and the colour indicating if we observed the female mating with one or two distinct males (i.e., in case of multiple clutches at a given age). The light grey buffer around unknown-age females indicates the 95% CrI of the ages for an individuals' observed period (i.e., lower limit indicates the minimum age the individual could have entered the population and the upper limit indicates the maximum age of an individual's last observation based on BaSTA's birth year posterior). The dark grey lines indicate the period for which an individual was observed alive (i.e., in some cases we encountered an individual in the field and confirmed its survival, but we did not observe its nest. Note also that the age at first encounter of all known-aged individuals is 0).

1050　　**Figure S2.** Homoscedastic spread of residuals from the A) 'egg volume' and B) 'lay date'
1051　　models.

1052   **Figure S3.** Relationship between a female's structural size (i.e., her tarsus length) and A) the
1053   lay date of her first nest of the season, and B) the volume of her eggs.

Figure S4. Visualization of the effect of uncertainty in the age estimate provided by BaSTA on age-dependent trends in egg volume (top row) and lay date (bottom row) dynamics. Black trends show the model predictions for the 1000 bootstraps of the BaSTA age estimate posteriors (see *Methods*). Panels A and D show the within-individual trends of the 'age-deviation' score – as expected, these measures are not impacted by uncertainty in the BaSTA age estimate because they are centered for each individual (i.e., the absolute age is irrelevant). Panels B and E show the between-individual trend of the 'age at first breeding' (i.e., selective appearance), and panels C and F show the between-individual trend of the 'age at last breeding' (i.e., selective disappearance). Yellow trends and grey ribbons visualize the 95% CI of the model predictions when using the median age estimate provided by BaSTA (i.e., the effect sizes of the 'egg volume' and 'lay date' models shown in Fig. 1).

1065   **Figure S5.** The total observed number of chicks produced by females in years when they
1066   were polyandrous was 2.18 (1.83–2.55 95% CI) chicks more than in years when were
1067   monogamous (Fig. S9; $N = 14$ years, 424 females, 836 nests).