Article Type: Original Article

Title: **Egg size variation and breeding phenology in a long-lived sequentially polygamous shorebird**

Short Running Title: Egg size variation in a polygamous shorebird

Authors: Luke J. Eberhart-Hertel[*†1], Lourenço Falcão Rodrigues[*†2,3], Johannes Krietsch[4], Anne G. Hertel[5,6], Medardo Cruz-López[7], Karina Alejandra Vázquez-Rojas[1], Erick González-Medina[8], Julia Schroeder[3], and Clemens Küpper[*1]

Affiliations:

[1]Research Group Behavioural Genetics and Evolutionary Ecology, Max Planck Institute for Ornithology, Seewiesen, Germany

[2]Department of Ecology, Autonomous University of Madrid, Madrid, Spain

[3]Department of Life Sciences, Imperial College London, Silwood Park Campus, United Kingdom

[4]Department of Behavioural Ecology and Evolutionary Genetics, Max Planck Institute for Ornithology, Seewiesen, Germany

[5]Research Group for Movement Ecology, Senckenberg Biodiversity and Climate Research Centre, Frankfurt, Germany

[6]Behavioural Ecology, Department of Biology, Ludwig-Maximilians University of Munich, Planegg-Martinsried, Germany

[7]Institute of Ocean Sciences and Limnology, Universidad Nacional Autónoma de México, Ciudad de Mexico, Mexico

[8]Conservation Biology Research Group, Department of Anatomy, Cell Biology & Zoology, University of Extremadura, Badajoz, Spain

[*]Corresponding authors: luke.eberhart@orn.mpg.de, lourenco.falcao@uam.es, ckuepper@orn.mpg.de,

[†]These authors contributed equally.

AUTHOR CONTRIBUTIONS

LJE-H, CK, and JS: conceived and supervised the study
LJE-H, LF, CK, MC-L, KAV-R, and EG-M: collected the data
LJE-H, LF, and AGH: conducted the analysis and wrote the accompanying RMarkdown document (Supplementary File 1)
LJE-H, LF, JK, AGH, and CK: interpreted the results
LJE-H, LF, and CK: wrote the manuscript
All authors helped to revise the manuscript and approved the final version.

ACKNOWLEDGEMENTS

We dedicate this article to our colleague and friend René Beamonte-Barrientos who tragically passed away in March 2016. René worked on senescence in birds including snowy plovers and inspired us to conduct this study. We thank Salvador Ángel del Gómez, Miguel Guevara-Medina and Ivan Guardado for assistance with fieldwork, and Daniel Galindo-Espinosa for logistic support. We also thank Martin Bulla, Niels Dingemanse, Wolfgang Forstmeier, Bart Kempenaers, James Tolliver, and Martin Stoffel for their constructive discussions and statistical advice. Fieldwork was conducted under permit from SEMARNAT (SGPA/DVGS/2631/19). A full list of funding for fieldwork is provided on www.chorlito.org. LJE-H was funded by the German Science Foundation (DFG Eigene Stelle grant: 415037502). AGE-H was funded by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 793077 and the German Science Foundation (DFG Eigene Stelle grant: 448621093). MCL was supported by a doctoral grant from CONACyT (248125/378124) and the Cornell Lab of Ornithology 'Coastal Solutions Fellowship'. CK was supported by the Max Planck Society. We declare no conflicts of interest.

CONFLICT OF INTEREST

We declare no conflict of interest in regards to our submitted work.

DATA ACCESSIBILITY

We provide all computer code and documentation as a html vignette written in Rmarkdown (Supplementary File 1) together with all the raw datasets needed to reproduce our modeling and analyses – these files can be found in this project's Open Science Framework repository (https://doi.org/10.17605/OSF.IO/UCW6J) and GitHub page (https://leberhartphillips.github.io/snowy_plover_eggs/).

Article Type: Original Article

Title: **Egg size variation and breeding phenology in a long-lived sequentially**
polygamous shorebird

Short Running Title: Egg size variation in a polygamous shorebird

Keywords: Charadrius nivosus, female-female competition, mating system, season,
reproductive investment, snowy plover

Abstract:
Gamete size variation *between* the sexes is central to the concept of sex roles, however, to what
extent gamete size variation *within* sexes relates to sex role variation remains unclear.
Comparative and theoretical studies suggest that, when clutch size is invariable, female
polygamy leads to a reduction of egg size, whereas increased female-female competition for
mates favors early breeding. To understand whether and how breeding phenology, egg size,
and mating behavior are related at the individual level, we studied the reproductive histories of
426 snowy plover females collected over 15 years. Egg size but not mating strategy were highly
repeatable for individual females. Consistent with theoretical predictions, we found that
polyandrous females were the earliest breeders and that early clutches contained smaller eggs
than clutches initiated later, however, the latter effect was only moderate. Neither egg size nor
mating behavior showed clear signs of an age-related deterioration, on the contrary, prior
experience acquired either through age or local recruitment enabled females to nest early.
Taken together, these results suggest that gamete size variation is not linked to female mating
behavior at the individual level.

Teaser Text:
Comparative avian analyses have linked female polyandry and sex-role reversal to the
production of smaller eggs: the notion being that smaller eggs allow polyandrous females to
lay multiple sequential clutches quickly. Our research, however, has found that this process is
more complex at the individual level. We found that females who started breeding early in the
season were more likely to be polyandrous and produce smaller eggs. Experience was also
found to give older and local females a competitive advantage over younger and inexperienced
females vying for early breeding opportunities. Despite this, we found limited evidence that
egg size reduction gives a competitive advantage in the scramble for mates. Instead, seasonal

variations in egg size can be attributed to other factors, such as the survival of early-hatching

chicks, resource limitations during egg production, and individual differences. We found that

senescence had little impact on egg size or mating behavior. Future research should examine

how seasonality, age, and mating behavior impact the reproductive fitness of males, who

provide more parental care than females in many sex role reversed systems. By studying these

effects in concert, we can gain a deeper understanding of sex roles in natural populations.

INTRODUCTION

Anisogamy is a fundamental prerequisite for the concept of sex roles (e.g., Darwin 1871;

Bateman 1948; Wade and Shuster 2002; Schärer et al. 2012; Lehtonen et al. 2016). In species

with conventional sex roles, males, the producers of many small gametes, invest into acquiring

matings to sire as many offspring as possible, whereas females, the producers of few large

gametes, aim to improve the survival prospects of their offspring by obtaining critical resources

and providing parental care. However, there are many species where sex roles do not conform

to this conventional paradigm (reviewed in Janicke et al., 2016). Moreover, there can also be

considerable gamete size variation within the sexes – however, the causes and consequences

of this intrasexual variation remains unclear (Bernardo, 1996; Immler et al., 2011). For

example, the largest eggs produced by some females are typically at least 50% larger than the

smallest eggs produced in the same population (Christians, 2002) and occasionally the largest

egg can be more than twice as heavy as the smallest viable egg (Svagelj & Quintana, 2011).

Such among-individual variation in egg size may have important consequences for individual

fitness and could be linked to different reproductive strategies of females. Egg size is an

important life history trait under maternal control that is positively associated with offspring

fitness (Smith & Fretwell, 1974; Williams, 1994; Christians, 2002; Krist, 2011). Egg size

typically reflects hatchling quality: the amount of nutrients provided to the developing

offspring are especially important during early life (Williams, 1994; Krist, 2011). Because

resources for maternal investment during oogenesis are typically limited, among- and within-

female variation in egg size may signal a trade-off between offspring quality and quantity.

In oviparous organisms, egg size represents a basic measure of maternal investment (Kaplan,

1980; Fox, 1994; Williams, 1994, 2012; Starck & Ricklefs, 1998; Moran & Emlet, 2001; Xu

et al., 2019) and is shown to be related to inter- and intra-specific variation in several life

history traits. For example, egg size is associated with developmental mode, with precocial

species typically producing larger eggs than altricial species (Smith & Fretwell, 1974;

Williams, 1994; Christians, 2002; Krist, 2011). Within species a substantial proportion of the

egg size variation can be explained by consistent individual differences between females which

are highly repeatable across breeding attempts (Christians, 2002). In some cases, intraspecific

variation in egg size is related to among-female variation in reproductive strategies (Slagsvold

et al., 1984) and is occasionally linked to discrete demographic classes such as reproductive

morphs (Beck *et al.*, 2022; Giraldo-Deck *et al.*, 2022).

Given the strong association between egg size variation and life-history traits or maternal

investment it is not surprising that there are strong temporal components related to egg size

variation, such as seasonality and senescence. Reproductive output is often age-dependent

(Bouwhuis et al., 2009; Dingemanse et al., 2020; Hammers et al., 2012; Jankowiak et al., 2018;

Lemaître et al., 2015; Reznick et al., 2004; Riecke et al., 2023; Salguero-Gómez et al., 2016;

Zhang et al., 2015) whereby an individual's performance increases over early life to a

maximum, followed by senescent decline. Senescence is a consequence of age-dependent

trade-offs between energy investments in reproduction at the expense of somatic repair

(Kirkwood & Rose, 1991). Decreasing egg and/or clutch size can indicate reproductive

senescence that reduces female reproductive performance with older ages (Reid *et al.*, 2003;

Beamonte-Barrientos *et al.*, 2010; Dingemanse *et al.*, 2020; Fay *et al.*, 2021). Conversely,

polyandry has been associated to female age, with older more experienced females being more

likely to be polyandrous than younger females (Oring *et al.*, 1994). Such observations of higher

mating rates at older ages suggest that differences in individual quality or experience are related

to female mating behavior (Whittingham & Dunn, 2010), which could in turn be linked to

among- and within-female variation in egg size.

Another source for temporal variation in egg size is seasonality. Egg size often changes

throughout the reproductive season, especially in birds (Dittmann & Hötker, 2001; Skrade &

Dinsmore, 2013; Weiser *et al.*, 2018; Kubelka *et al.*, 2020; Verhoeven *et al.*, 2020). At the

population level, reproductive performance often declines with season which may coincide

with declining egg sizes (Varpe, 2017; Weiser *et al.*, 2018). Several processes may contribute

to this trend. First, in environments with limited or seasonally dependent resources, breeding

early may provide favorable conditions for raising offspring reducing the amount of parental

care required and maximizing offspring survival prospects (Varpe, 2017; Weiser *et al.*, 2018).

Consistent with this, offspring that are born early in the season often have higher survival

prospects than offspring born late in the season (Festa-Bianchet, 1988; Reznick *et al.*, 2006;

Plard *et al.*, 2015; Cruz-López *et al.*, 2017; Varpe, 2017). Second, only high quality and
experienced females may be able to reproduce early in the season because of potential
increased physiological stress for early egg-laying females, and/or the increased likelihood of
brood failure in environments with high stochasticity, such as low food availability, inclement
weather, or frequency-dependent predation risk (Borgmann *et al.*, 2013; Ockendon *et al.*,
2013). Therefore, less experienced or low-quality females may need to postpone their
reproduction to a later date when conditions are more benign. Third, alternation between
extended breeding and non-breeding periods may contribute to differences in egg and,
consequently, offspring quality: when breeding is restricted to a certain period of the year,
females can use the preceding non-reproductive period to amass resources for investment into
egg or clutch size of their initial breeding attempt (Yohannes *et al.*, 2010). As the breeding
season advances, less time is available for successful breeding attempts and as a consequence
females may acquire fewer resources for egg production resulting in fewer and or/smaller eggs
in late clutches (Wallander & Andersson, 2003).

An underappreciated evolutionary process explaining egg size variation within species is
sexual selection acting on females. Although sexual selection is considered to shape
predominantly male-male competition, it is increasingly acknowledged that sexual selection
also operates on females with strong female-female competition for mates (Clutton-Brock,
2007; Kraaijeveld *et al.*, 2007; Fromonteil *et al.*, 2023). In some species, competition over
mating opportunities is stronger in females than in males, leading to a reversal of conventional
sex roles (Oring & Lank, 1982; Rosenqvist, 1990; Andersson, 2004; Goymann *et al.*, 2015;
Fritzsche *et al.*, 2021). Theoretical models have linked intrasexual competition to variation in
gamete size, with more intense competition leading to smaller gametes (Wade & Shuster, 2002;
Lehtonen *et al.*, 2016). Notably, this relationship is not restricted to males, as there is some
support for female-female competition over mates, the hallmark of social polyandry, resulting
in a reduction of egg size. For instance, at the species level, a comparative analysis of birds
showed that a lineage's egg size tends to decrease following the evolution of polyandry (Oring
& Lank, 1982; Rosenqvist, 1990; Andersson, 2004; Goymann *et al.*, 2015; Fritzsche *et al.*,
2021) – supposedly due to the selective advantages that producing smaller eggs has on
minimizing the laying period between matings (Goymann *et al.*, 2015).

Andersson (2004) presented a theoretical quantitative genetics model in which intense female
scramble competition (i.e., whereby females jostle for available males to commence breeding)

shapes breeding phenology and egg size variation in females. According to the model, early
breeding provides substantial advantages as it allows these females enough time to pursue
sequential breeding attempts or to replace a failed first attempt (i.e., "re-nesting"; Morrison *et*
al., 2019). Early breeding should therefore be tightly correlated with female polyandry and,
ultimately enhance the fitness of polyandrous females. Laying smaller eggs can provide
another competitive edge in female-female competition over males: when clutch sizes are
uniform, females that produce small eggs can complete their clutch faster than those that
produce larger eggs and hence gain a competitive edge in the competition for mates especially
when breeding synchrony is high and additional males are limited. A lower boundary for egg
size is provided by reduced offspring viability because smaller eggs have fewer nutrients
(Williams, 1994; Blomqvist *et al.*, 1997; Krist, 2011; Giraldo-Deck *et al.*, 2022). However,
despite its highly persuasive character, the predictions on breeding phenology, egg size, and
their relationship with female mating behavior provided in Andersson's (2004) model remain
to be tested empirically at the individual level.

Among oviparous animals, shorebirds (part of the order Charadriiformes) produce some of the
largest eggs in relation to body mass due to the needs of their precocial and (predominantly)
nidifugous young (Lack, 1968; Rahn *et al.*, 1975). Clutch size is highly uniform within
shorebird species and rarely exceeds four eggs (Rahn *et al.*, 1975) meaning that egg size
represents an adequate proxy for maternal investment. As a clade, shorebirds also exhibit a
disproportionately high prevalence of polyandry (Andersson, 2005; Colwell, 2010; Oring,
1986). Here, we investigate breeding phenology, egg size variation, and their relation with
female mating strategy in snowy plovers (*Charadrius nivosus*) using a 15-year longitudinal
mark-recapture dataset. The snowy plover is a long-lived shorebird (longevity record: 20 years;
Colwell et al., 2017) exhibiting flexible sex-specific parental roles with sequential social
monogamy and polygamy occurring in the same population (Warriner *et al.*, 1986; Eberhart-
Phillips *et al.*, 2017). In parts of the species' range, the breeding season can extend up to six
months allowing for multiple nesting attempts following successful fledging of young or
replacing failed nests due to depredation or flooding (Eberhart-Phillips, 2019; Plaschke *et al.*,
2019). On average, females have higher polygamy potential and more partners per season than
males, who in turn, provide more parental care than females, specifically for boods (Carmona-
Isunza *et al.*, 2017; Cruz-López *et al.*, 2017; Eberhart-Phillips *et al.*, 2017). The reason for the
higher polygamy potential in females is a surplus of males in the adult breeding population
(Stenzel *et al.*, 2011; Carmona-Isunza *et al.*, 2017; Eberhart-Phillips *et al.*, 2017).

Consequently, sequential polyandry, whereby females regularly desert their broods after

hatching to start a new breeding attempt with another male, is commonly observed in snowy

plovers (Warriner *et al.*, 1986). Female desertion for re-mating declines with season in

polyandrous plovers (Amat *et al.*, 1999; Székely *et al.*, 1999; Cruz-López *et al.*, 2017; Kupán

et al., 2021) suggesting that female-female competition for mates should be most pronounced

early in the season.

To disentangle the relationships between gamete size, breeding phenology, and female mating

behavior, we examined the reproductive histories of 426 snowy plover females. First, we

investigated the correlates of re-nesting and female polyandry. We hypothesized that early

nesting females would have higher probabilities of re-nesting or becoming sequentially

polyandrous: early nesting females can rejoin the mating pool faster and therefore have higher

potential to secure an additional breeding attempt compared to later nesting females. Secondly,

we explored the predictors of clutch initiation date for a given female in a season. We

hypothesized that early nesting females would have a competitive advantage over late nesters

– either due to a direct body size advantage or an indirect advantage gained through prior

experience as a previous breeder or local natal recruit. To evaluate this, we explored how body

size, age, and origin (i.e., immigrant or natal recruit) were related to female nesting phenology

and the likelihood of re-nesting or polyandry. Third, we examined the pattern of egg size

variation in relation to season, age, and female body size. We predicted that early-laying

females would produce the smallest eggs compared to later laying females due to the costs

associated with intense female-female competition at the start of the season, i.e., the 'race' to

breed early and maximize re-nesting or polyandry potential (Andersson, 2004). If reproductive

senescence affects egg size variation, we predicted egg size to follow a quadratic relationship

with female age. Alternatively, if female polyandry increases with age, we predicted a negative

linear relationship of egg size with female age.

MATERIALS AND METHODS

Data collection

We studied the reproductive effort and breeding schedules of snowy plovers at Bahía de Ceuta

– an important breeding site located on the coast of Sinaloa, western Mexico (23°54'N,

106°57'W). Details on the study site and population are provided elsewhere (e.g., Cruz-López

et al., 2017; Eberhart-Phillips et al., 2020). In brief, we annually monitored breeding birds from

mid-April until early July and collected mark-recapture data following the methods described

in Székely et al. (2011) . We searched for nests using telescopes and mobile hides to minimize

disturbance. Upon finding a nest, we measured the length and width of each egg to the nearest

tenth of a mm to determine their size (Figs. S1a, b). Using these egg dimensions, we calculated

egg volume (Fig. S1c) following (Hoyt, 1979) as:

Eq. 1 $$egg\ volume = K \times length \times width^2,$$

where K is 0.486, a volume-index constant for snowy plovers determined by (Székely *et al.*,

1994) through the use of an egg volumeter (Hanson, 1954). The modal clutch size of snowy

plovers is three (86.2%) and is the maximum number of eggs we have observed in this

population (Eberhart-Phillips *et al.*, 2020a). We regularly checked incomplete nests until the

clutch was complete and assigned the age of these nests according to the date of the last egg

laid (Plaschke *et al.*, 2019). If the clutch was complete upon discovery and had not been

incubated longer than 10 days (92.5%; $N = 778$ nests), we determined its lay date by floating

the egg and estimating the stage of embryonic development. For successful clutches that had

been incubated for more than 10 days (5.4%, $N = 45$ nests), we back-calculated the laying date

based on the hatching date assuming an incubation period of 25 days (Plaschke *et al.*, 2019).

In the rare case that the nest did not hatch (2.1%, $N = 18$ nests) and we discovered it after day

10 of incubation, we assumed that the nest was 17 days old upon discovery (i.e., the midpoint

between the minimum age of 11 days and the 25-day incubation period).

We identified previously marked nesting adults based on their unique colour ring combination.

We captured unmarked adults on their nests during incubation using funnel traps (Hall &

Cavitt, 2012) and assigned a unique colour ring combination for subsequent recognition. In the

rare circumstance when we were unable to identify parents before hatching, we attempted to

capture parents while they tended the chicks. As snowy plovers only show a small degree of

sexual dimorphism, we determined the sex of all captured plovers in the field through a

combination of plumage characteristics (Argüelles-Tico *et al.*, 2015), time of capture (i.e.

females typically incubate during the day and males during the night; Vincze *et al.*, 2017), and

other behavioral cues (e.g., sex-specific brood care; Kupán et al., 2021). For a subset of adults

(57.5%), we confirmed sex molecularly from DNA extracted from blood samples by PCR

amplification of Z and W specific DNA regions with two sex-typing markers: P2/P8 and Calex-

31 (Griffiths *et al.*, 1998; Küpper *et al.*, 2009b; dos Remedios *et al.*, 2010).

We visited known active nests every four or five days to determine the status of the nest (e.g.,
active, depredated, etc.) until the 20th day after egg laying and thereafter daily until the eggs
hatched or failed. We weighed the chicks shortly after hatching (879 [84.7%] within 24 hours
after hatching, 159 [15.3%] during the second day after hatching) and marked them with an
alphanumeric metal and a single color ring for subsequent identification in the chance that these
individuals would be recruited into the breeding population as adults in future years.

For the years 2006 to 2016 all longitudinal data collected has been compiled as part of the
CeutaOPEN project – an open-access database for individual-based field studies in
evolutionary ecology and conservation biology (Eberhart-Phillips *et al.*, 2020a). We accessed
these data directly from the open source repository (Eberhart-Phillips *et al.*, 2020b) and
supplemented them with data from four additional field seasons: 2017–2020. Please refer to
our RMarkdown vignette that connects to *CeutaOPEN* and reproduces all analytical methods
and results presented below (Supplementary File 1). All of our field activities were performed
and permitted in accordance with the approved ethical guidelines outlined by SEMARNAT
(Secretariat of Environment and Natural Resources, Mexico).

Statistical Analyses
To evaluate the relationship between female breeding behavior, nesting phenology, and egg
size, our investigation implemented three step-wise analyses that: 1) established the factors
related to a female's potential to have multiple breeding attempts either through polyandry or
re-nesting, 2) identified the key female traits linked to lay date, and 3) dissected the elements
explaining within- and among-female variation in egg volume investment. A central trait of
interest in these three analyses was female age, as we needed to acknowledge age-dependent
variation in the repeated measures of individuals over the 15-year period of data collection. We
first present our statistical methods for estimating the age of females that entered the population
as adults, then we describe our models for investigating the interplay between female breeding
behavior, nesting phenology, and egg size.

Statistical estimation of age for females first encountered as adults—To estimate the ages of
unknown individuals in our marked population, we conducted a capture-mark-recapture
analysis using the 'Bayesian Survival Trajectory Analysis' (BaSTA) package in R (v1.9.4,
Colchero et al., 2012), which uses a Bayesian hierarchical framework to fit parametric survival
functions of the marked population while accounting for imperfect detection (see

Supplementary Text A for details). In short, the capture histories of unknown-age individuals
(89.9%, $N = 383$) are cast against the survival function of the known-age population (10.1%,
$N = 43$) to produce posterior age estimates. To acknowledge uncertainty in the BaSTA age
estimate of an unknown age individual: we ran bootstrap simulations for each of the following
models 1000 times, with every iteration randomly drawing a birth year estimate for unknown
aged individuals from their posterior distributions provided by BaSTA. For each simulation,
we evaluated the influence of birth year uncertainty by examining the effect size distribution
of the 1000 bootstraps in relation to the 95% confidence interval for effect sizes of the original
model that used the median birth year estimate from BaSTA. Lastly, we considered a time-
since marking capture-mark-recapture model and determined that the recapture probability of
our dataset was not biased by individual heterogeneity in post-marking processes (see
Supplementary Text B for details).

A key methodological issue for studying age related processes in wild populations is that
stochastic extrinsic mortality reduces the frequency of individuals in older age classes, hence
making it challenging to disentangle between- vs. within-individual age-dependent variation –
a phenomenon known as "selective disappearance" (van de Pol & Verhulst, 2006a; Nussey *et*
al., 2008). Investigations using longitudinal data to test for age-related processes are
particularly suitable for this task, as they can control for the confounding effects of selective
disappearance through repeated measures of individuals as they age (van de Pol & Verhulst,
2006a; Nussey *et al.*, 2008; Dingemanse *et al.*, 2020). We controlled for selective appearance
and disappearance of females differing in the trait of interest by fitting 'first observed age' and
'last observed age' as fixed effects – a method that estimates between-individual age effects
introduced by selective disappearance and appearance (Dingemanse et al., 2020; van de Pol &
Verhulst, 2006a, 2006b). We also modelled within-individual age effects by fitting linear and
quadratic forms of a within-group deviation score for age (henceforth 'age-deviance'),
calculated for individual i at age j as: age_{ij} – [*first observed age*]$_i$ (van de Pol & Verhulst,
2006a; Snijders & Bosker, 2011) – essentially describing the number of years since the bird's
arrival to the population. To ensure that intercepts of our age-dependent models represented
the reproductive performance for the earliest age at reproduction (i.e., age 1 in snowy plovers;
Page et al., 2009), we fitted age as 'age – 1' – otherwise it would represent reproduction at age
0, which is an empirically meaningless estimate. Moreover, we followed common statistical
approaches to investigate age-related processes in birds (e.g., Bouwhuis et al., 2009, 2010;
Dingemanse et al., 2020; Graham et al., 2019; Herborn et al., 2016; Schroeder et al., 2012) by

fitting a quadratic function of age to model age-specific trends. Lastly, in all models tarsus
length was included as a fixed effect proxy of female structural size, and was averaged over an
individual's lifetime of repeated measurements, i.e., our *a priori* expectation was that tarsus
length is static throughout adult life and that any variation in this trait was due to measurement
error.

Modelling seasonal variation in polyandry potential ("Polyandry model")—Our sample for
studying seasonal polyandry dynamics included 426 females for which the identity of their
mates had been verified through observation. We defined observed polyandry as a binomial
variable that scored an individual as being monogamous or polyandrous each year based on
our observations of them having one or multiple breeding partners, respectively (see Fig. S2
for an example of the sampling distribution). Overall, we observed 92 cases of polyandry from
76 females over the 15-year period (annual average incidence of observed polyandry: 11.8%,
range: 0–25.6%). To assess the relationship between the likelihood of polyandry and lay date,
age, and female structural size, we fitted a binomial linear mixed effects model that tested the
likelihood of polyandry predicted by the fixed effects of first lay date, age-deviance (see
above), first observed age, and tarsus length. We included individual ID and year as random
effects.

Modelling seasonal variation in re-nesting potential ("Re-nesting model")—Our sample for
studying seasonal re-nesting dynamics included 177 females for which the fate of their first
nest of the season had been verified as a failure. We defined re-nesting as a binomial variable
that scored an individual as being a re-nester or a single-nester each year based on our
observations of them attempting to re-nest after the loss of their first clutch or not, respectively.
Overall, we observed 64 cases of re-nesting from 55 females over the 15-year period following
a failed attempt (annual average incidence of observed re-nesting: 27%, range: 0–58.3%). To
evaluate re-nesting dynamics, we fitted a binomial linear mixed effects model that tested the
likelihood of re-nesting predicted by the fixed effects of first lay date, age-deviance, first
observed age, and tarsus length. We included individual ID and year as random effects.

Modelling individual variation in lay date ("Lay date model")— We modeled the effect of age
on a female's first lay date using a univariate mixed-effect structure that included age-deviance,
age-deviance-squared, first observed age, and last observed age as fixed covariates, and
individual and year as random intercepts. Furthermore, we included average tarsus length was

as a fixed covariate in addition to recruitment status as a two-level fixed effect that described

whether a breeding female hatched locally ("local recruit") or was first encountered as an adult

of unknown origin ("immigrant"). Our sample for studying lay date dynamics used the same

nest-level sample as the polyandry model above, however, as we were interested in how the

recruitment status of an individual influenced breeding phenology, we excluded data from 2006

as this was the first year when we began individual marking of all birds in the population. This

subset resulted in 568 nests from 376 females. We visualized the distribution of lay dates to

confirm normality and to assess the population-level variance in breeding schedule – an

indication of inter-female breeding asynchrony and the intensity of competition for mates

(Andersson, 2004).

Modelling individual variation in egg volume ("Egg volume model")—Our sample for

studying egg volume dynamics included 2392 eggs from 841 nests belonging to 426 females.

56 (13.1%) females had three or more years of repeated measures (Fig. S2), 83 (19.5%) had

two years of repeated measures, and 286 (67.4%) were measured in a single year. Furthermore,

43 (10.1%) individuals in our sample were marked as hatchlings but later recruited as breeding

adults in subsequent years ("known age"), with the remaining 383 (89.9%) individuals being

initially marked as adults ("unknown age"). We modelled within-individual age effects on egg

volume by fitting a univariate mixed-effect model that included linear and quadratic forms of

age-deviance, and assessed between-individual age effects by additionally incorporating first

observed age and last observed age as fixed covariates. Female tarsus length was also included

as a covariate to control for female body size, and we incorporated a quadratic function of lay

date to assess seasonal variation in egg volume as several shorebird studies report seasonal

increases (Skrade & Dinsmore, 2013; Kwon *et al.*, 2018) or decreases in egg volume (Dittmann

& Hötker, 2001; Skrade & Dinsmore, 2013; Kwon *et al.*, 2018; Kubelka *et al.*, 2020;

Verhoeven *et al.*, 2020). To disentangle within- from between-individual effects of lay date on

egg size, we used the same logic as with age above: first lay dates of all individuals each year

represented the between-individual seasonal effect, whereas the deviation in first lay dates of

an individual relative across all successive breeding attempts for a given season represented

the within-individual seasonal effect. We included random intercepts for nest, individual ID,

and year, and assumed a Gaussian error distribution for egg volume.

Evaluating effect sizes and uncertainty—We used packages "lme4" (Bates *et al.*, 2015), "rptR"

(Stoffel *et al.*, 2017) and "partR2" (Stoffel *et al.*, 2020) in the R version "Bunny-Wunnies Freak

Out" (R Core Team, 2020) to conduct our statistical modelling and assessed homoscedasticity

by visually examining residuals (see Fig. S3). For each of the four mixed-effect models

described above, we evaluated the uncertainty in our parameter estimates by simulating 1000

parametric bootstraps via the "partR2::partR2" function (Stoffel *et al.*, 2020). Likewise, we

derived nest-, individual-, and year-level repeatabilities (i.e., intra-class correlations) by

simulating 1000 parametric bootstraps of the four mixed-effect models using "rptR::rpt". We

report fixed effects as standardized regression coefficients (i.e., beta weights) and repeatability

as the 'adjusted repeatability' – interpreted as the repeatability of a given hierarchical group

after controlling for fixed effects (Nakagawa & Schielzeth, 2010).

RESULTS

Sample summary

The modal clutch size of the 841 clutches was 3 eggs (724 nests, 86.1%; 2-eggs: 103 nests,

12.3%, 1-egg: 14 nests, 1.6%). Average egg length was 3.09 cm (0.10 cm SD, range: 2.72–

3.49 cm, Fig. S1a) and width was 2.24 cm (0.05 cm SD, range: 2.02–2.40 cm, Fig. S1b), which

translated into an average egg volume of 7.59 cm^3 (0.46 cm^3 SD, range: 6.00–8.98 cm^3). The

average egg volume of a clutch strongly predicted the average hatch weight of the subsequent

brood (β [95% CIs]: 0.63 [0.55–0.70]; $R^2_{marginal}$ = 0.370 [0.310–0.436]; Figs. S1c and S4, Table

S1, see Supplementary Text D for methods). Based on BaSTA's estimated birth year, 185 of

the 383 unknown-age females in our sample were first observed nesting at age 1 (48.3%), 120

at age two (31.3%), 72 at age three (18.8%), five at age 4 (1.3%), and one at age 5 (0.3%). Of

the 43 locally hatched females in our sample, 29 first nested at age one (67.4%), six were first

observed nesting at age two (14.0%), two at age 3 (4.7%), three at age 4 (7.0%), three at ages

5, 7, and 8, respectively (7.0%). Female tarsus size was not related to their annual apparent

survival (see Supplementary Text C for details). The average local tenure of all females in the

sample was 1.57 years (2.15 SD) with an average age span of 3.12 years (2.03 SD, median: 3,

range: 1–14 years) and an average of 1.56 years of observed ages per female (1.04 SD, median:

1, range: 1–8 age-specific observations). Females in our sample were typically observed

nesting every consecutive year since their first observation, however, some individuals skipped

years (Fig. S2, average yearly interval between nesting attempts = 1.07, 0.27 SD). On average,

made 1.43 (0.56 SD) nesting attempts per season (median = 1, range 1 to 3). The lifetime

number of chicks produced by females in years when they were polyandrous, was 2.18 (1.83–

2.53 95% CI) chicks more than in years when were monogamous (Fig. S9; *N* = 14 years, 426

females, 841 nests).

Seasonal variation in polyandry and re-nesting potential

A female's probability of being polyandrous in a given season was strongly dependent on the date she laid her first nest, with early breeders being more likely to be polyandrous than late breeders (β [95% CIs]: -2.10 [-2.77, -1.69]; $R^2_{marginal}$ = 0.33 [0.23, 0.45]; Figs. 2a and S5, Table S2). Likewise, a female's likelihood of re-nesting following a failed attempt decreased with advancing first lay date (β: -1.79 [-5.79, -1.31]; $R^2_{marginal}$ = 0.40 [0.27, 0.6]; Figs. 2e and S6, Table S3). We found no effect of female tarsus length on the likelihood of being polyandrous (β: 0.22 [-0.05, 0.54]; $R^2_{marginal}$ = 0.001 [0, 0.18]; Fig. S5, Table S2) or re-nesting (β: -0.09 [-0.47, 0.29]; $R^2_{marginal}$ = 0.001 [0, 0.43]; Fig. S6, Table S3). We found a clear effect of age at first breeding on the likelihood of re-nesting, with females that initiated local breeding at an older age being the most likely to re-nest (β: 0.41 [0.06, 1.7]; $R^2_{marginal}$ = 0.03 [0, 0.44]; Fig. S6, Table S3). This was the same for polyandry, albeit the effect was weaker (β: 0.21 [-0.07, 0.48]; $R^2_{marginal}$ = 0.001 [0, 0.19]; Fig. S5, Table S2). We found no evidence that polyandry or re-nesting potential was related to age since first breeding (i.e., age-deviance; Figs. S5 and S6). The lay date distribution of polyandrous females was bimodal, with peaks in the first and second nests occurring 11.1 days before and 29.1 days after the unimodal seasonal peak for monogamous females (Fig. 2b). Similarly, the lay date distribution of re-nesting females was also bimodal, with peaks in the first and replacement clutches occurring 9.7 days before and 28.4 days after the unimodal seasonal peak for single nesters (Fig. 2d). Females had low repeatability among years in polyandry (adjusted individual cross-year repeatability, r [95% CIs] = 0.01 [0, 0.15]; Fig. S5, Table S2) and re-nesting (r = 0.03 [0, 0.14]; Fig. S6, Table S3). When pooled together to assess multi-clutching behavior (i.e., the probability of simply laying two nests or one regardless of the outcome of the first nest or the identity of the second male), females had higher repeatability than when polyandry or re-nesting were assessed separately (r = 0.35 [0, 0.86]; Fig. S10, Table S7).

Individual variation in lay date

Females showed moderate repeatability in their first lay date among years (r = 0.19 [0.06, 0.32] 95%CI; Fig. 1, Table S4). We found strong support for the effect of origin on first lay date: females that locally hatched and later recruited into the breeding population initiated nests 7.80 days earlier (95% CI: [5.09, 10.50]) on average compared to conspecifics whose origin was unknown (Figs. 1 and 3b). The next strongest fixed effect was within-individual age: young females laid later nests compared to older females with lay date advancing by ~2.17 days per

year until age six (95% CI: [1.41, 2.93]; Fig. 3a), after which the uncertainty in the trend

became unwieldly in the oldest age classes of our sample (Fig. 3a). Notably, female size did

not affect lay date (Fig. 1 and S7a), nor did between-individual effects of first or last age at

breeding (Fig. 1).

Individual variation in egg volume

Overall, our model accounted for 71.2% ([68.2, 74.3] 95%CIs) of variation in egg volume,

with fixed effects explaining 7.9% ([5.1, 12.3] 95%CIs) of this variation (Table S5). Females

were highly repeatable in their egg volumes between clutches: $r = 0.47$ ([0.41, 0.53] 95%CIs;

Fig. 1, Table S5). Furthermore, eggs within the same clutch were moderately repeatable in

volume ($r = 0.18$ [0.14, 0.22]; Fig. 1, Table S5). The strongest fixed effect explaining egg

volume variation was the structural size of the mother (β_{tarsus} [95% CI]: 0.23 [0.15, 0.29]; semi-

partial R^2 of female tarsus = 0.05 [0.02, 0.09]; Figs. 1 and S3b, Table S5): larger females laid

larger eggs than smaller females (model predicted difference: 0.58 cm^3 [0.34, 0.81] 95%CI).

The second strongest effect was the between-individual quadratic season function (Fig. 2c):

eggs were smallest at the start of the season (model prediction: 7.18 cm^3 [6.98, 7.37] 95%CI)

and largest shortly after the middle of the season (model prediction: 7.68 cm^3 [7.60, 7.76]

95%CI). Average egg volume also increased between sequential clutches within individuals

but with smaller magnitude than the population-level trend (β_{within} [95%CI]: 0.10 [0.06, 0.14],

predicted increase of 0.21 cm^3 [0.04, 0.38]). Senescence in egg volume was not supported (β_{age}

[95% CIs]: 0.00 [-0.06, 0.6], β_{age}^2: -0.05 [-0.09, -0.01]; semi-partial R^2 of senescence function

= 0.003 [0, 0.05]; Fig. 1, Table S5). Furthermore, we found no support for selective

(dis)appearance of individuals according to egg volume, as the 95% CIs for first and last

observed ages of reproduction overlapped zero (Fig. 1, Table S5).

To further acknowledge uncertainty in the age-estimates provided by BaSTA, we re-analyzed

the egg volume model using only known-aged females. Notably, this strengthened the effect

sizes of 'mother tarsus length' and the within-individual age effects (i.e., 'linear age' and

'quadratic age'). For example, the 'mother tarsus length' effect (β_{tarsus}) increased to 0.33 [0.09,

0.51 95% CI], the 'linear age' effect (β_{age}) increased to -0.05 [-0.21, 0.11], and the 'quadratic

age' effect (β_{age}^2) increased to -0.10 [-0.24, 0.05] (Table S6). However, given that the sample

size of the known-aged female data set was greatly reduced, this analysis is arguably

underpowered: the 95% confidence intervals bounding the effect sizes increased and, in many

cases, overlapped zero. Furthermore, the bootstrap simulations incorporating the individual

birth-year posteriors estimated from BaSTA confirmed the results of all age-dependent models

(Fig. S8).

DISCUSSION

Increased competition for mating opportunities in species with conventional sex roles is

typically seen as a consequence of anisogamy where the production of many small gametes

allows polygamous males to maximize their reproductive potential (Darwin, 1871; Bateman,

1948; Wade & Shuster, 2002; Schärer *et al.*, 2012; Lehtonen *et al.*, 2016). Applying this

paradigm to females, an increasing competition over mates should therefore be associated with

the evolution of smaller eggs. Andersson (2004) developed a theoretical quantitative genetic

model of female sexual selection by scramble competition over males in an attempt to explain

reduced egg size in polyandrous species. Andersson's (2004) model predicted that scramble

competition for males should lead to egg size reduction and early breeding in sequentially

polyandrous females of species with largely invariable clutch sizes. We tested the predictions

of Andersson's (2004) model empirically at the individual level: examining the links between

breeding phenology, individual mating strategies, and egg size variation in a population of

snowy plovers that contains monogamous and polyandrous females (Cruz-López *et al.*, 2017;

Eberhart-Phillips *et al.*, 2017; Kupán *et al.*, 2021). Consistent with Andersson's (2004)

predictions, we found that female polyandry was associated with early breeding and that early

clutches typically contained smaller eggs than clutches laid later in the season – both between

and within individual females. In our population, early breeding females whose nesting

attempts were successful invariably deserted the brood shortly after hatching (Cruz-López *et*

al., 2017; Kupán *et al.*, 2021) and the resulting collinearity between mating behavior and first

lay date prevented us from explicitly testing whether mating behavior was directly related to

egg size.

Although the statistical associations between breeding phenology, egg size and female mating

behavior are intriguing, we note that their impact on the egg size variation observed in this

population is rather limited. First, despite snowy plovers' unusual mating system with

sequential polyandry, monogamy, and occasionally polygyny occurring in the same population

(Warriner *et al.*, 1986; Eberhart-Phillips *et al.*, 2017), the egg size variation we report is not

more pronounced than in other birds: the proportional difference between the largest and

smallest egg in our population was approximately 50%, which is of very similar magnitude to

the intraspecific range observed within other bird populations (Christians, 2002). Other

constraints may provide limitations on egg size variation in snowy plovers. Waders produce
relatively large eggs that provide vital reserves for their precocial young and the moderate egg
size variation observed in snowy plovers could indicate that females are unable to reduce egg
size further without jeopardizing chick viability. Second, we note that structural size of females
had a stronger effect on egg size variation than breeding phenology. This is consistent with
other studies (Christians, 2002; Giraldo-Deck *et al.*, 2022) showing that larger females produce
larger eggs than smaller females. Third, we found that, as demonstrated in other studies (e.g.,
Christians, 2002), female identity explained the largest part of the variation in egg size and
repeatability of egg size in females was very high. The high repeatability suggests that egg size
may have considerable genetic variation, although it has been suggested that many loci with
small effects seem to encode this life history trait (Santure *et al.*, 2013).

In contrast to egg size, female mating behavior showed a very low individual repeatability
across years. The observed low individual consistency is most likely related to the high
stochastic nature of nest survival where the main sources of clutch/brood failure in our
population are flooding, predation, and chick mortality through starvation (Cruz-López *et al.*,
2017). Early breeding females whose nests failed, typically stayed together with their mate for
rapid renesting. These females may have intended to become polygamous but since their
replacement clutch hatched later in the season when conditions for chicks or remating are less
favorable, they switched to a monogamous strategy that involves longer brood care. Snowy
plover females can adjust their brood care efforts flexibly according to the value and needs of
the young (Kupán *et al.*, 2021). Female care and re-mating efforts are inversely related to each
other and show strong seasonality. Early in the breeding season when survival prospects for
the chicks and re-mating opportunities are high, and their mate can successfully rear the chicks
alone, females typically desert their broods right after hatching of the chicks to quickly start a
new breeding attempt with another male (Székely *et al.*, 1999; Cruz-López *et al.*, 2017; Kupán
et al., 2021). Females whose chicks hatch later in the season, typically remain with the brood
and provide care jointly with their mate. However, desertion during brood care is tightly
associated with chick mortality (Kupán *et al.*, 2021). When survival prospects of their offspring
are deteriorating and chicks die despite the joint care, initially caring females desert their
broods and presumably remate elsewhere where environmental conditions are better. Such high
plasticity in female care and mating strategies allows females to flexibly respond to nest and
chick survival, and is an important ecological modulator of breeding behavior in snowy
plovers.

Most studies conducted on temperate or high latitude breeding shorebirds have found a negative association between time of the season and egg size (Byrkjedal & Kalas, 1985; Sandercock *et al.*, 1999; Kubelka *et al.*, 2020) although in polyandrous red-necked phalaropes (*Phalaropus lobatus*) egg size increased throughout the breeding season (Kwon et al., 2018, n.b., the presented effect size is small). However, many investigations of seasonal egg size dynamics did not disentangle whether the observed changes were due to within- or between-individual effects. A study that included 15 arctic shorebirds suggested that indeed between-individual variation may account for more of the seasonal variation in egg size than within-individual variation (Weiser *et al.*, 2018). In our population, early season nesters produced smaller eggs on average than females initiating nests at a later date. These seasonal changes were driven by within individual changes: females generally increased egg volume between consecutive nesting attempts (albeit the effect size was small), suggesting that maternal investment during early breeding attempts is impaired by physiological limitations. Mutually non-exclusive constraints may include low food availability (Steigerwald *et al.*, 2015), and carry-over costs associated with intense competition (Duckworth, 2006) or migratory status (Crossin *et al.*, 2010). In contrast, delayed breeders can take advantage of a slower pace to gather more resources and maximize investment prior to clutch initiation – whilst having only enough time for a single breeding attempt. Although we did not collect any direct empirical data on temporal changes in food availability during our study (e.g., with pit-fall traps), the invertebrate prey communities of salina ecosystems (i.e., like that of our study population) are well documented to have dynamic emergence and quiescent phenologies (Britton & Johnson, 1987), providing important ecological context and support of our notion that resource availability for egg-laying female snowy plovers is highly. The observed decrease of egg size in clutches produced at the end of season is consistent with such constraints for breeding females as females are in a rush to complete their clutches and finish incubation before the onset of catastrophic floods caused by increased precipitation and tide height at the end of the breeding season (Plaschke *et al.*, 2019).

Despite being long-lived – the oldest documented female in our sample was at least 13 years old (longevity record: 20 years; Colwell et al., 2017) – and investing substantially in reproduction year-after-year, we found no evidence of age-dependent trade-offs in egg size or polyandry potential. Contrary to our predictions derived from senescence theory, older females initiated nests earlier in the season compared to their younger conspecifics – indicating age-

dependent competitive ability that may initially increase the polygamy potential of females
(Oring *et al.*, 1994). Age-dependent variation in lay date followed a non-linear pattern: lay date
advanced with each year of age until a peak at age six, after which the age-lay date trajectory
was unclear because of limited sampling in older age classes. In snowy plovers, occasionally
females re-mate with or retain their last mate in the following season (Halimubieke *et al.*,
2019), presumably to skip the time consuming mate search and start breeding early – a process
that could also explain the age-dependent advance in lay date we observed. Furthermore,
locally recruited females (i.e., hatched locally) bred earlier than immigrant females indicating
a role for prior experience giving competitive advantage for breeders in our population.

Egg size did not change clearly with female age. Several studies of oviparous organisms have
observed age-dependent variation in egg size, with some studies finding a positive relationship
(Cooch *et al.*, 1992; Flint & Sedinger, 1992; Robertson *et al.*, 1994; Warner *et al.*, 2016;
Verhoeven *et al.*, 2020) and others observing a negative relationship (Reid, 1988; Potti, 1993;
Ito, 1997). Recent longitudinal studies of wild avian systems showed a quadratic relationship
with a prominent increase in egg size in early life, followed by a peak and then a late-life
decline (Bouwhuis *et al.*, 2009; Jankowiak *et al.*, 2018). In captive avian systems, the early life
increase in egg size is also observed, but not the late-life decline likely due to ad libitum food
(Woodard & Abplanalp, 1971). A quadratic age relationship provided the best fit for a possible
relationship between egg size and age in our study, but the effect was not statistically clear.
This finding did not change when we repeated the analysis including only the 45 females of
known age (10% of the original sample). We acknowledge that our relatively small sample of
known-age individuals presents a limitation to our study – we cannot exclude a moderate effect
of senescence on egg size. Nevertheless, one potential explanation for the lack of a clear
senescence trend observed in egg size is that there is little room for further reduction because
of viability constraints for the chicks. In shorebirds, chicks are not fed by the parents but rather
must forage for themselves immediately after hatching and larger chicks that hatch from larger
eggs typically survive better than smaller chicks (Blomqvist *et al.*, 1997; Giraldo-Deck *et al.*,
2022). The extra nutrients provided by a large egg may be crucial for the chick survival during
the first few days of life when foraging efficiency is reduced as chicks still learn how to catch
prey (Ricklefs, 1968). Consequently, additional reductions of egg size may not be feasible as
the knock-on effects for chick survival are too costly (Williams, 1994; Starck & Ricklefs,
1998), therefore limiting a females' potential to save resources by reducing egg volume. The

chick viability constraints may also provide an explanation why the variation in egg size at the
population level is relatively low.

Conclusions
Past species-level studies have linked polyandry and sex-role reversal to reduced female
gamete size (Slotow, 1996; Andersson, 2004), as smaller eggs would permit females to produce
several clutches in rapid succession (Liker *et al.*, 2001). In our individual-level investigation
we found that early nesting females were more likely to be polyandrous and produced clutches
containing smaller eggs than females that started their first breeding attempt later in the season.
Starting to breed early in the season is an important pre-determinant of female polygamy. Prior
experience can provide both older and local females a competitive advantage over younger and
naïve conspecifics scrambling for early breeding opportunities that are at the heart of the
polyandrous mating strategy (Andersson, 2004). However, evidence supporting an adaptive
egg-size adjustment to obtain a competitive advantage in the scramble competition for mates
was weak. Rather we found that the potential for egg size reduction is limited in snowy plovers
and the observed seasonal variation in egg size can also be explained by other constraints that
include a survival advantage for early hatching chicks, resource constraints during the egg-
producing phase and, most importantly, between individual differences. Senescence did not
seem to have a major impact on egg size variation nor on the mating behavior of females.
Future research of polyandrous systems should also examine the impact of seasonality, age,
and mating behavior on the reproductive fitness of males, as males provide a larger share of
parental care than females and therefore should have a more direct impact on the survival
prospects of the young. Examining the effects of season- and age-dependent trade-offs on
individual fitness and variation in mating tactics will enable a deeper understanding about
selective advantages and constraints of diverse and plastic sex roles in natural populations.

DATA ACCESSIBILITY

We provide all computer code and documentation as a html vignette written in Rmarkdown

(click here: File S1) together with all the raw datasets needed to reproduce our modeling and

analyses – these files can be found in this project's Open Science Framework repository:

doi.org/ 10.17605/OSF.IO/UCW6J

LITERATURE CITED

Amat, J.A., Fraga, R.M. & Arroyo, G.M. 1999. Brood desertion and polygamous breeding in the Kentish plover Charadrius alexandrinus. *Ibis* **141**: 596–607.

Andersson, M. 2005. Evolution of Classical Polyandry: Three Steps to Female Emancipation. *Ethology* **111**: 1–23.

Andersson, M. 2004. Social polyandry, parental Investment, sexual selection, and evolution of reduced female gamete size. *Evolution* **58**: 24–34. [Society for the Study of Evolution, Wiley].

Argüelles-Tico, A., Küpper, C., Kelsh, R.N., Kosztolányi, A., Székely, T. & van Dijk, R.E. 2015. Geographic variation in breeding system and environment predicts melanin-based plumage ornamentation of male and female Kentish plovers. *Behavioral Ecology and Sociobiology* **70**: 49–60.

Bateman, A.J. 1948. Intra-sexual selection in Drosophila. *Heredity* **2**: 349–368. Essex.

Bates, D., Mächler, M., Bolker, B. & Walker, S. 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* **67**.

Beamonte-Barrientos, R., Velando, A., Drummond, H. & Torres, R. 2010. Senescence of Maternal Effects: Aging Influences Egg Quality and Rearing Capacities of a Long-Lived Bird. *The American Naturalist* **175**: 469–480.

Beck, S.V., Räsänen, K., Kristjánsson, B.K., Skúlason, S., Jónsson, Z.O., Tsinganis, M., *et al.* 2022. Variation in egg size and offspring phenotype among and within seven Arctic charr morphs. *Ecology and Evolution* **12**: e9427.

Bernardo, J. 1996. The Particular Maternal Effect of Propagule Size, Especially Egg Size: Patterns, Models, Quality of Evidence and Interpretations1. *American Zoologist* **36**: 216–236.

Blomqvist, D., Johansson, O.C. & Götmark, F. 1997. Parental quality and egg size affect chick survival in a precocial bird, the lapwing Vanellus vanellus. *Oecologia* **110**: 18–24.

Borgmann, K.L., Conway, C.J. & Morrison, M.L. 2013. Breeding phenology of birds: mechanisms underlying seasonal declines in the risk of nest predation. *PLoS ONE* **8**: e65909.

Bouwhuis, S., Charmantier, A., Verhulst, S. & Sheldon, B.C. 2010. Trans-generational effects on ageing in a wild bird population. *Journal of Evolutionary Biology* **23**: 636–642.

Bouwhuis, S., Sheldon, B.C., Verhulst, S. & Charmantier, A. 2009. Great tits growing old: selective disappearance and the partitioning of senescence to stages within the breeding cycle. *Proceedings of the Royal Society B: Biological Sciences* **276**: 2769–2777.

Britton, R.H. & Johnson, A.R. 1987. An ecological account of a Mediterranean salina: The
Salin de Giraud, Camargue (S. France). *Biological Conservation* **42**: 185–230.

Byrkjedal, I. & Kalas, J.A. 1985. Seasonal variation in egg size in golden plover Pluvialis
apricaria and dotterel Charadrius morinellus populations. *Ornis Scandinavica* **16**:
108–112. WileyNordic Society Oikos.

Carmona-Isunza, M.C., Ancona, S., Székely, T., Ramallo-González, A.P., Cruz-López, M.,
Serrano-Meneses, M.A., *et al.* 2017. Adult sex ratio and operational sex ratio exhibit
different temporal dynamics in the wild. *Behavioral Ecology* **28**: 523–532.

Christians, J.K. 2002. Avian egg size: variation within species and inflexibility within
individuals. *Biological reviews* **77**: 1–26. Cambridge University Press.

Clutton-Brock, T. 2007. Sexual Selection in Males and Females. *Science* **318**: 1882–1885.
American Association for the Advancement of Science.

Colchero, F., Jones, O.R. & Rebke, M. 2012. BaSTA: an R package for Bayesian estimation
of age-specific survival from incomplete mark–recapture/recovery data with
covariates. *Methods in Ecology and Evolution* **3**: 466–470.

Colwell, M.A. 2010. *Shorebird Ecology, Conservation, and Management*. University of
California Press.

Colwell, M.A., Feucht, E.J., McAllister, S.E. & Transou, A.N. 2017. Lessons learned from
the oldest Snowy Plover. *Wader Study* **124**: 157–159.

Cooch, E.G., Lank, D.B., Rockwell, R.F. & Cooke, F. 1992. Is There a Positive Relationship
between Body Size and Fecundity in Lesser Snow Geese? *The Auk* **109**: 667–673.

Crossin, G.T., Trathan, P.N., Phillips, R.A., Dawson, A., Le Bouard, F. & Williams, T.D.
2010. A carryover effect of migration underlies individual variation in reproductive
readiness and extreme egg size dimorphism in macaroni penguins. *The American
Naturalist* **176**: 357–366.

Cruz-López, M., Eberhart-Phillips, L.J., Fernández, G., Beamonte-Barrientos, R., Székely,
T., Serrano-Meneses, M.A., *et al.* 2017. The plight of a plover: Viability of an
important snowy plover population with flexible brood care in Mexico. *Biological
Conservation* **209**: 440–448.

Darwin, C. 1871. *The descent of man and selection in relation to sex*. Murray, London.

Dingemanse, N.J., Moiron, M., Araya-Ajoy, Y.G., Mouchet, A. & Abbey-Lee, R.N. 2020.
Individual variation in age-dependent reproduction: Fast explorers live fast but
senesce young? *Journal of Animal Ecology* **89**: 601–613.

Dittmann, T. & Hötker, H. 2001. Intraspecific Variation in the Egg Size of the Pied Avocet.
Waterbirds: The International Journal of Waterbird Biology **24**: 83–88.

dos Remedios, N., Lee, P.L., Székely, T., Dawson, D.A. & Küpper, C. 2010. Molecular sex-
typing in shorebirds: a review of an essential method for research in evolution,
ecology and conservation. *Wader Study Group Bulletin* **117**: 109–118.

Duckworth, R.A. 2006. Aggressive behavior affects selection on morphology by influencing
settlement patterns in a passerine bird. *Proceedings of the Royal Society B: Biological
Sciences* **273**: 1789–1795.

Eberhart-Phillips, L.J. 2019. Plover Breeding Systems. In: *The Population Ecology and
Conservation of Charadrius Plovers* (M. A. Colwell & S. M. Haig, eds), pp. 65–88.
New York.

Eberhart-Phillips, L.J., Cruz-López, M., Lozano-Angulo, L., del Ángel, S.G., Rojas-Abreu,
W., Bucio-Pacheco, M., *et al.* 2020a. CeutaOPEN, individual-based field observations
of breeding snowy plovers Charadrius nivosus. *Sci Data* **7**: 149.

Eberhart-Phillips, L.J., Cruz-López, M., Lozano-Angulo, L., Gómez del Ángel, S., Rojas-
Abreu, W., Bucio-Pacheco, M., *et al.* 2020b. CeutaOPEN v1.5. Open Science
Framework.

Eberhart-Phillips, L.J., Küpper, C., Miller, T.E.X., Cruz-López, M., Maher, K.H., Dos
Remedios, N., *et al.* 2017. Sex-specific early survival drives adult sex ratio bias in
snowy plovers and impacts mating system and population growth. *Proceedings of the
National Academy of Sciences of the United States of America* **114**: e5474–e5481.

Fay, R., Ravussin, P.-A., Arrigo, D., von Rönn, J.A.C. & Schaub, M. 2021. Age-specific
reproduction in female pied flycatchers: evidence for asynchronous aging. *Oecologia*
196: 723–734.

Festa-Bianchet, M. 1988. Birthdate and survival in bighorn lambs (Ovis canadensis). *Journal
of Zoology* **214**: 653–661.

Flint, P.L. & Sedinger, J.S. 1992. Reproductive Implications of Egg-Size Variation in the
Black Brant. *The Auk* **109**: 896–903.

Fox, C.W. 1994. The Influence of Egg Size on Offspring Performance in the Seed Beetle,
Callosobruchus maculatus. *Oikos* **71**: 321–325.

Fritzsche, K., Henshaw, J.M., Johnson, B.D. & Jones, A.G. 2021. The 150th anniversary of
The Descent of Man: Darwin and the impact of sex-role reversal on sexual selection
research. *Biological Journal of the Linnean Society* **134**: 525–540.

Fromonteil, S., Marie-Orleach, L., Winkler, L. & Janicke, T. 2023. Sexual selection in
females and the evolution of polyandry. *PLOS Biology* **21**: e3001916. Public Library
of Science.

Giraldo-Deck, L.M., Loveland, J.L., Goymann, W., Tschirren, B., Burke, T., Kempenaers,
B., *et al.* 2022. Intralocus conflicts associated with a supergene. *Nat Commun* **13**:
1384. Nature Publishing Group.

Goymann, W., Makomba, M., Urasa, F. & Schwabl, I. 2015. Social monogamy vs.
polyandry: ecological factors associated with sex roles in two closely related birds
within the same habitat. *Journal of Evolutionary Biology* **28**: 1335–1353.

Graham, J.L., Bauer, C.M., Heidinger, B.J., Ketterson, E.D. & Greives, T.J. 2019. Early-
breeding females experience greater telomere loss. *Molecular Ecology* **28**: 114–126.

Griffiths, R., DOUBLE, M.C., ORR, K. & DAWSON, R.J.G. 1998. A DNA test to sex most
birds. *Molecular Ecology* **7**: 1071–1075.

Halimubieke, N., Valdebenito, J.O., Harding, P., Cruz-López, M., Serrano-Meneses, M.A.,
James, R., *et al.* 2019. Mate fidelity in a polygamous shorebird, the snowy plover
(Charadrius nivosus). *Ecology and Evolution* **9**: 10734–10745.

Hall, L.K. & Cavitt, J.F. 2012. Comparative Study of Trapping Methods for Ground-Nesting
Shorebirds. *Waterbirds* **35**: 342–346.

Hammers, M., Richardson, D.S., Burke, T. & Komdeur, J. 2012. Age-dependent terminal
declines in reproductive output in a wild bird. *PLoS ONE* **7**: e40413.

Hanson, H.C. 1954. Apparatus for the Study of Incubated Bird Eggs. *The Journal of Wildlife*
Management **18**: 191–198.

Herborn, K.A., Daunt, F., Heidinger, B.J., Wilding, H.M.V.G., Burthe, S.J., Newell, M.A., *et*
al. 2016. Age, oxidative stress exposure and fitness in a long-lived seabird.
Functional Ecology **30**: 913–921.

Hoyt, D.F. 1979. Practical Methods of Estimating Volume and Fresh Weight of Bird Eggs.
The Auk **96**: 73–77.

Immler, S., Pitnick, S., Parker, G.A., Durrant, K.L., Lüpold, S., Calhim, S., *et al.* 2011.
Resolving variation in the reproductive tradeoff between sperm size and number.
Proceedings of the National Academy of Sciences **108**: 5325–5330. National Acad
Sciences.

Ito, K. 1997. Egg-size and -number variations related to maternal size and age, and the
relationship between egg size and larval characteristics in an annual marine
gastropod, Haloa japonica (Opisthobranchia; Cephalaspidea). *Marine Ecology*
Progress Series **152**: 187–195.

Janicke, T., Häderer, I.K., Lajeunesse, M.J. & Anthes, N. 2016. Darwinian sex roles
confirmed across the animal kingdom. *Science Advances* **2**: e1500983. American
Association for the Advancement of Science.

Jankowiak, \Lukasz, Zyskowski, D. & Wysocki, D. 2018. Age-specific reproduction and
disposable soma in an urban population of Common Blackbirds Turdus merula. *Ibis*
160: 130–144.

Kaplan, R.H. 1980. The Implications of Ovum Size Variability for Offspring Fitness and
Clutch Size within Several Populations of Salamanders (Ambystoma). *Evolution* **34**:
51–64.

Kirkwood, T.B.L. & Rose, M.R. 1991. Evolution of senescence: late survival sacrificed for
reproduction. *Philosophical Transactions of the Royal Society B-Biological Sciences*
332: 15–24.

Kraaijeveld, K., Kraaijeveld-Smit, F.J.L. & Komdeur, J. 2007. The evolution of mutual
ornamentation. *Animal Behavior* **74**: 657–677.

Krist, M. 2011. Egg size and offspring quality: a meta-analysis in birds. *Biological Reviews*
86: 692–716.

Kubelka, V., Sládeček, M., Zámečník, V., Vozabulová, E. & Šálek, M. 2020. Seasonality
Predicts Egg Size Better Than Nesting Habitat in a Precocial Shorebird. *Ardea* **107**:
239.

Kupán, K., Székely, T., Cruz-López, M., Seymour, K. & Küpper, C. 2021. Offspring
desertion with care? Chick mortality and plastic female desertion in Snowy Plovers.
Behavioral Ecology.

Küpper, C., Augustin, J., Kosztolányi, A., Burke, T., Flguerola, J. & Székely, T. 2009a.
Kentish Versus Snowy Plover: Phenotypic and Genetic Analyses of Charadrius
alexandrinus Reveal Divergence of Eurasian and American Subspecies. *The Auk* **126**:
839–852.

Küpper, C., Augustin, J., Kosztolányi, A., Burke, T., Flguerola, J. & Székely, T. 2009b.
Kentish Versus Snowy Plover: Phenotypic and Genetic Analyses of Charadrius
alexandrinus Reveal Divergence of Eurasian and American Subspecies. *The Auk* **126**:
839–852.

Kwon, E., English, W.B., Weiser, E.L., Franks, S.E., Hodkinson, D.J., Lank, D.B., *et al.*
2018. Delayed egg-laying and shortened incubation duration of Arctic-breeding
shorebirds coincide with climate cooling. *Ecology and Evolution* **8**: 1339–1351.

Lack, D. 1968. *Ecological adaptations for breeding in birds*. Methuen Ltd, London.

Lehtonen, J., Parker, G.A. & Schärer, L. 2016. Why anisogamy drives ancestral sex roles.
Evolution **70**: 1129–1135.

Lemaître, J.-F., Berger, V., Bonenfant, C., Douhard, M., Gamelon, M., Plard, F., *et al.* 2015.
Early-late life trade-offs and the evolution of ageing in the wild. *Proceedings of the
Royal Society B: Biological Sciences* **282**: 20150209.

Liker, A., Reynolds, J.D. & Székely, T. 2001. The evolution of egg size in socially
polyandrous shorebirds. *Oikos* **95**: 3–14.

Moran, A.L. & Emlet, R.B. 2001. Offspring size and performance in variable environments:
field studies on a marine snail. *Ecology* **82**: 1597–1612.

Morrison, C.A., Alves, J.A., Gunnarsson, T.G., Þórisson, B. & Gill, J.A. 2019. Why do
earlier-arriving migratory birds have better breeding success? *Ecology and Evolution*
9: 8856–8864.

Nakagawa, S. & Schielzeth, H. 2010. Repeatability for Gaussian and non-Gaussian data: a
practical guide for biologists. *Biological Reviews* **85**: 935–956.

Nosály, G. & Székely, T. 1993. Clutch and egg-size variation in the Kentish Plover
(\emphCharadrius alexandrinus) during the breeding season. *Aquila* **100**: 161–179.

Nussey, D.H., Coulson, T., Bianchet, M.F. & Gaillard, J.M. 2008. Measuring senescence in
wild animal populations: towards a longitudinal approach. *Functional Ecology* **22**:
393–406.

Ockendon, N., Leech, D. & Pearce-Higgins, J.W. 2013. Climatic effects on breeding grounds
are more important drivers of breeding phenology in migrant birds than carry-over
effects from wintering grounds. *Biology Letters* **9**: 20130669.

Oring, L.W. 1986. Avian Polyandry. In: *Current Ornithology, vol 3* (R. F. Johnston, ed), pp.
309–351.

Oring, L.W. & Lank, D.B. 1982. Sexual selection, arrival times, philopatry and site fidelity in
the polyandrous spotted sandpiper. *Behavioral Ecology and Sociobiology* **10**: 185–
191.

Oring, L.W., Reed, M.J. & Alberico, A.J. 1994. Mate acquisition tactics in polyandrous
spotted sandpipers (Actitis macularia): the role of age and experience. *Behavioral*
Ecology **5**: 9–16.

Page, G.W., Stenzel, L.E., Warriner, J.S., Warriner, J.C. & Paton, P.W. 2009. Snowy Plover
(Charadrius nivosus).

Plard, F., Gaillard, J.-M., Coulson, T., Hewison, A.J.M., Douhard, M., Klein, F., *et al.* 2015.
The influence of birth date via body mass on individual fitness in a long-lived
mammal. *Ecology* **96**: 1516–1528.

Plaschke, S., Bulla, M., Cruz-López, M., Gómez del Ángel, S. & Küpper, C. 2019. Nest
initiation and flooding in response to season and semi-lunar spring tides in a ground-
nesting shorebird. *Frontiers in Zoology* **16**: 15–11.

Potti, J. 1993. Environmental, ontogenetic, and genetic variation in egg size of Pied
Flycatchers. *Canadian Journal of Zoology-revue Canadienne De Zoologie - CAN J*
ZOOL **71**: 1534–1542.

R Core Team. 2013. R: A language and environment for statistical computing. Vienna.

R Core Team. 2020. *R: A Language and Environment for Statistical Computing*. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Rahn, H., Paganelli, V.C. & Ar, A. 1975. Relation of Avian Egg Weight to Body Weight.
The Auk **92**: 750–765.

Reid, J.M., Bignal, E.M., Bignal, S., McCracken, D.I. & Monaghan, P. 2003. Age-specific
reproductive performance in red-billed choughs Pyrrhocorax pyrrhocorax: patterns
and processes in a natural population. *Journal of Animal Ecology* **72**: 765–776.

Reid, V.W. 1988. Age-Specific Patterns of Reproduction in the Glaucous-Winged Gull: Increased Effort with Age. *Ecology* **69**: 1454–1465.

Reznick, D., Schultz, E., Morey, S. & Roff, D. 2006. On the virtue of being the first born: the influence of date of birth on fitness in the mosquitofish, Gambusia affinis. *Oikos* **114**: 135–147.

Reznick, D.N., Bryant, M.J., Roff, D., Ghalambor, C.K. & Ghalambor, D.E. 2004. Effect of extrinsic mortality on the evolution of senescence in guppies. *Nature* **431**: 1095–1099. Nature Publishing Group.

Ricklefs, R.E. 1968. Patterns of growth in birds. *Ibis* **110**: 419–451.

Riecke, T.V., Hegelbach, J. & Schaub, M. n.d. Reproductive senescence and mating tactic interact and conflict to drive reproductive success in a passerine. *Journal of Animal Ecology* **n/a**.

Robertson, G.J., Cooch, E.G., Lank, D.B., Rockwell, R.F. & Cooke, F. 1994. Female Age and Egg Size in the Lesser Snow Goose. *Journal of Avian Biology* **25**: 149.

Rosenqvist, G. 1990. Male mate choice and female-female competition for mates in the pipefish Nerophis ophidion. *Animal Behavior* **39**: 1110–1115.

Salguero-Gómez, R., Jones, O.R., Jongejans, E., Blomberg, S.P., Hodgson, D.J., Mbeau-Ache, C., *et al.* 2016. Fast-slow continuum and reproductive strategies structure plant life-history variation worldwide. *Proceedings of the National Academy of Sciences of the United States of America* **113**: 230–235.

Sandercock, B.K., Lank, D.B. & Cooke, F. 1999. Seasonal Declines in the Fecundity of Arctic-Breeding Sandpipers: Different Tactics in Two Species with an Invariant Clutch Size. *Journal of Avian Biology* **30**: 460. JSTOR.

Santure, A.W., De Cauwer, I., Robinson, M.R., Poissant, J., Sheldon, B.C. & Slate, J. 2013. Genomic dissection of variation in clutch size and egg mass in a wild great tit (Parus major) population. *Molecular Ecology* **22**: 3949–3962.

Schärer, L., Rowe, L. & Arnqvist, G. 2012. Anisogamy, chance and the evolution of sex roles. *Trends in Ecology & Evolution* **27**: 260–264.

Schroeder, J., Burke, T., Mannarelli, M.E., Dawson, D.A. & Nakagawa, S. 2012. Maternal effects and heritability of annual productivity. *Journal of Evolutionary Biology* **25**: 149–156.

Skrade, P. & Dinsmore, S.J. 2013. Egg-size investment in a bird with uniparental incubation by both sexes. *The Condor* **115**: 508–514.

Slagsvold, T., Sandvik, J., Rofstad, G., Lorentsen, Ö. & Husby, M. 1984. On the Adaptive Value of Intraclutch Egg-Size Variation in Birds. *The Auk* **101**: 685–697.

Slotow, R. 1996. Black coucal Centropus grillii egg volume predicts their polyandrous mating system. *Journal of Avian Biology* 171–173.

862 Smith, C.C. & Fretwell, S.D. 1974. The Optimal Balance between Size and Number of
863   Offspring. *The American Naturalist* **108**: 499–506. The University of Chicago Press.

864 Snijders, T. & Bosker, R.J. 2011. \emphMultilevel analysis: An introduction to basic and
865   *advanced multilevel modeling*, 2nd ed. Sage.

866 Starck, J.M. & Ricklefs, R.E. 1998. *Avian growth and development. Evolution within the
867   altricial-precocial spectrum*. New York.

868 Steigerwald, E.C., Igual, J.-M., Payo-Payo, A. & Tavecchia, G. 2015. Effects of decreased
869   anthropogenic food availability on an opportunistic gull: evidence for a size-mediated
870   response in breeding females. *Ibis* **157**: 439–448.

871 Stenzel, L.E., Page, G.W., Warriner, J.C., Warriner, J.S., Neuman, K.K., George, D.E., *et al.*
872   2011. Male-skewed adult sex ratio, survival, mating opportunity and annual
873   productivity in the snowy plover \emphCharadrius alexandrinus. *Ibis* **153**: 312–322.

874 Stoffel, M.A., Nakagawa, S. & Schielzeth, H. 2020. partR2: Partitioning R2 in generalized
875   linear mixed models. *bioRxiv* 2020.07.26.221168.

876 Stoffel, M.A., Nakagawa, S. & Schielzeth, H. 2017. rptR: repeatability estimation and
877   variance decomposition by generalized linear mixed-effects models. *Methods in
878   Ecology and Evolution* **8**: 1639–1644.

879 Svagelj, W.S. & Quintana, F. 2011. Egg-size variation in the Imperial Cormorant: on the
880   importance of individual effects. *The condor* **113**: 528–537. Oxford University Press.

881 Székely, T., Argüelles-Tico, A., Kosztolányi, A. & Küpper, C. 2011. Practical guide for
882   investigating breeding ecology of Kentish plover. 1–16.

883 Székely, T., Cuthill, I.C. & Kis, J. 1999. Brood desertion in Kentish plover sex differences in
884   remating opportunities. *Behavioral Ecology* **10**: 185–190.

885 Székely, T., Kosztolányi, A. & Küpper, C. 2008. *Practical guide for investigating breeding
886   ecology of Kentish plover Charadrius alexandrinus*. University of Bath, Bath, UK.

887 Székely, T., Kozma, J. & Piti, A. 1994. The Volume of Snowy Plover Eggs (El Volumen de
888   los Huevos de Charadrius alexandrinus). *Journal of Field Ornithology* **65**: 60–64.

889 van de Pol, M. & Verhulst, S. 2006a. Age-Dependent Traits: A New Statistical Model to
890   Separate Within- and Between-Individual Effects. *The American Naturalist* **167**: 766–
891   773.

892 van de Pol, M. & Verhulst, S. 2006b. Age-Dependent Traits: A New Statistical Model to
893   Separate Within- and Between-Individual Effects. *The American Naturalist* **167**: 766–
894   773. The University of Chicago Press.

895 Varpe, Ø. 2017. Life History Adaptations to Seasonality. *Integrative and Comparative
896   Biology* **57**: 943–960.

Verhoeven, M.A., Loonstra, A.H.J., McBride, A.D., Tinbergen, J.M., Kentie, R.,
Hooijmeijer, J.C.E.W., *et al.* 2020. Variation in Egg Size of Black-Tailed Godwits.
Ardea **107**: 291.

Vincze, O., Kosztolányi, A., Barta, Z., Küpper, C., Alrashidi, M., Amat, J.A., *et al.* 2017.
Parental cooperation in a changing climate: fluctuating environments predict shifts in
care division. *Global Ecology and Biogeography* **26**: 347–358.

Wade, M.J. & Shuster, S.M. 2002. The evolution of parental care in the context of sexual
selection: a critical reassessment of parental investment theory. *The American*
Naturalist **160**: 285–292. The University of Chicago Press.

Wallander, J. & Andersson, M. 2003. Reproductive tactics of the ringed plover Charadrius
hiaticula. *Journal of Avian Biology* **34**: 259–266.

Warner, D.A., Miller, D.A.W., Bronikowski, A.M. & Janzen, F.J. 2016. Decades of field data
reveal that turtles senesce in the wild. *Proceedings of the National Academy of*
Sciences of the United States of America **113**: 6502–6507.

Warriner, J.S., Warriner, J.C., Page, G.W. & Stenzel, L.E. 1986. Mating System and
Reproductive Success of a Small Population of Polygamous Snowy Plovers. *The*
Wilson Bulletin **98**: 15–37. Wilson Ornithological Society.

Weiser, E.L., Brown, S.C., Lanctot, R.B., Gates, H.R., Abraham, K.F., Bentzen, R.L., *et al.*
2018. Life-history tradeoffs revealed by seasonal declines in reproductive traits of
Arctic-breeding shorebirds. *Journal of Avian Biology* **49**: jav-01531.

Whittingham, L.A. & Dunn, P.O. 2010. Fitness benefits of polyandry for experienced
females. *Molecular Ecology* **19**: 2328–2335.

Williams, T.D. 1994. Intraespecific Variation in Egg Size and Egg Composition in Birds:
Effects on Offspring Fitness. *Biological Reviews of the Cambridge Philosophical*
Society **69**: 35–59.

Williams, T.D. 2012. *Physiological adaptations for breeding in birds*. Princenton.

Woodard, A.E. & Abplanalp, H. 1971. Longevity and Reproduction in Japanese Quail
Maintained Under Stimulatory Lighting. *Poultry Science* **50**: 688–692.

Xu, F., Yang, W. & Li, Y. 2019. Enlarged Egg Size Increases Offspring Fitness of a Frog
Species on the Zhoushan Archipelago of China. *Scientific Reports* **9**: 11653.

Yohannes, E., Valcu, M., Lee, R.W. & Kempenaers, B. 2010. Resource use for reproduction
depends on spring arrival time and wintering area in an arctic breeding shorebird.
Journal of Avian Biology **41**: 580–590.

Zhang, H., Vedder, O., Becker, P.H. & Bouwhuis, S. 2015. Age-dependent trait variation: the
relative contribution of within-individual change, selective appearance and
disappearance in a long-lived seabird. *Journal of Animal Ecology* **84**: 797–807.

Figure 1. Sources of egg volume (left) and lay date (right) variation. Top row: standardized effect sizes (± 95% CI) of fixed effects. Middle row: variance explained by fixed effects (note: the term 'Senescence' describes the collective variation explained by the linear and quadratic within-individual age effects in the top row; the term 'Selective appearance' and 'Selective disappearance' describe the variation explained by the between individual first- and last-breeding age fixed effects of the top row, respectively; the term 'Within ind. seasonality' describes the variation explained by the within individual lay date effect in the top row, and the term 'Between ind. seasonality' describes the collective variation explained by the linear and quadratic lay date effects in the top row). Bottom row: adjusted repeatabilities (± 95% CI) of random effects.

Figure 2. Phenology of mating behaviors, re-nesting, egg laying, and egg size in 426 female snowy plovers breeding at Bahía de Ceuta. A) Relationship between polyandry potential and lay date of a female's first nest of the season. Each datum is the lay date of an individual's first nest and their observed local mating behavior of each year. B) Lay date distributions of all nests for females that were polyandrous (yellow; 92 nests from 76 females) or monogamous (green; 573 nests from 403 females). C) Seasonal variation in egg volume – trend shows the between-individual polynomial function of the model prediction. Each datum is an egg's

volume (cm^3) and lay date. D) Lay date distributions of all nests for females that laid
replacement clutches (orange; 64 nests from 127 females) or did not re-nest (purple; 146 nests
from 134 females) following failure of their first nest. E) Relationship between re-nesting
potential and lay date of a female's first nest of the season. As with panel (A), each datum is
the lay date of an individual's first nest and their observed local re-nesting activity of each year.
Late date is standardized for each year across all panels.

Figure 3. Age- and origin-dependent breeding phenology of female snowy plovers. A) Within-
individual variation in age-specific nest initiation date – as females gained more experience in
the local population, they started nesting earlier, however this trend reversed at older ages
(albeit with high uncertainty). Each datum represents an individual's 'age-deviance' (i.e., a
within-group centred measure of the number of years since the individual's first observed local
breeding attempt, see *Methods* for more details) and the lay date of its first nest each year. B)
Origin-specific variation in nest initiation date – females that hatched locally and recruited into
the breeding population (orange) tended to nest earlier than birds originating from elsewhere
(purple). Inner-most distributions show the model estimates and 95% CI, outer-most box plots
show the inter-quartile ranges of the raw data (point-cloud).

 SUPPLEMENTARY TABLES

 Table S1. Relationship between the average egg volume of a clutch and the average chick weight at hatching. See Supplementary Text D for methods.

Chick hatch weight	Mean estimate	95% confidence interval
Fixed effects β (standardized)		
Average egg volume of clutch	0.60	[0.55, 0.66]
Partitioned R^2		
Total Marginal R^2	0.37	[0.3, 0.43]
Total Conditional R^2	0.47	[0.38, 0.56]
Random effects σ^2		
Mother identity	0.01	[0, 0.02]
Year	0.01	[0, 0.02]
Residual	0.09	[0.08, 0.11]
Adjusted repeatability r		
Mother identity	0.05	[0, 0.12]
Year	0.05	[0.01, 0.11]
Residual	0.90	[0.81, 0.98]
Sample sizes n		
Year	14	
Individuals	425	
Observations (i.e., Nests)	840	

972 **Table S2.** Relationship between the lay date of an individual's first nest and their likelihood of
973 polyandry each year. Fixed effect size of 'first nest lay date' is the standardized estimate on the
974 logit scale.

Polyandry probability	Mean estimate	95% confidence interval
Fixed effects β (logit standardized)		
First nest lay date	−2.10	[-2.77, -1.69]
Age since first breeding	−0.01	[-0.29, 0.25]
First breeding age	0.21	[-0.07, 0.48]
Mother tarsus length	0.22	[-0.05, 0.54]
Partitioned R^2		
Total Marginal R^2	0.34	[0.24, 0.45]
First nest lay date	0.33	[0.23, 0.45]
Age since first breeding	0.00	[0, 0.18]
First breeding age	0.00	[0, 0.19]
Mother tarsus length	0.00	[0, 0.18]
Total Conditional R^2	0.36	[0.25, 0.52]
Random effects σ^2		
Individual	0.08	[0, 1.55]
Year	0.11	[0, 0.39]
Adjusted repeatability r		
Individual	0.01	[0, 0.15]
Year	0.01	[0, 0.05]
Sample sizes n		
Years	14	
Individuals	426	
Observations (i.e., Nests)	665	

975 **Table S3.** Relationship between the lay date of an individual's first nest and their likelihood of
976 re-nesting following breeding failure in a given year. Fixed effect size of 'first nest lay date' is
977 the standardized estimate on the logit scale.

Re-nesting probability	Mean estimate	95% confidence interval
Fixed effects β (logit standardized)		
First nest lay date	−1.79	[−5.79, −1.31]
Age since first breeding	0.20	[−0.19, 0.73]
First breeding age	0.41	[0.06, 1.7]
Mother tarsus length	−0.09	[−0.47, 0.29]
Partitioned R^2		
Total Marginal R^2	0.42	[0.29, 0.61]
First nest lay date	0.40	[0.27, 0.6]
Age since first breeding	0.01	[0, 0.43]
First breeding age	0.03	[0, 0.44]
Mother tarsus length	0.00	[0, 0.43]
Total Conditional R^2	0.44	[0.3, 0.96]
Random effects σ^2		
Individual	0.15	[0, 1153.43]
Year	0.00	[0, 2.8]
Adjusted repeatability r		
Individual	0.03	[0, 14.04]
Year	0.00	[0, 0.21]
Sample sizes n		
Years	13	
Individuals	177	
Observations (i.e., Nests)	209	

Table S4. Sources of lay date variation. Note: within the *Partitioned R²* sub-table the term 'Senescence' describes the collective variation explained by the linear and quadratic within-individual age effects in the top sub-table, the term 'Selective appearance' and 'Selective disappearance' describe the variation explained by the between individual first- and last-breeding age fixed effects of the top sub-table, respectively.

Lay date of first nest	Mean estimate	95% confidence interval
Fixed effects β (standardized)		
Mother tarsus length	0.02	[-0.07, 0.1]
Local recruit	−0.41	[-0.68, -0.15]
Within ind. linear age	−0.26	[-0.41, -0.11]
Within ind. quadratic age	0.04	[0.01, 0.08]
Between ind. first breeding age	0.06	[-0.04, 0.17]
Between ind. last breeding age	−0.04	[-0.19, 0.11]
Partitioned R^2		
Total Marginal R^2	0.06	[0.04, 0.12]
Total Conditional R^2	0.23	[0.13, 0.38]
Mother tarsus length	0.00	[0, 0.06]
Local recruit	0.02	[0, 0.08]
Senescence	0.02	[0, 0.07]
Selective appearance	0.01	[0, 0.06]
Selective disappearance	0.00	[0, 0.06]
Random effects σ^2		
Individual	60.69	[18.94, 117.67]
Year	2.00	[0, 12.89]
Residual	283.38	[229.95, 326.82]
Adjusted repeatability r		
Individual	0.19	[0.06, 0.32]
Year	0.01	[0, 0.03]
Residual	0.80	[0.67, 0.93]
Sample sizes n		
Years	13	
Individuals	376	
Observations (i.e., Nests)	568	

Table S5. (*displayed on next page*) Sources of egg size variation. Note: within the *Partitioned*
R^2 sub-table the term 'Senescence' describes the collective variation explained by the linear
and quadratic within-individual age effects in the top sub-table, the term 'Selective appearance'
and 'Selective disappearance' describe the variation explained by the between individual first-
and last-breeding age fixed effects of the top sub-table, respectively; the term 'Within ind.
seasonality' describes the variation explained by the within individual lay date effect in the top
sub-table, and the term 'Between ind. seasonality' describes the collective variation explained
by the linear and quadratic lay date effects in the top sub-table.

 Table S5. (*caption on previous page*)

Egg volume	Mean estimate	95% confidence interval
Fixed effects β (standardized)		
Mother tarsus length	0.23	[0.15, 0.29]
Within ind. linear age	−0.00	[-0.06, 0.06]
Within ind. quadratic age	−0.05	[-0.09, -0.01]
Between ind. first breeding age	0.04	[-0.04, 0.14]
Between ind. last breeding age	0.02	[-0.1, 0.15]
Within ind. lay date	0.10	[0.06, 0.14]
Between ind. linear lay date	0.16	[0.11, 0.21]
Between ind. quadratic lay date	−0.07	[-0.12, -0.01]
Partitioned R^2		
Total Marginal R^2	0.08	[0.05, 0.12]
Total Conditional R^2	0.71	[0.68, 0.74]
Mother tarsus length	0.05	[0.02, 0.09]
Senescence	0.00	[0, 0.05]
Selective appearance	0.00	[0, 0.05]
Selective disappearance	0.00	[0, 0.05]
Between ind. seasonality	0.01	[0, 0.06]
Within ind. seasonality	0.01	[0, 0.05]
Random effects σ^2		
Nest / Individual	0.04	[0.03, 0.04]
Individual	0.10	[0.08, 0.12]
Year	0.01	[0, 0.02]
Residual	0.06	[0.06, 0.07]
Adjusted repeatability r		
Nest / Individual	0.18	[0.14, 0.22]
Individual	0.47	[0.41, 0.53]
Year	0.04	[0.01, 0.08]
Residual	0.31	[0.28, 0.35]
Sample sizes n		
Years	14	
Individuals	426	
Nests	841	
Observations (i.e., Eggs)	2392	

992   **Table S6.** Sources of egg volume variation when using only known-aged females. See
993   caption of Table S1 for definitions of table terms.

Egg volume (known-age analysis)	Mean estimate	95% confidence interval
Fixed effects β (standardized)		
Mother tarsus length	0.33	[0.09, 0.51]
Within ind. linear age	−0.05	[-0.21, 0.11]
Within ind. quadratic age	−0.10	[-0.24, 0.05]
Between ind. first breeding age	0.05	[-0.32, 0.38]
Between ind. last breeding age	−0.03	[-0.41, 0.36]
Within ind. lay date	0.09	[-0.03, 0.22]
Between ind. linear lay date	0.14	[-0.02, 0.3]
Between ind. quadratic lay date	−0.17	[-0.34, 0]
Partitioned R^2		
Total Marginal R^2	0.13	[0.06, 0.32]
Total Conditional R^2	0.74	[0.67, 0.83]
Mother tarsus length	0.09	[0.02, 0.29]
Senescence	0.01	[0, 0.23]
Selective appearance	0.00	[0, 0.22]
Selective disappearance	0.00	[0, 0.22]
Between ind. seasonality	0.04	[0, 0.25]
Within ind. seasonality	0.01	[0, 0.22]
Random effects σ^2		
Nest / Individual	0.05	[0.03, 0.08]
Individual	0.07	[0.03, 0.14]
Residual	0.05	[0.04, 0.06]
Adjusted repeatability r		
Nest / Individual	0.29	[0.17, 0.46]
Individual	0.41	[0.19, 0.58]
Residual	0.30	[0.21, 0.41]
Sample sizes n		
Years	14	
Individuals	43	
Nests	112	
Observations (i.e., Eggs)	319	

994　**Table S7.** Relationship between the lay date of an individual's first nest and their likelihood
995　of re-nesting or being polyandrous (i.e., "multi-clutching") in a given year. Fixed effect size
996　of 'first nest lay date' is the standardized estimate on the logit scale.
997

multiclutching probability	Mean estimate	95% confidence interval
Fixed effects β (logit standardized)		
First nest lay date	−1.83	[-2.3, -1.52]
Age since first breeding	−0.04	[-0.26, 0.19]
First breeding age	0.11	[-0.12, 0.33]
Female tarsus length	−0.03	[-0.25, 0.19]
Partitioned R^2		
Total Marginal R^2	0.37	[0.28, 0.47]
First nest lay date	0.37	[0.28, 0.46]
Age since first breeding	0.00	[0, 0.17]
First breeding age	0.00	[0, 0.17]
Female tarsus length	0.00	[0, 0.17]
Total Conditional R^2	0.40	[0.3, 0.51]
Random effects σ^2		
Individual	0.35	[0, 0.86]
Year	0.40	[0, 0.69]
Adjusted repeatability r		
Individual	0.02	[0, 0.12]
Year	0.03	[0, 0.08]
Sample sizes n		
Years	14	
Individuals	426	
Observations (i.e., Nests)	665	

SUPPLEMENTARY FIGURES

Figure S1. Egg size variation for snowy plovers breeding in Bahía de Ceuta, Mexico between 2006 and 2020. Egg volume was estimated using an allometric equation combining length (A) and width (B) measurements collected in the field (see *Methods*). C) Average egg volume of a clutch was highly correlated with the clutch's average chick weight at hatch (β [95% CIs]: 0.628 [0.552–0.704]; $R^2_{marginal}$ = 0.370 [0.310–0.436]). Ribbon shows the 95% CI of the model prediction, vertical and horizontal error bars visualize within clutch variation in egg volume and chick hatch weight (1 SD); see Supplementary Text D for modelling methods.

Figure S2. Mating strategy and clutch number of female snowy plovers according to age. Each row shows an individual female in the population for which we have at least three years of

observations (note that our analysis also includes females with one or two years of observation, but given space constraints only individuals with a minimum of three years are plotted in this graph). Panel A) shows known-aged females which were born locally, whereas B) shows females that were initially captured as adults and are therefore of unknown age. Points illustrate the age at which we collected observations of egg volume, with the size of the point corresponding to the number of clutches measured at a given age, and the colour indicating if we observed the female mating with one or two distinct males (i.e., in case of multiple clutches at a given age). The light grey buffer around unknown-age females indicates the 95% CrI of the ages for an individuals' observed period (i.e., lower limit indicates the minimum age the individual could have entered the population and the upper limit indicates the maximum age of an individual's last observation based on BaSTA's birth year posterior). The dark grey lines indicate the period for which an individual was observed alive (i.e., in some cases we encountered an individual in the field and confirmed its survival, but we did not observe its nest. Note also that the age at first encounter of all known-aged individuals is 0).

1021   **Figure S3.** Homoscedastic spread of residuals from the A) 'egg volume' and B) 'lay date'
1022   models.

chick model

1023 **Figure S4.** Sources of variation in the average hatch weight of chicks.

Polyandry model

A

Fixed effects

First nest lay date

First breeding age

Age since first breeding

Mother tarsus length

Standardized effect size (logit β) ± 95% CI

B

Semi-partial R^2

Marginal R^2

First nest lay date

First breeding age

Age since first breeding

Mother tarsus length

Variance explained (R^2) ± 95% CI

C

Intra-class correlation

Individual

Year

Adjusted repeatability (r) ± 95% CI

1024   **Figure S5.** Sources of variation in a female's probability of being polyandrous each year.

Re-nesting model

 Figure S6. Sources of variation in a female's probability of re-nesting after a failed attempt
 each year.

1027 **Figure S7.** Relationship between a female's structural size (i.e., her tarsus length) and A) the
1028 lay date of her first nest of the season, and B) the volume of her eggs.

 Figure S8. Visualization of the effect of uncertainty in the age estimate provided by BaSTA
on age-dependent trends in egg volume (top row) and lay date (bottom row) dynamics. Black
trends show the model predictions for the 1000 bootstraps of the BaSTA age estimate posteriors
(see *Methods*). Panels A and D show the within-individual trends of the 'age-deviation' score
– as expected, these measures are not impacted by uncertainty in the BaSTA age estimate
because they are centred for each individual (i.e., the absolute age is irrelevant). Panels B and
E show the between-individual trend of the 'age at first breeding' (i.e., selective appearance),
and panels C and F show the between-individual trend of the 'age at last breeding' (i.e.,
selective disappearance). Yellow trends and grey ribbons visualize the 95% CI of the model
predictions when using the median age estimate provided by BaSTA (i.e., the effect sizes of
the 'egg volume' and 'lay date' models shown in Fig. 1).
1040

Figure S9. The lifetime number of chicks produced by females in years when they were polyandrous was 2.18 (1.83–2.53 95% CI) chicks more than in years when were monogamous (Fig. S9; $N = 14$ years, 426 females, 841 nests).

Multiclutching model

SUPPLEMENTARY TEXTS

Supplementary Text A

Age estimation of individuals with unknown origin—Investigating age-dependent processes in the wild is challenging, as analyses often involve a mix of individuals of known or unknown age (Colchero *et al.*, 2012) – with the former being initially marked at birth (i.e., uncensored), and the latter being immigrants of unknown age or those that were born before the study's first marking occasion (i.e., left-truncated). To estimate the ages of unknown individuals in our marked population, we employed a capture-mark-recapture analysis using the 'Bayesian Survival Trajectory Analysis' (BaSTA) package in R (v1.9.4, Colchero *et al.*, 2012), which uses a Bayesian hierarchical framework to fit parametric survival functions of the marked population while accounting for imperfect detection. Furthermore, BaSTA derives estimates of birth year of left-truncated individuals from the population mean of the projected survival function. As snowy plovers show sex differences in survival (Eberhart-Phillips *et al.*, 2017; 2018), we used female-specific survival functions for this study. Due to high natal dispersal, we could not confidently determine the fate of juveniles marked in our population. To acknowledge this uncertainty, our capture-mark-recapture sample only included individuals that survived their first breeding season, i.e., we constrained first-year survival probability to 1.

In total, our capture-mark-recapture data comprised records of 450 uniquely marked females, of which 45 hatched locally and subsequently recruited into the adult population as known-age individuals (Fig. S2a), and the remaining 405 females were adults of unknown age and origin. We monitored the presence or absence of marked individuals by recapturing or observing them in the field in all study years except for 2014, amounting to a total of 916 post-birth annual detections of the 450 females in our sample (median annual detections per adult = 2; mean = 2.04, 1.45 SD). We discarded annual detections if a bird was only encountered once via a resighting of a color combination, as single-observations are known to influence capture-mark-recapture analyses (Tucker et al. 2019). A logistic bathtub-shaped mortality model had the best fit to our data – revealing that female mortality rate increased until age 5 years, after which it became constant (Fig. S2b; see Appendix S1 for detailed methods). Using this model, we extracted the birth year estimate posteriors for each unknown-age individual in the capture-mark-recapture sample. Note that three individuals (one first encountered as an adult [CA1579] and two locally hatched recruits [CA2036 and CA1526]; Fig. S3) had been already marked two

years prior to the start of our monitoring period (i.e., pre-2006) and were thus added to our

sample after running BaSTA on capture-mark-recapture data for 2006–2020.

To explain how mortality risk is associated with age, we compared four commonly used

survival functions: exponential, Gompertz, logistic, and Weibull models. The exponential

function keeps survival constant across age (Cox & Oakes, 1984), whereas the other functions

allow for age-dependent variation in survival: the Gompertz model is an exponential function

in which age-specific mortality is scaled by baseline mortality (Gompertz, 1812; Pletcher,

1999) and the Weibull model is a power function which assumes that baseline and age-specific

mortality rates are independent (Pinder *et al.*, 1978). In addition to the stand-alone versions of

these functions, we considered two alternative forms of the Gompertz, logistic, and Weibull

functions – "Makeham" and "bathtub" shapes. The Makeham shape constrains the survival

function to converge to a constant, rather than zero, as age increases (Pletcher, 1999), whereas

the bathtub shape enables concavity in the function, such that mortality could decrease at early

ages, but increase later in life (Siler, 1979).

We used four parallel simulations to run the Markov chain Monte Carlo (MCMC) optimization

procedure in BaSTA with 800,000 iterations, a 100,000 burn-in period, and a thinning interval

of 2000 to minimize serial autocorrelation in the chain (see Fig. S2 for simulation diagnostics).

We ranked the survival models according to their deviance information criterion (*DIC*) and

determined that the logistic model with a bathtub shape fitted our data best (i.e., lowest *DIC*,

(Spiegelhalter *et al.*, 2002); Table S1). This logistic model revealed that female mortality rate

increased until age 5 years, after which it became constant (Fig. 1b). Using the top model, we

extracted the point estimate and 95% credible interval birth year for each individual in the

capture-mark-recapture sample.

Literature Cited:

Colchero, F., Jones, O.R. & Rebke, M. (2012) BaSTA: An R package for Bayesian

estimation of age-specific survival from incomplete mark-recapture/recovery data with

covariates. *Methods in Ecology and Evolution*, **3**, 466–470.

Cox, D.R. & Oakes, D. (1984) *Analysis of Survival Data*. Chapman and Hall, London.

Eberhart-Phillips, L.J., Küpper, C., Carmona-Isunza, M.C., Vincze, O., Zefania, S., Cruz-

López, M., *et al.* (2018) Demographic causes of adult sex ratio variation and their

consequences for parental cooperation. *Nature Communications*, **9**, 1–8.

Eberhart-Phillips, L.J., Küpper, C., Miller, T.E.X., Cruz-López, M., Maher, K.H., Remedios,

dos, N., *et al.* (2017) Sex-specific early survival drives adult sex ratio bias in snowy

plovers and impacts mating system and population growth. *Proceedings of the National*

Academy of Sciences of the United States of America, **114**, e5474–e5481.

Gompertz, B. (1812) On the nature of the function expressive of the law of human mortality,

and on a new mode of determining the value of life contingencies. *Philosophical*

Transactions of the Royal Society of London B: Biological Sciences, **115**, 513–583.

Pinder, J.E., Wiener, J.G. & Smith, M.H. (1978) The Weibull distribution: a new method of

summarizing survivorship data. *Ecology*, **59**, 175–179.

Pletcher, S. (1999) Model fitting and hypothesis testing for age-specific mortality data.

Journal of Evolutionary Biology, **12**, 430–439.

Siler, W. (1979) A Competing-Risk Model for Animal Mortality. *Ecology*, **60**, 750–757.

Spiegelhalter, D.J., Best, N.G., Carlin, B.P. & Van Der Linde, A. (2002) Bayesian measures

of model complexity and fit. *Journal of the Royal Statistical Society: Series B*

(Statistical Methodology), **64**, 583–639.

Tucker, A.M., McGowan, C.P., Robinson, R.A., Clark, J.A., Lyons, J.E., DeRose-Wilson, A.,

Du Feu, R., Austin, G.E., Atkinson, P.W. & Clark, N.A., Effects of individual

misidentification on estimates of survival in long-term mark–resight studies, *The*

Condor, **1211**, duy017

Table S-A1. Hierarchical capture-mark-recapture models used to describe mortality patterns
of snowy plovers at Bahía de Ceuta, Mexico, between 2006 and 2020. *k:* number of modelled
parameters; Δ*DIC:* difference in deviance information criterion (*DIC*) between a given model
and the top model. *DIC* value of top model was 5279.1.

Mortality function	Shape	k	ΔDIC
Logistic	Bathtub	7	0.00
Logistic	Simple	4	90.91
Logistic	Makeham	5	97.01
Weibull	Bathtub	6	139.36
Weibull	Simple	3	211.98
Weibull	Makeham	4	226.01
Gompertz	Bathtub	6	328.12
Gompertz	Simple	3	401.06
Gompertz	Make	4	445.12
Exponential	Simple	2	679.94

Figure S-A1. Logistic-bathtub mortality function for snowy plover females: a) frequency
distribution of age-specific observations of 45 known-aged females and b) age-dependent
mortality hazard.

Figure S-A2. Bayesian model diagnostics of the Logistic bathtub shaped survival model. a)
Serial autocorrelation of iterations within chains, b) chain convergence (*R-hat*) of model, and
c) trace plots of chain dynamics.

Supplementary Text B

Considering the effect of individual heterogeneity in post-marking processes on mark-

recapture models of survival

Time-since-marking models control for losses to mortality or permanent emigration that occur

during the interval after the first capture occasion (Pradel *et al.* 1997). To code a time-since-

marking variable into the design matrix of the mark-recapture model, a time-varying covariate

needs to be added that distinguishes the first occasion from all subsequent occasions.

Unfortunately, time-varying covariates are unable to be implemented in the latest version of

BaSTA, so we evaluated a time-since-marking Cormack–Jolly–Seber (CJS) model with the R

package "RMark" (Laake 2013) and demographic parameters were estimated via maximum

likelihood implemented in program MARK (White & Burnham 1999). We limited this analysis

to unknown-aged individuals (i.e., only included individuals that were first captured as adults)

because the first-year survival probability for individuals first marked as juveniles is 100% in

our dataset because our dataset is limited to adults that have breeding information.

First, we evaluated the best model capturing variation in recapture probability while keeping

apparent survival constant. In addition to the time-since-marking model, we tested an intercept

only model, and a factorial time (i.e., year) model. This revealed that the factorial time model

was far superior to the other models tested based on AICc statistics (Table S-B1). Furthermore,

a comparison between the time-since-marking model and the intercept-only model shows that

the DeltaAICc scores differ by <2 units – providing additional support that the time-since-

marking model does not improve model fit. We interpret this as encouraging evidence that our

inference about recapture probability is not hindered by individual heterogeneity in post-

marking processes.

Second, we evaluated the if there was evidence of transience in our mark-recapture dataset that

could bias our ability to appropriately estimate survival probability. In RMark we tested a time-

since-marking survival model that incorporated annual variation in encounter probability

structure shown above (i.e., Phi(~tsm)p(~time), Table S-B1). The results of this analysis

revealed no difference in the survival probability of unknown-aged adults transitioning from

their first or between subsequent capture occasions (Figure S-B1). Importantly, the estimated

survival probability of the first occasion is actually larger than that of subsequent occasions

(confidence intervals aside), indicating that there is little evidence that transient adults are

biasing our mark-recapture analyses.

Table S-B1. AIC comparisons for three CJS models evaluating sources of variation in
encounter probability. A time-since-marking model did not improve model fit over an
intercept-only model, suggesting that our inference about recapture probability based on the
BaSTA analyses (i.e., Supplementary Text A) are not hindered by individual heterogeneity in
post-marking processes**.**

model	MARK model	npar	DeltaAICc	weight
Year	Phi(~1)p(~time)	15	0.00	1.00
constant	Phi(~1)p(~1)	2	35.39	0.00
time-since-marking	Phi(~1)p(~tsm)	3	36.86	0.00

130   **Figure S-B1.** Results of a time-since-marking survival model that incorporated annual
131   variation in encounter probability revealed no evidence that transient adults are biasing our
132   mark-recapture-based inference.
133
Literature cited:
JL Laake *RMark: An R Interface for Analysis of Capture-Recapture Data with MARK. AFSC*
Processed Rep 2013-01 (NOAA, Alaska Fisheries Science Center, National Marine
Fisheries Service, Seattle, 2013).
Pradel, R., Hines, J.E., Lebreton, J.D. and Nichols, J.D., 1997. Capture-recapture survival
models taking account of transients. *Biometrics*, pp.60-72.

GC White, KP Burnham, Program MARK: Survival estimation from populations of marked

animals. *Bird Study* **46**, S120–S139 (1999).

Supplementary Text C

Considering the effect of body size on female survival

To evaluate if individual quality predicts survival, we included tarsus length as a covariate in

the RMark mark-recapture model (as described in Supplementary Text B above, individual

covariates are not able to be incorporated into the latest version of BaSTA). Our analysis

showed that there was no effect of tarsus length on survival probability (Figure S-C1). An

intercept-only model had > 2 DeltaAICc units than the tarsus model (Table S-C1) and the

graphical relationship between tarsus and survival unconvincing (Figure S-C1).

Table S-C1. AIC comparisons of two CJS models evaluating the effect of female tarsus length

on apparent survival, while acknowledging annual variation in encounter probability.

MARK model	npar	DeltaAICc	weight
Phi(~1)p(~time)	15	0.00	0.74
Phi(~tarsi)p(~time)	16	2.06	0.26

Figure S-C1. Model predictions for a relationship between apparent survival and female

tarsus length reveal no relationship, while controlling for imperfect detection and annual

variation in encounter probability.

Supplementary Text D

In theory, the selective benefits of larger eggs is that the subsequent hatchlings will be larger

and have higher survival owing to more intrinsic reserves provided by the mother (Blomqvist

et al., 1997). To link egg size variation to potential fitness consequences of subsequent

offspring we evaluated the predicted positive relationship between egg volume and chick

weight using the egg dataset described above but reduced observations to the nest level and

filtered to only include nests that had chicks measured within one day of hatching, resulting in

456 nests from 276 females. As it was unclear which chick came from which egg, each datum

represented the nest-level average of chick weights and egg volumes. We included random

intercepts for mother identity and year, and assumed a Gaussian error distribution of egg

volume.

Literature Cited:

Blomqvist, D., Johansson, O.C. & Götmark, F. (1997) Parental quality and egg size affect

chick survival in a precocial bird, the lapwing *Vanellus vanellus*. *Oecologia*, **110**, 18–

24.

SUPPLEMENTARY FILES

Supplementary File 1. Commented RMarkdown vignette detailing the analytical steps needed

to reproduce all our models and results. Click here to view on your internet browser:

https://raw.githack.com/leberhartphillips/snowy_plover_eggs/main/Rmd/Supplementary_File

_1/Supplementary_File_1.html

175