Effect of oxidized β-carotene-oxygen copolymer compounds on health and performance of pre- and post-weaned pigs

La Van Kinh[1], William W. Riley[2], James G. Nickerson[3], Doan Vinh[1], Nguyen Van Phu[1], Nguyen Thanh Van[1], La Thi Thanh Huyen[1], and Graham W. Burton[1*]

[1] Institute of Animal Sciences for Southern Vietnam, Binh Duong, Vietnam

[2] International School, Jinan University, Guangzhou, China

[3] Avivagen, Inc., Ottawa, ON, Canada

*Corresponding author

Email: g.burton@avivagen.com (GB)

All authors contributed equally to this work

Abstract

The discovery of naturally occurring, biologically active β-carotene-oxygen copolymer compounds in spontaneously oxidized β-carotene has stimulated interest in their potential health benefits. The copolymers, formed in Nature or synthetically by the air-oxidation of β-carotene, possess beneficial immune modulating activities that previously had been attributed to β-carotene itself. Support for these benefits is provided by previous studies showing that supplementation in feed with low parts-per-million levels of copolymer-rich, fully oxidized β-carotene (OxBC) helped reduce the negative impact of subclinical necrotic enteritis in broilers and improved growth in weaned piglets. To further assess these potential benefits, two trials were conducted in swine raised under commercial conditions in Vietnam. Trial 1, a 140-day full-grow, post-wean study with 500 28-day-old pigs, compared 2, 4 or 8 ppm OxBC against both an unsupplemented and an antibiotic control group. OxBC and antibiotics each improved growth rate, feed efficiency, and body weight compared to the control. Animals receiving 4 and 8 ppm OxBC performed better than did animals on antibiotics. In starter pigs, OxBC reduced the occurrence of diarrhea dose-dependently (4 and 8 ppm) and to a greater extent than did antibiotics. Trial 2, a 49-day study with 420 piglets, was conducted in two-stages. In Stage 1 (pre-wean), OxBC in the transition (creep) feed produced a dose-dependent trend toward increased body weight over 21 days, reaching significance at the highest inclusion level (16 ppm). In Stage 2 (post-wean), body weight gain showed a dose-dependent trend and was significant for both 8 ppm OxBC and the antibiotics at 28 days post-wean. Feed conversion was better at 8 ppm OxBC and for the antibiotic group. These findings support the concept that β-carotene-oxygen copolymers help optimize immune function, and

38   provide validation for the effectiveness of this strategy in enhancing animal performance in

39   the absence of in-feed antibiotics.

Introduction

We have previously reported that the spontaneous non-enzymatic oxidation of β-carotene produces predominantly an oxygen-rich family of copolymer compounds with immunomodulatory properties [1, 2]. Fully oxidized β-carotene, termed OxBC and rich in copolymer compounds, exerts its actions on the immune system through pathways that are distinct from either vitamin A or intact β-carotene, which are both absent. We have proposed that copolymer compounds are in fact the actual agents responsible for many of the provitamin A-independent activities of β-carotene and other carotenoids [2].

OxBC exhibits dual immunological activities relating to: a) enhancing innate immune detection and response to pathogens [2] and, b) an anti-inflammatory/pro-resolution action that limits the extent of over-zealous immune responses and reduces the level of background inflammation [3, 4].

The utility of OxBC as a feed additive and alternative to antibiotic growth promoters has been demonstrated in studies with piglets [5], sows [4] and broiler chickens [6]. In piglets, dietary supplementation with OxBC improved growth performance and prevented the vaccine-induced growth-lag associated with the PRRS (porcine reproductive and respiratory syndrome) vaccination [5]. In sows, supplementation with OxBC, begining at late gestation and continuing through lactation, resulted in reduced proinflammatory cytokine levels in colostriumn and milk concurrent with increased colostral and milk immunoglobulin levels [4]. In broilers, dietary supplementation with OxBC reduced the level of pathogen (*Clostridium perfringens)* recovered from the gut and protected against the reduction in growth performance associated with induction of subclinical necrotic enteritis (NE) [6].

Several authors have proposed that the search for suitable alternatives to antibiotic

growth promoters should focus on substances that achieve effects similar to the antibiotics

that they are intended to replace, namely, reduction in both bacterial load and inflammation

[5-8]. The results from trials with pigs and poultry highlight the utility of OxBC in achieving

both of these outcomes. The benefits observed with piglets and gestating/lactating sows likely

are consistent with the anti-inflammatory actions of OxBC, whereas, the reduction in *C.*

perfringens in the poultry study supports the benefits of innate immune priming. Note that

OxBC has no direct anti-microbial effect and that the reduction in *C. perfringens* in the

poultry study reflects actions on the host's immune system, which is better able to detect and

respond to the presence of pathogens.

In the current paper, we report the results of two dietary supplementation trials with

OxBC in swine raised under commercial production conditions in Vietnam. The trials

evaluate the utility of OxBC as a feed additive supporting the health and productivity of pigs.

The first trial evaluated OxBC over an entire 140-day post-wean growth cycle. The second

trial evaluated the effect of OxBC on the health and growth performance of pre-wean and

post-wean pigs. These trials provide the first assessment of OxBC utility as a substitute for

in-feed antibiotics in real-world, commercial, swine-production conditions.

Materials and Methods

All experimental protocols were approved by the animal care committee of the

Institute of Animal Sciences for southern Vietnam.

Trial 1. Full Grow Study

The trial was carried out at a commercial farm, Thai My Pig farm, Thai My commune,

in the Cu Chi district, Ho Chi Minh City, from 17 May to 30 November 2014.

Five hundred weaned barrows and gilts (28 days of age) were used in a 140-day

complete wean-to-finish feeding trial. The animals were reared on site and were the progeny

of Landrace x Yorkshire females and Duroc boars. The herd was asymptomatic for PRRS

(Porcine Reproductive and Respiratory Syndrome) and was vaccinated against hog cholera

(Coglapest™, Ceva-Phylaxia Veterinary Biologicals Co. Ltd, Hungary), foot-and-mouth

disease (Biotaftogen™, Biogénesis Bagó, Argentina) and PRRS (Tai Xanh™, Navetco,

Vietnam).

Animals were randomized by weight with each pen containing 20 pigs with the same

ratio of barrows to gilts. The trial was comprised of five dietary treatment groups with five

replicate pens per treatment as follows: Control (basal diet with no antibiotics or OxBC), AB

(basal diet with antibiotics; no OxBC) and OxBC (basal diet supplemented with 2, 4 or 8 ppm

(mg/kg) OxBC; no antibiotics). The compositions of the basal diets for the Starter, Grower

and Finisher phases of the study are shown in Table 1. No in-feed or water-administered

medications or feed additives were employed with the exception of the antibiotics used in the

AB group and OxBC used in the three OxBC groups. OxBC was provided in-feed as the OxC-

beta™ Livestock 10% commercial premix product (Avivagen Inc. Ottawa, Canada).

Chlortetracycline and colistin sulfate were purchased from Zhumadian Huazhong Chia Tai

Co. Ltd (China) and Lifecome Biochemistry Co. Ltd (China), respectively.

Table 1. **Composition of basal diets for Starter, Grower and Finisher phases of the Full**

Grow Study.

No.	Ingredient name	Post-Wean (g/kg)	Grower (g/kg)	Finisher (g/kg)
1	Corn	481.0	700.00	513.32
2	Rice bran	-	58.08	123.07

3	Cassava root	-	10.00	157.86
4	Soybean meal 47.5% CP	237.0	136.92	100.00
6	Whey powder 11% CP	100	-	-
7	Prelac 38% CP	100	-	-
8	Fish meal 50% CP	-	75.81	74.00
9	Fish meal 60% CP	20.0	-	-
10	Soybean oil	27.1	11.83	25.72
11	Premix Min-Vit	3.0	2.50	2.50
12	DCP	18.0	-	-
13	Seashell powder	3.4	-	-
14	Enzyme	1.0	-	-
15	Salt	2.9	2.45	2.51
16	Sweetener	0.3	-	-
17	L-Lysine	3.2	1.57	0.63
18	DL-Methionine	1.6	0.26	0.13
19	L-Threonine	1.3	0.48	0.26
20	DL-Tryptophan	0.2	0.11	0.0
	Total	1,000.0	1,000.0	1,000.0

Nutrient composition

1	Dry Matter (%)	88.15	87.85	88.04
2	ME (Kcal/kg)	3200	3,265	3,265
3	Protein (%)	22.0	18.0	15.5

4	Fat (%)	6.23	5.84	7.03
5	Fiber (%)	2.34	2.41	2.77
6	Calcium (%)	0.90	0.60	0.60
7	Total Phosphorus (%)	0.70	0.57	0.57
8	Avail. P (%)	0.52	0.48	0.48
9	Lysine (%)	1.42	1.07	0,92
10	Methionine (%)	0.82	0.39	0,32
11	Meth + Cystine (%)	0.47	0.69	0,61
12	Threonine (%)	0.91	0.81	0,70
13	Tryptophan (%)	0.27	0.21	0,19

For logistical and space purposes, the trial was conducted in a time-replicated fashion

with replicates one and two for all five treatment groups beginning on Study Day 1; replicates

three and four for all five treatment groups began the trial 28 days later, and replicate five for

all five treatment groups began the trial 56 days after the first cohort.

Each pen was equipped with a dry hopper feeder with six feed access holes, each

measuring 13 cm x 13 cm, aligned parallel to the pen front. Feeders were located at the front

of the pen, and a water drinker was located at the back of each pen. The pigs had *ad libitum*

access to feed and water throughout the trial.

Pigs were individually weighed at the start of the study (Day 1) and at Days 28 (~20

kg), 84 (~50 kg) and 140 (~100 kg). All feed given was weighed daily, the feeders were

emptied weekly and the remaining feed weighed.

Growth performance parameters were calculated for each stage of the production

cycle (Starter: Days 1-28, Grower: Days 29-84, and Finisher: Days 85-140) as well as for the

overall study period (Days 1-140).

Statistical analyses were performed using the Statistical Analysis System for Personal

Computers (SAS) Version 9.1 (SAS Institute, Cary, NC). Treatment effects were tested with

a combination of analysis of variance for continuous variables using the MIXED procedure

and Chi-square analysis, including the Cochrane-Armitage trend test. Dichotomous variables

(pigs clinically treated or removed) were created using the post-mortem and clinical results

from the data provided to analyze the number of pigs removed from or treated during the

study. The data were analyzed by logistic regression using the GLIMMIX procedure.

127 Trial 2. Pre-and Post-Wean Piglet Study

The trial was carried out at a commercial farm, Thai My Pig farm, Thai My commune,

in the Cu Chi district, Ho Chi Minh City from March 15 to May 30, 2016.

Forty-two (42) sows and their offspring were used in the two-stage, 49-day feeding

trial. Stage 1 evaluated the benefits of supplementation with OxBC in creep feed in pre-

weaned piglets. Stage 2 evaluated the benefits of continuing OxBC supplementation through

the post-wean Starter phase.

All piglets were the progeny of Landrace x Yorkshire females and Duroc boars and

were reared on site. The piglets were vaccinated against PRRS (Tai Xanh™, Navetco

Vietnam) at 14 days old.

After one day post-farrowing, sows with minimum litter sizes of ten piglets and

similar weights were selected. Litter sizes were also standardized by cross-fostering between

sows within treatment groups within three days post-farrowing. As most sows had litter sizes

greater than ten, surplus piglets were cross-fostered to sows that were not part of the trial.

Stage 1. Pre-Wean (Study Days 0-21)

In Stage 1, 42 lactating sows (one week post-farrow) and their one-week-old offspring

were randomly and evenly assigned to one of six treatments with seven replicate pens per

treatment. Each pen contained one sow and her litter of ten piglets. The mean piglet body

weight was 2.44 kg. As there were insufficient sows to carry out seven replicates

simultaneously, the trial was run in a time-replicated fashion, with two cohorts of sows. The

first cohort contained 18 sows with 180 piglets (six treatments and three replicates) and the

second cohort contained 24 sows with 240 piglets (six treatments and four replicates). The

second cohort began the trial one week after the first cohort.

The diets were introduced as soon as the piglets were ready to begin consuming creep

feed. Sows in all groups received a basal commercial lactation diet without OxBC

supplementation. Dietary treatments were delivered to piglets via supplementation of the

creep feed as follows: Control (basal diet with no antibiotic or OxBC), AB (basal diet with

antibiotics) and OxBC (basal diet with 2, 4, 8 or 16 ppm OxBC; no antibiotic). The

composition of the basal diet is shown in Table 2.

Table 2. Composition of the basal diets for the Creep and Starter phases of the Pre- and

Post-Wean Study.

Ingredient Name	Pre-Wean (g/kg)	Post-Wean (g/kg)
Corn	573.2	636.5
DDGS	-	30.0

Soybean meal 47%CP	120.0	200.0
Soya oil	30.0	31.5
Skimmed milk replacer	93.7	-
Whey powder	50.0	-
Blood Plasma Protein	43.0	39.3
Fish meal	70.0	33.5
Dicalcium phosphate	5.6	09.9
Calcium carbonate	02.5	06.8
Salt	-	02.5
Vitamin-mineral premix	5.0	03.0
L-Lysine HCl	2.7	03.6
Threamine	1.0	1.0
Methionine	1.8	1.6
Trypthophan	1.2	0.8
Sweetener	0.3	-
Total	**1000**	**1000**
Chemical composition		
Crude protein, %	22.21	21.29
Crude fiber, %	2.01	2.85
Crude fat, %	6.32	6.42
NE swine, kcal/kg	2557	2486
ME swine, kcal/kg	3275	3271
Calcium, %	0.85	0.75

Phosphorus, %	0.70	0.65
Available P, %	0.47	0.40
Sodium, %	0.29	0.28
Lysine, %	1.42	1.32
Methionine, %	0.54	0.45
Cystine, %	0.30	0.32
Met+Cys, %	0.85	0.79
Threonine, %	0.89	0.83
Tryptophan, %	0.31	0.29
Arginine, %	1.10	1.18
Isoleucine, %	0.79	0.73
Leucine, %	1.74	1.68
Valine, %	0.97	0.90
Histidine, %	0.52	0.52
Phenylalanine, %	0.92	0.92

Feed and water were provided *ad libitum*. Piglets were individually weighed at the

start of the study (Day 1) and at weaning (Day 21). Feed given was weighed daily and unused

feed was weighed at the end of each week.

Stage 2. Post-Wean (Days 22-49)

Stage 2 began at Study Day 22, one day post-wean, when the piglets were 29 days old.

Three hundred and sixty (360) piglets were used in a random block design of six treatments

with six replicate pens of ten piglets each. Piglets were randomized within their prior

treatment groups before being equally assigned into pens. Treatments were as follows:

Control (basal diet with no antibiotics or OxBC), AB (basal diet with antibiotics, no OxBC)

and OxBC (basal diet with 2, 4, 8 or 4 (16 in Stage 1) ppm OxBC; no antibiotics). The

composition of the basal diet is shown in Table 2. The piglets were provided feed and water

ad libitum. Feed given was weighed daily and any unused feed was weighed at the end of

each week.

The following measurements were performed: mean body weight (BW) at Days 1, 21

and 49 for each animal; average daily feed intake (ADFI): total feed intake per pen divided

by the product of the number of live animals and days on diet; ADG: total BW gained per pen

divided by the product of the number of live animals and days on diet; FCR: total feed

consumption divided by BW increase, defined as: (sum of final body weights of surviving

animals plus weight of mortalities and removals) – (sum of initial body weights including

initial bodyweight of animals dead or removed during the specified period); mortality and

diarrhea incidence, for each treatment group for both Stage 1 and Stage 2 of the study.

Diarrhea rate was determined as the percentage of animals presenting with loose or

watery stools.

Data were analyzed by ANOVA using Minitab Statistical Software (Minitab, State

College, PA, USA). Differences between means were tested using Fisher's multiple range

test when the F value was significant at $P < 0.05$. Data were considered significantly different

at $P<0.05$.

Results

Trial 1. Full Grow Study

The effect of dietary supplementation with OxBC on the growth performance of pigs is shown in Table 3. On Day 1, there were no significant differences in BWs among any of the groups (P>0.05). Dietary supplementation with all three levels of OxBC during the Starter period (Days 1 to 28) resulted in significant improvements in body weight (BW), average daily weight gain (ADG), average daily feed intake (ADFI) and feed/gain ratio (F/G) compared to the control group (P<0.001). Animals in the OxBC groups also performed better than or equivalent to those receiving antibiotics during the starter period.

During the Grower period (Days 29-84), supplementation with all levels of OxBC significantly increased ADG relative to the Control (P<0.001). The ADG of the AB group was also higher than the Control (P<0.001), but there were no differences among the OxBC and the AB groups. There was an apparent dose dependent decrease in feed consumption across the three OxBC groups, with the ADFI of the 8 ppm OxBC group being significantly lower than the Control (P<0.05). The F/G ratios for all OxBC groups and the AB group were significantly improved compared to the Control (P<0.001). The 8 ppm OxBC group had the lowest F/G ratio, which was significantly lower than the ratios for the Control, AB and 2 ppm OxBC groups (P<0.001).

 Table 3. Effect of dietary OxBC on ADG, ADFI, F/G and BW of pigs in the Starter, Grower, Finisher and Overall periods.[1,2]

Treatment:	Control	AB[3]	OxBC 2 ppm	OxBC 4 ppm	OxBC 8 ppm	Pooled SEM	P-value
Starter Period: Days 1-28							
BW Day 1	7.7±0.1	7.8±0.1	7.6±0.1	7.6±0.1	7.8±0.1	0.11	0.152
BW Day 28	17.7±0.3[b]	18.4±0.3[b]	19.8±0.5[a]	20.0±0.5[a]	20.2±0.5[a]	0.41	<0.001
ADG	358±8[b]	381±12[b]	435±20[a]	440±19[a]	442±15[a]	15	<0.001
(% Difference)[4]	-	6	22	23	23	-	-
ADFI	601±10[b]	607±20[b]	689±30[a]	676±27[a]	684±21[a]	23	<0.001
(% Difference)[4]	-	1	15	13	14	-	-
Feed/Gain	1.68±0.03[a]	1.59±0.03[b]	1.59±0.01[bc]	1.54±0.02[d]	1.55±0.01[cd]	0.02	<0.001
(% Difference)[4]	-	-5.4	-5.4	-8.3	-7.7	-	-

Grower Period: Days 29-84

BW Day 84	49.6±0.3[c]	51.9±0.8[b]	53.2±0.4[a]	53.6±0.5[a]	53.8±0.5[a]	0.51	<0.001
ADG	569±2[b]	598±17[a]	596±5[a]	601±14[a]	600±11[a]	11	0.001
(% Difference)[4]	-	5.1	4.7	5.6	5.4	-	-
ADFI	1645±78[a]	1581±35[ab]	1625±33[ab]	1590±34[ab]	1542±29[b]	46	0.018
(% Difference)[4]	-	-3.9	-1.2	-3.3	-6.3	-	-
Feed/Gain	2.89±0.14[a]	2.73±0.03[b]	2.73±0.05[b]	2.64±0.04[bc]	2.57±0.08[c]	0.08	<0.001
(% Difference)[4]	-	-5.5	-5.5	-8.7	-11.1	-	-

Finisher Period: Days 85-140

BW Day 140	99.5±0.5[c]	101.7±0.5[b]	103.6±0.5[a]	104.2±0.3[a]	104.4±0.5[a]	0.44	<0.001
(% Difference)[4]	-	2.2	4.1	4.7	4.9	-	-
ADG	892±13	889±8	901±9	902±7	904±9	9.3	0.056

(% Difference)[4]	-	0.3	1.0	1.1	1.3	-	-
ADFI	3085±114[a]	3031±102[ab]	2959±87[ab]	2897±47[b]	2876±49[b]	84	0.004
(% Difference)[4]	-	-1.8	-4.1	-6.1	-6.8	-	-
Feed/Gain	3.46±0.14[a]	3.41±0.13[a]	3.28±0.08[ab]	3.21±0.05[b]	3.18±0.03[b]	0.09	<0.001
(% Difference)[4]	-	-1.4	-3.8	-2.1	-0.9	-	-

Overall: Days 1-140

ADG	656±4[c]	671±3[b]	686±4[a]	690±1[a]	690±3[a]	3.2	<0.001
(% Difference)[4]	-	2.3	4.6	5.2	5.2	-	-
ADFI	2012±77[a]	1966±50[ab]	1971±41[ab]	1930±25[ab]	1904±30[b]	48	0.021
(% Difference)[4]	-	-2.3	-2.0	-4.1	-5.4	-	-
Feed/Gain	3.07±0.12[a]	2.93±0.06[b]	2.87±0.05[bc]	2.80±0.04[c]	2.76±0.04[c]	0.07	<0.001
(% Difference)[4]	-	-4.6	-6.5	-8.8	-10.1	-	-

[1] Values represent means ± SD. Values in the same row with different superscripts, a,b,c, are significantly different (P<0.05).

[2] Units: BW: kg/head; ADG, ADFI: g/head/d.

[3] Antibiotics were included in the feed as follows: Pre-wean (nursery): 100 ppm colistin sulfate; Starter (days 0-28): 150 ppm

chlortetracycline and 100 ppm colistin sulfate; Grower (days 29-84): 150 ppm chlortetracycline; Finisher (days 85-140): no antibiotics.

[4] Percent difference compared to Control.

During the Finisher stage (Days 85-140), the ADGs of animals in the OxBC groups

were greater than those of either the Control or AB groups, although the differences did not

reach statistical significance (P>0.05). Supplementation with 4 or 8 ppm OxBC resulted in

significant improvements in the F/G ratio compared to the Control and AB groups (P<0.001).

As was observed in the grower period, animals receiving OxBC had lower feed intake during

the finisher phase when compared to the Control. The feed intake was significantly lower

(P<0.005) in the 4 ppm OxBC and 8 ppm OxBC groups compared to the Control group. Note

that during the finisher stage, animals in the AB group did not receive antibiotics, which is a

common practice and legal requirement in many jurisdictions, including Vietnam.

Overall (Study Days 1-140), dietary supplementation with OxBC at 2, 4 or 8 ppm led

to significant increases in final BW and ADG, relative to both the Control and the AB groups

(P<0.001). OxBC also reduced the overall F/G (P<0.001) and ADFI (P<0.05) compared to

the Control with the 4 and 8 ppm groups also outperforming the AB group on F/G (P<0.001).

In terms of clinical health, the results presented in Table 4 show that the highest

incidences of both diarrhea and mortality were observed in the Control group for all study

periods, with the lowest being observed in the OxBC groups. The highest incidence of

diarrhea was during the Starter period, and OxBC significantly and dose-dependently reduced

the diarrhea rate (P<0.001). Diarrhea was the lowest (less than half that of the Control group)

in the 8 ppm OxBC group, and both the 4 and 8 ppm OxBC groups had significantly reduced

incidences relative to the AB group (P<0.001).

Table 4. **Effect of dietary OxBC on diarrhea incidence and overall mortality of pigs in the Starter, Grower, Finisher and Overall**

periods.[1]

Treatment:	Control	AB[2]	OxBC 2 ppm	OxBC 4 ppm	OxBC 8 ppm	Pooled SEM	P-value
Starter Period: Days 1-28							
Diarrhea (%)	7.9±0.6ª	6.6±0.5ᵇ	6.0±0.2ᵇ	4.9±0.4ᶜ	3.7±0.6ᵈ	0.47	<0.001
(% Difference)[3]	-	-17	-25	-39	-54	-	-
Grower Period: Days 29-84							
Diarrhea (%)	2.7±0.2ª	2.1±0.1ᵇ	2.0±0.2ᵇ	1.6±0.1ᶜ	1.4±0.1ᶜ	0.16	<0.001
(% Difference)[3]	-	-21	-26	-41	-48	-	-
Finisher Period: Days 85-140							
Diarrhea (%)	0.87±0.06ª	0.70±0.06ᵇ	0.64±0.07ᵇᶜ	0.67±0.06ᵇ	0.55±0.08ᶜ	0.06	<0.001
(% Difference)[3]	-	-20	-26	-23	-37	-	-

Overall: Days 1-140

Diarrhea (%)	3.1±0.1[a]	2.5±0.1[b]	2.3±0.1[c]	1.9±0.1[d]	1.5±0.1[e]	0.10	<0.001
(% Difference)[3]	-	-19	-27	-39	-50	-	-
Mortality (%)	5.0±3.5	3.0±2.7	2.0±2.7	2.0±2.7	2.0±2.7	2.9	0.431

[1] Values represent means ± SD. Values in the same row with different superscripts, a,b,c, etc., are significantly different ($P<0.05$).

[2] Antibiotics were included in the feed as follows: Pre-wean (nursery): 100 ppm colistin sulfate; Starter (Days 1-28): 150 ppm chlortetracycline and 100 ppm colistin sulfate; Grower (Days 29-84): 150 ppm chlortetracycline; Finisher (Days 85-140): no antibiotics.

[3] Percent difference compared to Control.

The occurrence of diarrhea during the Grower and Finisher periods was markedly less

compared to the Starter period. Despite the lower background incidence of diarrhea in these

later growth phases, OxBC still continued to provide significant and dose-dependent

protection from the condition.

Overall, animals in the Control group had the highest incidence, whereas animals in

the 8 ppm OxBC group had the lowest incidence. OxBC significantly and dose-dependently

reduced the incidence relative to both the Control and AB groups (P<0.001).

Mortality was relatively low, overall, with few pigs dying during the study. The lowest

number of deaths occurred in the OxBC groups, but differences were not significant between

any of the groups.

250   Trial 2: Pre-and Post-Wean Piglet Study

251   Stage 1 Pre-Weaning (Days 1-21)

The effects of OxBC and antibiotics on the growth performance of pre-weaning

piglets are provided in Table 5.

Inclusion of OxBC in the creep feed resulted in a dose-dependent trend towards

increased ADG and BW that reached significance for 16 ppm OxBC compared to the Control

over the 21 days of Stage 1. The ADG and BW of the AB group were also higher than the

Control. Antibiotics and 16 ppm OxBC improved the F/G ratio compared to the Control,

while the ADFI was not significantly different across the 6 treatment groups (P>0.05).

The very low F/G ratio observed for all groups is attributed to the small amounts of

feed ingested by the piglets, which were still mainly consuming milk from the sows.

Table 5. BW, ADG, ADFI and F/G of pre-weaning piglets in Stage 1 of Trial 2 (Days 1-21)[1,2]

Treatment:	Control	AB[3]	OxBC 2 ppm	OxBC 4 ppm	OxBC 8 ppm	OxBC 16 ppm	SEM	P-Value
BW Day 0	2.45±0.07	2.44±0.09	2.43±0.08	2.45±0.10	2.46±0.05	2.45±0.07	0.078	0.976
(% Difference)[4]	-	-0.4	-0.8	0.0	0.8	0.4		
BW Day 21	6.88±0.07^c	7.21±0.07^a	6.92±0.17^{bc}	6.94±0.12^{bc}	6.98±0.08^{bc}	7.10±0.13^{ab}	0.11	0.001
(% Difference)[4]	-	4.8	0.6	0.9	1.3	3.2		
ADG	211±2.3^c	227±2.8^a	214±7.5^{bc}	214±2.9^{bc}	215±4.2^{bc}	222±7.8^{ab}	5.1	0.001
(% Difference)[4]	-	8.1	1.9	1.9	2.4	5.2		
ADFI	79.9±0.9	80.1±0.9	79.9±0.9	79.7±1.2	80.5±1.1	79.5±1.5	1.1	0.592
(% Difference)[4]	-	0.9	0.1	-0.3	0.9	-0.4		
Feed/Gain	0.38±0.01^a	0.35±0.01^c	0.37±0.01^{ab}	0.37±0.01^{ab}	0.37±0.01^{ab}	0.36±0.02^{bc}	0.01	0.001
(% Difference)[4]	-	-7.9	-2.6	-2.6	-2.6	-5.3		
Intake of Active[5]	0	8.05	0.16	0.32	0.64	1.27	-	-

[1] Values represent means ± SD. Values in the same row with different superscripts, a,b,c, are significantly different (P<0.05).

[2] Units: BW: kg/head; ADG, ADFI: g/head/d; Intake of Active: mg/d.

[3] The AB diet contained 100 ppm colistin sulfate.

[4] Percent difference compared to Control.

[5] Average daily intake (mg) of antibiotic or OxBC.

No evidence of diarrhea or mortality was observed in Stage 1.

Trial 2: Stage 2 (Days 22-49)

The effects of OxBC and antibiotics on BW, ADG, ADFI and F/G for weaned piglets

are shown in Table 6.

At the beginning of Stage 2, a subset of select piglets from each treatment group was

randomized within treatment groups to continue the trial. Piglets were selected such that the

average BW among the six treatment groups did not differ on Day 22 (the beginning of Stage

2). After 28 days (Day 49, the end of Stage 2), average BWs were highest for the AB group

followed by the four OxBC groups and the Control. Increases in average BW reached

statistical significance for both the AB and the 8 ppm OxBC groups (P<0.001).

The increases in BW were reflected in the ADG. The AB group had the highest ADG,

which was significantly higher than the Control (P<0.001). ADG showed a trend towards a

dose-dependent increase in the OxBC groups, reaching statistical significance for the 8 ppm

OxBC group compared to the Control. However, there was no statistical difference in ADG

across the four OxBC groups.

There were no significant differences in ADFI among the treatment groups. F/G ratios

were significantly improved for both the AB group and the 8 ppm OxBC group compared to

all other groups (P<0.001).

The effects on the incidence of diarrhea and mortality are given in Table 7.

Table 6. BW, ADG, ADFI and F/G of weaned piglets in Stage 2 of Trial 2 (Days 22-49).[1,2]

Treatment:	Control	AB[3]	OxBC 2 ppm	OxBC 4 ppm	OxBC 8 ppm	OxBC 4 ppm[4]	SEM	P-Value
BW Day 21	6.91±0.07	6.95±0.08	6.93±0.06	6.94±0.09	6.93±0.06	6.98±0.07	0.07	0.697
(% Difference)[5]	-	0.4	0.3	0.4	0.1	1.0		
BW Day 49	17.74±0.12^c	18.80±0.25^a	17.94±0.50^{bc}	18.00±0.47^{bc}	18.32±0.09^{ab}	18.23±0.07^{bc}	0.31	0.001
(% Difference)[5]	-	6.0	1.1	1.5	3.3	2.8		
ADG	387±5^c	423±7^a	393±16^{bc}	395±16^{bc}	407±3^{ab}	402±2^{bc}	10	0.001
(% Difference)[5]	-	9.6	1.8	2.3	5.4	3.9		
ADFI	639.7±7.4	652.4±7.5	640.2±15	639.5±3.7	637.5±7.1	643.7±5.7	2.2	0.087
(% Difference)[5]	-	2.0	0.1	0.0	-0.3	0.6		
Feed/Gain	1.65±0.04^a	1.54±0.03^c	1.63±0.08^{ab}	1.62±0.06^{ab}	1.57±0.01^{bc}	1.60±0.01^{ab}	0.035	0.001
(% Difference)[5]	-	-6.7	-1.2	-1.8	-5.5	-3.0		
Intake of Active[6]	0	59.2 CS 88.8 CTC	1.16	2.32	4.62	2.33	-	-

[1] Values represent means ± SD. Values in the same row with different superscripts, a,b,c, are significantly different (P<0.05).

[2] Units: BW: kg/head; ADG, ADFI: g/head/d; Intake of Active: mg/d.

[3] The AB diet contained 100 ppm colistin sulfate (CS) and 150 ppm chlortetracycline (CTC).

[4] The treatment provided 16 ppm OxBC during Stage 1 and 4 ppm OxBC during Stage 2. The supplementation was higher during Stage

1 to account for the low feed intake of nursing piglets.

[5] Percent difference compared to the Control.

[6] Average daily intake (mg) of antibiotic or OxBC.

Table 7. Diarrhea incidence and mortality in weaned piglets during Stage 2 (Days 22-49)

Treatment:	Control	AB[1]	OxBC 2 ppm	OxBC 4 ppm	OxBC 8 ppm	OxBC 4 ppm[2]
Diarrhea (%)	4.7±0.8	3.4±1.1	4.6±1.3	4.1±0.9	3.6±1.1	3.9±0.8
(% Difference)[3]	-	-28	-1	-12	-24	-18
Mortality (%)	5.0	0	3.4	3.4	0	0
% of Control	(100)	0	67	67	0	0

[1] The AB diet contained 100 ppm colistin sulfate and 150 ppm chlortetracycline.

[2] The treatment provided 16 ppm OxBC during Stage 1 and 4 ppm OxBC during Stage 2. The supplementation was higher during Stage 1 to account for the low feed intake of nursing piglets.

[3] Percent difference compared to the Control.

Diarrhea and mortality incidence were generally low in all groups, with the highest

incidence of diarrhea and mortality being observed in the Control group. Treatment with 8

ppm OxBC or antibiotics decreased diarrhea incidence by 24% and 28%, respectively.

No mortalities were recorded in the AB group, the 8 ppm OxBC group or the group

fed 16 ppm OxBC in Stage 1 and 4 ppm OxBC in Stage 2. The differences in mortalities

between the groups were not significant.

Discussion

Results from the two trials reported here demonstrate that dietary supplementation

with low parts-per-million levels of OxBC improves the health and productivity of pigs reared

under commercial production conditions. In Trial 1, conducted over the entire 140-day post-

wean grow-out period, inclusion of OxBC in the feed led to improvements in growth

performance during each of the three stages of the production cycle and also significantly

reduced the incidence of diarrhea in post-wean starter piglets. Results from the second trial

confirm the benefits of dietary OxBC on growth performance of post-wean starter piglets and

indicate that further benefits on piglet performance are possible when OxBC is added to the

transition (creep) feed of nursing piglets prior to weaning.

In Trial 1, the observation that the benefits of OxBC were most apparent during the

starter period compared to the grower and finisher periods is not surprising, given that OxBC

plays a role in supporting immunity [1-3] and the fact that young piglets experience high

levels of stress associated with weaning and that they are immunologically immature. The

highest incidence rates of diarrhea occurred during the starter period, consistent with the well-

known susceptibility of young piglets to post-wean diarrhea [7]. The reduced incidence of

diarrhea observed in the OxBC supplemented groups relative to controls is consistent with

the immune priming actions of OxBC, which center upon an ability to increase the level of

pathogen pattern recognition receptors (PPRRs) and enhanced down-stream innate immune

response to receptor activation [2].

More specifically, OxBC has been shown to increase the level of toll-like receptor

subtypes 2 and 4 (TLR-2, TLR-4) and their coreceptor CD14 in both *in vitro* and *in vivo*

studies [2]. TLR-4 and CD14 are well-characterized co-receptors responsible for recognizing

the lipopolysaccharide (LPS) moiety present in gram negative bacteria, such as *E. coli,* while

TLR-2 recognizes the lipoteichoic acid (LTA) moiety present on gram positive bacteria, such

as *Clostridium perfringens* [8-10]. Binding of bacterial LPS or LTA to the TLR-4/CD14 or

TLR-2 receptor triggers an innate immune response aimed at clearing the bacteria [11, 12].

A study comparing *E. coli* resistant versus susceptible lineages of pigs found that the

genetic basis for the resistance to *E. coli* colonization of the gut was higher levels of CD14

expression [13]. The authors concluded that the TLR-4/CD14 pathway plays a significant role

in reducing *E. coli* colonization and the incidence of diarrhea in piglets. While we did not

measure intestinal receptor or *E. coli* levels in the current trials, an increase in intestinal CD14

and TLR-4 content (as we have observed in mice fed OxBC [2]) and the subsequent reduction

in *E. coli* colonization offers a plausible explanation for the reduced incidence of diarrhea

observed in the OxBC-supplemented piglets. Results from an infectious challenge study in

broilers, where dietary supplementation with OxBC reduced *Clostridium perfringens* levels

by 2 to 3 log units, were consistent with the concept that OxBC supports the intestinal immune

system and allows the host to resist pathogen colonization [6].

The improvement in clinical health (reduced incidence of diarrhea) likely explains, in

part, the improved growth performance of piglets in the OxBC-supplemented versus control

groups during the Starter period in Trial 1. As would be expected for older, more

immunocompetent pigs, there were fewer incidences of diarrhea in grower and finisher pigs

compared to starter pigs across all treatment groups in the first trial. While there tended to be

a lower incidence of diarrhea in the OxBC supplemented groups compared to the control

during the grower and finisher periods, it seems unlikely that these small reductions in disease

incidence would explain the improved growth performance observed in the OxBC groups

during the latter stages of the trial. Instead, the ability of OxBC to reduce inflammatory tone

represents the most plausible mechanism to explain the improved growth of older pigs in the

OxBC groups.

The intestine has been described by some authors as a tissue in a constant state of

controlled inflammation, and there is a need for organisms to tightly control the level of

intestinal inflammation [14, 15]. Commercial production practices are known to further

exacerbate the inflammatory state of the intestine, as stresses associated with weaning, *ad*

libitum feeding of high energy diets, and the presence of anti-nutritional components in the

diet all contribute to increased inflammation of the gut [16]. The contribution that reduced

inflammation plays in improving growth performance is further highlighted by the fact that

an anti-inflammatory action has been proposed as an underlying mechanism contributing to

the growth promoting effects of subtherapeutic antibiotic growth promoters [15].

Results from studies in an experimental model of bovine respiratory disease (BRD)

and in gestating/lactating sows demonstrate an anti-inflammatory action for OxBC. In the

experimental BRD trial, animals that received dietary supplementation with OxBC had a

significant increase in anti-inflammatory or pro-resolution activity in the lung [3] while

OxBC-supplementation in gestating/lactating sows led to reductions in the concentration of

pro-inflammatory cytokines, TNFα and IL-8 in colostrum and milk [4]. Reduced

inflammation may also be partly responsible for the observed reduction in intestinal lesion

severity in *C. perfringens*-challenged broilers [6].

The performance benefits of OxBC in weaned piglets observed in Trial 1 are

supported by the results from Trial 2 in which inclusion of OxBC in the Starter feed led to an

apparent dose-dependent trend for improved growth performance and feed efficiency, with

the highest inclusion level of OxBC (8 ppm) reaching significance. In terms of clinical health,

the results from Trial 2 indicate that, while the incidences of diarrhea were numerically lower

in the antibiotic and OxBC supplemented groups compared to the control, there was no

significant treatment effect (Table 7). There were considerably fewer cases of diarrhea

observed in the control group in Trial 2 (4.7%) compared to Trial 1 (7.9%). This difference

in the background incidence of diarrhea in the two trials likely reflects a lower level of

endemic pathogen pressure during Tial 2 relative to Trial 1. With a relatively low incidence

of diarrhea observed in the control group in Trial 2, further treatment-induced reductions may

not have been possible. Thus, in Trial 2 the benefits of OxBC on growth performance of post-

wean piglets is more plausibly due to anti-inflammatory actions rather than a reduction in the

incidence of diarrhea. It is interesting to note that the magnitude of the performance benefits

in piglets was much larger in Trial 1 compared to Trial 2. The difference in the magnitude of

the benefits may have been due to a combined benefit of reducing diarrhea and background

inflammation in the gut in Trial 1, while in the second trial, where there was lower endemic

pathogen pressure, only the anti-inflammatory activity may have factored in the benefit.

The results from the pre-wean period (Stage 1) of Trial 2 indicate that inclusion of

OxBC in transition or creep feed can also improve weight gain and body weight of nursing

piglets. OxBC appeared to dose-dependently increase ADG and body weight at weaning with

the highest inclusion rate of 16 ppm OxBC reaching statistical significance (Table 6). The

lack of a statistically detectable effect at the lower OxBC inclusion rates likely reflects the

very low amounts of OxBC received by nursing piglets that consumed relatively little feed.

Given the apparent dose-dependent improvements in growth performance, it would be useful

to investigate higher inclusion rates of OxBC in transition feed or OxBC-supplementation of

milk replacer. There were no mortalities or incidences of diarrhea observed during the pre-

wean stage of Trial 2, consistent with the presence of a relatively low level of endemic

pathogen pressure.

Overall, the results of the two trials reported here demonstrate the benefits of dietary

supplementation with OxBC on growth performance, feed efficiency, and the health of pre-

and post-weaned piglets. These findings have significance for both the commercial feed

industry and the field of nutrition. For the feed industry, the findings support the application

of OxBC as an effective alternative to antibiotic growth promoters. The performance and

health benefits obtained with OxBC were comparable to those of the antibiotic growth

promoters in both trials. For the field of nutrition, the results lend further support to the

identification of the copolymers present in OxBC as a source of the benefits of dietary β-

carotene. The concept that β-carotene is a source of beneficial compounds beyond vitamin A

is not new; however, there is a dearth of reproducible evidence as to the identity and biological

actions of these compounds. A scientific panel of the European Food Safety Association

(EFSA) recently concluded there was insufficient scientific evidence to support a proposed

benefit of β-carotene, as a feed additive, on host immunity [17]. The EFSA panel cited an

inconsistency of results with β-carotene as the major problem. The inconsistent results cited

by EFSA likely stem from the fact that the proposed immunological benefits arise not from

vitamin A or a direct action of β-carotene itself but rather from the presence of copolymers,

which were unknown at the time. Dietary supplementation with β-carotene potentially

provides variable levels of copolymers, depending upon the extent of adventitious oxidation.

β-Carotene's susceptibility to oxidation is recognized by manufacturers who nowadays

formulate the product to help prevent its oxidation.

In the trials reported here, the source of the β-carotene copolymers is a commercial

product manufactured by a highly reproducible process under stringent quality control

standards. However, the copolymers have also been shown to exist naturally at varying levels

in various food and feedstuffs [18, 19]. The presence of naturally occurring copolymers may

provide an explanation, at least in part, for the health and productivity benefits associated

with the use of certain traditional forages. For example, alfalfa, a forage rich in β-carotene,

has been proposed to contain unidentified growth factors that benefit the growth and

reproductive performance of various species [20-22]. The mechanisms of action of the so-

called unidentified growth factors remains largely unknown; however the presence of

naturally occurring copolymers arising from oxidation of β-carotene when alfalfa is dried [18]

represents one possible growth factor.

We propose that β-carotene is the source of two classes of compounds that are

biologically active and which both benefit health, the first being vitamin A and the second

being β-carotene oxygen copolymers. Driven by the desire to reduce costs, the modern feed

industry has largely replaced its use of β-carotene, whether synthetic or natural, with synthetic

vitamin A. Removal of β-carotene, particularly natural sources, such as alfalfa, from diets has

resulted in the simultaneous, albeit unintended, removal of the naturally present β-carotene

copolymers and their associated benefits. The positive effects on growth performance and

health observed with OxBC in these trials coupled with similar findings in previous trials in

swine [4, 5], poultry [6] and dairy calves [3] demonstrate that re-introduction of very low

amounts of β-carotene oxygen copolymer into feeds can beneficially support the health and

productivity of livestock and substantially reduce the need for in-feed antibiotics.

References

1.   Burton GW, Daroszewski J, Nickerson JG, Johnston JB, Mogg TJ, Nikiforov GB. ß-

Carotene autoxidation: oxygen copolymerization, non-vitamin A products and

immunological activity. Can J Chem. 2014;92: 305-316. doi: 10.1139/cjc-2013-0494.

2.   Johnston JB, Nickerson JG, Daroszewski J, Mogg TJ, Burton GW. Biologically

active polymers from spontaneous carotenoid oxidation. A new frontier in carotenoid

activity. PLoS ONE. 2014;9: e111346. doi: doi:10.1371/journal.pone.0111346.

3.   Duquette SC, Fischer CD, Feener TD, Muench GP, Morck DW, Barreda DR, et al.

Anti-inflammatory benefits of retinoids and carotenoid derivatives: retinoic acid and fully

oxidized β-carotene induce caspase-3-dependent apoptosis and promote efferocytosis of

bovine neutrophils. Am J Vet Res. 2014;75: 1064-1075. doi: 10.2460/ajvr.75.12.1064.

4.   Chen J, Zhang Y, Lu Y, Tian M, Cheng L, Chen F, et al. Effects of dietary

supplementation with fully oxidized β-carotene during late gestation and lactation on

productivity and immune status of sows. Brit J Nutr. 2020. doi:

10.1017/S0007114520002652.

5.   Hurnik D, Daroszewski J, Burton GW, editors. Determination of the effect of a fully

oxidized β -carotene dietary supplement on the immune system and growth performance of

weaned pigs. American Association of Swine Veterinarians 42nd Annual Meeting; 2011;

Phoenix, Arizona.

6.   Kang M, Oh JY, Cha SY, Kim WI, Cho HS, Jang HK. Efficacy of polymers from

spontaneous carotenoid oxidation in reducing necrotic enteritis in broilers. Poultry Sci.

2018;97: 3058-3062. doi: 10.3382/ps/pey180.

7.   Rhouma M, Fairbrother JM, Beaudry F, Letellier A. Post weaning diarrhea in pigs:

risk factors and non-colistin-based control strategies. Acta Vet Scand. 2017;59: 31. doi:

10.1186/s13028-017-0299-7.

8.   Dessing MC, Schouten M, Draing C, Levi M, von Aulock S, van der Poll T. Role

Played by Toll-Like Receptors 2 and 4 in Lipoteichoic Acid—Induced Lung Inflammation

and Coagulation. J Infect Dis. 2008;197: 245-252. doi: 10.1086/524873.

9.   Park BS, Lee JO. Recognition of lipopolysaccharide pattern by TLR4 complexes.

Exp Mol Med. 2013;45: e66. Epub 2013/12/07. doi: 10.1038/emm.2013.97. PubMed

PMID: 24310172; PubMed Central PMCID: PMCPMC3880462.

10.   Takehara M. [Host Defense against Bacterial Infection and Bacterial Toxin-induced

Impairment of Innate Immunity]. Yakuga Zasshi. 2018;138: 1249-1253. Epub 2018/10/03.

doi: 10.1248/yakushi.18-00102. PubMed PMID: 30270267.

11.   Mukherjee S, Karmakar S, Babu SP. TLR2 and TLR4 mediated host immune

responses in major infectious diseases: a review. Braz J Infect Dis. 2016;20: 193-204. Epub

2016/01/19. doi: 10.1016/j.bjid.2015.10.011. PubMed PMID: 26775799.

12.   Vijay K. Toll-like receptors in immunity and inflammatory diseases: Past, present,

and future. Int Immunopharmacol. 2018;59: 391-412. Epub 2018/05/08. doi:

10.1016/j.intimp.2018.03.002. PubMed PMID: 29730580; PubMed Central PMCID:

PMCPMC7106078.

13.   Wu Z, Liu Y, Dong W, Zhu GQ, Wu S, Bao W. CD14 in the TLRs signaling

pathway is associated with the resistance to E. coli F18 in Chinese domestic weaned piglets.

Sci Rep-UK. 2016;6: 24611. Epub 2016/04/22. doi: 10.1038/srep24611. PubMed PMID:

27098998; PubMed Central PMCID: PMCPMC4838916.

14.   Biancone L, Monteleone I, Del Vecchio Blanco G, Vavassori P, Pallone F. Resident

bacterial flora and immune system. Digest Liver Dis. 2002;34 Suppl 2: S37-43. Epub

2002/11/01. doi: 10.1016/s1590-8658(02)80162-1. PubMed PMID: 12408438.

15.   Niewold TA. The nonantibiotic anti-inflammatory effect of antimicrobial growth

promoters, the real mode of action? A hypothesis. Poultry Sci. 2007;86: 605-609. PubMed

PMID: 17369528.

16.   Pluske JR, Turpin DL, Kim JC. Gastrointestinal tract (gut) health in the young pig.

Anim Nutr. 2018;4: 187-196. Epub 2018/08/25. doi: 10.1016/j.aninu.2017.12.004. PubMed

PMID: 30140758; PubMed Central PMCID: PMCPMC6104527.

17.   EFSA. Scientific Opinion on the safety and efficacy of beta-carotene as a feed

additive for all animal species and categories. EFSA Journal. 2012;10: 2737. doi:

10.2903/j.efsa.2012.2737.

18.   Burton GW, Daroszewski J, Mogg TJ, Nikiforov GB, Nickerson JG. Discovery and

Characterization of Carotenoid-Oxygen Copolymers in Fruits and Vegetables with Potential

Health Benefits. J Agric Food Chem. 2016;64: 3767-3777. doi: 10.1021/acs.jafc.6b00503.

PubMed PMID: 27111491.

19.   Schaub P, Wust F, Koschmieder J, Yu Q, Virk P, Tohme J, et al. Nonenzymatic

beta-Carotene Degradation in Provitamin A-Biofortified Crop Plants. J Agric Food Chem.

2017;65: 6588-6598. Epub 2017/07/14. doi: 10.1021/acs.jafc.7b01693. PubMed PMID:

28703588.

20.   Scott HM, Fisher H, Snyder JM. Alfalfa Meal as a Source of Unidentified Growth

Factors. Poultry Sci. 1953;32: 555-558. doi: https://doi.org/10.3382/ps.0320555.

21.   Lakhanpal RK, Davis JR, Typpo JT, Briggs GM. Evidence for an unidentified

growth factor(s) from alfalfa and otherplant sources for young guinea pigs. J Nutr. 1966;89:

341-346. Epub 1966/07/01. doi: 10.1093/jn/89.3.341. PubMed PMID: 5950206.

22.   Hertrampf JW, Piedad-Pascual F. Unidentified Growth Factors. In: Handbook on

Ingredients for Aquaculture Feeds: Springer, Dordrecht; 2000. https://doi.org/10.1007/978-

94-011-4018-8_44