Memory Trace Imbalance in Reinforcement and Punishment Systems Can Reinforce

Implicit Choices Leading to Obsessive-Compulsive Behavior

Yuki Sakai[a,b,†], Yutaka Sakai[c,†], Yoshinari Abe[b], Jin Narumoto[b], Saori C. Tanaka[a,*]

[a]ATR Brain Information Communication Research Laboratory Group, 2-2-2 Hikaridai Seika-Cho, Soraku-Gun, Kyoto 619-0288, Japan

[b]Department of Psychiatry, Graduate School of Medical Science, Kyoto Prefectural University of Medicine, 465 Kajii-Cho, Kawaramachi-Hirokoji, Kamigyo-Ku, Kyoto 602-8566, Japan

[c]Brain Science Institute, Tamagawa University, 6-1-1, Tamagawa-gakuen, Machida, Tokyo 194-8610, Japan

[†]These authors contributed equally and should be considered co-first authors.

*Corresponding author: Saori C. Tanaka

Computational Neuroscience Laboratories, Advanced Telecommunications Research Institute International, 2-2-2 Hikaridai, Seika-Cho, Soraku-Gun, Kyoto 619-0288, Japan.

Tel: +81-774-95-1250; Fax: +81-774-95-1236.

E-mail: xsaori@atr.jp

Abstract

We may view most of our daily activities as rational action selections; however, we sometimes

reinforce maladaptive behaviors despite having explicit environmental knowledge. In this study,

we modeled obsessive-compulsive disorder (OCD) symptoms as implicitly learned maladaptive

behaviors. Simulations in the reinforcement learning framework showed that agents implicitly

learn to respond to intrusive thoughts when the memory trace signal for past actions decays

differently for positive and negative prediction errors. Moreover, our model extends our

understanding of therapeutic effects of behavioral therapy in OCD. Using empirical data, we

confirmed that patients with OCD showed extremely imbalanced traces, which were normalized

by serotonin enhancers. We found that healthy participants also varied in their obsessive-

compulsive tendencies, consistent with the degree of imbalanced traces. These behavioral

characteristics can be generalized to variations in the healthy population beyond the spectrum of

clinical phenotypes.

Introduction

We live in a complex and dynamic environment and choose better behaviors based upon trial and error. Although we may view most of our daily activities as rational action selections based on our environmental knowledge, in reality, this is not always the case. For example, we may repeatedly study hard even if we are already prepared enough for an exam. We may also find ourselves abiding by superstitions, even though we know they make no sense. In these situations, there is a minor discrepancy between explicit environmental understanding, i.e., action-outcome knowledge, and implicit learning. In cases where these discrepancies are significant, they are accompanied by "ego-dystonic" anxiety, which is a characteristic of obsessive-compulsive disorder (OCD)[1, 2]. In patients with OCD, an "ego-dystonic" anxiety becomes lodged in their minds[3]. To neutralize and relieve the anxiety, they tend to perform compulsive actions, despite their cost, e.g., excessively washing their hands or repeatedly checking their keys, even though they realize that their behavior is excessive and unreasonable[4].

We must consider its theoretical and biological implementation when investigating implicit learning leading to ego-dystonic anxiety. Implicit trial and error learning requires repetitive experiences, and reinforcement learning theory provides a framework for learning appropriate choices through repetitive experiences[5]. Reinforcement learning enhances or punishes recent choices based on differences between actual outcomes and predictions, called prediction errors. Positive prediction errors reinforce recent choices, whereas negative errors punish them.

When examining the biological implementation of implicit learning, it is essential to consider previous work demonstrating that the activity of dopamine-projecting neurons in the midbrain resembles prediction errors[6]. Dopaminergic neurons project widely to cortical and

subcortical areas. One of the main targets of dopamine is the striatum, which is located in the basal ganglia[7]. In the striatum, distinct types of dopamine receptors—D1 and D2 receptors—are expressed in exclusive neuron groups, which are thought to be involved in distinct circuit patterns, the direct and indirect pathways[7]. Previous research implies that reinforcement and punishment of recent actions may be reflected in the direct and indirect pathways, respectively[8, 9].

Since abundant evidence suggests that imbalance between direct and indirect pathways is central to the pathophysiology of OCD[10-12], we hypothesized that obsessive-compulsive behaviors are generated by imbalanced implicit learning between reinforcement and punishment. First, we propose a computational model that enables implementation of imbalanced learning between reinforcement and punishment. We conducted a simulation showing that imbalanced learning reinforces implicit learning, leading to a spiral of repetitive obsession and compulsion, even if agents have intact environmental understanding. Second, we tested our hypothesis in a behavioral task for participants with OCD and healthy participants. In addition, we evaluated mechanisms of first-line treatments, i.e., behavioral therapy and psychotropic medication for OCD, in our computational model and behavioral data. Finally, we investigated whether these behavioral characteristics could be generalized to variations in the healthy population, beyond the spectrum of clinical phenotypes.

Results

Theoretical model of OCD

We described mental states involved in obsessive-compulsive symptoms into stochastic transitions between two states: a relief state and an obsessive state (Figure 1a). We assumed an

action to relieve anxiety as an option in the obsessive state, which could stochastically permit a

transition from the obsessive state to the relief state, at a certain cost. Such an action might

become compulsive in OCD, and we labeled it as a "compulsion" (Figure 1a). We combined any

other options in the obsessive state, which could relieve anxiety with lower probability without

any cost into the option "other" (Figure 1a). Even in the relief state, we sometimes experience

idle thoughts called intrusive thoughts[3]. Healthy people, as well as OCD patients experience

intrusive thoughts[3]. Therefore, an essential difference between OCD patients and healthy people

is whether they reinforce an abnormal reaction to intrusive thoughts[3]. Accordingly, we labeled

the transition from the relief state to the obsessive state "abnormal reaction to intrusive thoughts"

mainly seen in OCD[3] (Figure 1a). We also combined other options that maintain the relief state

into "other" (Figure 1a). We assumed that every stay in the obsessive state would produce a

negative outcome representing anxiety. No positive outcome was introduced in the state

transition diagram imitating obsessive compulsive symptoms (Figure 1a). Hence, maintaining

"other" in the relief state is clearly optimal. Normal reinforcement learning should reduce

abnormal reactions to intrusive thoughts. However, if the learning system has some defects, it

may fail to reduce the reaction to intrusive thoughts and may fall into a spiral of obsession and

compulsion. This is the discrepancy between explicit environmental understanding and implicit

learning, since agents fall into a spiral of obsession and compulsion even though they know that

they will experience only loss.

We implemented imbalanced learning with a reinforcement learning model separated into

reinforcement and punishment systems, which we call *a separate eligibility trace model*. In

general, the outcome of a certain choice is available after a certain delay. Therefore,

reinforcement and punishment should be assigned to recent choices within a certain time scale.

This is called credit assignment, which is implemented as eligibility traces in reinforcement learning[5]. Each trace represents the recent frequency of action in a specific state within a time scale determined by the trace decay factor, ν (see Online Methods). When a prediction error occurs, the choice probability of each action in each state is updated by the product of the eligibility trace and the prediction error (Figure 1b). Thus, a positive prediction error reinforces recent actions within a time scale, whereas a negative error punishes them. An eligibility trace can be implemented as a memory trace in each synapse[13-15]. In this case, the trace time scales for reinforcement and punishment should be separately controlled in direct and indirect pathways (Figure 1b; Online Methods). Theoretically, the trace factors in the two systems, $\nu^{\pm}$, should be balanced: $\nu^{+} = \nu^{-}$. However, separate neural implementations makes it difficult to maintain the balance perfectly. In our separate eligibility trace model, we assumed that an imbalance in the trace factors can occur: $\nu^{+} \neq \nu^{-}$.

To show whether imbalanced learning generates OCD-like behavior, we simulated actor-critic learning (Online Methods), a typical method of reinforcement learning models[5] in the state transition diagram (Figure 1a), incorporating a separate eligibility trace (Figure 1b). When imbalance in the trace factors was moderate, actor-critic learning reduced the reaction rate to intrusive thoughts, even when the simulation started at a high reaction rate to intrusive thoughts (Figure 1c). In contrast, extreme imbalance increased the reaction rate to intrusive thoughts such that it culminated in a spiral of repetitive obsession and compulsion (Figure 1d). Such OCD-like behavior was reduced by preventing the compulsion during the obsessive state (Figure 1d). This is the behavioral therapy of exposure and response prevention (ERP) and is one of the first-line treatments for OCD[16].

We identified the condition for the trace factors $\nu^{\pm}$ in which OCD-like behavior might

emerge by numerical simulations in a representative set of other model parameters (Figure 1e;

Online Methods). In addition to actor-critic learning (Figure 1e), we simulated SARSA (State-

Action-Reward-State-Action) and Q-learning (Figure 1e), other typical reinforcement learning

models[5]. The color map represents the fraction of 100 simulation runs in which abnormal

reactions to intrusive thoughts were reinforced at each pair of (ν^{+}, ν^{-}). We identified conditions

of OCD-like behavior in the region $\nu^{+} > \nu^{-}$ (Figure 1e). In such cases, the trace scale for

reinforcement was longer than that for punishment. We also identified conditions in which

behavioral therapy of ERP would not work and localized it to a region with a more robust

imbalance (Figure 1f).

We derived the theoretical conditions as functions of other parameters (Supplementary

Note). Each OCD patient presents a certain compulsive action for specific anxieties among many

types of anxiety. Therefore, corresponding parameters of the state transition (Figure 1a) should

differ by the kind of anxiety. Next, we derived conditions for learning parameters in which

OCD-like behavior emerges in some anxiety in actor-critic, SARSA, and Q-learning, and we

found the common condition that $\nu^{+} > \nu^{-}$ (Section 3.3 in the Supplementary Note). This result

suggests that a person with $\nu^{+} > \nu^{-}$ has a risk of OCD.

Figure 1. OCD-like behavior caused by imbalanced implicit learning. (a) State transition diagram imitating obsessive compulsive symptoms (b) Schema of the separate eligibility trace model. (c) Simulation of the separate eligibility trace model in the state transition diagram with a moderate imbalance in trace factors, $\nu^{\pm}$. Black polylines represent the temporal pattern of state transitions. Red and magenta triangles represent occurrences of compulsions and abnormal reactions to intrusive thoughts, respectively. The lower plots show temporal patterns of compulsion (red curve) and abnormal-reaction (magenta curve) rates on a logarithmic scale. (d) As in (c), but in the case of extreme imbalance. In the latter half, we demonstrated behavioral therapy by preventing the compulsion in the obsessive state (green bar). (e) Conditions of OCD-like behavior in the space of the trace factors, $\nu^{\pm}$ with different types of learning algorithms: actor-critic, SARSA, and Q-learning. The magenta color scale represents the percentage increase of the abnormal-reaction rate among 100 simulations around optimal behavior (zero reaction rate to intrusive thoughts). Solid blue lines represent the theoretically derived boundary

(Supplementary Note). The square and triangle represent the trace factors $v^{\pm}$ used in (c) and (d). (f) As in (e), but in this condition, behavioral therapy did not work.

Empirical evaluation of the theoretical model of OCD

To empirically test the suggestion that $v^+ > v^-$, derived from our computational model of OCD, we subjected patients with OCD (n = 45) and healthy controls (HCs) (n = 168) to a behavioral task. We excluded 3 OCDs due to Intelligence Quotient (IQ) < 80 using the Wechsler Adult Intelligence Scale - Third Edition (WAIS-III). Seventeen HCs were excluded for medical or experimental reasons. The subsequent analysis was conducted with 42 OCD patients and 151 HCs (Online Methods and Supplementary Table 1).

We used the delayed feedback task to evaluate the trace factors $v^{\pm}$, as in our previous research[17], except for presented stimuli (abstract cues). Briefly, participants chose one of two options displayed on a monitor by pressing a left or right button in each trial (Figure 2a). Each stimulus represented different outcomes (+10, +40, −10, or −40 yen) and a delay to the feedback (immediately after a button press or three trials later) (Figure 2b and c). For example, +40(0) represented a gain of 40 yen during the current trial (immediate reward), and −10(3) represented a loss of 10 yen after three trials (delayed punishment). Participants were not told about the stimulus-outcome associations shown in Figure 2b. They received money after the experiment in proportion to the total outcome obtained in the delayed feedback task. To maximize the total outcome, participants needed to learn appropriate stimulus choices by correctly assigning the credit of the current feedback to the recent choices that caused the current feedback. The difference in learning effects between immediate and delayed feedback reflected the trace factors, $v^{\pm}$.

Figure 2. Delayed feedback task. (a) Two abstract cues were displayed in each trial. When participants heard an auditory cue (beep), they chose one of the stimuli within 1 s. A single trial lasted 2.5 s. Sequences of stimulus pairs were pseudorandom. (b) Outcome-delay mapping for each stimulus. Mapping differed among participants. (c) An example of a delayed feedback task. If participants chose the stimulus with a delay [+40(3)] at trial *t*, that outcome was not displayed immediately. If they chose the stimulus with no delay [+40(0)] at trial *t*+3, the sum of the delayed and immediate outcomes was displayed at trial *t*+3 (in this case +80 yen).

Based on our simulation of OCD-like behavior with trace factors $v^+ > v^-$ (Figure 1e), patients with OCD showed impaired learning in stimuli with delayed feedback. To confirm the relationships between the parameters (v^+ and v^-) and the total outcome from stimuli with immediate or delayed feedback, we conducted a simulation (Online Methods). The total outcome

from stimuli with immediate feedback did not depend on the parameters, v^+ and v^-, except for extremely high values (Figure 3a). In contrast, we confirmed that more balanced and higher values of v^+ and v^- were needed to achieve a greater total outcome from stimuli with delayed feedback (Figure 3a). Therefore, the total outcome from stimuli with immediate or delayed feedback was compared between groups.

Consistent with our trace factor hypothesis, the OCD group showed impaired learning of stimuli with delays (Figure 3b). We used the ANOVA-type statistic (ATS) for nonparametric repeated measures ANOVA[18] with a within-participant factor of sessions (1–6 sessions) and a between-participant factor of groups (OCD and HC). There were significant interactions in learning stimuli with delays ($F(2.71, \infty)=5.35$, $p=0.0017$). Bonferroni-Holm–corrected post-hoc comparisons using the Brunner-Munzel test confirmed OCD < HC in session 4 (statistic = −3.72, $p_{adjusted} = 0.0017$), session 5 (statistic = −4.93, $p_{adjusted} < 0.001$), and session 6 (statistic = −3.94, $p_{adjusted} < 0.001$) (Figure 3b). In contrast, patients with OCD showed intact learning of stimuli with immediate feedback (Figure 3b). There were no significant interactions ($F(3.01, \infty)=1.04$, $p > 0.05$) and no main effect of groups ($F(1, 56.5)=2.57$, $p > 0.05$). To better clarify the effect of delay on learning, we conducted an ATS analysis of a within-participant factor of delays (immediate and delayed) and a between-participant factor of groups using the total outcome in sessions 5 and 6. We confirmed that there were significant interactions ($F(1, \infty)=5.17$, $p=0.023$); that is, a delay in receiving feedback impaired learning more in patients with OCD than in HC (Figure 3c).

Considering the therapeutic effect of serotonin reuptake inhibitors (SRIs) in OCD[18], SRIs may normalize impaired learning of stimuli with delayed feedback. To investigate effects of SRIs on learning, we conducted a correlation analysis between the dose equivalence of

SRIs[20] (Online Methods and Supplementary Table 2) and the total outcome from stimuli with

immediate or delayed feedback in sessions 5 and 6. Consistent with our prediction, SRI dose was

significantly correlated with the total outcome from stimuli with delayed feedback [Spearman's

rank correlation (27); r = 0.41, p = 0.027; Figure 3d], but not with immediate feedback

[Spearman's rank correlation (27); r = 0.061, p > 0.05; Figure 3d]. Since there was no significant

correlation between total outcome and depressive symptoms evaluated using the 17-item

Hamilton Depression Rating Scale (HDRS)[19] [Immediate: Spearman's rank correlation (26); r =

0.14, p > 0.05; Delayed: Spearman's rank correlation (26); r = -0.20, p > 0.05], a simple

antidepressant effect of SRIs could not explain this improved learning caused by SRIs.

        In summary, patients with OCD demonstrated the hypothesized impaired learning in

stimuli with delayed feedback and SRIs improved it in a dose-dependent manner.

Figure 3. Learning of stimuli with immediate or delayed feedback and its relationship with the dose
equivalence of SRIs. (a) The total simulated outcome from stimuli with immediate (left) or delayed
feedback (right) projected in v^+/v^- space. The color represents the amount of total outcome in sessions 5
and 6. (b) The total outcome from stimuli with immediate (left) or delayed feedback (right) (red cross, OCD
patients; blue circle, HCs). The line and colored area represent the mean and standard error. The horizontal
dashed line at 0 yen. *$p_{adjusted} < 0.05$ with Bonferroni-Holm–corrected post-hoc comparisons of the simple

main effects of group (OCD patients < HCs). (c) Significant interactions between delays (immediate and delayed) and groups (red, OCD patients; blue, HCs) in the total outcome in sessions 5 and 6. The line and error bar represent the mean and standard error. The horizontal dashed line at 0 yen. (d) Correlation between the SRI dose and the total outcome in sessions 5 and 6 (left, immediate feedback; right, delayed feedback). The line and colored areas are the regression line and the 95% confidence interval (+: each OCD patient taking SRIs). (e) As in (b), but in the case of simulated data using estimated parameters (red cross, simulated OCD patients; blue circle, simulated HCs). (f) As in (c), but in the case of simulated data using estimated parameters (red simulated OCD patients; blue, simulated HCs).

Estimation of model parameters from empirical data

We fitted behavioral data to the actor-critic learning model using the learning rate (α),

exploration-exploitation degree (β), and separation of the eligibility trace for reinforcement and

punishment ($\nu^{\pm}$) (Online Methods). Because we were specifically interested in the individual

variance represented by reinforcement learning parameters, we fitted parameters in data from

each participant independently, using maximum a posteriori estimation rather than pooled data of

a group[20].

We simulated behavioral data of each group with the estimated parameters to validate the

actor-critic learning model with separate eligibility traces (see Online Methods). We evaluated

whether all-important behavioral characteristics are qualitatively and quantitatively captured [21].

We confirmed that the specific impairment in learning from stimuli with delayed feedback in the

simulated OCD group (Figure 3e and 3f) resembles the experimental data (Figure 3b and 3c).

Therefore, our model actually captured essential features of human behaviors in this task.

To clarify the ν^{+}/ν^{-} distribution of each group, we projected estimated parameters from

each participant to ν^{+}/ν^{-} space and visualized them using kernel density estimation (Figure 4a).

Consistent with our computational model simulation of OCD, patients with OCD were

distributed in the $\nu^{+} > \nu^{-}$ imbalanced area, whereas the distribution was balanced in HCs

(Figure 4a). To compare the ν^{+}/ν^{-} distribution among groups, we confirmed the multivariate

homogeneity of group dispersions [$F_{(1, 191)} = 0.25$, $p = 0.62$] and applied permutational

multivariate analysis of variance (PERMANOVA). The ν^+/ν^- distribution differed significantly

between groups [PERMANOVA: F (1, 191) = 5.99, p = 0.0060] (Figure 4a). Parameters α and β

did not differ significantly between groups (Brunner-Munzel test, p > 0.05). Among OCD

participants taking SRIs (n = 29), the equivalent dose of SRIs was significantly correlated with

the imbalanced settings of ν ($\nu^+ - \nu^-$) [Spearman's rank correlation (27); r = −0.48, p = 0.0086;

Figure 4b]. That is, higher doses of SRIs normalized the imbalanced settings of ν in participants

with OCD.

Although the ν^+/ν^- distribution was more balanced in HCs compared with OCD

patients, we found some diversity among HCs. We conducted clustering analysis using

hierarchical density-based spatial clustering of applications with noise (HDBSCAN)[22] to

evaluate diversity among HCs. HDBSCAN revealed two clusters of HCs: a balanced ν cluster

(Figure 4a, blue circle; n = 83) and an imbalanced ν cluster (Figure 4a, white circle; n = 57), with

11 HCs that were not clustered. Obsessive-compulsive trends and a propensity to adhere to fine-

grained details were compared between clusters using five subscales of the Padua Inventory (PI;

"Checking", "Dirt", "Doubt", "Impulse", "Precision")[23] and the Attention to Detail subscale of

the Autism Quotient (AQ)[24], respectively. The imbalanced HC cluster showed significantly

higher PI Checking scores (Brunner-Munzel test, statistic = −2.28, p = 0.012), PI Dirt scores

(Brunner-Munzel test, statistic = −1.71, p = 0.045), and AQ Attention to Detail scores (Brunner-

Munzel test, statistic = −2.82, p = 0.0028) than the balanced HC cluster. Significant differences

were still found in the PI Checking score and the AQ Attention to Detail score after false

discovery rate correction for multiple comparisons among scales (Figure 4c) (PI Checking,

$p_{adjusted}$ = 0.025; AQ Attention to Detail, $p_{adjusted}$ = 0.011).

Figure 4. Estimated parameters and their relationships with clinical characteristics. (a) Distribution of participants in each group projected in ν^+/ν^- space using the kernel density estimation (red, OCD patients; blue, HCs). The horizontal axis represents ν^+ and the vertical axis represents ν^-. The diagonal line represents the balanced line in the ν^+/ν^- space. Blue and white circles in the right figure represent balanced and imbalanced clusters in HCs, respectively. Other participants were depicted using +. (b) Significant correlation between the SRI dose and $\nu^+ - \nu^-$ [Spearman's rank correlation (27); r = −0.48, p = 0.0086]. The line and colored areas are the regression line and the 95% confidence interval (+: each OCD patient taking SRIs). The dashed line represents the balanced settings of $\nu^\pm$. (c) PI Checking and AQ Attention to Detail scores in the balanced and imbalanced clusters in HCs. The half-violin, box, and strip plots represent the probability density, interquartile range and median, and raw data, respectively. Blue and white circles represent balanced and imbalanced clusters, respectively, whereas * and ** represent significant differences (Brunner-Munzel test, * statistic = −2.28, p = 0.012; ** statistic = −2.82, p = 0.0028).

Discussion

This study showed that imbalanced implicit learning between reinforcement and punishment can

induce a spiral of repetitive obsession and compulsion, even despite explicit environmental

understanding. We constructed a computational model of OCD using a separate eligibility trace

model (Figure 1a–e) and found extremely imbalanced trace factors $v^+ > v^-$ in OCD (Figure 4a).

In addition, effects of behavioral therapy (ERP) and psychotropic medication (SRIs), which are

the first-line treatments for OCD, were reflected in our computational model (Figure 1d) and

behavioral results (Figure 3d and 4b), respectively.

While the theoretical framework of the eligibility trace has long been conceptualized in

the field of reinforcement learning[5], empirical evidence for it has been reported only relatively

recently[13, 25-27]. The latest theories regarding synaptic plasticity have proposed that co-activation

of pre- and postsynaptic neurons sets a flag at the synapse (eligibility trace) that leads to a weight

change only if additional factors, i.e., reinforcement or punishment, are present while the flag is

set[15]. These additional factors could be implemented by the phasic activity of some

neuromodulators, such as dopamine, which is supposed to represent the prediction error[6].

Dopamine-dependent synaptic plasticity in corticostriatal synapses, which underlies

reinforcement learning, shows opposing dopamine dependence for D1 and D2 neurons[13, 28].

Although detailed mechanisms concerning how trace time scales for reinforcement and

punishment are modulated remain unclear, our theoretical consideration of the state transition

diagram imitating obsessive compulsive symptoms showed that imbalanced trace factors $v^+ >$

v^- could lead to a condition of repetitive choices of abnormal reaction to intrusive thoughts and

neutralization of anxiety, similar to obsession and compulsion in OCD (Figure 1a–e). Our

experimental data from OCD patients and HCs strongly support predictions from our

computational model, i.e., apparent impairment in learning with delayed feedback (Figure 3b, 3c,

and 3d) and extremely imbalanced trace factors $\nu^+ > \nu^-$ in OCD (Figure 4a). It is noteworthy

that imbalanced trace factors $\nu^+ > \nu^-$ are quite convincing because the conventional

pathophysiological model of OCD suggests excess tone in the direct pathway over the indirect

pathway[10-12], which are supposed to be related to ν^+ and ν^-, respectively.

There have been several computational models of OCD, all of which focused primarily

on compulsive behaviors[1, 2, 29]. Compulsive actions are thought to be reinforced by the rewarding

effect of relief from anxiety[4]. Excessive anxiety induces excessive reinforcement of compulsion

leading to habit formation[2]. This excessively habitual compulsion is considered a cause of OCD[2, 30].

However, obsessive thoughts that drive anxiety also increase with the severity of OCD

symptoms[4]. Although the essence of OCD symptoms is the abnormal repetition of obsession and

compulsion[4], thus far there has been no unified model to explain growth of both obsession and

compulsion. Our unified model successfully represents the vicious circle of obsession and

compulsion, which are the phenomenological characteristics of OCD[4] (Figure 1a–1e). Moreover,

this model extends our understanding of therapeutic effects of first-line treatments of OCD.

While ERP seems to promote appropriate choices to prevent abnormal reactions to intrusive

thoughts, even under imbalanced trace factors (Figure 1d), SRIs resolve the imbalance itself

(Figure 3d and 4b). In practice guidelines for treatment of OCD[16], combination therapy of ERP

and SRI is recommended if patients do not respond to ERP monotherapy. We identified such a

condition where ERP fails in our computational model (Figure 1f). SRI add-on therapy in this

ERP-refractory condition can be viewed as assisting normalization of trace factors. Regarding

relationships between the neuromodulator, serotonin, and time scales of the eligibility trace,

serotonin seems to modulate synaptic plasticity in many brain regions directly or indirectly

through its regulatory effects on other neuromodulatory systems[15, 25, 31, 32]. In our previous study,

we also found similar modulatory effects of the trace factor time scales, that is, depletion of the

serotonin precursor, tryptophan, increased the $\nu^+ > \nu^-$ imbalance compared with the control

condition[17]. There is still abundant scope for further research aimed at illuminating how

serotonin modulates the time scale of trace factors.

Beyond the range of clinical phenotypes seen in patients with OCD, a broader continuum

of the obsessive-compulsive trait is also observed in behavioral characteristics. Specifically, the

imbalanced ($\nu^+ > \nu^-$) HC cluster showed a significantly greater propensity for Checking and

Attention to Detail (Figure 4c). These results increase the reliability of our clinical findings and

support the generalizability of our findings to the broader population on the obsessive-

compulsive continuum. Further study with a greater focus on OCD patients and their unaffected

healthy first-degree relatives should be performed to evaluate the potential of our findings as an

endophenotype of OCD[33].

This study provided evidence that one's own imbalanced implicit learning between

reinforcement and punishment can induce a vicious circle of obsession and compulsion even

despite explicit environmental understanding. The discrepancy between explicit environmental

knowledge and implicit learning could lead to ego-dystonic anxiety[2]. We believe that these

results are not limited to OCD, but capture characteristics of an obsessive-compulsive type of

anxiety widely present in the general population (Figure 4a and 4c). Anxiety can manifest itself

in many different forms. In some situations, anxiety may motivate people to take action or to

strive to achieve a goal. In other situations, people may experience various forms of anxiety as a

symptom of an anxiety disorder, such as social anxiety disorder or generalized anxiety disorder.

Because we cannot cover all forms of anxiety in this research, further study should be performed with a greater focus on constructing computational models of other types of anxiety.

Conclusion

Although we think that we always make rational decisions, our computational model proves that we sometimes implicitly reinforce maladaptive behaviors and may feel intense anxiety. Psychiatric symptoms are often regarded as mental alterations that are not directly quantifiable, but they can be directly assessed by creating appropriate computational models. Although it is currently difficult to identify treatment-resistant patients based upon their clinical symptoms, our computational model suggests that patients with highly imbalanced trace scale parameters may not respond to behavioral therapy alone. These results suggest that our findings could one day be applied to appropriate selection of OCD treatments. In addition, psychiatric symptoms have been regarded in recent years as a symptom dimension common to various mental diseases rather than to a specific disease. In this study, we focused on patients with OCD and healthy participants, but our approach could be applied to assess the obsessive-compulsive dimension in various populations[34]. Future research is needed to address these hypotheses in prospective longitudinal cohorts or in larger cohorts with various psychiatric symptoms.

Online Methods

Separate eligibility trace model

Eligibility traces determine the weight assignment of outcome prediction errors for recent actions (Figure 1b). If eligibility traces for positive and negative prediction errors are implemented in

distinct neural systems, the respective eligibility traces $\eta_{as}^+(t)$ and $\eta_{as}^-(t)$ at time step t for action

a ($a = 0, 1, \ldots$) in state s ($s = 0, 1, \ldots$) obey the following equation[5],

$$\eta_{as}^\pm(t+1) = \nu^\pm \eta_{as}^\pm(t) + X_{as}(t),$$

where $X_{as}(t) = 1$ when action a is chosen in state s at time step t and $X_{as}(t) = 0$ otherwise. The

factors $\nu^\pm$ ($0 < \nu^\pm < 1$) determine the decaying time scales of the eligibility traces. The policy

parameters $q_{as}(t)$ ($a = 0, 1, \ldots$) determine choice probability in state s as a soft-max function,

$$P(a|s) = e^{\beta q_{as}(t)} \Big/ \sum_{\acute{a}} e^{\beta q_{\acute{a}s}(t)},$$

where β represents the degree of the exploration-exploitation balance. Each policy parameter is

updated as below,

$$q_{as}(t+1) = q_{as}(t) + \alpha[\eta_{as}^+(t)\max\{0, \varepsilon(t)\} + \eta_{as}^-(t)\min\{0, \varepsilon(t)\}\,],$$

where $\varepsilon(t)$ denotes the outcome prediction error and $\max\{0, \varepsilon(t)\}$ and $\min\{0, \varepsilon(t)\}$ represent

positive and negative components of the prediction error, respectively (Figure 1b). Theoretically,

the trace factors $\nu^\pm$ should be balanced as $\nu^+ = \nu^-$. However, separate neural implementation

makes it difficult to maintain the balance perfectly. Here, we assumed that the balance could be

broken: $\nu^+ \neq \nu^-$.

The outcome prediction error $\varepsilon(t)$ is determined as a function of the current outcome

$r(t)$ and the policy parameters $\{q_{as}\}$. The form depends on learning algorithms. In actor-critic

learning, the outcome prediction is based on the state value $v_s = \sum_a q_{as}$, and the prediction error

$\varepsilon(t) = r(t) + \gamma v_{s(t+1)} - v_{s(t)}$, where $r(t)$ denotes the current outcome and $s(t)$ and $s(t+1)$

denote the current and next states. The parameter γ denotes the discount factor that determines

the weight of future prediction. In SARSA and Q-learning, the outcome prediction is based on

the action value estimated as each policy parameter q_{as}, and the prediction error $\varepsilon(t) = r(t) +$

1    $\gamma V_{s(t+1)} - q_{a(t)s(t)}$. The difference is in the term of the next state value: $V_{s(t+1)} = q_{a(t+1)s(t+1)}$

2   in SARSA and $V_{s(t+1)} = \max_{a} q_{as(t+1)}$ in Q-learning.

State transition diagram imitating obsessive compulsive symptoms

5   We described mental states involved in obsessive-compulsive symptoms in stochastic transitions

6   between two states: a relief state (s=0) and an obsessive state (s=1) (Figure 1a). We assumed two

7   options in each state: "compulsion" (a=1) and "other" (a=0) in the obsessive state, and "abnormal

8   reaction to intrusive thoughts" (a=1) and "other" (a=0) in the relief state. We defined a matrix b

9   to determine the state-transition probabilities,

$$P(s(t+1) = 1|a(t) = i, s(t) = j) = b_{ij},$$

$$P(s(t+1) = 0|a(t) = i, s(t) = j) = 1 - b_{ij}.$$

12   We assumed that every stay in the obsessive state produced a negative outcome normalized to

13   -1. The relative cost of compulsion was denoted as c. Although b_{00}>0 allows passive anxiety,

14   we focused on b_{00}=0 in the main article. More general cases, including passive anxiety, are

15   considered in the Supplementary Note.

Demonstration of OCD-like behavior

For the example in Figure 1, we fixed the parameters of the state transition as b_{00}=0,

b_{10}=1, b_{01}=0.9, b_{11}=0.5, and c=0.01 and the learning parameters as α=0.1, β=1, $\gamma = 0.5$, and

v^{+}=0.8, except for v^{-}. For Figure 1c, the trace factor for punishment was set as v^{-}=0.6 and the

initial values of variables as $\eta_{as}^{\pm}(0) = q_{as}(0) = s(0) = 0$. The simulation consisted of 100,000-

time steps. For Figure 1d, we set the trace factors as v^{-}=0.1 and the initial values as $\eta_{as}^{\pm}(0) =$

$q_{a1}(0) = s(0) = 0$, $q_{00}(0) = 3$, and $q_{10}(0) = -3$, to show that reinforcement of the abnormal

reaction to intrusive thoughts started from even a low reaction rate. After 50,000-time steps,

compulsion ($a=1$ in $s=1$) was always prevented, and other ($a=0$ in $s=1$) was forced regardless of

the choice probability, demonstrating the behavioral therapy of ERP.

For Figure 1e and f, we evaluated the fraction of 100 simulation runs in which the

reaction rate to intrusive thoughts was reinforced from a low reaction rate on 40×40 grids in

(ν^+, ν^-) space. In each simulation, 200 instances of forced abnormal reaction were intermittently

caused in the relief state because spontaneous abnormal reactions scarcely occurred at a low

reaction rate. The simulation was judged to be abnormal-reaction-reinforced if the choice

probability of reaction to intrusive thoughts became larger than the initial value. The initial

values of variables were set as $\eta_{as}^\pm(0) = q_{a1}(0) = s(0) = 0$, $q_{00}(0) = 5$, and $q_{10}(0) = -5$.

Each forced abnormal reaction was caused when $\eta_{as}^\pm(t)$ sufficiently approached zero in the relief

state ($\eta_{as}^\pm(t) < 0.001$).

Participants

Forty-five patients with OCD and 168 HCs participated in the behavioral task. The Medical

Committee on Human Studies at Kyoto Prefectural University of Medicine (KPUM) and the

Ethics Committee at Advanced Telecommunications Research Institute International (ATR)

approved all procedures in this study. All participants gave written informed consent after

receiving a complete description of the study. All methods were carried out following approved

guidelines and regulations. Trained, experienced clinical psychiatrists and psychologists assessed

all participants.

All patients were primarily diagnosed using the Structured Clinical Interview for DSM-

IV Axis I Disorders-Patient Edition (SCID)[35]. Experienced clinical psychiatrists or psychologists

applied the Yale-Brown Obsessive-Compulsive Scale (Y-BOCS)[36], the 17-item Hamilton

Depression Rating Scale (HDRS)[19], and Wechsler Adult Intelligence Scale - Third Edition

(WAIS-III) for evaluation of obsessive-compulsive symptoms, depressive symptoms, and IQ in

patients with OCD, respectively. There was one missing value for the HDRS. Exclusion criteria

were 1) a cardiac pacemaker or other metallic implants or artifacts; 2) significant disease,

including neurological diseases, disorders of the pulmonary, cardiac, renal, hepatic, or endocrine

systems, or metabolic disorders; 3) prior psychosurgery; 4) IQ < 80 using WAIS-III; 5) DSM-IV

diagnosis of mental retardation and pervasive developmental disorders based on a clinical

interview and psychosocial history, and 6) pregnancy. We excluded 3 OCDs due to IQ < 80

using the WAIS-III. Seventeen HCs were excluded for medical or experimental reasons. The

subsequent analysis was conducted in 42 OCD patients and 151 HCs. There were no significant

differences in age, sex, or handedness (Supplementary Table 1). To the extent possible, we

selected patients without a current DSM-IV Axis I diagnosis of any significant psychiatric illness

except OCD. Only two patients with trichotillomania, one patient with a tic disorder, two

patients with panic disorders, and one patient with bulimia nervosa were included as patients

with comorbidities. Handedness was classified based on a modified 25-item version of the

Edinburgh Handedness Inventory. Thirteen patients with OCD were drug-free for all types of

psychotropic medication. SRI and imipramine equivalent doses[37] of the remaining 29 patients are

summarized in Supplementary Table 2.

Behavioral task and statistical analysis

The delayed feedback task was similar to our previous research[17], except for the

presented stimuli. Participants chose one of the two options (abstract cues) displayed on the

screen by pressing a left or right button in each trial within 1 s after an auditory cue (Figure 2a).

Depending on the selected stimulus, monetary feedback with different outcomes (+10, +40, −10,

or −40 yen) was displayed either immediately after the button press or three trials later (Figure

2b and c). We did not offer monetary feedback for the first five trials because participants can

learn the relationships between stimuli and outcomes or delays quickly. If participants pressed a

button before the auditory cue or more than 1 s passed without any button press, −50 yen was

displayed as a punishment. Such trials were considered error trials, and delayed outcomes were

not considered.

Two abstract cues were displayed at each trial side by side on the screen (Figure 2a). We

prepared 16 pairs (Figure 2b), counterbalancing the number of appearances of each stimulus. The

16 pairs of stimuli were presented in pseudorandom order. Each pair was presented as the

scheduled number of trials in each session: each of six pairs [+10(0) vs. +40(0); +10(3) vs.

+40(3); −10(0) vs. −40(0); −10(3) vs. −40(3); +10(0) vs. +40(3); −10(0) vs. −40(3)] was

presented in 10 trials during a single session, and each of 10 pairs [+40(0) vs. +40(3); +10(0) vs.

+10(3); −10(0) vs. −10(3); −40(0) vs. −40(3); +10(3) vs. +40(0); −10(3) vs. −40(0); +10(0) vs.

−10(0); +40(0) vs. −40(0); +10(3) vs. −10(3); +40(3) vs. −40(3)] was presented in five trials

during a single session. Each participant performed 110 trials during a single session and six

sessions in each experiment. About 28 min was required for participants to complete six

sessions. At the beginning of each session, the session number was displayed on the screen for

2.5 s. Before the task, each participant practiced the test session under the same task settings

except for stimuli, and we confirmed that all participants understood the task set.

The total outcome (except for punishments related to button press errors), reaction time,

and the number of error trials were compared between the OCD and HC groups. Patients with

OCD showed a significantly lower total monetary outcome than HCs [median (interquartile

range): OCD patients, 2705 (1510–3542) yen; HCs, 4700(3095–5955) yen; Brunner-Munzel test,

statistic = 5.31, p < 0.001], whereas there were no group differences in reaction time [median

(interquartile range): OCD patients, 545.9 (503.4–614.8) ms; HCs, 522.6 (475.9–574.8) ms;

Brunner-Munzel test, p > 0.05)] and numbers of button press errors [median (interquartile

range): OCD patients, 4.5 (2.25–14.75); HC, 4.0 (2.0–8.0); Brunner-Munzel test, p > 0.05).

These results suggested impaired learning in OCD individuals compared with HCs.

Based on our hypothesis of trace factors v^+/v^-, patients with OCD were expected to

show impaired learning in stimuli with delayed feedback. To confirm relationships between the

parameters (v^+ and v^-) and the total outcome from stimuli with immediate or delayed feedback,

we conducted the simulation. We evaluated the total outcome from stimuli with immediate or

delayed feedback using actor-critic learning with separate eligibility traces in the same

experimental settings as the actual experiment. We created 500 simulated cases using each

combination of v^+ and v^- (0-0.99 in 0.01 increments). Values of α and β were fixed as the

median values of HC estimated parameters (α, 0.0011; β, 2.96). We projected the mean values

of total simulated outcome from stimuli with immediate or delayed feedback in v^+/v^- space

(Figure 3a).

An ANOVA-type statistic (ATS) for nonparametric repeated measures ANOVA[18] was

conducted to clarify between-group differences. Bonferroni-Holm–corrected post-hoc

comparisons using Brunner-Munzel test were conducted when needed. Therapeutic effects of

SRIs were evaluated using Spearman's rank correlation between the dose equivalence of SRIs[20]

(Supplementary Table 2) and the total outcome from stimuli with immediate or delayed feedback

in sessions 5 and 6.

Model validation and parameter estimation for the delayed feedback task

We fitted behavioral data with actor-critic learning with separate eligibility traces for

positive and negative prediction errors. We defined 16 states for possible pairs of stimuli

presented at each trial. Available actions at each state involved choosing alternative stimuli. To

facilitate model-fitting in the face of limited experimental data from each participant, we used

regularizing priors that favored realistic values and maximum a posteriori estimation rather than

maximum likelihood estimation[20]. The learning rate α and trace factor ν were constrained to the

range of $0 \leq \alpha \leq 0.95$ and $0 \leq \nu \leq 0.95$ with a uniform prior. The exploration-exploitation degree

β was constrained to the range of $0 \leq \beta \leq 100$ with a gamma (2,3) prior distribution that favored

relatively lower values. We fixed the discount factor $\gamma = 0$, because the term of the next state

value was just noise in our delayed feedback task in which the state of each trial was randomly

selected. We optimized parameters by minimizing the negative log posterior of the data with

different parameter settings using the hyperopt package[38]. Likelihood ratio tests to assess the

contribution of the additional parameter in our model (four parameters: $\alpha, \beta, \nu^+, \nu^-$) compared

with the standard actor-critic learning model (three parameters: α, β, ν) showed that the

additional parameter was justified in 133 of 193 participants (X^2 test with one degree of freedom,

$p < 0.05$). To further validate the actor-critic learning model with separate eligibility traces, we

simulated the behavioral data of each group using the estimated parameters. We randomly chose

100 participants from each group with duplicates and simulated the total outcome using the

estimated parameters of the selected participants. We illustrated the mean and standard errors of

the simulated total ouctome (Figure 3e, f) and compared them with the experimental results

(Figure 3b and 3c). We confirmed that the specific impairment in learning from stimuli with

delayed feedback in the simulated OCD group (Figure 3e and 3f) is similar to the experimental

data (Figure 3b and 3c). Because our target model was validated with behavioral data, we

applied the separate eligibility trace model in the substantive analysis.

We evaluated the performance of parameter estimation in actor-critic learning with

separate eligibility traces. We created 200 simulation data points using a model with the

following parameters: α, 0.1 ± 0.05 (mean $\pm$ SD); β, 1 ± 0.2; ν^+ and ν^-, randomly selected

within 0.01–0.95. Parameter estimation was quite accurate[39] (Supplementary Figure 1).

Specifically, Pearson's r and the mean absolute error between the true and estimated ν^+ or ν^-

were 0.99 and 0.03, respectively (Supplementary Figure 1).

We compared the ν^+/ν^- distribution in our model between groups using PERMANOVA

with the ADNOIS function and 10,000 permutations using the Euclidean distance implemented

in the statistical package R[40]. The multivariate homogeneity of group dispersions was confirmed

with the *betadisper* function with 10,000 permutations in R[40]. The learning rate α and inverse

temperature β were compared using the Brunner-Munzel test. To further investigate

relationships between clinical characteristics and estimated parameters in the HC group, we

conducted clustering analysis using HDBSCAN[22]. We detected two clusters (balanced cluster, n

= 83; imbalanced cluster, n = 57; the remaining 11 HCs were not clustered) and evaluated their

obsessive-compulsive trait using the five PI subscales: "Checking", "Dirt", "Doubt", "Impulse",

"Precision"[23]. In addition, the propensity to adhere to fine-grained details was evaluated using

the Attention to Detail subscale of AQ[24]. There were 14 missing values in PI (n = 126) and 1

missing value in AQ (n = 140). Therapeutic effects of SRIs were evaluated using Spearman's

rank correlation between the SRI dose and imbalanced settings of ν ($\nu^+ - \nu^-$).

References

1. Vaghi, M.M.*, et al.* Compulsivity Reveals a Novel Dissociation between Action and Confidence. *Neuron* (2017).
2. Gillan, C.M. & Robbins, T.W. Goal-directed learning and obsessive-compulsive disorder. *Philos Trans R Soc Lond B Biol Sci* **369** (2014).
3. Salkovskis, P.M. Understanding and treating obsessive-compulsive disorder. *Behav Res Ther* **37 Suppl 1**, S29-52 (1999).
4. Cavedini, P., Gorini, A. & Bellodi, L. Understanding obsessive-compulsive disorder: focus on decision making. *Neuropsychol Rev* **16**, 3-15 (2006).
5. Sutton, R.S. & Barto, A.G. Reinforcement Learning: An Introduction. *MIT Press, Cambridge, MA.* (1998).
6. Schultz, W., Dayan, P. & Montague, P.R. A neural substrate of prediction and reward. *Science* **275**, 1593-1599 (1997).
7. Graybiel, A.M. The basal ganglia. *Curr Biol* **10**, R509-511 (2000).
8. Kravitz, A.V., Tye, L.D. & Kreitzer, A.C. Distinct roles for direct and indirect pathway striatal neurons in reinforcement. *Nat Neurosci* **15**, 816-818 (2012).
9. Nonomura, S.*, et al.* Monitoring and updating of action selection for goal-directed behavior through the striatal direct and indirect pathways. *Neuron* **99**, 1302-1314. e1305 (2018).
10. Pauls, D.L., Abramovitch, A., Rauch, S.L. & Geller, D.A. Obsessive-compulsive disorder: an integrative genetic and neurobiological perspective. *Nat Rev Neurosci* **15**, 410-424 (2014).
11. Sakai, Y.*, et al.* Corticostriatal functional connectivity in non-medicated patients with obsessive-compulsive disorder. *Eur Psychiatry* **26**, 463-469 (2011).
12. Abe, Y.*, et al.* Hyper-influence of the orbitofrontal cortex over the ventral striatum in obsessive-compulsive disorder. *Eur Neuropsychopharmacol* **25**, 1898-1905 (2015).
13. Yagishita, S.*, et al.* A critical time window for dopamine actions on the structural plasticity of dendritic spines. *Science* **345**, 1616-1620 (2014).
14. Izhikevich, E.M. Solving the distal reward problem through linkage of STDP and dopamine signaling. *Cereb Cortex* **17**, 2443-2452 (2007).
15. Gerstner, W., Lehmann, M., Liakoni, V., Corneil, D. & Brea, J. Eligibility Traces and Plasticity on Behavioral Time Scales: Experimental Support of NeoHebbian Three-Factor Learning Rules. *Frontiers in neural circuits* **12**, 53 (2018).
16. Koran, L.M.*, et al.* Practice guideline for the treatment of patients with obsessive-compulsive disorder. *Am J Psychiatry* **164**, 5-53 (2007).
17. Tanaka, S.C.*, et al.* Serotonin affects association of aversive outcomes to past actions. *J Neurosci* **29**, 15669-15674 (2009).
18. Noguchi, K., Gel, Y.R., Brunner, E. & Konietschke, F. nparLD: an R software package for the nonparametric analysis of longitudinal data in factorial experiments. *Journal of Statistical software* **50** (2012).
19. Hamilton, M. Development of a rating scale for primary depressive illness. *The British journal of social and clinical psychology* **6**, 278-296 (1967).
20. Daw, N.D. Trial-by-trial data analysis using computational models. *Decision making, affect, and learning: Attention and performance XXIII* **23**, 3-38 (2011).
21. Wilson, R.C. & Collins, A.G.E. Ten simple rules for the computational modeling of behavioral data. *eLife* **8**, e49547 (2019).
22. Campello, R.J., Moulavi, D., Zimek, A. & Sander, J. Hierarchical density estimates for data clustering, visualization, and outlier detection. *ACM Transactions on Knowledge Discovery from Data (TKDD)* **10**, 5 (2015).
23. Sugiura, Y. & Tanno, Y. *Self-report inventory of obsessive-compulsive symptoms: Reliability and validity of the Japanese version of the Padua Inventory* (2000).
24. Wakabayashi, A., Tojo, Y., Baron-Cohen, S. & Wheelwright, S. [The Autism-Spectrum Quotient (AQ) Japanese version: evidence from high-functioning clinical group and normal adults]. *Shinrigaku Kenkyu* **75**, 78-84 (2004).
25. He, K.*, et al.* Distinct Eligibility Traces for LTP and LTD in Cortical Synapses. *Neuron* **88**, 528-538 (2015).
26. Brzosko, Z., Schultz, W. & Paulsen, O. Retroactive modulation of spike timing-dependent plasticity by dopamine. *eLife* **4**, e09685 (2015).

27. Brzosko, Z., Zannone, S., Schultz, W., Clopath, C. & Paulsen, O. Sequential neuromodulation of Hebbian plasticity offers mechanism for effective reward-based navigation. *eLife* **6**, e27756 (2017).

28. Shen, W., Flajolet, M., Greengard, P. & Surmeier, D.J. Dichotomous dopaminergic control of striatal synaptic plasticity. *Science* **321**, 848-851 (2008).

29. Hauser, T.U.*, et al.* Increased decision thresholds enhance information gathering performance in juvenile Obsessive-Compulsive Disorder (OCD). *PLoS Comput Biol* **13**, e1005440 (2017).

30. Gillan, C.M. & Sahakian, B.J. Which is the driver, the obsessions or the compulsions, in OCD? *Neuropsychopharmacology* **40**, 247-248 (2015).

31. Brzosko, Z., Mierau, S.B. & Paulsen, O. Neuromodulation of Spike-Timing-Dependent Plasticity: Past, Present, and Future. *Neuron* **103**, 563-581 (2019).

32. Cavaccini, A.*, et al.* Serotonergic Signaling Controls Input-Specific Synaptic Plasticity at Striatal Circuits. *Neuron* **98**, 801-816 e807 (2018).

33. Chamberlain, S.R. & Menzies, L. Endophenotypes of obsessive-compulsive disorder: rationale, evidence and future potential. *Expert Rev Neurother* **9**, 1133-1146 (2009).

34. Gillan, C.M.*, et al.* Comparison of the Association Between Goal-Directed Planning and Self-reported Compulsivity vs Obsessive-Compulsive Disorder Diagnosis. *JAMA psychiatry*, 1-10 (2019).

35. First, M.B., Spizer, R.L., Gibbon, M. & Williams, J.B.W. Structures Clinical Interview for Axis I DSM-IV Disorders-Patient Edition (CID-I/P). New York: Biometrics Research Department, New York State Psychiatric Institute. (1994).

36. Nakajima, T.*, et al.* Reliability and validity of the Japanese version of the Yale-Brown Obsessive-Compulsive Scale. *Psychiatry Clin Neurosci* **49**, 121-126 (1995).

37. Inada, T. & Inagaki, A. Psychotropic dose equivalence in Japan. *Psychiatry Clin Neurosci* **69**, 440-447 (2015).

38. Bergstra, J.S., Bardenet, R., Bengio, Y. & Kégl, B. Algorithms for hyper-parameter optimization. in *Advances in neural information processing systems* 2546-2554 (2011).

39. Yazdani, S., Vahabie, A.-H., Araabi, B.N. & Ahmadabadi, M.N. Better than maximum likelihood estimation of model-based and model-free learning style. *bioRxiv*, 296335 (2018).

40. Anderson, M.J., Walsh, D.C.I., Robert Clarke, K., Gorley, R.N. & Guerra-Castro, E. Some solutions to the multivariate Behrens-Fisher problem for dissimilarity-based analyses. *Australian & New Zealand Journal of Statistics* **59**, 57-79 (2017).

Data Availability

Patient data supporting conclusions of this paper are not publicly available because they contain

information that could compromise research participant privacy or consent. The theoretical

derivation of our computational model is in the Supplementary Note along with the MATLAB

source code.

Acknowledgments

All authors had full access to data used in this study. The corresponding author, S. C. Tanaka,

assumes responsibility for data integrity and accuracy of the data analysis. This study was mainly

supported by JSPS KAKENHI Grant Number JP21H05172 (S.C. Tanaka). The following partly

supported this study: "Research and development of technology for enhancing functional

recovery of elderly and disabled people based on non-invasive brain imaging and robotic

assistive devices", the Commissioned Research of National Institute of Information and

Communications Technology (NICT), JAPAN (S.C. Tanaka); JSPS KAKENHI Grant Number

JP16K01958 and JP16H06396 (S.C. Tanaka); JP16H01516 and JP18H05524 (Yutaka Sakai);

16H01512 and the Sakamoto Research Foundation of Psychiatric Diseases (Yuki Sakai); the

Joint Usage/Research Center ("Behavioral economics") of Institute of Social and Economic

Research, Osaka University (J. Narumoto).

Author Contributions

Yuki S, Yutaka S, JN, and SCT designed the study. Yutaka S developed the theory and

performed computational modeling. Yuki S, YA, JN, and SCT collected the data. Yuki S

conducted the computational modeling and statistical analysis of the data. Yuki S and Yutaka S

1   wrote the manuscript, which all authors edited. SCT acquired funding to support theory

2   development and data analysis.

4   **Competing Interests Statement**

5   The authors declare no competing financial interests.

1 **Supplementary information**

2
3 Supplementary Figure 1. Parameter estimation. The horizontal and vertical axes represent the

4 true and estimated parameters, respectively. The line and colored areas are the regression line

5 and the 95% confidence interval, respectively.

1 **Supplementary Table 1. Demographic information about the participants.**

2 All demographic distributions are matched among groups (p > 0.05).

	OCD (n = 42)	Healthy controls (n = 151)
Demographic characteristics		
Age, years	33.2 ± 9.8	29.4 ± 10.3
Male sex, No.	21	99
Clinical characteristics		
With/without Psychotropic drug, No.	29 / 13	NA
Total Y-BOCS score[†]	21.0 ± 6.7	NR
Total HDRS score[†]	4.6 ± 3.9	NR

Values represent the mean ± standard deviation.

Abbreviations: NA, not applicable; NR, not recorded; OCD, obsessive-compulsive disorder; SRI,

serotonin reuptake inhibitor; Y-BOCS, Yale-Brown Obsessive-Compulsive Scale[36]; HDRS, 17-

item Hamilton Depression Rating Scale[19].

[†]Classification of handedness was based on a modified 25-item version of the Edinburgh

Handedness Inventory. Except for one missing value of the HDRS in the OCD group, all patients

were surveyed for obsessive-compulsive and depressive symptoms based upon the Japanese

version of Y-BOCS and HDRS, respectively.

1   **Supplementary Table 2. SRI doses and imipramine equivalent doses of each patient (n =**

2   **29).**

SRI dose	Imipramine equivalent dose
Paroxetine 5 mg	18.75 mg
Paroxetine 15 mg	56.25 mg
Paroxetine 20 mg	75 mg
Paroxetine 25 mg	93.75 mg
Paroxetine 30 mg	112.5 mg
Paroxetine 30 mg	112.5 mg
Escitalopram 10 mg	75 mg
Escitalopram 10 mg	75 mg
Escitalopram 10 mg	75 mg
Clomipramine 50 mg	62.5 mg
Paroxetine 40 mg	150 mg
Paroxetine 40 mg	150 mg
Paroxetine 50 mg	187.5 mg
Paroxetine 50 mg	187.5 mg
Paroxetine 50 mg	187.5 mg
Fluvoxamine 150 mg	150 mg
Fluvoxamine 150 mg	150 mg
Fluvoxamine 150 mg	150 mg
Sertraline 75 mg + Escitalopram 10 mg	187.5 mg

Clomipramine 150 mg	187.5 mg
Clomipramine 75 mg	93.75 mg
Escitalopram 10 mg	75 mg
Fluvoxamine 100 mg	100 mg
Paroxetine 30 mg	112.5 mg
Paroxetine 50 mg	187.5 mg
Escitalopram 20 mg	150 mg
Paroxetine 50 mg	187.5 mg
Paroxetine 50 mg	187.5 mg
Paroxetine 30 mg	112.5 mg

<div align="center">**Supplementary Note**</div>

Theoretical derivation of conditions for OCD-like behavior

Here, we analytically derived the condition in which the separate eligibility trace model (Figure 1b in the main text) exhibits OCD-like behavior in the obsession-relief state transition (Figure 1a in the main text). In the following derivations, our manual calculations were confirmed using MATLAB Symbolic Math Toolbox (Mathworks, version R2017a). Hence, we have skipped the derivation passages, because they are too long and complicated to include in the manuscript. Instead, at the end of the Supplementary Text, we provide the MATLAB source code for confirming the equations (see Sec. 4).

1   Stationary analysis of the obsession-relief state transition

The state transition diagram imitating obsessive compulsive symptoms (Figure 1a) consists of a relief state ($s_0 = 0$) and an obsessive state ($s_1 = 1$). Available options are 'abnormal reaction to intrusive thoughts' ($a_1 = 1$) and 'other' ($a_0 = 0$) in s_0 and 'compulsion' ($a_1 = 1$) and 'other' ($a_0 = 0$) in s_1. Define a binary vector variable $\boldsymbol{x}(t)$ to represent the combination of state and action, $(s(t), a(t))$, at time step t as

$$\boldsymbol{x}(t) \equiv \boldsymbol{X}(a(t), s(t)),$$

where $\boldsymbol{X}(a, s)$ is a binary vector function defined as

$$\boldsymbol{X}(0,0) \equiv \begin{pmatrix} 1 & 0 & 0 & 0 \end{pmatrix}^T, \quad \boldsymbol{X}(0,1) \equiv \begin{pmatrix} 0 & 0 & 1 & 0 \end{pmatrix}^T,$$
$$\boldsymbol{X}(1,0) \equiv \begin{pmatrix} 0 & 1 & 0 & 0 \end{pmatrix}^T, \quad \boldsymbol{X}(1,1) \equiv \begin{pmatrix} 0 & 0 & 0 & 1 \end{pmatrix}^T, \tag{1}$$

where M^T means transposition of the matrix M. The expectation value of vector $\boldsymbol{x}$, $E[\boldsymbol{x}]$ represents the probability distribution of the combination of state and action. Throughout the Supplementary Note, all four-dimensional vectors are denoted as column vectors, each of which consisted of components corresponding to state-action patterns in the order $a_0 s_0$, $a_1 s_0$, $a_0 s_1$, and $a_1 s_1$.

Likewise, let $\boldsymbol{s}(t)$ be a binary vector variable to represent only state:

$$\boldsymbol{s}(t) \equiv B\boldsymbol{x}(t), \quad B \equiv \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

The combination of state and action at time t determines the state transition probabilities,

$$E[\boldsymbol{s}(t+1)|\boldsymbol{x}(t)] = M\boldsymbol{x}(t), \quad M \equiv \begin{pmatrix} \boldsymbol{1}^T - \boldsymbol{p}^T \\ \boldsymbol{p}^T \end{pmatrix},$$
$$\boldsymbol{1} \equiv \begin{pmatrix} 1 & 1 & 1 & 1 \end{pmatrix}^T,$$
$$\boldsymbol{p} \equiv \begin{pmatrix} p_{00} & p_{10} & p_{01} & p_{11} \end{pmatrix}^T = \begin{pmatrix} 0 & 1 & p_{01} & p_{11} \end{pmatrix}^T,$$
$$p_{as} \equiv P(s(t+1) = 1 | a(t) = a, s(t) = s).$$

Let $c_{as}^{s'}$ be a given cost at a state transition: $a(t) = a$ and $s(t) = s$ to $s(t+1) = s'$:

$$\boldsymbol{c}^0 \equiv \begin{pmatrix} c_{00}^0 & c_{10}^0 & c_{01}^0 & c_{11}^0 \end{pmatrix}^T = \begin{pmatrix} 0 & 0 & 0 & c \end{pmatrix}^T,$$

$$\boldsymbol{c}^1 \equiv \begin{pmatrix} c_{00}^1 & c_{10}^1 & c_{01}^1 & c_{11}^1 \end{pmatrix}^T = \begin{pmatrix} 0 & 0 & 1 & 1+c \end{pmatrix}^T.$$

We focused on the case in which the compulsive action in the obsessive state minimizes immediate cost. The expected cost for the compulsion is $c_{11}^1 p_{11} + c_{11}^0(1 - p_{11}) = p_{11} + c$, and the expected cost for "other" is $c_{01}^1 p_{01} + c_{01}^0(1 - p_{01}) = p_{01}$. Thus, we focused on the condition:

$$p_{11} + c < p_{01}. \tag{2}$$

The conditional expectation value of $\boldsymbol{x}(t)$ on state $\boldsymbol{s}(t)$ is described with the state-dependent choice probabilities,

$$E[\boldsymbol{x}(t)|\boldsymbol{s}(t)] = C\boldsymbol{s}(t), \quad C \equiv \begin{pmatrix} 1 - \pi_0 & \pi_0 & 0 & 0 \\ 0 & 0 & 1 - \pi_1 & \pi_1 \end{pmatrix}^T,$$

$$\pi_s \equiv P(a(t) = 1 | s(t) = s).$$

Markov chains for variables $\boldsymbol{s}$ and $\boldsymbol{x}$ are respectively described as

$$E[\boldsymbol{s}(t+1)] = MC\,E[\boldsymbol{s}(t)], \quad E[\boldsymbol{x}(t+1)] = CM\,E[\boldsymbol{x}(t)].$$

The stationary distribution of $\boldsymbol{s}$ is obtained as the eigenvector of 2×2 matrix MC satisfying the recursive equation $E[\boldsymbol{s}] = MC\,E[\boldsymbol{s}]$. The stationary distribution of $\boldsymbol{x}$ is obtained as $E[\boldsymbol{x}] = C\,E[\boldsymbol{s}]$. The average outcome is described as

$$E[r(t+1)] = -\left(\text{diag}(\boldsymbol{1} - \boldsymbol{p})\boldsymbol{c}^0 + \text{diag}(\boldsymbol{p})\boldsymbol{c}^1\right)^T E[\boldsymbol{x}(t)].$$

The partial derivative for π_0 is obtained as (execute script in Sec. 4)

$$\frac{\partial E[r(t+1)]}{\partial \pi_0} = -\frac{\left((1 - p_{01})(1 - \pi_1) + (1 - p_{11})\pi_1\right)\left(p_{01}(1 - \pi_1) + (c + p_{11})\pi_1\right)}{(\pi_0 - p_{01} + p_{01}\pi_1 - p_{11}\pi_1 + 1)^2} < 0. \tag{3}$$

Thus, the optimal reaction rate to intrusive thoughts is always $\pi_0 = 0$. The partial derivative for π_1 is obtained as (execute script in Sec. 4)

$$\frac{\partial E[r(t+1)]}{\partial \pi_1} = \frac{(p_{01} - c - p_{11})(1 + \pi 0) + cp_{01}}{(\pi_0 - p_{01} + p_{01}\pi_1 - p_{11}\pi_1 + 1)^2}\pi_0 \geq 0, \tag{4}$$

because we focused on the condition (2). Thus, the optimal compulsion rate is $\pi_1 = 1$.

2   Stationary analysis of the separate eligibility trace model

In the separate eligibility trace model, the trace patterns are described as a decaying summation of the past $\boldsymbol{x}$ (see Online Methods).

$$\boldsymbol{\eta}^{\pm}(t) = \sum_{\tau=0}^{\infty} (\nu^{\pm})^{\tau} \boldsymbol{x}(t - \tau),$$

where the trace factors are in the range $0 \leq \nu^{\pm} < 1$. We define the policy vector $\boldsymbol{q}$ to represent the policy parameters,

$$\boldsymbol{q} = \begin{pmatrix} q_{00} & q_{10} & q_{01} & q_{11} \end{pmatrix}^T,$$

and the conditional expectation value of the policy update $\Delta \boldsymbol{q}(t)$ on $\boldsymbol{x}(t)$ is described as

$$E[\Delta \boldsymbol{q}(t)|\boldsymbol{x}(t)] = \alpha E\left[\max\left(0, \varepsilon(t)\right)\boldsymbol{\eta}^+(t) + \min\left(0, \varepsilon(t)\right)\boldsymbol{\eta}^-(t)\Big|\boldsymbol{x}(t)\right], \tag{5}$$

where the outcome prediction error $\varepsilon(t)$ depends on learning algorithms,

$$\varepsilon(t) = \begin{cases} r(t+1) + \gamma(q_{0s(t+1)} + q_{1s(t+1)}) - (q_{0s(t)} + q_{1s(t)}), & \text{(Actor-Critic)}, \\ r(t+1) + \gamma \max(q_{0s(t+1)}, q_{1s(t+1)}) - q_{a(t)s(t)}, & \text{(Q-learning)}, \\ r(t+1) + \gamma q_{a(t+1)s(t+1)} - q_{a(t)s(t)}, & \text{(SARSA)}, \end{cases} \tag{6}$$

where $r(t)$ is the outcome at time step t. The parameter γ denotes the discount factor ($0 \leq \gamma < 1$) in the temporal difference (TD) learning framework. The average prediction error should approach zero at the stationary phase after sufficient learning. Therefore, the transition to the relief state s_0 produces a positive error, and the transition to the obsessive state s_1 produces a negative error. Accordingly, the conditional expectation values of the positive and negative prediction errors are described as

$$E\left[\max\left(0, \varepsilon(t)\right)|a(t) = a, s(t) = s\right] = E[\varepsilon(t)|s(t+1) = 0, a(t) = a, s(t) = s](1 - p_{as}),$$
$$E\left[\min\left(0, \varepsilon(t)\right)|a(t) = a, s(t) = s\right] = E[\varepsilon(t)|s(t+1) = 1, a(t) = a, s(t) = s]p_{as}.$$

Define vector $\boldsymbol{\varepsilon}^{s'}$ as

$$\boldsymbol{\varepsilon}^{s'} \equiv \begin{pmatrix} \varepsilon_{00}^{s'} & \varepsilon_{10}^{s'} & \varepsilon_{01}^{s'} & \varepsilon_{11}^{s'} \end{pmatrix}^T,$$
$$\varepsilon_{as}^{s'} \equiv E[\varepsilon(t)|s(t+1) = s', a(t) = a, s(t) = s].$$

Using these vector forms, the expectation value of the policy update $E[\Delta \boldsymbol{q}]$ is obtained by averaging Eq. (5) over the stationary distribution $E[\boldsymbol{x}(t)] = E[\boldsymbol{x}]$ as

$$E[\Delta \boldsymbol{q}^T] = \alpha \boldsymbol{\varepsilon}^{0T} \mathrm{diag}(\mathbf{1} - \boldsymbol{p}) E[\boldsymbol{x}(t)\boldsymbol{\eta}^+(t)^T] + \alpha \boldsymbol{\varepsilon}^{1T} \mathrm{diag}(\boldsymbol{p}) E[\boldsymbol{x}(t)\boldsymbol{\eta}^-(t)^T],$$

where matrices $E[\boldsymbol{x}(t)\boldsymbol{\eta}^{\pm}(t)^T]$ are calculated as

$$E[\boldsymbol{x}(t)\boldsymbol{\eta}^{\pm}(t)^T] = \sum_{\tau=0}^{\infty}(\nu^{\pm})^{\tau} E[\boldsymbol{x}(t)\boldsymbol{x}(t-\tau)^T] = \sum_{\tau=0}^{\infty}(\nu^{\pm}CM)^{\tau} E[\boldsymbol{x}(t-\tau)\boldsymbol{x}(t-\tau)^T]$$
$$= \sum_{\tau=0}^{\infty}(\nu^{\pm}CM)^{\tau} \mathrm{diag}(E[\boldsymbol{x}]) = (I - \nu^{\pm}CM)^{-1}\mathrm{diag}(E[\boldsymbol{x}]),$$

because vector $\boldsymbol{x}(t - \tau)$ consists of exclusive binary components. The matrix I denotes the identity matrix. Thus, we obtain the expectation value of the policy update as

$$E[\Delta \boldsymbol{q}] = Z\left(H^0 \boldsymbol{\varepsilon}^0 + H^1 \boldsymbol{\varepsilon}^1\right), \tag{7}$$
$$H^0 \equiv (I - \nu^+ M^T C^T)^{-1}\mathrm{diag}(\mathbf{1} - \boldsymbol{p}),$$
$$H^1 \equiv (I - \nu^- M^T C^T)^{-1}\mathrm{diag}(\boldsymbol{p}), \quad Z \equiv \alpha \, \mathrm{diag}(E[\boldsymbol{x}]).$$

3 Stability of optimal behavior

3.1 Q-learning and SARSA

The abnormal-reaction and compulsion rates, π_0 and π_1, are described as soft-max functions of the policy parameters (see Online Methods),

$$\pi_s \equiv P(a(t) = 1|s(t) = s) = \frac{e^{\beta q_{1s}}}{e^{\beta q_{0s}} + e^{\beta q_{1s}}} = \frac{1}{1 + e^{-\beta(q_{1s} - q_{0s})}}. \tag{8}$$

In Q-learning and SARSA, the policy parameters represent action values. The balance of exploration and exploitation is explicitly controlled, and the parameter β increased to the limit $\beta \to \infty$ in order to reach the final behavior. Therefore, the final behavior is determined by the difference in the action values,

$$E[D\boldsymbol{q}] = E\left[\begin{pmatrix} q_{10} - q_{00} \\ q_{11} - q_{01} \end{pmatrix}\right], \quad D \equiv \begin{pmatrix} -1 & 1 & 0 & 0 \\ 0 & 0 & -1 & 1 \end{pmatrix}. \tag{9}$$

In order to reach $(\pi_0, \pi_1) = (0, 1)$ as the final behavior, the stationary action values should satisfy the inequalities, $q_{00} > q_{10}$ and $q_{01} < q_{11}$. Regarding the assumptions of the inequalities, prediction error patterns in Q-learning are described as $\varepsilon_{as}^0 = \gamma q_{00} - q_{as} - c_{as}^0$, and $\varepsilon_{as}^1 = \gamma q_{11} - q_{as} - c_{as}^1$. These can be described in vector forms by using the binary vector function $\boldsymbol{X}(a, s)$ in Eq.(1), the unit vector $\boldsymbol{1}$, and the identity matrix I,

$$\begin{aligned}
\boldsymbol{\varepsilon}^s &= G_{\mathrm{QL}}^s \boldsymbol{q} - \boldsymbol{c}^s, \\
G_{\mathrm{QL}}^0 &\equiv \gamma \boldsymbol{1}\, \boldsymbol{X}(0, 0)^T - I^{(4)}, \quad G_{\mathrm{QL}}^1 \equiv \gamma \boldsymbol{1}\, \boldsymbol{X}(1, 1)^T - I^{(4)}.
\end{aligned}$$

In SARSA, the prediction error patterns are described as

$$\varepsilon_{as}^{s'} = \gamma E[q_{a(t+1)s'}] - q_{as} - c_{as}^{s'} = \gamma(q_{a_0 s'}(1 - \pi_{s'}) + q_{a_1 s'}\pi_{s'}) - q_{as} - c_{as}^{s'},$$

which can be described in vector forms,

$$\begin{aligned}
\boldsymbol{\varepsilon}^s &= G_{\mathrm{SA}}^s \boldsymbol{q} - \boldsymbol{c}^s, \\
G_{\mathrm{SA}}^s &\equiv \gamma \boldsymbol{1}\left(\boldsymbol{X}(1, s)^T(1 - \pi_s) + \boldsymbol{X}(0, s)^T \pi_s\right) - I^{(4)}.
\end{aligned}$$

The stationary solution of $E[\Delta \boldsymbol{q}] = \boldsymbol{0}$ in Eq. (7) is obtained as

$$\begin{aligned}
E[\Delta \boldsymbol{q}] &= Z(H^0 \boldsymbol{\varepsilon}^0 + H^1 \boldsymbol{\varepsilon}^1) = ZH^0\left(G^0 \boldsymbol{q} - \boldsymbol{c}^0\right) + ZH^1\left(G^1 \boldsymbol{q} - \boldsymbol{c}^1\right) \\
&= Z(H^0 G^0 + H^1 G^1)\boldsymbol{q} - Z(H^0 \boldsymbol{c}^0 + H^1 \boldsymbol{c}^1) = \boldsymbol{0}, \\
\boldsymbol{q}^* &= \left(H^0 G^0 + H^1 G^1\right)^{-1}(H^0 \boldsymbol{c}^0 + H^1 \boldsymbol{c}^1). \tag{10}
\end{aligned}$$

For both Q-learning and SARSA, differences in the action values in the optimal behavior $(\pi_0, \pi_1) = (0, 1)$ are obtained as (execute script in Sec. 4)

$$\lim_{\pi_0 \to 0, \pi_1 \to 1} E[D\boldsymbol{q}^*] = \frac{1}{N} \left(J - K \quad L - c(J - K) \right)^T, \tag{11}$$

$$J \equiv p_{11}(1 - p_{11})(\nu^+ - \nu^-),$$

$$K \equiv \gamma(p_{11} + c)(1 - \nu^+ p_{11}) \geq 0,$$

$$L \equiv (1 + c\gamma)(p_{01} - p_{11} - c)(1 - \nu^+ p_{11}) > 0,$$

$$N \equiv (1 - p_{11})(1 - \nu^- p_{11}) + p_{11}(1 - \gamma)(1 - \nu^+ p_{11}) > 0.$$

If $J < K$, then the stationary action values satisfy the inequalities, $q_{00} > q_{10}$ and $q_{01} < q_{11}$ and the optimal behavior $(\pi_0, \pi_1) = (0, 1)$ would be preserved. But if $J > K$, then the zero abnormal-reaction rate to intrusive thoughts $\pi_0 = 0$ is broken. Thus, the condition in which the optimal behavior might be broken is described as

$$\text{OCD}^+ : \quad J > K. \tag{12}$$

Because $K \geq 0$, a necessary condition is that $J > 0$, namely, a trace imbalance: $\nu^+ > \nu^-$.

In behavioral therapy [exposure and response prevention (ERP)], compulsion is prevented in the obsessive state, that is, $\pi_1 = 0$. For Q-learning, the action value q_{11} can be used even if the compulsive action $(a = 1)$ is prevented in the obsessive state $(s = 1)$ by the definition of the algorithm. However, response prevention is equivalent to the infinite response cost, which leads to inequality in action values, $q_{11} < q_{01}$. Thus, the matrix G_{QL}^0 is replaced in the behavioral therapy,

$$G_{\text{QL}}^1 = \gamma \boldsymbol{1} \, \boldsymbol{X}(0, 1)^T - I^{(4)}.$$

For both Q-learning and SARSA, differences in action values at the optimal behavior in the behavioral therapy, $(\pi_0, \pi_1) = (0, 0)$, are obtained as (execute script in Sec. 4)

$$\lim_{\pi_0 \to 0, \pi_1 \to 0} E[D\boldsymbol{q}^*] = \frac{1}{\tilde{N}} \left(\tilde{J} - \tilde{K} \quad \tilde{L} - c(\tilde{J} - \tilde{K}) \right)^T, \tag{13}$$

$$\tilde{J} \equiv p_{01}(1 - p_{01})(\nu^+ - \nu^-),$$

$$\tilde{K} \equiv \gamma p_{01}(1 - \nu^+ p_{01}) \geq 0,$$

$$\tilde{L} \equiv (p_{01} - p_{11} - c)(1 - \nu^+ p_{01}) > 0,$$

$$\tilde{N} \equiv (1 - p_{01})(1 - \nu^- p_{01}) + (1 - \gamma)p_{01}(1 - \nu^+ p_{11}) > 0.$$

In the same logic as the derivation of the condition around $(\pi_0, \pi_1) = (0, 1)$, the condition in which the behavioral therapy (ERP) might be ineffective is

$$\text{ERP}^- : \quad \tilde{J} > \tilde{K}. \tag{14}$$

3.2 Actor-critic learning

In actor-critic learning, the policy parameters diverge to approach an optimal deterministic behavior, whereas the state values $\boldsymbol{v} = B\boldsymbol{q}$ converge. The prediction error patterns in actor-critic learning are described as

$$\boldsymbol{\varepsilon}^s = G_{\text{AC}}^s \boldsymbol{v} - \boldsymbol{c}^s,$$

$$G_{\text{AC}}^0 \equiv \begin{pmatrix} \gamma - 1 & 0 \\ \gamma - 1 & 0 \\ \gamma & -1 \\ \gamma & -1 \end{pmatrix}, \quad G_{\text{AC}}^1 \equiv \begin{pmatrix} -1 & \gamma \\ -1 & \gamma \\ 0 & \gamma - 1 \\ 0 & \gamma - 1 \end{pmatrix}.$$

The solution of $E[\Delta \boldsymbol{v}] = E[\Delta B\boldsymbol{q}] = \boldsymbol{0}$ in Eq. (7) is obtained as

$$\begin{aligned}
E[\Delta \boldsymbol{v}] &= E[\Delta B\boldsymbol{q}] = BZ\left(H^0 \boldsymbol{\varepsilon}^0 + H^1 \boldsymbol{\varepsilon}^1\right) \\
&= BZ\left(H^0 \left(G_{\text{AC}}^0 \boldsymbol{v} - \boldsymbol{c}^0\right) + H^1 \left(G_{\text{AC}}^1 \boldsymbol{v} - \boldsymbol{c}^1\right)\right) \\
&= BZ(H^0 G_{\text{AC}}^0 + H^1 G_{\text{AC}}^1)\boldsymbol{v} - BZ(H^0 \boldsymbol{c}^0 + H^1 \boldsymbol{c}^1) = \boldsymbol{0}, \\
\boldsymbol{v}^* &= \left(BZH^0 G_{\text{AC}}^0 + BZH^1 G_{\text{AC}}^1\right)^{-1} BZ(H^0 \boldsymbol{c}^0 + H^1 \boldsymbol{c}^1). \quad (15)
\end{aligned}$$

The signs of the updates in the abnormal-reaction and compulsion rates for the stationary $\boldsymbol{v}^*$ determine the stability of the optimal behavior $(\pi_0, \pi_1) = (0, 1)$; hence, signs of the updates in the abnormal-reaction and compulsion rates coincide with the signs of the difference between the policy updates $\Delta q_{1s} - \Delta q_{0s}$. To examine learning stability, the difference update is evaluated,

$$E[D\Delta \boldsymbol{q}] = DZH^0 \left(G_{\text{AC}}^0 \boldsymbol{v}^* - \boldsymbol{c}^0\right) + DZH^1 \left(G_{\text{AC}}^1 \boldsymbol{v}^* - \boldsymbol{c}^1\right). \quad (16)$$

The difference update $E[D\Delta \boldsymbol{q}]$ is zero at the optimal behavior $(\pi_0, \pi_1) = (0, 1)$ (execute script in Sec. 4). Therefore, we derived the Taylor expansion up to the first non-zero order (execute script in Sec. 4),

$$E[D\Delta \boldsymbol{q}] \simeq \frac{2\alpha\pi_0}{L} \begin{pmatrix} J - K & 0 \end{pmatrix}^T + O(|\boldsymbol{\pi}|^2), \quad (17)$$

For a sufficiently small abnormal-reaction rate to intrusive thoughts $\pi_0 \simeq 0$, the update of the abnormal-reaction rate $O(\pi_0)$ is dominant over the compulsion update $O(|\boldsymbol{\pi}|^2)$. Therefore, the stability of the optimal behavior $(\pi_0, \pi_1) = (0, 1)$ would be broken if $E[\Delta q_{10} - \Delta q_{00}] > 0$. Thus, the condition for the stability broken in actor-critic learning is equivalent to those in Q-learning and SARSA, $J > K$, as Eq.(12).

In behavioral therapy (ERP) where $\pi_1 = 0$ is forced, the difference update $E[D\Delta \boldsymbol{q}]$ is also zero at $\pi_0 = 0$ (see Sec. 4). Therefore, we derived the Taylor expansion up to the first non-zero order around $\pi_0 = 0$ (execute script in Sec. 4),

$$E[D\Delta \boldsymbol{q}] \simeq \frac{2\alpha\pi_0}{\tilde{L}} \begin{pmatrix} \tilde{J} - \tilde{K} & 0 \end{pmatrix}^T + O(\pi_0^2). \quad (18)$$

Thus, the condition in which behavioral therapy (ERP) might be ineffective in actor-critic learning is equivalent to that in Q-learning and SARSA, $\tilde{J} > \tilde{K}$, as Eq.(14).

3.3 Necessary condition for OCD patients

The state transition diagram (Figure 1a in the main text) represents a certain obsessive-compulsive situation. State-transition probabilities p_{01}, p_{11} and the relative cost of compulsive action c depend on the concrete contents of the abnormal reaction to intrusive thoughts, obsessive state, and compulsive action. Patients with OCD exhibit various symptoms in response to many possible situations. Hence, it is significant to derive the condition for the learning parameters in which some state-transition probabilities p_{01}, p_{11} and cost c exist to satisfy the condition for OCD-like behavior (12).

$$
\begin{aligned}
K &= \gamma(p_{11} + c)(1 - \nu^+ p_{11}) < J = p_{11}(1 - p_{11})(\nu^+ - \nu^-), \\
c &< \frac{p_{11}(1 - p_{11})(\nu^+ - \nu^-)}{\gamma(1 - \nu^+ p_{11})} - p_{11} = \frac{(1 - p_{11})(\nu^+ - \nu^-) - \gamma(1 - \nu^+ p_{11})}{\gamma(1 - \nu^+ p_{11})} p_{11}.
\end{aligned}
$$

Therefore, the condition for the existence of $c > 0$ is

$$
\begin{aligned}
0 &< (1 - p_{11})(\nu^+ - \nu^-) - \gamma(1 - \nu^+ p_{11}) \\
&= (1 - p_{11})(\nu^+ - \nu^- - \gamma \nu^+) - \gamma(1 - \nu^+) \equiv f(p_{11}).
\end{aligned}
$$

The right-hand value at $p_{11} = 1$ is negative, $f(1) = -\gamma(1 - \nu^+) < 0$; hence, the condition for the existence of p_{11}, such that $0 < p_{11} < 1$ and $f(p_{11}) > 0$ is $f(0) = \nu^+ - \nu^- - \gamma > 0$, that is,

$$
\gamma < \nu^+ - \nu^-. \tag{19}
$$

If the individual's learning parameters satisfy the condition (19), the individual might exhibit OCD symptoms in some situations.

4 MATLAB script to confirm derivations

Copy and execute the following MATLAB script, and you can confirm the equations above mentioned. This script requires Symbolic Math Toolbox.

```matlab
1  % derivation.m
2  %
3  % MATLAB script to confirm derivations in Supplementary Text of the article:
4  %    Sakai and Sakai et al. "Memory Trace Imbalance in Reinforcement and ...
       Punishment Systems Can Reinforce Implicit Choices Leading to Obsessive-...
       Compulsive Behavior"
5  %
6  % This requires Symbolic Math Toolbox
7  %
8  %%%%% Definitions
9  syms p01 p11 pi0 pi1 gam np nm al c;
10 B=[1 1 0 0;0 0 1 1]; p=[0 1 p01 p11]; M=[1-p;p]; C=[1-pi0 0;pi0 0;0 1-pi1;0 pi1...
       ];
11 D=[-1 1 0 0;0 0 -1 1]; c0=[0;0;0;c]; c1=[0;0;1;1+c];
12
13 %%%%% Stationary probability distribution
14 MC=M*C; s=[MC(1,2);1-MC(1,1)]; s=s/sum(s); x=C*s; Z=al*diag(x);
15 H0=inv(eye(4)-np*M.'*C.')*diag(1-p); H1=inv(eye(4)-nm*M.'*C.')*diag(p);
16 %H0=(eye(4)+np*M.'*inv(eye(2)-np*MC.')*C.')*diag(1-p); H1=(eye(4)+nm*M.'*inv(...
       eye(2)-nm*MC.')*C.')*diag(p); %%% Forms using 2x2
17
18 %%%%%% Optimal Behavior
19 r=simplify(-(diag(1-p)*c0+diag(p)*c1)'*x);
20 dr0=-((1-p01)*(1-pi1)+(1-p11)*pi1)*(p01*(1-pi1)+(c+p11)*pi1)/(pi0-p01+p01*pi1-...
       p11*pi1+1)^2; %%% Partial derivative of E[r] for pi0
21 dr1=pi0*((p01-c-p11)*(1+pi0) +c*p01)/(pi0-p01+p01*pi1-p11*pi1+1)^2; ...
                            %%% Partial derivative of E[r] for pi1
22 checkEq3=simplify(dr0-diff(r,pi0))
23 checkEq4=simplify(dr1-diff(r,pi1))
24
25 %%%%%% Stationary analysis of SARSA
26 Gsa0=gam*ones(4,1)*[1-pi0 pi0 0 0]-eye(4); Gsa1=gam*ones(4,1)*[0 0 1-pi1 pi1]-...
       eye(4); F=H0*Gsa0+H1*Gsa1;
27 Dq=D*simplify(inv(subs(F,[pi0 pi1],[0 1]))*subs(H0*c0+H1*c1,[pi0 pi1],[0 1]));
28 J=p11*(1-p11)*(np-nm); K=gam*(p11+c)*(1-np*p11); L=(1-np*p11)*(1+c*gam)*(p01-...
       p11-c); N=(1-p11)*(1-nm*p11)+p11*(1-gam)*(1-np*p11);
29 checkEq11=simplify(Dq-[J-K; L-c*(J-K)]/N)
30
31 %%%%% Behaviral therapy
32 Dq=D*simplify(inv(subs(F,[pi0 pi1],[0 0]))*subs(H0*c0+H1*c1,[pi0 pi1],[0 0]));
33 tJ=p01*(1-p01)*(np-nm); tK=gam*p01*(1-np*p01); tL=(1-np*p01)*(p01-p11-c); tN...
```

```matlab
        =(1-p01)*(1-nm*p01)+p01*(1-gam)*(1-np*p01);
34  checkEq13=simplify(Dq-[tJ-tK; tL-c*(tJ-tK)]/tN)
35
36  %%%%%% Stationary analysis of Q-learning
37  Gql0=gam*ones(4,1)*[1 0 0 0]-eye(4); Gql1=gam*ones(4,1)*[0 0 0 1]-eye(4); F=H0*...
        Gql0+H1*Gql1;
38  Dq=D*simplify(inv(subs(F,[pi0 pi1],[0 1])))*subs(H0*c0+H1*c1,[pi0 pi1],[0 1]));
39  checkEq11=simplify(Dq-[J-K; L-c*(J-K)]/N)
40
41  %%%%% Behaviral therapy
42  Gql1=gam*ones(4,1)*[0 0 1 0] - eye(4); F=H0*Gql0+H1*Gql1;
43  Dq=D*simplify(inv(subs(F,[pi0 pi1],[0 0])))*subs(H0*c0+H1*c1,[pi0 pi1],[0 0]));
44  checkEq13=simplify(Dq-[tJ-tK; tL-c*(tJ-tK)]/tN)
45
46  %%%%%% Stationary analysis of Actor-Critic
47  Gac0=[gam-1 0;gam-1 0;gam -1;gam -1]; Gac1=[-1 gam;-1 gam;0 gam-1;0 gam-1];
48
49  v=simplify(inv( B*Z*(H0*Gac0+H1*Gac1) )*B*Z*(H0*c0+H1*c1));  %%% Stationary ...
        state value
50  Ddq=simplify(D*Z*(H0*(Gac0*v-c0)+H1*(Gac1*v-c1)));          %%% Policy update ...
        difference for stationary state value
51  %%%%% Tailor expansion of E[D Delta q]
52  Ddq_opt=simplify(subs(Ddq,[pi0 pi1],[0 1]))    %%% Policy update differences ...
        at pi=[0,1] are 0
53  %%% Partial derivertives of policy update difference at pi=[0,1]
54  Ddq_0=simplify(subs(diff(Ddq,pi0),[pi0 pi1],[0 1])); Ddq_1=simplify(subs(diff(...
        Ddq,pi1),[pi0 pi1],[0 1]))
55  checkEq17=simplify(Ddq_0-[2*al/N*(J-K); 0])
56
57  %%%%% Behaviral therapy
58  Ddq_beh=subs(Ddq,pi1,0);
59  %%%%% Tailor expansion of E[D Delta q]
60  Ddq_opt=simplify(subs(Ddq_beh,pi0,0))           %%% Policy update differences ...
        at pi0=0 are 0
61  %%% Derivertives of policy update difference at pi0=0
62  Ddq_0=simplify(subs(diff(Ddq_beh,pi0),pi0,0));
63  checkEq18=simplify(Ddq_0-[2*al/tN*(tJ-tK); 0])
```