1 **AtHDA15 attenuates COP1 *via* transcriptional quiescence, direct binding, and sub-**

2 **compartmentalization during photomorphogenesis**

3 Malona V. Alinsug[1,2,3*] and Custer C. Deocaris [4,5]

4
5 [1] Institute of Plant Biology, College of Life Science, National Taiwan University, Taipei, Taiwan
6
7 [2] Center for Food & Bio convergence, College of Agriculture & Life Sciences, Seoul National
8 University, South Korea
9
10 [3] Science Department, College of Natural Sciences & Mathematics, Mindanao State University-
11 General Santos City, Philippines
12
13 [4] Biomedical Research Section, Philippine Nuclear Research Institute, Department of Science
14 and Technology, Commonwealth Avenue, Diliman, Quezon City, Philippines
15
16 [5] Technological Institute of the Philippines, Cubao, Quezon City, Philippines
17
18
19 * Corresponding author
20
21 Science Department, College of Natural Sciences & Mathematics,
22 Mindanao State University-General Santos City, Philippines
23 E-mail: malinsug@yahoo.com
24

30 **SHORT TITLE: Mechanisms for HDA15 attenuation of COP1**

Abstract

Light is an essential environmental cue that determines the overall growth and development of plants. However, the molecular mechanisms underpinning the light signaling network are obscured by the epigenetic machinery where reversible acetylation and deacetylation play crucial roles in modulating light-regulated gene expression. In this paper, we demonstrate that HDA15 represses COP1, the master switch in the light signaling network, by deacetylation, protein interaction, and sub-compartmentalization. *hda15* T-DNA mutant lines exhibited light hyposensitivity with significantly reduced HY5 and PIF3 transcript levels leading to long-hypocotyl phenotypes in the dark while its overexpression exhibited elevated HY5 transcripts and short hypocotyl phenotypes. *In vivo* and *in vitro* binding assays further show that HDA15 directly interacts with COP1 inside the nucleus modulating COP1's repressive activities. Crossing *hda15-t27* with *cop1-4* mutants resulted in short-hypocotyl and dwarfed phenotypes, reminiscent of *cop1-4* mutants suggesting COP1 is epistatic to HDA15. Although light signals the nucleocytoplasmic shuttling of HDA15, the presence of COP1 triggers its nuclear localization. A working model is presented elucidating the concerted interplay between HDA15 and COP1 under light and dark conditions.

Keywords: histone deacetylases, photomorphogenesis, protein interaction, light signal transduction, gene expression regulation

Introduction

Plants adapt extensively to the dynamic changes in their light environment as they are critically dependent on light for photosynthesis, growth, and development. In general, light induces massive changes in a plant's transcriptional activity resulting in distinct expression profiles during seedling development (Jiao et al., 2007; Ma et al., 2013). Approximately 15% of these transcriptional changes have been attributed to histone modifications, specifically the acetylation in H3K9K27, which coincides with the upregulation of key photomorphogenesis gene expression, such as HY5 and its homolog, HYH (Charron et al., 2009). H3K9ac is one of the most characterized epigenetic marks invariably correlated with transcriptional activation in all the species studied (Ausin et al., 2004; Kurdistani et al., 2004; Schübeler et al., 2004; Benhamed et al., 2008; Guo et al., 2008; Charron et al., 2009; Zhou et al., 2009; Schenke et al., 2014; Liu et al., 2017). However, most of these light-induced pathways remain obscure, while little is understood about histone deacetylases' involvement in regulating these genes.

Studies on histone active marks such as H3K9K27ac have been limited to positive photomorphogenic regulators such as CABs, LHBIB1, LHCA1, HYH and HY5 and its downstream targets CHS, IAA3, RBCS-1a (Benhamed et al., 2006; Guo et al., 2008; Charron et al., 2009). Among the 12 members of the RPD3/HDA1 superfamily in *Arabidopsis*, HDA19, HDA6, and HDA15 have been ascribed active roles in light-regulated gene expression. Based on studies, HDA19 and HDA6 play a negative regulatory role by repressing some of these light-regulated genes (Benhamed et al., 2006; Tessadori et al., 2009). On the other hand, HDA15 has recently been shown to have a regulatory role in photomorphogenesis. As human Class II HDACs are known to undergo nucleocytoplasmic shuttling, we previously demonstrated that HDA15 shuttles to the nucleus under light (Alinsug et al., 2012). Moreover, phytochrome

interacting factor 3 (PIF3) associates with HDA15 repressing chlorophyll biosynthesis and

photosynthesis in etiolated seedlings (Liu et al., 2013). Similarly, PIF1 recruits HDA15 at the

promoter sites repressing the transcription of light-responsive genes involved in multiple

hormonal signaling pathways and cellular processes in germinating seeds in the dark (Gu et al.,

2017). Furthermore, HDA15 forms a complex with nuclear factor homologs (NF-YC)

functioning as a transcriptional co-repressor via deacetylation inhibiting hypocotyl elongation in

photomorphogenesis (Tang et al., 2017). A recent study describes that long hypocotyl 5 (HY5)

protein interacts with HDA15 repressing hypocotyl cell elongation during photomorphogenesis

(Zhao et al., 2019).

HY5 and PIFs are key transcription factors playing crucial regulatory roles in

photomorphogenesis and skotomorphogenesis, respectively. Along with the light signaling

network, both HY5 and PIFs are repressed by COP1, an E3 ubiquitin ligase that targets both

photoreceptors HY5 and HYH for ubiquitination and proteasome degradation in the dark

(Osterlund et al., 2000).   However, the molecular interplay between COP1, HY5, PIFs, and

HDA15 remains unclear. We do not know how or when HDA15 precisely interjects within the

light response molecular network. As epigenetic regulation further adds complexity to the light

signaling pathway, we attempt to explore the multiple roles HDA15 plays during

photomorphogenesis. In this paper, we demonstrate that HDA15 represses COP1, the master

switch in the light signaling network, by transcriptional silencing via promoter deacetylation,

protein-protein association, and sub-compartmentalization. A working model is presented

elucidating the interplay of the multiple actions exerted by HDA15 on COP1 under light and

dark conditions.

Materials & Methods

Plant Material

Seeds of T-DNA knockout lines *hda5-t24* (SALK_030624)*, hda15-t27* (SALK_004027C), and *hda18-t* (SALK_006938C) in Col-0 background were purchased from ABRC and screened via PCR using specific primers. Homozygous mutant plants with positive T-DNA inserts were grown under long day and assessed for the expression of the corresponding gene compared to wildtype via RT- PCR using specifically designed primers. Seeds of homozygous T-DNA knockout lines were then collected and further used for this study. To establish the double mutant line, we crossed the homozygous *hda15-t27* mutant with the *cop1-4* mutant. The resultant F2 lines were then finally screened by PCR.

Protoplast PEG Transfection

Leaves of 3-week old Col-0 wildtype, T2 transgenic lines, and mutant lines *phyA, phyB, cry1/cry2,* and *cop1-4* were used for protoplast transient expression assay. Isolation of Arabidopsis mesophyll protoplasts and PEG transfection were done as described previously (Yoo et al., 2007) with some modifications. Mesophyll protoplasts were isolated from 3-week old Col-0 plants and Arabidopsis PSB-D cell lines. Twenty µg of HDA15-YFP/GFP fusion plasmid and VirD2-NLS, a nuclear marker, were co-transfected into protoplasts using polyethylene glycol (PEG) solution (0.4 g/ml PEG 4000, 0.8 M mannitol, 125 mM CaCl2). After incubation for 5-15 min at room temperature, the protoplasts were washed and resuspended in W5 solution (154 mM NaCl, 125 mM CaCl2, 5 mM KCl, 2 mM MES at pH5.7), and further incubated under white light for 16–24 h before imaging using Leica SP5 confocal microscope. For the localization of HDA15 in different light treatments, transfected protoplasts were

125 incubated under white light for 18 h then transferred to E30LEDL3 growth chambers (Percival

126 Scientific) with far-red, red, and blue light-emitting diode sources for 3 h. Low light intensities

127 used as treatment were measured at 2.77 μmol m^{-2} s^{-2} (FR), 1.77 μmol m^{-2} s^{-2} (R), and 3.84 μmol

128 m^{-2} s^{-2} (B).

Light treatments and hypocotyl measurements

Arabidopsis seeds were surface sterilized, cold treated for 3-4 days and sown on 1/2 MS

plates with 0.1% sucrose then subjected to different light treatments at 21°C for 5-7 consecutive

days. White light intensity was approximately 130 μmol m^{-2}s^{-1}. Plated seeds were incubated in

E30LEDL3 colored light chambers (Percival Scientific) with the corresponding low (L) and high

(H) wavelength measurements as follows: far-red at 18.065 (H) and 2.77 (L) μmol m^{-2} s^{-1}, red

light at 22.938 (H) and 1.77 (L) μmol m^{-2} s^{-1}, and blue light at 11.095 (H) and 3.84 (L) μmol m^{-2}

s^{-1}, respectively.   On the other hand, dark treated seedlings were wrapped in foil and placed

inside a dark growth chamber. Hypocotyl length was measured using UTHSCA Image Tool

using 30-50 seedlings for each line and further analyzed for statistical tests.

Total RNA extraction & qRT-PCR

Gene expression was initially assessed by semi-quantitative RT-PCR. Total RNA was

extracted from plant samples weighing 0.25 to 0.3 g using TRIZOL reagent (Invitrogen).

Oligo(dT) primed reverse transcription of first-strand cDNA synthesis was carried out with 7 mg

total RNA using Super ScriptTM III (Invitrogen). Equal volumes of each first-strand reaction

were amplified with gene-specific primer pairs. Thermocycling conditions were 95°C for 4 mins

followed by 30 cycles of 95°C for 30 s, 55–60°C for 30 s, and 74°C for 1–2 min. cDNAs

obtained from RT were then used as templates to run real-time PCR. The following components

were added to a reaction tube: 9 μL of iQ SYBR Green Supermix solution (Bio-Rad), 1 μL of 5

mM specific primers, and 8 µL of the diluted template. Thermocycling conditions were as

follows: 95ºC for 3 min followed by 40 cycles of 95ºC for 30 s, 60ºC for 30 s, and 72ºC for 20 s,

with a melting curve detected at 95ºC for 1 min, 55ºC for 1 min, and detection of the denature

time from 55ºC to 95ºC. Each sample was quantified in triplicate and normalized using

Ubiquitin10 as the internal control. Gene-specific primer pairs were designed and are available

upon request.

Chromatin Immunoprecipitation (ChIP) assay

ChIP assay was performed as described (Gendrel et al., 2005). Chromatin extracts were

prepared from light treated 4-day old seedlings treated with formaldehyde. The chromatin was

sheared to an average length of 500 bp by sonication and immunoprecipitated with anti-

acetylated histone H3K9K14 (catalog no. 06-599; Millipore). The DNA cross-linked to

immunoprecipitated proteins was analyzed by real-time PCR. Relative enrichments of various

regions of COP1 over Col-0 were calculated after normalization to Ubiquitin10. Each of the

immunoprecipitations was tested and analyzed in triplicates, and each sample was quantified at

least three times.

Protein Extraction and Gel-Blot Analysis

Total protein extraction was conducted as described (Hsieh et al., 2000). Four-day old

seedlings of wild type Col-0, all mutant, and overexpression lines were continuously grown

under different light treatments and duration. After the treatment, the seedlings were harvested

and frozen immediately for protein extraction. One hundred micrograms of total protein were

separated on a 12% SDS-polyacrylamide gel, blotted on immuno-blot PVDF (Bio-Rad), and

detected using anti-HY5, anti-COP1, and anti-RPN6 specific antibodies.

Protein-Protein Interaction

For the bifluorescence complementation (BiFC) assay, HDA15 and COP1 coding

sequences were cloned into the pCR8/GW/TOPO vector and used for recombination into its

corresponding destination vectors, pEG201-N-YFP and pEG202-C-YFP, using LR mix.

Kanamycin was used for bacterial selection. Purified plasmids were then analyzed for DNA

sequencing for confirmation and further used for protoplast PEG transfection and imaging.

Pull-down assay was carried out as described (Yang et al., 2008) with some

modifications. For GST pull-down, GST and GST-HDA15 recombinant proteins were incubated

in 30 ㎖ of GST resin with binding buffer (50 mM Tris-Cl, pH 7.5, 100 mM NaCl, 0.25%

Triton X-100, and 35 mM β-mercaptoethanol) for two h at 4℃. The binding reaction was

washed three times with the binding buffer, and then the COP1-His recombinant protein was

added and incubated for an additional two h at 4℃. For His pull-down, His and COP1-His

recombinant proteins were incubated in 30 ㎖ of His resin in a phosphate buffer (10 mM

Na2HPO4, 10 mM NaH2PO4, 500 mM NaCl, and 10 mM imidazole) for two h at 4℃, the

binding reaction was washed three times with the phosphate buffer, and then GST-HDA15

recombinant protein was added and incubated for an additional two h at 4℃. After extensive

washing (for at least eight times), the "pulled-down" proteins were eluted by boiling, separated

by 10% SDS-PAGE, and detected by western blotting using anti-His, anti-GST, or anti-COP1

specific antibody.

Bioinformatics analysis

The protein sequences of AtHDA15, COP1, and HY5 (ACCN #: Q8GXJ1, P43254, and

O24646) were retrieved from the NCBI's protein sequence repository database

(www.ncbi.nlm.nih.gov/). Putative protein ubiquitylation sites were predicted by UbPred (Kumar

and Kumar, 2019), BDM-PUB (Li et al., 2009), and Ubi-Site (Akimov et al., 2018). ProtParam

server (http://us.expasy.org/tools/protparam.html) was used to compute the instability

index (Guruprasad et al., 1990).

Statistical analysis

Statistical analyses were conducted using SigmaStat 11 software. One-way ANOVA

using the Holm-Sidak method was performed for multiple pairwise comparisons. Similarly,

Dunnette's method was conducted for multiple comparisons using WT as control. However,

Kruskal-Wallis One-way ANOVA on Ranks was used to analyze data that failed to pass the

Normality test. Significance testing was set at an α of 0.05 or 0.01. Graphs were presented in

mean values with error bars set from its standard deviation.

Results

1. Class II HDA mutant lines display light hyposensitivity.

207   T-DNA SALK lines of Class II histone deacetylases were screened, analyzed, and grown

under various light conditions where long hypocotyl phenotypes were initially observed. To

determine the maximum phenotypic expression of the mutant lines, we exposed the plants to

varying light fluence rates, light intensities, and durations. Seeds of knockout lines were grown

under white light and dark treatment for seven consecutive days. Class II mutant lines were

likewise grown under continuous far-red, red, and blue light at high and low light intensities for

five consecutive days. Hypocotyl lengths were measured on the 3rd, 5th, and 7th days (Figure 1).

214   Results indicate that loss-of-function lines of HDA5, HDA15, and HDA18 were

generally normal under white light treatment. However, in the absence of light, we observed

significantly longer hypocotyls starting on the 3rd day for *hda15-t27* and 4th day for *hda5-t24* and

hda18-t. The hyper-skotomorphogenic phenotype exhibited by Class II HDA mutants after four

days of dark treatment was noticeably pronounced as those displayed by the *phyA* and *phyB*

mutants. This behavior suggests that HDA15 and HDA18 may act downstream of the

phytochrome signaling pathway.

221   Furthermore, light hyposensitive phenotypes of T-DNA knockout mutants were

significantly prominent at low light intensities after 3 to 4 days of continuous light treatment.

Under far-red, red, and blue light treatments, four days of constant light treatment not only

induced long hypocotyls but also yielded bigger, more developed open cotyledons with longer

petioles indicating that Class II histone deacetylases might act as positive photomorphogenic

regulators. Hypocotyl length of mutant lines eventually leveled off with the wildtype on the fifth

day of light treatment, indicating that these histone deacetylases could be active at the early

stages of growth and development.

All in all, among all the Class II HDAs tested, *hda15-t27* mutants had the most prominent

hyposensitive phenotype in both light and dark conditions. Thus, this study focused on

elucidating the function of HDA15 in photomorphogenesis.

2. HDA15 targets the deacetylation of COP1 promoter

Like the long-hypocotyl phenotype displayed by the *hy5* mutant, we hypothesize that

HDA15 acts as a positive regulator of photomorphogenesis. To determine if HDA15 is indeed

involved in photomorphogenesis, expression levels of downstream positive regulators of

photomorphogenesis were assessed *via* RT-PCR and qPCR. As presented in Figure 2, among the

key positive regulators, which are known to be targeted by acetylation upon light exposure,

include HY5, LHB1B1, LHCA1, RBCS-1a, and 1b, CAB1, and CAB2 (Guo et al., 2008;

Charron et al., 2009). However, most of these genes, including HY5, were significantly reduced

in all the light treatments, which explain the long-hypocotyl phenotype suggesting that a negative

regulator catapulted the downregulation of these genes, possibly targeted by HDA15.

To further assess the regulatory role of HDA15, mRNA expression profiles of key

regulators of photomorphogenesis in *hda15* mutant and overexpression lines were conducted

(Figure 3). Compared to white light and far-red light treated plants, dark-treatment has eminently

amplified HY5 transcripts levels up to 7-folds in HDA15 overexpression lines. Concomitantly,

mRNA levels of COP1 were significantly elevated 3-folds in *hda15* mutants, while its

overexpression leads to COP1's extreme downregulation compared to wild type. ChIP assay

using anti-H3K9K14ac further reveals that the promoter and start sites of COP1 were highly

acetylated in the absence of HDA15 in dark-treated plants. The comparative acetylation profiles

suggest that HDA15 targets the deacetylation of COP1 in the dark.

3. Light signals HDA15 nuclear localization, dark treatment induces cytoplasmic

translocation

Hypocotyls of 4-day old HDA15-GFP transformants were used to determine its

localization (Figure 4). Under white light conditions, HDA15 remained to be prominently

nuclear. When grown under far-red light, HDA15-GFP concentrated in the nucleus, which is

similar to blue and red light treated lines. Consequently, seedlings grown in the dark exhibited

strong signals at the cytoplasmic area, although small quantities were still detected at the

nucleus. Similar results were observed when HDA15-GFP were transiently expressed in

protoplasts under various light treatment.

As mammalian Class II HDAs undergo nucleocytoplasmic shuttling, there is a possibility

that HDA15 may exhibit similar functional regulatory mechanisms. To illustrate whether

HDA15 undergoes nucleocytoplasmic shuttling, we determined the response of transfected

protoplasts after successive incubation under white light overnight, dark treatment, and light re-

exposure. As shown in Figure 5, HDA15-GFP transfected protoplasts exhibited nuclear

localization after 18 hours of white light incubation. Further dark treatment for 3 hours elicited

partial cytoplasmic translocation of HDA15-GFP. After one hour, re-exposure of the protoplasts

to white light led to its complete nuclear import. The observed response demonstrates that light

indeed drives the nucleocytoplasmic shuttling of HDA15.

4. COP1 triggers the nuclear localization of HDA15

Scanning through the amino acid sequence of HDA15, it contains a type par4 nuclear

localization signal (NLS) near the amino end, two overlapping bipartite NLS near the carboxyl

end, and a nuclear export signal (NES) near the C terminal half. Our previous study has

indicated that these signals navigate the subcellular compartmentalization of HDA15, and that

light, in general, drives the nucleocytoplasmic shuttling of HDA15 (Alinsug et al., 2012). Albeit

the exact mechanisms controlling its NLS and NES signals upon light exposure remains obscure.

Thus, we speculated that the only way for HDA15 to detect light is through the photoreceptors;

their inactivation upon the termination of light may have initiated HDA15's nuclear export.

To test this hypothesis, HDA15-GFP/YFP was transiently expressed in protoplasts of

mutant photoreceptors *phyA, phyB*, and *cry1xcry2*. As shown in Figure 6, all these mutant

photoreceptor protoplasts displayed nuclear localization of HDA15 suggesting that the nuclear

import of HDA15 is not dependent on light quality nor controlled singly by the

phytochrome/cryptochrome signaling cascades, but implicates the involvement of all

photoreceptors. The downstream target of all four photoreceptors, COP1, appears to have a

significant impact on HDA15's nuclear localization, where *cop1-4* mutant protoplasts exhibited

speckled distribution of HDA15 in the cytoplasm immediately near the vicinity of the nucleus.

Overall, our results suggested that the activation of photoreceptors did not trigger HDA15

nuclear localization. Instead, the shuttling observed was likely attributed to the presence of

COP1.

5. HDA15 directly interacts with COP1 inside the nucleus

The human Class II HDAC6 has been well studied to catalyze non-histone proteins such

as α-tubulin, cortactin, and HSP90, as well as, bind to ubiquitinated proteins inhibiting their

proteosomal degradation via its zinc finger ubiquitin-binding protein (ZnF-UBP) domain (Hook

et al., 2002; Kovacs et al., 2005; Luxton and Gundersen, 2007; Valenzuela-Fernandez et al.,

2008). Although the corresponding plant ortholog of HDAC6 is unclear, HDA15 contains a

RanBP zinc finger analogous to ZnF-UBP of HDAC6. To determine if HDA15 can associate

with non-histone proteins such as COP1, pull-down and BiFC assays were conducted (Figure 7).

Direct binding between HDA15 and COP1 was illustrated via pull-down assay using anti-His,

anti-GST, and anti-COP1 antibodies where HDA15 and COP1 were fused with GST- and His-

tags, respectively. This interaction was further confirmed *in vivo* using BiFC demonstrating their

direct binding inside the nucleus both under white light and dark conditions. Although it has

been consistently demonstrated that HDA15 is nuclear under all the light treatments, the

abrogation of COP1 in *cop1-4* mutant protoplasts renders it cytoplasmically speckled.

6. HDA15 positively regulates photomorphogenesis by repressing COP1

It has long been hypothesized that COP1 may recruit deacetylases or chromatin

remodeling factors that can repress transcription from target promoters (Holm and Deng, 1999).

If HDA15 functions as a co-repressor of COP1, then the mutant lines of HDA15 should exhibit

cop1-like phenotypes. On the contrary, *hda15-t27* lines manifested light hyposensitivity similar

to *hy5* mutants with down-regulated levels of HY5 and other positive regulators of

photomorphogenesis. Thus, HDA15 may indirectly regulate HY5 by repressing COP1, making

it a positive regulator of photomorphogenesis.

To establish the genetic relationship between HDA15 and COP1, we generated the

double mutant *cop1-4xhda15-t27*. As shown in Figure 8, the cross between *hda15-t27* and *cop1-*

4 mutants exhibited short hypocotyl phenotypes under white light and dark treatments and

dwarfed phenotypes in 3-week old plants all reminiscent of *cop1-4* mutants. These responses

suggest that COP1 is epistatic to HDA15. Therefore, HDA15 targets the repression of COP1 via

direct binding, thereby attenuating its capacity to ubiquitinate HY5 and its targets.

7. Ubiquitinated sites of HY5 and COP1 are less stable than HDA15

Considering that COP1 is an E3 ligase, the binding of HDA15 with COP1 raises pertinent

questions as to whether COP1 ubiquitinates HDA15 or HDA15 represses COP1.   To address

this, bioinformatics analysis using three algorithms, namely: Ub-Pred, BDM-PUB, and UbiSite,

were used to predict ubiquitination sites comparing HY5, COP1, and HDA15. Based on the

algorithms' consensus, as presented in Table 1, HY5 was found to contain five residues that are

actively targeted for ubiquitination, rendering it the most unstable among the three, with

instability index of 68.21. COP1 follows HY5 with an instability index of 47.13, with four

consensus residues prone to ubiquitination. On the other hand, it appears that HDA15 is the

most stable among the three proteins with an instability index of 37.11, noting that it only has

two residues as prospective sites for ubiquitination. This bioinformatics analysis hence reaffirms

our prior notion that HY5 is targeted for ubiquitination by COP1 in the dark during

skotomorphogenesis. With HDA15 being more stable than COP1, the direct interaction between

HDA15 and COP1 leads us to conclude that HDA15 represses COP1.

Discussion

Histone active marks such as H3K9K27ac signals transcriptional activation whereby light

induces the acetylation of positive photomorphogenic regulators functioning as chlorophyll a/b-

binding protein, peroxidase, citrate synthase, RBCS small chain 1B, ribosomal protein, PSII-

light-harvesting complex, HYH, HY5 and its downstream targets CHS, IAA3, RBCS-1a during

photomorphogenesis (Benhamed et al., 2006; Guo et al., 2008; Charron et al., 2009).   On the

other hand, histone deacetylation stimulates chromatin compaction triggering transcriptional

quiescence. Among the RPD3/HDA1-like superfamily of HDAs, Class I histone deacetylases

HDA6 and HDA19 display negative regulatory roles in photomorphogenesis. Together with

PHYB, Tessadori *et al.* have identified HDA6 as a regulator of light-induced chromatin

condensation (Tessadori et al., 2009). Similarly, HDA19 is suspected of repressing many light-

regulated genes by deacetylating H3K9K14 at the transcriptional start site of the phyA locus

(Benhamed et al., 2006; Jang et al., 2011).

On the other hand, it appears that Class II HDAs play a positive regulatory role in the

light signaling network with HDA15 repressing COP1 at its locus and modulating its activities

via direct interaction. As HDA5 and HDA18 also exhibit long-hypocotyl phenotypes at varying

light treatments, it remains a conundrum of how these and other Class II HDAs function during

photomorphogenesis. These findings fine-tune our prior understanding of the role of histone

modifications in light-regulated gene expression. Light not only induces the acetylation of key

positive regulators of photomorphogenesis but also renders the simultaneous repression of vital

negative regulators such as COP1 via direct interaction. Taken together, our results indicated

that histone acetyltransferases and deacetylases act as multi-functional and potent chromatin-

modifying enzymes during photomorphogenesis.

356   As light signals the nucleocytoplasmic shuttling of HDA15, its nuclear import is

357 primarily dependent on COP1. Since HDA15 have contrasting localization with COP1 in the

358 presence and absence of light, it is yet to be determined if a concentration gradient on either of

359 the proteins influences the movement of the other, assuming these enzymes are not degraded; or,

360 if a dominant ligand induces the degradation of the other. It would be interesting to unravel the

361 molecular underpinnings of the opposing nucleocytoplasmic translocation of HDA15 & COP1

362 upon light and dark signaling cues.   Although a significant amount of HDA15 translocates into

363 the cytoplasm, this nucleocytoplasmic shuttling suggests a functional regulatory mechanism for

364 deactivation both as a histone deacetylase and a direct repressor of COP1. As HDA15 have been

365 demonstrated to shuttle in and out of the nucleus with a functionally active nuclear export signal

366 and two nuclear localization signals, the direct association of HDA15 and COP1 may also

367 implicate the possibility of HDA15 acting as a chaperone to transport COP1 to the cytoplasm as

368 it lacks a nuclear export signal.

369   While other mutant Class II HDAs also manifest light hyposensitivity, only HDA15 is

370 driven by light to shuttle in and out of the nucleus. The environmental cue or stress that would

371 signal the nuclear import of the other Class II HDAs and their functional characterization remain

372 to be investigated as most of them have been observed to be cytoplasmic (Alinsug et al., 2012;

373 Tran et al., 2012).

374   Histone deacetylases constitute a family of nuclear enzymes known primarily for their

375 role in transcriptional repression. Class II HDAs, in particular, are known to be versatile

376 regulators catalyzing the deacetylation of both histones and non-histone proteins.   The

377 mammalian HDAC6 modulates a wide variety of cellular activities involving complex

378 interactions with α-tubulin, cortactin, HSP90, and other proteins (Matthias et al., 2008;

Valenzuela-Fernandez et al., 2008). Similarly, HDAC1 complexes with MDM2, an E3 ubiquitin

ligase, deacetylate p53, and promote its ubiquitination (Ito et al., 2002). In the case of

AtHDA15, it attenuates COP1 modulating its repressive activities via direct interaction,

subcellular compartmentalization, and chromatin compaction at its transcriptional start site.

Based on our results, this versatile regulator not only functions as a traditional histone

deacetylase silencing key regulators of photomorphogenesis, specifically COP1, but it also

directly interacts with COP1 likely attenuating its E3 ligase activities inside the nucleus.

The specific molecular mechanisms for the degradation and/or cytoplasmic export of

COP1 in the nucleus under light conditions remain obscure. The nuclear interaction of HDA15

with COP1 under light conditions may support the notion that COP1's depletion may be

enhanced by the binding of HDA15 to induce auto-ubiquitination. However, based on the

Western blot, the mutant nor overexpression lines of HDA15 neither influenced COP1's protein

levels.   Instead, HDA15 attenuates the E3 ligase activity of COP1 that prevents the

ubiquitination of HY5. This action explains the downregulation of HY5 protein concentrations

in *hda15* lines and the elevated levels of the protein in HDA15-OE lines.

On a similar note, in the dark, where COP1 is active and nuclear, partial concentrations of

HDA15 are inactivated via cytoplasmic importation. HDA15 in the nucleus may modulate the

activities of COP1 and target the deacetylation of its transcriptional start site. Presented in

Figure 9 is a working model depicting the sub-compartmentalization and modularized activities

of COP1 and HDA15 under light and dark conditions.

In light, COP1 is mainly cytoplasmic, although some trace amounts are left inside the

nucleus. When HDA15 shuttles into the nucleus, it binds to COP1 and conceivably leads to the

abrogation of HY5's ubiquitination regulating photomorphogenesis. In the dark, contrarily,

COP1 exclusively localizes inside the nucleus and targets the ubiquitination of HY5 to advance

skotomorphogenesis. While the majority of HDA15 translocates into the cytoplasm, partial

amounts left in the nucleus target the deacetylation of the transcriptional start site of COP1 locus

and/or interact with COP1 to modulate its repressive activities. The molecular switches in

apportioning the exact amounts of HDA15 to remain active inside the nucleus while others are

deactivated inducing their nuclear export remains elusive and requires further investigation.

We also attempted to determine the binding sites of the two repressors by generating

different truncated constructs of COP1 and HDA15 containing various combinations of potential

protein-binding domains. Surprisingly, only the full-length HDA15 was able to bind the full-

length COP1 suggesting strict, proper protein conformation for such interaction.

Previous studies on the conserved domains of RPD3/HDA1-like superfamily show that

Zn finger domains are imprinted on several Class II HDAs (Pandey et al., 2002; Alinsug et al.,

2009). The monocot ortholog of AtHDA15 in maize, ZmHDA110, has a C3HC4 type RING Zn

finger that binds with ubiquitin while the ZnF-UBP of the mammalian Class II HDAC6 binds to

ubiquitin specifically at the C termini diglycine motif catalyzed by a deubiquitinase, ataxin-3

(Ouyang et al., 2012). RanBP-ZnF in HDA15 may similarly interact with ubiquitin, such as

those found in Npl4 (Alam et al., 2004). Based on sequence analyses on Zn fingers conducted

by Wang *et al.*, the molecules have highly conserved cysteine residues and other key motifs in

binding ubiquitin (Wang et al., 2003). As illustrated in Figure 10, sequence comparison between

HDA15's RanBP2-ZnF to the human Np14 (NZF) reveals the presence of the two canonical

highly-conserved cysteine residues and a key ubiquitin-binding motif, Thr_{95}. Considering the

RING-ZnF domain of COP1 catalyzes the transfer of ubiquitin units from E2 transferases to

target proteins such as HY5, we speculate that ubiquitin may constitute a recognition site for

HDA15 to identify COP1 as an active E3 ligase.

Nevertheless, unlike HDAC6, which binds and transports ubiquitinated proteins to

aggresomes, the binding of HDA15 to COP1 undertakes a regulatory role via the attenuation of

the latter's E3 ligase activity. Based on bioinformatics analysis comparing the ubiquitination

sites of HY5, COP1, and HDA15, it is predicted that HDA15 is more stable than COP1. As

expected, HY5 is the least stable making it more prone to ubiquitination. Still, questions remain

on the exact nature of binding of HDA15 to ubiquitin and/or polyubiquitin, as well as the

properties of the molecular switches downstream of the HDA15-COP1 complex.

Acknowledgments

The authors are grateful to the unwavering support of the various laboratory professors, mentors,

colleagues, and staff for the conduct of this research. Special mention goes to Prof. Keqiang Wu

and Prof. Hsu-Liang Hsieh of NTU and Prof. Jeong Joo Cheong of SNU for their guidance and

access to their facilities. MVA is equally indebted to the scholarship and research fellowship

grants funded by the Democratic Pacific Union, National Science Council of Taiwan, National

Taiwan University, and Seoul National University.

Author Contribution

MVA conceptualized the study, screened and generated double mutant lines for phenotypes,

conducted light experiments, ChIP and BiFC assay, and wrote the manuscript. CD performed

pull down assay, mRNA extraction, bioinformatics analyses, and wrote a portion of the

manuscript.

Figure 1. Light hyposensitivity in Class II HDAs (A) Hypocotyl lengths of WT, *hda5*, *hda15* and *hda18* mutants under dark and white light were measured for 7 consecutive days. (B) Hypocotyl lengths of Class II HDA mutants under far red, red and blue light treatments at different light intensities were observed for 5 days. Light intensity treatments were designated as low (L, upper panel) or high (H, lower panel): far red, H at 18.065 and L at 2.77 μmol m^{-2} s^{-1}; red light, H at 22.938 and L at 1.77 μmol m^{-2} s^{-1}; and blue light, H at 11.095 and L at 3.84 μmol m^{-2} s^{-1}. Data presented are mean values with error bars set from its standard deviation.

Figure 2. Down-regulation of positive regulators to photomorphogenesis in AtHDA15 knockout lines. mRNA expression levels of positive regulators of photomorphogenesis were assessed in *A. thaliana* exposed to (A) white light, (B) far-red light, and (C) the dark. Expression levels reflected were normalized to ubiquitin.

Figure 3. mRNA expression profiles of key regulators of photomorphogenesis in HDA15 mutant and overexpression lines. (A) Schematic diagram of repressive action of HDA15 on COP1 locus. (B) ChIP assay using anti-H3K9K14ac further reveals that the promoter & start sites of COP1 were highly acetylated in the absence of HDA15 in dark-treated plants. (C) Compared to white light and far red light treated plants, dark-treatment has eminently amplified HY5 transcripts in HDA15 overexpression lines. Concomitantly, mRNA levels of COP1 were significantly elevated in *hda15* mutants while its overexpression lead to COP1's extreme downregulation.

Figure 4. Subcellular localization of AtHDA15 under different light radiance with wildtype and mutant photoreceptors. Localization HDA15-GFP/YFP was localized in the nuclei of hypocotyls and protoplasts of transgenic lines grown under white light and different colored light treatments of A. thaliana with wild-type photoreceptors. Note: In dark treated plants, HDA15 has a pan-cytoplasmic distribution. Hoechst was used as a nuclear marker.

Figure 5. Time-course monitoring of nucleocytoplasmic shuttling and pan-cytoplasmic scattering of HDA15. Transfected protoplasts were incubated under white light overnight then covered with foil for 3 hours and re-exposed to white light. HDA15-GFP transfected protoplasts exhibited nuclear localization after 18h of white light incubation. Further dark treatment for 3h elicited partial cytoplasmic translocation of HDA15-GFP. Re-exposition to white light after 1h lead to its complete nuclear import. VirD2NLS was co-transfected as nuclear marker (blue).

Figure 6. COP1 activates nuclear localization of HDA15. HDA15 localization in protoplasts of mutant photoreceptors were all nuclear. However, when transiently expressed in *cop1-4* mutant protoplasts, HDA15 exhibited cytoplasmic localization. VirD2-NLS was co-transfected as nuclear marker (blue).

Figure 7. Binding to COP1 is associated with the nucleocytoplasmic shutting of AtHDA15. **(A)** *In vitro* and **(B)** i*n vivo* interaction between COP1 and AtHDA15 indicates their direct binding inside the nucleus. AtHDA15-COP1 interaction was further confirmed via pull down assay using anti-GST antibody where HDA15 was fused with GST and COP1 with His tags. Interaction of COP1 and AtHDA15 in isolated protoplasts was visualized 18–24 hr after transfection using bifluorescence complementation assay under laser confocal microscopy. AtHDA15 and COP1 genes were recombined with N-YFP and C-YFP. VirD2-NLS was used as nuclear marker.

Figure 8. *hda15-t27xcop1-4* phenotype. The cross between *hda15-t27* and *cop1-4* mutants exhibited short hypocotyl phenotypes under white light (A) and dark (B) treatment for 4 days and dwarfed phenotypes in 3-week old plants (C) which are reminiscent of *cop1-4* mutants.

Figure 9. Working hypothesis illustrating sub-compartmentalization and modular activities of COP1 and HDA15 under light and dark conditions. In the presence of light, COP1 is cytoplasmic with some trace amounts left inside the nucleus. On the contrary, HDA15 concentrates inside the nucleus binding to nuclear COP1, which prevents the ubiquitination of HY5 thereby positively regulating photomorphogenesis. In the dark, however, COP1 exclusively localizes inside the nucleus targeting the ubiquitination of HY5 proceeding to skotomorphogenesis. On the other hand, majority of HDA15 translocates into the cytoplasm with some minimal amounts left in the nucleus. HDA15 directly associates with COP1 to modulate its repressive activities and targets its deacetylation at the promoter and start sites leading to transcriptional quiescence.

NZF $_3$TSAMWACQHC*TFMNQPGTGHCEMCSLPRT$_{31}$

RanBP2   KEGQWDCSACLVQNEGSSTKCAACQNPRK

HDA15-RanBP2 $_{86}$FVKWCCVNCTMSNPGDMVHCCICGEHKES$_{115}$

Figure 10. Sequence comparison between Zn fingers in human Npl4 (NZF), wild type RanBP2 which has minimal Ubq binding, and RanBP2 in AtHDA15. Conserved residues in NZF was defined by Wang *et al* (2003) where blue letters indicate semi-conserved residues, green cysteine residues are highly conserved and contact Ubq, and orange residues as key motifs in binding Ubq.

Parsed Citations

Akimov, V., Barrio-Hernandez, I., Hansen, S.V., Hallenborg, P., Pedersen, A.-K., Bekker-Jensen, D.B., Puglia, M., Christensen, S.D., Vanselow, J.T., and Nielsen, M.M. (2018). UbiSite approach for comprehensive mapping of lysine and N-terminal ubiquitination sites. Nature structural & molecular biology 25, 631-640.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Alam, S.L., Sun, J., Payne, M., Welch, B.D., Blake, B.K., Davis, D.R., Meyer, H.H., Emr, S.D., and Sundquist, W.I. (2004). Ubiquitin interactions of NZF zinc fingers. The EMBO journal 23, 1411-1421.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Alinsug, M.V., Yu, C.W., and Wu, K. (2009). Phylogenetic analysis, subcellular localization, and expression patterns of RPD3/HDA1 family histone deacetylases in plants. BMC plant biology 9, 37.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Alinsug, M.V., Chen, F.F., Luo, M., Tai, R., Jiang, L., and Wu, K. (2012). Subcellular localization of class II HDAs in Arabidopsis thaliana: nucleocytoplasmic shuttling of HDA15 is driven by light. PloS one 7, e30846.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ausin, I., Alonso-Blanco, C., and Martinez-Zapater, J.-M. (2004). Environmental regulation of flowering. International Journal of Developmental Biology 49, 689-705.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Benhamed, M., Bertrand, C., Servet, C., and Zhou, D.X. (2006). Arabidopsis GCN5, HD1, and TAF1/HAF2 interact to regulate histone acetylation required for light-responsive gene expression. The Plant cell 18, 2893-2903.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Benhamed, M., Martin-Magniette, M.L., Taconnat, L., Bitton, F., Servet, C., De Clercq, R., De Meyer, B., Buysschaert, C., Rombauts, S., Villarroel, R., Aubourg, S., Beynon, J., Bhalerao, R.P., Coupland, G., Gruissem, W., Menke, F.L., Weisshaar, B., Renou, J.P., Zhou, D.X., and Hilson, P. (2008). Genome-scale Arabidopsis promoter array identifies targets of the histone acetyltransferase GCN5. The Plant journal : for cell and molecular biology 56, 493-504.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Charron, J.B., He, H., Elling, A.A., and Deng, X.W. (2009). Dynamic landscapes of four histone modifications during deetiolation in Arabidopsis. The Plant cell 21, 3732-3748.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gendrel, A.V., Lippman, Z., Martienssen, R., and Colot, V. (2005). Profiling histone modification patterns in plants using genomic tiling microarrays. Nature methods 2, 213-218.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gu, D., Chen, C.Y., Zhao, M., Zhao, L., Duan, X., Duan, J., Wu, K., and Liu, X. (2017). Identification of HDA15-PIF1 as a key repression module directing the transcriptional network of seed germination in the dark. Nucleic acids research 45, 7137-7150.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Guo, L., Zhou, J., Elling, A.A., Charron, J.B., and Deng, X.W. (2008). Histone modifications and expression of light-regulated genes in Arabidopsis are cooperatively influenced by changing light conditions. Plant physiology 147, 2070-2083.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Guruprasad, K., Reddy, B.B., and Pandit, M.W. (1990). Correlation between stability of a protein and its dipeptide composition: a novel approach for predicting in vivo stability of a protein from its primary sequence. Protein Engineering, Design and Selection 4, 155-161.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Holm, M., and Deng, X.W. (1999). Structural organization and interactions of COP1, a light-regulated developmental switch. Plant molecular biology 41, 151-158.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hook, S.S., Orian, A, Cowley, S.M., and Eisenman, R.N. (2002). Histone deacetylase 6 binds polyubiquitin through its zinc finger (PAZ domain) and copurifies with deubiquitinating enzymes. Proceedings of the National Academy of Sciences of the United States of America 99, 13425-13430.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hsieh, H.L., Okamoto, H., Wang, M., Ang, L.H., Matsui, M., Goodman, H., and Deng, X.W. (2000). FIN219, an auxin-regulated gene, defines a link between phytochrome A and the downstream regulator COP1 in light control of Arabidopsis development. Genes & development 14, 1958-1970.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ito, A, Kawaguchi, Y., Lai, C.H., Kovacs, J.J., Higashimoto, Y., Appella, E., and Yao, T.P. (2002). MDM2-HDAC1-mediated deacetylation of p53 is required for its degradation. The EMBO journal 21, 6236-6245.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Jang, I.C., Chung, P.J., Hemmes, H., Jung, C., and Chua, N.H. (2011). Rapid and reversible light-mediated chromatin modifications of Arabidopsis phytochrome A locus. The Plant cell 23, 459-470.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Jiao, Y., Lau, O.S., and Deng, X.W. (2007). Light-regulated transcriptional networks in higher plants. Nature reviews. Genetics 8, 217-230.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kovacs, J.J., Cohen, T.J., and Yao, T.P. (2005). Chaperoning steroid hormone signaling via reversible acetylation. Nuclear receptor signaling 3, e004.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kumar, D., and Kumar, P. (2019). An In-Silico Investigation of Key Lysine Residues and Their Selection for Clearing off Aβ and Holo-AβPP Through Ubiquitination. Interdisciplinary Sciences: Computational Life Sciences 11, 584-596.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kurdistani, S.K., Tavazoie, S., and Grunstein, M. (2004). Mapping global histone acetylation patterns to gene expression. Cell 117, 721-733.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Li, B., Krishnan, V.G., Mort, M.E., Xin, F., Kamati, K.K., Cooper, D.N., Mooney, S.D., and Radivojac, P. (2009). Automated inference of molecular mechanisms of disease from amino acid substitutions. Bioinformatics 25, 2744-2750.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Liu, X., Wei, W., Zhu, W., Su, L., Xiong, Z., Zhou, M., Zheng, Y., and Zhou, D.X. (2017). Histone Deacetylase AtSRT1 Links Metabolic Flux and Stress Response in Arabidopsis. Molecular plant 10, 1510-1522.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Liu, X., Chen, C.Y., Wang, K.C., Luo, M., Tai, R., Yuan, L., Zhao, M., Yang, S., Tian, G., Cui, Y., Hsieh, H.L., and Wu, K. (2013). PHYTOCHROME INTERACTING FACTOR3 associates with the histone deacetylase HDA15 in repression of chlorophyll biosynthesis and photosynthesis in etiolated Arabidopsis seedlings. The Plant cell 25, 1258-1273.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Luxton, G.G., and Gundersen, G.G. (2007). HDAC6-pack: cortactin acetylation joins the brew. Developmental cell 13, 161-162.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ma, X., Lv, S., Zhang, C., and Yang, C. (2013). Histone deacetylases and their functions in plants. Plant cell reports 32, 465-478.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Matthias, P., Yoshida, M., and Khochbin, S. (2008). HDAC6 a new cellular stress surveillance factor. Cell cycle 7, 7-10.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Osterlund, M.T., Hardtke, C.S., Wei, N., and Deng, X.W. (2000). Targeted destabilization of HY5 during light-regulated development of Arabidopsis. Nature 405, 462-466.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ouyang, H., Ali, Y.O., Ravichandran, M., Dong, A, Qiu, W., MacKenzie, F., Dhe-Paganon, S., Arrowsmith, C.H., and Zhai, R.G. (2012). Protein aggregates are recruited to aggresome by histone deacetylase 6 via unanchored ubiquitin C termini. The Journal of biological chemistry 287, 2317-2327.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pandey, R., Muller, A, Napoli, C.A, Selinger, D.A, Pikaard, C.S., Richards, E.J., Bender, J., Mount, D.W., and Jorgensen, R.A. (2002). Analysis of histone acetyltransferase and histone deacetylase families of Arabidopsis thaliana suggests functional diversification of chromatin modification among multicellular eukaryotes. Nucleic acids research 30, 5036-5055.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Schenke, D., Cai, D., and Scheel, D. (2014). Suppression of UV-B stress responses by flg22 is regulated at the chromatin level via histone modification. Plant, cell & environment 37, 1716-1721.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Schübeler, D., MacAlpine, D.M., Scalzo, D., Wirbelauer, C., Kooperberg, C., van Leeuwen, F., Gottschling, D.E., O'Neill, L.P., Turner, B.M., and Delrow, J. (2004). The histone modification pattern of active genes revealed through genome-wide chromatin analysis of a higher eukaryote. Genes & development 18, 1263-1271.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Tang, Y., Liu, X., Liu, X., Li, Y., Wu, K., and Hou, X. (2017). Arabidopsis NF-YCs Mediate the Light-Controlled Hypocotyl Elongation via Modulating Histone Acetylation. Molecular plant 10, 260-273.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Tessadori, F., van Zanten, M., Pavlova, P., Clifton, R., Pontvianne, F., Snoek, L.B., Millenaar, F.F., Schulkes, R.K., van Driel, R., Voesenek, L.A, Spillane, C., Pikaard, C.S., Fransz, P., and Peeters, A.J. (2009). Phytochrome B and histone deacetylase 6 control light-induced chromatin compaction in Arabidopsis thaliana. PLoS genetics 5, e1000638.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Tran, H.T., Nimick, M., Uhrig, R.G., Templeton, G., Morrice, N., Gourlay, R., DeLong, A, and Moorhead, G.B. (2012). Arabidopsis thaliana histone deacetylase 14 (HDA14) is an alpha-tubulin deacetylase that associates with PP2A and enriches in the microtubule fraction with the putative histone acetyltransferase ELP3. The Plant journal : for cell and molecular biology 71, 263-272.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Valenzuela-Fernandez, A, Cabrero, J.R., Serrador, J.M., and Sanchez-Madrid, F. (2008). HDAC6: a key regulator of cytoskeleton, cell migration and cell-cell interactions. Trends in cell biology 18, 291-297.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Wang, B., Alam, S.L., Meyer, H.H., Payne, M., Stemmler, T.L., Davis, D.R., and Sundquist, W.I. (2003). Structure and ubiquitin interactions of the conserved zinc finger domain of Npl4. The Journal of biological chemistry 278, 20225-20234.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yang, J.Y., Iwasaki, M., Machida, C., Machida, Y., Zhou, X., and Chua, N.H. (2008). betaC1, the pathogenicity factor of TYLCCNV, interacts with AS1 to alter leaf development and suppress selective jasmonic acid responses. Genes & development 22, 2564-2577.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yoo, S.D., Cho, Y.H., and Sheen, J. (2007). Arabidopsis mesophyll protoplasts: a versatile cell system for transient gene expression analysis. Nature protocols 2, 1565-1572.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zhao, L., Peng, T., Chen, C.Y., Ji, R., Gu, D., Li, T., Zhang, D., Tu, Y.T., Wu, K., and Liu, X. (2019). HY5 Interacts with the Histone Deacetylase HDA15 to Repress Hypocotyl Cell Elongation in Photomorphogenesis. Plant physiology 180, 1450-1466.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zhou, R., Han, L., Li, G., and Tong, T. (2009). Senescence delay and repression of p16INK4a by Lsh via recruitment of histone deacetylases in human diploid fibroblasts. Nucleic acids research 37, 5183-5196.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title