*Research Article*
*Prophylactic efficacy of riluzole against anxiety- and depressive-like behaviours in two rodent stress models*

Yashika Bansal[¥,a], Corey Fee[¥,a,b], Keith A. Misquitta[a,b], Sierra Codeluppi[a,b], Etienne Sibille, PhD[a,b,c], Robert M. Berman, MD[d], Vladimir Coric, MD[d], Gerard Sanacora, PhD, MD[e], and Mounira Banasr, PhD*[a,b,c,e]

[a]Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, Toronto, ON, Canada
[b]Department of Pharmacology and Toxicology, University of Toronto, Toronto, ON, Canada
[c]Department of Psychiatry, University of Toronto, Toronto, ON, Canada
[d]BioHaven Pharmaceuticals, New Haven, CT, USA
[e]Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA

**Running title**: Prophylactic riluzole in two stress models
[¥]**Shared First Author**
*****Corresponding author:** Mounira Banasr, PhD, CAMH, 250 College street, Toronto, ON M5T 1R8, Canada Tel: 416-535-8501, ext 32319; Email: Mounira.banasr@camh.ca

**Word Count:** 4949

**Number of Figures:** 3

**Number of Tables:** 0

**Keywords:** Riluzole, Chronic Stress, Prophylactic, Depression, Antidepressants

**Abstract**

**Background**: Chronic stress-related illnesses, such as major depressive disorder and post-traumatic stress disorder share symptomatology, including anxiety, anhedonia, and helplessness. Across disorders, neurotoxic dysregulated glutamate (Glu) signaling may underlie symptom emergence. Current first-line antidepressant drugs, which do not directly target Glu signaling, fail to provide adequate benefit for many patients and are associated with high relapse rates. Riluzole modulates glutamatergic neurotransmission by increasing metabolic cycling and modulating signal transduction. Clinical studies exploring riluzole's efficacy in stress-related disorders have provided varied results. However, the utility of riluzole for treating specific symptom dimensions or as a prophylactic treatment has not been comprehensively assessed.

**Methods:** We investigated whether chronic prophylactic riluzole (~12-15/kg/day p.o.) could prevent the emergence of <u>behavioural</u> deficits induced by unpredictable chronic mild stress (UCMS) in mice. We assessed: i) anxiety-like behaviour using the elevated-plus maze, open field test, and novelty-suppressed feeding, ii) mixed anxiety/anhedonia-like behaviour in the novelty-induced hypophagia test and, iii) anhedonia-like behaviour using the sucrose consumption test. Z-scoring summarized changes across tests measuring similar dimensions. In a separate learned helplessness (LH) cohort, we investigated whether chronic prophylactic riluzole treatment could block the development of helplessness-like behaviour.

**Results**: UCMS induced an elevation in anxiety-, anhedonia-like behaviour, and overall behavioural emotionality that was blocked by prophylactic riluzole. In the LH cohort, prophylactic riluzole blocked the development of helplessness-like behaviour.

**Conclusion**: This study supports the utility of riluzole as a prophylactic medication for preventing anhedonia, anxiety, and helplessness symptoms associated with stress-related disorders.

**Introduction**

Chronic stress is the primary risk factor for a variety of psychiatric disorders including major depressive disorder (MDD), bipolar disorder (BPD), schizophrenia, and post-traumatic stress disorder (PTSD) [1–3]. Together, these disorders which share several common symptoms including low mood, anhedonia and helplessness [4] account for one-third of global years of productivity loss due to disability [5]. Notably, the burden of these disorders has substantially increased over the last couple of years due do to the global COVID-19 pandemic [6, 7]. selective serotonin reuptake inhibitors (SSRIs) serve as first-line treatment approaches for several of these disorders [8–10]. Indeed, antidepressant drugs (ADs) have demonstrated efficacy in treating mood and anxiety related symptoms across these disorders and provide neuroprotection in stress-related illnesses [11]. However, they require several rounds of treatment to achieve remission in a majority of patients, thus exhibiting lengthy delays until remission is achieved [12, 13]. Moreover, in PTSD only 59% of individuals receiving SSRI treatment responded after fourteen weeks [14]. Other pharmacotherapies for stress-related illnesses include selective noradrenergic reuptake inhibitors, tricyclic antidepressants and anti-adrenergic agents [15–17]. Unfortunately, approximately a third of MDD patients do not achieve remission despite multiple medication trials [18]. The limited therapeutic efficacy of these drugs points to the need for more effective therapeutics or potential strategies to prevent the onset of mental illnesses. Individuals respond differently to stressful life events [3, 19–21] with some showing resiliency successfully. Adapting to the stressful situations), and others show susceptibility developing pathological states such as MDD or PTSD. Similar resiliency and susceptibility to stress can be found in rodent models of chronic stress exposure [22–25]. Preclinical studies have focused on leveraging these individual differences to address the maladaptive response to stress and the associated symptom emergence by developing pharmaceuticals with prophylactic properties [26–29].

Therefore, several drugs have been investigated for potential prophylactic effects. One such drug is fluoxetine but unfortunately, its administration has little to no prophylactic efficacy in rodent stress models [30]. However, pretreatment with serotonin receptor 4 agonists or ketamine showed effective prophylactic properties [31]. Specifically, pretreatment with ketamine, the glutamate (Glu) *N*-methyl-D-aspartate (NMDA) receptor antagonist was shown to attenuate fear response in the contextual fear conditioning [29] and prevent induction of depressive-like behaviour in the chronic social-defeat stress, learned helplessness (LH), and chronic corticosterone models [30, 32–34]. Theoretically, the use of prophylactic drugs could prevent chronic stress-related pathogenesis in individuals and be beneficial in clinical settings, in particular in patients who suffered one depressive or PTSD episode as well as MDD or BPD patients who discontinue their treatment therefore patients that are at high risk for relapse [35–38]. Other populations that could benefit from prophylactic treatment include individuals exposed to acute trauma and who are at high risk for developing PTSD [39]. This is accordance with findings of the literature suggesting that some treatments such as lithium maintenance therapy can increase resilience in clinical MDD and BPD patients [40].

Extensive evidence from both preclinical and clinical studies indicate abnormalities of glutamatergic neurotransmission in the development of psychiatric disorders [32, 40–43]. Specifically, altered regulation of Glu metabolism and neurotransmission and changed expression/function of Glu receptors have been reported in MDD and PTSD patients, as well as in postmortem subjects [44–47]. Similar glutamatergic dysfunctions are found in rodent models of chronic stress [48]. Specifically, stress exposure induce Glu release in corticolimbic brain regions, including the prefrontal cortex (PFC), hippocampus (HPC), and amygdala [49, 50]. Elevated or sustained Glu levels induced by chronic stress can causes excitotoxic effects via

enhanced extrasynaptic signaling activity [48, 51, 52]. These alterations have been linked to maladaptive changes in the structure and function of neurons [53, 54] and glial cells [55–58]. Glial cells such as astrocytes play a pivotal role in the homeostatic response to the stress-induced increases in glutamatergic neurotransmission [48, 57, 59, 60] as they are responsible for over 80% of Glu reuptake, Glu synthesis and metabolism [61–63]. In light of this, drugs that enhance Glu clearance may prevent the deleterious effects of excess Glu [64–67].

Riluzole, a FDA-approved drug for amyotrophic lateral sclerosis, provides neuroprotection by enhancing synaptic reuptake of elevated excitatory amino acid transporters (EAATs) present on the astrocytes; thereby increasing the ratio of synaptic to extra-synaptic signaling which further promotes neurotrophic factor expression [40, 68, 69]. Indeed, in chronic stress- or glucocorticoid-exposed rodents, chronic riluzole treatment reversed depressive-like behaviour [56, 70]. Chronic riluzole administration also attenuated the structural and functional deficits in PFC and HPC as indicated by markers of synaptic Glu signaling and increased levels of brain derived neurotropic factor (BDNF) [56, 70]. Riluzole shares neurotrophic effects with conventional antidepressants and its modulation of Glu signaling with rapid-acting anti-depressants, such as the NR2B-containing NMDAR antagonist ketamine [71]. Moreover, early open-label studies in MDD patients suggested riluzole possessed antidepressant properties when used either as monotherapy or as adjunctive treatment of monoaminergic antidepressants [72–75]. Further, a randomized, double-blind, placebo-controlled (RDBPC) clinical trial in hospitalised MDD patients showed the combination of riluzole plus citalopram to be significantly more efficacious and to have a faster onset of action compared to placebo plus [76]. However it is important to mention that the largest randomized placebo-controlled study examining the efficacy of rilzuole augmentation of standard antidepressants in the treatment of

MDD did not find evidence of riluzole benefit and a recent study of 74 patients diagnosed PTSD failed to demonstrate efficacy on the Clinicin Administered PTSD Scale, the primary outcome measure in the study [77-78]. This discrepancy may be due to difference in study design, patient history (i.e. length, severity, prior drug administration) or the timing of treatment intervention in relation to the illness, or the result of under-powered studies.

Recent studies in rodent models of other pathologies such as spinal cord injury [79] or autoimmune encepthalomyelitis [80] suggest that riluzole treatment might be more effective at reducing behavioural and cellular deficits associated with these models when administered prophylactically. Thus, in this study we investigated if riluzole confers enhanced stress resilience when administered prophylactically, i.e., concurrent with unpredictable chronic mild stress (UCMS), while specifically assessing anxiety-, anhedonia-like behaviours, and overall behavioural emotionality. In a separate cohort, we also investigated whether chronic preventative riluzole treatment could block the development of helplessness, a core feature of PTSD and other mood disorders. Given that chronic stress and learned helplessness paradigms are well-documented models that appear recapitulate the behavioural and glutamatergic tripartite synapse deficits akin to those observed in humans with stress-related illnesses [48], we predicted that the Glu-modulating drug riluzole would prevent the emergence of anxiety-, anhedonia-, and helplessness-like behaviour.

**Methods**

**Animals**

Eight-weeks old male C57BL/6 mice (Jackson Laboratory, Saint Constant, QC) were housed under a 12hr light/dark cycle at constant temperature (20-23°C) with *ad libitum* access to food and water, except when subjected to light disturbances or food/water deprivation during

unpredictable chronic mild stress (UCMS) or learned helplessness (LH) procedures. Following a 2-week facility habituation, animals were single-housed and randomly assigned to 1 of 4 treatment groups: control+vehicle, control+riluzole, UCMS+vehicle, or UCMS+riluzole (*n*=10/group). A separate cohort of 5-week-old male C57BL/6 mice (Jackson Laboratory, Bar Harbor, ME) were habituated to the facility under group-housed conditions throughout the experiment. LH mice were randomly assigned to 1 of 2 treatment groups: LH+vehicle or LH+riluzole (*n*=10/group). All testing was performed during the animal's light cycle and in accordance with Institutional and Canadian Council on Animal Care (CCAC) guidelines.

**Drug Administration**

To increase palatability of oral drug treatment, riluzole was diluted into 0.3% (v/v) saccharin (Sigma Aldrich, St. Louis, MO) in drinking water at a concentration of 60ug/mL, translating to ~12-15mg/kg/day p.o. for mice weighing 22-30g and drinking 6-8mL/day, following a previous study [70]. Riluzole was dissolved by stirring in the saccharin solution at room temperature. Solutions were changed every 48-72hrs. Mice not treated with riluzole consumed saccharin alone (vehicle). To assess the prophylactic efficacy of riluzole in UCMS, drug administration commenced on the same day as stressors. To assess preventative efficacy in LH, riluzole commenced 2 weeks prior to the inescapable footshock session. Both cohorts were maintained on riluzole throughout the duration of the experiments, except when fluid deprivation was required for the sucrose and water consumption tests.

**Unpredictable Chronic Mild Stress (UCMS) Procedure**

UCMS mice were subjected to 5 weeks of randomized stressors (2-4/day) and maintained for 2 weeks during testing (1-2/day) based on previously validated methods [81–83]. Stressors included: forced bath (~2cm of water in rat cage for 15min), wet bedding (30min-1h), aversive

predator odor (20min-1h exposure to fox urine), light cycle reversal or disruption, social stress (rotate mice between cages within drug groups), tilted cage ($45^0$ tilt up to 2h), reduced space (dividers in cages up to 2h), acute restraint (15-30min in 50mL falcon tube with air/tail holes), bedding change (replacing soiled bedding with clean bedding), bedding or nestlet removal (up to 2h). Control animals underwent regular handling.

**Behavioural Tests**

For the UCMS cohort, behavioural testing commenced in week 3. Tests assessed locomotor activity (LA), anxiety-like behaviours in the elevated plus-maze (EPM), open field test (OFT) and novelty-suppressed feeding (NSF) tests, anhedonia-like behaviour in the sucrose consumption test (SCT), or both in the novelty-induced hypophagia test (NIH). Physiological deficits were tracked with weekly fur coat assessment during 5 weeks of UCMS exposure and drug treatment. Testing order was randomized, and experimenters were blinded to the treatment history.

      **Locomotor Activity (LA):** On day 21, LA was measured for 20min in a regular home cage (26x15x12cm) using an aerial-mounted video camera. ANY-maze tracking software (Stoelting Co., Wood Dale, IL) extracted distance traveled. LA was assessed to exclude the possibility that changes observed in other behavioural tests were due to bias introduced by treatment-related alterations in ambulatory movement.

      **Elevated Plus Maze (EPM):** On day 22, animal behaviour was assessed in a plus shaped maze consisting of 4 white Plexiglas arms, two open arms (27x5cm) and two enclosed arms (27x5x15cm) situated 55cm high with similar arms faced opposite each other. Animals freely explored the EPM for 10min in a 20-lux room. An aerial-mounted camera/ANY-maze software measured the time spent (sec) and number of entries in open and closed arms.

**Open Field Test (OFT):** On day 23, behaviour was assessed in an open field (70x70x33cm) placed on the floor. An aerial-mounted camera/ANY-maze software divided the field into 3 concentric square zones of equal area, and measured time spent and number of entries into the innermost zone (40x40cm).

**Novelty-suppressed Feeding (NSF):** On day 25, following 16h food deprivation, animals were introduced to a novel arena (62x31x48cm) containing a single food pellet. An experimenter blind to treatment conditions measured latency to feed on the pellet during a 12min period. As a control for differences in appetitive drive, latency to feed was then measured in the animal's home cage during a 6 min period.

**Novelty-Induced Hypophagia (NIH):** On days 25-26, animals were habituated to a daily liquid reward (1mL of 1/3 v/v sweetened condensed milk). On day 26, latency to drink the reward was measured in the home cage as a control for potential treatment effects on appetite or activity. The following day, the animals were moved to a new cage (26x15x12cm) in a brightly lit (500-600 lux) room and the latency to drink the milk was measured over a 6min trial as a measure for hyponeophagia.

**Sucrose Consumption Test (SCT):** On days 27-28, home cage water bottles were switched to sucrose solution (1%; Sigma, St. Louis, MO) ± riluzole. Animals were habituated to sucrose for 48h and then fluid deprived overnight (~14h). On day 29, sucrose intake (mL) was measured over a 1h test. After the SCT, animals were returned to regular solutions (saccharine±riluzole) for 24h. The same test was repeated the next day i.e. fluid deprivation (14h) and water intake (1h) to control for potential treatment effects on fluid intake.

**Fur Coat Assessment:** The coat state of each animal was assessed on 7 anatomical areas by attributing a score of 0, 0.5, or 1 for each, from maintained to unkempt. Coat assessment was scored by an experimenter blind to treatment conditions every week for 5 weeks of UMCS.

## Learned Helplessness (LH) Procedure

A separate cohort of animals was habituated to a shuttle box (Med Associates, St. Albans, VT) for 5min and underwent a session consisting of 60 randomized, unpredictable, and inescapable footshocks (0.35 mA) administered over ~1h. Two days after the footshock session, mice were tested in the active avoidance (AA) paradigm. Mice underwent 30 trials of randomized footshocks (0.35 mA). Each trial lasted a maximum of 1min and included a minimum of 5sec before each footshock. Then, the door separating 2-halves of the chamber was opened (2sec before footshock initiation) and remained open for the duration of the shock. The first 5 trials required one chamber crossing (FR1) to terminate the footshock, whereas the next 25 trials required two crossings (FR2). The AA session lasted ~30min, during which the number of escape failures was recorded.

## Statistical analysis

All data was analyzed using the Statistical Product and Service Solutions software (SPSS; IBM, North Castle, NY). Based on previously validated methods [84, 85], parameters were z-scored and averaged across tests reflecting anxiety- (EPM, OFT, NSF, NIH) and anhedonia-like behaviour (NIH, SCT), as well as overall emotionality (all previous + fur coat state in the last week). Normalized z-scores integrate behavioural tests measuring similar phenotypes to increase the consistency and reliability of these characteristically inconsistent tests [81, 84, 85]. Behavioural data was analyzed using a two-way (stress x drug) analysis of variance (ANOVA), followed by Bonferroni-corrected *post-hoc* comparisons. Weekly coat state measures were

analyzed using repeated-measures ANOVA. AA escape failures were analyzed using *student's t-test* comparing data between vehicle and riluzole groups. Data is presented as mean ± standard error of the mean (SEM).

**Results**

**Riluzole prevents the emergence of UCMS-induced anhedonia-like behaviours**

We first sought to determine whether chronic prophylactic riluzole treatment (i.e., administered concurrent with UCMS) could block the emergence of behavioural deficits induced by UCMS. Mice were subjected to 3 weeks of daily handling (control) or UCMS ± vehicle or riluzole and tested for 2 weeks, including for LA and in the OFT, EPM, NSF, NIH, and SCT (**Fig 1*a***). Fur coat deterioration was tracked weekly (**Fig. 1*a***). In the LA assessment, distance travelled in a home cage-like environment was not significantly affected by stress ($F_{1,36}$=0.025; $p$=0.87), drug treatment ($F_{1,36}$=0.183; $p$=0.67), or stress*drug interaction ($F_{1,36}$=3.02; $p$=0.09), confirming that subsequent tests were not biased by alterations in ambulatory movement (*not shown*).

In the EPM, a two-way ANOVA of open arm entries revealed no main effect of stress ($F_{1,36}$=0.009; $p$=0.92), drug ($F_{1,36}$=0.084; $p$=0.77), or a stress*drug interaction ($F_{1,36}$=2.68; $p$=0.11; shown in **Fig.1.*b***). For time spent in the open arms, a two-way ANOVA revealed a significant stress*drug interaction ($F_{1,36}$=4.49; $p$=0.041; shown in **Fig. 1.*b***); however, group-wise differences did not survive *post-hoc* correction.

In the OFT, no significant differences were found for the effects of stress, drug treatment, or their interaction on the number of entries or time spent in the center zone (shown in **Fig. 1.*c***).

In the NSF, a two-way ANOVA revealed a trend towards a main effect of stress on latency to feed in the novel environment ($F_{1,36}$=3.71; $p$=0.061; shown in **Fig. 1.*d***). However, this

may have been confounded by an UCMS-induced increase in appetitive drive, since feeding latency in the home cage control test was significantly reduced for both stress groups ($F_{1,36}$=12.68; $p$=0.0011) (CTR+vehicle: 71.7±10.9; CTR+riluzole: 77.1±8.3; UCMS+vehicle: 49.3±6; UCMS+riluzole: 42.4±5.5). The UCMS paradigm includes brief food deprivation experiences which may bias animals towards increased motivation for feeding after longer deprivation periods such as with the NSF. Indeed, in the NIH, animals were not food-deprived and underwent a similar assessment for reward approach. In the NIH, two-way ANOVA detected a significant main effect of stress ($F_{1,36}$=19.29; $p$<0.001), drug treatment ($F_{1,36}$=19.13; $p$<0.001), and a stress*drug interaction ($F_{1,36}$=10.15; $p$=0.003) for latency to drink the milk in a novel environment (shown in **Fig. 1.*e***). *Post-hoc* analysis revealed that this was driven by 3-fold increase in latency to drink induced by UCMS ($p$<0.0001) that was prevented by riluzole ($p$<0.0001). Home cage latency was not affected by stress, drug treatment, or their interaction, suggesting that novel environment differences in approach latency were not due to appetite (CTR+vehicle: 23.2±4.8; CTR+riluzole: 29.2±5.2; UCMS+vehicle: 27.4±11.2; UCMS+riluzole: 46.1±9.4).

In the SCT, two-way ANOVA revealed a significant main effect of stress ($F_{1,36}$=39.61; $p$<0.0001), drug treatment ($F_{1,36}$=7.880; $p$=0.008), and stress*drug interaction ($F_{1,36}$=14.76; $p$=0.0005) for 1hr sucrose consumption following overnight deprivation (shown in **Fig. 1.*f***). Further *post-hoc* analysis revealed a significant decrease in sucrose consumption ($p$<0.0001) that was prevented by riluzole ($p$=0.0002). After a 24hr recovery, mice were again deprived and measured for water consumption under identical conditions. Although we detected a main effect of stress on water consumption ($F_{1,36}$=5.89; $p$=0.02), reflecting reduced water intake among UCMS-exposed animals, group-wise differences did not survive *post-hoc* analyses, confirming

that sucrose consumption differences were reliably due to stress and/or drug treatment effects (CTR+vehicle: 1.6±0.2; CTR+riluzole: 1.5±0.1; UCMS+vehicle: 1.1±0.1; UCMS+riluzole: 0.9±0.2).

Repeated-measures ANOVA of weekly fur coat state assessments from week 0 to 5 revealed significant main effect of stress ($F_{1,36}$= 125.46; p<0.0001), time ($F_{5,180}$= 111.3; p<0.0001), stress*time interaction ($F_{5,180}$= 32.03; p<0.0001) and stress*drug*time interaction ($F_{5,180}$= 5.023; p=0.002), but no main effect of drug and no drug*stress or drug*time interactions. *Post-hoc* analysis showed that UCMS induced persistent fur coat deterioration starting on week 3 which was also significant for each following week (p<0.0001; Fig 1g) and not modified by riluzole treatment.

**Subsequent z-score normalization confirms that riluzole prevents the emergence of anhedonia-like behaviour, and composite emotionality-like behaviour in UCMS-exposed mice**

One issue with rodent chronic stress paradigms and classical testing methods, noted by us and others, is insufficient reliability to consistently measure depressive-like features across tests, cohorts, and between labs [82, 86, 87]. To address this, we employed multiple tests assessing common dimensional phenotypes (anxiety- and anhedonia-like behaviour, and emotionality) and analyzed both individual test parameters and normalized z-scores averaged across tests, based on validated methods [84].

Two-way ANOVA of normalized Z-anxiety scores (reflecting EPM, OFT, NSF, NIH) revealed a marginally significant stress*drug interaction ($F_{1,36}$=3.996; *p*=0.0532; shown in **Fig. 2.a**) with no main effect of stress or drug. However, post-hoc analysis did not reveal group

differences. Z-anhedonia (reflecting NIH, SCT) was significantly affected by stress ($F_{1,36}=36.74$; $p<0.001$), drug treatment ($F_{1,36}=5.899$; $p=0.0203$), and their interaction ($F_{1,36}=9.032$; $p=0048$), wherein UCMS induced an elevation in anhedonia-like behaviour ($p<0.0001$) that was prevented by riluzole ($p=0.0029$; shown in **Fig. 2.b**). Summary of behavioural emotionality via Z-emotionality scores analysis (Z-anxiety and Z-anhedonia) revealed significant effects of stress ($F_{1,36}=11.93$; $p=0.0014$), drug ($F_{1,36}=2.68$; $p=0.0109$) and a stress*drug interaction ($F_{1,36}=11.46$; $p=0.0017$; shown in **Fig. 2.c**). Overall, behavioural emotionality was significantly elevated by UCMS ($p=0.0001$) and prevented by riluzole treatment ($p=0.0065$).

**Riluzole prevents the development of learned helplessness**

Next, we investigated whether preventative riluzole treatment could block the development of helplessness-like behaviour induced by inescapable footshock. A separate cohort of mice ($n=10$/group) were administered riluzole or vehicle for 2 weeks and submitted to unpredictable inescapable footshocks as part of the LH protocol (shown in **Fig. 3.a**). In the AA paradigm, *t-test* analysis revealed a significant effect of drug treatment on the number of escape failures ($t=4.55$, df=18; $p=0.0001$), wherein compared to vehicle, riluzole decreased the number of escape failures (shown in **Fig. 3.b**).

**Discussion**

This study provides evidence that chronic prophylactic riluzole treatment can increase stress resilience and prevent the development of specific behavioural deficits induced by UCMS and LH. UCMS and LH are relevant rodent models of stress which induce behavioural deficits, such as anxiety, anhedonia and helplessness, commonly associated with stress-related illnesses including depression and PTSD [88–90]. Animals exposed to UCMS exhibited elevated anhedonia-like behaviours in the novelty-induced hypophagia and sucrose consumption tests,

which were prevented by chronic concomitant riluzole treatment. In this study, UCMS induced only slight non-significant anxiety-like behaviours making it difficult to assess the effects of riluzole treatment in these tests. Overall, z-score normalization confirms that riluzole prevents UCMS-induced elevation of behavioural emotionality, without affecting the physical state of the animals as we found no blockade of the UCMS effects on coat state in riluzole treated-animals. In a separate cohort, we also demonstrated that 2 week treatment with riluzole prior to inescapable footshocks blocked the development of helplessness-like behaviour in the AA test. These findings suggest that treatment with the Glu-modulating drug riluzole may enhance stress resilience and prevent the development of anhedonia and helplessness.

Chronic stress models such as UCMS or chronic restraint stress are classically used to model anxiety and anhedonia-like behaviour in rodents. However some tests such as open field or elevated plus maze have been proven to be inconsistent for detecting the anxiogenic effects of UCMS [82, 91]. Similar variations have been reported in our previous lab papers [81–83, 92] and outside labs as well [93, 94]. However, top-down dimension reduction via Z-scoring consistently increases the sensitivity and reliability of behavioural tests [84]. Although this approach may improve the quality of conclusions about preclinical behaviour changes, it is also important to mention that repeated testing is itself a source of variability [95]. In this study, EPM, OFT, or NSF tests failed to individually detect significant elevation of anxiety-like behaviour while marginal significance was observed across tests using the z-anxiety score. In contrast, UCMS induced robust anhedonia-like behaviours in both the NIH and SC tests, confirming previous findings from our lab [55, 96, 97] and others [98]. This effect was prevented by treatment with riluzole in individual tests and across multiple tests (z-anhedonia score). UCMS also induced coat state deterioration that was unaffected by riluzole. This is consistent

with the evidence that fur coat quality tracks the induction of stress-related deficits, but does not respond to most antidepressant treatments [82, 83]. Finally, in consideration of the potential sedative effects of high doses of riluzole [69], we confirmed that behavioural outcomes were not influenced by locomotor changes, in agreement with previous studies using similar dosing regimens [56, 99]. Altogether, our results demonstrate that chronic riluzole treatment prevents the development of UCMS-induced anxiety and anhedonia deficits.

The LH paradigm models helplessness-like behaviour associated with exposure to inescapable and uncontrollable stressors [100], and are used to assess the predictive validity of antidepressant treatment [101]. The acute/intense LH stressor may be more closely related to PTSD given that it reproduces aspects of avoidance, anxiety, and hyperarousal [102]. In rodents, escape failures in the active avoidance paradigm reflect helplessness-like behaviour and was shown to partially correlate with other dimensions including anhedonia-like behaviour [103]. In a previous study, we demonstrated that chronic riluzole administration reversed anhedonia- and helplessness-like behaviour in UCMS-exposed rats [56]. The antidepressant action of riluzole was also shown in other models, such as following olfactory bulbectomy [104] or chronic corticosterone treatment [99]. Here, we identified for the first time that chronic pre-treatment with riluzole significantly reduced the number of escape failures in the active avoidance test. This is in accordance with another study demonstrating that pre-treatment with anti-glutamatergic agents with antidepressant properties such as ketamine, an antagonist of the NMDA receptor prevented behavioural deficits induced by social defeat and LH [99]. Our results suggest that prophylactic administration of riluzole may prevent the development of helplessness-like behaviour.

Although outside the scope of this study, we can speculate on the potential cellular mechanisms underlying the prophylactic effects of chronic riluzole administration in preventing the effects of stress. Riluzole's neuroprotective and antidepressant actions have been associated with several neuronal and non-neuronal cellular mechanisms including modulation of glutamate release and enhancement of glutamate uptake and neurotrophic factors [56, 69, 99]. In previous work, we demonstrated that riluzole exerts its antidepressant-like efficacy through increased glutamate transporter-1 (GLT-1) expression as well as increased glutamate cycling in neurons and glial cells in chronically stressed and non-stressed rats [56, 105]. Anti-glutamatergic effects of riluzole were also shown to involve inhibition of the release of glutamate through a blockade of voltage-activated sodium channels [106, 107]. Finally, neuroprotective effects of riluzole have also been associated with increased BDNF expression *in vivo* [68, 99, 108, 109] and *in vitro* by neurons and astrocytes [68, 110]. Interestingly, prophylactic effects of riluzole were also demonstrated in aging studies, where pretreatment was reported to prevent cognitive decline in aged rodents through increased GLT-1 expression and maintenance neuronal morphology/complexity [111, 112]. More recently, prodromal riluzole treatment was described to induce long-term pro-cognitive effects in a genetic mouse model of Alzheimer's disease. This effect was associated with attenuated glutamatergic neurotransmission which was observed even six months after riluzole treatment discontinuation [113]. Collectively, these findings implicate Glu homeostasis, modulation, and subsequent neuroprotective or neurotrophic effects in the prophylactic and therapeutic actions of riluzole.

Initial open-label clinical studies have observed significant symptom improvement following oral riluzole administration in patients with treatment resistant depression [72, 74]. Follow-up clinical studies have confirmed or found limited efficacy of riluzole monotherapy/

adjunctive therapy to monoaminergic antidepressant or ketamine [69, 78, 114–116]. While some of these trials presented promising evidence for riluzole antidepressant action, double-blind placebo studies have demonstrated that adjunctive riluzole in MDD and riluzole monotherapy in bipolar depression, failed to induce clinically relevant improvement of depressive symptoms [115, 116]. It is possible that the reported mixed efficacy of riluzole in clinical studies is due to variance in treatment protocol or patient characteristics. For example, extension of an 8-week open-label riluzole trial to 12 weeks achieved response in one-fourth of non-responders [117]. Riluzole may also be more useful in patients with deficits in specific biomarkers. Indeed, it was demonstrated that patients with lower BDNF plasma concentrations at baseline (pre-treatment) show greater treatment response to riluzole [118]. A potential clinical implication of our findings (i.e. enhanced stress resilience from prophylactic treatment) suggests that riluzole treatment may instead be well-suited for currently remitted MDD or BPD patients who are at high risk for relapse. Several populations may also benefit from potential prophylactic riluzole treatment. These include remitted patients who still display residual symptoms or at high risk of relapse after monoaminergic antidepressant treatment discontinuation [35, 36][37, 38]

Consistent with existing evidence from preclinical and some clinical studies, we demonstrated that riluzole could prevent or block the emergence of specific behaviours related to symptom dimensions of stress-related illnesses, including anxiety-, anhedonia-, and helplessness-like behaviour, as well as overall behavioural emotionality. A major limitation of the present work is that it is a male only mouse study. Given that MDD is more prevalent among females [119], and postmortem studies identify more severe cellular deficits in females [120–122], future studies should investigate potential sex-differences in response to riluzole treatment. Although we did not investigate cellular mediators involved in riluzole prophylactic action, this study

provides a promising starting point for the use of riluzole for enhancing stress resilience through prophylactic or preventative treatment regimens. Our work raises the possibility of new therapeutic uses for riluzole including as a preventative or prophylactic treatment in individuals experiencing not only chronic stress, but also those subjected to severe acute stressors with high likelihood to develop depressive symptoms.

## Acknowledgements

Biohaven Pharmaceuticals provided funding for the experiments. C.F. was supported by a CAMH Discovery Fund fellowship and Ontario Graduate Scholarship during the studies. Y.B. was supported by Labatt Family Network for Research on the Biology of Depression. M.B. is supported by the CAMH Discovery Seed Fund and the Canadian Institutes of Health Research (PJT-165852). E.S. was supported by the Brain & Behaviour Research Foundation (#25637) and Canadian Institutes of Health Research (PJT-153175). The project was also supported by the Campbell Family Mental Health Research Institute.

## Conflicts of Interest Disclosure

G.S. has served as consultant to Allergan, Alkermes, Aptiny, , Axsome Therapeutics, Biogen, Biohaven Pharmaceuticals, Boehringer Ingelheim International GmbH, Bristol-Myers Squibb, Cowen, EMA Wellness, Engrail Therapeutics, Clexio, Denovo Biopharma, Gilgamesh, Hoffman La-Roche, Intra-Cellular Therapies, Janssen, Levo, Lundbeck, Merck, Navitor Pharmaceuticals, Neurocrine, Novartis, Noven Pharmaceuticals, Otsuka, Perception Neuroscience, Praxis Therapeutics, Sage Pharmaceuticals, Servier Pharmaceuticals, Seelos Pharmaceuticals, Taisho Pharmaceuticals, Teva, Valeant, Vistagen Therapeutics, and XW Labs; and received research

contracts from AstraZeneca, Bristol-Myers Squibb, Eli Lilly, Johnson & Johnson, Merck, Naurex, and the Usona Institute over the past 36 months. Dr. Sanacora holds equity in BioHaven Pharmaceuticals and is a co-inventor on a US patent (#8,778,979) held by Yale University and a co-inventor on US Provisional Patent Application No. 047162-7177P1 (00754) filed on August 20, 2018, by Yale University Office of Cooperative Research. Yale University has a financial relationship with Janssen Pharmaceuticals and may in the future receive financial benefits from this relationship. The University has put multiple measures in place to mitigate this institutional conflict of interest. Questions about the details of these measures should be directed to Yale University's Conflict of Interest office. RMB and VC are employees and stockholders of Biohaven Pharmaceuticals.

**Author Contributions**

**Data Availability Statement**

The data that support the findings of this study are not publicly available but are available from the corresponding author [MB] upon request.

**References**

1.  Yang L, Zhao Y, Wang Y, et al. The Effects of Psychological Stress on Depression. Curr Neuropharmacol. 2015; 13: 494–504.
2.  Kim EY, Miklowitz DJ, Biuckians A, et al. Life stress and the course of early-onset bipolar disorder. J Affect Disord. 2007; 99: 37–44.
3.  McEwen BS, Akil H. Revisiting the Stress Concept: Implications for Affective Disorders. J Neurosci. 2020; 40: 12–21.
4.  APA. *AMERICAN PSYCHIATRIC ASSOCIATION. Manual Diagnóstico e Estatístico de Transtornos Mentais - DSM-5. Porto Alegre: Artmed, 5ª edição, 2014.* 2014.
5.  Vigo D, Thornicroft G, Atun R. Estimating the true global burden of mental illness. The lancet Psychiatry. 2016; 3: 171–8.
6.  Ettman CK, Abdalla SM, Cohen GH, et al. Prevalence of Depression Symptoms in US Adults Before and During the COVID-19 Pandemic. JAMA Netw open. 2020; 3: e2019686.
7.  Xiong J, Lipsitz O, Nasri F, et al. Impact of COVID-19 pandemic on mental health in the general population: A systematic review. J Affect Disord. 2020; 277: 55–64.
8.  Clevenger SS, Malhotra D, Dang J, et al. The role of selective serotonin reuptake inhibitors in preventing relapse of major depressive disorder. Ther Adv Psychopharmacol. 2018; 8: 49–58.
9.  Garnock-Jones KP, McCormack PL. Escitalopram: a review of its use in the management of major depressive disorder in adults. CNS Drugs. 2010; 24: 769–96.
10. Huang Z-D, Zhao Y-F, Li S, et al. Comparative Efficacy and Acceptability of Pharmaceutical Management for Adults With Post-Traumatic Stress Disorder: A Systematic Review and Meta-Analysis. Front Pharmacol.; 11. Epub ahead of print 8 May 2020. DOI: 10.3389/fphar.2020.00559.
11. Yulug B. Neuroprotective treatment strategies for poststroke mood disorders: A minireview on atypical neuroleptic drugs and selective serotonin re-uptake inhibitors. Brain Res Bull. 2009; 80: 95–99.
12. Gaynes BN, Warden D, Trivedi MH, et al. What did STAR*D teach us? Results from a large-scale, practical, clinical trial for patients with depression. Psychiatr Serv. 2009; 60: 1439–45.
13. Holmes SE, Girgenti MJ, Davis MT, et al. Altered metabotropic glutamate receptor 5 markers in PTSD: In vivo and postmortem evidence. Proc Natl Acad Sci U S A. 2017; 114: 8390–8395.
14. Stein DJ, Ipser JC, Seedat S, et al. Pharmacotherapy for post traumatic stress disorder (PTSD). Cochrane Database Syst Rev. 2006; CD002795.
15. Henry M, Fishman JR, Youngner SJ. Propranolol and the prevention of post-traumatic stress disorder: is it wrong to erase the 'sting' of bad memories? Am J Bioeth. 2007; 7: 12–20.
16. Banzi R, Cusi C, Randazzo C, et al. Selective serotonin reuptake inhibitors (SSRIs) and serotonin-norepinephrine reuptake inhibitors (SNRIs) for the prevention of tension-type headache in adults. Cochrane database Syst Rev. 2015; CD011681.
17. Tran K, McGill SC, Horton J. Bupropion for Treatment-Resistant Depression. Can J Heal Technol.; 1. Epub ahead of print 8 April 2021. DOI: 10.51731/cjht.2021.55.
18. Rush AJ, Trivedi MH, Wisniewski SR, et al. Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report. Am J

Psychiatry. 2006; 163: 1905–17.

19. Sokratous S, Merkouris A, Middleton N, et al. The association between stressful life events and depressive symptoms among Cypriot university students: a cross-sectional descriptive correlational study. BMC Public Health. 2013; 13: 1121.

20. Kessler RC, Aguilar-Gaxiola S, Alonso J, et al. The associations of earlier trauma exposures and history of mental disorders with PTSD after subsequent traumas. Mol Psychiatry. 2018; 23: 1892–1899.

21. Kessler RC. The effects of stressful life events on depression. Annu Rev Psychol. 1997; 48: 191–214.

22. Sillivan SE, Joseph NF, Jamieson S, et al. Susceptibility and Resilience to Posttraumatic Stress Disorder-like Behaviors in Inbred Mice. Biol Psychiatry. 2017; 82: 924–933.

23. Peña CJ, Nestler EJ, Bagot RC. Environmental Programming of Susceptibility and Resilience to Stress in Adulthood in Male Mice. Front Behav Neurosci. 2019; 13: 40.

24. Colucci P, Marchetta E, Mancini GF, et al. Predicting susceptibility and resilience in an animal model of post-traumatic stress disorder (PTSD). Transl Psychiatry. 2020; 10: 243.

25. Torrisi SA, Lavanco G, Maurel OM, et al. A novel arousal-based individual screening reveals susceptibility and resilience to PTSD-like phenotypes in mice. Neurobiol Stress. 2021; 14: 100286.

26. Howlett JR, Stein MB. Prevention of Trauma and Stressor-Related Disorders: A Review. Neuropsychopharmacology. 2016; 41: 357–69.

27. Horn SR, Charney DS, Feder A. Understanding resilience: New approaches for preventing and treating PTSD. Exp Neurol. 2016; 284: 119–132.

28. Bernardini F, Attademo L, Cleary SD, et al. Risk Prediction Models in Psychiatry: Toward a New Frontier for the Prevention of Mental Illnesses. J Clin Psychiatry. 2017; 78: 572–583.

29. Spangler PT, West JC, Dempsey CL, et al. Randomized Controlled Trial of Riluzole Augmentation for Posttraumatic Stress Disorder: Efficacy of a Glutamatergic Modulator for Antidepressant-Resistant Symptoms. J Clin Psychiatry. 2020; 81: 1577–1589.

30. Brachman RA, McGowan JC, Perusini JN, et al. Ketamine as a Prophylactic Against Stress-Induced Depressive-like Behavior. Biol Psychiatry. 2016; 79: 776–786.

31. Chen BK, Mendez-David I, Luna VM, et al. Prophylactic efficacy of 5-HT4R agonists against stress. Neuropsychopharmacology. 2020; 45: 542–552.

32. Amat J, Dolzani SD, Tilden S, et al. Previous Ketamine Produces an Enduring Blockade of Neurochemical and Behavioral Effects of Uncontrollable Stress. J Neurosci. 2016; 36: 153–61.

33. Soumier A, Carter RM, Schoenfeld TJ, et al. New Hippocampal Neurons Mature Rapidly in Response to Ketamine But Are Not Required for Its Acute Antidepressant Effects on Neophagia in Rats. eNeuro. 2016; 3: ENEURO.0116-15.2016.

34. Mastrodonato A, Cohensedgh O, LaGamma CT, et al. Prophylactic (R,S)-ketamine selectively protects against inflammatory stressors. Behav Brain Res. 2020; 378: 112238.

35. Iovieno N, van Nieuwenhuizen A, Clain A, et al. Residual symptoms after remission of major depressive disorder with fluoxetine and risk of relapse. Depress Anxiety. 2011; 28: 137–144.

36. Nierenberg AA, Husain MM, Trivedi MH, et al. Residual symptoms after remission of major depressive disorder with citalopram and risk of relapse: a STAR*D report. Psychol Med. 2010; 40: 41–50.

37. Mueller TI, Leon AC, Keller MB, et al. Recurrence after recovery from major depressive disorder during 15 years of observational follow-up. Am J Psychiatry. 1999; 156: 1000–1006.

38. Judd LL. The Clinical Course of Unipolar Major Depressive Disorders. Arch Gen Psychiatry. 1997; 54: 989.

39. Roque AP. Pharmacotherapy as Prophylactic Treatment of Post-Traumatic Stress Disorder: A Review of the Literature. Issues Ment Health Nurs. 2015; 36: 740–751.

40. Sanacora G, Treccani G, Popoli M. Towards a glutamate hypothesis of depression: an emerging frontier of neuropsychopharmacology for mood disorders. Neuropharmacology. 2012; 62: 63–77.

41. Li C-T, Yang K-C, Lin W-C. Glutamatergic Dysfunction and Glutamatergic Compounds for Major Psychiatric Disorders: Evidence From Clinical Neuroimaging Studies. Front Psychiatry.; 9. Epub ahead of print 24 January 2019. DOI: 10.3389/fpsyt.2018.00767.

42. Henter ID, de Sousa RT, Zarate CA. Glutamatergic Modulators in Depression. Harv Rev Psychiatry. 2018; 26: 307–319.

43. Jiang X, Lin W, Cheng Y, et al. mGluR5 Facilitates Long-Term Synaptic Depression in a Stress-Induced Depressive Mouse Model. Can J Psychiatry. 2020; 65: 347–355.

44. Feyissa AM, Chandran A, Stockmeier CA, et al. Reduced levels of NR2A and NR2B subunits of NMDA receptor and PSD-95 in the prefrontal cortex in major depression. Prog Neuropsychopharmacol Biol Psychiatry. 2009; 33: 70–5.

45. Duman RS, Sanacora G, Krystal JH. Altered Connectivity in Depression: GABA and Glutamate Neurotransmitter Deficits and Reversal by Novel Treatments. Neuron. 2019; 102: 75–90.

46. Holmes SE, Girgenti MJ, Davis MT, et al. Altered metabotropic glutamate receptor 5 markers in PTSD: In vivo and postmortem evidence. Proc Natl Acad Sci U S A. 2017; 114: 8390–8395.

47. Sanacora G, Zarate CA, Krystal JH, et al. Targeting the glutamatergic system to develop novel, improved therapeutics for mood disorders. Nat Rev Drug Discov. 2008; 7: 426–437.

48. Popoli M, Yan Z, McEwen BS, et al. The stressed synapse: the impact of stress and glucocorticoids on glutamate transmission. Nat Rev Neurosci. 2011; 13: 22–37.

49. Musazzi L, Milanese M, Farisello P, et al. Acute Stress Increases Depolarization-Evoked Glutamate Release in the Rat Prefrontal/Frontal Cortex: The Dampening Action of Antidepressants. PLoS One. 2010; 5: e8566.

50. Bagley J, Moghaddam B. Temporal dynamics of glutamate efflux in the prefrontal cortex and in the hippocampus following repeated stress: effects of pretreatment with saline or diazepam. Neuroscience. 1997; 77: 65–73.

51. Hardingham GE, Bading H. Synaptic versus extrasynaptic NMDA receptor signalling: implications for neurodegenerative disorders. Nat Rev Neurosci. 2010; 11: 682–96.

52. Pál B. Involvement of extrasynaptic glutamate in physiological and pathophysiological changes of neuronal excitability. Cell Mol Life Sci. 2018; 75: 2917–2949.

53. Lener MS, Niciu MJ, Ballard ED, et al. Glutamate and Gamma-Aminobutyric Acid Systems in the Pathophysiology of Major Depression and Antidepressant Response to Ketamine. Biol Psychiatry. 2017; 81: 886–897.

54. Musazzi L, Treccani G, Popoli M. Functional and Structural Remodeling of Glutamate Synapses in Prefrontal and Frontal Cortex Induced by Behavioral Stress. Front Psychiatry.

2015; 6: 60.

55. Codeluppi SA, Chatterjee D, Prevot TD, et al. Chronic Stress Alters Astrocyte Morphology in Mouse Prefrontal Cortex. Int J Neuropsychopharmacol. 2021; pyab052.

56. Banasr M, Chowdhury GMI, Terwilliger R, et al. Glial pathology in an animal model of depression: reversal of stress-induced cellular, metabolic and behavioral deficits by the glutamate-modulating drug riluzole. Mol Psychiatry. 2010; 15: 501–11.

57. Sanacora G, Banasr M. From Pathophysiology to Novel Antidepressant Drugs: Glial Contributions to the Pathology and Treatment of Mood Disorders. Biol Psychiatry. 2013; 73: 1172–1179.

58. Mayegowda SB, Thomas C. Glial pathology in neuropsychiatric disorders: a brief review. J Basic Clin Physiol Pharmacol.; 30. Epub ahead of print 26 July 2019. DOI: 10.1515/jbcpp-2018-0120.

59. Abbink MR, van Deijk A-LF, Heine VM, et al. The involvement of astrocytes in early-life adversity induced programming of the brain. Glia. 2019; 67: 1637–1653.

60. Rajkowska G, Stockmeier CA. Astrocyte pathology in major depressive disorder: insights from human postmortem brain tissue. Curr Drug Targets. 2013; 14: 1225–36.

61. Chen J-X, Yao L-H, Xu B-B, et al. Glutamate transporter 1-mediated antidepressant-like effect in a rat model of chronic unpredictable stress. J Huazhong Univ Sci Technolog Med Sci. 2014; 34: 838–844.

62. Baek JH, Vignesh A, Son H, et al. Glutamine Supplementation Ameliorates Chronic Stress-induced Reductions in Glutamate and Glutamine Transporters in the Mouse Prefrontal Cortex. Exp Neurobiol. 2019; 28: 270–278.

63. Fullana N, Gasull-Camós J, Tarrés-Gatius M, et al. Astrocyte control of glutamatergic activity: Downstream effects on serotonergic function and emotional behavior. Neuropharmacology. 2020; 166: 107914.

64. Grant P, Song JY, Swedo SE. Review of the Use of the Glutamate Antagonist Riluzole in Psychiatric Disorders and a Description of Recent Use in Childhood Obsessive-Compulsive Disorder. J Child Adolesc Psychopharmacol. 2010; 20: 309–315.

65. Peterson AR, Binder DK. Post-translational Regulation of GLT-1 in Neurological Diseases and Its Potential as an Effective Therapeutic Target. Front Mol Neurosci. 2019; 12: 164.

66. Wilkie CM, Barron JC, Brymer KJ, et al. The Effect of GLT-1 Upregulation on Extracellular Glutamate Dynamics. Front Cell Neurosci. 2021; 15: 661412.

67. Fontana ACK. Current approaches to enhance glutamate transporter function and expression. J Neurochem. 2015; 134: 982–1007.

68. Yoshizumi M, Eisenach JC, Hayashida K. Riluzole and gabapentinoids activate glutamate transporters to facilitate glutamate-induced glutamate release from cultured astrocytes. Eur J Pharmacol. 2012; 677: 87–92.

69. Pittenger C, Coric V, Banasr M, et al. Riluzole in the treatment of mood and anxiety disorders. CNS Drugs. 2008; 22: 761–786.

70. Gourley SL, Kedves AT, Olausson P, et al. A history of corticosterone exposure regulates fear extinction and cortical NR2B, GluR2/3, and BDNF. Neuropsychopharmacology. 2009; 34: 707–716.

71. Deutschenbaur L, Beck J, Kiyhankhadiv A, et al. Role of calcium, glutamate and NMDA in major depression and therapeutic application. Prog Neuro-Psychopharmacology Biol Psychiatry. 2016; 64: 325–333.

72.     Zarate CA, Payne JL, Quiroz J, et al. An Open-Label Trial of Riluzole in Patients With Treatment-Resistant Major Depression. Am J Psychiatry. 2004; 161: 171–174.
73.     Zarate CA, Quiroz JA, Singh JB, et al. An open-label trial of the glutamate-modulating agent riluzole in combination with lithium for the treatment of bipolar depression. Biol Psychiatry. 2005; 57: 430–432.
74.     Sanacora G, Kendell SF, Levin Y, et al. Preliminary Evidence of Riluzole Efficacy in Antidepressant-Treated Patients with Residual Depressive Symptoms. Biol Psychiatry. 2007; 61: 822–825.
75.     Brennan BP, Hudson JI, Jensen JE, et al. Rapid Enhancement of Glutamatergic Neurotransmission in Bipolar Depression Following Treatment with Riluzole. Neuropsychopharmacology. 2010; 35: 834–846.
76.     Salardini E, Zeinoddini A, Mohammadinejad P, et al. Riluzole combination therapy for moderate-to-severe major depressive disorder: A randomized, double-blind, placebo-controlled trial. J Psychiatr Res. 2016; 75: 24–30.
77.     Spangler PT, West JC, Dempsey CL, et al. Randomized Controlled Trial of Riluzole Augmentation for Posttraumatic Stress Disorder. J Clin Psychiatry. 2020; 81: 20m13233.
78.     Mathew SJ, Gueorguieva R, Brandt C, et al. A Randomized, Double-Blind, Placebo-Controlled, Sequential Parallel Comparison Design Trial of Adjunctive Riluzole for Treatment-Resistant Major Depressive Disorder. Neuropsychopharmacology. 2017; 42: 2567–2574.
79.     Shimizu EN, Seifert JL, Johnson KJ, et al. Prophylactic Riluzole Attenuates Oxidative Stress Damage in Spinal Cord Distraction. J Neurotrauma. 2018; 35: 1319–1328.
80.     Rotolo RA, Demuro J, Drummond G, et al. Prophylactic exposure to oral riluzole reduces the clinical severity and immune-related biomarkers of experimental autoimmune encephalomyelitis. J Neuroimmunol. 2021; 356: 577603.
81.     Nikolova YS, Misquitta KA, Rocco BR, et al. Shifting priorities: highly conserved behavioral and brain network adaptations to chronic stress across species. Transl Psychiatry. 2018; 8: 26.
82.     Prevot TD, Misquitta KA, Fee C, et al. Residual avoidance: A new, consistent and repeatable readout of chronic stress-induced conflict anxiety reversible by antidepressant treatment. Neuropharmacology. 2019; 153: 98–110.
83.     Fee C, Prevot T, Misquitta K, et al. Chronic Stress-induced Behaviors Correlate with Exacerbated Acute Stress-induced Cingulate Cortex and Ventral Hippocampus Activation. Neuroscience. 2020; 440: 113–129.
84.     Guilloux J-P, Seney M, Edgar N, et al. Integrated behavioral z-scoring increases the sensitivity and reliability of behavioral phenotyping in mice: Relevance to emotionality and sex. J Neurosci Methods. 2011; 197: 21–31.
85.     Piantadosi SC, French BJ, Poe MM, et al. Sex-Dependent Anti-Stress Effect of an α5 Subunit Containing GABAA Receptor Positive Allosteric Modulator. Front Pharmacol. 2016; 7: 446.
86.     Andreatini R, Bacellar LFS. Animal models: Trait or state measure? The test-retest reliability of the elevated plus-maze and behavioral despair. Prog Neuro-Psychopharmacology Biol Psychiatry. 2000; 24: 549–560.
87.     Wahlsten D, Bachmanov A, Finn DA, et al. Stability of inbred mouse strain differences in behavior and brain size between laboratories and across decades. Proc Natl Acad Sci U S A. 2006; 103: 16364–9.

88.    Mineur YS, Belzung C, Crusio WE. Effects of unpredictable chronic mild stress on anxiety and depression-like behavior in mice. Behav Brain Res. 2006; 175: 43–50.

89.    Hill MN, Hellemans KGC, Verma P, et al. Neurobiology of chronic mild stress: parallels to major depression. Neurosci Biobehav Rev. 2012; 36: 2085–117.

90.    Jesberger JA, Richardson JS. Brain Output Dysregulation Induced by Olfactory Bulbectomy: An Approximation in the Rat of Major Depressive Disorder in Humans? Int J Neurosci. 1988; 38: 241–265.

91.    Troubat R, Leman S, Pinchaud K, et al. Brain immune cells characterization in UCMS exposed P2X7 knock-out mouse. Brain Behav Immun. 2021; 94: 159–174.

92.    Maluach AM, Misquitta KA, Prevot TD, et al. Increased Neuronal DNA/RNA Oxidation in the Frontal Cortex of Mice Subjected to Unpredictable Chronic Mild Stress. Chronic Stress. 2017; 1: 247054701772474.

93.    Ramos A. Animal models of anxiety: do I need multiple tests? Trends Pharmacol Sci. 2008; 29: 493–498.

94.    Willner P. Reliability of the chronic mild stress model of depression: A user survey. Neurobiol Stress. 2017; 6: 68–77.

95.    Voikar V, Vasar E, Rauvala H. Behavioral alterations induced by repeated testing in C57BL/6J and 129S2/Sv mice: implications for phenotyping screens. Genes, Brain Behav. 2004; 3: 27–38.

96.    Duric V, Banasr M, Franklin T, et al. Cariprazine Exhibits Anxiolytic and Dopamine D 3 Receptor-Dependent Antidepressant Effects in the Chronic Stress Model. Int J Neuropsychopharmacol. 2017; 20: 788–796.

97.    Misquitta KA, Miles A, Prevot TD, et al. Reduced anterior cingulate cortex volume induced by chronic stress correlates with increased behavioral emotionality and decreased synaptic puncta density. Neuropharmacology. 2021; 190: 108562.

98.    Willner P. Chronic Mild Stress (CMS) Revisited: Consistency and Behavioural-Neurobiological Concordance in the Effects of CMS. Neuropsychobiology. 2005; 52: 90–110.

99.    Gourley SL, Espitia JW, Sanacora G, et al. Antidepressant-like properties of oral riluzole and utility of incentive disengagement models of depression in mice. Psychopharmacology (Berl). 2012; 219: 805–814.

100.    Maier SF. Learned helplessness and animal models of depression. Prog Neuropsychopharmacol Biol Psychiatry. 1984; 8: 435–46.

101.    Zazpe A, Artaiz I, Labeaga L, et al. Reversal of learned helplessness by selective serotonin reuptake inhibitors in rats is not dependent on 5-HT availability. Neuropharmacology. 2007; 52: 975–984.

102.    Schöner J, Heinz A, Endres M, et al. Post-traumatic stress disorder and beyond: an overview of rodent stress models. J Cell Mol Med. 2017; 21: 2248–2256.

103.    Landgraf D, Long J, Der-Avakian A, et al. Dissociation of learned helplessness and fear conditioning in mice: a mouse model of depression. PLoS One. 2015; 10: e0125892.

104.    Takahashi K, Murasawa H, Yamaguchi K, et al. Riluzole rapidly attenuates hyperemotional responses in olfactory bulbectomized rats, an animal model of depression. Behav Brain Res. 2011; 216: 46–52.

105.    Choudary P V., Molnar M, Evans SJ, et al. Altered cortical glutamatergic and GABAergic signal transmission with glial involvement in depression. Proc Natl Acad Sci U S A. 2005; 102: 15653–8.

106. Chéramy A, Barbeito L, Godeheu G, et al. Riluzole inhibits the release of glutamate in the caudate nucleus of the cat in vivo. Neurosci Lett. 1992; 147: 209–212.
107. Hebert T, Drapeau P, Pradier L, et al. Block of the rat brain IIA sodium channel alpha subunit by the neuroprotective drug riluzole. Mol Pharmacol. 1994; 45: 1055 LP – 1060.
108. Fumagalli E, Funicello M, Rauen T, et al. Riluzole enhances the activity of glutamate transporters GLAST, GLT1 and EAAC1. Eur J Pharmacol. 2008; 578: 171–176.
109. Katoh R, Asano T, Ueda H, et al. Riluzole enhances expression of brain⬜derived neurotrophic factor with consequent proliferation of granule precursor cells in the rat hippocampus. FASEB J. 2002; 16: 1328–1330.
110. Mizuta I, Ohta M, Ohta K, et al. Riluzole stimulates nerve growth factor, brain-derived neurotrophic factor and glial cell line-derived neurotrophic factor synthesis in cultured mouse astrocytes. Neurosci Lett. 2001; 310: 117–120.
111. Pereira AC, Lambert HK, Grossman YS, et al. Glutamatergic regulation prevents hippocampal-dependent age-related cognitive decline through dendritic spine clustering. Proc Natl Acad Sci. 2014; 111: 18733–18738.
112. Pereira AC, Gray JD, Kogan JF, et al. Age and Alzheimer's disease gene expression profiles reversed by the glutamate modulator riluzole. Mol Psychiatry. 2017; 22: 296–305.
113. Hascup KN, Findley CA, Britz J, et al. Riluzole attenuates glutamatergic tone and cognitive decline in AβPP/PS1 mice. J Neurochem. 2021; 156: 513–523.
114. Yao R, Wang H, Yuan M, et al. Efficacy and safety of riluzole for depressive disorder: A systematic review and meta-analysis of randomized placebo-controlled trials. Psychiatry Res. 2020; 284: 112750.
115. Ibrahim L, Diazgranados N, Franco-Chaves J, et al. Course of improvement in depressive symptoms to a single intravenous infusion of ketamine vs add-on riluzole: results from a 4-week, double-blind, placebo-controlled study. Neuropsychopharmacology. 2012; 37: 1526–33.
116. Mathew SJ, Murrough JW, aan het Rot M, et al. Riluzole for relapse prevention following intravenous ketamine in treatment-resistant depression: a pilot randomized, placebo-controlled continuation trial. Int J Neuropsychopharmacol. 2010; 13: 71–82.
117. Sakurai H, Dording C, Yeung A, et al. Longer-term open-label study of adjunctive riluzole in treatment-resistant depression. J Affect Disord. 2019; 258: 102–108.
118. Wilkinson ST, Kiselycznyk C, Banasr M, et al. Serum and plasma brain-derived neurotrophic factor and response in a randomized controlled trial of riluzole for treatment resistant depression. J Affect Disord. 2018; 241: 514–518.
119. World Health Organization. *Depression and Other Common Mental Health Disorders: Global Health Estimates*. Geneva, 2017.
120. Seney ML, Chang L-C, Oh H, et al. The Role of Genetic Sex in Affect Regulation and Expression of GABA-Related Genes Across Species. Front Psychiatry. 2013; 4: 104.
121. Halstead JM, Lionnet T, Wilbertz JH, et al. An RNA biosensor for imaging the first round of translation from single cells to living animals. Science (80- ). 2015; 347: 1367–1671.
122. Behan ÁT, van den Hove DLA, Mueller L, et al. Evidence of female-specific glial deficits in the hippocampus in a mouse model of prenatal stress. Eur Neuropsychopharmacol. 2011; 21: 71–79.

**Fig. 1. Riluzole prevents the emergence of anhedonia-like behaviours induced by unpredictable chronic mild stress (UCMS).** (**A**) study design for behavioural analysis of Fur/Weight (FW0-5), Locomotor Activity (LA), Elevated-Plus Maze (EPM), Open Field Test (OFT), Novelty-Suppressed Feeding (NSF), Novelty-Induced Hypophagia (NIH), and Sucrose Consumption Test (SCT) beginning on day 21 of Control (Ctrl)/UCMS ± Vehicle (Veh)/Riluzole 13.2mg/kg/day p.o. ($n = 10$/group). (**B**) Number of open arm entries (left) and time spent in open arms in the EPM. (**C**) Number of center zone entries and time spent in the center zone in the OFT. (**D**) Novel environment latency to feed in the NSF. (**E**) Novel environment latency to drink a milk reward in the NIH. (**F**) Sucrose consumed in the SCT. (**G**) Weekly fur coat deterioration ratings. ****$p<0.0001$ UCMS + Veh vs. Ctrl + Veh; $^{\$\$\$\$}p<0.0001$, $^{\$\$\$}p<0.001$ UCMS + Riluzole vs. UCMS + Veh.

**Fig. 2. Z-score normalization reveals that riluzole prevents the emergence of anxiety- and anhedonia-like behaviours, and emotionality-like behaviour induced by unpredictable chronic mild stress (UCMS).** (**A**) Z-anxiety scores (reflecting elevated-plus maze, open field test, novelty-suppressed feeding, and novelty-induced hypophagia behavioural performances). (**B**) Z-anhedonia scores (reflecting novelty-induced hypophagia, sucrose consumption test performances). (**C**) Z-emotionality (reflecting all previous plus fur coat deterioration). ($n=10$/group) ****$p<0.0001$, ***$p<0.001$ UCMS + Veh vs. Ctrl+Veh; $^{\$\$\$}p<.001$, $^{\$\$}p<.01$ UCMS + Riluzole vs. UCMS + Veh.

**Fig. 3. Riluzole blocks the development of learned helplessness (LH)**. (**A**) Study design for chronic preventative riluzole treatment prior to learned helplessness conditioning including 60 inescapable footshocks with active avoidance (AA) testing 2 days later. (**B**) Number of escape failures in the active avoidance session ($n=10$/group). ***$p<.001$ Riluzole vs. Vehicle (Veh).

**Figure 1**

**Figure 2**

**Figure 3**

**A**

**B**

Active Avoidance