1 **TITLE:** An ultra-potent synthetic nanobody neutralizes SARS-CoV-2 by locking Spike into an
2 inactive conformation
3

Michael Schoof[1,2#*], Bryan Faust[1,2,3,4#], Reuben A. Saunders[1,5#], Smriti Sangwan[1,2#], Veronica
Rezelj[6#], Nick Hoppe[3,4], Morgane Boone[1,2], Christian B. Billesbølle[3,4], Cristina Puchades[4],
Caleigh M. Azumaya[4], Huong T. Kratochvil[4], Marcell Zimanyi[1,2], Ishan Deshpande[3,4], Jiahao
Liang[3], Sasha Dickinson[4], Henry C. Nguyen[4], Cynthia M. Chio[4], Gregory E. Merz[4], Michael C.
Thompson[4], Devan Diwanji[4], Kaitlin Schaefer[4], Aditya A. Anand[1,2], Niv Dobzinski[1,2], Beth
Shoshana Zha[7], Camille R. Simoneau[10,11,12], Kristoffer Leon[10,11,12], Kris M. White[13,14], Un Seng
Chio[4], Meghna Gupta[4], Mingliang Jin[4], Fei Li[4], Yanxin Liu[4], Kaihua Zhang[4], David Bulkley[4], Ming
Sun[4], Amber M. Smith[4], Alexandrea N. Rizo[4], Frank Moss[4], Axel F. Brilot[4], Sergei Pourmal[4],
Raphael Trenker[4], Thomas Pospiech[4], Sayan Gupta[9], Benjamin Barsi-Rhyne[3], Vladislav
Belyy[1,2], Andrew W. Barile-Hill[8], Silke Nock[1,2], Yuwei Liu[1,2], Nevan J. Krogan[4,5,10,11], Corie Y.
Ralston[9], Danielle L. Swaney[4,5,10,11], Adolfo García-Sastre[13,14,15,16], Melanie Ott[10,11,12], Marco
Vignuzzi[6], QCRG Structural Biology Consortium[4‡], Peter Walter[1,2*], Aashish Manglik[3,4,10,17*]
16
AFFILIATIONS:
[1]Howard Hughes Medical Institute, University of California at San Francisco, San Francisco, CA,
USA.
[2]Department of Biochemistry and Biophysics, University of California at San Francisco, San
Francisco, CA, USA.
[3]Department of Pharmaceutical Chemistry, University of California at San Francisco, San
Francisco, CA, USA.
[4]Quantitative Biosciences Institute (QBI) Coronavirus Research Group Structural Biology
Consortium, University of California, San Francisco, CA, USA.
[5]Department of Cellular and Molecular Pharmacology, University of California at San Francisco,
San Francisco, CA, USA.
[6]Viral Populations and Pathogenesis Unit, CNRS UMR 3569, Institut Pasteur, 75724, Paris,
Cedex 15, France
[7]Department of Pulmonary, Critical Care, Allergy and Sleep Medicine, University of California
San Francisco, San Francisco, CA 94158, USA.
[8]Cytiva Life Sciences, Marlborough, MA, USA.
[9]Molecular Biophysics and Integrated Bioimaging and the Molecular Foundry, Lawrence
Berkeley National Laboratory, Berkeley, CA, USA.
[10]Quantitative Biosciences Institute (QBI), University of California San Francisco, San
Francisco, CA, USA.
[11]J. David Gladstone Institutes, San Francisco, CA, USA.
[12]Department of Medicine, University of California San Francisco, San Francisco, CA, USA.
[13]Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY, USA.
[14]Global Health and Emerging Pathogens Institute, Icahn School of Medicine at Mount Sinai,
New York, NY, USA.
[15]Department of Medicine, Division of Infectious Diseases, Icahn School of Medicine at Mount
Sinai, New York, NY, USA
[16]The Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA
[17]Department of Anesthesia and Perioperative Care, University of California at San Francisco,
San Francisco, CA, USA.
47
48 # These authors contributed equally
49 ‡ Full listing in Supplementary Information
50 * To whom correspondence should be addressed; Email: michael@walterlab.ucsf.edu,
51 peter@walterlab.ucsf.edu, Aashish.Manglik@ucsf.edu

ABSTRACT

Without an effective prophylactic solution, infections from SARS-CoV-2 continue to rise worldwide with devastating health and economic costs. SARS-CoV-2 gains entry into host cells via an interaction between its Spike protein and the host cell receptor angiotensin converting enzyme 2 (ACE2). Disruption of this interaction confers potent neutralization of viral entry, providing an avenue for vaccine design and for therapeutic antibodies. Here, we develop single-domain antibodies (nanobodies) that potently disrupt the interaction between the SARS-CoV-2 Spike and ACE2. By screening a yeast surface-displayed library of synthetic nanobody sequences, we identified a panel of nanobodies that bind to multiple epitopes on Spike and block ACE2 interaction via two distinct mechanisms. Cryogenic electron microscopy (cryo-EM) revealed that one exceptionally stable nanobody, Nb6, binds Spike in a fully inactive conformation with its receptor binding domains (RBDs) locked into their inaccessible down-state, incapable of binding ACE2. Affinity maturation and structure-guided design of multivalency yielded a trivalent nanobody, mNb6-tri, with femtomolar affinity for SARS-CoV-2 Spike and picomolar neutralization of SARS-CoV-2 infection. mNb6-tri retains stability and function after aerosolization, lyophilization, and heat treatment. These properties may enable aerosol-mediated delivery of this potent neutralizer directly to the airway epithelia, promising to yield a widely deployable, patient-friendly prophylactic and/or early infection therapeutic agent to stem the worst pandemic in a century.

MAIN TEXT:

Over the last two decades, three zoonotic β-coronaviruses have entered the human population,

causing severe respiratory symptoms with high mortality (*1-3*). The ongoing COVID-19

pandemic is caused by SARS-CoV-2, the most readily transmissible of these three

coronaviruses (*4-7*). SARS-CoV-2 has wrecked the world's economy and societies to an

unprecedented extent, to date (Aug. 14, 2020) causing 751,154

reported deaths around the globe (*8*). Although public health measures have slowed its spread

in many regions, infection hotspots keep reemerging. No successful vaccine or preventive

treatment has yet been manufactured for any coronavirus, and the time to develop an effective

and broadly available vaccine for SARS-CoV-2 remains uncertain. The development of novel

therapeutic and prophylactic approaches thus remains essential, both as temporary stopgaps

until an effective vaccine is generated and as permanent solutions for those segments of the

population for which vaccination proves ineffective or contraindicated.

Coronavirus virions are bounded by a membrane envelope that contains ~25 copies of the

homotrimeric transmembrane spike glycoprotein (Spike) responsible for virus entry into the host

cell (*9*). The surface-exposed portion of Spike is composed of two domains, S_1 and S_2 (*10*). The

S_1 domain mediates the interaction between virus and its host cell receptor, the angiotensin

converting enzyme 2 (ACE2), while the S_2 domain catalyzes fusion of the viral and host cell

membranes (*3, 11-13*). During its biogenesis, the Spike protein is proteolytically cleaved

between the S_1 and S_2 domains, which primes the virus for cellular entry (*10*). Contained within

S_1 is the receptor binding domain (RBD), which directly binds to ACE2. The RBD is attached to

the body of Spike by a flexible region and can exist in an inaccessible down-state or an

accessible up-state (*14, 15*). Binding to ACE2 requires the RBD in the up-state and enables

cleavage by host proteases TMPRSS2 or cathepsin, triggering a dramatic conformational

change in S_2 that enables viral entry (*16*). In SARS-CoV-2 virions, Spike oscillates between an

active, open conformation with at least one RBD in the up-state and an inactive, closed

conformation with all RBDs in the down-state (*9, 11, 14, 15*).

By screening a high-complexity yeast surface-displayed library of synthetic nanobodies, we

have uncovered a collection of nanobodies that block the Spike-ACE2 interaction. Biochemical

and structural studies revealed that two classes of these nanobodies act in distinct ways to

prevent ACE2 binding. Combining affinity maturation and structure-guided multimerization, we

optimized these agents and generated Spike binders that match or exceed the potency of most

monoclonal antibodies disclosed to date. Our lead neutralizing molecule, mNb6-tri, blocks

SARS-CoV-2 entry in human cells at picomolar efficacy and withstands aerosolization,

lyophilization, and elevated temperatures. mNb6-tri provides a promising approach to deliver a

potent SARS-CoV-2 neutralizing molecule directly to the airways for prophylaxis or therapy.

RESULTS

Synthetic nanobodies that disrupt Spike-ACE2 interaction

To isolate nanobodies that neutralize SARS-CoV-2, we screened a yeast surface-displayed

library of $>2x10^9$ synthetic nanobody sequences. Our strategy was to screen for binders to the

full Spike protein ectodomain, in order to capture not only those nanobodies that would compete

by binding to the ACE2-binding site on the RBD directly but also those that might bind

elsewhere on Spike and block ACE2 interaction through indirect mechanisms. We used a

mutant form of SARS-CoV-2 Spike (Spike*,) as the antigen (*15*). Spike* lacks one of the two

activating proteolytic cleavage sites between the S_1 and S_2 domains and introduces two

mutations to stabilize the pre-fusion conformation. Spike* expressed in mammalian cells binds

ACE2 with a K_D = 44 nM (Supplementary Fig. 1), consistent with previous reports (*17*). Next, we

labeled Spike* with biotin or with fluorescent dyes and selected nanobody-displaying yeast over

multiple rounds, first by magnetic bead binding and then by fluorescence-activated cell sorting

(Fig. 1A).

Three rounds of selection yielded 21 unique nanobodies that bound Spike* and showed

decreased Spike* binding in the presence of ACE2. Closer inspection of their binding properties

revealed that these nanobodies fall into two distinct classes. One group (Class I) binds the RBD

and competes with ACE2 (Fig. 1B). A prototypical example of this class is nanobody Nb6, which

binds to Spike* and to RBD alone with a K_D of 210 nM and 41 nM, respectively (Fig. 1C; Table

1). Another group (Class II), exemplified by nanobody Nb3, binds to Spike* (K_D = 61 nM), but

displays no binding to RBD alone (Fig. 1C, Table 1). In the presence of excess ACE2, binding of

Nb6 and other Class I nanobodies is blocked entirely, whereas binding of Nb3 and other Class II

nanobodies is decreased only moderately (Fig. 1B). These results suggest that Class I

nanobodies target the RBD to block ACE2 binding, whereas Class II nanobodies target other

epitopes and decrease ACE2 interaction with Spike allosterically or through steric interference.

Indeed, surface plasmon resonance (SPR) experiments demonstrate that Class I and Class II

nanobodies can bind Spike* simultaneously (Fig. 1D).

Analysis of the kinetic rate constants for Class I nanobodies revealed a consistently greater

association rate constant (k_a) for nanobody binding to the isolated RBD than to full-length Spike*

(Table 1), which suggests that RBD accessibility influences the K_D. We next tested the efficacy

of our nanobodies, both Class I and Class II, to inhibit binding of fluorescently labeled Spike* to

ACE2-expressing HEK293 cells (Table 1, Fig. 1E). Class I nanobodies emerged with highly

variable activity in this assay with Nb6 and Nb11 as two of the most potent clones with IC_{50}

values of 370 and 540 nM, respectively (Table 1). For unexplained reasons, Class II nanobodies

showed little to no activity in this assay (Table 1, Fig. 1E).

Going forward, we prioritized two Class I nanobodies, Nb6 and Nb11, that combine potent

Spike* binding with relatively small differences in k_a between binding to Spike* or RBD. We

reasoned that the epitopes recognized by Nb6 and Nb11 would be more readily accessible in

the Spike protein on intact virions. For Class II nanobodies we prioritized Nb3 because of its

optimal stability and yield during purification.

Nb6 and Nb11 target the RBD and directly compete with ACE2

To define the binding sites of Nb6 and Nb11, we determined their cryogenic electron

microscopy (cryo-EM) structures bound to Spike* (Fig. 2A-B, Supplementary Fig. 2-4,

Supplementary Table 1). Both nanobodies recognize RBD epitopes that overlap the ACE2

binding site (Fig. 2E). For Nb6 and Nb11, we resolved nanobody binding to both the open and

closed conformations of Spike*. We obtained a 3.0 Å map of Nb6 bound to closed Spike*, which

enabled modeling of the Nb6-Spike* complex (Fig. 2A), including the complementarity

determining regions (CDRs). We also obtained lower resolution maps for Nb6 bound to open

Spike* (3.8 Å), Nb11 bound to open Spike* (4.2 Å), and Nb11 bound to closed Spike* (3.7 Å).

For these lower resolution maps, we could define the nanobody's binding orientation but not

accurately model the CDRs.

Nb6 bound to closed Spike* straddles the interface between two adjacent RBDs. The majority of

the contacting surfaces are contributed by CDR1 and CDR2 of Nb6 (Fig. 2C). CDR3 contacts

the adjacent RBD that is counterclockwise positioned when viewed from the top of Spike* (Fig.

2C). The binding of one Nb6 therefore stabilizes two adjacent RBDs in the down-state. We

surmise that this initial binding event pre-organizes the binding site for a second and third Nb6

molecule to stabilize the closed Spike* conformation. Indeed, binding of two Nb6 molecules

would lock all three RBDs into the down-state, thus highly favoring binding of a third Nb6

because binding would not entail any further entropic cost. By contrast, Nb11 bound to down-

state RBDs only contacts a single RBD (Fig. 2D).

Nb3 interacts with the Spike S$_1$ domain external to the RBD

Our attempts to determine the binding site of Nb3 by cryo-EM proved unsuccessful. We

therefore turned to radiolytic hydroxyl radical footprinting to determine potential binding sites for

Nb3. Spike*, either apo or bound to Nb3, was exposed to 5-50 milliseconds of synchrotron X-ray

radiation to label solvent-exposed amino acids with hydroxyl radicals. Radical-labeled amino

acids were subsequently identified and quantified by mass spectrometry of trypsin/Lys-C or Glu-

C protease digested Spike*[18]. Two neighboring surface residues on the S$_1$ domain of Spike

(M177 and H207) emerged as highly protected sites in the presence of Nb3 (Supplementary

Fig. 5). The degree of protection is consistent with prior observations of antibody-antigen

interactions by hydroxyl radical footprinting [19]. Both M177 and H207 are greater than 40 Å

distant from the ACE2 binding site on the RBD, suggesting that Nb3 may inhibit Spike-ACE2

interactions through allosteric means.

Rationally engineered multivalency increases potency

The structure of Nb6 bound to closed Spike* enabled us to engineer bivalent and trivalent

nanobodies predicted to lock all RBDs in the down-state. To this end, we inserted flexible Gly-

Ser linkers of either 15 or 20 amino acids to span the 52 Å distance between adjacent Nb6

monomers bound to down-state RBDs in closed Spike* (Supplementary Fig. 6). Both linker

lengths are too short to span the distance (72 Å) between Nb6 bound to a down-state RBD and

an up-state RBD that would co-exist in an open Spike. Moreover, binding of three RBDs in the

previously reported conformation of Nb6-bound open Spike* would be physically impossible

even with longer linker length because of steric clashes (Supplementary Fig. 6). By contrast, the

minimum distance between adjacent Nb11 monomers bound to either open or closed Spike* is

68 Å (Supplementary Fig. 6). We therefore predicted that multivalent binding by Nb6 constructs

would display significantly slowed dissociation rates due to the enhanced avidity afforded by

Spike's trimeric architecture.

We assessed multivalent Nb6 binding to Spike* by SPR. Both bivalent Nb6 with a 15 amino acid

linker (Nb6-bi) and trivalent Nb6 with two 20 amino acid linkers (Nb6-tri) dissociate from Spike*

in a biphasic manner. The dissociation phase can be fitted to two components: a fast phase with

kinetic rate constants k_{d1} of 2.7×10^{-2} s^{-1} for Nb6-bi and 2.9×10^{-2} s^{-1} for Nb6-tri, which are of the

same magnitude as that observed for monovalent Nb6 ($k_d = 5.6 \times 10^{-2}$ s^{-1}) and a slow phase that

is dependent on avidity ($k_{d2} = 3.1 \times 10^{-4}$ for Nb6-bi and $k_{d2} < 1.0 \times 10^{-6}$ s^{-1} for Nb6-tri, respectively)

(Fig. 3A). The relatively similar k_d for the fast phase suggests that a fraction of the observed

binding for the multivalent constructs is nanobody binding to a single Spike* RBD. By contrast,

the slow dissociation phase of Nb6-bi and Nb6-tri indicates engagement of two or three RBDs.

We observed no dissociation for the slow phase of Nb6-tri over 10 minutes, indicating an upper

boundary for k_{d2} of 1×10^{-6} s^{-1} and subpicomolar affinity. This measurement remains an upper-

bound estimate rather than an accurate measurement because the technique is limited by the

intrinsic dissociation rate of Spike* from the chip imposed by the chemistry used to immobilize

Spike*.

We reasoned that the biphasic dissociation behavior could be explained by a slow

interconversion between up- and down-state RBDs, with conversion to the more stable down-

state required for full trivalent binding. According to this view, a single domain of Nb6-tri

engaged with an up-state RBD would dissociate rapidly. The system would then re-equilibrate

as the RBD flips into the down-state, eventually allowing Nb6-tri to trap all RBDs in closed

Spike*. To test this notion directly, we varied the time allowed for Nb6-tri binding to Spike*.

Indeed, we observed an exponential decrease in the percent fast-phase with a $t_{1/2}$ of 65 s (Fig.

3B), which, we surmise, reflects the timescale of conversion between the RBD up- and down-

states in Spike*. Taken together, dimerization and trimerization of Nb6 afforded 750-fold and

>200,000-fold gains in K_D, respectively.

Class I and II nanobodies prevent SARS-CoV-2 infection

We next tested the neutralization activity of trivalent versions of our top Class I (Nb6 and Nb11)

and Class II (Nb3) nanobodies against SARS-CoV-2 pseudotyped lentivirus. In this assay,

SARS-CoV-2 Spike is expressed as a surface protein on a lentiviral particle that contains a

ZsGreen reporter gene, which is integrated and expressed upon successful viral entry into cells

harboring the ACE2 receptor ([20]). Nb6 and Nb11 inhibited pseudovirus infection with IC$_{50}$

values of 2.0 μM and 2.4 μM, respectively, and Nb3 inhibited pseudovirus infection with an IC$_{50}$

of 3.9 μM (Fig. 3C, Table 1). Nb6-tri shows a 2000-fold enhancement of inhibitory activity, with

an IC$_{50}$ of 1.2 nM, whereas trimerization of Nb11 and Nb3 resulted in more modest gains of 40-

and 10-fold (51 nM and 400 nM), respectively (Fig. 3C).

We next confirmed these neutralization activities with a viral plaque assay using live SARS-

CoV-2 virus infection of VeroE6 cells. Consistent with its activity against pseudotyped lentivirus,

Nb6-tri proved exceptionally potent, neutralizing SARS-CoV-2 with an average IC_{50} of 160 pM

(Fig. 3D). Nb3-tri neutralized SARS-CoV-2 with an average IC_{50} of 140 nM (Fig. 3D).

Affinity maturation yields a femtomolar K_D Spike inhibitor

We further optimized the potency of Nb6 by selecting high-affinity variants. To this end, we

prepared a new library, starting with the Nb6 coding sequence, in which we varied each amino

acid position of all three CDRs by saturation mutagenesis (Fig. 4A). After two rounds of

magnetic bead-based selection, we isolated a population of high-affinity clones. Sequencing

revealed two highly penetrant mutations: I27Y in CDR1 and P105Y in CDR3. We incorporated

these two mutations into Nb6 to generate matured Nb6 (mNb6), which binds with 500-fold

increased affinity to Spike* as measured by SPR (Fig. 4B). As a monomer, mNb6 inhibits both

pseudovirus and live SARS-CoV-2 infection with low nanomolar potency, a ~200-fold

improvement compared to Nb6 (Fig. 4I-J, Table 1).

A 2.9 Å cryo-EM structure of mNb6 bound to Spike* shows that, like the parent nanobody Nb6,

mNb6 binds to closed Spike (Fig. 4C, Supplementary Fig. 7). The higher resolution map allowed

us to build a model with high confidence and determine the effects of the I27Y and P105Y

substitutions. mNb6 induces a slight rearrangement of the down-state RBDs as compared to

both previously determined structures of apo-Spike* and Spike* bound to Nb6, inducing a 9°

rotation of the RBD away from the central three-fold symmetry axis (Fig. 4H) (*14, 15*). This

deviation likely arises from a different interaction between CDR3 and Spike*, which nudges the

RBDs into a new resting position. While the I27Y substitution optimizes local contacts between

CDR1 in its original binding site on the RBD, the P105Y substitution leads to a marked

rearrangement of CDR3 in mNb6 (Fig. 4F-G). This conformational change yields a different set

of contacts between mNb6 CDR3 and the adjacent RBD (Fig. 4D). Remarkably, an X-ray crystal

structure of mNb6 alone revealed dramatic conformational differences in CDR1 and CDR3

between free and Spike*-bound mNb6, suggestive of significant conformational heterogeneity

for the unbound nanobodies and induced-fit rearrangements upon binding to Spike* (Fig. 4E).

The binding orientation of mNb6 is similar to that of Nb6, supporting the notion that our

multivalent design would likewise enhance binding affinity. Unlike Nb6-tri, trivalent mNb6

(mNb6-tri) bound to Spike with no observable fast-phase dissociation and no measurable

dissociation over ten minutes, yielding an upper bound for the dissociation rate constant k_d of

1.0×10^{-6} s^{-1} ($t_{1/2}$ > 8 days) and a K_D of <1 pM (Fig. 4B). As above, more precise measurements

of the dissociation rate are precluded by the surface chemistry used to immobilize Spike*.

mNb6-tri displays further gains in potency in both pseudovirus and live SARS-CoV-2 infection

assays with IC_{50} values of 120 pM (5.0 ng/mL) and 54 pM (2.3 ng/mL), respectively (Fig. 4H-I,

Table 1). Given the sub-picomolar affinity observed by SPR, it is likely that these viral

neutralization potencies reflect the lower limit of the assays. mNb6-tri is therefore an

exceptionally potent SARS-CoV-2 neutralizing antibody, among the most potent molecules

disclosed to date.

Nb6, Nb6-tri, mNb6, and mNb6-tri are robust proteins

One of the most attractive properties that distinguishes nanobodies from traditional monoclonal

antibodies is their extreme stability (*21*). We therefore tested Nb6, Nb6-tri, mNb6, and mNb6-tri

for stability regarding temperature, lyophilization, and aerosolization. Temperature denaturation

experiments using circular dichroism measurements to assess protein unfolding revealed

melting temperatures of 66.9, 62.0, 67.6, and 61.4 °C for Nb6, Nb6-tri, mNb6 and mNb6-tri,

respectively (Fig 5A). Aerosolization and prolonged heating of Nb6, mNb6, and mNb6-tri for 1

hour at 50°C induced no loss of activity (Fig 5B). Moreover, mNb6 and mNb6-tri were stable to

lyophilization and to aerosolization using a mesh nebulizer, showing no aggregation by size

exclusion chromatography and preserved high affinity binding to Spike* (Fig. 5C-D).

DISCUSSION

There is a pressing need for prophylactics and therapeutics against SARS-CoV-2 infection.

Most recent strategies to prevent SARS-CoV-2 entry into the host cell aim at blocking the

ACE2-RBD interaction. High-affinity monoclonal antibodies, many identified from convalescent

patients, are leading the way as potential therapeutics (*22-29*). While highly effective *in vitro*,

these agents are expensive to produce by mammalian cell expression and need to be

intravenously administered by healthcare professionals (*30*). Moreover, large doses are likely to

be required for prophylactic viral neutralization, as only a small fraction of systemically

circulating antibodies cross the epithelial cell layers that line the airways (*31*). By contrast, single

domain antibodies (nanobodies) provide significant advantages in terms of production and

deliverability. They can be inexpensively produced at scale in bacteria *(E. coli)* or yeast *(P.*

pastoris). Furthermore, their inherent stability enables aerosolized delivery directly to the nasal

and lung epithelia by self-administered inhalation (*32*).

Monomeric mNb6 is among the most potent single domain antibodies neutralizing SARS-CoV-2

discovered to date. Multimerization of single domain antibodies has been shown to improve

target affinity by avidity (*32, 33*). In the case of Nb6 and mNb6, however, our design strategy

enabled a multimeric construct that simultaneously engages all three RBDs, yielding profound

gains in potency. Furthermore, because RBDs must be in the up-state to engage with ACE2,

conformational control of RBD accessibility can serve as an added neutralization mechanism.

Indeed, our Nb6-tri and mNb6-tri molecules were designed with this functionality in mind. Thus,

when mNb6-tri engages with Spike, it prevents ACE2 binding by both directly occluding the

binding site and by locking the RBDs into an inactive conformation. Although a multitude of

other monoclonal and single-domain antibodies against SARS-CoV-2 Spike have been

discovered to date, there are few if any molecules as potent and stable as mNb6-tri (*33-43*).

Resistance to aerosolization, in particular, offers unprecedented opportunity for patient-friendly

nasal and pulmonary administration.

Our discovery of Class II neutralizing nanobodies demonstrates the presence of previously

unexplored mechanisms of blocking Spike binding to ACE2. For one Class II nanobody, Nb3,

we identified a likely binding site in the Spike S_1 domain external to the RBDs. Previously

discovered neutralizing antibodies from convalescent patients bind an epitope in a similar region

of Spike (*24, 26, 27*). Binding of Nb3 to this epitope may allosterically stabilize RBDs in the

down-state, thereby decreasing ACE2 binding. Pairing of Class I and Class II nanobodies in a

prophylactic or therapeutic cocktail could thus be a highly advantageous strategy for both potent

neutralization and prevention of escape variants. The combined stability, potency, and diverse

epitope engagement of our anti-Spike nanobodies therefore provide a unique potential

prophylactic and therapeutic strategy to limit the continued toll of the COVID-19 pandemic.

REFERENCES AND NOTES:

1. T. G. Ksiazek *et al.*, A novel coronavirus associated with severe acute respiratory syndrome. *N Engl J Med* **348**, 1953-1966 (2003).

2. A. M. Zaki, S. van Boheemen, T. M. Bestebroer, A. D. Osterhaus, R. A. Fouchier, Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* **367**, 1814-1820 (2012).

3. P. Zhou *et al.*, A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273 (2020).

4. J. F. Chan *et al.*, A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* **395**, 514-523 (2020).

5. C. Huang *et al.*, Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, 497-506 (2020).

6. F. Wu *et al.*, A new coronavirus associated with human respiratory disease in China. *Nature* **579**, 265-269 (2020).

7. N. Zhu *et al.*, A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* **382**, 727-733 (2020).

8. World Health Organization. (2020).

9. Z. Ke *et al.*, Structures, conformations and distributions of SARS-CoV-2 spike protein trimers on intact virions. *bioRxiv*, 2020.2006.2027.174979 (2020).

10. B. J. Bosch, R. van der Zee, C. A. M. de Haan, P. J. M. Rottier, The coronavirus spike protein is a class I virus fusion protein: structural and functional characterization of the fusion core complex. *J. Virol.* **77**, 8801-8811 (2003).

11. Y. Cai *et al.*, Distinct conformational states of SARS-CoV-2 spike protein. *Science*, (2020).

12. Q. Wang *et al.*, Structural and Functional Basis of SARS-CoV-2 Entry by Using Human ACE2. *Cell* **181**, 894-904.e899 (2020).

13. R. Yan *et al.*, Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science* **367**, 1444-1448 (2020).

14. A. C. Walls *et al.*, Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* **181**, 281-292.e286 (2020).

15. D. Wrapp *et al.*, Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* **367**, 1260-1263 (2020).

16. M. Hoffmann *et al.*, SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280.e278 (2020).

17. J. Shang *et al.*, Structural basis of receptor recognition by SARS-CoV-2. *Nature* **581**, 221-224 (2020).

18. S. Gupta, J. Feng, L. J. Chan, C. J. Petzold, C. Y. Ralston, Synchrotron X-ray footprinting as a method to visualize water in proteins. *J Synchrotron Radiat* **23**, 1056-1069 (2016).

19. Y. Zhang, A. T. Wecksler, P. Molina, G. Deperalta, M. L. Gross, Mapping the Binding Interface of VEGF and a Monoclonal Antibody Fab-1 Fragment with Fast Photochemical Oxidation of Proteins (FPOP) and Mass Spectrometry. *J Am Soc Mass Spectrom* **28**, 850-858 (2017).

20. K. H. D. Crawford *et al.*, Protocol and Reagents for Pseudotyping Lentiviral Particles with SARS-CoV-2 Spike Protein for Neutralization Assays. *Viruses* **12**, (2020).

21. P. Kunz *et al.*, The structural basis of nanobody unfolding reversibility and thermoresistance. *Sci Rep* **8**, 7934 (2018).

22. A. Baum *et al.*, Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with individual antibodies. *Science*, (2020).

386   23.   Y. Cao *et al.*, Potent Neutralizing Antibodies against SARS-CoV-2 Identified by High-
387   Throughput Single-Cell Sequencing of Convalescent Patients' B Cells. *Cell* **182**, 73-
388   84.e16 (2020).
389   24.   X. Chi *et al.*, A neutralizing human antibody binds to the N-terminal domain of the Spike
390   protein of SARS-CoV-2. *Science*, (2020).
391   25.   B. Ju *et al.*, Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature*,
392   (2020).
393   26.   L. Liu *et al.*, Potent neutralizing antibodies directed to multiple epitopes on SARS-CoV-2
394   spike. *Nature*, (2020).
395   27.   D. Pinto *et al.*, Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
396   antibody. *Nature* **583**, 290-295 (2020).
397   28.   T. F. Rogers *et al.*, Isolation of potent SARS-CoV-2 neutralizing antibodies and
398   protection from disease in a small animal model. *Science*, (2020).
399   29.   S. J. Zost *et al.*, Potently neutralizing and protective human antibodies against SARS-
400   CoV-2. *Nature*, (2020).
401   30.   H. Ledford, Antibody therapies could be a bridge to a coronavirus vaccine - but will the
402   world benefit? *Nature*, (2020).
403   31.   V. H. Leyva-Grado, G. S. Tan, P. E. Leon, M. Yondola, P. Palese, Direct administration
404   in the respiratory tract improves efficacy of broadly neutralizing anti-influenza virus
405   monoclonal antibodies. *Antimicrob Agents Chemother* **59**, 4162-4172 (2015).
406   32.   L. Detalle *et al.*, Generation and Characterization of ALX-0171, a Potent Novel
407   Therapeutic Nanobody for the Treatment of Respiratory Syncytial Virus Infection.
408   *Antimicrob Agents Chemother* **60**, 6-13 (2016).
409   33.   D. Wrapp *et al.*, Structural Basis for Potent Neutralization of Betacoronaviruses by
410   Single-Domain Camelid Antibodies. *Cell* **181**, 1004-1015.e1015 (2020).
411   34.   J. Huo *et al.*, Neutralizing nanobodies bind SARS-CoV-2 spike RBD and block
412   interaction with ACE2. *Nat. Struct. Mol. Biol.*, (2020).
413   35.   J. D. Walter *et al.*, Sybodies targeting the SARS-CoV-2 receptor-binding domain.
414   *bioRxiv*, (2020).
415   36.   Z. Sun *et al.*, Potent neutralization of SARS-CoV-2 by human antibody heavy-chain
416   variable domains isolated from a large library with a new stable scaffold. *MAbs* **12**,
417   1778435 (2020).
418   37.   Y. Wu *et al.*, Identification of Human Single-Domain Antibodies against SARS-CoV-2.
419   *Cell Host Microbe* **27**, 891-898.e895 (2020).
420   38.   T. J. Esparza, D. L. Brody, High Affinity Nanobodies Block SARS-CoV-2 Spike Receptor
421   Binding Domain Interaction with Human Angiotensin Converting Enzyme. *bioRxiv*,
422   2020.2007.2024.219857 (2020).
423   39.   L. Hanke *et al.*, An alpaca nanobody neutralizes SARS-CoV-2 by blocking receptor
424   interaction. *bioRxiv*, 2020.2006.2002.130161 (2020).
425   40.   T. F. Custódio *et al.*, Selection, biophysical and structural analysis of synthetic
426   nanobodies that effectively neutralize SARS-CoV-2. *bioRxiv*, 2020.2006.2023.165415
427   (2020).
428   41.   T. Li *et al.*, Potent synthetic nanobodies against SARS-CoV-2 and molecular basis for
429   neutralization. *bioRxiv*, 2020.2006.2009.143438 (2020).
430   42.   G. V. Nieto *et al.*, Fast isolation of sub-nanomolar affinity alpaca nanobody against the
431   Spike RBD of SARS-CoV-2 by combining bacterial display and a simple single-step
432   density gradient selection. *bioRxiv*, 2020.2006.2009.137935 (2020).
433   43.   X. Chi *et al.*, Humanized Single Domain Antibodies Neutralize SARS-CoV-2 by
434   Targeting Spike Receptor Binding Domain. *bioRxiv*, 2020.2004.2014.042010 (2020).

44. D. Stadlbauer *et al.*, SARS-CoV-2 Seroconversion in Humans: A Detailed Protocol for a
Serological Assay, Antigen Production, and Test Setup. *Curr. Protoc. Microbiol.* **57**, e100
(2020).
45. I. Lui *et al.*, Trimeric SARS-CoV-2 Spike interacts with dimeric ACE2 with limited intra-
Spike avidity. *bioRxiv*, 2020.2005.2021.109157 (2020).
46. C. McMahon *et al.*, Yeast surface display platform for rapid discovery of conformationally
selective nanobodies. *Nat. Struct. Mol. Biol.* **25**, 289-296 (2018).
47. D. N. Mastronarde, Automated electron microscope tomography using robust prediction
of specimen movements. *J Struct Biol* **152**, 36-51 (2005).
48. S. Q. Zheng *et al.*, MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017).
49. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for
rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296 (2017).
50. J. Zivanov *et al.*, New tools for automated high-resolution cryo-EM structure
determination in RELION-3. *Elife* **7**, (2018).
51. T. Grant, A. Rohou, N. Grigorieff, cisTEM, user-friendly software for single-particle image
processing. *Elife* **7**, (2018).
52. J. Lan *et al.*, Structure of the SARS-CoV-2 spike receptor-binding domain bound to the
ACE2 receptor. *Nature* **581**, 215-220 (2020).
53. S. G. Rasmussen *et al.*, Structure of a nanobody-stabilized active state of the β(2)
adrenoceptor. *Nature* **469**, 175-180 (2011).
54. B. Frenz, A. C. Walls, E. H. Egelman, D. Veesler, F. DiMaio, RosettaES: a sampling
strategy enabling automated interpretation of difficult cryo-EM maps. *Nat Methods* **14**,
797-800 (2017).
55. P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta
Crystallogr D Biol Crystallogr* **60**, 2126-2132 (2004).
56. T. I. Croll, ISOLDE: a physically realistic environment for model building into low-
resolution electron-density maps. *Acta Crystallogr D Struct Biol* **74**, 519-530 (2018).
57. P. D. Adams *et al.*, PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010).
58. D. P. Staus *et al.*, Allosteric nanobodies reveal the dynamic range and diverse
mechanisms of G-protein-coupled receptor activation. *Nature* **535**, 448-452 (2016).
59. S. Gupta, M. Sullivan, J. Toomey, J. Kiselar, M. R. Chance, The Beamline X28C of the
Center for Synchrotron Biosciences: a national resource for biomolecular structure and
dynamics experiments using synchrotron footprinting. *J Synchrotron Radiat* **14**, 233-243
(2007).
60. F. Yu *et al.*, Fast quantitative analysis of timsTOF PASEF data with MSFragger and
IonQuant. *Mol Cell Proteomics*, (2020).
61. M. Choi *et al.*, MSstats: an R package for statistical analysis of quantitative mass
spectrometry-based proteomic experiments. *Bioinformatics* **30**, 2524-2526 (2014).
62. W. Kabsch, Automatic processing of rotation diffraction data from crystals of initially
unknown symmetry and cell constants. *Journal of Applied Crystallography* **26**, 795-800
(1993).
63. A. J. McCoy *et al.*, Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-674
(2007).
64. B. E. Bricogne G., Brandl M., Flensburg C., Keller P., Paciorek W.,, S. A. Roversi P,
Smart O.S., Vonrhein C., Womack T.O. , BUSTER version 1.10.0. . *Cambridge, United
Kingdom: Global Phasing Ltd.*, (2017).
483

ACKNOWLEDGEMENTS

We thank the entire Walter and Manglik labs for facilitating the development and rapid execution

of this large-scale collaborative effort. We thank Sebastian Bernales and Tony De Fougerolles

for advice and helpful discussion, and Jonathan Weissman for input into the project and reagent

and machine use. We thank Jim Wells for providing the ACE2 ECD-Fc construct, Jason

McLellan for providing Spike, RBD, and ACE2 constructs, and Florian Krammer for providing an

RBD construct. We thank Jesse Bloom for providing the ACE2 expressing HEK293T cells as

well as the plasmids for pseudovirus work. We thank George Meigs and other Beamline staff at

ALS, 8.3.1 for their help in data collection. We thank Randy A. Albrecht for oversight of the

conventional BSL3 biocontainment facility at the Icahn School of Medicine at Mount Sinai.

Funding

This work was supported by the UCSF COVID-19 Response Fund, a grant from Allen &

Company, and supporters of the UCSF Program for Breakthrough Biomedical Research

(PBBR), which was established with support from the Sandler Foundation. Further support was

provided by the National Institutes of Health (NIH) grant DP5OD023048 (A.Manglik). Cryo-EM

equipment at UCSF is partially supported by NIH grants S10OD020054 and S10OD021741.

Work by M.Vignuzzi was funded by the Laboratoire d'Excellence grant ANR-10-LABX-62-IBEID

and the URGENCE COVID-19 Institut Pasteur fundraising campaign. The radiolytic hydroxyl

radical footprinting is supported by NIH 1R01GM126218. The Advanced Light Source is

supported by the Office of Science, Office of Biological and Environmental Research, of the

U.S. DOE under contract DE-AC02-05CH11231. S.Sangwan was supported by a Helen Hay

Whitney postdoctoral fellowship. C.B.Billesbølle acknowledges support from the Alfred Benzon

Foundation. K.Leon was funded by NIH/NINDS award F31NS113432 and a UCSF Discovery

Fellowship from the Otellini Family. C.Puchades and V.Belyy are Fellows of the Damon Runyon

Cancer Research Foundation. H. Kratochvil was supported by a Ruth L. Kirschstein NRSA

Postdoctoral Fellowship (F32GM125217). This research was also partly funded by CRIP

(Center for Research for Influenza Pathogenesis), a NIAID supported Center of Excellence for

Influenza Research and Surveillance (CEIRS, contract # HHSN272201400008C), by DARPA

grant HR0011-19-2-0020, by an administrative supplement to NIAID grant U19AI142733, and

by the generous support of the JPB Foundation and the Open Philanthropy to A.Garcia-Sastre.

M.Ott acknowledges support through a gift from the Roddenberry Foundation. P.Walter is an

Investigator of the Howard Hughes Medical Institute. A.Manglik acknowledges support from the

Author Contributions

M.Schoof purified Spike*, RBD, and ACE2 proteins, performed yeast display selections to

identify and affinity mature nanobodies, expressed and purified nanobodies, tested activity in

cell-based assays, cloned, expressed, and purified multivalent nanobody constructs, and

coordinated live virus experiments. B.Faust purified and characterized Spike* protein and

candidate nanobodies, developed, performed and analyzed SPR experiments for Spike* and

RBD-nanobody affinity determination, developed, performed and analyzed SPR binning,

experiments, determined optimal freezing conditions for cryo-EM experiments, processed,

refined and generated figures for Nb6, Nb11, and mNb6 EM datasets. R.A. Saunders expressed

and purified ACE2 and nanobodies, developed and performed cell-based assays for inhibition of

Spike* binding and pseudovirus assays for determining nanobody efficacy. S.Sangwan

expressed and purified Spike*, RBD, ACE2-Fc, and nanobodies, processed cryo-EM data,

optimized RBD-nanobody complexes for crystallography, grew crystals of mNb6, collected

diffraction data, and refined the X-ray crystal structure of mNb6. V.Rezelj tested efficacy of

nanobody constructs in live SARS-CoV-2 infection assays under the guidance of M.Vignuzzi.

N.Hoppe purified nanobodies, developed, performed and analyzed SPR binning experiments,

developed performed and analyzed variable Nb6-bi and Nb6-tri association experiments, and

performed thermal melting stability assays for nanobody constructs. M.Boone developed

approaches to express and purify nanobodies from *Pichia pastoris* and developed, performed,

and analyzed approaches to quantify nanobody efficacy in live virus assays. C.B.Billesbølle

expressed and purified Spike*, generated affinity maturation library for Nb6, performed yeast

display selections to identify mNb6, and built the synthetic yeast nanobody library with J.Liang.

I.Deshpande expressed and purified nanobody constructs. B.S.Zha performed live SARS-CoV-2

virus assays to test nanobody efficacy with guidance from O.Rosenberg. C.R.Simoneau and

K.Leon performed live SARS-CoV-2 virus assays to test nanobody efficacy with guidance from

M.Ott. K.M.White performed live SARS-CoV-2 virus assays to test nanobody efficacy with

guidance from A.Garcia-Sastre. A.W.Barile-Hill performed SPR experiments. A.A.Anand,

N.Dobzinski, B.Barsi-Rhyne, and Y.Liu. assisted in cloning, expression, and purification of

nanobody and pseudovirus constructs. V.Belyy performed single-molecule nanobody-Spike*

interaction studies. S.Nock prepared media and coordinated lab usage during UCSF's partial

shutdown. M.Zimanyi and S.Gupta performed radiolytic footprinting experiments with guidance

Competing Interests

Data and Materials Availability

All data generated or analyzed during this study are included in this published article and its
Supplementary Information. Crystallographic coordinates and structure factors for mNb6 have
been deposited in the Protein Data Bank under accession code XXXX. Coordinates for
Spike*:Nb6 and Spike*:mNb6 complexes have been deposited in the Protein Data Bank under
accession codes XXXX and XXXX, respectively. Maps for Spike*:Nb6, Spike*:Nb11, and
Spike*:mN6 have been deposited in the Electron Microscopy Data Bank under accession codes
XXXXX (Spike*-Nb6 Open), XXXXX (Spike*-Nb6 Closed), XXXXX (Spike*-Nb11 Open), XXXXX
(Spike*-Nb11 Closed), XXXXX (Spike*-mNb6 Closed). Plasmids for nanobody constructs used
in this study are available under a Material Transfer Agreement with the University of California,
San Francisco.
582

     MAIN TEXT FIGURES

Figure 1

Figure 1. Discovery of two distinct classes of anti-Spike nanobodies. A, Selection strategy

for identification of anti-Spike nanobodies that disrupt Spike-ACE2 interactions using magnetic

bead selections (MACS) or fluorescence activated cell sorting (FACS). **B,** Flow cytometry of

yeast displaying Nb6 (a Class I nanobody) or Nb3 (a Class II nanobody). Nb6 binds Spike*-

Alexa 647 and receptor binding domain (RBD-Alexa 647). Nb6 binding to Spike* is completely

disrupted by an excess (1.4 µM) of ACE2-Fc. Nb3 binds Spike*, but not the RBD. Nb3 binding

to Spike* is partially decreased by ACE2-Fc. **C,** SPR of Nb6 and Nb3 binding to either Spike* or

RBD. Red traces are raw data and global kinetic fits are shown in black. Nb3 shows no binding

to RBD. **D,** SPR experiments with immobilized Spike* show that Class I and Class II nanobodies

can bind Spike* simultaneously. By contrast, two Class I nanobodies or Class II nanobodies do

not bind simultaneously. **E,** Nanobody inhibition of 1 nM Spike*-Alexa 647 binding to ACE2

expressing HEK293T cells. n = 3 (ACE2, Nb3) or 5 (Nb6, Nb11) biological replicates. All error

bars represent s.e.m.

Figure 2

599
600

Figure 2. Cryo-EM structures of Nb6 and Nb11 bound to Spike. A, Cryo-EM maps of Spike*-
Nb6 complex in either closed (left) or open (right) Spike* conformation. **B,** Cryo-EM maps of
Spike*-Nb11 complex in either closed (left) or open (right) Spike* conformation. The top views
show receptor binding domain (RBD) up- or down-states. **C,** Nb6 straddles the interface of two
down-state RBDs, with CDR3 reaching over to an adjacent RBD. **D,** Nb11 binds a single RBD in
the down-state (displayed) or similarly in the up-state. No cross-RBD contacts are made by
Nb11 in either RBD up- or down-state. **E,** Comparison of RBD epitopes engaged by ACE2
(purple), Nb6 (red), or Nb11 (green). Both Nb11 and Nb6 directly compete with ACE2 binding.

Figure 3

Figure 3. Multivalency improves nanobody affinity and inhibitory efficacy. **A,** SPR of Nb6 and multivalent variants. Red traces show raw data and black lines show global kinetic fit for Nb6 and independent fits for association and dissociation phases for Nb6-bi and Nb6-tri. **B,** Dissociation phase SPR traces for Nb6-tri after variable association time ranging from 4 to 520 s. Curves were normalized to maximal signal at the beginning of the dissociation phase. Percent fast phase is plotted as a function of association time (right) with a single exponential fit. n = 3 independent biological replicates. **C,** Inhibition of pseudotyped lentivirus infection of ACE2 expressing HEK293T cells. n = 3 biological replicates for all but Nb11-tri (n = 2) **D,** Inhibition of live SARS-CoV-2 virus. Representative biological replicate with n = 3 (right panel) or 4 (left panel) technical replicates per concentration. n = 3 biological replicates for all but Nb3 and Nb3-tri (n = 2). All error bars represent s.e.m.

Figure 4. Affinity maturation of Nb6 yields a picomolar SARS-CoV-2 neutralizing molecule. A, A saturation mutagenesis library of Nb6 was subjected to two rounds of selection to identify consensus mutations I27Y and P105Y. **B,** SPR of mNb6 and mNb6-tri binding to immobilized Spike*. Red traces show raw data and black lines show global kinetic fit. No dissociation was observed for mNb6-tri over 10 minutes. **C,** Cryo-EM structure of Spike*-mNb6 complex. **D,** Comparison of receptor binding domain (RBD) engagement by Nb6 and mNb6. One RBD was used to align both structures (RBD align), demonstrating changes in Nb6 and mNb6 position and the adjacent RBD. **E,** Comparison of mNb6 complementarity determining regions in either the cryo-EM structure of the Spike*-mNb6 complex or an X-ray crystal structure of mNb6 alone. **F,** CDR1 of Nb6 and mNb6 binding to the RBD. As compared to I27 in Nb6, Y27 of mNb6 hydrogen bonds to Y453 and optimizes pi-pi and pi-cation interactions with the RBD. **G,** CDR3 of Nb6 and mNb6 binding to the RBD demonstrating a large conformational rearrangement of the entire loop in mNb6. **H,** Comparison of closed Spike* bound to mNb6 and Nb6. Rotational axis for RBD movement is highlighted. **I,** Inhibition of pseudotyped lentivirus infection of ACE2 expressing HEK293T cells by mNb6 and mNb6-tri. n = 3 biological replicates

639   **J,** mNb6 and mNb6-tri inhibit SARS-CoV-2 infection of VeroE6 cells in a plaque assay.

640   Representative biological replicate with n = 4 technical replicates per concentration. n = 3

641   biological replicates for all samples. All error bars represent s.e.m.

Figure 5

Figure 5. Nb6 and its derivates are robust proteins. A, Thermal denaturation of nanobodies assessed by circular dichroism measurement of molar ellipticity at 204 nm. Apparent melting temperatures (T_m) for each nanobody are indicated. **B,** Nanobody inhibition of 1 nM Spike*-Alexa 647 binding to ACE2 expressing HEK293T cells after incubation at either 25 °C or 50 °C for 1 hour or after aerosolization. **C,** Size exclusion chromatography of nanobodies after lyophilization or aerosolization. **D,** Summary table of SPR kinetics data and affinities for aerosolized or lyophilized mNb6 and mNb6-tri.

652 **Table 1. Anti-Spike nanobody affinity and neutralization potency**

Nanobody	Class	Spike* Binding			RBD Binding			Spike* Competition IC$_{50}$ (s.e.m) (M)[a]	SARS-CoV-2 Pseudovirus IC$_{50}$ (s.e.m.) (M)[b]	Live SARS-CoV-2 IC$_{50}$ (s.e.m.) (M)[c]
		k_a (M^{-1}s^{-1})	k_d (s^{-1})	K$_D$ (M)	k_a (M^{-1}s^{-1})	k_d (s^{-1})	K$_D$ (M)			
Nb2	I	9.0×10^5	5.3×10^{-1}	5.9×10^{-7}	1.0×10^6	9.9×10^{-1}	9.7×10^{-7}	8.3×10^{-6} (1.7×10^{-6})	NP	NP
Nb3	II	1.8×10^6	1.1×10^{-1}	6.1×10^{-8}	NB			NC	3.9×10^{-6} (7.9×10^{-7})	3.0×10^{-6} (3.2×10^{-7})
Nb6	I	2.7×10^5	5.6×10^{-2}	2.1×10^{-7}	2.1×10^6	8.7×10^{-2}	4.1×10^{-8}	3.7×10^{-7} (4.9×10^{-8})	2.0×10^{-6} (3.5×10^{-7})	3.3×10^{-6} (7.2×10^{-7})
Nb8	I	1.4×10^5	8.1×10^{-1}	5.8×10^{-6}	6.6×10^5	3.3×10^{-1}	5.1×10^{-7}	4.8×10^{-6} (4.9×10^{-7})	NP	NP
Nb11	I	1.2×10^6	1.6×10^{-1}	1.4×10^{-7}	3.2×10^6	2.4×10^{-1}	7.6×10^{-8}	5.4×10^{-7} (1.2×10^{-7})	2.4×10^{-6} (5.4×10^{-7})	NP
Nb12	I	1.2×10^2	2.0×10^{-4}	1.6×10^{-6}	Biphasic	Biphasic	Biphasic	2.5×10^{-7} (5.5×10^{-8})	1.2×10^{-6} (9.0×10^{-7})	NP
Nb15	I	1.7×10^5	2.3×10^{-1}	1.3×10^{-6}	6.0×10^5	2.2×10^{-1}	3.6×10^{-7}	2.2×10^{-6} (2.5×10^{-7})	6.7×10^{-6} (3.6×10^{-6})	NP
Nb16	I	1.1×10^5	1.3×10^{-1}	1.3×10^{-6}	NP			9.5×10^{-7} (1.1×10^{-7})	NP	NP
Nb17	II	7.3×10^5	2.0×10^{-1}	2.7×10^{-7}	NB			NC	7.6×10^{-6} (1.0×10^{-6})	NP
Nb18	II	1.4×10^5	6.4×10^{-3}	4.5×10^{-8}	NB			5.2×10^{-5} (1.5×10^{-5})	NP	NP
Nb19	I	2.4×10^4	1.1×10^{-1}	4.5×10^{-6}	1.0×10^5	8.9×10^{-2}	8.8×10^{-7}	4.1×10^{-6} (4.9×10^{-7})	2.4×10^{-5} (7.7×10^{-6})	NP
Nb24	I	9.3×10^5	2.7×10^{-1}	2.9×10^{-7}	2.4×10^6	3.5×10^{-1}	1.5×10^{-7}	7.5×10^{-7} (1.0×10^{-7})	NP	NP
ACE2	N/A	2.7×10^5	1.2×10^{-2}	4.4×10^{-8}	NP	NP	NP	1.7×10^{-7} (6.6×10^{-8})	6.2×10^{-7} (1.7×10^{-7})	NP
mNb6	I	1.0×10^6	4.5×10^{-4}	4.5×10^{-10}	1.1×10^6	6.4×10^{-4}	5.6×10^{-10}	1.3×10^{-9} (4.1×10^{-10})	6.3×10^{-9} (1.6×10^{-9})	1.2×10^{-8} (2.5×10^{-9})
Nb3-bi	II	NP	NP	NP	NP	NP	NP	NP	3.6×10^{-7} (1.5×10^{-7})	1.8×10^{-7} (1.2×10^{-8})
Nb3-tri	II	Biphasic	Biphasic	Biphasic	NP	NP	NP	4.1×10^{-8} (1.6×10^{-8})	4.0×10^{-7} (1.6×10^{-7})	1.4×10^{-7} (4.9×10^{-8})
Nb6-bi	I	Biphasic	Biphasic	Biphasic	NP	NP	NP	NP	6.3×10^{-8} (1.5×10^{-8})	NP
Nb6-tri	I	Biphasic	Biphasic	Biphasic	NP	NP	NP	1.5×10^{-9} (5.2×10^{-10})	1.2×10^{-9} (2.5×10^{-10})	1.6×10^{-10} (2.6×10^{-11})
Nb11-tri	I	Biphasic	Biphasic	Biphasic	NP	NP	NP	NP	5.1×10^{-8} (1.6×10^{-8})	NP
ACE2-Fc	N/A	NP	NP	NP	NP	NP	NP	5.3×10^{-9} (2.5×10^{-9})	4.0×10^{-8} (8.8×10^{-9})	2.6×10^{-8} (8.5×10^{-9})
mNb6-tri	I	1.4×10^6	$<1.0 \times 10^{-6}$	$<1.0 \times 10^{-12}$	NP	NP	NP	4.0×10^{-10} (1.4×10^{-10})	1.2×10^{-10} (2.8×10^{-11})	5.4×10^{-11} (1.0×10^{-11})

[a]Average values from n = 5 biological replicates for Nb6, Nb11, Nb15, Nb19 are presented, all

others were tested with n = 3 biological replicates.

[b]Average values from n = 2 biological replicates for Nb12, Nb17, and Nb11-tri are presented, all

others were tested with n = 3 biological replicates.

[c]Average values from n = 2 biological replicates for Nb3, Nb3-bi, and Nb3-tri. n = 3 biological

replicates for all others.

NB – no binding

NC – no competition

NP – not performed