An ultra-high affinity synthetic nanobody blocks SARS-CoV-2 infection by locking Spike into an inactive conformation

Michael Schoof[1,2#*], Bryan Faust[1,2,3,4#], Reuben A. Saunders[1,5#], Smriti Sangwan[1,2#], Veronica Rezelj[6#], Nick Hoppe[3,4], Morgane Boone[1,2], Christian Bache Billesbølle[3,4], Marcell Zimanyi[1,2,3], Ishan Deshpande[3,4], Jiahao Liang[3], Aditya A. Anand[1,2], Niv Dobzinski[1,2], Beth Shoshana Zha[7], Benjamin Barsi-Rhyne[3], Vladislav Belyy[1,2], Andrew W. Barile-Hill[8], Sayan Gupta[9], Camille R. Simoneau[10,11,12], Kristoffer Leon[10,11,12], Kris M. White[13,14], Silke Nock[1,2], Yuwei Liu[1,2], Nevan J. Krogan[4,5,10,11], Corie Y. Ralston[9], Danielle L. Swaney[4,5,10,11], Adolfo García-Sastre[13,14], Melanie Ott[10,11,12], Marco Vignuzzi[6], QCRG Structural Biology Consortium[4], Peter Walter[1,2*], Aashish Manglik[3,4,10,15*]

[1]Howard Hughes Medical Institute, University of California at San Francisco, San Francisco, CA, USA.
[2]Department of Biochemistry and Biophysics, University of California at San Francisco, San Francisco, CA, USA.
[3]Department of Pharmaceutical Chemistry, University of California at San Francisco, San Francisco, CA, USA.
[4]Quantitative Biosciences Institute (QBI) Coronavirus Research Group Structural Biology Consortium, University of California, San Francisco, CA, USA.
[5]Department of Cellular and Molecular Pharmacology, University of California at San Francisco, San Francisco, CA, USA.
[6]Viral Populations and Pathogenesis Unit, CNRS UMR 3569, Institut Pasteur, 75724, Paris, cedex 15, France
[7]Department of Pulmonary, Critical Care, Allergy and Sleep Medicine, University of California San Francisco, San Francisco, CA 94158, USA.
[8]Cytiva Life Sciences, Marlborough, MA, USA.
[9]Molecular Biophysics and Integrated Bioimaging and the Molecular Foundry, Lawrence Berkeley National Laboratory, Berkeley, CA, USA.
[10]Quantitative Biosciences Institute (QBI), University of California San Francisco, San Francisco, CA, USA.
[11]J. David Gladstone Institutes, San Francisco, CA, USA.
[12]Department of Medicine, University of California San Francisco, San Francisco, CA, USA.
[13]Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY, USA.
[14]Global Health and Emerging Pathogens Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA.
[15]Department of Anesthesia and Perioperative Care, University of California at San Francisco, San Francisco, CA, USA.

[#] These authors contributed equally
[*] To whom correspondence should be addressed; Email: michael@walterlab.ucsf.edu, peter@walterlab.ucsf.edu, Aashish.Manglik@ucsf.edu

Abstract

Without an effective prophylactic solution, infections from SARS-CoV-2 continue to rise
worldwide with devastating health and economic costs. SARS-CoV-2 gains entry into host cells
via an interaction between its Spike protein and the host cell receptor angiotensin converting
enzyme 2 (ACE2). Disruption of this interaction confers potent neutralization of viral entry,
providing an avenue for vaccine design and for therapeutic antibodies. Here, we develop single-
domain antibodies (nanobodies) that potently disrupt the interaction between the SARS-CoV-2
Spike and ACE2. By screening a yeast surface-displayed library of synthetic nanobody
sequences, we identified a panel of nanobodies that bind to multiple epitopes on Spike and
block ACE2 interaction via two distinct mechanisms. Cryogenic electron microscopy (cryo-EM)
revealed that one exceptionally stable nanobody, Nb6, binds Spike in a fully inactive
conformation with its receptor binding domains (RBDs) locked into their inaccessible down-
state, incapable of binding ACE2. Affinity maturation and structure-guided design of
multivalency yielded a trivalent nanobody, mNb6-tri, with femtomolar affinity for SARS-CoV-2
Spike and picomolar neutralization of SARS-CoV-2 infection. mNb6-tri retains stability and
function after aerosolization, lyophilization, and heat treatment. These properties may enable
aerosol-mediated delivery of this potent neutralizer directly to the airway epithelia, promising to
yield a widely deployable, patient-friendly prophylactic and/or early infection therapeutic agent to
stem the worst pandemic in a century.

Introduction

Over the last two decades, three zoonotic β-coronaviruses have entered the human population, causing severe respiratory symptoms with high mortality [1-3]. The ongoing COVID-19 pandemic is caused by SARS-CoV-2, the most readily transmissible of these three coronaviruses [4-7]. SARS-CoV-2 has wrecked the world's economy and societies to an unprecedented extent, to date (Aug. 3, 2020) causing 691,320 reported deaths around the globe [8]. Although public health measures have slowed its spread in many regions, infection hotspots keep reemerging. No successful vaccine or preventive treatment has yet been manufactured for any coronavirus, and the time to develop an effective and broadly available vaccine for SARS-CoV-2 remains uncertain. The development of novel therapeutic and prophylactic approaches thus remains essential, both as temporary stopgaps until an effective vaccine is generated and as permanent solutions for those segments of the population for which vaccination proves ineffective or contraindicated.

Coronavirus virions are bounded by a membrane envelope that contains ~25 copies of the homotrimeric transmembrane spike glycoprotein (Spike) responsible for virus entry into the host cell [9]. The surface-exposed portion of Spike is composed of two domains, S_1 and S_2 [10]. The S_1 domain mediates the interaction between virus and its host cell receptor, the angiotensin converting enzyme 2 (ACE2), while the S_2 domain catalyzes fusion of the viral and host cell membranes [3, 11-13]. During its biogenesis, the Spike protein is proteolytically cleaved between the S_1 and S_2 domains, which primes the virus for cellular entry [10]. Contained within S_1 is the receptor binding domain (RBD), which directly binds to ACE2. The RBD is attached to the body of Spike by a flexible region and can exist in an inaccessible down-state or an accessible up-state [14, 15]. Binding to ACE2 requires the RBD in the up-state and enables cleavage by host proteases TMPRSS2 or cathepsin, triggering a dramatic conformational change in S_2 that enables viral entry [16]. In SARS-CoV-2 virions, Spike oscillates between an active, open conformation with at least one RBD in the up-state and an inactive, closed conformation with all RBDs in the down-state [9, 11, 14, 15].

By screening a high-complexity yeast surface-displayed library of synthetic nanobodies, we have uncovered a collection of nanobodies that block the Spike-ACE2 interaction. Biochemical and structural studies revealed that two classes of these nanobodies act in distinct ways to prevent ACE2 binding. Combining affinity maturation and structure-guided multimerization, we optimized these agents and generated Spike binders that match or exceed the potency of most

monoclonal antibodies disclosed to date. Our lead neutralizing molecule, mNb6-tri, blocks

SARS-CoV-2 entry in human cells at picomolar efficacy and withstands aerosolization,

lyophilization, and elevated temperatures. mNb6-tri provides a promising approach to deliver a

potent SARS-CoV-2 neutralizing molecule directly to the airways for prophylaxis or therapy.

Results

Synthetic nanobodies that disrupt Spike-ACE2 interaction

To isolate nanobodies that neutralize SARS-CoV-2, we screened a yeast surface-displayed

library of >2×10^9 synthetic nanobody sequences. Our strategy was to screen for binders to the

full Spike protein ectodomain, in order to capture not only those nanobodies that would compete

by binding to the ACE2-binding site on the RBD directly but also those that might bind

elsewhere on Spike and block ACE2 interaction through indirect mechanisms. We used a

mutant form of SARS-CoV-2 Spike, Spike*, as the antigen [15]. Spike* lacks one of the two

activating proteolytic cleavage sites between the S_1 and S_2 domains and introduces two

mutations to stabilize the pre-fusion conformation. Spike* expressed in mammalian cells binds

ACE2 with a K_D = 44 nM (Supplementary Fig. 1), consistent with previous reports [17]. Next, we

labeled Spike* with biotin or with fluorescent dyes and selected nanobody-displaying yeast over

multiple rounds, first by magnetic bead binding and then by fluorescence-activated cell sorting

(Fig. 1A).

Three rounds of selection yielded 21 unique nanobodies that bound Spike* and showed

decreased Spike* binding in the presence of ACE2. Closer inspection of their binding properties

revealed that these nanobodies fall into two distinct classes. One group (Class I) binds the RBD

and competes with ACE2 (Fig. 1B). A prototypical example of this class is nanobody Nb6, which

binds to Spike* and to RBD alone with a K_D of 210 nM and 41 nM, respectively (Fig. 1C; Table

1). Another group (Class II), exemplified by nanobody Nb3, binds to Spike* (K_D = 61 nM), but

displays no binding to RBD alone (Fig. 1C, Table 1). In the presence of excess ACE2, binding of

Nb6 and other Class I nanobodies is blocked entirely, whereas binding of Nb3 and other Class II

nanobodies is decreased only moderately (Fig. 1B). These results suggest that Class I

nanobodies target the RBD to block ACE2 binding, whereas Class II nanobodies target other

epitopes and decrease ACE2 interaction with Spike allosterically or through steric interference.

Indeed, surface plasmon resonance (SPR) experiments demonstrate that Class I and Class II

nanobodies can bind Spike* simultaneously (Fig. 1D).

Analysis of the kinetic rate constants for Class I nanobodies revealed a consistently greater

association rate constant (k_a) for nanobody binding to the isolated RBD than to full-length Spike*

(Table 1), which suggests that RBD accessibility influences the K_D. We next tested the efficacy

of our nanobodies, both Class I and Class II, to inhibit binding of fluorescently labeled Spike* to

ACE2-expressing HEK293 cells (Table 1, Fig. 1E). Class I nanobodies emerged with highly

variable activity in this assay with Nb6 and Nb11 as two of the most potent clones with IC_{50}

values of 370 and 540 nM, respectively (Table 1). For unexplained reasons, Class II nanobodies

showed little to no activity in this assay (Table 1, Fig. 1E).

Going forward, we prioritized two Class I nanobodies, Nb6 and Nb11, that combine potent

Spike* binding with relatively small differences in k_a between binding to Spike* or RBD. We

reasoned that the epitopes recognized by Nb6 and Nb11 would be more readily accessible in

the Spike protein on intact virions. For Class II nanobodies we prioritized Nb3 because of its

optimal stability and yield during purification.

Nb6 and Nb11 target the RBD and directly compete with ACE2

To define the binding sites of Nb6 and Nb11, we determined their cryogenic electron

microscopy (cryo-EM) structures bound to Spike* (Fig. 2A-B, Supplementary Fig. 2-4,

Supplementary Table 1). Both nanobodies recognize RBD epitopes that overlap the ACE2

binding site (Fig. 2E). For Nb6 and Nb11, we resolved nanobody binding to both the open and

closed conformations of Spike*. We obtained a 3.0 Å map of Nb6 bound to closed Spike*, which

enabled modeling of the Nb6-Spike* complex (Fig. 2A), including the complementarity

determining regions (CDRs). We also obtained lower resolution maps for Nb6 bound to open

Spike* (3.8 Å), Nb11 bound to open Spike* (4.2 Å), and Nb11 bound to closed Spike* (3.7 Å).

For these lower resolution maps, we could define the nanobody's binding orientation but not

accurately model the CDRs.

Nb6 bound to closed Spike* straddles the interface between two adjacent RBDs. The majority of

the contacting surfaces are contributed by CDR1 and CDR2 of Nb6 (Fig. 2C). CDR3 contacts

the adjacent RBD that is counterclockwise positioned when viewed from the top of Spike* (Fig.

2C). The binding of one Nb6 therefore stabilizes two adjacent RBDs in the down-state. We

surmise that this initial binding event pre-organizes the binding site for a second and third Nb6

molecule to stabilize the closed Spike* conformation. Indeed, binding of two Nb6 molecules

would lock all three RBDs into the down-state, thus highly favoring binding of a third Nb6

because binding would not entail any further entropic cost. By contrast, Nb11 bound to down-

state RBDs only contacts a single RBD (Fig. 2D).

Nb3 interacts with the Spike S$_1$ domain external to the RBD

Our attempts to determine the binding site of Nb3 by cryo-EM proved unsuccessful. We

therefore turned to radiolytic hydroxyl radical footprinting to determine potential binding sites for

Nb3. Spike*, either apo or bound to Nb3, was exposed to 5-50 milliseconds of synchrotron X-ray

radiation to label solvent exposed amino acids with hydroxyl radicals. Radical-labeled amino

acids were subsequently identified and quantified by mass spectrometry of trypsin/Lys-C or Glu-

C protease digested Spike*[18]. Two neighboring surface residues on the S$_1$ domain of Spike

(M177 and H207) emerged as highly protected sites in the presence of Nb3 (Supplementary

Fig. 5). The degree of protection is consistent with prior observations of antibody-antigen

interactions by hydroxyl radical footprinting [19]. Both M177 and H207 are greater than 40 Å

distant from the ACE2 binding site on the RBD, suggesting that Nb3 may inhibit Spike-ACE2

interactions through allosteric means.

Rationally engineered multivalency increases potency

The structure of Nb6 bound to closed Spike* enabled us to engineer bivalent and trivalent

nanobodies predicted to lock all RBDs in the down-state. To this end, we inserted flexible Gly-

Ser linkers of either 15 or 20 amino acids to span the 52 Å distance between adjacent Nb6

monomers bound to down-state RBDs in closed Spike* (Supplementary Fig. 6). Both linker

lengths are too short to span the distance (72 Å) between Nb6 bound to a down-state RBD and

an up-state RBD that would co-exist in an open Spike. Moreover, binding of three RBDs in the

previously reported conformation of Nb6-bound open Spike* would be physically impossible

even with longer linker length because of steric clashes (Supplementary Fig. 6). By contrast, the

minimum distance between adjacent Nb11 monomers bound to either open or closed Spike* is

68 Å (Supplementary Fig. 6). We therefore predicted that multivalent binding by Nb6 constructs

would display significantly slowed dissociation rates due to the enhanced avidity afforded by

Spike's trimeric architecture.

We assessed multivalent Nb6 binding to Spike* by SPR. Both bivalent Nb6 with a 15 amino acid

linker (Nb6-bi) and trivalent Nb6 with two 20 amino acid linkers (Nb6-tri) dissociate from Spike*

in a biphasic manner. The dissociation phase can be fitted to two components: a fast phase with

kinetic rate constants k_{d1} of 2.7x10^{-2} s^{-1} for Nb6-bi and 2.9x10^{-2} s^{-1} for Nb6-tri, which are of the

same magnitude as that observed for monovalent Nb6 (k_d = 5.6x10^{-2} s^{-1}) and a slow phase that

is dependent on avidity (k_{d2} = 3.1x10^{-4} for Nb6-bi and k_{d2} < 1.0x10^{-6} s^{-1} for Nb6-tri, respectively)

(Fig. 3A). The relatively similar k_d for the fast phase suggests that a fraction of the observed

binding for the multivalent constructs is nanobody binding to a single Spike* RBD. By contrast,

the slow dissociation phase of Nb6-bi and Nb6-tri indicates engagement of two or three RBDs.

We observed no dissociation for the slow phase of Nb6-tri over 10 minutes, indicating an upper

boundary for k_{d2} of 1x10^{-6} s^{-1} and subpicomolar affinity. This measurement remains an upper-

bound estimate rather than an accurate measurement because the technique is limited by the

intrinsic dissociation rate of Spike* from the chip imposed by the chemistry used to immobilize

Spike*.

We reasoned that the biphasic dissociation behavior could be explained by a slow

interconversion between up- and down-state RBDs, with conversion to the more stable down-

state required for full trivalent binding. According to this view, a single domain of Nb6-tri

engaged with an up-state RBD would dissociate rapidly. The system would then re-equilibrate

as the RBD flips into the down-state, eventually allowing Nb6-tri to trap all RBDs in closed

Spike*. To test this notion directly, we varied the time allowed for Nb6-tri binding to Spike*.

Indeed, we observed an exponential decrease in the percent fast-phase with a $t_{1/2}$ of 65 s (Fig.

3B), which, we surmise, reflects the timescale of conversion between the RBD up- and down-

states in Spike*. Taken together, dimerization and trimerization of Nb6 afforded 750-fold and

>200,000-fold gains in K_D, respectively.

Class I and II nanobodies prevent SARS-CoV-2 infection

We next tested the neutralization activity of trivalent versions of our top Class I (Nb6 and Nb11)

and Class II (Nb3) nanobodies against SARS-CoV-2 pseudotyped lentivirus. In this assay,

SARS-CoV-2 Spike is expressed as a surface protein on a lentiviral particle that contains a

ZsGreen reporter gene, which is integrated and expressed upon successful viral entry into cells

harboring the ACE2 receptor [20]. Nb6 and Nb11 inhibited pseudovirus infection with IC$_{50}$

values of 2.0 µM and 2.4 µM, respectively, and Nb3 inhibited pseudovirus infection with an IC$_{50}$

of 3.9 µM (Fig. 3C, Table 1). Nb6-tri shows a 2000-fold enhancement of inhibitory activity, with

an IC$_{50}$ of 1.2 nM, whereas trimerization of Nb11 and Nb3 resulted in more modest gains of 40-

and 10-fold (51 nM and 400 nM), respectively (Fig. 3C).

We next confirmed these neutralization activities with a viral plaque assay using live SARS-

CoV-2 virus infection of VeroE6 cells. Consistent with its activity against pseudotyped lentivirus,

Nb6-tri proved exceptionally potent, neutralizing SARS-CoV-2 with an average IC_{50} of 160 pM

(Fig. 3D). Nb3-tri neutralized SARS-CoV-2 with an average IC_{50} of 140 nM (Fig. 3D).

Affinity maturation yields a femtomolar K_D Spike inhibitor

We further optimized the potency of Nb6 by selecting high-affinity variants. To this end, we

prepared a new library, starting with the Nb6 coding sequence, in which we varied each amino

acid position of all three CDRs by saturation mutagenesis (Fig. 4A). After two rounds of

magnetic bead-based selection, we isolated a population of high-affinity clones. Sequencing

revealed two highly penetrant mutations: I27Y in CDR1 and P105Y in CDR3. We incorporated

these two mutations into Nb6 to generate matured Nb6 (mNb6), which binds with 500-fold

increased affinity to Spike* as measured by SPR (Fig. 4B). As a monomer, mNb6 inhibits both

pseudovirus and live SARS-CoV-2 infection with low nanomolar potency, a ~200-fold

improvement compared to Nb6 (Fig. 4I-J, Table 1).

A 2.9 Å cryo-EM structure of mNb6 bound to Spike* shows that, like the parent nanobody Nb6,

mNb6 binds to closed Spike (Fig. 4C, Supplementary Fig. 7). The higher resolution map allowed

us to build a model with high confidence and determine the effects of the I27Y and P105Y

substitutions. mNb6 induces a slight rearrangement of the down-state RBDs as compared to

both previously determined structures of apo-Spike* and Spike* bound to Nb6, inducing a 9°

rotation of the RBD away from the central three-fold symmetry axis (Fig. 4H) [14, 15]. This

deviation likely arises from a different interaction between CDR3 and Spike*, which nudges the

RBDs into a new resting position. While the I27Y substitution optimizes local contacts between

CDR1 in its original binding site on the RBD, the P105Y substitution leads to a marked

rearrangement of CDR3 in mNb6 (Fig. 4F-G). This conformational change yields a different set

of contacts between mNb6 CDR3 and the adjacent RBD (Fig. 4D). Remarkably, an X-ray crystal

structure of mNb6 alone revealed dramatic conformational differences in CDR1 and CDR3

between free and Spike*-bound mNb6, suggestive of significant conformational heterogeneity

for the unbound nanobodies and induced-fit rearrangements upon binding to Spike* (Fig. 4E).

The binding orientation of mNb6 is similar to that of Nb6, supporting the notion that our

multivalent design would likewise enhance binding affinity. Unlike Nb6-tri, trivalent mNb6

(mNb6-tri) bound to Spike with no observable fast-phase dissociation and no measurable

dissociation over ten minutes, yielding an upper bound for the dissociation rate constant k_d of

1.0×10^{-6} s^{-1} ($t_{1/2} > 8$ days) and a K_D of <1 pM (Fig. 4B). As above, more precise measurements

of the dissociation rate are precluded by the surface chemistry used to immobilize Spike*.

mNb6-tri displays further gains in potency in both pseudovirus and live SARS-CoV-2 infection

assays with IC_{50} values of 120 pM (5.0 ng/mL) and 54 pM (2.3 ng/mL), respectively (Fig. 4H-I,

Table 1). Given the sub-picomolar affinity observed by SPR, it is likely that these viral

neutralization potencies reflect the lower limit of the assays. mNb6-tri is therefore an

exceptionally potent SARS-CoV-2 neutralizing antibody, among the most potent molecules

disclosed to date.

Nb6, Nb6-tri, mNb6, and mNb6-tri are robust proteins

One of the most attractive properties that distinguishes nanobodies from traditional monoclonal

antibodies is their extreme stability [21]. We therefore tested Nb6, Nb6-tri, mNb6, and mNb6-tri

for stability regarding temperature, lyophilization, and aerosolization. Temperature denaturation

experiments using circular dichroism measurements to assess protein unfolding revealed

melting temperatures of 66.9, 62.0, 67.6, and 61.4 °C for Nb6, Nb6-tri, mNb6 and mNb6-tri,

respectively (Fig 5A). Aerosolization and prolonged heating of Nb6, mNb6, and mNb6-tri for 1

hour at 50°C induced no loss of activity (Fig 5B). Moreover, mNb6 and mNb6-tri were stable to

lyophilization and to aerosolization using a mesh nebulizer, showing no aggregation by size

exclusion chromatography and preserved high affinity binding to Spike* (Fig. 5C-D).

Discussion

There is a pressing need for prophylactics and therapeutics against SARS-CoV-2 infection.

Most recent strategies to prevent SARS-CoV-2 entry into the host cell aim at blocking the

ACE2-RBD interaction. High-affinity monoclonal antibodies, many identified from convalescent

patients, are leading the way as potential therapeutics [22-29]. While highly effective *in vitro*,

these agents are expensive to produce by mammalian cell expression and need to be

intravenously administered by healthcare professionals. Moreover, large doses are likely to be

required for prophylactic viral neutralization, as only a small fraction of systemically circulating

antibodies cross the epithelial cell layers that line the airways [30]. By contrast, single domain

antibodies (nanobodies) provide significant advantages in terms of production and deliverability.

They can be inexpensively produced at scale in bacteria *(E. coli)* or yeast *(P. pastoris)*.

Furthermore, their inherent stability enables aerosolized delivery directly to the nasal and lung

epithelia by self-administered inhalation [31].

Monomeric mNb6 is among the most potent single domain antibodies neutralizing SARS-CoV-2

discovered to date. Multimerization of single domain antibodies has been shown to improve

target affinity by avidity [31, 32]. In the case of Nb6 and mNb6, however, our design strategy

enabled a multimeric construct that simultaneously engages all three RBDs, yielding profound

gains in potency. Furthermore, because RBDs must be in the up-state to engage with ACE2,

conformational control of RBD accessibility can serve as an added neutralization mechanism.

Indeed, our Nb6-tri and mNb6-tri molecules were designed with this functionality in mind. Thus,

when mNb6-tri engages with Spike, it prevents ACE2 binding by both directly occluding the

binding site and by locking the RBDs into an inactive conformation. Although a multitude of

other monoclonal and single-domain antibodies against SARS-CoV-2 Spike have been

discovered to date, there are few if any molecules as potent and stable as mNb6-tri [32-42].

Resistance to aerosolization, in particular, offers unprecedented opportunity for patient-friendly

nasal and pulmonary administration.

Our discovery of Class II neutralizing nanobodies demonstrates the presence of previously

unexplored mechanisms of blocking Spike binding to ACE2. For one Class II nanobody, Nb3,

we identified a likely binding site in the Spike S_1 domain external to the RBDs. Previously

discovered neutralizing antibodies from convalescent patients bind an epitope in a similar region

of Spike [24, 26, 27]. Binding of Nb3 to this epitope may allosterically stabilize RBDs in the

down-state, thereby decreasing ACE2 binding. Pairing of Class I and Class II nanobodies in a

prophylactic or therapeutic cocktail could thus be a highly advantageous strategy for both potent

neutralization and prevention of escape variants. The combined stability, potency, and diverse

epitope engagement of our anti-Spike nanobodies therefore provide a unique potential

prophylactic and therapeutic strategy to limit the continued toll of the COVID-19 pandemic.

MAIN TEXT FIGURES

Figure 1

Figure 1. Discovery of two distinct classes of anti-Spike nanobodies. A, Selection strategy

for identification of anti-Spike nanobodies that disrupt Spike-ACE2 interactions using magnetic

bead selections (MACS) or fluorescence activated cell sorting (FACS). **B,** Flow cytometry of

yeast displaying Nb6 (a Class I nanobody) or Nb3 (a Class II nanobody). Nb6 binds Spike*-

Alexa 647 and receptor binding domain (RBD-Alexa 647). Nb6 binding to Spike* is completely

disrupted by an excess (1.4 μM) of ACE2-Fc. Nb3 binds Spike*, but not the RBD. Nb3 binding

to Spike* is partially decreased by ACE2-Fc. **C,** SPR of Nb6 and Nb3 binding to either Spike* or

RBD. Red traces are raw data and global kinetic fits are shown in black. Nb3 shows no binding

to RBD. **D,** SPR experiments with immobilized Spike* show that Class I and Class II nanobodies

can bind Spike* simultaneously. By contrast, two Class I nanobodies or Class II nanobodies do

not bind simultaneously. **E,** Nanobody inhibition of 1 nM Spike*-Alexa 647 binding to ACE2

expressing HEK293T cells. n = 3 (ACE2, Nb3) or 5 (Nb6, Nb11) biological replicates. All error

bars represent s.e.m.

Figure 2

Figure 2. Cryo-EM structures of Nb6 and Nb11 bound to Spike. A, Cryo-EM maps of Spike*-
Nb6 complex in either closed (left) or open (right) Spike* conformation. **B,** Cryo-EM maps of
Spike*-Nb11 complex in either closed (left) or open (right) Spike* conformation. The top views
show receptor binding domain (RBD) up- or down-states. **C,** Nb6 straddles the interface of two
down-state RBDs, with CDR3 reaching over to an adjacent RBD. **D,** Nb11 binds a single RBD in
the down-state (displayed) or similarly in the up-state. No cross-RBD contacts are made by
Nb11 in either RBD up- or down-state. **E,** Comparison of RBD epitopes engaged by ACE2
(purple), Nb6 (red), or Nb11 (green). Both Nb11 and Nb6 directly compete with ACE2 binding.

Figure 3

Figure 3. Multivalency improves nanobody affinity and inhibitory efficacy. A, SPR of Nb6 and multivalent variants. Red traces show raw data and black lines show global kinetic fit for Nb6 and independent fits for association and dissociation phases for Nb6-bi and Nb6-tri. **B,** Dissociation phase SPR traces for Nb6-tri after variable association time ranging from 4 to 520 s. Curves were normalized to maximal signal at the beginning of the dissociation phase. Percent fast phase is plotted as a function of association time (right) with a single exponential fit. n = 3 independent biological replicates. **C,** Inhibition of pseudotyped lentivirus infection of ACE2 expressing HEK293T cells. n = 3 biological replicates for all but Nb11-tri (n = 2) **D,** Inhibition of live SARS-CoV-2 virus. Representative biological replicate with n = 3 (right panel) or 4 (left panel) technical replicates per concentration. n = 3 biological replicates for all but Nb3 and Nb3-tri (n = 2). All error bars represent s.e.m.

Figure 4

Figure 4. Affinity maturation of Nb6 yields a picomolar SARS-CoV-2 neutralizing

molecule. A, A saturation mutagenesis library of Nb6 was subjected to two rounds of selection

to identify consensus mutations I27Y and P105Y. **B,** SPR of mNb6 and mNb6-tri binding to

immobilized Spike*. Red traces show raw data and black lines show global kinetic fit. No

dissociation was observed for mNb6-tri over 10 minutes. **C,** Cryo-EM structure of Spike*-mNb6

complex. **D,** Comparison of receptor binding domain (RBD) engagement by Nb6 and mNb6.

One RBD was used to align both structures (RBD align), demonstrating changes in Nb6 and

mNb6 position and the adjacent RBD. **E,** Comparison of mNb6 complementarity determining

regions in either the cryo-EM structure of the Spike*-mNb6 complex or an X-ray crystal structure

of mNb6 alone. **F,** CDR1 of Nb6 and mNb6 binding to the RBD. As compared to I27 in Nb6, Y27

of mNb6 hydrogen bonds to Y453 and optimizes pi-pi and pi-cation interactions with the RBD.

G, CDR3 of Nb6 and mNb6 binding to the RBD demonstrating a large conformational

rearrangement of the entire loop in mNb6. **H,** Comparison of closed Spike* bound to mNb6 and

Nb6. Rotational axis for RBD movement is highlighted. **I,** Inhibition of pseudotyped lentivirus

infection of ACE2 expressing HEK293T cells by mNb6 and mNb6-tri. n = 3 biological replicates

J, mNb6 and mNb6-tri inhibit SARS-CoV-2 infection of VeroE6 cells in a plaque assay.

Representative biological replicate with n = 4 technical replicates per concentration. n = 3

biological replicates for all samples. All error bars represent s.e.m.

Figure 5

Figure 5. Nb6 and its derivates are robust proteins. A, Thermal denaturation of nanobodies assessed by circular dichroism measurement of molar ellipticity at 204 nm. Apparent melting temperatures (T_m) for each nanobody are indicated. **B,** Nanobody inhibition of 1 nM Spike*-Alexa 647 binding to ACE2 expressing HEK293T cells after incubation at either 25 °C or 50 °C for 1 hour or after aerosolization. **C,** Size exclusion chromatography of nanobodies after lyophilization or aerosolization. **D,** Summary table of SPR kinetics data and affinities for aerosolized or lyophilized mNb6 and mNb6-tri.

396 **Table 1. Anti-Spike nanobody affinity and neutralization potency**

Nanobody	Class	Spike* Binding			RBD Binding			Spike* Competition IC$_{50}$ (s.e.m) (M)[a]	SARS-CoV-2 Pseudovirus IC$_{50}$ (s.e.m.) (M)[b]	Live SARS-CoV-2 IC$_{50}$ (s.e.m.) (M)[c]
		k_a (M^{-1}s^{-1})	k_d (s^{-1})	K$_D$ (M)	k_a (M^{-1}s^{-1})	k_d (s^{-1})	K$_D$ (M)			
Nb2	I	9.0×10^5	5.3×10^{-1}	5.9×10^{-7}	1.0×10^6	9.9×10^{-1}	9.7×10^{-7}	8.3×10^{-6} (1.7×10^{-6})	NP	NP
Nb3	II	1.8×10^6	1.1×10^{-1}	6.1×10^{-8}	NB			NC	3.9×10^{-6} (7.9×10^{-7})	3.0×10^{-6} (3.2×10^{-7})
Nb6	I	2.7×10^5	5.6×10^{-2}	2.1×10^{-7}	2.1×10^6	8.7×10^{-2}	4.1×10^{-8}	3.7×10^{-7} (4.9×10^{-8})	2.0×10^{-6} (3.5×10^{-7})	3.3×10^{-6} (7.2×10^{-7})
Nb8	I	1.4×10^5	8.1×10^{-1}	5.8×10^{-6}	6.6×10^5	3.3×10^{-1}	5.1×10^{-7}	4.8×10^{-6} (4.9×10^{-7})	NP	NP
Nb11	I	1.2×10^6	1.6×10^{-1}	1.4×10^{-7}	3.2×10^6	2.4×10^{-1}	7.6×10^{-8}	5.4×10^{-7} (1.2×10^{-7})	2.4×10^{-6} (5.4×10^{-7})	NP
Nb12	I	1.2×10^2	2.0×10^{-4}	1.6×10^{-6}	Biphasic	Biphasic	Biphasic	2.5×10^{-7} (5.5×10^{-8})	1.2×10^{-6} (9.0×10^{-7})	NP
Nb15	I	1.7×10^5	2.3×10^{-1}	1.3×10^{-6}	6.0×10^5	2.2×10^{-1}	3.6×10^{-7}	2.2×10^{-6} (2.5×10^{-7})	6.7×10^{-6} (3.6×10^{-6})	NP
Nb16	I	1.1×10^5	1.3×10^{-1}	1.3×10^{-6}	NP			9.5×10^{-7} (1.1×10^{-7})	NP	NP
Nb17	II	7.3×10^5	2.0×10^{-1}	2.7×10^{-7}	NB			NC	7.6×10^{-6} (1.0×10^{-6})	NP
Nb18	II	1.4×10^5	6.4×10^{-3}	4.5×10^{-8}	NB			5.2×10^{-5} (1.5×10^{-5})	NP	NP
Nb19	I	2.4×10^4	1.1×10^{-1}	4.5×10^{-6}	1.0×10^5	8.9×10^{-2}	8.8×10^{-7}	4.1×10^{-6} (4.9×10^{-7})	2.4×10^{-5} (7.7×10^{-6})	NP
Nb24	I	9.3×10^5	2.7×10^{-1}	2.9×10^{-7}	2.4×10^6	3.5×10^{-1}	1.5×10^{-7}	7.5×10^{-7} (1.0×10^{-7})	NP	NP
ACE2	N/A	2.7×10^5	1.2×10^{-2}	4.4×10^{-8}	NP	NP	NP	1.7×10^{-7} (6.6×10^{-8})	6.2×10^{-7} (1.7×10^{-7})	NP
mNb6	I	1.0×10^6	4.5×10^{-4}	4.5×10^{-10}	1.1×10^6	6.4×10^{-4}	5.6×10^{-10}	1.3×10^{-9} (4.1×10^{-10})	6.3×10^{-9} (1.6×10^{-9})	1.2×10^{-8} (2.5×10^{-9})
Nb3-bi	II	NP	NP	NP	NP	NP	NP	NP	3.6×10^{-7} (1.5×10^{-7})	1.8×10^{-7} (1.2×10^{-8})
Nb3-tri	II	Biphasic	Biphasic	Biphasic	NP	NP	NP	4.1×10^{-8} (1.6×10^{-8})	4.0×10^{-7} (1.6×10^{-7})	1.4×10^{-7} (4.9×10^{-8})
Nb6-bi	I	Biphasic	Biphasic	Biphasic	NP	NP	NP	NP	6.3×10^{-8} (1.5×10^{-8})	NP
Nb6-tri	I	Biphasic	Biphasic	Biphasic	NP	NP	NP	1.5×10^{-9} (5.2×10^{-10})	1.2×10^{-9} (2.5×10^{-10})	1.6×10^{-10} (2.6×10^{-11})
Nb11-tri	I	Biphasic	Biphasic	Biphasic	NP	NP	NP	NP	5.1×10^{-8} (1.6×10^{-8})	NP
ACE2-Fc	N/A	NP	NP	NP	NP	NP	NP	5.3×10^{-9} (2.5×10^{-9})	4.0×10^{-8} (8.8×10^{-9})	2.6×10^{-8} (8.5×10^{-9})
mNb6-tri	I	1.4×10^6	$<1.0\times10^{-6}$	$<1.0\times10^{-12}$	NP	NP	NP	4.0×10^{-10} (1.4×10^{-10})	1.2×10^{-10} (2.8×10^{-11})	5.4×10^{-11} (1.0×10^{-11})

397

398 [a]Average values from n = 5 biological replicates for Nb6, Nb11, Nb15, Nb19 are presented, all

399 others were tested with n = 3 biological replicates.

[b]Average values from n = 2 biological replicates for Nb12, Nb17, and Nb11-tri are presented, all

others were tested with n = 3 biological replicates.

[c]Average values from n = 2 biological replicates for Nb3, Nb3-bi, and Nb3-tri. n = 3 biological

replicates for all others.

NB – no binding

NC – no competition

NP – not performed

Supplementary Figures

Supplementary Fig. 1. Validation of purified Spike*. A, Size exclusion chromatogram of
purified Spike* from ExpiCHO cells. **B,** SPR of immobilized Spike* binding to monomeric ACE2
extracellular domain (ECD).

Supplementary Fig. 2. Cryo-EM workflow for Nb6

A flowchart representation of the classification workflow for Spike*-Nb6 complexes yielding open and closed Spike* conformations. From top to bottom, particles were template picked with a set of 20 Å low-pass filtered 2D backprojections of apo-Spike* in the closed conformation. Extracted particles in 2D classes suggestive of various Spike* views were subject to a round of heterogenous refinement in cryoSPARC with two naïve classes generated from a truncated *Ab initio* job, and a 20 Å low-pass filtered volume of apo-Spike* in the closed conformation. Particles in the Spike* 3D class were subject to 25 iterations of 3D classification into 6 classes without alignment in RELION, using the same input volume from cryoSPARC 3D classification,

425 low pass filtered to 60 Å, T = 8. Particles in classes representing the open and closed Spike*

426 conformations were imported into cisTEM for automatic refinement. Half maps from refinement

427 were imported into cryoSPARC for local resolution estimation as shown in Supplementary Fig.

428 4.

Supplementary Fig. 3. Cryo-EM workflow for Nb11

A flowchart representation of the classification workflow for Spike*-Nb11 complexes yielding

open and closed Spike* conformations. From top to bottom, particles were template picked from

two separate collections with a set of 20 Å low-pass filtered 2D backprojections of apo-Spike* in

the closed conformation. Extracted particles were Fourier cropped to 128 pixels prior to

extensive heterogenous refinement in cryoSPARC, using a 20 Å low-pass filtered volume of

apo-Spike* in the closed conformation and additional naïve classes for removal of non-Spike*

particles. After cryoSPARC micrograph curation and heterogenous refinement, Spike* density

corresponding to all regions outside of the ACE2 RBD::Nanobody interface were subtracted. A

439   mask around the ACE2 RBD::Nanobody interface was generated, and used for multiple rounds

440   of 3D classification without alignment in RELION. Particles in classes representing open and

441   closed Spike* conformations were selected, unsubtracted and unbinned prior to refinement in

442   RELION. Half maps from refinement were imported into cryoSPARC for local resolution

443   estimation as shown in Supplementary Fig. 4.

Supplementary Fig. 4. Local resolution of cryo-EM maps

Local resolution estimates of Spike* complexes with A-B) Nb6, C-D) Nb11, and E) mNb6 as
generated in cryoSPARC. All maps (except mNb6) are shown with the same enclosed volume.
All maps are colored on the same scale, as indicated.

449
Supplementary Fig. 5. Radiolytic hydroxyl radical footprinting of Spike*.

A, Change in oxidation rate between Spike* and Nb3-Spike* complexes at all residues. A

cluster of highly protected residues in the Spike*-Nb3 complex is observed in the N-terminal

domain. **B,** Oxidation rate plots of the two (M177, H207) most heavily protected residues upon

Nb3 binding to Spike*. Data points labeled with an asterisk are excluded from rate calculations

as these values fall outside of the first order reaction, likely due to extensive oxidation-mediated

damage. **C,** Change in oxidation rate mapped onto Spike in the all RBD down conformation.

Supplementary Fig. 6. Modeling of distances for multimeric nanobody design. A, Model of

Spike*:Nb6 complex in the closed state. The minimal distance between adjacent Nb6 N- and C-

termini is 52 Å (dashed line). **B,** Model of Spike*:Nb6 complex in the open state with Nb6

docked into the cryo-EM density for up-state RBD. Minimal distance between N- and C-termini

of both nanobodies is 72 Å**.** Nb6 cannot bind RBD2 in open Spike*, as this would sterically clash

with RBD3. **C,** Model of Spike*:Nb11 complex in the closed state. The minimal distance

between adjacent Nb6 N- and C-termini is 71 Å (dashed line). **D,** Model of Spike*:Nb11 complex

in the open state. The minimal distance between adjacent Nb6 N- and C-termini is 68 Å

between Nb11 bound to RBD2 in the down-state and RBD3 in the up-state. For B, the model of

Nb6 from A was docked into the cryo-EM map to enable modeling of distance between N- and

C-termini. For C and D, a generic nanobody was docked into cryo-EM maps to model the

distance between N- and C-termini.

Supplementary Fig. 7. CryoEM workflow for mNb6

A flowchart representation of the classification workflow for the Spike*-mNb6 complex yielding a closed Spike* conformation. From top to bottom, particles were template picked from two separate collections with a set of 20Å low-pass filtered 2D backprojections of apo-Spike* in the

closed conformation. Extracted particles were Fourier cropped to 96 pixels prior to 2D

classification. Particles in Spike* 2D classes were selected for a round of heterogeneous

refinement in cryoSPARC using a 20 Å low-pass filtered volume of apo-Spike* in the closed

conformation and additional naïve classes for removal of non-Spike* particles. In RELION,

particles in the Spike* 3D class were subject to two rounds of 3D classification without

alignment into 6 classes using the same input volume from cryoSPARC 3D classification, low

pass filtered to 60 Å, T = 8. Unbinned particles in the Spike*-closed conformation were exported

into cisTEM for automatic refinement, followed by local refinement using a mask around the

ACE2 RBD::Nanobody interface. Half maps from refinement were imported into cryoSPARC for

local resolution estimation as shown in Supplementary Fig. 4.

489 Supplementary Table 1. CryoEM datasets

Sample:	Spike*-Nb6		Spike*-Nb11		Spike*-mNb6
Spike* conformation:	Open	Closed	Open	Closed	Closed
EMDB:	XXXX	XXXX	XXXX	XXXX	XXXX
PDB:		XXXX			XXXX
Data collection and processing					
Microscope/Detector	Titan Krios/Gatan K3 with Gatan Bioquantum Energy Filter				
Imaging software and collection	SerialEM, 3x3 image shift				
Magnification	105,000				
Voltage (kV)	300				
Electron exposure (e–/Å^2)	66				
Dose rate (e–/pix/sec)	8				
Frame exposure (e–/Å^2)	0.55				
Defocus range (μm)	-0.8 to -2.0				
Pixel size (Å)	0.834 (physical)				
Micrographs	5,317		4,103		1,609
Reconstruction					
Autopicked particles	2,033,067		1,204,855		585,250
(template-based in cryosparc)					
Particles in final refinement	40,125	58,493	21,570	27,611	53,690
	(cisTEM)	(cisTEM)	(cisTEM)	(RELION)	(cisTEM)
Symmetry imposed	C1	C3	C1	C1	C3
Map sharpening B factor (Å^2)		-90			-140
Map resolution, global FSC (Å)					
FSC 0.5, unmasked/masked	7.8/4.6	4.1/3.4	7.0/4.4	7.6/5.3	3.9/3.3
FSC 0.143, unmasked/masked	4.7/3.8	3.5/3.0	4.3/3.7	5.1/4.2	3.2/2.9
Refinement					
Initial model used (PDB code)		6VXX, 3P0G			6VXX, 3P0G
Model resolution (Å)					
FSC 0.5, unmasked/masked		3.5/3.1			3.2/2.9
Model composition					
Non-hydrogen atoms		26904			27015
Protein residues		3360			3360
B factors (Å^2)					
Protein		97.0			57.5
Ligand		107.4			85.7
R.m.s. deviations					
Bond lengths (Å)		0.014			0.007
Bond angles (°)		1.379			1.027
Validation					
MolProbity score		1.99			1.71
Clashscore		12.70			6.46
Poor rotamers (%)		0.45			0.41
EMRinger score		2.98			4.01
CaBLAM score		3.11			2.95
Ramachandran plot					
Favored (%)		94.49			94.92
Allowed (%)		5.51			5.08
Disallowed (%)		0			0

490
491

492 **Supplementary Table 2. X-ray data collection and refinement statistics**

	mNb6 (PDB XXXX)
Data collection	
Space group	$P2_1$
Cell dimensions	
a, b, c (Å)	44.56, 71.25, 46.43
α, β, γ (°)	90.0, 114.93, 90.0
Molecules in asymmetric unit	2
Resolution (Å)	71.25 - 2.05 (2.09 - 2.05)[a]
R_{sym} or R_{merge}	0.13 (0.94)[b]
$I / \sigma I$	7.2 (0.9)
Completeness (%)	97.8 (96.6)
Redundancy	6.4 (5.7)
CC (1/2) (%)	99.8 (64.4)
Refinement	
Resolution (Å)	71.25 – 2.05
No. reflections	104195
R_{work} / R_{free} (%)	21.16 / 24.75
No. atoms	
Protein	1798
Ligand/ion	21
Water	131
B-factors	
Protein	33.1
Ligand/ion	76.1
Water	42.2
R.m.s. deviations	
Bond lengths (Å)	0.07
Bond angles (°)	0.826
Ramachandran plot	
Allowed (%)	99.06
Generous (%)	0.94
Disallowed (%)	0

493 [a] Values in parentheses correspond to the highest resolution shell.

494 [b] $R_{merge} = \Sigma|I - <I>|/\Sigma I$

495 [c] $R_{work} = \Sigma|F_O - F_C|/\Sigma F_O$

496 [d] $R_{free} = \Sigma|F_O - F_C|/\Sigma F_O$, calculated using a random set containing 5% reflections that were not included throughout

497 structure refinement.

498 **Supplementary Table 3. Nanobody expression plasmids**
499

Plasmid	Nanobody	Plasmid backbone	Resistance Marker
pPW3544	Nb2	pet-26b(+)	kanamycin
pPW3545	Nb3	pet-26b(+)	kanamycin
pPW3546	Nb6	pet-26b(+)	kanamycin
pPW3547	Nb8	pet-26b(+)	kanamycin
pPW3548	Nb11	pet-26b(+)	kanamycin
pPW3549	Nb12	pet-26b(+)	kanamycin
pPW3550	Nb15	pet-26b(+)	kanamycin
pPW3551	Nb16	pet-26b(+)	kanamycin
pPW3552	Nb17	pet-26b(+)	kanamycin
pPW3553	Nb18	pet-26b(+)	kanamycin
pPW3554	Nb19	pet-26b(+)	kanamycin
pPW3555	Nb24	pet-26b(+)	kanamycin
pPW3557	Trivalent Nb6, 20AA length GS linker	pet-26b(+)	kanamycin
pPW3558	Trivalent Nb3, 15AA length GS linker	pet-26b(+)	kanamycin
pPW3559	Trivalent Nb11, 15AA length GS linker	pet-26b(+)	kanamycin
pPW3560	Bivalent Nb3, 15AA length GS linker	pet-26b(+)	kanamycin
pPW3561	Bivalent Nb6, 15AA length GS linker	pet-26b(+)	kanamycin
pPW3563	Trivalent mNb6, 20AA length GS linker	pet-26b(+)	kanamycin
pPW3564	mNb6	pet-26b(+)	kanamycin

500
501
502

METHODS

Expression and purification of SARS-CoV-2 Spike, RBD, and ACE2.

We used a previously described construct to express and purify the pre-fusion SARS-CoV-2

Spike ectodomain (Spike*) [15]. ExpiCHO or Expi293T cells (ThermoFisher) were transfected

with the Spike* construct per the manufacturer's instructions for the MaxTiter protocol and

harvested between 3-9 days after transfection. Clarified cell culture supernatant was loaded

onto Ni-Excel beads (Cytiva) followed by extensive washes in 20 mM HEPES pH 8.0, 200 mM

sodium chloride, and 10 mM imidazole and elution in the same buffer supplemented with 500

mM imidazole. Spike* was concentrated using a 100 kDa MWCO spin concentrator (Millipore)

and further purified by size exclusion chromatography over a Superose 6 Increase 10/300

column (GE Healthcare) in 20 mM HEPES pH 8.0 and 200 mM sodium chloride. All purification

steps were performed at room temperature. The resulting fractions for trimeric Spike* were

pooled and either used directly for cryo-EM studies or concentrated and flash frozen in liquid

nitrogen with 15% glycerol for other biochemical studies.

We used a previously described construct to express and purify the SARS-CoV-2 Receptor

binding domain (RBD) [43]. Expi293T cells (ThermoFisher) were transfected with the RBD

construct per the manufacturer's instructions and harvested between 3-6 days after transfection.

Clarified cell culture supernatant was loaded onto Ni-Excel beads (Cytiva) or a His-Trap Excel

column (GE Healthcare) followed by washes in 20 mM HEPES pH 8.0, 200 mM sodium

chloride, and 10 mM imidazole and elution using the same buffer supplemented with 500 mM

imidazole. RBD was concentrated using a 30 kDa MWCO spin concentrator (Millipore) and

further purified by size exclusion chromatography over a Superdex 200 Increase 10/300 GL

column (GE Healthcare) in 20 mM HEPES pH 8.0 and 200 mM sodium chloride. The resulting

fractions were pooled, concentrated, and flash frozen in liquid nitrogen with 10% glycerol.

For biochemical and yeast display experiments, Spike* and RBD were labeled with freshly

prepared stocks of Alexa 647-NHS, Alexa 488-NHS, or Biotin-NHS (ThermoFisher) with a 5-fold

stoichiometry for 1 hour at room temperature followed by quenching of NHS with 10 mM Tris pH

8.0 for 60 minutes. Labeled proteins were further purified by size exclusion chromatography,

concentrated using a spin concentrator (Millipore), and flash frozen in liquid nitrogen with 10-

15% glycerol.

We used an ACE2-ECD (18-614) Fc fusion expression plasmid to express and purify Fc tagged

ACE2-ECD [44]. Expi293T cells (ThermoFisher) were transfected with the ACE2-Fc construct

per the manufacturer's instructions and harvested between 5-7 days after transfection. Clarified

cell culture supernatant was loaded onto a MabSelect Pure 1 mL Column (GE Healthcare).

Column was washed with Buffer A (20 mM HEPES pH 7.5, 150 mM NaCl) and protein was

eluted with Buffer B (100 mM Sodium Citrate pH 3.0, 150 mM NaCl) into a deep well block

containing 1 M HEPES pH 7.5 to neutralize the acidic elution. ACE2-Fc was concentrated using

a 30 kDa MWCO spin concentrator (Millipore) and further purified by size exclusion

chromatography over a Superdex 200 Increase 10/300 GL column (GE Healthcare) in SEC

Buffer (20 mM HEPES pH 7.5, 150 mM NaCl, 5% v/v Glycerol). The resulting fractions were

pooled, concentrated, and flash frozen in liquid nitrogen. To obtain monomeric ACE2, 1:50

(w/w) His-tagged TEV protease was added to ACE2-Fc and incubated at 4 °C overnight. This

mixture was then purified by size exclusion chromatography in SEC Buffer. Monomeric ACE2

fractions were pooled and washed with His-resin (1 mL of 50% slurry) to remove excess TEV.

The resulting supernatant was pooled, concentrated, and flash frozen in liquid nitrogen.

Identification of anti SARS-CoV2 Spike nanobodies

To identify nanobodies against the SARS-CoV-2 Spike ECD, we used a yeast surface displayed

library of synthetic nanobody sequences that recapitulate amino acid position specific-variation

in natural llama immunological repertoires. This library encodes a diversity of >$2x10^9$ variants,

and uses a synthetic stalk sequence for nanobody display, as described previously in a modified

vector encoding nourseothricin (NTC) resistance [45]. For the first round of selection, $2x10^{10}$

yeast induced in YPG (Yeast Extract-Peptone-Galactose) supplemented with NTC were washed

repeatedly in selection buffer (20 mM HEPES, pH 7.5, 150 mM sodium chloride, 0.1% (w/v) low

biotin bovine serum albumin, BSA) and finally resuspended in 10 mL of selection buffer

containing 200 nM biotinylated-Spike*. Yeast were incubated for 30 minutes at 25 °C, then

washed repeatedly in cold selection buffer, and finally resuspended in 10 mL of cold selection

buffer containing 200 µL of Miltenyi anti-Streptavidin microbeads. After 30 minutes of incubation

at 4 °C, yeast were again washed with cold selection buffer. Spike* binding yeast were captured

on a Miltenyi MACS LS column and recovered in YPD (Yeast Extract-Peptone-Dextrose)

medium supplemented with NTC.

For round 2, $4x10^8$ induced yeast from Round 1 were incubated with 100 nM Spike* labeled with

Alexa647 in 1 mL of selection buffer for 1 hr at 25 °C. After extensive washes with cold selection

buffer, Spike* binding yeast were isolated by fluorescence activated cell sorting (FACS) on a

Sony SH800 instrument. A similar approach was used for round 3, with substitution of 10 nM

Spike* labeled with Alexa647. Post round 3 yeast were plated on YPD+NTC solid media and

768 individual colonies were induced with YPG+NTC media in 2 mL deep well plates. Each

individual clone was tested for binding to 4 nM Spike*-Alexa488 by flow cytometry on a

Beckman Coulter Cytoflex. To identify nanobodies that disrupt Spike-ACE2 interactions, Spike*

binding was repeated in the presence of 0.5-1 µM ACE2-Fc. Out of 768 clones, we identified 21

that strongly bind Spike* and are competitive with ACE2 (Supplementary Table 3).

Expression and purification of nanobodies

Nanobody sequences were cloned into the pET26-b(+) expression vector using In-Fusion HD

cloning (Takara Bio), transformed into BL21(DE3) *E. coli*, grown in Terrific Broth at 37 °C until

OD 0.7-0.8, followed by gene induction using 1 mM IPTG for 18-22 hours at 25°C. *E. Coli* were

harvested and resuspended in SET Buffer (200 mM Tris, pH 8.0, 500 mM sucrose, 0.5 mM

EDTA, 1X cOmplete protease inhibitor (Roche)) for 30 minutes at 25 °C before a 45 minute

osmotic shock with a two-fold volume addition of water. NaCl, MgCl2, and imidazole were

added to the lysate to 150 mM, 2 mM, and 40 mM respectively before centrifugation at 17-

20,000xg for 15 minutes to separate cell debris from the periplasmic fraction. For every liter of

bacterial culture, the periplasmic fraction was then incubated with 4 mL of 50% HisPur Ni-NTA

resin (Thermo Scientific) which had been equilibrated in Nickel Wash Buffer (20 mM HEPES,

pH 7.5, 150 mM NaCl, 40 mM imidazole). This mixture was incubated for 1 hr with rotation at

RT before centrifugation at 50xg to collect the resin. The resin was then washed with 5 volumes

of Nickel Wash buffer 3 times, each time using centrifugation to remove excess wash buffer.

Bound proteins were then eluted using three washes with Elution Buffer (20 mM HEPES, pH

7.5, 150 mM NaCl, 500 mM imidazole). The eluted protein was concentrated using a 3.5 kDa

MWCO centrifugal filter unit (Amicon) before injection onto a Superdex 200 Increase 10/300 GL

column equilibrated with 20 mM HEPES, pH 7.5, 150 mM NaCl. Nanobody constructs were

concentrated again using a 3.5k MWCO centrifugal filter unit, and flash frozen in liquid nitrogen.

Affinity determination by surface plasmon resonance

Nanobody (Nb) affinity determination experiments were performed on Biacore T200 and 8K

instruments (Cytiva Life Sciences) by capturing the StreptagII-tagged Spike* at 10 µg/mL on a

StreptactinXT-immobilized (Iba Life Sciences) CM5 Series S sensor chip (Cytiva Life Sciences)

to achieve maximum response (Rmax) of approximately 30 response units (RUs) upon

nanobody binding. 2-fold serial dilutions of purified nanobody from 1 µM to 31.25 nM (for

monovalent constructs) or from 50 nM to 1.56 nM (for affinity matured and multimeric

constructs) were flowed over the captured Spike* surface at 30 µL/minute for 60 seconds

followed by 600 seconds of dissociation flow. Following each cycle, the chip surface was

regenerated with 3 M guanidine hydrochloride.

Separately, biotinylated SARS-CoV-2 RBD at 8 µg/mL was loaded onto a preconditioned Series

S Sensor Chip CAP chip (Cytiva Life Sciences) to achieve an Rmax of approximately 60 RUs

upon nanobody binding. 2-fold serial dilutions in the same running buffer and sample series

(parent or affinity matured clone) as the Spike* runs were flowed over the RBD surface at 30

µL/minute for 60 seconds followed by 600 seconds of dissociation flow. Chip surface

regeneration was performed with a guanidine hydrochloride/sodium hydroxide solution.

The resulting sensorgrams for all monovalent clones were fit to a 1:1 Langmuir binding model

using the Biacore Insight Evaluation Software (Cytiva Life Sciences) or the

association/dissociation model in GraphPad Prism 8.0. For determination of kinetic parameters

for Nb6-bi and Nb6-tri binding, the dissociation phase was fit to a biexponential decay

constrained to two dissociation rate constants shared between each concentration. The

association phase was fit separately using an association kinetics model simultaneously fitting

the association rate constant for each concentration.

For nanobody competition experiments, Spike* was loaded onto a StreptactinXT-immobilized

CM5 sensor chip as previously described. As in the kinetics experiments, the primary nanobody

was flowed over the captured Spike* surface for 60 seconds at 30 µL/minute to achieve

saturation. Immediately following this, a second injection of a mixture of primary and variable

nanobody at the same concentration as in the primary injection was performed.

ACE2 cellular surface binding competition assays

A dilution series of nanobody was generated in PBE (PBS + 0.5% (w/v) BSA + 2 mM EDTA and

mixed with Spike*-Alexa647 or RBD-Alexa647. ACE2 expressing HEK293T cells were

dissociated with TrypLE Express (ThermoFisher) and resuspended in PBE [20]. The cells were

mixed with the Spike*-nanobody solution and incubated for 45 minutes, washed in PBE, and

then resuspended in PBE. Cell surface Alexa647 fluorescence intensity was assessed on an

Attune Flow Cytometer (ThermoFisher).

Affinity maturation of Nb6

A site saturation mutagenesis library of Nb6 was generated by assembly PCR of overlapping

oligonucleotides encoding the Nb6 sequence. Individual oligos for each position in CDR1,

CDR2, and CDR3 were designed with the degenerate "NNK" codon. The assembled gene

product was amplified with oligonucleotides with overlapping ends to enable homologous

recombination with the yeast surface display vector as previously described and purified with

standard silica-based chromatography [45]. The resulting insert DNA was transformed into

Saccharomyces cerevisiae strain BJ5465 along with the yeast display vector pYDS2.0 to

generate a library of $2x10^8$ transformants. After induction in YPD+NTC medium at 20 °C for 2

days, $2x10^9$ yeast were washed in selection buffer (20 mM HEPES, pH 8.0, 150 mM sodium

chloride, 0.1% (w/v) low biotin BSA) and incubated with 1 nM biotin-Spike* for 1 hour at 25 °C.

Yeast were subsequently washed in selection buffer, resuspended in 1 mL selection buffer, and

incubated with 10 µL streptavidin microbeads (Miltenyi) for 15 min. at 4 °C. Yeast were washed

again with cold selection buffer and Spike*-binding yeast were isolated by magnetic separation

using an LS column (Miltenyi). Recovered yeast were grown in YPD+NTC at 37 °C and induced

in YPG+NTC at 20 °C. A second round of selection was performed as above, substituting 100

pM RBD-Alexa647 as the antigen. Yeast displaying high affinity clones were selected by

magnetic separation using Anti-Cy5 microbeads (Miltenyi) and an LS column. Analysis of the

library after the second round of selection revealed a population of clones with clear binding of

10 pM RBD-Alexa647. Therefore, 96 individual clones were screened for binding to 10 pM RBD-

Alexa647 by flow cytometry. Sequence analysis of eight clones that showed robust binding to

10 pM RBD-Alexa647 revealed two consensus mutations, I27Y and P105Y, which were used to

generate the affinity matured clone mNb6.

Structures of Spike-nanobody complexes by cryo-EM

Sample preparation and microscopy

To prepare Spike*-nanobody complexes, each nanobody was incubated on ice at a 3-fold molar

excess to Spike* at 2.5 µM for 10 minutes. 3 µL of Spike*-nanobody complex was added to a

300 mesh 1.2/1.3R Au Quantifoil grid previously glow discharged at 15 mA for 30 seconds.

Blotting was performed with a blot force of 0 for 4 seconds at 4°C and 100% humidity in a FEI

Vitrobot Mark IV (ThermoFisher) prior to plunge freezing into liquid ethane.

For each complex, 120-frame super-resolution movies were collected with a 3x3 image shift

collection strategy at a nominal magnification of 105,000x (physical pixel size: 0.834 Å/pix) on a

Titan Krios (ThermoFisher) equipped with a K3 camera and a Bioquantum energy filter (Gatan)

set to a slit width of 20 eV. Collection dose rate was 8 e$^-$/pixel/second for a total dose of 66 e$^-$

/Å^2. Each collection was performed with semi-automated scripts in SerialEM [46].

Image Processing

For all datasets, dose fractionated super-resolution movies were motion corrected with

MotionCor2 [47]. Contrast transfer function determination was performed with cryoSPARC patch

CTF [48]. Particles were picked with a 20 Å low-pass filtered apo Spike 2D templates generated

from a prior data collection.

Nb6-Spike* and mNb6-Spike* particles were extracted with a 384 pixel box, binned to 96 pixels

and subject to single rounds of 2D and 3D classification prior to unbinning for homogenous

refinement in cryoSPARC [48]. Refined particles were then imported into Relion3.1 for 3D

classification without alignment using the input refinement map low pass filtered to 40 Å [49].

Particles in classes representing the closed conformation of Spike were imported into cisTEM

and subject to autorefinement followed by local refinement within a RBD::nanobody masked

region [50]. Following local refinement, a new refinement package symmetrized to the C3 axis

was created for a final round of local refinement without masking. Final particle counts for each

map are as follows: Nb6-Open: 40,125, Nb6-Closed: 58,493, mNb6: 53,690.

Nb11-Spike* particles were extracted with a 512 pixel box, binned to 128 pixels for multiple

rounds of 3D classification as described in Figure S4. Following homogenous refinement,

particles were exported to Relion3.1. Particle density roughly corresponding to RBD-nanobody

complexes was retained post-particle subtraction. 3D classification without alignment was

performed on the particle subtracted stacks. Particles in classes with robust RBD-nanobody

density were selected, unsubtracted and refined in Relion followed by post-processing. 21,570

particles contributed to the final maps. Final particle counts for each map are as follows: Nb11-

Open: 21,570, Nb11-Closed: 27,611. For all maps, final local resolution estimation and GSFSC

determination was carried out in cryoSPARC.

Structure modeling

Models of Nb6-Spike* and mNb6-Spike* were built using a previously determined structure of

closed Spike* (PDB: 6VXX) [14]. A composite model incorporating resolved regions of the RBD

was made using a previously determined X-ray crystal structure of the SARS-CoV-2 RBD (PDB:

6M0J) [51]. For Nb6, the beta2-adrenergic receptor nanobody Nb80 (PDB: 3P0G) was used as

a template to first fit the nanobody into the cryo-EM density map for the Nb6-Spike* complex

[52]. Complementarity determining loops were then truncated and rebuilt using RosettaES [53].

The final structure was inspected and manually adjusted in COOT and ISOLDE, followed by real

space refinement in PHENIX [54-56]. The higher resolution structure of mNb6 enabled manual

building of nanobody CDR loops *de novo*, and therefore the Rosetta-based approach was not

used for modeling. Final models were analyzed in PHENIX, with statistics reported in

Supplementary Table 1.

For models of Nb11-Spike* complexes presented here, the closest nanobody by sequence in

the PDB (beta2-adrenergic receptor Nb60, PDB ID: 5JQH) was fit by rigid-body refinement in

COOT into the cryo-EM density map using only the framework regions [57]. While the lower

resolution of these maps precluded confident assignment of loop conformations, the overall

orientation of Nb11 relative to Spike* was well constrained, enabling accurate modeling of

distances between the N- and C- termini of two Nb11 molecules bound to Spike*.

Radiolytic hydroxyl radical footprinting and mass-spectrometry of Spike* and Nb3-Spike*

Spike* and Nb3 samples were buffer exchanged into 10 mM phosphate buffer (pH 7.4) by

extensive dialysis at 25 °C. A 1.5-fold molar excess of Nb3 was added to 5 μM Spike* and the

complex was incubated for >24 hr at 25 °C. For radiolytic footprinting, protein concentrations

and beam parameters were optimized using an Alexa-488 fluorophore assay [58]. Apo Spike*

and Spike*-Nb3 complex at concentrations of 1-3 μM were exposed to a synchrotron X-ray

white beam at 6 timepoints between 0-50 ms at beamline 3.2.1 at the Advanced Light Source in

Berkeley, CA and were quenched with 10 mM methionine amide immediately post-exposure.

Glycans were removed by treatment with 5% SDS, 5 mM DTT at 95 °C for five minutes and

subsequent PNGase (Promega) digestion at 37°C for 2 hours. Samples were buffer exchanged

into ammonium bicarbonate (ABC) buffer (pH 8.0) using ZebaSpin columns (Thermo Fisher).

Alkylation of cysteines was achieved by treatment with 8 M urea and 5 mM DTT at 37°C for 30

minutes followed by an incubation with 15 mM iodoacetamide at 25 °C in the dark for 30

minutes. All samples were further buffer exchanged to ABC pH 8.0 using ZebaSpin columns

and digested with either Trypsin/Lys-C or Glu-C (Promega) at an enzyme:protein ratio of 1:20

(w/w) at 37 °C for 8 hours.

Samples were lyophilized and resuspended in 1% formic acid at 200 fmol/µL concentration. For

each MS analysis, 1 µL of sample was injected onto a 5 mm Thermo Trap C18 cartridge, and

then separated over a 15 cm column packed with 1.9 µm Reprosil C18 particles (Dr. Maisch

HPLC GmbH) by a nanoElute HPLC (Bruker). Separation was performed at 50 °C and a flow

rate of 400 µL/min by the following gradient in 0.1% formic acid: 2% to 17% acetonitrile from 0

to 20 min, followed by 17% to 28% acetonitrile from 20 to 40 min. The eluent was electrospray

ionized into a Bruker timsTOF Pro mass spectrometer and data was collected using data-

dependent PASEF acquisition. Database searching and extraction of MS1 peptide abundances

was performed using the FragPipe platform with either trypsin or GluC enzyme specificity, and

all peptide and protein identifications were filtered to a 1% false-discovery rate [59]. Searches

were performed against a concatenated protein database of the Spike protein, common

contaminant proteins, and the *Saccharomyces cerevisiae* proteome (downloaded July 23,

2020). Note, the *Saccharomyces cerevisiae* proteome was included to generate a sufficient

population of true negative identifications for robust false discovery rate estimation of peptide

and protein identifications. Lastly, the area under the curve MS1 intensities reported from

FragPipe were summarized for each peptide species using MSstats [60].

The peak areas of extracted ion chromatograms and associated side-chain modifications were

used to quantify modification at each timepoint. Increasing beamline exposure time decreases

the fraction of unmodified peptide and can be represented as a site-specific dose-response plot

(Supplementary Fig. 5B). The rate of hydroxyl radical reactivity (k_{fp}) is dependent on both the

intrinsic reactivity of each residue and its solvent accessibility and was calculated by fitting the

dose-response to a pseudo-first order reaction scheme in Graphpad Prism Version 8. The ratio

of k_{fp} between apo Spike* and the Spike-Nb3 complex at specific residues gave information on

solvent accessibility changes between the two samples. These changes were mapped onto the

SARS-CoV-2 Spike (PDB 6XR8) [11]. In some cases, heavily modified residues show a

flattening of dose-response at long exposures which we interpret as radical induced damage.

These over-exposed timepoints were excluded from the calculation of k_{fp}.

mNb6 crystallography and structure determination

Purified mNb6 was concentrated to 18.7 mg/mL and filtered using 0.1 µm hydrophilic PVDF

filters (Millipore). mNb6 crystal screens were set up in 96 well plates in hanging drop format at

2:1 protein:reservoir in Index and AmSO4 screens (Hampton Research, Aliso Viejo, CA).

Crystals in over 60 different screening conditions with various morphologies appeared overnight

at ambient temperature and were obtained directly from the screens without further optimization.

The crystals were cryoprotected by quick dipping in a solution containing 80% reservoir and

20% PEG400 or 20% Glycerol, then mounted in CrystalCap HT Cryoloops (Hampton Research,

Aliso Viejo, CA) and flash cooled in a cryogenic nitrogen stream (100 K). All data were collected

at the Advanced Light Source (Berkeley, CA) beam line 8.3.1. A single crystal of mNb6 that

grew in 0.1 M Tris.HCl pH 8.5, 1.0 M Ammonium sulfate diffracted to 2.05 Å. Integration, and

scaling were performed with Xia2, using XDS for indexing and integration and XSCALE for

scaling and merging [61]. The structure was solved molecular replacement using PHASER

using the structure of nanobody, Nb.b201 (PDB 5VNV) as search model [45, 62]. Model building

was performed with COOT and refined with PHENIX and BUSTER[54, 56, 63].

Pseudovirus assays for nanobody neutralization

ZsGreen SARS-CoV-2-pseudotyped lentivirus was generated according to a published protocol

[20]. The day before transduction, 50,000 ACE2 expressing HEK293T cells were plated in each

well of a 24-well plate. 10-fold serial dilutions of nanobody were generated in complete medium

(DMEM + 10% FBS + PSG) and pseudotyped virus was added to a final volume of 200 µL.

Media was replaced with nanobody/pseudotyped virus mixture for four hours, then removed.

Cells were washed with complete medium and then incubated in complete medium at 37 °C.

Three days post-transduction, cells were trypsinized and the proportion of ZsGreen+ cells was

measured on an Attune flow cytometer (ThermoFisher).

Authentic SARS-CoV-2 neutralization assay

SARS-CoV-2, isolate France/IDF0372/2020, was supplied by the National Reference Centre for

Respiratory Viruses hosted by Institut Pasteur (Paris, France) and headed by Pr. Sylvie van der

Werf. Viral stocks were prepared by propagation in Vero E6 cells in Dulbecco's modified Eagle's

medium (DMEM) supplemented with 2% (v/v) fetal bovine serum (FBS, Invitrogen). Viral titers

were determined by plaque assay. All plaque assays involving live SARS-CoV-2 were

performed at Institut Pasteur Paris (IPP) in compliance with IPP's guidelines following Biosafety

Level 3 (BSL-3) containment procedures in approved laboratories. All experiments were

performed in at least three biologically independent samples.

Neutralization of infectious SARS-CoV-2 was performed using a plaque reduction neutralization

test in Vero E6 cells (CRL-1586, ATCC). Briefly, nanobodies (or ACE2-Fc) were eight-fold

serially diluted in DMEM containing 2% (v/v) FBS and mixed with 50 plaque forming units (PFU)

of SARS-CoV-2 for one hour at 37°C, 5% CO_2. The mixture was then used to inoculate Vero E6

cells seeded in 12-well plates, for one hour at 37 °C, 5% CO2. Following this virus adsorption

time, a solid agarose overlay (DMEM, 10% (v/v) FBS and 0.8% agarose) was added. The cells

were incubated for a further 3 days prior to fixation using 4% formalin and plaques visualized by

the addition of crystal violet. The number of plaques in quadruplicate wells for each dilution was

used to determine the half maximal inhibitory concentrations (IC_{50}) using 3-parameter logistic

regression (GraphPad Prism version 8).

Nanobody stability studies

Nanobody thermostability by circular dichroism was assessed using a Jasco J710 CD

spectrometer equipped with a Peltier temperature control. Individual nanobody constructs were

diluted to 5 µM in phosphate buffered saline. Mollar ellipticity was measured at 204 nm (2 nm

bandwidth) between 25 °C and 80 °C with a 1 °C/min heating rate. The resulting molar ellipticity

values were normalized and plotted in GraphPad Prism 8.0 after applying a nearest neighbor

smoothing function.

For nanobody competition experiments on ACE2 expressing HEK293T cells, nanobodies were

incubated at either 25°C or 50°C for one hour. Alternatively, each nanobody was aerosolized

with a portable mesh nebulizer producing 2-5 µm particles at a final concentration of 0.5 mg/mL.

The resulting aerosol was collected by condensation into a 50 mL tube cooled on ice. Samples

were then treated as indicated above to determine IC50 values for binding to Spike*-Alexa647.

Further experiments assessing mNb6 and mNb6-tri stability to aerosolization and lyophilization

used a starting concentration of 0.5 mg/mL of each construct. Aerosolization was performed as

described above. For lyophilization, nanobodies were first flash frozen in liquid nitrogen and the

solution was dried to completion under vacuum. The resulting dried material was resuspended

in 20 mM HEPES pH 7.5, 150 mM NaCl. Size exclusion chromatography of the unstressed,

post-aerosolization, and post-lyophilization samples were performed an a Superdex 75 Increase

10/300 column in 20 mM HEPES pH 7.5, 150 mM NaCl. SPR experiments to assess binding to

Spike* were performed as described above.

Acknowledgements

This work was supported by the UCSF Program for Breakthrough Biomedical Research

(PBBR), which is partially funded by the Sandler Foundation. Further support was provided by

the National Institutes of Health (NIH) grant DP5OD023048 (A.Manglik). Cryo-EM equipment at

UCSF is partially supported by NIH grants S10OD020054 and S10OD021741. Work by

M.Vignuzzi was funded by the Laboratoire d'Excellence grant ANR-10-LABX-62-IBEID and the

URGENCE COVID-19 Institut Pasteur fundraising campaign. The radiolytic hydroxyl radical

footprinting is supported by NIH 1R01GM126218. The Advanced Light Source is supported by

the Office of Science, Office of Biological and Environmental Research, of the U.S. DOE under

contract DE-AC02-05CH11231. S.Sangwan was supported by a Helen Hay Whitney

postdoctoral fellowship. C.Billesbølle acknowledges support from the Alfred Benzon Foundation.

K.Leon was funded by NIH/NINDS award F31NS113432 and a UCSF Discovery Fellowship

from the Otellini Family. M.Ott acknowledges support through a gift from the Roddenbury

Foundation. P.Walter is an Investigator of the Howard Hughes Medical Institute. A.Manglik

acknowledges support from the Pew Charitable Trusts, the Esther and A. & Joseph Klingenstein

Fund and the Searle Scholars Program.

We thank the entire Walter and Manglik labs for facilitating the development and rapid execution

of this large-scale collaborative effort. We thank Sebastian Bernales and Tony De Fougerolles

for advice and helpful discussion, and Jonathan Weissman for input into the project and reagent

and machine use. We thank Jim Wells for providing the ACE2 ECD-Fc construct, Jason

McLellan for providing Spike, RBD, and ACE2 constructs, and Florian Krammer for providing an

RBD construct. We thank George Meigs and other Beamline staff at ALS, 8.3.1 for their help in

data collection.

QCRG Structural Biology Consortium.

In addition to those listed explicitly in the author contributions, the structural biology portion of

this work was performed by the QCRG (Quantitative Biosciences

Institute Coronavirus Research Group) Structural Biology Consortium. Listed below are the

contributing members of the consortium listed by teams in order of team relevance to the

published work. Within each team the team leads are italicized (responsible for organization of

each team, and for the experimental design utilized within each team), then the rest of team

members are listed alphabetically. *CryoEM grid freezing/collection team: Caleigh M. Azumaya,*

Cristina Puchades, Ming Sun, Julian R. Braxton, Axel F. Brilot, Meghna Gupta, Fei Li, Kyle E.

Lopez, Arthur Melo, Gregory E. Merz, Frank Moss, Joana Paulino, Thomas H. Pospiech,

Jr., Sergei Pourmal, Alexandrea N. Rizo, Amber M. Smith, Paul V. Thomas, Feng Wang, Zanlin

Yu. *CryoEM data processing team: Miles Sasha Dickinson, Henry C. Nguyen*, Daniel Asarnow,

Julian R. Braxton, Melody G. Campbell, Cynthia M. Chio, Un Seng Chio, Devan Diwanji, Bryan

Faust, Meghna Gupta, Nick Hoppe, Mingliang Jin, Fei Li, Junrui Li, Yanxin Liu, Gregory E.

Merz, Joana Paulino, Thomas H. Pospiech, Jr., Sergei Pourmal, Smriti Sangwan, Raphael

Trenker, Donovan Trinidad, Eric Tse, Kaihua Zhang, Fengbo Zhou. *Crystallography team:*

Nadia Herrera, Huong T. Kratochvil, Ursula Schulze-Gahmen, Michael C. Thompson, Iris D.

Young, Justin Biel, Ishan Deshpande, Xi Liu. *Mammalian cell expression team: Christian Bache*

Billesbølle, Melody G. Campbell, Devan Diwanji, Carlos Nowotny, Amber M. Smith, Jianhua

Zhao, Caleigh M. Azumaya, Alisa Bowen, Nick Hoppe, Yen-Li Li, Phuong Nguyen, Cristina

Puchades, Mali Safari, Smriti Sangwan, Kaitlin Schaefer, Raphael Trenker, Tsz Kin Martin Tsui,

Natalie Whitis. *Protein purification team: Daniel Asarnow, Michelle Moritz, Tristan W. Owens,*

Sergei Pourmal, Caleigh M. Azumaya, Cynthia M. Chio, Amy Diallo, Bryan Faust, Meghna

Gupta, Kate Kim, Joana Paulino, Jessica K. Peters, Kaitlin Schaefer, Tsz Kin Martin

Tsui. *Bacterial expression team: Amy Diallo, Meghna Gupta*, *Erron W. Titus*, Jenny

Chen, Roberto Efraín Díaz, Loan Doan, Sebastian Flores, Mingliang Jin, Huong T. Kratochvil,

Victor L. Lam, Yang Li, Megan Lo, Gregory E. Merz, Joana Paulino, Aye C. Thwin, Stephanie

Wankowicz, Zanlin Yu, Yang Zhang, Fengbo Zhou. *Infrastructure team:* David Bulkley, Arceli

Joves, Almarie Joves, Liam McKay, Mariano Tabios, Eric Tse. *Leadership team: Oren S*

Rosenberg, Kliment A Verba, David A Agard, Yifan Cheng, James S Fraser, Adam Frost,

Natalia Jura, Tanja Kortemme, Nevan J Krogan, Aashish Manglik, Daniel R. Southworth, Robert

M Stroud. The QCRG Structural Biology Consortium has received support from: Quantitative

Biosciences Institute, Defense Advanced Research Projects Agency HR0011-19-2-0020 (to

D.A.Agard and K.A.Verba; B. Shoichet PI), FastGrants COVID19 grant (K.A.Verba PI),

Laboratory For Genomics Research (O.S.Rosenberg PI) and Laboratory for Genomics

Research LGR-ERA (R.M.Stroud PI). R.M.Stroud is supported by NIH grants AI 50476,

GM24485.

Author Contributions

M.Schoof purified Spike*, RBD, and ACE2 proteins, performed yeast display selections to

identify and affinity mature nanobodies, expressed and purified nanobodies, tested activity in

cell-based assays, cloned, expressed, and purified multivalent nanobody constructs, and

949   **Competing Interests**

950   M.Schoof, B.Faust, R.Saunders, N.Hoppe, P.Walter, and A.Manglik are inventors on a

951   provisional patent describing anti-Spike nanobodies described in this manuscript.

REFERENCES

1. Ksiazek, T.G., et al., *A novel coronavirus associated with severe acute respiratory syndrome.* N Engl J Med, 2003. **348**(20): p. 1953-66.
2. Zaki, A.M., et al., *Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia.* N Engl J Med, 2012. **367**(19): p. 1814-20.
3. Zhou, P., et al., *A pneumonia outbreak associated with a new coronavirus of probable bat origin.* Nature, 2020. **579**(7798): p. 270-273.
4. Chan, J.F., et al., *A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster.* Lancet, 2020. **395**(10223): p. 514-523.
5. Huang, C., et al., *Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China.* Lancet, 2020. **395**(10223): p. 497-506.
6. Wu, F., et al., *A new coronavirus associated with human respiratory disease in China.* Nature, 2020. **579**(7798): p. 265-269.
7. Zhu, N., et al., *A Novel Coronavirus from Patients with Pneumonia in China, 2019.* N Engl J Med, 2020. **382**(8): p. 727-733.
8. World Health Organization. *WHO Coronavirus Disease (COVID-19) Dashboard*. 2020; Available from: https://covid19.who.int/.
9. Ke, Z., et al., *Structures, conformations and distributions of SARS-CoV-2 spike protein trimers on intact virions.* bioRxiv, 2020: p. 2020.06.27.174979.
10. Bosch, B.J., et al., *The coronavirus spike protein is a class I virus fusion protein: structural and functional characterization of the fusion core complex.* J. Virol., 2003. **77**(16): p. 8801-8811.
11. Cai, Y., et al., *Distinct conformational states of SARS-CoV-2 spike protein.* Science, 2020.
12. Wang, Q., et al., *Structural and Functional Basis of SARS-CoV-2 Entry by Using Human ACE2.* Cell, 2020. **181**(4): p. 894-904.e9.
13. Yan, R., et al., *Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2.* Science, 2020. **367**(6485): p. 1444-1448.
14. Walls, A.C., et al., *Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein.* Cell, 2020. **181**(2): p. 281-292.e6.
15. Wrapp, D., et al., *Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.* Science, 2020. **367**(6483): p. 1260-1263.
16. Hoffmann, M., et al., *SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor.* Cell, 2020. **181**(2): p. 271-280.e8.
17. Shang, J., et al., *Structural basis of receptor recognition by SARS-CoV-2.* Nature, 2020. **581**(7807): p. 221-224.
18. Gupta, S., et al., *Synchrotron X-ray footprinting as a method to visualize water in proteins.* J Synchrotron Radiat, 2016. **23**(Pt 5): p. 1056-69.
19. Zhang, Y., et al., *Mapping the Binding Interface of VEGF and a Monoclonal Antibody Fab-1 Fragment with Fast Photochemical Oxidation of Proteins (FPOP) and Mass Spectrometry.* J Am Soc Mass Spectrom, 2017. **28**(5): p. 850-858.
20. Crawford, K.H.D., et al., *Protocol and Reagents for Pseudotyping Lentiviral Particles with SARS-CoV-2 Spike Protein for Neutralization Assays.* Viruses, 2020. **12**(5).
21. Kunz, P., et al., *The structural basis of nanobody unfolding reversibility and thermoresistance.* Sci Rep, 2018. **8**(1): p. 7934.
22. Baum, A., et al., *Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with individual antibodies.* Science, 2020.
23. Cao, Y., et al., *Potent Neutralizing Antibodies against SARS-CoV-2 Identified by High-Throughput Single-Cell Sequencing of Convalescent Patients' B Cells.* Cell, 2020. **182**(1): p. 73-84.e16.

24. Chi, X., et al., *A neutralizing human antibody binds to the N-terminal domain of the Spike*
protein of SARS-CoV-2. Science, 2020.
25. Ju, B., et al., *Human neutralizing antibodies elicited by SARS-CoV-2 infection.* Nature,
2020.
26. Liu, L., et al., *Potent neutralizing antibodies directed to multiple epitopes on SARS-CoV-*
2 spike. Nature, 2020.
27. Pinto, D., et al., *Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-*
CoV antibody. Nature, 2020. **583**(7815): p. 290-295.
28. Rogers, T.F., et al., *Isolation of potent SARS-CoV-2 neutralizing antibodies and*
protection from disease in a small animal model. Science, 2020.
29. Zost, S.J., et al., *Potently neutralizing and protective human antibodies against SARS-*
CoV-2. Nature, 2020.
30. Leyva-Grado, V.H., et al., *Direct administration in the respiratory tract improves efficacy*
of broadly neutralizing anti-influenza virus monoclonal antibodies. Antimicrob Agents
Chemother, 2015. **59**(7): p. 4162-72.
31. Detalle, L., et al., *Generation and Characterization of ALX-0171, a Potent Novel*
Therapeutic Nanobody for the Treatment of Respiratory Syncytial Virus Infection.
Antimicrob Agents Chemother, 2016. **60**(1): p. 6-13.
32. Wrapp, D., et al., *Structural Basis for Potent Neutralization of Betacoronaviruses by*
Single-Domain Camelid Antibodies. Cell, 2020. **181**(5): p. 1004-1015.e15.
33. Huo, J., et al., *Neutralizing nanobodies bind SARS-CoV-2 spike RBD and block*
interaction with ACE2. Nat. Struct. Mol. Biol., 2020.
34. Walter, J.D., et al., *Sybodies targeting the SARS-CoV-2 receptor-binding domain.*
bioRxiv, 2020.
35. Sun, Z., et al., *Potent neutralization of SARS-CoV-2 by human antibody heavy-chain*
variable domains isolated from a large library with a new stable scaffold. MAbs, 2020.
12(1): p. 1778435.
36. Wu, Y., et al., *Identification of Human Single-Domain Antibodies against SARS-CoV-2.*
Cell Host Microbe, 2020. **27**(6): p. 891-898.e5.
37. Esparza, T.J. and D.L. Brody, *High Affinity Nanobodies Block SARS-CoV-2 Spike*
Receptor Binding Domain Interaction with Human Angiotensin Converting Enzyme.
bioRxiv, 2020: p. 2020.07.24.219857.
38. Hanke, L., et al., *An alpaca nanobody neutralizes SARS-CoV-2 by blocking receptor*
interaction. bioRxiv, 2020: p. 2020.06.02.130161.
39. Custódio, T.F., et al., *Selection, biophysical and structural analysis of synthetic*
nanobodies that effectively neutralize SARS-CoV-2. bioRxiv, 2020: p.
2020.06.23.165415.
40. Li, T., et al., *Potent synthetic nanobodies against SARS-CoV-2 and molecular basis for*
neutralization. bioRxiv, 2020: p. 2020.06.09.143438.
41. Nieto, G.V., et al., *Fast isolation of sub-nanomolar affinity alpaca nanobody against the*
Spike RBD of SARS-CoV-2 by combining bacterial display and a simple single-step
density gradient selection. bioRxiv, 2020: p. 2020.06.09.137935.
42. Chi, X., et al., *Humanized Single Domain Antibodies Neutralize SARS-CoV-2 by*
Targeting Spike Receptor Binding Domain. bioRxiv, 2020: p. 2020.04.14.042010.
43. Stadlbauer, D., et al., *SARS-CoV-2 Seroconversion in Humans: A Detailed Protocol for a*
Serological Assay, Antigen Production, and Test Setup. Curr. Protoc. Microbiol., 2020.
57(1): p. e100.
44. Lui, I., et al., *Trimeric SARS-CoV-2 Spike interacts with dimeric ACE2 with limited intra-*
Spike avidity. bioRxiv, 2020: p. 2020.05.21.109157.
45. McMahon, C., et al., *Yeast surface display platform for rapid discovery of*
conformationally selective nanobodies. Nat. Struct. Mol. Biol., 2018. **25**(3): p. 289-296.

46.   Mastronarde, D.N., *Automated electron microscope tomography using robust prediction*
1056   *of specimen movements.* J Struct Biol, 2005. **152**(1): p. 36-51.
1057   47.   Zheng, S.Q., et al., *MotionCor2: anisotropic correction of beam-induced motion for*
1058   *improved cryo-electron microscopy.* Nat Methods, 2017. **14**(4): p. 331-332.
1059   48.   Punjani, A., et al., *cryoSPARC: algorithms for rapid unsupervised cryo-EM structure*
1060   *determination.* Nat Methods, 2017. **14**(3): p. 290-296.
1061   49.   Zivanov, J., et al., *New tools for automated high-resolution cryo-EM structure*
1062   *determination in RELION-3.* Elife, 2018. **7**.
1063   50.   Grant, T., A. Rohou, and N. Grigorieff, *cisTEM, user-friendly software for single-particle*
1064   *image processing.* Elife, 2018. **7**.
1065   51.   Lan, J., et al., *Structure of the SARS-CoV-2 spike receptor-binding domain bound to the*
1066   *ACE2 receptor.* Nature, 2020. **581**(7807): p. 215-220.
1067   52.   Rasmussen, S.G., et al., *Structure of a nanobody-stabilized active state of the β(2)*
1068   *adrenoceptor.* Nature, 2011. **469**(7329): p. 175-80.
1069   53.   Frenz, B., et al., *RosettaES: a sampling strategy enabling automated interpretation of*
1070   *difficult cryo-EM maps.* Nat Methods, 2017. **14**(8): p. 797-800.
1071   54.   Emsley, P. and K. Cowtan, *Coot: model-building tools for molecular graphics.* Acta
1072   Crystallogr D Biol Crystallogr, 2004. **60**(Pt 12 Pt 1): p. 2126-32.
1073   55.   Croll, T.I., *ISOLDE: a physically realistic environment for model building into low-*
1074   *resolution electron-density maps.* Acta Crystallogr D Struct Biol, 2018. **74**(Pt 6): p. 519-
1075   530.
1076   56.   Adams, P.D., et al., *PHENIX: a comprehensive Python-based system for*
1077   *macromolecular structure solution.* Acta Crystallogr D Biol Crystallogr, 2010. **66**(Pt 2): p.
1078   213-21.
1079   57.   Staus, D.P., et al., *Allosteric nanobodies reveal the dynamic range and diverse*
1080   *mechanisms of G-protein-coupled receptor activation.* Nature, 2016. **535**(7612): p. 448-
1081   52.
1082   58.   Gupta, S., et al., *The Beamline X28C of the Center for Synchrotron Biosciences: a*
1083   *national resource for biomolecular structure and dynamics experiments using*
1084   *synchrotron footprinting.* J Synchrotron Radiat, 2007. **14**(Pt 3): p. 233-43.
1085   59.   Yu, F., et al., *Fast quantitative analysis of timsTOF PASEF data with MSFragger and*
1086   *IonQuant.* Mol Cell Proteomics, 2020.
1087   60.   Choi, M., et al., *MSstats: an R package for statistical analysis of quantitative mass*
1088   *spectrometry-based proteomic experiments.* Bioinformatics, 2014. **30**(17): p. 2524-6.
1089   61.   Kabsch, W., *Automatic processing of rotation diffraction data from crystals of initially*
1090   *unknown symmetry and cell constants.* Journal of Applied Crystallography, 1993. **26**(6):
1091   p. 795-800.
1092   62.   McCoy, A.J., et al., *Phaser crystallographic software.* J Appl Crystallogr, 2007. **40**(Pt 4):
1093   p. 658-674.
1094   63.   Bricogne G., B.E., Brandl M., Flensburg C., Keller P., Paciorek W., and S.A. Roversi P,
1095   Smart O.S., Vonrhein C., Womack T.O. , *BUSTER version 1.10.0. .* Cambridge, United
1096   Kingdom: Global Phasing Ltd., 2017.
1097