DEER-PREdict: Software for Efficient Calculation of Spin-Labeling EPR and NMR Data from Conformational Ensembles

Giulio Tesei[1,⊘], João M. Martins[1,⊘], Micha B. A. Kunze[1], Yong Wang[1], Ramon Crehuet[1,2], Kresten Lindorff-Larsen[1,*]

1 Structural Biology and NMR Laboratory & the Linderstrøm-Lang Centre for Protein Science, Department of Biology, University of Copenhagen, Copenhagen N, Denmark **2** CSIC-Institute for Advanced Chemistry of Catalonia (IQAC), Jordi Girona 18-26, 08034, Barcelona, Spain

⊘These authors contributed equally to this work.

* lindorff@bio.ku.dk

Abstract

Owing to their plasticity, intrinsically disordered and multidomain proteins require descriptions based on multiple conformations, thus calling for techniques and analysis tools that are capable of dealing with conformational ensembles rather than a single protein structure. Here, we introduce DEER-PREdict, a software to predict Double Electron-Electron Resonance distance distributions as well as Paramagnetic Relaxation Enhancement rates from ensembles of protein conformations. DEER-PREdict uses an established rotamer library approach to describe the paramagnetic probes which are bound covalently to the protein. DEER-PREdict has been designed to operate efficiently on large conformational ensembles, such as those generated by molecular dynamics simulation, to facilitate the validation or refinement of molecular models as well as the interpretation of experimental data. The performance and accuracy of the software is demonstrated with experimentally characterized protein systems: HIV-1 protease, T4 Lysozyme and Acyl-CoA-binding protein. DEER-PREdict is open source

(GPLv3) and available at github.com/KULL-Centre/DEERpredict and as a Python PyPI package pypi.org/project/DEERPREdict.

Introduction

A detailed understanding of protein function often requires an accurate description of the structure and dynamics of a protein. The characterization of protein complexes as well as multi-domain and disordered proteins is typically achieved by combining experimental techniques of distinct spatial resolution [1]. Among the many different experimental techniques that may be used, we focus here on (i) a pulsed electron paramagnetic resonance (EPR) technique called double electron-electron resonance (DEER) and (ii) a nuclear magnetic resonance (NMR) method called paramagnetic relaxation enhancement (PRE). While the two methods differ substantially in their physics and applications, they have in common that they generally involve adding so-called spin-labels to the protein of interest.

DEER, also sometimes known as pulsed electron-electron double resonance (PELDOR), [2–6] relies on probing magnetic dipole-dipole interactions that are sensitive to distributions of residue-residue distances ranging from ~1.8 nm to ~8 nm, and up to 16 nm in deuterated soluble proteins [7–10]. For proteins, DEER generally requires site-directed spin labeling (SDSL) to functionalize a pair of selected residues with paramagnetic probes, e.g. 1-Oxyl-2,2,5,5-tetramethylpyrroline-3-methyl methanethiosulfonate (MTSSL) [4].

PRE NMR also makes use of SDSL to provide information on the average proximity of protein backbone nuclei up to ~3.5 nm away from the unpaired electron of the paramagnetic probe [11]. The dependence of the rate of relaxation enhancement on the electron-proton distance, r, scales as $\langle r^{-6} \rangle$, making the measurement particularly sensitive to contributions from different probe conformations [11].

Since spin labels are conformationally dynamic, both protein and paramagnetic probes need to be described by conformational ensembles to obtain accurate predictions of DEER and PRE observables from molecular models [12–14]. Molecular dynamics (MD) simulations are one approach to obtain conformational ensembles that model the structure and dynamics of spin-labels for the calculation of EPR and NMR data [15–18].

While such analyses can provide unique insight into the motions of and interactions between protein and spin-label [19], they may be relatively expensive computationally. Further, many studies integrate results from multiple probe positions, or pairs thereof, which may be difficult to represent in a single MD simulation with explicit representations of the probes.

Another approach is to use conformational analysis of the spin-label combined with modelling of the dynamics [20–23]. Such analyses suggest that the conformational variation of spin-labelled sites is rotameric, i.e. it can be relatively well described by a finite number of defined structures. Thus, in the calculation of DEER data, rapid modeling of dynamic paramagnetic probes was made possible with the introduction of the rotamer library approach (RLA) applied to the MTSSL probe by Polyhach *et al.* [24].

Here, building and expanding on earlier work [3, 24–27], we developed a software tool for fast predictions of DEER and PRE observables from large conformational ensembles using the RLA. We present our implementation, distributed as the DEER-PREdict software, and test it against experimental data on HIV-1 Protease, T4 Lysozyme and the Acyl-CoA-Binding Protein. This software has been previously used for the calculation of both intra- and intermolecular DEER and PRE NMR data [28, 29], and has some overlap with the features in RotamerConvolveMD [25] (github.com/MDAnalysis/RotamerConvolveMD). DEER-PREdict is open-source, documented (deerpredict.readthedocs.io) and open to contributions from the community.

Design and Implementation

DEER-PREdict is written in Python and is available as a Python API, which facilitates its integration within larger data pipelines. Predictions of DEER and PRE data are carried out via the DEERpredict and PREpredict classes. Both classes are initialized with protein structures (provided as MDAnalysis [30] Universe objects) and spin-labeled positions (residue numbers and chain IDs). As shown in the *Results* section, the calculations are triggered by the *run* function, which also sets additional attributes such as the paths of input and output files as well as experiment-specific parameters. Per-frame data is saved in compressed binary files (HDF5 and pickle files) to allow for

fast calculations of ensemble averages in reweighting schemes. 59

For the presented software, we adopt a procedure of rotamer placement and 60
evaluation of labeled sites which is analogous to the RLA of Polyhach *et al.* [24], and we 61
build on this previous work to implement fast calculations of DEER and PRE 62
observables from large structural ensembles, such as MD trajectories. 63

Rotamer Library Approach 64

Rotamer libraries have a long history in protein structural analysis [31], with an early 65
application being to study side-chain packing [32]. Several other applications of this 66
approach were later employed, e.g. in homology modeling and protein design [33, 34]. In 67
our implementation, the RLA is used to insert the rotamer conformations of a 68
paramagnetic probe at the spin-labeled site and to calculate the Boltzmann weight of 69
each conformer. By default, we use the MTSSL 175 K rotamer library by Polyhach et 70
al. [24], which was filtered to include only the $\chi_1\chi_2$ conformations that are most 71
commonly found in crystal structures of T4 Lysozyme [35]. As shown by Klose *et* 72
al. [26], this selection criterion increases the accuracy of the calculated electron-electron 73
distributions. The code is, however, general and it is possible to add new 74
rotamer libraries by providing a text file containing the Boltzmann weights of each 75
rotamer state p_i^{int}, a topology file (PDB format) and a trajectory file (DCD format) 76
where rotamers are aligned with respect to the the plane defined by Cα atom and C–N 77
peptide bond. These files should be included in the *lib* folder and listed in the yaml file 78
DEERPREdict/lib/libraries.yml. The default MTSSL 298 K MC/UFF CαSδ rotamer 79
libraries of the Matlab-based MMM modeling toolbox [13] are also provided in the 80
DEER-PREdict package. 81

Following the alignment of the rotamer to the protein backbone (Cα, C and N 82
atoms), the calculation of the Boltzmann weights is based on the sum of internal, ϵ_i^{int}, 83
and external, ϵ_i^{ext}, energy contributions. The internal contribution is taken from 84
Polyhach et al. [24] and results from the clustering of representative dihedral 85
combinations from MD simulations. The normalized frequency of each cluster 86
throughout the MD trajectory was used to determine the Boltzmann probability, p_i^{int}, 87
of a given i^{th} state, which readily can be converted into an internal energy contribution, 88

ϵ_i^{int}, via Boltzmann inversion. On the other hand, the external energy contribution is $\qquad$ 89

calculated on the fly as the dispersion interaction energy between heavy atoms of $\qquad$ 90

rotamer and protein residues within a 1-nm cutoff, using the pairwise 6-12 $\qquad$ 91

Lennard-Jones potential of the CHARMM36 force field. $\qquad$ 92

The overall probability of the i^{th} rotamer state is then calculated as $\qquad$ 93

$$p_i = p_i^{int} p_i^{ext} = p_i^{int} \frac{\exp\left(-\epsilon_i^{ext}/kT\right)}{Z} \qquad (1)$$

where $Z = \sum_i p_i^{int} \exp\left(-\epsilon_i^{ext}/kT\right)$ is the steric partition function quantifying the fit of $\qquad$ 94

the rotamer in the embedding protein conformation. Low values of Z result from large $\qquad$ 95

probe-protein van der Waals interaction energies, suggesting a tight placement of the $\qquad$ 96

spin label either due to a displacement of the rotamers or indicative of a wild-type $\qquad$ 97

conformation made inaccessible by the presence of the MTSSL probe. Especially in $\qquad$ 98

folded proteins, probes located in closely packed regions are likely to induce changes in $\qquad$ 99

the ensemble of the spin-labeled protein compared to the native form, and should be $\qquad$ 100

avoided in designing SDSL experiments. Therefore, in the calculation of DEER or PRE $\qquad$ 101

NMR observables, frames with $Z < 0.05$ are discarded to preclude spurious conformers $\qquad$ 102

from contributing to the ensemble average [24]. For the MTSSL 175 K rotamer library, $\qquad$ 103

a Z cutoff of 0.05 is compatible with distributions of ϵ_i^{ext} values where at most one of $\qquad$ 104

the 46 rotamers has $\epsilon_i^{ext} \approx 3\ k_BT$ while the rest has $\epsilon_i^{ext} \leq 7\ k_BT$. We observed that $\qquad$ 105

the results shown in this paper are virtually insensitive to the choice of the Z cutoff $\qquad$ 106

between 0.05 and 0.5 (see Figure S1), therefore, in DEER-PREdict the default Z cutoff $\qquad$ 107

can be conveniently replaced by a user-provided value. $\qquad$ 108

Predicting the DEER signal from structural ensembles $\qquad$ 109

Electron-electron distance distributions extracted from DEER experiments, e.g. using $\qquad$ 110

the DeerLab package [36], have previously routinely been compared with distributions $\qquad$ 111

predicted using the RLA implemented in the Matlab-based MMM modeling toolbox $\qquad$ 112

(http://www.epr.ethz.ch/software) [13]. Since MMM intrinsically operates on single $\qquad$ 113

structures, we and others had to resort to wrapper scripts to compute distance $\qquad$ 114

distributions of large ensembles, such as MD trajectories [3, 25, 37]. With the program $\qquad$ 115

presented herein, we provide a tool to conveniently predict DEER distance distributions $\qquad$ 116

from large conformational ensembles, which can be easily integrated in reweighting   117

schemes such as the Bayesian/maximum entropy procedure [1, 14, 38, 39].   118

For each trajectory frame or conformation of a given ensemble, the rotamers from   119

the library are placed at the spin-labeled position (Figure 1A) and the distances   120

between all pair combinations of N-O paramagnetic centers are calculated. The   121

resulting matrix of pair-wise distances is then used to compute the distance distribution   122

weighted by the combined probability of each probe conformation, $p_i \times p_j$, with p_i and   123

p_j being the conformation probabilities of rotamers i and j. After averaging over all the   124

frames, a low-pass filter is applied to the distance distribution for noise reduction [40],   125

$$P(r) = \mathcal{F} \left\{ \mathcal{F}^{-1} \left[P(r) \right] \times \mathcal{F}^{-1} \left[\exp \left(-\frac{r^2}{2\sigma^2} \right) \right] \right\} \tag{2}$$

where $\mathcal{F}$ and $\mathcal{F}^{-1}$ are the Fourier transform and inverse Fourier transform operators,   126

respectively, whereas σ is the standard deviation of the low-pass filter. The resulting   127

$P(r)$ is a smooth curve even for the analysis of a single protein conformation   128

(Figure 1B). The standard deviation of the low-pass filter can readily be provided by the   129

user through the option *filter_stdev* of the *run* function in the *DEERpredict* class,   130

overriding the default value of 0.5 Å. The average over the trajectory frames can be   131

weighted by a user-specified list of weights e.g. to remove the bias from enhanced   132

sampling simulations.   133

The dipolar modulation signal can be back-calculated from the distance distribution,   134

$P(r)$, via the following integral [41]   135

$$S(t) = \int_0^\infty \mathrm{d}r \ P(r) K(r, t). \tag{3}$$

$K(r, t)$ is the DEER kernel   136

$$K(r, t) = \sqrt{\frac{\pi}{6\omega t}} \left[\cos(\omega) \, \mathrm{Fr} \, \mathrm{C} \left(\sqrt{\frac{6\omega t}{\pi}} \right) + \sin(Dt) \, \mathrm{Fr} \, \mathrm{S} \left(\sqrt{\frac{6\omega t}{\pi}} \right) \right] \tag{4}$$

where FrC and FrS are Fresnel cosine and sine functions, and ω is the dipolar frequency   137

$$\omega = \frac{\mu_0}{4\pi\hbar} \frac{\mu_B^2 g^2}{r^3} \tag{5}$$

where μ_0 is the permeability of free space, μ_B is the Bohr magneton and g is the electron 138

g-factor. The inter-probe distances and the time range from $[0, r_{max}]$ and $[t_{min}, t_{max}]$ 139

with increments $\mathrm{d}r = 0.05$ nm and $\mathrm{d}t$, respectively. The default values $r_{max} = 12$ nm, 140

$t_{min} = 0.01$ μs, $t_{max} = 5.5$ μs and $\mathrm{d}t = 0.01$ μs can be overridden by the user. 141

Following the correction of the experimental DEER time trace for the intermolecular 142

background [36, 42], the resulting form factor can directly be compared with 143

$$V(t) = 1 + \lambda\left[S(t) - 1\right] \tag{6}$$

where $0.02 \leq \lambda \leq 0.5$ is the modulation depth of the experimental signal [43], 144

quantifying the efficiency of the DEER pump pulse [8].

Fig 1. Probe placement scheme and comparison to DEER data. (A) A pool of 46 conformations of the MTSSL probe from the rotamer library are aligned to the backbone of residues K55 and K55' of HIV-1 protease. The color code represent the Boltzmann weights of each rotamer, increasing from blue to red. (B) Electron-electron distance distribution for HIV-1 protease spin labeled at residues K55 and K55'. The blue line is the experimental data from Torbeev *et al.* [44] whereas the red line is the prediction using DEER-PREdict and a crystal structure of HIV-1 protease (PDB code 3BVB).

Prediction of PRE rates and intensity ratios

In analogy to the calculations of electron-electron distances to predict DEER 147

distributions, we extended the use of the RLA to electron-proton separations to improve 148

the accuracy of PRE predictions. We focus here is on PRE NMR experiments that 149

probe the increase in transverse relaxation rates of any backbone proton due to the 150

dipolar interaction with the unpaired electron of the paramagnetic probe:

$$R_2^{ox} = R_2^{red} + \Gamma_2 \tag{7}$$

where R_2^{ox} and R_2^{red} are the transverse relaxation rates in the presence of the spin label in the oxidized or reduced (diamagnetic) state, respectively. We note that it is also possible to measure PREs on other atoms and to probe longitudinal relaxation enhancement, and it would be possible to include such measurements in future versions of DEER-PREdict.

A description of the enhancement of the transverse relaxation due to dipole-dipole interactions in paramagnetic solutions was first proposed by Solomon and Bloembergen [45, 46]

$$\Gamma_2 = \frac{1}{15} \left(\frac{\mu_0}{4\pi}\right)^2 \gamma_I^2 g^2 \mu_B^2 s_e \left(s_e + 1\right) \left[4J(0) + 3J(\omega_I)\right], \tag{8}$$

where γ_I and ω_I are the gyromagnetic ratio and the Larmor frequency of the proton, respectively, whereas s_e is the electron spin quantum number, equal to $1/2$ for nitroxide probe systems. The spectral density function $J(\omega_I)$ can be described using a model-free formalism [47–50], which takes into account the overall molecular tumbling in the external magnetic field as well as the internal motion of the spin label:

$$J(\omega_I) = \langle r^{-6} \rangle \left[\frac{S^2 \tau_c}{1 + \omega_I^2 \tau_c^2} + \frac{(1 - S^2)\tau_t}{1 + \omega_I^2 \tau_t^2}\right] \tag{9}$$

where

$$\tau_c = \left(\frac{1}{\tau_r} + \frac{1}{\tau_s}\right)^{-1} \tag{10}$$

and

$$\tau_t = \left(\frac{1}{\tau_r} + \frac{1}{\tau_s} + \frac{1}{\tau_i}\right)^{-1}. \tag{11}$$

τ_r is the rotational correlation time of the protein, τ_s is the effective electron correlation rate and τ_i is the correlation time of the internal motion (effective correlation time of the spin label). For MTSSL probes, $\tau_s \gg \tau_r$ and $\tau_c \approx \tau_r$ [51]. The value of τ_c depends on protein size and structure and is generally of the order of 1–10 ns [27, 52–55]. For τ_i, values between 100 to 500 ps can be assumed, based on e.g. ^{15}N spin relaxation rates

and MD simulations [56,57]. In general, τ_c and τ_i can be specified as user input in   172

DEER-PREdict.   173

For the generalized order parameter, S, we use the factorization into contributions   174
from radial and angular internal motions introduced by Brüschweiler *et al.* [49],   175
$S^2 = S^2_{radial}S^2_{angular}$. The expressions for S^2_{radial} and $S^2_{angular}$ were derived from a   176
jump model that treats the N conformers of the rotamer library as N discrete states   177
with equal probabilities $(1/N)$ [50]. In reality, the various dihedral angles of the spin   178
label have different free energy barriers, resulting in residence times between jumps   179
ranging from less than 1 to several ns [17].   180

$$S^2_{radial} = \frac{\langle r^{-3} \rangle^2}{\langle r^{-6} \rangle} \tag{12}$$

where r is the proton-electron distance and the brackets denote averages over the   181
conformers weighted by the respective Boltzmann weights, p_i, i.e. $\langle r^{-3} \rangle = \sum_i^N r_i^{-3} p_i$ 182
and $\langle r^{-6} \rangle = \sum_i^N r_i^{-6} p_i$.   183

$$S^2_{angular} = \left\langle \frac{3}{2} \cos^2 \Omega - \frac{1}{2} \right\rangle = \sum_{i,j}^N \left[\frac{3}{2} \left(\frac{\mathbf{r}_i \cdot \mathbf{r}_j}{r_i r_j} \right)^2 - \frac{1}{2} \right] p_i p_j \tag{13}$$

where Ω is the angle between the vectors $\mathbf{r_i}$ and $\mathbf{r_j}$, connecting a backbone proton with   184
the ith and jth rotamer states, respectively. The relaxation enhancement rate for a   185
single protein structure is calculated using Equation 8, and assuming that the motion of   186
the paramagnetic label is much faster than the protein conformational changes, the   187
ensemble average is estimated as   188

$$\langle \Gamma_2 \rangle = \sum_k^M w_k \Gamma_{2,k}. \tag{14}$$

where M is the number of configurations or frames of the simulation trajectory. In the   189
case of unbiased simulations, the statistical weights, w_l, are simply $1/M$. Optionally, a   190
list of weights can be provided by the user, e.g. to reweight a biased MD   191
simulation [58,59] or to incorporate the prediction of the PRE rates into a   192
Bayesian/maximum entropy reweighting scheme [1].   193

For samples with particularly high PRE rates it can be infeasible to obtain Γ_2 from   194

multiple time-point measurements [60]. In such and other cases, the PRE is sometimes   195
probed indirectly from the ratio of the peak intensities in ^{1}H,^{15}N-HSQC spectra of the   196
spin-labeled protein in the oxidized and reduced state. Assuming that the intensity of   197
the proton magnetization decays exponentially — by transverse relaxation only —   198
during the total INEPT time of the HSQC measurement [61], t_d, the intensity ratio is   199
estimated as   200

$$\frac{I_{para}}{I_{dia}} = \frac{R_2^{red} \exp\left(-\Gamma_2 t_d\right)}{R_2^{red} + \Gamma_2}. \tag{15}$$

Requirements and Installation   201

The main requirements are Python 3.6–3.8 and MDAnalysis 1.0 [30, 62]. In an   202
environment with Python 3.6–3.8, DEER-PREdict can readily be installed through the   203
package manager PIP by executing   204

```
1    pip install DEERPREdict
```
    205

Package Stability   206

Tests reproducing DEER and PRE data for the protein systems studied in this article,   207
as well as for a nanodisc [29], are performed automatically using Travis CI   208
(travis-ci.com/github/KULL-Centre/DEERpredict) every time the code is modified on   209
the GitHub repository. The same tests can also be run locally using the test running   210
tool pytest.   211

Results   212

In the following, we present applications of our tool to the prediction of DEER distance   213
distributions and PRE intensity ratios of three folded proteins.   214

    The code snippets reported in this section pertain to DEER-PREdict version 0.1.4.   215
A Jupyter Notebook to reproduce the results shown below (*article.ipynb*) can be found   216
in the *tests/data* folder on the GitHub repository. The up-to-date documentation is   217
available at deerpredict.readthedocs.io.   218

Case study 1: DEER data for HIV-1 Protease

HIV-1 protease (HIV-1PR) is a homodimeric aspartic hydrolase involved in the cleavage of the gag-pol polyprotein complex. The inhibition of this process affects the life cycle of the HIV-1 virus, rendering it noninfectious [63]. The HIV-1PR monomer is composed of 99 residues and presents a structurally stable core region (residues 1-43 and 58-99) and a dynamic region characterized by a β-hairpin turn, called the flap (residues 44-57). The active site is located at the interstice between the core regions of the two monomers, in proximity to the catalytic D25 residues. This cavity is closed off by the dynamic flap regions, which are considered to act as a gate controlling the access to the active site. The dynamics of the flap regions are of utmost importance for the development of inhibitors, and have been extensively studied, both experimentally and *in silico* [44, 64–69]. Based on the relative position of the flaps, three main conformational states have been proposed. In X-ray crystallography, the closed state is typically observed for the ligand-bound enzyme (e.g. PDB codes 3BVB [70] and 2BPX [71]), the semi-open state is predominant for the apo form (e.g. PDB code 1HHP [72]) whereas the wide-open state has been observed for variants (e.g. PDB codes 1TW7 [73] and 1RPI [74]) [69]. In DEER measurements, these conformational states can be resolved by spin-labeling sites K55 and K55' (see Figure S2).

To assess the predictive ability of DEER-PREdict, we generated conformational ensembles of the HIV-1PR homodimer via two different approaches: (a) a single 500-ns unbiased MD simulation, and (b) four independent 125-ns MD simulations restrained with experimental residual dipolar couplings (RDC) data [58, 75] from Roche *et al.* [65, 66] (see SI for methodological details). The initial configuration of our simulations is the X-ray crystal structure of the active-site mutant D25N bound to the inhibitor Darunavir (PDB code 3BVB).

Figure 2 presents a comparison of experimental DEER distance distributions and echo intensity curves with predictions from simulation trajectories of 1,000 frames sampled every 0.5 ns. The echo intensity curves are calculated using Equation 6, where the λ is estimated to 0.0922 by fitting the experimental dipolar evolution function to the corresponding curve derived from the experimental $P(r)$ via Equation 3. For a single trajectory, the analysis is performed in 40 s on a 1.7 GHz processor by running

the following code: 250

```
1  import MDAnalysis                                                          251
2  from DEERPREdict.DEER import DEERpredict                                   252
3  u = MDAnalysis.Universe('conf.pdb','traj.xtc')                            253
4  DEER = DEERpredict(u,residues =[55,55],chains=['A', 'B'],temperature=298)  254
5  DEER.run()                                                                 255
```

The third line generates the MDAnalysis Universe object from an XTC trajectory and a 256
PDB topology. The fourth line initializes the DEERpredict object with the spin-labeled 257
residue numbers and the respective chain IDs. The fifth line runs the calculations and 258
saves per-frame and ensemble-averaged data to *res-55-55.hdf5* and *res-55-55.dat*, 259
respectively, as well as the steric partition functions of sites K55 and K55' to the file 260
res-Z-55-55.dat. 261

Fig 2. Comparing experiments and simulations for HIV1-PR. DEER distance
distributions (A) and echo intensity curves (B) obtained by Torbeev *et al.* [44] from
DEER experiments (blue), and calculated using DEER-PREdict from unbiased (orange)
and RDC ensemble-biased MD simulations (red).

In the experimental distance distribution, the main peak at ~ 3.3 nm corresponds to 262
the closed state whereas the second peak between 4 and 5 nm is characteristic of the 263
wide-open state. The shoulder peak at ~ 2.8 nm has been identified as an open-like 264
state known as the curled/tucked conformation [9, 76, 77]. The results of our unbiased 265
and restrained simulations are in substantial agreement with the findings of Roche *et* 266
al. [65, 66], indicating that the flaps of the inhibitor-free HIV-1PR are predominantly in 267
closed conformation. Compared to the distance distribution calculated from the starting 268
configuration of PDB code 3BVB (see Figure 1), predictions based on MD trajectories 269

more accurately reproduce the shape of the shoulder and the main peak of the 270
experimental $P(r)$. Moreover, using the RDC data as restraints leads to a significant 271
improvement in the agreement between simulations and experiments, with the RMSD 272
decreasing from 0.07 for the unbiased to 0.03 for the RDC ensemble-biased simulations. 273
However, in the simulations we do not observe the wide-open state. This discrepancy 274
could be due to insufficient sampling or could be attributed to the difference in 275
sequence between the simulated protein and the experimental construct. 276

Case study 2: DEER data for T4 Lysozyme 277

Lysozyme from the T4 bacteriophage (T4L) has long been used as a model system in 278
the study of protein structure and dynamics [78–83]. Here, we focus on the L99A and 279
the triple L99A-G113A-R119P mutants which are structurally similar and mainly differ 280
in the relative populations of their major conformational states. The L99A variant 281
presents a $150\,\text{Å}^3$ hydrophobic pocket capable of binding hydrophobic ligands and has 282
been thoroughly studied to further our understanding of the dynamics and selectivity of 283
the binding pocket [78, 84]. The L99A variant occupies two distinct conformational 284
states: the ground state (G) and the transient excited state (E), amounting for 97% and 285
3% of the population, respectively. The large-scale motions converting the G into the E 286
state occur on the millisecond time scale and result in the occlusion of the cavity, which 287
is occupied by the side chain of F114 in the E state [82]. The additional G113A and 288
R119P mutations in the triple-mutant variant interconvert the populations of the 289
conformational states to 4% for the G state and 96% for the E state [82] — note that, 290
here and in the following, we refer to the G and E states based on their structural 291
similarity to the L99A variant rather than on their relative populations. These 292
conformational equilibria have been studied by DEER for various pairs of spin-labeled 293
sites, which effectively resolve the G and E states as separate peaks of the $P(r)$ [83]. 294

Here, we compare DEER distance distributions calculated with DEER-PREdict for 295
two pairs of probe positions (D89C–T109C and T109C–N140C) with the corresponding 296
experimental data by Lerch *et al.* [83]. First, we calculate the $P(r)$ of the single states 297
using PDB code 3DMV for the G states and PDB codes 2LCB and 2LC9 for the E 298
states of single and triple mutants, respectively. Second, the $P(r)$'s are linearly 299

combined based on the experimentally derived ratios of G and E populations (97:3 for L99A and 4:96 for L99A-G113A-R119P) [82]. Additionally, we predict DEER distance distributions from previously reported metadynamics MD simulations of L99A and L99A-G113A-R119P [80]. In these calculations, the average over the trajectory is weighted by $\exp\left(F_{bias}/k_B T\right)$, where F_{bias} is the final static bias for each frame and $k_B T$ is the thermal energy. The analysis of a trajectory of 6,670 frames is performed in 11 min on a 1.7 GHz processor executing the following lines of code:

```
1   import MDAnalysis
2   from DEERPREdict.DEER import DEERpredict
3   import numpy as np
4   u = MDAnalysis.Universe('conf.pdb','traj.xtc')
5   for residues in [[89, 109],[109, 140]]:
6       DEER = DEERpredict(u,residues=residues,temperature=298,z_cutoff=0.1)
7       DEER.run(weights=np.exp(Fbias/(0.298*8.3145)))
```

In line six, we specify the positions of the spin-labels, the temperature at which the metadynamics simulations were performed and a non-default value for the Z cutoff. In line seven, we provide the weights of each trajectory frame, generated from the array of F_{bias} values.

Figure 3 shows a comparison between the experimental distance distributions obtained by Lerch *et al.* [83] and our predictions. In general, the calculated distributions fall within the experimental ranges of inter-probe distances and are particularly accurate for the D89C–T109C spin-labeled pair in metadynamics simulations. The sharper shape of the experimental $P(r)$'s, relative to the calculated distributions, could be due to the cryogenic temperatures at which DEER experiments are conducted, whereas simulations were performed at room temperature. For the T109C–N140C spin-labeled pair of the triple variant, the discrepancy between predicted and calculated $P(r)$'s might be explained by considering that distances shorter than 1.5 nm fall below the range probed by DEER experiments. On the other hand, the inaccurate predictions of the T109C–N140C $P(r)$ for the single (L99A) variant is greater than expected. Such discrepancies may be due both to errors in the protein structure or in the DEER-calculations. While our results cannot distinguish between these scenarios, we follow previous work [14] by examining whether the discrepancies can can be attributed to the error on the Boltzmann probabilities of the rotamer states,

p_i^{int}. We thus use a Bayesian/maximum entropy procedure to show that a small change ₃₃₃ in the original rotamer weights can lead to a substantial improvement of the agreement ₃₃₄ with the experimental data (see Figure S3). ₃₃₅

Fig 3. Comparing experiments with simulations and structures of T4 Lysozyme variants. DEER distance distributions for probe positions (A) D89C–T109C and (B) T109C–N140C of the single (blue) and the triple variant (red). Solid lines are the experimental data by Lerch et al. [83], dotted lines are calculated from PDB codes and dashed lines are predictions from metadynamics (MTD) simulations by Wang and coworkers [80].

Case study 3: PRE data for Acyl-CoA-Binding Protein ₃₃₆

The RLA is well known in the EPR community and generally favored over e.g. a ₃₃₇ Cα-based approach as discussed elsewhere [3, 13, 26]. In the presented software, we ₃₃₈ apply the same improved modeling of the probe flexibility also to the prediction of PRE ₃₃₉ rates and intensity ratios. ₃₄₀

Our test data is the PRE data for the bovine Acyl-coenzyme A Binding Protein ₃₄₁ (ACBP) reported by Teilum *et al.* [53]. In this study the structural behavior of ACBP ₃₄₂ under native and mildly-denaturing conditions was investigated via the SDSL of five ₃₄₃ positions in the amino acid the sequence: T17C, V36C, M46C, S65C and I86C. Here, ₃₄₄ we focus on the native state of ACBP for which an NMR structure comprising 20 ₃₄₅ conformers has been refined from residual dipolar couplings (RDC) and deposited in the ₃₄₆ Protein Data Bank (PDB code 1NTI). Figure 4 shows a comparison between the ₃₄₇ experimental data and the intensity ratios calculated from the Γ_2 values averaged over ₃₄₈ the 20 conformations of the PDB entry. A good overall agreement is achieved across the ₃₄₉

different probe positions. Notably, using the RDC-refined structure, we reproduce most ⁣ ₃₅₀
of the structural features observed in the PRE experiments, including the proximity of ⁣ ₃₅₁
residues 24, 27, 31 and 34 to the spin-labeled residue 86, which is consistent with a ⁣ ₃₅₂
helix-turn-helix motif. The predicted intensity ratios are generated in 2 s on a 1.7 GHz ⁣ ₃₅₃
processor executing the following code: ⁣ ₃₅₄

```
1  import MDAnalysis
2  from DEERPREdict.PRE import PREpredict
3  u = MDAnalysis.Universe('1nti.pdb')
4  for res in [17,36,46,65,86]:
5      PRE = PREpredict(u,res,temperature=298,atom_selection='H')
6      PRE.run(tau_c=2e-09,tau_t=2*1e-10,delay=1e-2,r_2=12.6,wh=750)
```

At line three, we load PDB code 1NTI as an MDAnalysis Universe object. We then use ⁣ ₃₆₁
a for loop to calculate the PRE data from the distances between amide protons and the ⁣ ₃₆₂
spin-label N-O group at five different positions along the amino acid sequence. In the ⁣ ₃₆₃
last line we specify $\tau_c = 2$ ns, $\tau_t = 0.2$ ns, $t_d = 10$ ms, $R_2 = 12.6 \, \text{s}^{-1}$ and ⁣ ₃₆₄
$\omega_I = 2\pi \times 750$ MHz. Per-frame and ensemble-averaged PRE data are automatically ⁣ ₃₆₅
saved to files named *res-*.pkl* and *res-*.dat*, respectively, whereas per-frame steric ⁣ ₃₆₆
partition functions are saved to *res-Z-*.dat*. ⁣ ₃₆₇

As detailed in Figure S4, the steric partition functions provided by DEER-PREdict ⁣ ₃₆₈
can be used to predict whether a position in the sequence is likely to accommodate the ⁣ ₃₆₉
paramagnetic probe within the wild-type structure. Besides aiding the interpretation of ⁣ ₃₇₀
experimental data, this feature can be instrumental to designing and enhancing the ⁣ ₃₇₁
success-rate of time- and labor-intensive SDSL experiments. ⁣ ₃₇₂

As previously discussed, the explicit treatment of the paramagnetic probe may be ⁣ ₃₇₃
crucial for the accurate back-calculation of DEER data, and even more so for PRE ⁣ ₃₇₄
predictions, due to the $\langle r^{-6} \rangle$-dependence of the PRE. A common way to restrain MD ⁣ ₃₇₅
simulations or to back-calculate PRE experimental data without explicitly simulating ⁣ ₃₇₆
the paramagnetic probe is to approximate the electron location to the position of the ⁣ ₃₇₇
Cβ atom of the spin-labeled residue [85]. The advantage of this approach is that ⁣ ₃₇₈
(a) multiple labeling sites can be analyzed in a single simulation and (b) the explicit ⁣ ₃₇₉
atom is present in the simulation making the calculation of PREs straightforward. ⁣ ₃₈₀
Cβ-based calculations may, however, be prone to over- or underestimating ⁣ ₃₈₁

Fig 4. Calculated and experimental PRE HSQC intensity ratios for the T17C, V36C, M46C, S65C and I86C mutants of ACBP. Blue lines represent the experimental data [53], with the associated ± 0.1 error shown by the blue shaded areas. Red lines represent intensity ratios calculated from PDB code 1NTI with $\tau_c = 2\,\text{ns}$, $\tau_t = 0.2\,\text{ns}$, $t_d = 10\,\text{ms}$, $R_2 = 12.6\,\text{s}^{-1}$.

electron-proton distances by several Å, thereby introducing a systematic error. The impact of the Cβ-approximation on the accuracy of PRE predictions is illustrated in Figure S5 and Figure S6 for the case of ACBP.

Conclusion

We have introduced an open-source software solution with a fast implementation of the RLA in tandem with protein ensemble averaging, for the calculation of DEER and PRE data. Using three examples, we have highlighted the capabilities of our implementation: (a) the extension of the RLA for DEER data from a protein ensemble and (b) the calculation of PRE rates and intensity ratios with the same approach.

The structural interpretation of DEER and PRE measurements requires an accurate treatment of the structure and conformational heterogeneity of the spin labels. In the presented software, this is achieved using the RLA and, in the case of the PRE, a model-free approach to describe the dynamics. Relative to simulations of the explicitly spin-labeled mutants, the RLA presents the particular advantage of enabling the prediction for multiple SDSL experiments from a single simulation of the wild type

sequence.

Availability and Future Directions

The software is implemented using the popular trajectory analysis package MDAnalysis, version 1.0 [30] and is available on GitHub at github.com/KULL-Centre/DEERpredict. DEER-PREdict is also distributed as a PyPI package (pypi.org/project/DEERPREdict) and archived on Zenodo (DOI: 10.5281/zenodo.3968394). DEER-PREdict and MDAnalysis are published and distributed under GPL licenses, version 3 and 2, respectively.

DEER-PREdict has a general framework and can be readily extended to encompass non-protein biomolecules as well as additional rotamer libraries of paramagnetic groups. Moreover, the software can be augmented with a module to predict Förster resonance energy transfer data, combining the insertion routines already implemented for MTSSL probes with rotamer libraries for fluorescent dyes.

Acknowledgments

We thank Robert Best for help with RDC-restrained simulations as well as work on extending DEER-PREdict to use for prediction of FRET experiments. M.B.A.K. acknowledges funding from the Lundbeck Foundation. R.C. acknowledges funding from MINECO (CTQ2016-78636-P). K.L.-L. acknowledges funding via a Sapere Aude Starting Grant from the Danish Council for Independent Research and the Lundbeck Foundation BRAINSTRUC initiative in structural biology.

Author Contributions

Conceptualization: Giulio Tesei, João M. Martins, Micha B. A. Kunze, Kresten Lindorff-Larsen

Data curation: Giulio Tesei, João M. Martins, Micha B. A. Kunze, Yong Wang

Funding acquisition: Ramon Crehuet, Kresten Lindorff-Larsen

Software: Giulio Tesei, João M. Martins, Micha B. A. Kunze, Ramon Crehuet

Supervision: Micha B. A. Kunze, Kresten Lindorff-Larsen

Writing — original draft: Giulio Tesei, João M. Martins, Micha B. A. Kunze, Kresten Lindorff-Larsen

Writing — review & editing: Giulio Tesei, João M. Martins, Micha B. A. Kunze, Yong Wang, Ramon Crehuet, Kresten Lindorff-Larsen

Supporting Information

Molecular Dynamics Simulations

HIV-1 protease

All simulations were performed using GROMACS 5.1 [86] with the PLUMED 2 [85] plugin. Unbiased and RDC-biased metadynamics [75, 87] simulations where performed starting from a closed conformer after removal of the inhibitor from the X-ray crystal structure (PDB code 3BVB) [88]. The simulated protein is the wild type subtype B from isolate BRU/LAI, which differs from the construct of the reference experiments [44] by the following mutations: M36norleucine, S37N, R41pseudo-homoglutamine, M46norleucine, I63P, I64V, A67α-aminobutyric acid and A95α-aminobutyric acid. The protein was simulated in a cubic box with a side length of 8.677 nm containing 22,228 water molecules, 59 sodium cations, and 67 chloride anions. Although the K55C mutations have been shown to have a negligible impact on enzymatic activity [64], we maintained the lysine residues at the spin-labeled positions with the aim of capturing the conformational ensemble of the wild type.

The system was equilibrated for 10 ns with a 2-fs time step in the NPT ensemble, using the Berendsen barostat [89] with 0.5-ps coupling constant and isothermal compressibility of 4.5e-5 bar^{-1}. Starting from the equilibrated structure, a production run of 500 ns was performed in the NVT ensemble for the unbiased MD simulation. For the restrained simulations, we obtained the backbone N-H RDCs for the inhibitor-free HIV-1PR from Roche *et al.* [66] and applied a linear potential (force constant 25,000 kJ/mol) between the experimental data and the RDCs calculated as averages over 4 independent simulations. Each replica was simulated for 125 ns in the NVT ensemble. We used the AMBER ff99SB*-ILDN force field [90, 91] for all simulations. First, the structure was minimized with the steepest descent algorithm to a tolerance of 10 kJ mol^{-1} nm^{-1} with restrained water molecules. Second, we simulated the system the for 5 ns in the NVT ensemble using the leap-frog integrator with a time step of 1 fs. A 1.2 nm cutoff was used for van der Waals interactions, with a force-switch modification at 1.0 nm. Coulomb interactions were treated with the particle-mesh Ewald method [92] with direct space cutoff of 1.2 nm. The temperature was set to 298 K

using the v-rescale thermostat [93] with a 5 ps coupling constant. All the bonds   458

involving hydrogen atoms were constrained using the LINCS algorithm [94].   459

T4 Lysozyme   460

The simulations of the L99A single mutant and the L99A-G113A-R119P triple mutant   461

of T4 Lysozyme analysed in this study have been reported previously by Wang *et*   462

al. [80]. The X-ray crystal structure of PDB code 3DMV was used as the initial   463

configuration for the G states of both single and triple mutants. Simulations of the E   464

states were started from chemical-shift-derived Rosetta-calculated structures, i.e. PDB   465

code 2LCB and PDB code 2LC9 for single and triple mutant, respectively. Further   466

simulations details can be found in the original article [80].   467

Fig. S1   Influence of the Z cutoff on predicted DEER and PRE NMR   468

data.

(A) DEER distance distributions calculated from RDC ensemble-biased MD simulations
of HIV-1PR. (B) Predicted intensity ratios for ACBP spin-labeled at position 86
obtained from PDB code 1NTI with $\tau_c = 2\,\text{ns}$, $\tau_t = 0.2\,\text{ns}$, $t_d = 10\,\text{ms}$, $R_2 = 12.6\,\text{s}^{-1}$.
DEER and PRE predictions are performed using three different cutoff values of the
steric partition function, Z, namely 0.05 (blue lines), 0.5 (orange lines) and 0.8 (red
lines).

Fig. S2   Comparison of DEER data from Torbeev _et al._ [44] with X-ray   470
crystal structures deposited in the Protein Data Bank. This figure shows that   471
although the DEER data is calculated from single X-ray crystal structures, the RLA   472
results in multimodal distance distributions. For example, the K55-K55' separation   473
between the ammonium groups in 1TW7 is ~ 3.6 nm, which is consistent with the   474
semi-open conformation, however, the distances between the nitroxide groups of the   475
MTSSL conformers span a wide range between 3.3 and 4.4 nm.

DEER distance distributions (A) and echo intensity curves (B) obtained by Torbeev _et al._ [44] from DEER experiments (blue), and calculated using X-ray crystal structures representative of closed (PDB code 2BPX, orange), semi-open (PDB code 1HHP, green) and wide-open (PDB code 1TW7, red) HIV-1PR conformations.

Fig. S3   Optimization of rotamer weights using a Bayesian/maximum entropy procedure. We use a Bayesian/maximum entropy (BME) procedure to find a modified set of rotamer weights for the MTSSL 175 K rotamer library [24], $\boldsymbol{w}$, which improves the agreement between predicted and experimental T109C–N140C $P(r)$'s for the single variant of T4L. The prediction is based on the 97:3 linear combination of $P(r)$'s from PDB codes 3DMV and 2LCB (corresponding to the populations of these two states) whereas the experimental data is from Lerch *et al.* [83]. Simulated annealing is used to minimize the cost function $\mathcal{L}(\boldsymbol{w}) = \chi^2(\boldsymbol{w}) - \theta S(\boldsymbol{w})$ where $\chi^2(\boldsymbol{w})$ is the sum of the squared differences between predicted and experimental $P(r)$'s, θ quantifies the confidence in the original weights, $\boldsymbol{w^O}$, and $S(\boldsymbol{w}) = -\sum_{i=1}^{N} w_i \ln \frac{w_i}{w_i^O}$ is the relative entropy defined as the negative of the Kullback-Leibler divergence between the modified weights of the $N = 46$ rotamers, w_i, and the initial w_i^O. The effective fraction of rotamers used in the reweighted ensemble, compared to the original library, is quantified as $\phi_{eff}(w) = \exp\left[S(w)\right]$. We scan various values of θ and select the weights obtained using $\theta = 4$ as the smallest modification resulting in a substantial decrease in χ^2. It is noteworthy that the change in weights has a lesser impact on the triple variant than on the single variant. Moreover, albeit being optimized against the T109C–N140C $P(r)$ of the single variant, the modified weights lead to an overall improvement in accuracy, with the average χ^2 over the four $P(r)$'s decreasing from 7.5 to 5.9.

(A) χ^2 vs ϕ_{eff} for various values of the confidence parameter, θ. (B) Distance distributions calculated from PDB codes 3DMV and 2LCB, using optimized weights obtained for various θ values. (C) Original [24] and modified weights of the MTSSL 175 K rotamer library after BME reweighting with $\theta = 4$. DEER distance distributions for probe positions (D) D89C–T109C and (E) T109C–N140C of the single (blue) and the triple variant (red). Solid lines are the experimental data by Lerch *et al.* [83]; dotted and dashed lines are from PDB codes 3DMV, 2LC9 and 2LCB using the original and the BME-reweighted ($\theta = 4$) MTSSL 175 K rotamer library.

Fig. S4   Steric partition function quantifying the fitness of the rotamers   496

at the spin-labeled site. While for most spin-label sites of ACBP the steric partition   497

function, Z, varies between 1 and 1.5, for the placement of the probe at residue 86, Z 498

drops below the cutoff of 0.05 proposed by Polyhach *et al.* [24] in five conformers out of   499

20. The significantly lower Z values indicate that residue 86 is in a tightly packed   500

region in the protein structure and that spin-labeling position 86 can lead to structural   501

deformations or changes in the populations of the conformational ensemble. This   502

observation is consistent with stability experiments performed by Teilum *et al.* on wild   503

type and spin-labeled mutants [53], showing that I86C is the least stable of the studied   504

spin-labeled mutants. Although the RLA assumes that the overall protein conformation   505

is unaffected by the presence of the spin-label, the case of the I86C mutant of ACBP   506

highlights how DEER-PREdict can help to identify spin-labeled sites that are likely to   507

violate this assumption in folded proteins.

Steric partition function calculated from rotamer-protein van der Waals interactions for
five spin-labeled mutants of ACBP. The horizontal dashed line indicates the cutoff used
in the criterion for discarding protein conformations where the placement of the rotamer
is characterized by steric clashes with the surrounding residues.

Fig. S5 Comparison with Cβ-based PRE Predictions The Cβ approximation   509
may overestimate the effect of transient tertiary contacts on the PRE rates of ACBP,   510
yielding experimentally consistent predictions only when the time scales for the   511
reorientational dynamics is artificially made fast.

PRE intensity ratios for ACBP spin labeled at position 65 calculated for (A) $\tau_c = 2\,\text{ns}$
and (B) $\tau_c = 0.5\,\text{ns}$. Blue lines represent the experimental data [53], with the associated
± 0.1 error shown by the blue shaded areas. Orange and red lines represent Cβ-based
and RLA-based predictions, respectively.

Fig. S6   Comparison of optimal τ_c values for RLA vs. Cβ-based approach.
This figure illustrates the systematic time-scale difference when using explicit MTSSL probes instead of approximating the location of the unpaired electron with the position of the Cβ atom of the spin-labeled residue. Compared with the C_β-based approach, the RLA improves the accuracy in reproducing the experimental PRE data, as shown by the generally lower RMSD values. When using the RLA, the τ_c values that minimise the RMSD are found in the range of typical reorientational correlation time constants for proteins of ~ 100 residues (2–5 ns), and closer to the experimentally-derived value of 4 ns [53], whereas the Cβ-based approach underestimates the optimal τ_c.

Dependence on τ_c of the RMSD between experimental and predicted PRE ratios of ACBP. Red and blue lines are obtained using the RLA and approximating the electron location with the position of the Cβ atom, respectively. Solid and dashed lines represent the RMSD values calculated from all the data points and from intensity ratios in the dynamic range $0.1 < I_{para} / I_{dia} < 0.9$.

References

1. Orioli S, Larsen AH, Bottaro S, Lindorff-Larsen K. How to learn from inconsistencies: Integrating molecular simulations with experimental data. In: Computational Approaches for Understanding Dynamical Systems: Protein Folding and Assembly. Elsevier; 2020. p. 123–176.

2. Pannier M, Veit S, Godt A, Jeschke G, Spiess HW. Dead-Time Free Measurement of Dipole–Dipole Interactions between Electron Spins. Journal of Magnetic Resonance. 2000;142(2):331–340.

3. Bagnéris C, Rogala KB, Baratchian M, Zamfir V, Kunze MBA, Dagless S, et al. Probing the Solution Structure of IκB Kinase (IKK) Subunit γ and Its Interaction with Kaposi Sarcoma-associated Herpes Virus Flice-interacting Protein and IKK Subunit β by EPR Spectroscopy. Journal of Biological Chemistry. 2015;290(27):16539–16549.

4. Klare JP, Steinhoff HJ. Spin labeling EPR. Photosynthesis Research. 2009;102(2-3):377–390.

5. Phan G, Remaut H, Wang T, Allen WJ, Pirker KF, Lebedev A, et al. Crystal structure of the FimD usher bound to its cognate FimC–FimH substrate. Nature. 2011;474(7349):49–53.

6. Klose D, Voskoboynikova N, Orban-Glass I, Rickert C, Engelhard M, Klare JP, et al. Light-induced switching of HAMP domain conformation and dynamics revealed by time-resolved EPR spectroscopy. FEBS Letters. 2014;588(21):3970–3976.

7. Schmidt T, Wälti MA, Baber JL, Hustedt EJ, Clore GM. Long Distance Measurements up to 160 Å in the GroEL Tetradecamer Using Q-Band DEER EPR Spectroscopy. Angewandte Chemie International Edition. 2016;55(51):15905–15909.

8. Edwards TH, Stoll S. A Bayesian approach to quantifying uncertainty from experimental noise in DEER spectroscopy. J Magn Reson. 2016;270:87–97.

9. Casey TM, Fanucci GE. Spin labeling and Double Electron-Electron Resonance (DEER) to Deconstruct Conformational Ensembles of HIV Protease. In: Methods in Enzymology. Elsevier; 2015. p. 153–187.

10. Jeschke G. DEER Distance Measurements on Proteins. Annual Review of Physical Chemistry. 2012;63(1):419–446.

11. Clore GM, Iwahara J. Theory, practice, and applications of paramagnetic relaxation enhancement for the characterization of transient low-population states of biological macromolecules and their complexes. Chem Rev. 2009;109(9):4108–4139.

12. Fajer P, Fajer M, Zawrotny M, Yang W. Simulation of spin label structure and its implication in molecular characterization. Methods in enzymology. 2015;563:623.

13. Jeschke G. MMM: A toolbox for integrative structure modeling. Protein Science. 2017;27(1):76–85.

14. Reichel K, Stelzl LS, Köfinger J, Hummer G. Precision DEER distances from spin-label ensemble refinement. The journal of physical chemistry letters. 2018;9(19):5748–5752.

15. Budil DE, Sale KL, Khairy KA, Fajer PG. Calculating slow-motional electron paramagnetic resonance spectra from molecular dynamics using a diffusion operator approach. The Journal of Physical Chemistry A. 2006;110(10):3703–3713.

16. Ding F, Layten M, Simmerling C. Solution structure of HIV-1 protease flaps probed by comparison of molecular dynamics simulation ensembles and EPR experiments. Journal of the American Chemical Society. 2008;130(23):7184–7185.

17. Sezer D, Freed JH, Roux B. Parametrization, molecular dynamics simulation, and calculation of electron spin resonance spectra of a nitroxide spin label on a polyalanine α-helix. The journal of physical chemistry B. 2008;112(18):5755–5767.

18. Xue Y, Skrynnikov NR. Motion of a disordered polypeptide chain as studied by paramagnetic relaxation enhancements, 15N relaxation, and molecular dynamics

simulations: how fast is segmental diffusion in denatured ubiquitin? Journal of the American Chemical Society. 2011;133(37):14614–14628.

19. Sasmal S, Lincoff J, Head-Gordon T. Effect of a Paramagnetic Spin Label on the Intrinsically Disordered Peptide Ensemble of Amyloid-β. Biophysical journal. 2017;113(5):1002–1011.

20. Robinson B, Slutsky L, Auteri F. Direct simulation of continuous wave electron paramagnetic resonance spectra from Brownian dynamics trajectories. The Journal of chemical physics. 1992;96(4):2609–2616.

21. Steinhoff HJ, Hubbell WL. Calculation of electron paramagnetic resonance spectra from Brownian dynamics trajectories: application to nitroxide side chains in proteins. Biophysical journal. 1996;71(4):2201–2212.

22. Tombolato F, Ferrarini A, Freed JH. Dynamics of the nitroxide side chain in spin-labeled proteins. The Journal of Physical Chemistry B. 2006;110(51):26248–26259.

23. Tombolato F, Ferrarini A, Freed JH. Modeling the effects of structure and dynamics of the nitroxide side chain on the ESR spectra of spin-labeled proteins. The Journal of Physical Chemistry B. 2006;110(51):26260–26271.

24. Polyhach Y, Bordignon E, Jeschke G. Rotamer libraries of spin labelled cysteines for protein studies. Phys Chem Chem Phys. 2011;13(6):2356–2366.

25. Stelzl LS, Fowler PW, Sansom MSP, Beckstein O. Flexible Gates Generate Occluded Intermediates in the Transport Cycle of LacY. J Mol Biol. 2014;426(3):735–751.

26. Klose D, Klare JP, Grohmann D, Kay CWM, Werner F, Steinhoff HJ. Simulation vs. Reality: A Comparison of In Silico Distance Predictions with DEER and FRET Measurements. PloS One. 2012;7(6):e39492.

27. Salmon L, Nodet G, Ozenne V, Yin G, Jensen MR, Zweckstetter M, et al. NMR Characterization of Long-Range Order in Intrinsically Disordered Proteins. J Am Chem Soc. 2010;132(24):8407–8418.

28. Milkovic NM, Thomasen FE, Cuneo MJ, Grace CR, Martin EW, Nourse A, et al..   607
    Interplay of folded domains and the disordered low-complexity domain in   608
    mediating hnRNPA1 phase separation. BioRxiv [Preprint]. 2020; bioRxiv   609
    2020.05.15.096966 [cited 2020 Aug 12]. Available from: 610
    https://doi.org/10.1101/2020.05.15.096966.   611

29. Bengtsen T, Holm VL, Kjølbye LR, Midtgaard SR, Johansen NT, Tesei G, et al.   612
    Structure and dynamics of a nanodisc by integrating NMR, SAXS and SANS   613
    experiments with molecular dynamics simulations. eLife. 2020;9.   614

30. Michaud-Agrawal N, Denning EJ, Woolf TB, Beckstein O. MDAnalysis: A   615
    toolkit for the analysis of molecular dynamics simulations. Journal of   616
    Computational Chemistry. 2011;32(10):2319–2327.   617

31. Dunbrack Jr RL. Rotamer libraries in the 21st century. Current opinion in   618
    structural biology. 2002;12(4):431–440.   619

32. Ponder JW, Richards FM. Tertiary templates for proteins. J Mol Biol.   620
    1987;193(4):775–791.   621

33. Bower MJ, Cohen FE, Dunbrack RL. Prediction of protein side-chain rotamers   622
    from a backbone-dependent rotamer library: A new homology modeling tool. J   623
    Mol Biol. 1997;267(5):1268–1282.   624

34. Desjarlais JR, Handel TM. De novo design of the hydrophobic cores of proteins.   625
    Protein Sci. 1995;4(10):2006–2018.   626

35. Fleissner MR, Cascio D, Hubbell WL. Structural origin of weakly ordered   627
    nitroxide motion in spin-labeled proteins. Protein Sci. 2009;18(5):893–908.   628

36. Ibáñez LF, Jeschke G, Stoll S. DeerLab: A comprehensive toolbox for analyzing   629
    dipolar EPR spectroscopy data. Magnetic Resonance Discussions. 2020;.   630

37. Göddeke H, Timachi MH, Hutter CAJ, Galazzo L, Seeger MA, Karttunen M,   631
    et al. Atomistic Mechanism of Large-Scale Conformational Transition in a   632
    Heterodimeric ABC Exporter. Journal of the American Chemical Society.   633
    2018;140(13):4543–4551.   634

38. Köfinger J, Stelzl LS, Reuter K, Allande C, Reichel K, Hummer G. Efficient ensemble refinement by reweighting. Journal of chemical theory and computation. 2019;15(5):3390–3401.

39. Bottaro S, Bengtsen T, Lindorff-Larsen K. Integrating Molecular Simulation and Experimental Data: A Bayesian/Maximum Entropy Reweighting Approach. In: Methods in Molecular Biology. Springer US; 2020. p. 219–240.

40. Hayes J. Some observations on digital smoothing of electroanalytical data based on the Fourier Transformation. Analytical Chemistry. 1973;45(2):277–284.

41. Worswick SG, Spencer JA, Jeschke G, Kuprov I. Deep neural network processing of DEER data. Science Advances. 2018;4(8):eaat5218.

42. Jeschke G, Chechik V, Ionita P, Godt A, Zimmermann H, Banham J, et al. DeerAnalysis2006—a comprehensive software package for analyzing pulsed ELDOR data. Applied Magnetic Resonance. 2006;30(3-4):473–498.

43. del Alamo D, Tessmer MH, Stein RA, Feix JB, Mchaourab HS, Meiler J. Rapid Simulation of Unprocessed DEER Decay Data for Protein Fold Prediction. Biophysical Journal. 2020;118(2):366–375. doi:10.1016/j.bpj.2019.12.011.

44. Torbeev VY, Raghuraman H, Mandal K, Senapati S, Perozo E, Kent SBH. Dynamics of "Flap" Structures in Three HIV-1 Protease/Inhibitor Complexes Probed by Total Chemical Synthesis and Pulse-EPR Spectroscopy. J Am Chem Soc. 2009;131(3):884–885.

45. Solomon I. Relaxation Processes in a System of Two Spins. Phys Rev. 1955;99(2):559.

46. Bloembergen N. Proton Relaxation Times in Paramagnetic Solutions. J Chem Phys. 1957;27(2):572–573.

47. Lipari G, Szabo A. Model-free approach to the interpretation of nuclear magnetic resonance relaxation in macromolecules. 1. Theory and range of validity. Journal of the American Chemical Society. 1982;104(17):4546–4559.

48. Olejniczak ET, Dobson CM, Karplus M, Levy RM. Motional averaging of proton nuclear Overhauser effects in proteins. Predictions from a molecular dynamics simulation of lysozyme. J Am Chem Soc. 1983;106:1923–1930.

49. Brueschweiler R, Roux B, Blackledge M, Griesinger C, Karplus M, Ernst RR. Influence of rapid intramolecular motion on NMR cross-relaxation rates. A molecular dynamics study of antamanide in solution. J Am Chem Soc. 1991;114:2289–2302.

50. Iwahara J, Schwieters CD, Clore GM. Ensemble Approach for NMR Structure Refinement against 1H Paramagnetic Relaxation Enhancement Data Arising from a Flexible Paramagnetic Group Attached to a Macromolecule. J Am Chem Soc. 2004;126(18):5879–5896.

51. Clore GM, Iwahara J. Theory, Practice, and Applications of Paramagnetic Relaxation Enhancement for the Characterization of Transient Low-Population States of Biological Macromolecules and Their Complexes. Chemical Reviews. 2009;109(9):4108–4139.

52. Bibow S, Ozenne V, Biernat J, Blackledge M, Mandelkow E, Zweckstetter M. Structural Impact of Proline-Directed Pseudophosphorylation at AT8, AT100, and PHF1 Epitopes on 441-Residue Tau. Journal of the American Chemical Society. 2011;133(40):15842–15845.

53. Teilum K, Kragelund BB, Poulsen FM. Transient Structure Formation in Unfolded Acyl-coenzyme A-binding Protein Observed by Site-directed Spin Labelling. J Mol Biol. 2002;324(2):349–357.

54. Liu W, Liu X, Zhu G, Lu L, Yang D. A Method for Determining Structure Ensemble of Large Disordered Protein: Application to a Mechanosensing Protein. Journal of the American Chemical Society. 2018;140(36):11276–11285.

55. Gomes GNW, Krzeminski M, Namini A, Martin EW, Mittag T, Head-Gordon T, et al.. Integrating multiple experimental data to determine conformational ensembles of an intrinsically disordered protein. BioRxiv [Preprint]. 2020; bioRxiv 2020.02.05.935890 [cited 2020 Aug 12]. Available from: https://www.biorxiv.org/content/10.1101/2020.02.05.935890v2.

56. Khan SN, Charlier C, Augustyniak R, Salvi N, Déjean V, Bodenhausen G, et al. 692
    Distribution of Pico- and Nanosecond Motions in Disordered Proteins from 693
    Nuclear Spin Relaxation. Biophysical Journal. 2015;109(5):988–999. 694

57. Rezaei-Ghaleh N, Parigi G, Zweckstetter M. Reorientational Dynamics of 695
    Amyloid-β from NMR Spin Relaxation and Molecular Simulation. The Journal of 696
    Physical Chemistry Letters. 2019;10(12):3369–3375. 697

58. Camilloni C, Cavalli A, Vendruscolo M. Replica-Averaged Metadynamics. J 698
    Chem Theory Comput. 2013;9(12):5610–5617. 699

59. Bussi G, Donadio D, Parrinello M. Canonical sampling through velocity rescaling. 700
    J Chem Phys. 2007;. 701

60. Ryan VH, Dignon GL, Zerze GH, Chabata CV, Silva R, Conicella AE, et al. 702
    Mechanistic View of hnRNPA2 Low-Complexity Domain Structure, Interactions, 703
    and Phase Separation Altered by Mutation and Arginine Methylation. Molecular 704
    Cell. 2018;69(3):465–479.e7. 705

61. Battiste JL, Wagner G. Utilization of site-directed spin labeling and 706
    high-resolution heteronuclear nuclear magnetic resonance for global fold 707
    determination of large proteins with limited nuclear overhauser effect data. 708
    Biochemistry. 2000;39(18):5355–5365. 709

62. Gowers R, Linke M, Barnoud J, Reddy T, Melo M, Seyler S, et al. MDAnalysis: 710
    A Python Package for the Rapid Analysis of Molecular Dynamics Simulations. In: 711
    Proceedings of the 15th Python in Science Conference. SciPy; 2016. p. 98–105. 712
    Available from: https://doi.org/10.25080/majora-629e541a-00e. 713

63. Kohl NE, Emini EA, Schleif WA, Davis LJ, Heimbach JC, Dixon RA, et al. 714
    Active human immunodeficiency virus protease is required for viral infectivity. 715
    Proc Natl Acad Sci U S A. 1988;85(13):4686–4690. 716

64. Blackburn ME, Veloro AM, Fanucci GE. Monitoring inhibitor-induced 717
    conformational population shifts in HIV-1 protease by pulsed EPR spectroscopy. 718
    Biochemistry. 2009;48(37):8765–8767. 719

65. Roche J, Louis JM, Bax A. Conformation of inhibitor-free HIV-1 protease derived from NMR spectroscopy in a weakly oriented solution. ChemBioChem. 2015;16(2):214–218.

66. Roche J, Louis JM, Bax A, Best RB. Pressure-induced structural transition of mature HIV-1 Protease from a combined NMR/MD simulation approach. Proteins. 2015;83(12):2117–2123.

67. Hornak V, Okur A, Rizzo RC, Simmerling C. HIV-1 protease flaps spontaneously close to the correct structure in simulations following manual placement of an inhibitor into the open state. J Am Chem Soc. 2006;128(9):2812–2813.

68. Huang X, Britto MD, Kear-Scott JL, Boone CD, Rocca JR, Simmerling C, et al. The Role of Select Subtype Polymorphisms on HIV-1 Protease Conformational Sampling and Dynamics. J Biol Chem. 2014;289(24):17203–17214.

69. Liu Z, Casey TM, Blackburn ME, Huang X, Pham L, de Vera IMS, et al. Pulsed EPR characterization of HIV-1 protease conformational sampling and inhibitor-induced population shifts. Physical Chemistry Chemical Physics. 2016;18(8):5819–5831.

70. Sayer JM, Liu F, Ishima R, Weber IT, Louis JM. Effect of the Active Site D25N Mutation on the Structure, Stability, and Ligand Binding of the Mature HIV-1 Protease. Journal of Biological Chemistry. 2008;283(19):13459–13470.

71. Munshi S, Chen Z, Li Y, Olsen DB, Fraley ME, Hungate RW, et al. Rapid X-ray diffraction analysis of HIV-1 protease–inhibitor complexes: inhibitor exchange in single crystals of the bound enzyme. Acta Crystallographica Section D Biological Crystallography. 1998;54(5):1053–1060.

72. Spinelli S, Liu QZ, Alzari PM, Hirel PH, Poljak RJ. The three-dimensional structure of the aspartyl protease from the HIV-1 isolate BRU. Biochimie. 1991;73(11):1391 – 1396.

73. Martin P, Vickrey JF, Proteasa G, Jimenez YL, Wawrzak Z, Winters MA, et al. "Wide-Open" 1.3 Å Structure of a Multidrug-Resistant HIV-1 Protease as a Drug Target. Structure. 2005;13(12):1887–1895.

74. Logsdon BC, Vickrey JF, Martin P, Proteasa G, Koepke JI, Terlecky SR, et al. 749
    Crystal Structures of a Multidrug-Resistant Human Immunodeficiency Virus 750
    Type 1 Protease Reveal an Expanded Active-Site Cavity. Journal of Virology. 751
    2004;78(6):3123–3132. 752

75. Camilloni C, Vendruscolo M. A tensor-free method for the structural and 753
    dynamical refinement of proteins using residual dipolar couplings. J Phys Chem 754
    B. 2015;119(3):653–661. 755

76. Heaslet H, Rosenfeld R, Giffin M, Lin YC, Tam K, Torbett BE, et al. 756
    Conformational flexibility in the flap domains of ligand-free HIV protease. Acta 757
    Crystallographica Section D Biological Crystallography. 2007;63(8):866–875. 758

77. Huang X, Britto MD, Kear-Scott JL, Boone CD, Rocca JR, Simmerling C, et al. 759
    The Role of Select Subtype Polymorphisms on HIV-1 Protease Conformational 760
    Sampling and Dynamics. Journal of Biological Chemistry. 761
    2014;289(24):17203–17214. 762

78. Eriksson AE, Baase WA, Wozniak JA, Matthews BW. A cavity-containing 763
    mutant of T4 lysozyme is stabilized by buried benzene. Nature. 764
    1992;355(6358):371–373. 765

79. Kato H, Vu ND, Feng H, Zhou Z, Bai Y. The folding pathway of T4 lysozyme: 766
    an on-pathway hidden folding intermediate. J Mol Biol. 2007;365(3):881–891. 767

80. Wang Y, Papaleo E, Lindorff-Larsen K. Mapping transiently formed and sparsely 768
    populated conformations on a complex energy landscape. eLife. 2016;5. 769

81. Mulder FA, Hon B, Muhandiram DR, Dahlquist FW, Kay LE. Flexibility and 770
    ligand exchange in a buried cavity mutant of T4 lysozyme studied by 771
    multinuclear NMR. Biochemistry. 2000;39(41):12614–12622. 772

82. Bouvignies G, Vallurupalli P, Hansen DF, Correia BE, Lange O, Bah A, et al. 773
    Solution structure of a minor and transiently formed state of a T4 lysozyme 774
    mutant. Nature. 2011;477(7362):111–114. 775

83. Lerch MT, López CJ, Yang Z, Kreitman MJ, Horwitz J, Hubbell WL. Structure-relaxation mechanism for the response of T4 lysozyme cavity mutants to hydrostatic pressure. Proc Natl Acad Sci U S A. 2015;112(19):E2437–46.

84. Eriksson AE, Baase WA, Zhang XJ, Heinz DW, Blaber M, Baldwin EP, et al. Response of a protein structure to cavity-creating mutations and its relation to the hydrophobic effect. Science. 1992;255(5041):178–183.

85. Tribello GA, Bonomi M, Branduardi D, Camilloni C, Bussi G. PLUMED 2: New feathers for an old bird. Computer Physics Communications. 2014;185(2):604–613.

86. Abraham MJ, Murtola T, Schulz R, Páll S, Smith JC, Hess B, et al. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX. 2015;1-2:19–25.

87. Cavalli A, Camilloni C, Vendruscolo M. Molecular dynamics simulations with replica-averaged structural restraints generate structural ensembles according to the maximum entropy principle. J Chem Phys. 2013;138(9).

88. Sayer JM, Liu F, Ishima R, Weber IT, Louis JM. Effect of the active site D25N mutation on the structure, stability, and ligand binding of the mature HIV-1 protease. J Biol Chem. 2008;283(19):13459–13470.

89. Berendsen HJC, Postma JPM, van Gunsteren WF, DiNola A, Haak JR. Molecular dynamics with coupling to an external bath. The Journal of Chemical Physics. 1984;81(8):3684–3690.

90. Lindorff-Larsen K, Piana S, Palmo K, Maragakis P, Klepeis JL, Dror RO, et al. Improved side-chain torsion potentials for the Amber ff99SB protein force field. Proteins. 2010;78(8):1950–1958.

91. Best RB, Hummer G. Optimized Molecular Dynamics Force Fields Applied to the Helix−Coil Transition of Polypeptides. J Phys Chem B. 2009;113(26):9004–9015.

92. Darden T, York D, Pedersen L. Particle mesh Ewald: An N·log(N) method for Ewald sums in large systems. The Journal of Chemical Physics. 1993;98(12):10089–10092.

93. Bussi G, Donadio D, Parrinello M. Canonical Sampling Through Velocity
    Rescaling. J Chem Phys. 2007;126(1):014101.

94. Hess B. P-LINCS: A Parallel Linear Constraint Solver for Molecular Simulation.
    Journal of Chemical Theory and Computation. 2008;4(1):116–122.

805
806
807
808