Title

Neonatal LPS exposure reduces ATP8A2 level in the prefrontal cortex in mice via increasing IFN-γ level

Authors

Jiapeng Deng (ORCID: 0000-0002-0332-9296)[a] , Linyang Song (ORCID: 0000-0002-2770-9568)[b], Zhiqin Yang (ORCID: 0000-0002-3120-5944)[c], Sixie Zheng (ORCID: 0000-0001-6603-7947)[d], Zhuolin Du (ORCID: 0000-0003-0928-4081)[d], Li Luo[b,e], Jing Liu[b], Xiaobao Jin[e], Junhua Yang (ORCID: 0000-0001-8037-9644)[b,e*]

Affiliations

[a] Class 1, Grade 2018, School of Clinical Medicine, Guangdong Pharmaceutical University, Guangzhou, Guangdong, People's Republic of China.

[b] Department of anatomy, School of Biosciences & Biopharmaceutics, Guangdong Pharmaceutical University, Guangzhou, Guangdong, People's Republic of China.

[c] Aviation health center, China southern airlines company limited, Guangzhou, Guangdong, People's Republic of China.

[d] Class 4, Grade 2018, School of Clinical Medicine, Guangdong Pharmaceutical University, Guangzhou, Guangdong, People's Republic of China.

[e] Guangdong Key Laboratory of Pharmaceutical Bioactive Substances, Guangdong

Pharmaceutical University, Guangzhou, Guangdong, People's Republic of China.

Corresponding author

[*]Correspondence should be addressed to Junhua Yang at the following address, phone,

and email address:

Address: Junhua Yang, Room 510#, Building of Basic Medical Sciences, School of

Biosciences & Biopharmaceutics, Guangdong Pharmaceutical University, No. 230,

East Waihuan Street, Higher Education Mega Center, Guangzhou City, Guangdong

Province, Postcode 510006, P. R. China.

Tel.: + 86 20 39352189.

E-mail: jhyang2018@gdpu.edu.cn.

Abstract

Neonatal lipopolysaccharide (LPS) exposure can cause depressive-like behaviors in rodents involving elevated interferon (IFN)-γ. Studies have linked down-regulation of prefrontal cortex (PFC) ATPase phospholipid transporting 8A2(ATP8A2) expression to depressive-like behaviors. In non-neuronal cells, IFN-γ could reduce ATP8A2 expression. We therefore hypothesized that neonatal LPS exposure might induce PFC ATP8A2 down-regulation by increasing IFN-γ level. Here, C57BL6/J mice of both sexes received 3-dose-injections of LPS (50μg/kg bodyweight, i.p.) on postnatal day (PND)5, PND7 and PND9. LPS-treated mice showed a transiently decreased PFC ATP8A2 expression. Moreover, a negative correlation of PFC ATP8A2 expression was found with IFN-γ level. Using neutralizing mAb, IFN-γ was identified as the key mediator of LPS-induced PFC ATP8A2 decrease. Besides, neutralizing IFN-γ during neonatal LPS exposure attenuated the depressive-like behaviors in adulthood. In sum, neonatal LPS exposure reduced ATP8A2 level in PFC in mice via increasing IFN-γ level, maybe associated with mechanism underlying LPS-induced brain and behavior impairments.

Keywords: Immune activity; Cytokine; P4-ATPase; Brain; Depression

Introduction

ATP8A2, a member of the P4-ATPase family in mammals, is highly expressed in the brain, spinal cord, testis, and retina, which flipping phosphatidylserine and phosphatidylethanolamine across the cell membrane (Andersen et al.,2016). It plays an important role in maintaining the stability and normal function of the cell membrane (Coleman et al.,2009). Therefore, ATP8A2 in the brain is vital for normal neural development and function (Coleman and Molday,2011; Xu et al.,2012), since these processes involve cell proliferation, migration, synapses pruning of neurons (Brandon and Sawa,2011). ATP8A2 expressed in the brain has received extensive attention recently and researches have shown that reduced expression of ATP8A2 leads to axonal mutation and neurodegenerative diseases (Choi et al.,2019; Zhu et al.,2012). There is also research reporting that ATP8A2 expression decreases in conditions of Alzheimer's disease and bacterial infection (Aaron et al.,2018; Ross et al.,2011). Moreover, depressive-like behavior has been reported to be associated with decreased ATP8A2 expression in the prefrontal cortex (PFC) (Chen et al.,2017). Although ATP8A2 has been widely studied, little is known about how ATP8A2 expression is regulated.

Recently, IFN-γ was shown to significantly reduce the expression of ATP8A2 in non-neuron cells (Shulzhenko et al.,2018). IFN-γ produced in the periphery can penetrate brain parenchyma across the immature blood-brain barrier and even increase the permeability of the mature blood-brain barrier (Hansen-Pupp et al.,2005; Wong et

al.,2004). IFN-γ may mediate the neonatal lipopolysaccharide (LPS)

exposure-induced depressive-like behaviors in mice (Campos et al.,2014), which is

associated with the function of PFC. Besides, IFN-γ expression in PFC increased

more than twice after LPS treatment (Laumet et al.,2018). Based on the background

mentioned above, it was postulated that neonatal LPS exposure might induce ATP8A2

down-regulation in PFC in mice by increasing the IFN-γ level. There is yet no report

addressing this issue.

To test this hypothesis, we challenged neonatal mice with LPS and observed a

transiently decreased expression of ATP8A2 in PFC. Using a series of experiments,

we further identified IFN-γ as the key mediator of LPS-induced ATP8A2

down-regulation in PFC in mice. These findings may reveal a potential mechanism by

which early inflammation leads to impairment of the central nervous system

development and function.

2. Materials and methods

2.1. Animals designing

Litters of newborn C57BL6/J mice (one-day-old) consisting approximately evenly

of both sexes were ordered together with their mothers from Guangdong Medical

Laboratory Animal Center (Guangzhou, China). They were housed in a specific

pathogen-free condition under 12 h light-12 h dark conditions with food and water

available ad libitum.

In this study, a randomized block design was used to minimize the effects of

systematic error according to publications by statisticians who stated that if the

experimenters aim to focus exclusively on the differences between/among different

treatment conditions, the effects due to variations between the different blocks should

be eliminated by using a randomized block design ANOVA (Refer to

https://doi.org/10.1007/978-0-387-32833-1_344;http://www.r-tutor.com/elementary-st

atistics/analysis-variance/randomized-block-design). In such design, there is only one

primary factor under consideration in the experiment. Similar test subjects are

grouped into blocks. Then, subjects within each block are randomly assigned to

treatment conditions. Each block is tested against all treatment levels of the primary

factor at random order. Thus, possible influence by other extraneous factors will by

eliminated.

To be specific, a randomized block design was conducted for each of the

experiments for investigating potential differences between or among treatment

conditions. Each of the blocks consisted of two (for experiments reported in Fig.1 and

Fig.2), three (for experiment reported in Fig.5) or five (for experiments reported in

Fig.4 and Fig.6) same-sex pups from a same dam. The pups in each block were

randomly assigned to different treatment conditions (only one pup in each block for

receiving each kind of the different treatment conditions). The sex of n pups in

different blocks were same or different, while the litter background of pups in

different blocks were certainly different The detailed information indicating how

many litters were used and how many pups from each litter were included in each

group of each experiment in this study were shown in Supplementary talbe1 to table6

(Supplementary Material). Pups subjected to behavioral tests were weaned on

postnatal day (PND)21. Pups subjected to other tests were sacrificed on PND11. This

study was approved by the Institutional Animal Ethics Committee of Guangdong

Pharmaceutical University.

2.2. LPS treatment

Mice were intraperitoneally injected with 3 doses of LPS dissolved in 0.1 mol

phosphate balanced solution (PBS) (Escherichia coli O111: B4; Sigma-Aldrich) (50

µg/kg bodyweight for each dose;) on PND5, PND7 and PND9. The dosage and

schedule for LPS injection were determined based on previous studies (Dinel et

al.,2014; Doosti et al.,2013; Liang et al.,2019). The experiment was controlled by

intraperitoneal injection of PBS. Littermates were toe-clipped for identification

randomly assigned to each of the groups in each experiment as described in Results.

2.3 Administration of anti-IFN-γ neutralizing mAb and an isotype IgG1

For experiments conducted in this study as reported in Section 3.3 and Fig.4 to

Fig.6, anti-IFN-γ neutralizing mAb and/or an isotype IgG1 (Invitrogen, Thermo

Fisher Scientific, Waltham, MA, USA) were intraperitoneally injected daily to mice

from PND5 to PND10. By a special experiment, the optimal dosage was first

determined as 0.6 mg/kg body weight of anti-IFN-γ neutralizing mAb. The dosage of

isotype IgG1 was determined as 0.6 mg/kg body weight accordingly.

2.4 Western blot

After being over-anesthetized with 10% chloral hydrate, the mice were decapitated

and killed. The mouse PFC tissues were immediately taken on ice, and then

homogenized in RIPA lysate containing protease inhibitors (Beyotime, Wuhan,

China). After centrifuged at 13000g for 30 min at 4°C, the supernatant was taken. The

BCA protein analysis method was used to determine the total protein content of the

sample. After mixed with 5×SDS-PAGE loading buffer at the volume ratio of V

(protein sample solution): V (5×SDS-PAGE Loading Buffer) = 4:1, the samples were

boiled and denatured for 5 min, save for later use. The samples of each group were

subjected to SDS polyacrylamide gel electrophoresis. The gel used here is a

simplified hand-poured gradient gel, in which the ratio was 40% of the volume being

6% acrylamide and 60% of the volume being 12% acrylamide (using an SDS-PAGE

Gel Quick Preparation Kit, Beyotime, Shanghai, China). The procedure to make this

gradient gel was according to a previous study (Miller et al.,2016) with modification

with respect to the concentrations of acrylamide. When proteins were transferred to

PVDF membranes, the membranes were cut according to the indication of the color

marker (10-180 kDa, ThermoFisher, Shanghai, China). Two parts of each membrane,

one located in the marker's range 70-180 kDa and the other in the range 25-55 kDa,

containing the target proteins ATP8A2 (129 kDa) and the internal control proteins

β-actin (43 kDa) were subjected to the subsequent treatment. These membranes were

blocked with 5% skimmed milk powder at RT for 2 h. After this, these membranes

were put into the corresponding primary antibody incubation solutions, one

containing anti-ATP8A2 antibodies (1:500, Abcam, Cambridge, MA, USA) and the

other containing anti-β-actin antibodies (1:5000, Abcam, Cambridge, MA, USA) and

incubated at 4°C overnight. After rinsing with TBST, these membranes were put into

the corresponding secondary antibodies incubation solutions, one containing

horseradish peroxidase (HRP)-conjugated goat anti-rabbit antibodies (1:5000,

Bioworld, Atlanta, GA, USA) and the other containing HRP-conjugated goat

anti-rabbit antibodies (1:5000, Bioworld, Atlanta, GA, USA) and incubated at RT for

1 h. After rinsing with TBST, ECL luminescent solution is added on the membranes

in a dark room for exposure and development. Chemiluminescent images were

obtained with Carestream XBT X-ray Film (Rayco, Xiamen, Fujian, China), and

subsequently scanned and quantified by densitometry using ImageJ software.

2.5 Determination of cytokines levels

Forty-eight h after the last LPS injection, the mice were anesthetized deeply with

10% chloral hydrate before the blood was collected from the heart immediately. After

blood collection, it was allowed to stand at room temperature for 1 h, and after

centrifugation, the supernatant was taken and stored at -70°C for the experiment. Then

the mice were transcranial perfused with 0.9% NaCl and PFC was collected

immediately on ice. A mouse cytokine/chemokine magnetic bead panel

(MCYTOMAG-70K-06; Millipore, Billerica, MA, USA) was employed to detect the

levels of IFN-γ, tumor necrosis factor (TNF)-α, interleukin(IL)-1β, IL-6 both in the

serum and PFC (Yang et al.,2016). Each of the serum samples was diluted at 1:2 in

assay buffer. PFC tissue was prepared as homogenates before assayed. The total

protein concentration of each sample was adjusted to 4.5 mg/ml using a BCA protein

assay kit (Beyotime, Shanghai, China). Then, the prepared serum and PFC samples

were used strictly according to the manufacturer's protocols for the multiplex assays.

The data were collected on a Bio-Plex-200 system (Bio-Rad, Hercules, CA, USA) and

analyzed using professional software (Bio-Plex Manager).

2.6 Immunofluorescence and cell quantification

Forty-eight h after the last LPS injection, the mice were over-anesthetized with 10%

chloral hydrate and transcranial perfused with 0.9% NaCl followed by 4%

paraformaldehyde (PFA). After removed, the brains were immediately post-fixed in 4%

PFA overnight at 4 °C. Then, the brains were gradient dehydrated with 10%, 20%,

and 30% sucrose for 24 h each at 4°C. Free-floating, serial coronal sections (40 μm)

were collected on a Leica SM2000R freezing microtome (Leica Microsystems,

Richmond Hill, Ontario, Canada) and stored at 4°C for immunostaining. Sections

were washed in PBS three times and then blocked in PBS containing 1% bovine

serum albumin (BSA) and 0.25% Triton X-100 (Sigma-Aldrich, St. Louis, MO, USA)

for 1 h at 37°C. The slices were then incubated in the primary antibodies overnight at

4°C. The primary antibodies, including rabbit anti-ATP8A2 (1:200; Abcam,

Cambridge, MA, USA) and mouse anti-NeuN (1:1000; Abcam, Cambridge, MA,

USA), were diluted in PBS containing 1% BSA and 0.25% Triton X-100. Afterward,

the specimens were washed three times in PBS and then were incubated with

secondary antibodies, including Alexa Fluor 555-conjugated goat anti-rabbit and

Alexa Fluor 488-conjugated donkey anti-mouse for 2 h at 37°C. Both secondary

antibodies (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA) were diluted

to 1:400.

A Stereo Investigator stereological system (MicroBrightField, Williston, USA) was

used for quantitative analyses of the ATP8A2$^+$ cells in the unilateral PFC of each

mouse. Coronal sections of the PFC were collected through the rostrocaudal axis

spanning approximately from +3.20 mm to +2.10 mm relative to bregma (Xiong et

al.,2017)to count the interested cells. Measurements were recorded from an

equidistant series of six coronal sections. After the actual section thickness was

measured, appropriate guard zones at the top and bottom of each section were defined

to avoid oversampling. The 40× objective of a Nikon microscope was used for all

stereological analyses. The numbers of ATP8A2$^+$/NeuN$^+$ and NeuN$^+$ cells within the

traced region (PFC) in each of the six selected sections were straightly quantified

without using a grid or counting frames. A Zeiss LSM780 confocal laser-scanning

microscope (Carl Zeiss AG, Oberkochen, Germany) was used to capture the

representative confocal micrographs of the labeled cells.

For ATP8A2 in situ densitometric analysis, one 40-μm-thick coronal section at the

middle of the PFC (+2.10 mm relative to bregma) was collected from each animal.

Single labeling for ATP8A2 immunofluorescently was performed, rather than double

labeling for ATP8A2/NeuN, to minimize the possible interference with ATP8A2

positive immunofluorescence signal intensity during staining practice. A Zeiss

LSM780 confocal laser-scanning microscope (Carl Zeiss AG, Oberkochen, Germany)

was used to capture the micrographs of the same field as the double-labeled

micrographs shown in Fig.6D-F. The ATP8A2 signal was measured as a mean gray

value using the software ImageJ and the data were shown in Fig.6 as a relative

influence. The procedures were: 1) Image-Type-8bit; 2) Image-Adjust-Threshold; 3)

Image-Adjust-Auto Threshold; 4) Analyze-Set Measurements; 5) Analyze-Measure.

2.7. Forced swim test (FST)

FST is a standard test used as a screen for measuring depressive-like behaviors in

mice. The experiment was carried out on the PND90, the mice were brought to the

laboratory one week in advance to let them adapt to the environmental room. In this

experiment, mice were placed in a plastic cylinder (40 cm in-depth, 20 cm in diameter

and filled with water at 24±°C with the water's height at 25 cm above from the

bottom. Each mouse was put in the plastic cylinder in the same way and forced to

swim for 6 min in the plastic cylinder. The first two minutes of mouse behavior was

not recorded. The following 4 minutes were recorded by a video tracking system

EthoVision (Noldus Information Technology B.V., Wageningen, Netherlands) for the

analysis of the total time spent by each animal in staying on the water surface without

any struggling or swimming during the test. At the end of the experiment, the hair of

the mice was blown dry to prevent them from catching a cold.

2.8 Tail-suspension test (TST)

TST is also one of the most commonly used experiments to evaluate

depressive-like behavior in rodent models. In the present study, TST was performed

on the second day after the FST. During this test, each mouse was held below its tail

to the edge of the clip and suspended 60 cm above from the floor. The performances

of mice were recorded by a video tracking system EthoVision (Noldus Information

Technology B.V., Wageningen, Netherlands) four minutes with a 2-min-gap left prior

to the recording. The total time spent by each animal in not struggling during the test

was analyzed. After all behavioral tests were done, animals were killed by

over-anesthetized with 10% chloral hydrate.

2.9 Statistical analyses

The data were statistically analyzed using the SPSS 25.0 statistical software

(Chicago, IL, USA). *Pearson*'s correlation analysis was used for data shown in Fig.3.

Data shown in Fig.1, Fig.2, Fig.4, Fig.5 and Fig.6 were statistically analyzed using a

randomized block design ANOVA to account for the effects of treatment (such as LPS

injection and/or anti-IFN-γ neutralizing mAb injection) by modeling both litter factor

and sex factor as a fixed effect described as "block effect" in section Results and the

treatment factor as a random effect described as "treatment effect" in section Results.

For data obtained from more than two treatment groups, the performance of

randomized block design ANOVA were followed by Tukey's *post hoc* test. *p* values

are displayed as follows: n.s. = not significant, * *p* < 0.05, ** *p* < 0.01, *** *p* < 0.001.

3. Results

3.1 Neonatal LPS exposure induced a transiently down-regulated expression of

ATP8A2 in the PFC in mice

To determine whether neonatal LPS exposure influence the expression of ATP8A2

in the PFC in mice and how long this potential influence would last following LPS

exposure, the first experiment was carried out. Mice in the LPS group were injected

with LPS and those in the CON group with PBS of the same volume. Two groups of

mice were sacrificed after 1 day, 2 days, 4 days, or 10 days after the last LPS injection.

The left unilateral PFC tissue from every mouse was taken on ice to be homogenate

immediately for Western blot analyses of ATP8A2 level. The right unilateral PFC

tissue was taken on ice and stored at -70°C at once for ELISA analyses of

LPS-induced pro-inflammatory cytokines if necessary.

As shown in Fig.1, the ATP8A2 levels significantly decreased at the former three

test time points: 1 day (randomly block design ANOVA, block effect: $F_{(5,5)} = 1.120$, *p*

$= 0.452$, $n = 6$; treatment effect: $F_{(1,5)} = 44.835$, $p = 0.001$, $n = 6$), 2 days (randomly

block design ANOVA, block effect: $F_{(5,5)} = 1.369$, $p = 0.369$, $n = 6$; treatment effect:

$F_{(1,5)} = 102.221$, $p < 0.001$, $n = 6$) and 4 days (randomly block design ANOVA, block

effect: $F_{(5,5)} = 0.811$, $p = 0.588$, $n = 6$; treatment effect: $F_{(1,5)} = 37.424$, $p = 0.002$, $n =$

6) after the last LPS injection. However, such decrease was no longer detectable 10

days after the last LPS injection (randomly block design ANOVA, block effect: $F_{(5,5)}$

$= 0.489$, $p = 0.775$, $n = 6$; treatment effect: $F_{(1,5)} = 0.078$, $p = 0.791$, $n = 6$) (Fig.1). It

is worth noting that the largest extent of the decrease in PFC ATP8A2 expression was

observed in the sample obtained 48 h after the last LPS injection (Fig.1). These

findings verified the expected influence of neonatal LPS exposure on the PFC

ATP8A2 expression in mice and revealed the time curve of PFC ATP8A2 expression

following neonatal LPS exposure.

3.2 PFC ATP8A2 level was only significantly correlated to IFN-γ level among all four

elevated pro-inflammatory cytokines both in serum and in the PFC

Given the initial finding that PFC ATP8A2 expression was reduced to the largest

extent 48 h after the last LPS injection (Fig.1), the prepared serum and PFC samples

($n = 6$) obtained at the same time point in the last experiment were then subjected to

ELISA analyses for the levels of LPS-induced pro-inflammatory cytokines, including

IFN-γ, IL-1β, IL-6 and TNF-α.

Moreover, another experiment was performed in which the animal was treated all

the same way and the levels of PFC ATP8A2 as well as the levels pro-inflammatory

cytokines both in the serum and PFC were detected specifically 48 h after the last LPS

injection, using a larger sample size ($n = 9$) so as both to provide a repetitive

verification of the very novel finding of ATP8A2 in the brain and to make the total

sample size up to $n = 15$ (together with $n = 6$) for this chosen test time point.

Pearson's correlation analysis between PFC ATP8A2 level and the level of each of

the LPS-induced pro-inflammatory cytokines would get good confidence in the case

of $n = 15$.

As shown in Fig.2A-B, the PFC ATP8A2 level decreased by more than 90% in LPS

group (randomly block design ANOVA, block effect: $F_{(8,8)} = 1.062$, $p = 0.467$, $n = 9$;

treatment effect: $F_{(1,8)} = 112.127$, $p < 0.001$, $n = 9$). All the detected cytokines,

whether in serum (Fig.2C-F) or in PFC (Fig.2G-J), showed a dramatically increased

expression (serum IFN-γ: randomly block design ANOVA, block effect: $F_{(8,8)} = 2.310$,

$p = 0.129$, $n = 9$; treatment effect: $F_{(1,8)} = 68.475$, $p < 0.001$, $n = 9$; serum IL-6:

randomly block design ANOVA, block effect: $F_{(8,8)} = 0.983$, $p = 0.509$, $n = 9$;

treatment effect: $F_{(1,8)} = 170.514$, $p < 0.001$, $n = 9$; serum IL-1β: randomly block

design ANOVA, block effect: $F_{(8,8)} = 1.485$, $p = 0.295$, $n = 9$; treatment effect: $F_{(1,8)} =$

104.142, $p < 0.001$, $n = 9$. serum TNF-α: randomly block design ANOVA, block

effect: $F_{(8,8)} = 0.928$, $p = 0.541$, $n = 9$; treatment effect: $F_{(1,8)} = 133.012$, $p < 0.001$, $n =$

9; PFC IFN-γ: randomly block design ANOVA, block effect: $F_{(8,8)} = 0.722$, $p = 0.672$,

$n = 9$; treatment effect: $F_{(1,8)} = 107.626$, $p < 0.001$, $n = 9$; PFC IL-6: randomly block

design ANOVA, block effect: $F_{(8,8)} = 1.001$, $p = 0.499$, $n = 9$; treatment effect: $F_{(1,8)} =$

129.968, $p < 0.001$, $n = 9$; PFC IL-1β: randomly block design ANOVA, block effect:

$F_{(8,8)} = 1.130$, $p = 0.433$, $n = 9$; treatment effect: $F_{(1,8)} = 42.304$, $p < 0.001$, $n = 9$; PFC

TNF-α: randomly block design ANOVA, block effect: $F_{(8,8)} = 1.159$, $p = 0.420$, $n = 9$;

treatment effect: $F_{(1,8)} = 250.424$, $p < 0.001$, $n = 9$)

Taken all the data from 48 h time point samples from LPS-treated mice together,

Pearson's correlation analysis was performed between PFC ATP8A2 level and the

level of each of the LPS-induced pro-inflammatory cytokines ($n = 15$). There was a

significant negative correlation between ATP8A2 expression in PFC and IFN-γ level

both in serum ($p < 0.01$, *Pearson*'s correlation analysis, $n = 15$) (Fig.3 A) and in PFC

($p < 0.001$, *Pearson*'s correlation analysis, $n = 15$) (Fig.3E). No significant

correlations were found between ATP8A2 level and all the other pro-inflammatory

cytokines (all p values > 0.05) (Fig.3 B-D, F-H). These results suggest that IFN-γ

plays an important role in the PFC ATP8A2 down-regulation caused by neonatal LPS

exposure.

3.3 IFN-γ mediated the PFC ATP8A2 down-regulation caused by neonatal LPS

exposure

To further explore the potential role of IFN-γ in mediating the PFC ATP8A2

down-regulation following neonatal LPS exposure, an IFN-γ-blocking experiment

was conducted using anti-IFN-γ neutralizing mAb and an isotype IgG1. Before this

IFN-γ-blocking experiment, another experiment was carried out to determine the

optimal dosage of anti-IFN-γ neutralizing mAb. The dosage of 0.6 mg/kg body weight

restored the IFN-γ levels to normal physiological levels both in serum (randomly

block design ANOVA, block effect: $F_{(7,35)} = 0.449$, $p = 0.864$, $n = 8$; treatment effect:

$F_{(5,35)} = 78.511$, $p < 0.001$, $n = 8$; *post hoc test*, LPS+anti-IFN-γ(0.6) group vs. CON

group: $p = 1.000$) (Fig.4A) and in PFC (randomly block design ANOVA, block effect:

$F_{(7,35)} = 0.718$, $p = 0.657$, $n = 8$; treatment effect: $F_{(5,35)} = 35.581$, $p < 0.001$, $n = 8$;

post hoc test, LPS+anti-IFN-γ(0.6) group vs. CON group: $p = 0.821$) (Fig.4B). Then,

we proceeded to the IFN-γ-blocking experiment and five groups of mice were set. To

be specific, the CON group and LPS group were set as above. LPS+anti-IFN-γ(0.6)

group was set by injecting LPS combined with anti-IFN-γ neutralizing mAb. The

LPS+IgG1 group was set by injecting LPS combined with an isotype IgG1. The last

group was set by injecting mere anti-IFN-γ neutralizing mAb (anti-IFN-γ(0.6)).

Randomly block design ANOVA revealed significant differences in IFN-γ levels

among five groups (Fig.4C, serum IFN-γ level: randomly block design ANOVA,

block effect: $F_{(7,28)} = 0.611$, $p = 0.742$, $n = 8$; treatment effect: $F_{(4,28)} = 52.663$, $p <$

0.001, $n = 8$) (Fig.4D, PFC IFN-γ level: randomly block design ANOVA, block effect:

$F_{(7,28)} = 0.276$, $p = 0.958$, $n = 8$; treatment effect: $F_{(4,28)} = 95.873$, $p < 0.001$, $n = 8$).

We first confirmed that administration of anti-IFN-γ neutralizing mAb blocked the

increase of IFN-γ levels both in systemic blood (LPS+anti-IFN-γ(0.6) group vs. CON

group: $p = 0.998$) (Fig.4C) and PFC (LPS+anti-IFN-γ(0.6) group vs. CON group: $p =$

0.903) (Fig. 4D), while isotype IgG1 failed to block the LPS-induced IFN-γ increase

either in serum (LPS+IgG1 group vs. CON group: $p < 0.001$; LPS+IgG1 group vs.

LPS group: $p = 0.976$) (Fig.4C) or in PFC (LPS+IgG1 group vs. CON group: $p <$

0.001; LPS+IgG1 group vs. LPS group: $p = 0.563$) (Fig.4D). In the absence of LPS

exposure, the anti-IFN-γ neutralizing mAb significantly neutralized the physiological

IFN-γ both in blood (anti-IFN-γ(0.6) group vs. CON group: $p = 0.013$; LPS+IgG1

group vs. LPS group: $p = 0.976$) (Fig.4C) and in brain of mice (anti-IFN-γ(0.6) vs.

CON group: $p < 0.001$; LPS+IgG1 group vs. LPS group: $p = 0.563$) (Fig.4D).

Randomly block design ANOVA of Western blot data revealed significant

differences in PFC ATP8A2 levels among five groups (Fig.4E, randomly block design

ANOVA, block effect: $F_{(7,28)} = 1.274$, $p = 0.299$, $n = 8$; treatment effect: $F_{(4,28)} = $

122.680, $p < 0.001$, $n = 8$). We found that neutralization of IFN-γ blocked the

LPS-induced ATP8A2 decrease in PFC (LPS+anti-IFN-γ(0.6) group vs. CON group:

$p = 1$) (Fig.4E-F), while the isotype IgG1 showed no significant blocking effect on the

LPS-induced ATP8A2 decrease in PFC (LPS+IgG1 group vs. LPS group: $p = 0.922$;

LPS+IgG1 group vs. LPS+anti-IFN-γ(0.6) group: $p < 0.001$) (Fig.4E-F). Interestingly,

in the absence of LPS exposure, mere anti-IFN-γ neutralizing mAb administration led

to no significant influence on the PFC ATP8A2 level (anti-IFN-γ(0.6) group vs. CON

group: $p = 1$) (Fig.4E-F), although it neutralized the physiological IFN-γ both in the

blood (Fig.4C) and in the brain of mice (Fig.4D).

We next performed another experiment to observe the role of IFN-γ using

immunofluorescence detection, with a total of three groups set, CON group, LPS

group, and LPS+anti-IFN-γ(0.6) group. The groups by giving isotype IgG1 or mere

anti-IFN-γ neutralizing mAb were no longer set due to the confirmed results shown

above. The results of randomly block design ANOVA of immunofluorescence

detection of PFC ATP8A2 levels showed similar findings to Western blot.

Neutralization of IFN-γ blocked the LPS-induced ATP8A2 decrease in PFC, indicated

by the mean fluorescence intensity (randomly block design ANOVA, block effect:

$F_{(5,10)} = 0.473$, $p = 0.788$, $n = 6$; treatment effect: $F_{(2,10)} = 47.276$, $p < 0.001$, $n = 6$;

post hoc test, LPS+anti-IFN-γ(0.6) group vs. CON group: $p = 0.849$) (Fig.5K) and

ATP8A2[+]/NeuN[+] cells (randomly block design ANOVA, block effect: $F_{(5,10)} = 0.121$,

$p = 0.985$, $n = 6$; treatment effect: $F_{(2,10)} = 88.596$, $p < 0.001$, $n = 6$; *post hoc test*,

LPS+anti-IFN-γ(0.6) group vs. CON group: $p = 0.532$) (Fig.5L). Besides, there were

no significant differences among three groups in the number of neurons (NeuN[+] cells)

in PFC (data are not shown) although nearly all patchy ATP8A2[+] signals were

co-located with NeuN[+] signals (Fig.5G-I). The findings shown in this section

identified IFN-γ as the key mediator of the PFC ATP8A2 down-regulation caused by

neonatal LPS exposure. Moreover, ATP8A2 alterations were not accompanied by a

detectable change in the number of neurons (NeuN[+] cells) in PFC.

3.4 IFN-γ plays a role in depressive-like behaviors in adulthood induced by neonatal

LPS exposure

Abnormality in PFC, especially during the critical development period, is an

important pathophysiological basis of depression and it has been verified that

neonatal LPS exposure could result in depression in adulthood (Dinel et al.,2014;

Walker et al.,2013). Therefore, we investigated whether neutralization of IFN-γ

affects the behavioral performances in the FST and TST of the LPS-treated mice. The

floating immobility time was partially but significantly restored toward the levels of

CON group in FST task (randomly block design ANOVA, block effect: $F_{(11,44)} = 0.883$,

$p = 0.563$, $n = 12$; treatment effect: $F_{(4,44)} = 43.424$, $p < 0.001$, $n = 12$; *post hoc test*,

LPS+anti-IFN-γ(0.6) group vs. CON group: $p = 0.002$) (Fig.6A). In TST task,

neutralization of IFN-γ also resulted in an obvious restoring trend although Tukey

post hoc test reported the *p* value > 0.05 of the comparision between

LPS+anti-IFN-γ(0.6) group and CON group (randomly block design ANOVA, block

effect: $F_{(11,44)} = 0.466$, $p = 0.914$, $n = 12$; treatment effect: $F_{(4,44)} = 32.685$, $p < 0.001$,

$n = 12$; *post hoc test*, LPS+anti-IFN-γ(0.6) group vs. CON group: $p = 0.245$) (Fig.6B).

However, an LSD *post hoc test* after the same randomly block design ANOVA

reported a significant decrease in TST floating immobility time of

LPS+anti-IFN-γ(0.6) group than CON group ($p = 0.043$).

The isotype IgG1 failed to block the LPS-induced behavioral alterations either in

FST (LPS+IgG1 group vs. CON group: p < 0.001; LPS+IgG1 group vs. LPS group: p

$= 0.769$) (Fig.6A) or in TST (LPS+IgG1 group vs. CON group: $p < 0.001$; LPS+IgG1

group vs. LPS group: $p = 0.963$) (Fig.6B). In the absence of LPS exposure, the

anti-IFN-γ neutralizing mAb led to no significant behavioral alterations in FST

(anti-IFN-γ(0.6) group vs. CON group: $p = 0.319$; LPS+IgG1 group vs. LPS group: *p*

$= 0.769$)   (Fig.6A) or in TST (anti-IFN-γ(0.6) vs. CON group: $p = 0.999$) (Fig.6B).

These results suggested a role of IFN-γ in depressive-like behaviors in adulthood

induced by neonatal LPS exposure.

3.5 No sex dimorphism in ATP8A2 levels or cytokines levels despite a sex dimorphism

in depressive-like behaviors in mice

All the analyses described above were intended to exclusively detect the effects of different treatment conditions to test the hypothesis that neonatal LPS exposure might influence PFC ATP8A2 expression in mice involving an increased IFN-γ level and therefore the sex factor was not under consideration as stated in section 2.1. However, previous studies showed that sex may affect immune activation (Cai et al.,2016). So, we also analyzed the data collected from the experiments mentioned in section 3.1 and 3.2 shown in Fig.2 as well as in the bars in Fig.1 indicated as day2 using another statistical method to observe whether there were significant differences in PFC ATP8A2 expression and the levels of proinflammatory cytokines between two sexes either in the presence of LPS challenge or not. The results showed no significant differences in the levels of these molecules between male and female animals either in the presence of LPS challenge or not (all p values > 0.05, Supplementary Fig.1). Furthermore, data collected from behavioral tests and shown in section 3.4 (Fig.6) were also subjected to additional analysis for differences between two sexes. The results showed no significant differences in the behavioral task performances of LPS+anti-IFN-γ(0.6) group between male and female animals (FST: LPS+anti-IFN-γ(0.6) group: $p = 0.536$; TST: LPS+anti-IFN-γ(0.6) group: $p = 0.827$; Supplementary Fig.2). However, in all the other groups, significant differences between two sexes were found in FST (CON group: $p < 0.001$; LPS group: $p < 0.001$;

LPS+IgG1 group: $p = 0.018$; anti-IFN-γ(0.6) group: $p = 0.013$; Supplementary Fig.2A)

and TST (CON group: $p < 0.001$; LPS group: $p < 0.029$; LPS+IgG1 group: $p < 0.001$;

anti-IFN-γ(0.6) group: $p = 0.006$; Supplementary Fig.2B) task performances with

opposite directions of alterations in the presence of LPS challenge or not. Specifically,

female mice showed higher depressive-like behaviors in the absence of LPS challenge,

whereas they showed lower depressive-like behaviors in the presence of that.

Discussion

Our research revealed that neonatal LPS exposure reduced ATP8A2 level in PFC in

mice via increasing IFN-γ level. This finding, based on a series of studies concerning

the influence of LPS on brain development and behavior (Bilbo and Schwarz,2012;

Doosti et al.,2013), is the first to report the change of ATP8A2 in the central nervous

system and to further report the mechanism underlying how LPS affects PFC ATP8A2

expression. Moreover, this study further extends our understanding of the mechanism

underlying behavioral effects of early life immune activation.

Intraperitoneal LPS injection has been shown to cause a range of acute

physiological, pathological, and psychological disorders in rodents. Impairments both

in food intake and social exploratory behavior in rodents have been demonstrated in

rodents administered intraperitoneally with LPS (Haba et al.,2012; O'Reilly et

al.,1988). It has also been reported that LPS exposure can induce depressive and

anxiety-like behaviors (Depino,2015; Doosti et al.,2013). Moreover, LPS may reduce

acutely the level of prefrontal cortical neurogenesis in adult rodents (Wang et

al.,2016). The data from the LPS-treated mice in the present study enriched these

widely reported neurobehavioral impairments.

LPS challenge may induce a large extent release of cytokines both in the periphery

and brain, particularly pro-inflammatory cytokines, including IFN-γ, IL-1β, IL-6, and

TNF-α (Klimstra et al.,1999). These pro-inflammatory cytokines not only play an

important role in immunity but also may affect brain function and mediate

disease-like behaviors. For example, these pro-inflammatory factors have been shown

to cause depressive-like behavior in mice (Gupta et al.,2016; Hashimoto,2015; Kohler

et al.,2014; Pokryszko-Dragan et al.,2012). The results of cytokines in our study

showed clearly that their levels in the periphery and the brain were highly consistent.

this fact may partially due to the immature blood-brain barrier or increased

permeability of it during inflammation.

ATP8A2 is a protein located in the membrane, functioning to transport

phosphatidylserine into the inner layer of membrane, and thus it is important to

maintain the structural stability and normal function of the membrane (Andersen et

al.,2016; Coleman et al.,2009). Since the distribution of ATP8A2 in the brain has been

determined recently (Andersen et al.,2016), little has been known about the potential

factors involved in the regulation of its expression. ATP8A2 expression

down-regulation has been observed in certain pathological conditions including

Alzheimer's disease and bacterial infection (Aaron et al.,2018; Ross et al.,2011).

However, the study reported a mechanism underlying such regulation could not be

seen so far. Decreased level of ATP8A2 in the PFC also occurs in the presence of

stress and depressive situation (Chen et al.,2017), but the causal relationship between

ATP8A2 change and behavior change is unclear. The current study revealed that

ATP8A2 expression in the PFC could be down-regulated by a high concentration of

IFN-γ, which is consistent with another study verifying that IFN-γ down-regulated the

expression of ATP8A2 in non-neuron cells (Shulzhenko et al.,2018). This finding

deepens the understanding of how depression is induced during infections by

LPS-producing gram-negative bacteria.

LPS exposure can induce neuroinflammation in various brain regions and result in

functional abnormalities, such as hippocampal neuroinflammation and impaired

learning and memory (Lee et al.,2008; Shaw et al.,2001). In this study, the animal

model that simulates a depressive-like phenotype caused by early bacterial infection

was used to investigate whether LPS caused alteration in ATP8A2 expression.

Therefore, PFC was selected as the brain region to observe ATP8A2 expression that

it's the area of the brain mostly focused on by studies about depressive-like behavior

(Myers-Schulz and Koenigs,2012). In future studies, we will investigate the

expression of ATP8A2 in other pathological conditions and/or in other brain zones

such as the hippocampus.

As well-known, LPS-induced inflammation is transient. Likewise, the first

experiment in this study has shown the ATP8A2 levels decreased transiently and

restored within 10 days after the end of LPS administration (Fig.1). Therefore, the

animals were killed after finishing behavioral tasks by over-anesthetized without

investigating their PFC ATP8A2 levels or peripheral/cerebral cytokines in adulthood.

Understandably, a transient ATP8A2 decrease in PFC might mediate a delayed

depressive-like behavior phenotype due to the existing theory of early-life

programming (Dinel et al.,2014; Karrow,2006), saying that the brain is susceptible to

external stimuli, such as immune activation, which modulates the course of normal

brain development.

It has been reported that sex could affect the outcomes in behavior development by

itself or combining neonatal immune activation (Cai et al.,2016), which is confirmed

by the current study due to the observed between-sexes effects in FST and TST tasks

performances (Supplementary Fig.2). However, the PFC ATP8A2 expression and

cytokine levels both in blood and in PFC have not been significantly influenced by the

sex factor tested at PND11 (Supplementary Fig.1). This may be because the sex factor

exerts its effects mainly by gonadal hormones. These endocrine sex differences often

come to be obvious from the beginning of puberty and these hormonal disparities

contribute to the emerging sex differences in the brain (Cai et al.,2016). Further study

is required to address whether the sex factor makes a difference in ATP8A2

expression in the brain during puberty as well as adulthood.

In sum, neonatal LPS exposure reduced ATP8A2 level in PFC in mice via

increasing IFN-γ level, which may be associated with mechanism underlying brain

and behavior impairments induced by neonatal LPS exposure.

Acknowledgments

We thank Mr. Taoqi Tao (ORCID: 0000-0002-2770-9568, from GDPU), Mrs.

Yinyin Xie (ORCID: 0000-0002-5858-3873, from GDPU) for their valuable

discussions and help with this investigation.

Funding Statement

The funders had no role in study design, data collection and interpretation, or the

decision to submit the work for publication.

Funding Information

This paper was supported by the following grants: 1) the National Natural Science

Foundation of China (No.31600836) to Junhua Yang; 2) the starting fund for

high-level talent introduction into Guangdong Pharmaceutical University

(No.51355093) to Junhua Yang; 3) the Innovation and University Promotion Project

of Guangdong Pharmaceutical University Through No. 2017KCXTD020 to the

School of Biosciences & Biopharmaceutics of GDPU as a team project funding; 4)the

National Natural Science Foundation of China (No.81901524) to Li Luo; 5) the

Natural Science Foundation of Guangdong Province (No.2018A030313579) to Li

Luo.

Additional information

Competing interests

JD, LS, ZY, SZ, ZD, LL, JL, XJ and JY declare that there are no conflicts of

interest.

Author contributions

JD: Data curation, Investigation, Methodology, Writing-original draft, Writing-review

and editing.

LS: Formal analysis, Investigation, Methodology.

ZY: Data curation, Formal analysis, Investigation, Methodology.

SZ: Validation, Investigation.

ZD: Validation, Investigation.

LL: Resources, Funding acquisition.

JL: Resources.

XJ: Resources.

JY: Conceptualization, Supervision, Funding acquisition, Writing - original draft,

Writing - review and editing.

References

PA Aaron, Jamklang, M, Uhrig, JP, and Gelli, A(2018).The blood-brain barrier internalises

Cryptococcus neoformans via the EphA2-tyrosine kinase receptor.*Cell Microbiol.*20.Scholar

JP Andersen, Vestergaard, AL, Mikkelsen, SA, Mogensen, LS, Chalat, M, and Molday,

RS(2016).P4-ATPases as Phospholipid Flippases-Structure, Function, and Enigmas.*Front*

*Physiol.*7.275.Google Scholar

SD Bilbo, and Schwarz, JM(2012).The immune system and developmental programming of

brain and behavior.*Front Neuroendocrinol.*33.267-286.Google Scholar

NJ Brandon, and Sawa, A(2011).Linking neurodevelopmental and synaptic theories of mental

illness through DISC1.*Nat Rev Neurosci.*12.707-722.Google Scholar

KC Cai, van Mil, S, Murray, E, Mallet, JF, Matar, C, and Ismail, N(2016).Age and sex

differences in immune response following LPS treatment in mice.*Brain Behav*

*Immun.*58.327-337.Google Scholar

AC Campos, Vaz, GN, Saito, VM, and Teixeira, AL(2014).Further evidence for the role of

interferon-gamma on anxiety- and depressive-like behaviors: involvement of hippocampal

neurogenesis and NGF production.*Neurosci Lett.*578.100-105.Google Scholar

J Chen, Wang, Z, Zhang, S, Ai, Q, Chu, S, and Chen, N-h(2017).Possible target-related

proteins of stress-resistant rats suggested by label-free proteomic analysis.*RSC*

*Advances.*7.40957-40964.Google Scholar

H Choi, Andersen, JP, and Molday, RS(2019).Expression and functional characterization of

missense mutations in ATP8A2 linked to severe neurological disorders.*Hum*

*Mutat.*40.2353-2364.Google Scholar

JA Coleman, Kwok, MC, and Molday, RS(2009).Localization, purification, and functional

reconstitution of the P4-ATPase Atp8a2, a phosphatidylserine flippase in photoreceptor disc

membranes.*J Biol Chem.*284.32670-32679.Google Scholar

JA Coleman, and Molday, RS(2011).Critical role of the beta-subunit CDC50A in the stable

expression, assembly, subcellular localization, and lipid transport activity of the P4-ATPase

ATP8A2.*J Biol Chem*.286.17205-17216.Google Scholar

AM Depino(2015).Early prenatal exposure to LPS results in anxiety- and depression-related

behaviors in adulthood.*Neuroscience*.299.56-65.Google Scholar

AL Dinel, Joffre, C, Trifilieff, P, Aubert, A, Foury, A, Le Ruyet, P, and Laye,

S(2014).Inflammation early in life is a vulnerability factor for emotional behavior at

adolescence and for lipopolysaccharide-induced spatial memory and neurogenesis alteration

at adulthood.*J Neuroinflammation*.11.155.Google Scholar

MH Doosti, Bakhtiari, A, Zare, P, Amani, M, Majidi-Zolbanin, N, Babri, S, and Salari,

AA(2013).Impacts of early intervention with fluoxetine following early neonatal immune

activation on depression-like behaviors and body weight in mice.*Prog Neuropsychopharmacol*

Biol Psychiatry.43.55-65.Google Scholar

R Gupta, Gupta, K, Tripathi, AK, Bhatia, MS, and Gupta, LK(2016).Effect of Mirtazapine

Treatment on Serum Levels of Brain-Derived Neurotrophic Factor and Tumor Necrosis

Factor-alpha in Patients of Major Depressive Disorder with Severe

Depression.*Pharmacology*.97.184-188.Google Scholar

R Haba, Shintani, N, Onaka, Y, Wang, H, Takenaga, R, Hayata, A, Baba, A, and Hashimoto,

H(2012).Lipopolysaccharide affects exploratory behaviors toward novel objects by impairing

cognition and/or motivation in mice: Possible role of activation of the central amygdala.*Behav*

Brain Res.228.423-431.Google Scholar

I Hansen-Pupp, Harling, S, Berg, AC, Cilio, C, Hellstrom-Westas, L, and Ley,

D(2005).Circulating interferon-gamma and white matter brain damage in preterm

infants.*Pediatr Res.*58.946-952.Google Scholar

K Hashimoto(2015).Inflammatory biomarkers as differential predictors of antidepressant

response.*Int J Mol Sci.*16.7796-7801.Google Scholar

NA Karrow(2006).Activation of the hypothalamic-pituitary-adrenal axis and autonomic nervous

system during inflammation and altered programming of the neuroendocrine-immune axis

during fetal and neonatal development: lessons learned from the model inflammagen,

lipopolysaccharide.*Brain Behav Immun.*20.144-158.Google Scholar

WB Klimstra, Ryman, KD, Bernard, KA, Nguyen, KB, Biron, CA, and Johnston,

RE(1999).Infection of neonatal mice with sindbis virus results in a systemic inflammatory

response syndrome.*J Virol.*73.10387-10398.Google Scholar

O Kohler, Benros, ME, Nordentoft, M, Farkouh, ME, Iyengar, RL, Mors, O, and Krogh,

J(2014).Effect of anti-inflammatory treatment on depression, depressive symptoms, and

adverse effects: a systematic review and meta-analysis of randomized clinical trials.*JAMA*

*Psychiatry.*71.1381-1391.Google Scholar

G Laumet, Edralin, JD, Chiang, AC, Dantzer, R, Heijnen, CJ, and Kavelaars,

A(2018).Resolution of inflammation-induced depression requires T lymphocytes and

endogenous brain interleukin-10 signaling.*Neuropsychopharmacology.*43.2597-2605.Google

Scholar

JW Lee, Lee, YK, Yuk, DY, Choi, DY, Ban, SB, Oh, KW, and Hong,

JT(2008).Neuro-inflammation induced by lipopolysaccharide causes cognitive impairment

through enhancement of beta-amyloid generation.*J Neuroinflammation.*5.37.Google Scholar

M Liang, Zhong, H, Rong, J, Li, Y, Zhu, C, Zhou, L, and Zhou, R(2019).Postnatal

Lipopolysaccharide Exposure Impairs Adult Neurogenesis and Causes Depression-like

Behaviors Through Astrocytes Activation Triggering GABAA Receptor

Downregulation.*Neuroscience.*422.21-31.Google Scholar

AJ Miller, Roman, B, and Norstrom, E(2016).A method for easily customizable gradient gel

electrophoresis.*Anal Biochem.*509.12-14.Google Scholar

B Myers-Schulz, and Koenigs, M(2012).Functional anatomy of ventromedial prefrontal cortex:

implications for mood and anxiety disorders.*Mol Psychiatry.*17.132-141.Google Scholar

B O'Reilly, Vander, AJ, and Kluger, MJ(1988).Effects of chronic infusion of lipopolysaccharide

on food intake and body temperature of the rat.*Physiol Behav.*42.287-291.Google Scholar

A Pokryszko-Dragan, Frydecka, I, Kosmaczewska, A, Ciszak, L, Bilinska, M, Gruszka, E,

Podemski, R, and Frydecka, D(2012).Stimulated peripheral production of interferon-gamma is

related to fatigue and depression in multiple sclerosis.*Clin Neurol*

*Neurosurg.*114.1153-1158.Google Scholar

AE Ross, Marchionni, L, Vuica-Ross, M, Cheadle, C, Fan, J, Berman, DM, and Schaeffer,

EM(2011).Gene expression pathways of high grade localized prostate

cancer.*Prostate.*71.1568-1577.Google Scholar

KN Shaw, Commins, S, and O'Mara, SM(2001).Lipopolysaccharide causes deficits in spatial

learning in the watermaze but not in BDNF expression in the rat dentate gyrus.*Behav Brain*

*Res.*124.47-54.Google Scholar

N Shulzhenko, Dong, X, Vyshenska, D, Greer, RL, Gurung, M, Vasquez-Perez, S,

Peremyslova, E, Sosnovtsev, S, Quezado, M, Yao, M, *et al.*(2018).CVID enteropathy is

characterized by exceeding low mucosal IgA levels and interferon-driven inflammation

possibly related to the presence of a pathobiont.*Clin Immunol.*197.139-153.Google Scholar

AK Walker, Budac, DP, Bisulco, S, Lee, AW, Smith, RA, Beenders, B, Kelley, KW, and

Dantzer, R(2013).NMDA receptor blockade by ketamine abrogates

lipopolysaccharide-induced depressive-like behavior in C57BL/6J

mice.*Neuropsychopharmacology.*38.1609-1616.Google Scholar

Z Wang, Li, W, Chen, J, Shi, H, Zhao, M, You, H, Rao, C, Zhan, Y, Yang, Y, and Xie,

P(2016).Proteomic analysis reveals energy metabolic dysfunction and neurogenesis in the

prefrontal cortex of a lipopolysaccharide-induced mouse model of depression.*Mol Med*

*Rep.*13.1813-1820.Google Scholar

D Wong, Dorovini-Zis, K, and Vincent, SR(2004).Cytokines, nitric oxide, and cGMP modulate

the permeability of an in vitro model of the human blood-brain barrier.*Exp*

*Neurol.*190.446-455.Google Scholar

B Xiong, Li, A, Lou, Y, Chen, S, Long, B, Peng, J, Yang, Z, Xu, T, Yang, X, Li, X, *et*

al.(2017).Precise Cerebral Vascular Atlas in Stereotaxic Coordinates of Whole Mouse

Brain.*Front Neuroanat.*11.128.Google Scholar

Q Xu, Yang, GY, Liu, N, Xu, P, Chen, YL, Zhou, Z, Luo, ZG, and Ding, X(2012).P4-ATPase

ATP8A2 acts in synergy with CDC50A to enhance neurite outgrowth.*FEBS*

*Lett.*586.1803-1812.Google Scholar

J Yang, Qi, F, Gu, H, Zou, J, Yang, Y, Yuan, Q, and Yao, Z(2016).Neonatal BCG vaccination

of mice improves neurogenesis and behavior in early life.*Brain Res Bull.*120.25-33.Google

Scholar

X Zhu, Libby, RT, de Vries, WN, Smith, RS, Wright, DL, Bronson, RT, Seburn, KL, and John,

SW(2012).Mutations in a P-type ATPase gene cause axonal degeneration.*PLoS*

*Genet.*8.e1002853.Google Scholar

Legends

Fig.1. Neonatal LPS exposure induced a transiently down-regulated expression

of ATP8A2 in the PFC in mice. (A) Representative results for the Western blot

analysis of ATP8A2. **(B)** The relative quantification of ATP8A2 in each group of mice

was normalized using the level of β-actin. Data are expressed as means ± SEM.

Randomly block design ANOVA; $n = 6$/group; $* p < 0.05$; $*** p < 0.001$.

Fig.2. Neonatal LPS exposure induced a down-regulated expression of ATP8A2

accompanied by inflammation both in periphery and in PFC in mice.

(A) Representative results for the Western blot analysis of ATP8A2. **(B)** The relative

quantification of ATP8A2 in each group of mice was normalized using the level of

β-actin. **(C-J)** The bars represent the average levels of IFN-γ, IL-1β, IL-6, TNF-α in

the serum and IFN-γ , IL-1β, IL-6, TNF-α in the PFC. Data are expressed as means ±

SEM. Randomly block design ANOVA; $n = 9$/group; $* p < 0.05$; $**p < 0.01$; $*** p <$

0.001.

Fig.3. The correlation of PFC ATP8A2 level with the serum and PFC levels of

pro-inflammatory cytokines in mice. (A) The positive negative correlation of the

PFC IFN-γ level with the ATP8A2 expression level in LPS-treated mice. **(B-D)**

Correlation analyses between the PFC IL-1β, IL-6 and TNF-α levels and the level of

ATP8A2 LPS-treated mice**. (E)** The positive negative correlation of the serum IFN-γ

level with the ATP8A2 expression level in LPS-treated mice. **(F-H)** Correlation

analyses between the serum IL-1β, IL-6 and TNF-α levels and the level of ATP8A2

LPS-treated mice**.** $n = 15$ per analysis ; *Pearson*'s correlation analysis.

Fig.4. Western blot analyses showed that IFN-γ mediated the PFC ATP8A2

down-regulation caused by neonatal LPS exposure.

(A-B) The mean levels of IFN-γ in the serum and PFC were shown from the

experiment carried out to determine the optimal dosage of anti-IFN-γ neutralizing

mAb. **(C-D)** The mean levels of IFN-γ in the serum and PFC were shown from the

IFN-γ-blocking experiment. **(E)** The relative quantification of ATP8A2 in each group

of mice was normalized using the level of β-actin. **(F)** Representative results for the

Western blot analysis of ATP8A2. Data are expressed as means ± SEM. Randomly

block design ANOVA followed by Tukey's *post hoc* test; $n = 8$/group; $**p < 0.01$;

$*** \, p < 0.001$; n.s., no significant. anti-IFN-γ(0.6), the anti-IFN-γ dosage of 0.6

mg/kg body weight.

Fig.5. Immunofluorescence analyses showed that IFN-γ mediated the PFC

ATP8A2 down-regulation caused by neonatal LPS exposure. (A-C) Representative

immunofluorescence staining results in the PFC for ATP8A2(red)/NeuN(green)

co-labeling in CON group (A), LPS group (B) and LPS+anti-IFN-γ(0.6) group (C).

(E-F) Exhibition of the single red channel that shows ATP8A2 signal of the same

micrographs shown in (A), (B), (C), respectively. **(G-I)** Representative high

magnificent photos show the immunofluorescence staining results in the PFC for

ATP8A2/NeuN co-labeling. Thick arrows indicate ATP8A2/NeuN co-labeling cells.

Thin arrows indicate NeuN single labeling cells. **(J)** The larger scope shows the

location in PFC where the photos shown in (G-I) come from. **(K)** Bars represent mean

fluorescence intensity of the signal of immunofluorescence staining for ATP8A2 in

each group. **(I)** Bars represent mean numbers of ATP8A2/NeuN co-labeling cells in

unilateral PFC in each group. Data are expressed as means ± SEM. Randomly block

design ANOVA followed by Tukey's *post hoc* test; $n = 6$/group; *** $p < 0.001$;

anti-IFN-γ(0.6), the anti-IFN-γ dosage of 0.6 mg/kg body weight. Scale bar in (A-F),

300 μm; in (G-I), 10 μm; in (J), 200 μm.

Fig.6. IFN-γ plays an important role in depressive-like behaviors in adulthood

induced by neonatal LPS exposure. **(A)** Bars represent the mean immobility time of

mice in FST of each group. **(I)** Bars represent the mean immobility time of TST in

each group. The data represent the mean ± SEM; randomly block design ANOVA

followed by Tukey's *post hoc* test; $n = 12$/group; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$;

anti-IFN-γ(0.6), the anti-IFN-γ dosage of 0.6 mg/kg body weight.

NeuN/ATP8A2

○ CON
● LPS
● LPS+anti
-IFN-γ(0.6)